

Metabolic memory of $\Delta 9$ -tetrahydrocannabinol exposure in pluripotent stem cells and primordial germ cells-like cells

Reviewed Preprint

Revised by authors after peer review.

[About eLife's process](#)

Reviewed preprint version 2

November 20, 2023 (this version)

Reviewed preprint version 1

July 13, 2023

Sent for peer review

May 16, 2023

Posted to preprint server

March 15, 2023

Roxane Verdikt, Abigail A. Armstrong, Jenny Cheng, Young Sun Hwang, Amander T. Clark, Xia Yang, Patrick Allard 

Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA 90095, USA • Department of Obstetrics/Gynecology and Reproductive Endocrinology and Infertility, University of California, Los Angeles, CA, USA • Molecular, Cellular, and Integrative Physiology Graduate Program, University of California, Los Angeles, Los Angeles, CA 90095, USA • Department of Molecular Cell and Developmental Biology, University of California, Los Angeles, Los Angeles, CA, 90095, USA • Center for Reproductive Science, Health and Education, University of California, Los Angeles, Los Angeles, CA 90095, USA • Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California, Los Angeles, Los Angeles, CA, 90095, USA • Integrative Biology and Physiology Department, University of California, Los Angeles, CA, 90095, USA • Department of Molecular and Medical Pharmacology, University of California, Los Angeles, Los Angeles, CA 90095, USA • Molecular Biology Institute, University of California, Los Angeles, Los Angeles, CA 90095, USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Cannabis, the most consumed illicit psychoactive drug in the world, is increasingly used by pregnant women. However, while cannabinoid receptors are expressed in the early embryo, the impact of phytocannabinoids exposure on early embryonic processes is lacking. Here, we leverage a stepwise in vitro differentiation system that captures early embryonic developmental cascade to investigate the impact of exposure to the most abundant phytocannabinoid, $\Delta 9$ -tetrahydrocannabinol ($\Delta 9$ -THC). We demonstrate that $\Delta 9$ -THC increases the proliferation of naïve mouse embryonic stem cells (ESCs) but not of their primed counterpart. Surprisingly, this increased proliferation, dependent on the CB1 receptor binding, is only associated with moderate transcriptomic changes. Instead, $\Delta 9$ -THC capitalizes on ESCs' metabolic bivalence by increasing their glycolytic rates and anabolic capabilities. A memory of this metabolic rewiring is retained throughout differentiation to Primordial Germ Cell-Like Cells in the absence of direct exposure and is associated with an alteration of their transcriptional profile. These results represent the first in-depth molecular characterization of the impact of $\Delta 9$ -THC exposure on early stages of germline development.

eLife assessment

The study presents **valuable** findings demonstrating that physiologically relevant concentrations delta-9-tetrahydrocannabinol, which is found in cannabis, have metabolic effects on early mouse embryonic cell types. The evidence supporting the claims is **convincing**. The work will be of interest to researchers in stem cell and epigenetics fields.

Introduction

Cannabis is the most widely used illicit psychoactive drug in the world (U.N. Office on Drugs and Crime, 2022 [↗](#)). In the United States, an estimated 49.6 million people, roughly 18% of the population, consumed cannabis at least once in 2020, with indications that these numbers will likely increase in the coming years as attitudes and regulations change (Mennis et al., 2023 [↗](#); Substance Abuse and Mental Health Services Administration, 2020 [↗](#)). In particular, between 7-12% of expecting women report cannabis use, predominantly during the first trimester to alleviate the symptoms of morning sickness (Chabarria et al., 2016 [↗](#); Volkow et al., 2019 [↗](#); Young-Wolff et al., 2018 [↗](#)). These statistics indicate that a significant number of developing embryos are exposed to cannabis, with limited knowledge of the biological repercussions of such exposure.

Among the several hundred unique phytocannabinoids present in *Cannabis sativa*, (-)-trans- Δ^9 -tetrahydrocannabinol (Δ^9 -THC) is chiefly responsible for the psychoactivity of cannabis (Andre et al., 2016 [↗](#)). As a result, the level of Δ^9 -THC in recreational cannabis has increased over the last 10 years and now commonly accounts for 20% of total compounds (Chandra et al., 2019 [↗](#)). The psychoactive effects of Δ^9 -THC arise from its binding and subsequent activation of the G protein-coupled cannabinoid receptors CB1 largely expressed in the central nervous system (Pacher et al., 2020 [↗](#)). In this context, Δ^9 -THC exposure has been shown to durably alter metabolic, transcriptional and epigenetic programs in the brain (Bénard et al., 2012 [↗](#); Prini et al., 2017 [↗](#); Szutorisz and Hurd, 2018 [↗](#); Watson et al., 2015 [↗](#)). While over the last decades, significant attention has been paid to Δ^9 -THC's neurological effects, there is also evidence, albeit more limited, of its impact on reproductive functions (Lo et al., 2022 [↗](#)). Data shows CB1 expression in the male and the female reproductive tracts, in the pre-implantation embryo and in the placenta (Lo et al., 2022 [↗](#); Paria et al., 1995 [↗](#)). In animal models as well as in humans, exposure to cannabis is associated with reduced fertility, decreased testis weight and sperm count, and impairment of embryo implantation (Lo et al., 2022 [↗](#)). In males, these effects are correlated with an alteration of the sperm transcriptome and epigenome (Murphy et al., 2018 [↗](#); Osborne et al., 2020 [↗](#); Schrott and Murphy, 2020 [↗](#)). Epidemiological evidence also indicates that Δ^9 -THC exposure is associated with long-lasting adverse effects, with exposures in parents affecting the offspring (Smith et al., 2020 [↗](#); Szutorisz and Hurd, 2018 [↗](#)). Despite this accumulating evidence, the molecular impact and mechanisms of Δ^9 -THC exposure at the earliest stages of germ cells development remain to be determined.

Progression through states of pluripotency is controlled by metabolic reprogramming in the early mammalian embryo (Verdikt and Allard, 2021 [↗](#); Zhang et al., 2018 [↗](#)). Accordingly, cultured pluripotent stem cells (PSCs) exhibiting different developmental potentials are marked by specific metabolic signatures, similar to the ones displayed by their in vivo counterparts in the embryo. For instance, mouse embryonic stem cells (ESCs) are naïve PSCs that are functionally equivalent to the inner cell mass (ICM) of the E3.5 preimplantation mouse blastocyst (Nichols and Smith, 2009 [↗](#)). The extended developmental potential of mouse ESCs is associated with their metabolic bivalence, as these cells rely on both glycolysis and oxidative phosphorylation for energy

production. Differentiation of naïve ESCs into primed PSCs such as epiblast-like cells (EpiLCs) is accompanied by an important metabolic shift towards aerobic glycolysis, in link with a highly-proliferative phenotype and a more restricted developmental potential (Hayashi et al., 2017 [↗](#); Verdikt and Allard, 2021 [↗](#); Zhang et al., 2018 [↗](#)). Primordial germ cells (PGCs), the embryonic precursors of gametes in metazoans (Kurimoto and Saitou, 2018 [↗](#)), are considered dormant totipotent cells because they possess the unique ability to reacquire totipotency upon fertilization (Hayashi et al., 2017 [↗](#)). In the mouse embryo, the precursors to PGCs arise around embryonic day 6.25 (E6.25) from formative pluripotent cells in the epiblast (Kinoshita et al., 2021 [↗](#); Kurimoto and Saitou, 2018 [↗](#)). Progressive increase in oxidative phosphorylation correlates with the specification and differentiation from epiblast PSCs towards PGCs, a process that can be replicated in vitro by inducing PGC-like cells (PGCLCs) from EpiLCs (Hayashi et al., 2011 [↗](#)). In particular, the extensive metabolic, transcriptional, and epigenetic reprogramming that PGCs undergo during their development has been proposed to be uniquely sensitive to environmental insults, with potential consequences in the offspring (Verdikt et al., 2022 [↗](#)).

Here, we deployed this in vitro differentiation system to investigate the impact of $\Delta 9$ -THC exposure on early developmental stages. We demonstrate that exposure of ESCs and EpiLCs to $\Delta 9$ -THC durably alters their metabolome. We reveal that, in the absence of continuous exposure, metabolic memory of $\Delta 9$ -THC is passed onto the PGCLCs stage leading to transcriptional defects in these cells. Together, our findings highlight the role of metabolic reprogramming as a mechanism for early developmental $\Delta 9$ -THC exposure.

Results

$\Delta 9$ -THC induces cellular proliferation of mouse embryonic stem cells but not of mouse epiblast-like cells

To model the impact of early $\Delta 9$ -THC exposure on early embryonic events, we first tested three distinct developmental windows: 1) exposure of ESCs, 2) exposure of EpiLCs and 3) combined ESCs+EpiLCs exposure (Figure 1A [↗](#)). Cells were either exposed to the vehicle (mock) or exposed to $\Delta 9$ -THC in a wide dose range of 0.1nM-100 μ M, corresponding to the reported physiologically-relevant concentrations of $\Delta 9$ -THC in cannabis users (Fuchs Weizman et al., 2021 [↗](#); Hunault et al., 2008 [↗](#); Pacifici et al., 2019 [↗](#)).

The viability of ESCs exposed to increasing concentrations of $\Delta 9$ -THC for 48h was not significantly altered until the maximal dose of 100 μ M, subsequently serving as a positive control, at which only 13.17% of cells remained alive (Figure 1B [↗](#), $p=5.10^{-15}$, unpaired T-test). While no significant changes in viability were observed between 10nM and 1 μ M $\Delta 9$ -THC, the number of viable ESCs significantly increased by 1.69, 1.52 and 1.28-fold, respectively, compared to the mock-treated condition (Figure 1C [↗](#), $p=0.002$, $p=0.01$ and $p=0.03$ for 10nM, 100nM and 1 μ M of $\Delta 9$ -THC, unpaired T-test). Of note, the number of viable ESCs significantly increased for doses of $\Delta 9$ -THC as low as 1nM (Supplementary Figure 1 [↗](#), 1.59-fold increase, $p=0.0005$, unpaired T-test), corresponding to doses found in users of “light cannabis” products (Pacifici et al., 2019 [↗](#)). To determine whether the increased number of viable cells recovered after $\Delta 9$ -THC exposure was due to higher proliferation, we performed bromodeoxyuridine (BrdU) labeling experiments. Exposed cells were pulsed with BrdU for 30 minutes, and its incorporation in actively dividing cells was measured by flow cytometry. The percentage of BrdU-positive ESCs significantly increased between 10nM and 10 μ M of $\Delta 9$ -THC compared to the mock-treated condition (Figure 1D [↗](#), $p=0.01$, $p=0.001$, $p=0.05$ and $p=0.01$ for 10nM, 100nM, 1 μ M and 10 μ M of $\Delta 9$ -THC, unpaired T-test). Co-staining with DAPI (4',6-diamidino-2-phenylindole) to analyze the cell cycle showed that, at 100nM of $\Delta 9$ -THC, a significantly higher proportion of cells were in the G2/M phase (Supplementary Figure 2 [↗](#), 2.76-fold increase, $p=0.02$, unpaired T-test), consistent with cellular proliferation. Finally, whereas the




Figure 1

Δ9-THC exposure provokes the proliferation of ESCs but not EpiLCs

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. bFGF: basic fibroblast growth factor, ESCs, embryonic stem cells; EpiLCs, epiblast-like cells; LIF, leukemia inhibitory factor. **(B, E, H)** Whisker boxplot indicating the median cellular viability of stem cells exposed to the different Δ9-THC doses and associated errors. **(C, F, I)** Whisker boxplot indicating the median number of viable cells exposed to the different Δ9-THC doses indicated and associated errors. **(D, G, J)** Whisker boxplot indicating the median percentage of BrdU-stained cells exposed to the different Δ9-THC doses and associated errors. ESCs exposed cells are presented in (B, C and D). EpiLCs exposed cells deriving from unexposed ESCs are presented in (E, F and G). EpiLCs exposed cells deriving from exposed ESCs are presented in (H, I and J). At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical significance: * (p<0.05), ** (p<0.01), *** (p<0.001), **** (p<0.0001).

present data was obtained using female ESCs, the proliferative effect of $\Delta 9$ -THC exposure on ESCs was also observed in male ESCs (**Supplementary Figure 3**), suggesting that proliferation is sex-independent.

Next, we derived EpiLCs from unexposed ESCs and performed the same dose-response experiments. Akin to ESCs, $\Delta 9$ -THC exposure in EpiLCs did not significantly alter cellular viability until the dose of 100 μ M (**Figure 1E**, 11.56% of viable cells, $p=1.6^{-13}$, unpaired T-test). However, contrary to ESCs, $\Delta 9$ -THC exposure in EpiLCs did not significantly increase the number of viable cells nor the percentage of BrdU-positive cells (**Figure 1F** and **Figure 1G**). For the dose of 10 μ M of $\Delta 9$ -THC, viable EpiLCs numbers and BrdU-positive EpiLCs decreased compared to the mock-treated condition (**Figure 1F**, 1.53-fold decrease, $p=0.0007$, **Figure 1G**, 1.42-fold decrease, $p=0.0024$, unpaired T-test). When continuously exposing ESCs and EpiLCs, cell viability was more significantly and negatively impacted, except at the dose of 100nM of $\Delta 9$ -THC (**Figure 1H**). Deriving EpiLCs from exposed ESCs and exposing them to $\Delta 9$ -THC for 48h did not significantly affect either their cell number nor their incorporation of BrdU (**Figure 1I** and **Figure 1J**), indicating that the increased proliferation observed at the ESCs stage is not carried through the naïve-to-prime transition.

Finally, we further assessed the impact of $\Delta 9$ -THC exposure in human pluripotent stem cells. Human embryonic stem cells (hESCs) differ from their murine counterparts, particularly because they resemble a primed pluripotent state, akin to mouse EpiLCs (Weinberger et al., 2016). Cell viability of hESCs continuously exposed to 100nM of $\Delta 9$ -THC was not significantly impacted (**Supplementary Figure 4A**). However, hESCs number was significantly decreased upon $\Delta 9$ -THC exposure (**Supplementary Figure 4B**, 1.26-fold decrease, $p=0.004$, unpaired T-test), in a similar trend as observed with mouse EpiLCs (**Figure 1E** and **Figure 1I**).

Together, the systematic testing of different exposure schemes of $\Delta 9$ -THC in different pluripotent stem cell populations (mouse and human ESCs and mouse EpiLCs) revealed that physiologically relevant doses of $\Delta 9$ -THC (10nM-1 μ M) specifically stimulate the proliferation of mouse ESCs, but not of human ESCs, nor of mouse EpiLCs, whether the latter are derived from exposed ESCs or not.

Expression of the CB1 receptor does not explain differences in proliferative outcomes

We next sought to understand the source of variation in proliferative outcomes in response to $\Delta 9$ -THC between naïve mouse embryonic stem cells and primed pluripotent epiblast-like cells. Such differential effects have been previously reported, with $\Delta 9$ -THC eliciting the proliferation of neural progenitors (Galve-Roperh et al., 2013) and of human breast carcinoma cell lines (Takeda et al., 2008) but suppressing the proliferation of activated CD4⁺ T cells (Yang et al., 2016) and of non-small cell lung cancer cells (Preet et al., 2008). In these studies, the differential expression of cannabinoid receptors at the cell surface was proposed to primarily mediate the variation in cellular outcomes.

We therefore first tested whether expression levels of CB1 varied between ESCs and EpiLCs. Western-blot analysis of membrane proteins revealed however that CB1 was expressed at the same levels at the cell surface of both ESCs and EpiLCs (**Figure 2A** and **2B**). We next determined whether the $\Delta 9$ -THC-induced proliferative phenotype in ESCs was due to the engagement of the CB1 cannabinoid receptor. To do so, ESCs were pretreated for 1h with 1 μ M of SR141716 (also known as rimonabant, a specific CB1 blocker (Rinaldi-Carmona et al., 1994)) then exposed to 100nM or 100 μ M of $\Delta 9$ -THC for 48h. Rimonabant pre-treatment did not significantly alter the viability of ESCs compared to conditions exposed to $\Delta 9$ -THC only (**Figure 2C**) but abolished $\Delta 9$ -THC-induced ESCs increased cell number at 100nM $\Delta 9$ -THC (**Figure 2D**, 1.53-fold

decrease, $p=1.66^{-05}$, when comparing 100nM of $\Delta 9$ -THC +/- 1 μ M of SR141716, unpaired T-test). Notably, SR141716 pre-treatment, while not altering cell viability, reduced cell number compared to control, suggesting a basal role for CB1 in promoting proliferation.

Thus, the expression of CB1 at the cell surface does not explain the differential impact of $\Delta 9$ -THC on ESC and EpiLC proliferation even if CB1 engagement is a required event for this effect in ESCs.

$\Delta 9$ -THC exposure increases glycolysis in ESCs and EpiLCs

In the central nervous system, $\Delta 9$ -THC is a known metabolic perturbator which increases bioenergetic metabolism (Bartova and Birmingham, 1976 [↗](#); Bénard et al., 2012 [↗](#)). As mentioned above, the transition of naïve ESCs into the primed state of EpiLCs is accompanied by a switch to glycolysis for energy production (Hayashi et al., 2017 [↗](#); Verdikt and Allard, 2021 [↗](#)). Thus, to capture the impact of $\Delta 9$ -THC at every point of their transition between metabolic states, we used the continuous exposure scheme of ESCs and EpiLCs outlined in **Figure 1H-J** [↗](#). Similar to our other exposure schemes, at lower $\Delta 9$ -THC doses, the proliferation of ESCs was observed but not of EpiLCs. We performed these exposures in a wide $\Delta 9$ -THC dose range (10nM-10 μ M) followed by bioenergetics assessment (**Figure 3A** [↗](#)).




Figure 2

Implication of the CB1 receptor in the proliferative phenotype.

(A) Western blot analysis of transmembrane protein extracts of ESCs or EpiLCs. Antibodies raised against CB1 or β-actin serving as a loading control were used for immunoblotting. (B) Quantification of the gel presented in (A) was done using Image Studio (version 5.2). (C) Whisker boxplot indicating the median cellular viability of stem cells exposed to the different Δ9-THC and rimonabant doses indicated and their associated errors. (D) The median numbers of viable cells exposed to the different Δ9-THC and rimonabant doses indicated were normalized to their own control (+/- rimonabant). Median and associated errors were plotted in whisker boxplots. At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical significance: **($p < 0.01$), ****($p < 0.0001$).




Figure 3

$\Delta 9$ -THC exposure provokes an increase in glycolytic rates in ESCs and EpiLCs.

(A) Diagram illustrating $\Delta 9$ -THC exposure scheme and experimental strategy. **(B)** The NAD(P)⁺/NADPH ratio of stem cells exposed to the different $\Delta 9$ -THC doses was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots. **(C)** Mean fluorescence intensity (MFI) associated with the Mitotracker CMXRos stain was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots. **(D)** Median and associated error of the maximal extracellular acidification rate (ECAR) measured in cells exposed to the different $\Delta 9$ -THC doses and normalized to the protein content was plotted in whisker boxplots. **(E)** Median and associated error of the maximal oxygen consumption rate (OCR) measured in cells exposed to the different $\Delta 9$ -THC doses and normalized to the protein content was plotted in whisker boxplots. For (B and C), 5 technical repeats of 3 biological repeats (n=15) were plotted. One same representative experiment out of three independent experiments was used to plot results in (D and E). Statistical significance: *(p<0.05), **(p<0.01), *** (p<0.001), **** (p<0.0001).

First, we assessed the global energy metabolism of exposed cells by measuring the nicotinamide adenine dinucleotide (phosphate) couple ratios (NAD(P)⁺/NAD(P)H) using the WST-1 assay. In ESCs, the ratio of NAD(P)⁺/NAD(P)H significantly increased 1.57, 1.54, 1.29 and 1.38 -fold, for 10nM, 100nM, 1 μ M and 10 μ M of $\Delta 9$ -THC, respectively, compared to the mock-treated condition (**Figure 3B**, p=2.87⁻⁰⁶, p=8.01⁻⁰⁵, p=0.03, and p=0.0003 for 10nM, 100nM, 1 μ M and 10 μ M of $\Delta 9$ -THC, unpaired T-test). In contrast, no significant increase was observed in NAD(P)⁺/NAD(P)H ratios in exposed EpiLCs (**Figure 3B**). Consistent with the impact of continuous $\Delta 9$ -THC exposure on EpiLCs viability (**Figure 1H**), the NAD(P)⁺/NAD(P)H ratios significantly decreased at 10 μ M of $\Delta 9$ -THC in EpiLCs (**Figure 3B**, 59% decrease for 10 μ M of $\Delta 9$ -THC compared to the mock-treated condition, p=6.55⁻⁰⁹, unpaired T-test). Of note, the NAD(P)⁺/NAD(P)H ratios were slightly but significantly decreased at 100nM of $\Delta 9$ -THC in hESCs (**Supplementary Figure 5**, 1.09-fold decrease, p=0.04, unpaired T-test), in agreement with the deleterious effect of $\Delta 9$ -THC on cell number (**Supplementary Figure 4B**).

Because the elevated NAD(P)⁺/NAD(P)H levels in $\Delta 9$ -THC-exposed mouse ESCs could indicate increased mitochondrial activity in the context of oxidative phosphorylation (Locasale and Cantley, 2011), we next studied changes in mitochondrial membrane potential of exposed cells using the Mitotracker CMXRos fluorescent dye (Pendergrass et al., 2004). A significant increase in mean fluorescence intensity (MFI) associated with the mitochondrial stain was observed at 100nM of $\Delta 9$ -THC in ESCs (**Figure 3C**, p=0.02, unpaired T-test), indicating that, at this dose, the observed increase in NAD(P)⁺/NAD(P)H could be explained by higher mitochondrial membrane potential. By contrast, no change in EpiLCs mitochondrial activity was detected (**Figure 3C**), consistent with these cells relying on glycolysis for energy production (Hayashi et al., 2017; Verdikt and Allard, 2021).

Changes in mitochondrial activity in ESCs upon $\Delta 9$ -THC exposure, although significant, remained modest and are unlikely to be the sole contributor to the more significant increase in NAD(P)⁺/NAD(P)H upon exposure. Thus, we performed an in-depth analysis of the differential impact of $\Delta 9$ -THC on ESCs and EpiLCs bioenergetics by measuring both glycolysis (extracellular acidification rate, ECAR) and mitochondrial respiration (oxygen consumption rate, OCR) using a Seahorse bioanalyzer. At 100nM of $\Delta 9$ -THC, the maximal glycolytic capacity of both ESCs and EpiLCs increased significantly (**Figure 3D**, 15% increase, p=0.03 and 22% increase, p=0.03 for ESCs and EpiLCs, respectively, compared to the mock-treated condition, unpaired T-test). In both cell types, a significant decrease in glycolytic capacity was observed at 10 μ M of $\Delta 9$ -THC (**Figure 3D**, 39.8% reduction, p=0.0006 and 44.8% reduction, p=0.0001, for ESCs and EpiLCs, respectively, compared to the mock-treated condition, unpaired T-test). Of note, the maximal glycolytic capacity

of EpiLCs in the untreated condition was higher than the one of ESCs, in agreement with their metabolic shift towards aerobic glycolysis (**Figure 3D** [↗](#), 7.88% higher ECAR rate in mock-treated EpiLCs compared to mock-treated ESCs, $p=0.03$, unpaired T-test). As a consequence, $\Delta 9$ -THC exposure significantly impacted more glycolysis in EpiLCs than ESCs, both in basal capacity and upon mitochondrial inhibition by oligomycin (**Supplementary Figure 6A** [↗](#) and **Figure 6B** [↗](#)). In addition, at 100nM of $\Delta 9$ -THC, the maximal respiratory capacity of ESCs was significantly increased compared to the mock-treated condition (**Figure 3E** [↗](#), 21.8% increase, $p=0.03$, unpaired T-test). This increase was observed only for the maximal respiratory capacity of ESCs, but not for basal respiration, nor for ATP-linked respiration (**Supplementary Figure 6C** [↗](#)), suggesting that $\Delta 9$ -THC impact on mitochondrial respiration does not support increased energetic production. In agreement with EpiLCs metabolic shift towards a glycolytic phenotype, increasing doses of $\Delta 9$ -THC did not alter their maximal respiratory capacity (**Figure 3E** [↗](#)), nor their global oxygen consumption rate (**Supplementary Figure 6D** [↗](#)). In both cell types, a significant decrease in oxygen consumption rate was observed at 10 μ M of $\Delta 9$ -THC (**Figure 3E** [↗](#) and **Supplementary Figure 6C** [↗](#) and **Figure 1D** [↗](#)).

Together, our analysis of cellular bioenergetics following $\Delta 9$ -THC exposure showed an increased glycolytic rate in ESCs that was also observed in EpiLCs. However, the increased oxygen consumption and the associated increase in mitochondrial activity were observed only in ESCs following exposure to 100nM of $\Delta 9$ -THC, likely for the oxidization of the accumulating pyruvate generated from glycolysis.

$\Delta 9$ -THC-induced increase in glycolysis supports anabolism and ESCs proliferation

Because our data indicated that the impact of $\Delta 9$ -THC exposure on stem cells' bioenergetics did not result in greater ATP production, we next sought to characterize the global metabolic impact of $\Delta 9$ -THC in these cells. ESCs and EpiLCs were continuously exposed to 100nM $\Delta 9$ -THC and intracellular metabolites were detected and quantified by mass spectrometry (**Figure 4A-E** [↗](#)). To explore the metabolic signatures in the different samples, we performed a global principal component analysis (PCA) (**Figure 4B** [↗](#)). All samples clustered in well-defined groups of replicates, both by cell type on the first principal component (accounting for 65.81% of the variation) and by $\Delta 9$ -THC exposure on the second principal component (accounting for 20.83% of the variation). Of the 126 metabolites detected in ESCs, 39 were significantly upregulated (**Figure 4C** [↗](#) and **Supplementary Figure 7A** [↗](#)) and only two metabolites – NADPH and Adenine – were significantly downregulated. Of the 138 metabolites detected in EpiLCs, 95 were significantly upregulated (**Figure 4C** [↗](#) and **Supplementary Figure 7B** [↗](#)) and only one metabolite – NADPH – was significantly downregulated. In agreement with the PCA, the overlap of over-expressed metabolites in response to $\Delta 9$ -THC exposure was important between the two stem cell populations (**Figure 4C** [↗](#), accounting for 79.49% and 32.63% of all upregulated metabolites in ESCs and EpiLCs, respectively). The functional interpretation of the significantly upregulated metabolites confirmed the $\Delta 9$ -THC-associated increase in energy metabolism in the two stem cell populations. Indeed, amongst the 25 metabolic pathways upregulated, pyruvate metabolism and glycolysis were detected in both ESCs and EpiLCs (**Figure 4D** [↗](#) and **Figure 4E** [↗](#), respectively). Increased mitochondrial respiration was also seen in ESCs with the enrichment of (ubi)quinone metabolism, indicating an increased synthesis of ubiquinone that serves as an electron carrier in oxidative phosphorylation. Of note, metabolite measurements showed that the ratio of glutathione in its reduced to oxidized form (GSH/GSSG) was unchanged in both stem cell types in response to $\Delta 9$ -THC (**Supplementary Figure 7C** [↗](#)), suggesting that the increased mitochondrial respiration does not cause an overt elevation of oxidative stress. Importantly, and in agreement with the PCA, in both ESCs and EpiLCs, $\Delta 9$ -THC exposure elicited an increase in metabolic pathways that feed anabolic reactions, in particular contributing to the synthesis of amino acids (tyrosine, tryptophan, arginine, alanine, valine,

(iso)leucine, etc.), nucleotides (“Pyrimidine metabolism”, “Purine metabolism”), NAD(P)+ (“Nicotinate and nicotinamide metabolism”) and fatty acids (“Butanoate metabolism”) (**Figure 4D** [↗](#) and **Figure 4E** [↗](#)).

Extensive metabolic profiling of ESCs and EpiLCs upon $\Delta 9$ -THC exposure thus indicated that the increased glycolytic rates in both stem cell populations, rather than provoking an increased production of energy under the form of ATP, participated in increased anabolism. Such increased anabolism could explain the proliferation observed in ESCs upon $\Delta 9$ -THC exposure. To test this hypothesis, we exposed ESCs to 100nM of $\Delta 9$ -THC for 48h as above but 24h before the harvest, cells were exposed to 10mM of 2-deoxyglucose (2-DG), an inhibitor of glycolysis (Barban and Schulze, 1961 [↗](#)). Despite increasing the energy stress (**Supplementary Figure 8** [↗](#)), inhibition of glycolysis by 2-DG did not significantly impact viability over this shorter time frame and at this concentration (**Figure 4F** [↗](#)). Importantly, glycolytic inhibition by 2-DG abrogated the $\Delta 9$ -THC-induced increase in both cell number and NAD(P)+/NAD(P)H levels (**Figure 4G** [↗](#) and **Figure 4H** [↗](#), 1.39-fold reduction and $p=5.56 \cdot 10^{-5}$ and 1.68-fold reduction and $p=0.0064$, respectively, when comparing 100nM of $\Delta 9$ -THC +/- 10mM 2-DG, unpaired T-test). Thus, exposure to $\Delta 9$ -THC increases anabolism in both ESCs and EpiLCs, however, this increased anabolism only supports cellular proliferation in ESCs.

$\Delta 9$ -THC exposure is associated with the upregulation of genes involved in anabolic pathways in ESCs but not in EpiLCs

Our data shows that $\Delta 9$ -THC exposure increases anabolic pathways in both ESCs and EpiLCs and that this causes the proliferation of ESCs but not of EpiLCs. We thus next examined whether this differential impact of $\Delta 9$ -THC on ESCs and EpiLCs was mirrored by a change in these cells' transcriptomes. To this aim, we performed RNA-sequencing (RNA-seq) on ESCs and EpiLCs continuously exposed to 100nM $\Delta 9$ -THC or to the vehicle control (**Figure 5A** [↗](#)).

Unsupervised exploration of the global transcriptome by PCA revealed that the vast majority of data variation could be attributed to the cell type (PC1, accounting for 98% of the variation) rather than to $\Delta 9$ -THC exposure (PC2, accounting for 1% of the variation, **Figure 5B** [↗](#)). In addition, mining for the expression of pluripotency markers after $\Delta 9$ -THC exposure suggested that $\Delta 9$ -THC does not appear to influence the differentiation dynamics from ESCs to EpiLCs (**Supplementary Figure 9** [↗](#)). We identified a low number of differentially expressed genes (DEGs) in both ESCs and EpiLCs (**Figure 5C** [↗](#) and **Figure 5D** [↗](#), respectively), indicating that $\Delta 9$ -THC exposure only moderately impacts ESC and EpiLC transcriptomes. In ESCs, only 12 genes were significantly upregulated with a $\log_2(\text{fold-change}) > 0.5$ and only 9 were significantly downregulated at the same threshold (**Figure 5C** [↗](#), significance corresponds to adjusted $p\text{-value} \leq 0.05$). More genes were differentially expressed when looking at lower fold-changes ($|\log_2(\text{fold-change})| > 0.25$, **Figure 5C** [↗](#)), confirming that the magnitude of transcriptional effects due to $\Delta 9$ -THC exposure is moderate. This low transcriptional impact following $\Delta 9$ -THC exposure was also observed in EpiLCs (**Figure 5D** [↗](#)). Nevertheless, gene ontologies (GO) associated with $\Delta 9$ -THC-induced DEGs revealed the biological significance of these low transcriptional changes (**Figure 5E** [↗](#)). In particular, GO terms associated with metabolic pathways involved in anabolism were significantly over-represented for upregulated genes in ESCs following $\Delta 9$ -THC exposure (**Figure 5E** [↗](#)), such as: “Cellular aromatic compound metabolic process”, “Cellular nitrogen compound biosynthetic process”, “Organonitrogen compound metabolic process”. This suggests that the glycolytic rewiring elicited by $\Delta 9$ -THC exposure in ESCs has some transcriptional support. Indeed, when performing joint pathway integration between our transcriptomics data and our targeted metabolomics (Pang et al., 2022 [↗](#)), we observed that $\Delta 9$ -THC-induced perturbed genes and metabolites were associated with the observed anabolic effects (**Figure 5F** [↗](#)). In contrast, GO terms associated with metabolism were not found within the upregulated DEGs in EpiLCs. However, several GO terms relating to alterations in cellular components were enriched in EpiLCs (**Figure 5E** [↗](#)), such as: “Organelle organization”, “Cellular component organization or biogenesis”,




Figure 4

Δ9-THC-induced glycolysis sustain anabolism and ESCs proliferation

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. (B) PCA of the metabolomics profiling of either ESCs or EpiLCs mock-exposed or exposed to 100nM Δ9-THC. (C) Venn diagram showing the overlap in upregulated metabolites following Δ9-THC exposure in ESCs and EpiLCs. (D and E) KEGG metabolite sets enrichment analysis for upregulated metabolites in ESCs and EpiLCs, respectively, performed by MetaboAnalyst³⁸. KEGG, Kyoto Encyclopedia of Genes and Genomes. (F) Whisker boxplot indicating the median cellular viability of stem cells exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, and their associated errors. (G) The median numbers of viable cells exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, were normalized to their own control (+/- 2-DG). Median and associated errors were plotted in whisker boxplots. (H) The NAD(P)⁺/NAD(P)H ratio of stem cells exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, was normalized to the one measured in the mock-treated condition (+/- 2-DG). Median and associated errors were plotted in whisker boxplots. At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical significance: * (p<0.05), ** (p<0.01), *** (p<0.001), **** (p<0.0001).




Figure 5

Metabolic changes following Δ9-THC exposure in ESCs are transcriptionally encoded.

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** PCA of the transcriptomics profiling of either ESCs or EpiLCs mock-exposed or exposed to 100nM Δ9-THC. **(C and D)** Volcano plot in ESCs and EpiLCs, respectively, showing significance [expressed in $\log_{10}(\text{adjusted p-value or false-discovery rate, FDR})$] versus fold-change [expressed in $\log_2(\text{fold-change, FC})$]. Thresholds for significance ($\text{adjusted p-value} \leq 0.05$) and gene expression fold-change [$|\log_2(\text{FC})| > 0.25$ or $|\log_2(\text{FC})| > 0.5$] are shown as dashed lines. Color code is as follows: $\log_2(\text{FC}) > 0.5$ in red, $\log_2(\text{FC}) > 0.25$ in orange, $\log_2(\text{FC}) > 0$ in light orange, $\log_2(\text{FC}) > 0$ in light blue, $\log_2(\text{FC}) > 0.25$ in blue, $\log_2(\text{FC}) > 0.5$ in dark blue and $\text{p-value} < 0.01$ in pink. **(E)** Gene ontology (GO) terms associated with up- and down-regulated DEGs [$|\log_2(\text{FC})| > 0.25$ and $\text{p} < 0.01$] in ESCs and EpiLCs as determined by g:Profiler⁵⁵. **(F)** Joint pathway analysis performed by the multi-omics integration tool of MetaboAnalyst³⁸. The p-values were weighted based on the proportions of genes and metabolites at the individual pathway level.

“Microtubule-based process”. This indicates that $\Delta 9$ -THC exposure significantly upregulated genes in EpiLCs that impact organelles structure, integrity and position, in agreement with several reports in the literature (Lojpur et al., 2019 [↗](#); Miller et al., 2019 [↗](#)).

Multiple reports indicate that $\Delta 9$ -THC exposure alters the epigenome of sperm and brain tissue, both in terms of DNA methylation level and histone post-translational modifications (Murphy et al., 2018 [↗](#); Prini et al., 2017 [↗](#); Schrott and Murphy, 2020 [↗](#); Watson et al., 2015 [↗](#)). Despite the low transcriptional impact of $\Delta 9$ -THC exposure in ESCs and EpiLCs, we evaluated the expression of more than 100 genes encoding epigenetic modifiers in our RNA-seq datasets (Supplementary Table 1). Data shows that the expression levels of multiple epigenetic modifiers were significantly altered, albeit at a low levels, in either ESCs or EpiLCs upon $\Delta 9$ -THC exposure ($|\log_2(\text{fold-change})| > 0.25$, **Supplementary Figure 10** [↗](#)). In particular, the expression levels of the DNA dioxygenase *Tet2* were significantly increased in EpiLCs following $\Delta 9$ -THC exposure, which would contribute to changes in DNA methylation dynamics and differentiation potential (Sohni et al., 2015 [↗](#)). The expression of the two histone deacetylases *Hdac5* and *Hdac11* were significantly decreased in ESCs following $\Delta 9$ -THC exposure, while the expression of the histone phosphorylase *Rps6ka5* increased under the same conditions. In EpiLCs, the expression of the histone methyltransferase *Kmt2c* and of the regulators of histone ubiquitination *Dzip3* and *Mysm1* were all significantly increased upon $\Delta 9$ -THC exposure. Collectively, RNA-seq data suggests the existence of an epigenetic remodelling following $\Delta 9$ -THC exposure, although further analysis of the respective epigenetic marks associated with these modifiers is needed.

Together, our analysis of ESCs and EpiLCs transcriptomes reveals a difference in the response of these stem cell populations to $\Delta 9$ -THC exposure: the transcriptional alterations observed in ESCs supported their increased anabolism and proliferation, whereas changes in EpiLCs gene expression did not correlate with their metabolic changes.

Proliferation of Primordial Germ Cell-Like Cells stemming from prior $\Delta 9$ -THC exposure

PGCs display a distinct transcriptomic and metabolic profile compared to their cellular precursors that are recapitulated in vitro during the differentiation of ESCs into EpiLCs and then of EpiLCs into PGCLCs. Thus, we asked whether the metabolic alterations observed in ESCs and EpiLCs could lead to an altered differentiation program in PGCLCs. To this aim, we continuously exposed ESCs and EpiLCs to a $\Delta 9$ -THC dose range of 1nM–1 μ M (or mock control), before changing to a $\Delta 9$ -THC-free media and inducing PGCLCs differentiation (**Figure 6A** [↗](#)). In particular, we took advantage of ESCs that harbor two fluorescent reporters for germline markers, *Blimp1:mVenus* and *Stella:CFP* (Ohinata et al., 2008 [↗](#)). Thus, the induction efficiency of PGCLCs within 5-days embryoid bodies can be detected by monitoring the fluorescence associated with each cell in flow cytometry, allowing for the determination of a double-negative population (DN), a single-positive population (SP) wherein *Blimp1:mVenus* is expressed and a double-positive population (DP) expressing both *Blimp1:mVenus* and *Stella:CFP*, which represents the true specified PGCLC population (**Supplementary Figure 11** [↗](#)).

We first measured the impact of ESCs + EpiLCs $\Delta 9$ -THC exposure on PGCLC induction efficiency. Flow analyses revealed a dose-dependent increase in the induction efficiency of SP and DP cell populations (**Figure 6B** [↗](#)). Specifically, at 100nM $\Delta 9$ -THC, a significant decrease in DN was observed, with a corresponding significant increase of 1.14-fold in SP and of 1.64-fold in DP cells (**Figure 6C** [↗](#), $p=0.0002$, $p=0.05$, and $p=1.55 \times 10^{-6}$ for 100nM of $\Delta 9$ -THC in DN, SP and DP populations respectively compared to the mock-treated condition, unpaired T-test). The same pattern was observed when male ESCs and EpiLCs exposed to 100nM of $\Delta 9$ -THC were differentiated in PGCLCs (**Supplementary Figure 12** [↗](#)).




Figure 6

PGCLCs deriving from ESCs and EpiLCs exposed to 100nM of $\Delta 9$ -THC proliferate.

(A) Diagram illustrating $\Delta 9$ -THC exposure scheme and experimental strategy. (B) Representative flow contour plots showing distribution of live-gated events, gating strategy for Stella:CFP versus Blimp1:mVenus and percentages of cells in each subpopulations for ESCs and EpiLCs exposed to the different doses of $\Delta 9$ -THC indicated. DN: double negative, SP: single positive, DP: double positive subpopulations. (C) The percentage of events in the gates associated to each subpopulation was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots independently for each subpopulation. (D) Representative histograms showing CellTrace™ Yellow staining profile of cells arising from ESCs and EpiLCs exposed to the different doses of $\Delta 9$ -THC indicated. The Y-axis represents the average percentage of cells in each category of subpopulations undividing (purple), undergoing 1 division (blue), 2 divisions (green) or 3 divisions (orange). One representative experiment out of three is represented. Statistical significance: * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$), **** ($p < 0.0001$).

To determine if the increased proportion of PGCLCs generated from exposed precursors was due to higher proliferative kinetics, we performed a proliferation tracing assay (Tempany et al., 2018 [DOI](#)). The tracing dye was added to the cells on the day of aggregate formation, and fluorescence attenuation due to cell division was measured in each subpopulation on day 5. At 100nM $\Delta 9$ -THC, a smaller proportion of DN cells underwent two or three mitotic divisions compared to the control (**Figure 6D** [DOI](#), 1.14-fold fewer cells and 1.12-fold fewer cells, $p=0.05$ and $p=0.04$ for 2 divisions and 3 divisions, respectively, unpaired T-test). In parallel, for the same dose, a significantly higher proportion of SP and DP cells underwent three mitotic divisions compared to the control (**Figure 6D** [DOI](#), 1.24-fold and 1.11-fold, $p=0.03$ and $p=0.0035$, for 3 divisions in SP and DP cells, compared to the control, unpaired T-test). These results, therefore, indicate that the higher number of PGCLCs observed upon $\Delta 9$ -THC exposure originates from their increased proliferation during their specification and differentiation.

Finally, we sought to determine if the increased PGCLC proliferation was not due to residual intracellular $\Delta 9$ -THC persisting from EpiLCs over the span of PGCLCs differentiation. To do so, intracellular levels of $\Delta 9$ -THC were quantified by mass spectrometry in the cells on the day of aggregate formation and in day 5 embryoid bodies (referred as to “EpiLCs” and “PGCLCs”, respectively, in **Supplementary Figure 13** [DOI](#)). Data shows that no $\Delta 9$ -THC could be detected in day 5 embryoid bodies, indicating that $\Delta 9$ -THC does not persist to levels higher than the limit of detection of mass spectrometry (1ng/mL). This suggests that the proliferative effects are not due to residual $\Delta 9$ -THC persisting in the cells during differentiation towards PGCLCs. Thus, $\Delta 9$ -THC causes an alteration of the developmental kinetics that PGCLCs normally undergo, even in the absence of direct continuous exposure.

PGCLCs derived from $\Delta 9$ -THC-exposed cells present an altered metabolism and transcriptome

Since exposure to $\Delta 9$ -THC prior to their specification increased the number of PGCLCs and ESCs and PGCLCs share similar metabolic programs (Hayashi et al., 2017 [DOI](#); Verdikt and Allard, 2021 [DOI](#)), we next sought to characterize their associated metabolic and transcriptional changes. We, therefore, assessed the impact of exposure of ESCs + EpiLCs to 100nM $\Delta 9$ -THC on PGCLCs metabolism (**Figure 7A** [DOI](#)).




Figure 7

$\Delta 9$ -THC exposure prior to specification increases mitochondrial respiration in PGCLCs.

(A) Diagram illustrating $\Delta 9$ -THC exposure scheme and experimental strategy. (B) The NAD(P)⁺/NADPH ratio of embryoid bodies arising from ESCs and EpiLCs exposed to 100nM of $\Delta 9$ -THC was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplot. (C) Mean fluorescence intensity (MFI) associated with the Mitotracker CMXRos stain in each subpopulation was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots. (D) PCA of the transcriptomics profiling of DP PGCLCs deriving from ESCs and EpiLCs either mock-exposed or exposed to 100nM $\Delta 9$ -THC. (E) Volcano plot in DP PGCLCs showing significance [expressed in $\log_{10}$ (adjusted p-value or false-discovery rate, FDR)] versus fold-change [expressed in $\log_2$ (fold-change, FC)]. Thresholds for significance and different enrichment ratios [$|\log_2(FC)| > 0.25$ or $|\log_2(FC)| > 0.5$] are shown as dashed lines. Color code is as follows: $\log_2(FC) > 0.5$ in red, $\log_2(FC) > 0.25$ in orange, $\log_2(FC) > 0$ in light orange, $\log_2(FC) < 0$ in light blue, $\log_2(FC) < -0.25$ in blue, $\log_2(FC) < -0.5$ in dark blue and p-value < 0.01 in pink. (F and G) Gene ontology (GO) terms associated with up- and downregulated DEGs [$|\log_2(FC)| > 0.25$ and p < 0.01], respectively, as determined by g:Profiler⁵⁵. Statistical significance: * (p < 0.05), ** (p < 0.01).

First, NAD(P)⁺/NAD(P)H assessment revealed a modest but significant 1.17-fold increase in NAD(P)⁺/NAD(P)H ratio in whole day 5 embryoid bodies deriving from exposed ESCs + EpiLCs compared to those deriving from mock-treated cells (Figure 7B, p = 0.01, unpaired T-test). To garner cell type-specific information on whether these metabolic changes were related to mitochondrial activity and the differentiation of PGCLCs, we assessed the mitochondrial membrane potential of each subpopulation in day 5 embryoid bodies. Embryoid bodies were incubated with Mitotracker CMXRos (Pendergrass et al., 2004), and dissociated and analyzed by flow cytometry. The MFI associated with the mitochondrial stain was then measured in each subpopulation (Figure 7C). A significant increase in MFI was observed in DN, SP and DP populations deriving from exposed ESCs + EpiLCs compared to those deriving from mock controls (Figure 7C, 1.17, 1.16, 1.23 -fold, p = 0.006, p = 0.05 and p = 0.01 for DN, SP and DP, respectively, unpaired T-test). These results indicate that the metabolic changes induced by $\Delta 9$ -THC prior to PGCLCs induction and differentiation are not reset during the profound reprogramming that PGCLCs undergo.

Because our results indicated a sustained impact of $\Delta 9$ -THC beyond the period of direct exposure, we further examined PGCLCs by performing a transcriptomic analysis. In particular, day 5 embryoid bodies deriving from ESCs + EpiLCs, either exposed to 100nM of $\Delta 9$ -THC or mock-exposed, were sorted and the total RNA of DP subpopulations, representing true PGCLCs, was analyzed by RNA-seq (Figure 7A). Unsupervised analysis of the global transcriptome in DP PGCLCs by PCA delineated a transcriptional signature of prior $\Delta 9$ -THC exposure (Figure 7D, PC1 accounting for 59% of the variance and PC2 accounting for 24% of variance). Volcano plot of DEGs between DP PGCLCs deriving from mock- or 100nM $\Delta 9$ -THC-exposed ESCs and EpiLCs revealed that most of the significant transcriptional change was towards downregulation rather than upregulation (Figure 7E, 11 genes were significantly upregulated whereas 97 were significantly downregulated, $|\log_2(\text{fold-change})| > 0.25$ and adjusted p-value ≤ 0.05). Despite the low number of upregulated DEGs, the functional annotation of their associated GO terms showed that all terms enriched corresponded to metabolic processes involved in oxidative phosphorylation (Figure 7F, “Aerobic electron transport chain”, “Mitochondrial respiratory chain complex I assembly”, “Electron transport couple proton transport”). Thus, our data indicate that the metabolic changes induced by exposure to $\Delta 9$ -THC prior to PGCLCs specification are retained through transcriptional reprogramming. Importantly, while our results show that pre-specification $\Delta 9$ -THC exposure increases PGCLCs number and mitochondrial activity, the functional annotation of GO terms associated with downregulated DEGs suggests degradation of PGCLCs quality. Indeed, and

reminiscent of GO terms observed in EpiLCs, several GO terms relating to alterations in structural cellular components (“Anatomical structure morphogenesis”, “Cellular anatomical entity”), and in particular the interface with the extracellular environment (“External encapsulating structure organization”, “Membrane”, “Cell periphery”, “Extracellular region”, “Extracellular space”, “Extracellular matrix structural constituent”) were enriched (**Figure 7G** [↗](#)). Furthermore, GO terms associated with cell adhesion and junction (“Cell adhesion”, “Cell migration”, “Collagen metabolic process”, “Cell junction”, “Anchoring junction”, “Collagen trimer”) were also enriched in downregulated genes.

Together, our data show that $\Delta 9$ -THC exposure in ESCs and EpiLCs durably alters their metabolome and that these changes are carried through PGCLCs specification and differentiation, leading to an alteration of PGCLCs transcriptional program (**Figure 8** [↗](#)).

Discussion

With greater social acceptance and legalization, cannabis use has increased worldwide (Mennis et al., 2023 [↗](#); U.N. Office on Drugs and Crime, 2022 [↗](#)). Yet, the impact of such heightened use on reproductive functions, and in particular, on the earliest developmental stages is not well understood. Cannabis use directly alters adult male fertility and causes abnormal embryo implantation (Lo et al., 2022 [↗](#)). Using a well-characterized in vitro model of early embryonic differentiation events culminating in the differentiation of PGCLCs, our study is the first to shed light on the impact of $\Delta 9$ -THC at these stages which unfold during the first trimester in humans (Chabarria et al., 2016 [↗](#); Volkow et al., 2019 [↗](#); Young-Wolff et al., 2018 [↗](#)).

Our data revealed the differential effects of $\Delta 9$ -THC on naïve and primed pluripotent stem cells, respectively represented by mouse ESCs and EpiLCs. In particular, exposure to $\Delta 9$ -THC increased ESCs proliferation which was in a similar range to what has been previously reported for human breast carcinoma cell lines (about 30-50% between 10nM and 1 μ M of $\Delta 9$ -THC) (Takeda et al., 2008 [↗](#)). Differential expression and use of cannabinoid receptors on the surface of exposed cells have been shown to correlate with $\Delta 9$ -THC proliferative phenotypes (Galve-Roperh et al., 2013 [↗](#); Preet et al., 2008 [↗](#); Takeda et al., 2008 [↗](#); Yang et al., 2016 [↗](#)). However, our experiments demonstrated that despite being required for $\Delta 9$ -THC-induced proliferation in ESCs, CB1 expression did not significantly differ at the surface of ESCs and EpiLCs.

Because $\Delta 9$ -THC is a known perturbator of mitochondrial function as previously described in the central nervous system (Bartova and Birmingham, 1976 [↗](#); Bénard et al., 2012 [↗](#)), we studied the metabolic impact of its exposure in ESCs and EpiLCs. Our data indicate that, at 100nM, $\Delta 9$ -THC exposure increased the glycolytic rate in both ESCs and EpiLCs. Bioenergetics analyses and metabolite measurements showed that this increased glucose metabolism did not support increased energy production in the mitochondria, but rather, that it led to the accumulation of metabolic intermediates used in anabolic reactions for the synthesis of amino acids, nucleotides, and lipids. Thus, the metabolic signatures associated with $\Delta 9$ -THC exposure are reminiscent of those inherently occurring during naïve-to-prime transition, during which increased aerobic glycolytic rates feed anabolic reactions ultimately fueling proliferation (Lunt and Vander Heiden, 2011 [↗](#)). We verified this model by testing the requirement of increased glycolysis to support proliferation and indeed observed that ESCs proliferation upon $\Delta 9$ -THC exposure is abrogated in the presence of the glycolytic inhibitor 2-DG.

Transcriptomic analyses revealed that the metabolic reprogramming induced by $\Delta 9$ -THC exposure in ESCs was transcriptionally encoded, with increased expression of genes involved in anabolic pathways. In contrast, functional annotations of DEGs in EpiLCs did not show such transcriptional control of increased anabolism. Comparing the outputs of the metabolomic and transcriptomic analyses (i.e. PCA plots and volcano plots), the impact of $\Delta 9$ -THC at these early stages seems to be




Figure 8

Metabolic impact of $\Delta 9$ -THC exposure in pluripotent stem cells and primordial germ cells-like cells.

Diagram illustrating the impact of $\Delta 9$ -THC exposure on stem cells metabolism.

primarily metabolic, although the moderate effects on the transcriptome appear to support the metabolic outcome as revealed by our integrated analysis (**Figure 5F**). Together, we propose that $\Delta 9$ -THC exposure elicits a reprogramming of ESCs that (1) coaxes them to rely more on aerobic glycolysis, (2) drives anabolic pathways, and therefore (3) leads to their proliferation. In EpiLCs, the impact of $\Delta 9$ -THC exposure is not sufficient to override the cellular and metabolic programs of these already highly proliferative cells that are fully reliant on aerobic glycolysis (**Figure 8**).

Finally, we assessed the impact of $\Delta 9$ -THC exposure in ESCs and EpiLCs on the differentiation of PGCLCs. Our data indicate that at the physiologically relevant dose of 100nM of $\Delta 9$ -THC, a significant increase in PGCLCs was observed. In particular, during PGCLCs differentiation, metabolic reprogramming and increased oxidative phosphorylation play a critical role in the reacquisition of an extended developmental potential (Hayashi et al., 2017; Verdikt and Allard, 2021). Thus, we investigated whether the metabolic alterations observed in ESCs and EpiLCs upon $\Delta 9$ -THC exposure could be carried through PGCLCs differentiation. Metabolic characterization revealed that PGCLCs arising from exposed ESCs and EpiLCs showed increased mitochondrial respiration. Thus, in the absence of direct or indirect continuous exposure, $\Delta 9$ -THC still has lasting consequences on the metabolome of embryonic germ cells. A recent study in drosophila reported that nutrient stress induces oocyte metabolites remodelling that drives the onset of metabolic diseases in the progeny (Hocaoglu et al., 2021). This indicates that non-DNA-associated factors, such as germline metabolites, can act as factors of inheritance. Similarly, we show here that exposure to $\Delta 9$ -THC remodels ESCs and EpiLCs metabolome and that a metabolic memory of this exposure is retained during PGCLCs differentiation (**Figure 8**). In addition to metabolic remodelling, we show that the PGCLCs transcriptome is also altered. In particular, despite proliferation and a higher number of cells, the number of DEGs that were downregulated in PGCLCs deriving from $\Delta 9$ -THC-exposed ESCs and EpiLCs suggests a general degradation of PGCLCs' homeostasis. Functional annotation further indicated that these downregulated genes are related to structural cellular components, to the interaction with the extracellular environment and, specifically, to cell adhesion and junction. During the development of the central nervous system, perinatal $\Delta 9$ -THC exposure has also been associated with alteration in cell adhesion, with an impact on neuronal interactions and morphology (Gómez et al., 2007; Keimpema et al., 2011; Kittler et al., 2000). Cell-cell adhesion is crucial in PGCs' formation both in cell culture systems (Okamura et al., 2003) as well as in vivo where it controls PGCs motility during their migration towards the developing somatic gonad (Barton et al., 2016). Our results thus show that exposure to $\Delta 9$ -THC prior to specification affects embryonic germ cells' transcriptome and metabolome. This in turn could have adverse consequences on cell-cell adhesion with an impact on PGC normal development in vivo.

Despite epidemiological evidence of cannabis exposure in parents being associated with adverse effects in the offspring (Smith et al., 2020; Szutorisz and Hurd, 2018), the molecular mechanisms involved in the inheritance of exposure have not been extensively studied. In vitro models for germ cell development offer a unique opportunity for such studies (Verdikt et al., 2022) and our work in mouse PGCLCs identifies metabolites as relevant carriers of information across developmental stages. While in vitro gametogenesis, up to the reconstitution of fully functional spermatozoa and oocytes, is a reality in mouse (Hikabe et al., 2016; Komeya et al., 2018; Luo and Yu, 2022), transposition to humans has been hindered by ethical and technical considerations (Luo and Yu, 2022). For instance, protocols for human PGCLCs have been developed (Gell et al., 2020) but proper induction to viable, fertile offspring cannot be verified (Luo and Yu, 2022). Nevertheless, our preliminary results in hESCs show that $\Delta 9$ -THC exposure negatively impacts their global energy metabolism (**Supplementary Figure 5**). Future studies will need to confirm whether such metabolic reprogramming is also carried over developmental stages in human models for the germline.

Together, our studies reveal a moderate but significant impact of $\Delta 9$ -THC exposure on early embryonic processes. Our work also highlights the importance of the metabolic remodelling induced by $\Delta 9$ -THC and its potential role as a driver of exposure memory through differentiation stages.

Acknowledgements

We thank Jessica Scholes, Felicia Codrea and Jeffrey Calimlim from the UCLA BSCRC Flow Cytometry Core for their assistance in cell sorting. We thank Linsey Stiles (UCLA Mitochondria and Metabolism Core), Johanna ten Hoeve-Scott and Thomas Graeber (UCLA Metabolomics Center) for their technical support in the metabolomics analyses. We thank Xinmin Li and his team at the UCLA Technology Center for Genomics & Bioinformatics for their technical support in high-throughput sequencing. We thank Gazmend Elezi and Julian Whitelegge (UCLA Pasarow Mass Spectrometry Laboratory) for their help in $\Delta 9$ -THC quantification. We thank Luigi Bellocchio and Giovanni Marsicano for their recommendations in $\Delta 9$ -THC exposure setups. We would like to acknowledge the UCLA Cannabis Research Initiative (UCRI) and Dr Ziva Cooper for their continuous support and encouragements.

PA is supported by NIEHS R01 ES027487, the John Templeton Foundation Grant 60742, and the Iris Cantor-UCLA Women's Health Center and the NCATS UCLA CTSI Grant Number UL1TR001881. RV is a postdoctoral fellow and acknowledges support from the Belgian-American Educational Foundation (BAEF). ATC is supported by NIH/NICHD R01HD079546 (ATC).

Material and methods

Data availability

The RNA sequencing data from this study is made available at the Gene Expression Omnibus (GEO) under the following accession number GSE226955. All other data are available upon request.

Cell culture and PGCLCs model

Mouse ESCs containing the two fluorescent reporters *Blimp1::mVenus* and *Stella::ECFP* (BVSC cells) were described previously (Ohinata et al., 2008 [DOI](#)). The female BVSC clone H18 and male clone R8 were kindly provided by Mitinori Saitou. Cells were seeded on coated plates (Poly-L-ornithine [0.001%; A-004-C; Sigma-Aldrich] and laminin [300ng/mL; L2020; Sigma-Aldrich]) in 2i+LIF culture medium (N2B27 Media, CHIR99021 [30 μ M; NC9785126; Thermo Fisher], PD0325901 [10 μ M; NC9753132; Thermo Fisher], ESGRO® Leukemia Inhibitory Factor (LIF) [1,000 U/mL, ESG1106; Sigma-Aldrich]) for 48h. Differentiation of ESCs to EpiLCs was performed by seeding the cells on Human Plasma Fibronectin (HPF)-coated plates [16.7 μ g/mL; 33016015; Thermo Fisher] in the presence of EpiLC induction medium (N2B27 medium containing activin A [20ng/mL; 50-398-465; Thermo Fisher], basic fibroblast growth factor (bFGF) [12ng/mL; 3139FB025; R&D Systems], and KnockOut Serum Replacement [KSR, 1%; Thermo Fisher]). For PGCLCs induction, 44h EpiLCs were harvested using TrypLE™ Select (1X) (Thermo Fisher) and seeded either in 96-wells plate (Nunclon Sphera, Thermo Fisher) or in EZsphere plates for large-scale induction (Nacalai) in the presence of GK15 medium (Glasgow's Minimal Essential Medium [GMEM, 11710035, Thermo Fisher] supplemented with 15% KSR, 0.1 mM Minimal Essential Medium Nonessential Amino Acids [MEM-NEAA], 1 mM sodium pyruvate, 0.1 mM 2-mercaptoethanol, 100U/mL penicillin, 0.1mg/mL streptomycin, and 2 mM L-glutamine in the presence of bone morphogenetic protein 4 [BMP4; 500ng/mL; 5020-BP-010/CF; R&D Systems], LIF, stem cell factor [SCF; 100ng/mL; 50-399-595; R&D Systems], bone morphogenetic protein 8b [BMP8b; 500ng/mL; 7540-BP-025; R&D Systems], and epidermal growth factor [EGF; 50ng/mL; 2028EG200; R&D Systems]. Cells were cultured for 5d before collection, dissociation of embryoid bodies and downstream experiments.

The hESCs UCLA2 cells (Diaz Perez et al., 2012 [↗](#)) were cultured on plates coated with Recombinant laminin-511 E8 (iMatrix-511 Silk, 892 021, Amsbio) and were maintained under a feeder-free condition in the StemFit® Basic03 medium (SFB-503, Ajinomoto) containing bFGF (100-18B, Peprotech). Prior to passaging, hESC cultures were treated with a 1:1 mixture of TrypLE Select (12563011, Thermo Fisher) and 0.5 mM EDTA/PBS for 15 min at 37°C to dissociate them into single cells. For routine maintenance, hESCs were plated into a 6-well plate (3516, Corning) at a density of 2×10^3 cells/cm² with 10 μM ROCK inhibitor (Y-27632; Tocris, 1254) added in culture medium for 1 day after hESCs passaging. hESCs were plated into 24-wells plate (3526, Corning) at 10,000 cells per well for the viability and viable cell count. For WST-1 assays, hESCs were seeded in 96-wells plate (353072, Falcon) at 1,500 cells per well.

All cells were cultured in a humidified environment at 37°C under 5% CO₂. All cells were tested negative for mycoplasma by a PCR test (ATCC® ISO 9001:2008 and ISO/IEC 17025:2005 quality standards).

Δ9-THC exposures

To assess the impact of Δ9-THC exposure on the developmental trajectory of PGCLCs, three exposure schemes were tested: 1) ESCs exposure only, 2) EpiLCs exposure only, and 3) ESCs+EpiLCs exposure. The stock of Δ9-THC was obtained from the National Institute on Drug Abuse (7370-023 NIDA; Bethesda, MD). The stock was adjusted to a concentration of 200mM diluted in ethanol, aliquoted and stored according to the DEA's recommendations. The dose range of 0-100 μM was determined based on Δ9-THC physiological measurements in the blood, plasma, and follicular fluid (Fuchs Weizman et al., 2021 [↗](#); Hunault et al., 2008 [↗](#); Pacifici et al., 2019 [↗](#)). For each exposure, new aliquots of Δ9-THC were diluted in ESCs or EpiLCs culture media in coated tubes (Sigmacote, Sigma Aldrich). Exposure was performed for 48h. Solubility tests were performed and ethanol was added to reach the same amount for each Δ9-THC concentration (0.05% ethanol). Vehicle control corresponded to 0.05% ethanol added to the respective culture media for ESCs or EpiLCs. All experiments performed are authorized under DEA registration number RA0546828.

Δ9-THC quantification

Samples were treated with 500 μl of 1% Formic Acid (543804, Sigma Aldrich). Labelled Tetrahydrocannabinol (THC-d3), used as an internal standard, was added to every sample to account for compound loss during sample processing. Samples were then mixed vigorously and centrifuged at 16.000g for 5 min at room temperature. The supernatants were loaded into phospholipid removal cartridges (Phenomenex Phree) and the eluents were dried down in a vacuum concentrator. Samples were reconstituted in 30 μl of HPLC-grade water, vortexed rigorously, and centrifuged at 16.000g for 5 min at room temperature. The supernatant was transferred to HPLC vials and 15 μl were injected for analysis onto a hybrid linear ion trap/orbitrap mass spectrometer (Thermo Scientific LTQ Orbitrap XL, UCLA Pasarow Mass Spectrometry Lab). For specificity and accurate quantitative measurement, the mass spectrometer was set to fragment preselected precursor ions for THC and THC-d3 under standard MS/MS fragmentation conditions in positive ion mode. The mass spectrometer was coupled to a Dionex Ultimate 300 HPLC (Thermo Scientific) with a reversed phase Phenomenex analytical column (Kinetex 1.7 μm Polar C18 100 Å 100 x 2.1 mm) equilibrated in eluant A (water/formic acid, 100/0.1, v/v) and eluted (100 μl/min) with a linearly increasing concentration of eluant B (acetonitrile/formic acid, 100/0.1, v/v; min/%B, 0/5, 5/5, 8/95, 13/95, 14/5, 25/5). Data was collected and processed with instrument manufacturer-supplied software Xcalibur 2.07. A set of standard curve samples were prepared in cell culture media for each experiment. Samples and standards were prepared in duplicates. The standard curve was made by plotting the known concentration of THC per standard against the ratio of measured chromatographic peak areas corresponding to the THC over that of the IS THC-d3 (analyte/IS). The trendline equation was then utilized to calculate the absolute concentrations of the THC in cell culture.

PGCLCs induction efficiency

Changes in PGCLCs induction were calculated by flow cytometry. Practically, d5 aggregates were harvested, dissociated using TrypLE™ Select, and resuspended in fluorescence-activated cell sorting (FACS) buffer (1×Dulbecco's phosphate buffered saline [DPBS], 1% BSA, 1 mM EDTA, 25 mM HEPES). Quantification of subfractions of double-positive PGCLCs (Blimp1::mVenus+ and Stella::ECFP+), single-positive (Blimp1::mVenus+) and double-negative cells was performed on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). Cells were initially identified by forward- and side-scatter gating, with back-gating used to verify the accuracy by which target cell populations were identified. Cell populations of interest were identified by 2-D plots displaying the parameter of interest, using embryoid bodies cultured in GK15 medium without added cytokines and BMPs as a negative control. Fluorescent compensation beads were used as positive controls and to calculate the spectral overlap (Thermo Fisher, A10514 and 01-2222-42 adsorbed to a CD45 Pacific Blue antibody [OB180026], serving as compensation control for mVenus and ECFP, respectively). Manually defined gates as well as quadrants were used, as indicated. The FlowJo software was used to calculate the percentage of induction and generate the associated graphs (version 10, FlowJo, LLC).

Cell viability and proliferation studies

The viability and viable cell count of ESCs and EpiLCs were calculated using Trypan blue (0.4%, Thermo Fisher) on a Countess II FL Automated Cell Counter (Thermo Fisher). For BrdU incorporation studies, cells were permeabilized, fixed, and stained using the BrdU Flow Kit (PerCP-Cy™5.5 Mouse anti-BrdU, BD Biosciences) before analysis by flow cytometry on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). Quantification of PGCLCs proliferation was performed using CellTrace™ Yellow (5μM, added at the induction, Thermo Fisher), which binds to intracellular amines after diffusing through cell membranes. The overall fluorescent signal, which gradually decreases as cell division occurs, reflects the number of cell divisions occurring and was measured on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). The FlowJo software was used to calculate the percentage of induction, the number of cell divisions and generate the associated graphs (version 10, FlowJo, LLC).

CB1 antagonist treatment

To block the effects of Δ9-THC on the cannabinoid receptor CB1, ESCs were plated on 48-well plate and were pre-treated with 1μM of SR141716/Rimonabant (SML0800, Sigma Aldrich) for 1h before being exposed to the dose range of Δ9-THC, as above. After 24h incubation, this procedure was repeated and cells were harvested after 48h total incubation. The viability and viable cell count was calculated using Trypan blue (0.4%, Thermo Fisher) on a Countess II FL Automated Cell Counter (Thermo Fisher). The concentration of 1μM of Rimonabant was chosen based on previous experiments (Lojpur et al., 2019 [Lojpur et al., 2019](#)) and did not impact cell viability nor cell number on its own.

Western blotting

Membrane proteins were extracted from cell pellets using the Mem-PER™ Plus Membrane Protein Extraction Kit (89842, Thermo Fisher) according to the manufacturer's protocol. Western blotting was performed with 25μg of protein extracts. The immunodetection was assessed using primary antibodies targeting CB1 (101500, Cayman Chemical) or β-actin (3700, Cell Signaling Technology) as loading control. Horseradish peroxidase (HRP)-conjugated secondary antibodies were used for chemiluminescence detection (Amersham).

WST-1 assay

The colorimetric assay WST-1 was used according to the manufacturer's instructions (Roche). The tetrazolium salt WST-1 is reduced by mitochondrial dehydrogenases to formazan using NAD(P)H as co-substrates. Thus, the quantity of formazan is directly proportional to NAD(P)⁺.

Mitochondrial activity

Staining for mitochondria was performed by incubating cells at 37°C with 250nM MitoTracker CMXRos (M7512, Thermo Fisher) for 30min (Pendergrass et al., 2004 [\[link\]](#)). Cells were washed and analyzed by flow cytometry on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). The FlowJo software (version 10, FlowJo, LLC) was used to calculate the mean fluorescence intensity (MFI) corresponding to the average fluorescence intensity of each event of the selected cell population within the chosen fluorescence channel associated to MitoTracker CMXRos.

Seahorse experiments

The extracellular acidification rate (ECAR) and the oxygen consumption rate (OCR) are indicative of glycolysis and mitochondrial respiration, respectively. A total of 10×10^3 ESCs and 8×10^3 EpiLCs were seeded on Seahorse XF96 plates (101085-004, Agilent Technologies) and exposed to increasing doses of $\Delta 9$ -THC for 48h. On the day of the assay, cells were washed with assay medium (unbuffered DMEM assay medium [5030, Sigma Aldrich] supplemented with 31.6mM NaCl, 3mg/L phenol red, 5mM HEPES, 5mM glucose, 2mM glutamine and 1mM sodium pyruvate). For OCR measurement, compounds were injected sequentially during the assay resulting in final concentrations of 2 μ M oligomycin, 0.75 μ M and 1.35 μ M FCCP, 1 μ M rotenone and 2 μ M antimycin. ECAR was measured in parallel. The measured quantities were normalized to the protein content as measured by a BCA quantitation (23227, Thermo Fisher).

Mass spectrometry-based metabolomics analysis

To extract intracellular metabolites, cells were rinsed with cold 150mM ammonium acetate (pH 7.3) then incubated with 80% ice-cold methanol supplemented with 10 nmol D/L-norvaline for 1h. Following resuspension, cells were pelleted by centrifugation (15,000g, 4°C for 15min). The supernatant was transferred into a glass vial and metabolites were dried down under vacuum, then resuspended in 70% acetonitrile. Mass spectrometry analysis was performed at the UCLA Metabolomics Center with an UltiMate 3000RSLC (Thermo Scientific) coupled to a Q Exactive mass spectrometer (Thermo Scientific) in polarity-switching mode with positive voltage 3.0 kV and negative voltage 2.25 kV. Separation was achieved using a gradient elution with (A) 5mM NH₄AcO (pH 9.9) and (B) acetonitrile. The gradient ran from 15% (A) to 90% (A) over 18 min, followed by an isocratic step for 9 minutes and re-equilibration for 7 minutes. Metabolites were quantified as area under the curve based on retention times and using accurate mass measurements (≤ 3 ppm) with the TraceFinder 3.1 software (Thermo Scientific). For heatmap depiction, the relative amounts of metabolites were normalized to the mean value across all samples for the same condition and to the number of viable cells harvested in parallel on a control plate. Pathway enrichment for up- and downregulated KEGG metabolites ($|\log_2(\text{fold-change})| \geq 0.25$) was determined using the MetaboAnalyst 5.0 platform (www.metaboanalyst.ca [\[link\]](#)) (Pang et al., 2022).

RNA-sequencing

Total RNA was extracted from ESCs and EpiLCs pellets using the AllPrep DNA/RNA Micro Kit (Qiagen), according to the manufacturer's protocol. For PGCLCs, d5 embryoid bodies were harvested and cells were dissociated using TrypLE™ Select followed by resuspension in fluorescence-activated cell sorting (FACS) buffer (1×Dulbecco's phosphate buffered saline [DPBS], 1% BSA, 1 mM EDTA, 25 mM HEPES) and cell suspension were passed through a cell strainer (70 μ m). Cells were sorted on a BD Biosciences FACSaria III (UCLA BSCRC Flow Cytometry Core). Practically, cell populations of interest, being double-positive (Blimp1::mVenus+ and Stella::ECFP+) were sorted and collected in microtubes containing GK15 medium. Total RNA was extracted from double-positive PGCLCs using the AllPrep DNA/RNA Micro Kit (Qiagen). RNA concentration was measured using a NanoDrop™ 2000 UV spectrophotometer (Thermo Fisher). Libraries were prepared with the KAPA mRNA HyperPrep Kit (BioMek) or with the RNA library prep kit (ABClonal) following the manufacturers' protocols. Briefly, poly(A) RNA were selected, fragmented and double-stranded cDNA synthesized using a mixture of random and oligo(dT) priming,

followed by end repair to generate blunt ends, adaptor ligation, strand selection, and polymerase chain reaction (PCR) amplification to produce the final library. Different index adaptors were used for multiplexing samples in one sequencing lane. Sequencing was performed on an Illumina NovaSeq 6000 sequencers for paired end (PE), 2×150 base pair (bp) runs. Data quality check was performed using Illumina Sequencing Analysis Viewer (SAV) software. Demultiplexing was performed with Illumina Bcl2fastq2 program (version 2.19.1.403; Illumina Inc.).

Differential gene expression analysis

The quality of the reads was verified using FastQC ([Andrews, 2010](#)) before reads were aligned to the mm10 reference genome (GRCm39) using STAR ([Dobin et al., 2013](#)) with the following arguments: `-- readFilesCommand zcat --outSAMtype BAM SortedByCoordinate --quantMode GeneCounts -- outFilterMismatchNmax 5 --outFilterMultimapNmax 1`. The quality of the resulting alignments was assessed using QualiMap ([García-Alcalde et al., 2012](#)). The Python package HTseq was used for gene counts ([Anders et al., 2015](#)) using the following arguments: `-- stranded=no --idattr=gene_id --type=exon -- mode=union -r pos --format=bam`. Output files were filtered to remove genes with low count (≤ 10) then were used for differential gene expression analysis using DESeq2 ([Love et al., 2014](#)). The negative binomial regression model of ComBat-seq was used to correct unwanted batch effects ([Zhang et al., 2020](#)). For a gene to be classified as showing differential expression between treated and untreated cells, a threshold of $|\log_2(\text{fold-change})| \geq 0.5$ and Benjamini-Hochberg adjusted p-value ≤ 0.05 had to be met.


Gene Ontology (GO) Analysis

Lists of differentially expressed genes were generated from read counts using DESeq2 Bioconductor package ([Love et al., 2014](#)). Enrichment of GO terms in lists of up- and downregulated genes ($|\log_2(\text{fold-change})| \geq 0.25$) was determined using g:Profiler ([Raudvere et al., 2019](#)). Redundant GO terms were removed using reduce + visualize gene ontology (REVIGO) ([Supek et al., 2011](#)). Terms were included if the fold enrichment (frequency of DEGs in each GO term to the frequency of total genes in GO terms) was higher than 1.5 and if the Benjamini-Hochberg-adjusted p-value was less than 0.05. Plots for GO terms were generated using a custom R script ([Bonnot et al., 2019](#)).

Statistical Methods

Statistical analyses, when not otherwise specified, were performed using GraphPad Prism 9 software. For significance testing, two-tailed T-tests were performed on pairwise comparisons. In all cases, significance was determined by p-values less than or equal to 0.05. Each figure corresponds to at least three independent biological repeats with three technical replicates (N=3, n=3), unless otherwise specified. Number of asterisks on plots indicate level of statistical significance: * (p<0.05), ** (p<0.01), *** (p<0.001), **** (p<0.0001).


Supplementary figures



Supplementary Figure 1

$\Delta 9$ -THC induces ESCs proliferation for as low as 1nM.


(A) Whisker boxplot indicating the median cellular viability of ESCs exposed to the different $\Delta 9$ -THC doses and associated errors. (B) Whisker boxplot indicating the median number of viable cells exposed to the different $\Delta 9$ -THC doses indicated and associated errors. At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical significance: *($p < 0.05$), ***($p < 0.001$), ****($p < 0.0001$).



Supplementary Figure 2

$\Delta 9$ -THC induces alteration in ESCs cell cycle.


(A) Representative flow contour plots showing distribution of BrdU-stained and DAPI-stained cells, exposed to the different doses of $\Delta 9$ -THC indicated. The frequency of events in each gate is indicated. (B) The median percentage of events and associated errors for each cell cycle gate were plotted in histograms. At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical significance: *(p<0.05), **(p<0.01).



Supplementary Figure 3

$\Delta 9$ -THC exposure in male ESCs also provokes cell proliferation.


(A) Whisker boxplot indicating the median cellular viability of male ESCs (the R8 cell line, see Material and Methods section) exposed to the different $\Delta 9$ -THC doses and associated errors. (B) Whisker boxplot indicating the median number of viable cells exposed to the different $\Delta 9$ -THC doses indicated and associated errors. At least three independent biological repeats with three technical replicates ($N=3$, $n=3$). Statistical significance: * ($p<0.05$), ** ($p<0.01$), **** ($p<0.0001$).



Supplementary Figure 4

hESCs cell number decreases upon $\Delta 9$ -THC exposure.


(A) Whisker boxplot indicating the median cellular viability of human embryonic stem cells continuously exposed to 100nM $\Delta 9$ -THC doses over 6 days and associated errors. (B) Whisker boxplot indicating the median number of viable cells exposed to 100nM of $\Delta 9$ -THC doses indicated and associated errors. For (A and B), 6 technical repeats of 2 biological repeats ($n=12$) were plotted. Statistical significance: ** ($p<0.01$).



Supplementary Figure 5

hESCS metabolism is slightly but significantly impacted by Δ^9 -THC exposure.

The NAD(P)⁺/NADPH ratio of hESCs exposed to 100nM of Δ^9 -THC was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots. One representative experiment out of two independent experiments was used to plot results. Statistical significance: *(p<0.05).




Supplementary Figure 6

Extracellular acidification rates and oxygen consumption rates in ESCs and EpiLCs upon Δ9-THC exposure.

(A and B) Traces were plotted for the extracellular acidification rate (ECAR) measurements in ESCs and EpiLCs, respectively, exposed to the different Δ9-THC doses indicated and normalized to the protein content. The oligomycin injection time is indicated by an arrow and allows to differentiate basal glycolytic rate from maximal glycolytic rate (when mitochondria are inhibited). The datapoints used in the main figure correspond to the first timepoint in the maximal glycolytic capacity section.


(C and D) Traces were plotted for the oxygen consumption rate (OCR) measurements in ESCs and EpiLCs, respectively, exposed to the different Δ9-THC doses indicated and normalized to the protein content. The oligomycin, FCCP and AntimycinA/Rotenone injection times are indicated by arrows and allow to differentiate basal respiration from ATP-coupled respiration and maximal respiratory capacity. The datapoints used in the main figure correspond to the second timepoint in the maximal respiratory capacity section. FCCP: Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone. Statistical significance: *($p < 0.05$), **($p < 0.01$), ****($p < 0.0001$).



Supplementary Figure 7

Metabolite profiling in ESCs and EpiLCs upon Δ9-THC exposure.


(A and B) Heatmaps showing the log2 of the amount of each metabolite upregulated in ESCs and EpiLCs upon exposure to 100nM of Δ9-THC. The relative amounts of metabolites were normalized to the mean value across all samples for one same condition and to the number of viable cells harvested in parallel on a control plate. (C) Histograms showing the ratio of reduced to oxidized glutathione (GSH/GSSG) based on the amounts measured in the metabolomics profiling.



Supplementary Figure 8

Extracellular acidification rates and oxygen consumption rates in ESCs upon Δ9-THC and 2-DG exposure.


(A) Traces were plotted for the extracellular acidification rate (ECAR) measurements in ESCs exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, and normalized to the protein content. The oligomycin injection time is indicated by an arrow and allows to differentiate basal glycolytic rate from maximal glycolytic rate (when mitochondria are inhibited). (B) Traces were plotted for the oxygen consumption rate (OCR) measurements in ESCs exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, and normalized to the protein content. The oligomycin, FCCP and AntimycinA/Rotenone injection times are indicated by arrows and allow to differentiate basal respiration from ATP-coupled respiration and maximal respiratory capacity. FCCP: Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone. Statistical significance: *($p < 0.05$), **($p < 0.01$).



Supplementary Figure 9

$\Delta 9$ -THC exposure does not alter markers of pluripotency.


Gene expression profiles of markers for the inner cell mass (ICM) and epiblast. Histograms show the median and associated errors of normalized gene counts in each condition, as indicated.



Supplementary Figure 10

Δ9-THC exposure alter the expression of some epigenetic modifiers.


Histograms show the median and associated errors of normalized gene counts in each condition, as indicated. Only genes with $|\log_2(\text{FC})| > 0.25$ and $p\text{-value} < 0.01$ from Supplementary Table 1 were plotted. Statistical significance: **($p < 0.01$), ***($p < 0.001$), ****($p < 0.0001$).



Supplementary Figure 11

PGCLCs gating and sorting strategy.


(A) Representative flow contour plots showing distribution of events and gating based on embryoid bodies dissociation. (B and C) Representative flow contour plots to isolate singlets based on width to height ratios on the side scatter and front scatter, respectively. (D) Gating strategy for Stella:CFP versus Blimp1:mVenus on the negative control, corresponding to embryoid bodies obtained in an induction medium without cytokines and BMPs (GK15 only). (E) Gating strategy for Stella:CFP versus Blimp1:mVenus on mock-treated cells, corresponding to embryoid bodies obtained in an induction medium complemented with cytokines and BMPs. DN: double negative, SP: single positive, DP: double positive subpopulations.



Supplementary Figure 12

Male PGCLCs deriving from ESCs and EpiLCs exposed to 100nM of Δ9-THC proliferate.

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. (B) Representative flow contour plots showing distribution of live-gated events, gating strategy for Stella:CFP versus Blimp1:mVenus and percentages of cells in each subpopulations for ESCs and EpiLCs exposed to 100nM of Δ9-THC. DN: double negative, SP: single positive, DP: double positive subpopulations. (C) The percentage of events in the gates associated to each subpopulation was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots independently for each subpopulation. At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical significance: *(p<0.05).



Supplementary Figure 13

No residual $\Delta 9$ -THC is detected in day 5 embryoid bodies.

Intracellular levels of $\Delta 9$ -THC were quantified by mass spectrometry in EpiLCs on the day of aggregate formation and in day 5 embryoid bodies (referred as to "EpiLCs" and "PGCLCs"). Histograms show the median and associated errors of two independent quantifications. Statistical significance: ****($p < 0.0001$).

References

- Anders S, Pyl PT, Huber W (2015) **HTSeq-A Python framework to work with high-throughput sequencing data** *Bioinformatics* **31**:166–169 <https://doi.org/10.1093/bioinformatics/btu638>
- Andre CM, Hausman JF, Guerriero G (2016) **Cannabis sativa: The plant of the thousand and one molecules** *Front Plant Sci* **7**:1–17 <https://doi.org/10.3389/fpls.2016.00019>
- Andrews S. (2010) **Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.** <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Barban S, Schulze HO (1961) **The Effects of 2-Deoxyglucose on the Growth and Metabolism of Cultured Human Cells** *Journal of Biological Chemistry* **236**:1887–1890 [https://doi.org/10.1016/s0021-9258\(18\)64100-6](https://doi.org/10.1016/s0021-9258(18)64100-6)
- Barton LJ, LeBlanc MG, Lehmann R (2016) **Finding their way: themes in germ cell migration** *Curr Opin Cell Biol* **42**:128–137 <https://doi.org/10.1016/j.ceb.2016.07.007>
- Bartova A, Birmingham MK (1976) **Effect of Δ^9 tetrahydrocannabinol on mitochondrial NADH oxidase activity** *Journal of Biological Chemistry* **251**:5002–5006 [https://doi.org/10.1016/s0021-9258\(17\)33213-1](https://doi.org/10.1016/s0021-9258(17)33213-1)
- Bénard G *et al.* (2012) **Mitochondrial CB1 receptors regulate neuronal energy metabolism** *Nat Neurosci* **15**:558–564 <https://doi.org/10.1038/nn.3053>
- Bonnot T, Gillard M, Nagel D (2019) **A Simple Protocol for Informative Visualization of Enriched Gene Ontology Terms** *Bio Protoc* **9**:1–9 <https://doi.org/10.21769/bioprotoc.3429>
- Chabarria KC, Racusin DA, Antony KM, Kahr M, Suter MA, Mastrobattista JM, Aagaard KM (2016) **Marijuana use and its effects in pregnancy** *Am J Obstet Gynecol* **215**:506–506 <https://doi.org/10.1016/j.ajog.2016.05.044>
- Chandra S, Radwan MM, Majumdar CG, Church JC, Freeman TP, ElSohly MA (2019) **New trends in cannabis potency in USA and Europe during the last decade (2008–2017)** *Eur Arch Psychiatry Clin Neurosci* **269**:5–15 <https://doi.org/10.1007/s00406-019-00983-5>
- Diaz Perez S V., Kim R, Li Z, Marquez VE, Patel S, Plath K, Clark AT (2012) **Derivation of new human embryonic stem cell lines reveals rapid epigenetic progression in vitro that can be prevented by chemical modification of chromatin** *Hum Mol Genet* **21**:751–764 <https://doi.org/10.1093/hmg/ddr506>
- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) **STAR: Ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21 <https://doi.org/10.1093/bioinformatics/bts635>
- Fuchs Weizman N, Wyse BA, Szaraz P, Defer M, Jahangiri S, Librach CL (2021) **Cannabis alters epigenetic integrity and endocannabinoid signalling in the human follicular niche** *Human Reproduction* **0**:1–10 <https://doi.org/10.1093/humrep/deab104>

- Galve-Roperh I, Chiurchiù V, Díaz-Alonso J, Bari M, Guzmán M, Maccarrone M (2013) **Cannabinoid receptor signaling in progenitor/stem cell proliferation and differentiation** *Prog Lipid Res* **52**:633–650 <https://doi.org/10.1016/j.plipres.2013.05.004>
- García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, Dopazo J, Meyer TF, Conesa A (2012) **Qualimap: Evaluating next-generation sequencing alignment data** *Bioinformatics* **28**:2678–2679 <https://doi.org/10.1093/bioinformatics/bts503>
- Gell JJ, Liu W, Sosa E, Chialastri A, Hancock G, Tao Y, Wamaitha SE, Bower G, Dey SS, Clark AT (2020) **An Extended Culture System that Supports Human Primordial Germ Cell-like Cell Survival and Initiation of DNA Methylation Erasure** *Stem Cell Reports* **14** <https://doi.org/10.1016/j.stemcr.2020.01.009>
- Gómez M, Hernández M, Fernández-Ruiz J (2007) **The activation of cannabinoid receptors during early postnatal development reduces the expression of cell adhesion molecule L1 in the rat brain** *Brain Res* **1145**:48–55 <https://doi.org/10.1016/j.brainres.2007.01.102>
- Hayashi K, Ohta H, Kurimoto K, Aramaki S, Saitou M (2011) **Reconstitution of the Mouse Germ Cell Specification Pathway in Culture by Pluripotent Stem Cells** *Cell* **146**:519–532 <https://doi.org/10.1016/j.cell.2011.06.052>
- Hayashi Y, Otsuka K, Ebina M, Igarashi Kaori, Takehara A, Matsumoto M, Kanai A, Igarashi Kazuhiko, Soga T, Matsui Y (2017) **Distinct requirements for energy metabolism in mouse primordial germ cells and their reprogramming to embryonic germ cells** *Proc Natl Acad Sci U S A* **114**:8289–8294 <https://doi.org/10.1073/pnas.1620915114>
- Hikabe O *et al.* (2016) **Reconstitution in vitro of the entire cycle of the mouse female germ line** *Nature* **539**:299–303 <https://doi.org/10.1038/nature20104>
- Hocaoglu H, Wang L, Yang M, Yue S, Sieber M (2021) **Heritable shifts in redox metabolites during mitochondrial quiescence reprogramme progeny metabolism** *Nat Metab* **3**:1259–1274 <https://doi.org/10.1038/s42255-021-00450-3>
- Hunault CC, Mensinga TT, Leenders MEC, Meulenbelt J (2008) **Delta-9-tetrahydrocannabinol (THC) serum concentrations and pharmacological effects in males after smoking a combination of tobacco and cannabis containing up to 69 mg THC** *Psychopharmacology (Berl)* **201**:171–181 <https://doi.org/10.1007/s00213-008-1260-2>
- Keimpema E, MacKie K, Harkany T (2011) **Molecular model of cannabis sensitivity in developing neuronal circuits** *Trends Pharmacol Sci* **32**:551–561 <https://doi.org/10.1016/j.tips.2011.05.004>
- Kinoshita M, Barber M, Mansfield W, Cui Y, Spindlow D, Stirparo GG, Dietmann S, Nichols J, Smith A (2021) **Capture of Mouse and Human Stem Cells with Features of Formative Pluripotency II II Capture of Mouse and Human Stem Cells with Features of Formative Pluripotency** *Stem Cell* **28**:453–471 <https://doi.org/10.1016/j.stem.2020.11.005>
- Kittler JT, Grigorenko E V., Clayton C, Zhuang SY, Bunday SC, Trower MM, Wallace D, Hampson R, Deadwyler S (2000) **Large-scale analysis of gene expression changes during acute and chronic exposure to Δ9-THC in rats** *Physiol Genomics* **2000**:175–185 <https://doi.org/10.1152/physiolgenomics.2000.3.3.175>
- Komeya M, Sato T, Ogawa T (2018) **In vitro spermatogenesis: A century-long research journey still half way** *Reprod Med Biol* **17**:407–420 <https://doi.org/10.1002/rmb2.12225>

- Kurimoto K, Saitou M (2018) **Epigenome regulation during germ cell specification and development from pluripotent stem cells** *Curr Opin Genet Dev* **52**:57–64 <https://doi.org/10.1016/j.gde.2018.06.004>
- Lo JO, Hedges JC, Girardi G (2022) **Impact of cannabinoids on pregnancy, reproductive health, and offspring outcomes** *Am J Obstet Gynecol* **227**:571–581 <https://doi.org/10.1016/j.ajog.2022.05.056>
- Locasale JW, Cantley LC (2011) **Metabolic flux and the regulation of mammalian cell growth** *Cell Metab* **14**:443–451 <https://doi.org/10.1016/j.cmet.2011.07.014>
- Lojpur T, Easton Z, Raez-Villanueva S, Laviolette S, Holloway AC, Hardy DB (2019) **Δ9-Tetrahydrocannabinol leads to endoplasmic reticulum stress and mitochondrial dysfunction in human BeWo trophoblasts** *Reproductive Toxicology* **87**:21–31 <https://doi.org/10.1016/j.reprotox.2019.04.008>
- Love MI, Huber W, Anders S (2014) **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2** *Genome Biol* **15**:1–21 <https://doi.org/10.1186/s13059-014-0550-8>
- Lunt SY, Vander Heiden MG (2011) **Aerobic glycolysis: Meeting the metabolic requirements of cell proliferation** *Annu Rev Cell Dev Biol* **27**:441–464 <https://doi.org/10.1146/annurev-cellbio-092910-154237>
- Luo Y, Yu Y (2022) **Research Advances in Gametogenesis and Embryogenesis Using Pluripotent Stem Cells** *Front Cell Dev Biol* **9**:1–14 <https://doi.org/10.3389/fcell.2021.801468>
- Mennis J, Stahler GJ, Mason MJ (2023) **Cannabis Legalization and the Decline of Cannabis Use Disorder (CUD) Treatment Utilization in the US** *Curr Addict Rep* <https://doi.org/10.1007/s40429-022-00461-4>
- Miller ML *et al.* (2019) **Adolescent exposure to Δ 9-tetrahydrocannabinol alters the transcriptional trajectory and dendritic architecture of prefrontal pyramidal neurons** *Mol Psychiatry* **24**:588–600 <https://doi.org/10.1038/s41380-018-0243-x>
- Murphy SK *et al.* (2018) **Cannabinoid exposure and altered DNA methylation in rat and human sperm** *Epigenetics* **13**:1208–1221 <https://doi.org/10.1080/15592294.2018.1554521>
- Nichols J, Smith A (2009) **Naive and Primed Pluripotent States** *Cell Stem Cell* **4**:487–492 <https://doi.org/10.1016/j.stem.2009.05.015>
- Ohinata Y, Sano M, Shigeta M, Yamanaka K, Saitou M (2008) **A comprehensive, non-invasive visualization of primordial germ cell development in mice by the Prdm1-mVenus and Dppa3-ECFP double transgenic reporter** *Reproduction* **136**:503–514 <https://doi.org/10.1530/REP-08-0053>
- Okamura D, Kimura T, Nakano T, Matsui Y (2003) **Cadherin-mediated cell interaction regulates germ cell determination in mice** *Development* **130**:6423–6430 <https://doi.org/10.1242/dev.00870>
- Osborne AJ, Pearson JF, Noble AJ, Gemmell NJ, Horwood LJ, Boden JM, Benton MC, Macartney-Coxson DP, Kennedy MA (2020) **Genome-wide DNA methylation analysis of heavy cannabis exposure in a New Zealand longitudinal cohort** *Transl Psychiatry* **10**:1–10 <https://doi.org/10.1038/s41398-020-0800-3>

- Pacher P, Kogan NM, Mechoulam R (2020) **Beyond THC and Endocannabinoids** *Annu Rev Pharmacol Toxicol* **60** <https://doi.org/10.1146/annurev-pharmtox-010818-021441>
- Pacifici R, Pichini S, Pellegrini M, Rotolo MC, Giorgetti R, Tagliabracci A, Busardò FP, Huestis MA (2019) **THC and CBD concentrations in blood, oral fluid and urine following a single and repeated administration of “light cannabis** *Clin Chem Lab Med* :1–8
- Pang Z, Zhou G, Ewald J, Chang L, Hacariz O, Basu N, Xia J (2022) **Using MetaboAnalyst 5.0 for LC–HRMS spectra processing, multi-omics integration and covariate adjustment of global metabolomics data** *Nat Protoc* **17**:1735–1761 <https://doi.org/10.1038/s41596-022-00710-w>
- Paria BC, Das SK, Dey SK (1995) **The preimplantation mouse embryo is a target for cannabinoid ligand-receptor signaling** *Proc Natl Acad Sci U S A* **92**:9460–9464 <https://doi.org/10.1073/pnas.92.21.9460>
- Pendergrass W, Wolf N, Pool M (2004) **Efficacy of MitoTracker Green™ and CMXRosamine to measure changes in mitochondrial membrane potentials in living cells and tissues** *Cytometry Part A* **61**:162–169 <https://doi.org/10.1002/cyto.a.20033>
- Preet A, Ganju RK, Groopman JE (2008) **Δ9-Tetrahydrocannabinol inhibits epithelial growth factor-induced lung cancer cell migration in vitro as well as its growth and metastasis in vivo** *Oncogene* **27**:339–346 <https://doi.org/10.1038/sj.onc.1210641>
- Prini P, Rusconi F, Zamberletti E, Gabaglio M, Penna F, Fasano M, Battaglioli E, Parolaro D, Rubino T (2017) **Adolescent THC exposure in female rats leads to cognitive deficits through a mechanism involving chromatin modifications in the prefrontal cortex** *J Psychiatry Neurosci* **43** <https://doi.org/10.1503/jpn.170082>
- Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) **G:Profiler: A web server for functional enrichment analysis and conversions of gene lists (2019 update)** *Nucleic Acids Res* **47**:W191–W198 <https://doi.org/10.1093/nar/gkz369>
- Rinaldi-Carmona M *et al.* (1994) **SR141716A, a potent and selective antagonist of the brain cannabinoid receptor** *FEBS Lett* **350**:240–244 [https://doi.org/10.1016/0014-5793\(94\)00773-X](https://doi.org/10.1016/0014-5793(94)00773-X)
- Schrott R, Murphy SK (2020) **Cannabis use and the sperm epigenome: a budding concern?** *Environ Epigenet* **6**:1–10 <https://doi.org/10.1093/eep/dvaa002>
- Smith A, Kaufman F, Sandy MS, Cardenas A (2020) **Cannabis Exposure During Critical Windows of Development: Epigenetic and Molecular Pathways Implicated in Neuropsychiatric Disease** *Curr Environ Health Rep* **7**:325–342 <https://doi.org/10.1007/s40572-020-00275-4>
- Sohni A, Bartocetti M, Khoueiry R, Spans L, Vande Velde J, De Troyer L, Pulakanti K, Claessens F, Rao S, Koh KP (2015) **Dynamic Switching of Active Promoter and Enhancer Domains Regulates Tet1 and Tet2 Expression during Cell State Transitions between Pluripotency and Differentiation** *Mol Cell Biol* **35**:1026–1042 <https://doi.org/10.1128/mcb.01172-14>
- Substance Abuse and Mental Health Services Administration (2020) **Key substance use and mental health indicators in the United States: Results from the 2019 National Survey on Drug Use and Health**
- Supek F, Bošnjak M, Škunca N, Šmuc T (2011) **Revigo summarizes and visualizes long lists of gene ontology terms** *PLoS One* **6** <https://doi.org/10.1371/journal.pone.0021800>

- Szutorisz H, Hurd YL (2018) **High times for cannabis: Epigenetic imprint and its legacy on brain and behavior** *Neurosci Biobehav Rev* **85**:93–101 <https://doi.org/10.1016/j.neubiorev.2017.05.011>
- Takeda S, Yamaori S, Motoya E, Matsunaga T, Kimura T, Yamamoto I, Watanabe K (2008) **Δ9-Tetrahydrocannabinol enhances MCF-7 cell proliferation via cannabinoid receptor-independent signaling** *Toxicology* **245**:141–146 <https://doi.org/10.1016/j.tox.2007.12.019>
- Tempany JC, Zhou JHS, Hodgkin PD, Bryant VL (2018) **Superior properties of CellTrace YellowTM as a division tracking dye for human and murine lymphocytes** *Immunol Cell Biol* **96**:149–159 <https://doi.org/10.1111/imcb.1020>
- U.N. Office on Drugs and Crime (2022) **2022 World Drug Report**
- Verdikt R, Allard P (2021) **Metabolo-epigenetics: the interplay of metabolism and epigenetics during early germ cells development** *Biol Reprod* **105**:616–624 <https://doi.org/10.1093/biolre/ioab118>
- Verdikt R, Armstrong AA, Allard P (2022) **Transgenerational inheritance and its modulation by environmental cues** *Current Topics in Developmental Biology* Elsevier Inc :1–46 <https://doi.org/10.1016/bs.ctdb.2022.10.002>
- Volkow ND, Han B, Compton WM, McCance-Katz EF (2019) **Self-reported Medical and Nonmedical Cannabis Use Among Pregnant Women in the United States** *JAMA - Journal of the American Medical Association* **322**:167–169 <https://doi.org/10.1001/jama.2018.20391>
- Watson CT, Szutorisz H, Garg P, Martin Q, Landry JA, Sharp AJ, Hurd YL (2015) **Genome-Wide DNA Methylation Profiling Reveals Epigenetic Changes in the Rat Nucleus Accumbens Associated with Cross-Generational Effects of Adolescent THC Exposure** *Neuropsychopharmacology* **40**:2993–3005 <https://doi.org/10.1038/npp.2015.155>
- Weinberger L, Ayyash M, Novershtern N, Hanna JH (2016) **Dynamic stem cell states: Naive to primed pluripotency in rodents and humans** *Nat Rev Mol Cell Biol* **17**:155–169 <https://doi.org/10.1038/nrm.2015.28>
- Yang X, Bam M, Nagarkatti PS, Nagarkatti M (2016) **RNA-seq analysis of Δ9-tetrahydrocannabinol-treated T cells reveals altered gene expression profiles that regulate immune response and cell proliferation** *Journal of Biological Chemistry* **291**:15460–15472 <https://doi.org/10.1074/jbc.M116.719179>
- Young-Wolff KC, Tucker L, Alexeeff S, Anne M, Conway A, Weisner C, Goler N (2018) **Among Pregnant Females in California From 2009 – 2016** *JAMA - Journal of the American Medical Association* **318**:2490–2491 <https://doi.org/10.1001/jama.2017.17225.Trends>
- Zhang J, Zhao J, Dahan P, Lu V, Zhang C, Li H, Teitell MA (2018) **Metabolism in Pluripotent Stem Cells and Early Mammalian Development** *Cell Metab* **27**:332–338 <https://doi.org/10.1016/j.cmet.2018.01.008>
- Zhang Y, Parmigiani G, Johnson WE (2020) **ComBat-seq: Batch effect adjustment for RNA-seq count data** *NAR Genom Bioinform* **2**:1–10 <https://doi.org/10.1093/nargab/lqaa078>

Article and author information

Roxane Verdikt

Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA 90095, USA

ORCID iD: [0000-0002-1091-6720](https://orcid.org/0000-0002-1091-6720)

Abigail A. Armstrong

Department of Obstetrics/Gynecology and Reproductive Endocrinology and Infertility, University of California, Los Angeles, CA, USA

Jenny Cheng

Molecular, Cellular, and Integrative Physiology Graduate Program, University of California, Los Angeles, Los Angeles, CA 90095, USA

Young Sun Hwang

Department of Molecular Cell and Developmental Biology, University of California, Los Angeles, Los Angeles, CA, 90095, USA

Amander T. Clark

Department of Molecular Cell and Developmental Biology, University of California, Los Angeles, Los Angeles, CA, 90095, USA, Center for Reproductive Science, Health and Education, University of California, Los Angeles, Los Angeles, CA 90095, USA, Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California, Los Angeles, Los Angeles, CA, 90095, USA

Xia Yang

Integrative Biology and Physiology Department, University of California, Los Angeles, CA, 90095, USA, Department of Molecular and Medical Pharmacology, University of California, Los Angeles, Los Angeles, CA 90095, USA

ORCID iD: [0000-0002-3971-038X](https://orcid.org/0000-0002-3971-038X)

Patrick Allard

Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA 90095, USA, Molecular Biology Institute, University of California, Los Angeles, Los Angeles, CA 90095, USA

For correspondence: pallard@ucla.edu

ORCID iD: [0000-0001-7765-1547](https://orcid.org/0000-0001-7765-1547)

Copyright

© 2023, Verdikt et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Carmen Williams

National Institute of Environmental Health Sciences, Research Triangle Park, United States of America

Senior Editor

Wei Yan

University of California, Los Angeles, Torrance, United States of America

Reviewer #1 (Public Review):

The study is valuable because it shows that physiologically relevant $\Delta 9$ -THC concentrations have metabolic effects on early mouse embryonic cell types, which could cause developmental effects. Overall, the authors have convincing evidence showing that $\Delta 9$ -THC has metabolic effects on mouse embryonic stem cells (mESCs), and that these effects persist in mESC-derived primordial germ cell-like cells even after $\Delta 9$ -THC treatment has stopped. In this revised version, the authors included additional data to characterize the dose-dependence of the effects of $\Delta 9$ -THC. Furthermore, they supported their finding of metabolic memory in PGCLCs by ruling out the potential alternative explanation that $\Delta 9$ -THC persists in the cultured cells over the course of their experiment. This study has two significant implications: first, that $\Delta 9$ -THC may alter the metabolism of early mouse embryos, and second, that mouse primordial germ cell-like cells can have a memory of previous metabolic perturbations.

The authors investigated the metabolic effects of $\Delta 9$ -THC, the main psychoactive component of cannabis, on early mouse embryonic cell types. They found that $\Delta 9$ -THC increases proliferation in male and female mouse embryonic stem cells (mESCs) and upregulates glycolysis. Additionally, primordial germ cell-like cells (PGCLCs) differentiated from $\Delta 9$ -THC-exposed cells also show alterations to their metabolism. The study is valuable because it shows that physiologically relevant $\Delta 9$ -THC concentrations have metabolic effects on cell types from the early embryo, which may cause developmental effects. Intriguingly, these effects persist in PGCLCs even after withdrawal of $\Delta 9$ -THC.

The study shows that $\Delta 9$ -THC increases the proliferation rate of mESCs but not mEpiLCs, without substantially affecting cell viability, except at the highest dose of 100 μ M which shows toxicity (Figure 1). The dose required to cause increased proliferation was approximately 1 nM (Supplementary Figure 1), which is remarkably low. Treatment of mESCs with rimonabant (a CB1 receptor antagonist) blocks the effect of 100 nM $\Delta 9$ -THC on cell proliferation, showing that the proliferative effect is mediated by CB1 receptor signaling. Similarly, treatment with 2-deoxyglucose, a glycolysis inhibitor, also blocks this proliferative effect (Figure 4G-H). Therefore, the effect of $\Delta 9$ -THC depends on both CB1 signaling and glycolysis. This set of experiments strengthens the conclusions of the study by helping to elucidate the mechanism of the effects of $\Delta 9$ -THC.

The study also profiles the transcriptome and metabolome of cells exposed to 100 nM $\Delta 9$ -THC (Figure 4). Although the transcriptomic changes are modest overall, there is upregulation of anabolic genes, consistent with the increased proliferation rate in mESCs. Metabolomic profiling revealed a broad upregulation of metabolites in mESCs treated with 100 nM $\Delta 9$ -THC. Some metabolic effects were also observed at a lower dose of 10 nM (Figure 3B).

Additionally, the study shows that $\Delta 9$ -THC can influence germ cell specification. mESCs were differentiated to mEpiLCs in the presence or absence of $\Delta 9$ -THC, and the mEpiLCs were subsequently differentiated to mPGCLCs. mPGCLC induction efficiency was tracked using a BV:SC dual fluorescent reporter. $\Delta 9$ -THC treated cells had a moderate increase in the double

positive mPGCLC population, and decrease in the double negative population. A cell tracking dye showed that mPGCLCs differentiated from $\Delta 9$ -THC treated cells had undergone more divisions on average. As with the mESCs, these mPGCLCs also had altered gene expression and metabolism, consistent with an increased proliferation rate. Importantly, in the revised version, the authors supported their finding of metabolic memory in mPGCLCs by ruling out the potential alternative explanation that $\Delta 9$ -THC persists in the cultured cells over the course of their experiment (Supplementary Figure 13).

Finally, the authors also observed that $\Delta 9$ -THC decreases the proliferation of human ESCs (Supplementary Figure 4). These cells were in the primed pluripotent state, making them more similar to mEpiLCs than mESCs. Although this result may form the basis of follow-up experiments in a separate paper, it would be premature to conclude that the same effects will be observed in human cells as were observed in mouse cells.

Overall, this study provides good evidence for $\Delta 9$ -THC having metabolic effects on mouse ESCs, and additionally shows that these effects can persist during germ cell specification. Potential effects of $\Delta 9$ -THC exposure during early embryonic development are important for society to understand, and the results of this study are significant for public health.

<https://doi.org/10.7554/eLife.88795.2.sa1>

Reviewer #2 (Public Review):

Verdikt et al. focused on the influence of $\Delta 9$ -THC, the most abundant phytocannabinoid, on early embryonic processes. The authors chose an in vitro differentiation system as model, and compared the proliferation rate, metabolic status and transcriptional level in ESCs, exposure to $\Delta 9$ -THC. They also evaluated the change of metabolism and transcriptome in PGCLCs derived from $\Delta 9$ -THC-exposed cells. All the methods in this paper do not involve the differentiation of ESCs to lineage specific cells. So the results cannot demonstrate the impact of $\Delta 9$ -THC on postimplantation developmental stages. In brief, the authors want to explore the impact of $\Delta 9$ -THC on preimplantation developmental stages, but they only detected the change in ESCs and PGCLCs derived from ESCs, exposure to $\Delta 9$ -THC, which showed the molecular characterization of the impact of $\Delta 9$ -THC exposure on ESCs and PGCLCs.

<https://doi.org/10.7554/eLife.88795.2.sa0>

Author Response

The following is the authors' response to the original reviews.

We want to thank the Editor and Reviewers for their thorough assessment of the manuscript as well as their constructive critiques. We have collated below the public review and recommendations from each Reviewer as well as our responses to them.

eLife assessment

This study by Verdikt et al. provided solid evidence demonstrating the potential impacts of $\Delta 9$ -tetrahydrocannabinol ($\Delta 9$ -THC) on early embryonic development using mouse embryonic stem cells (mESCs) and in vitro differentiation. Their results revealed that $\Delta 9$ -THC enhanced mESCs proliferation and metabolic adaptation, possibly persisting through differentiation to Primordial Germ Cell-Like Cells (PGCLCs), though the evidence supporting this persistence was incomplete. Although the study is important, it was limited by being conducted solely in vitro and lacking parallel human model experiments.

Reviewer #1 (Public Review):

The authors investigated the metabolic effects of $\Delta 9$ -THC, the main psychoactive component of cannabis, on early mouse embryonic cell types. They found that $\Delta 9$ -THC increases proliferation in female mouse embryonic stem cells (mESCs) and upregulates glycolysis. Additionally, primordial germ cell-like cells (PGCLCs) differentiated from $\Delta 9$ -THC-exposed cells also show alterations to their metabolism. The study is valuable because it shows that physiologically relevant $\Delta 9$ -THC concentrations have metabolic effects on cell types from the early embryo, which may cause developmental effects. However, the claim of "metabolic memory" is not justified by the current data, since the effects on PGCLCs could potentially be due to $\Delta 9$ -THC persisting in the cultured cells over the course of the experiment, even after the growth medium without $\Delta 9$ -THC was added.

The study shows that $\Delta 9$ -THC increases the proliferation rate of mESCs but not mEpiLCs, without substantially affecting cell viability, except at the highest dose of 100 μ M which shows toxicity (Figure 1). Treatment of mESCs with rimonabant (a CB1 receptor antagonist) blocks the effect of 100 nM $\Delta 9$ -THC on cell proliferation, showing that the proliferative effect is mediated by CB1 receptor signaling. Similarly, treatment with 2-deoxyglucose, a glycolysis inhibitor, also blocks this proliferative effect (Figure 4G-H). Therefore, the effect of $\Delta 9$ -THC depends on both CB1 signaling and glycolysis. This set of experiments strengthens the conclusions of the study by helping to elucidate the mechanism of the effects of $\Delta 9$ -THC.

Although several experiments independently showed a metabolic effect of $\Delta 9$ -THC treatment, this effect was not dose-dependent over the range of concentrations tested (10 nM and above). Given that metabolic effects were observed even at 10 nM $\Delta 9$ -THC (see for example Figure 1C and 3B), the authors should test lower concentrations to determine the dose-dependence and EC50 of this effect. The authors should also compare their observed EC50 with the binding affinity of $\Delta 9$ -THC to cellular receptors such as CB1, CB2, and GPR55 (reported by other studies).

The study also profiles the transcriptome and metabolome of cells exposed to 100 nM $\Delta 9$ -THC. Although the transcriptomic changes are modest overall, there is upregulation of anabolic genes, consistent with the increased proliferation rate in mESCs. Metabolomic profiling revealed a broad upregulation of metabolites in mESCs treated with 100 nM $\Delta 9$ -THC.

Additionally, the study shows that $\Delta 9$ -THC can influence germ cell specification. mESCs were differentiated to mEpiLCs in the presence or absence of $\Delta 9$ -THC, and the mEpiLCs were subsequently differentiated to mPGCLCs. mPGCLC induction efficiency was tracked using a BV:SC dual fluorescent reporter. $\Delta 9$ -THC treated cells had a moderate increase in the double positive mPGCLC population and a decrease in the double negative population. A cell tracking dye showed that mPGCLCs differentiated from $\Delta 9$ -THC treated cells had undergone more divisions on average. As with the mESCs, these mPGCLCs also had altered gene expression and metabolism, consistent with an increased proliferation rate.

My main criticism is that the current experimental setup does not distinguish between "metabolic memory" vs. carryover of THC (or its metabolites) causing metabolic effects. The authors assume that their PGCLC induction was performed "in the absence of continuous exposure" but this assumption may not be justified. $\Delta 9$ -THC might persist in the cells since it is highly hydrophobic. In order to rule out the persistence of $\Delta 9$ -THC as an explanation of the effects seen in PGCLCs, the authors should measure concentrations of $\Delta 9$ -THC and THC metabolites over time during the course of their PGCLC induction experiment. This could be done by mass spectrometry. This is particularly important

because 10 nM of Δ^9 -THC was shown to have metabolic effects (Figure 1C, 3B, etc.). Since the EpiLCs were treated with 100 nM, if even 10% of the Δ^9 -THC remained, this could account for the metabolic effects. If the authors want to prove "metabolic memory", they need to show that the concentration of Δ^9 -THC is below the minimum dose required for metabolic effects.

Overall, this study is promising but needs some additional work in order to justify its conclusions. The developmental effects of Δ^9 -THC exposure are important for society to understand, and the results of this study are significant for public health.

**Reviewer #1 (Recommendations For The Authors):*

This has the potential to be a good study, but it's currently missing two key experiments:

What is the minimum dose of Δ^9 -THC required to see metabolic effects?

We would like to thank Reviewer 1 for their insightful comments. We have included exposures to lower doses of Δ^9 -THC in Supplementary Figure 1. Our data shows that Δ^9 -THC induces mESCs proliferation from 1nM onwards. However, when ESCs and EpiLCs were exposed to 1nM of Δ^9 -THC, no significant change in mPGCLCs induction was observed (updated Figure 6B). Of note, in their public review, Reviewer 1 mentioned that "The authors should also compare their observed EC50 with the binding affinity of Δ^9 -THC to cellular receptors such as CB1, CB2, and GPR55 (reported by other studies)." According to the literature, stimulation of non-cannabinoid receptors and ion channels (including GPR18, GPR55, TRPVs, etc.) occurs at 40nM-10 μ M of Δ^9 -THC (Banister et al., 2019). We therefore expect that at the lower nanomolar range tested, CB1 is the main receptor stimulated by Δ^9 -THC, as we showed for the 100nM dose in our rimonabant experiments (Fig. 2).

Is the residual THC concentration during the PGCLC induction below this minimum dose? Even if the effects are due to residual Δ^9 -THC, this would not undermine the overall study. There would simply be a different interpretation of the results.

This experiment was particularly important to distinguish between a "true" Δ^9 -THC metabolic memory or residual Δ^9 -THC leftover during PGCLCs differentiation. Our mass spectrometry quantification revealed that no significant Δ^9 -THC could be detected in day 5 embryoid bodies compared to treated EpiLCs prior to differentiation (Supplementary Figure 13). These results support the existence of Δ^9 -THC metabolic memory across differentiation.

You also do not mention whether you tested your cells for mycoplasma. This is important since mycoplasma contamination is a common problem that can cause artifactual results. Please test your cells and report the results.

All cells were tested negative for mycoplasma by a PCR test (ATCC® ISO 9001:2008 and ISO/IEC 17025:2005 quality standards). This information has been added in the Material and Methods section.

Minor points:

- 1. I don't think it's correct to say that cannabis is the most commonly used psychoactive drug. Alcohol and nicotine are more commonly used. See: <https://nida.nih.gov/research-topics/alcohol> and <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/psychoactive-substance> I looked at the UN drugs report [ref 1] and alcohol or nicotine were not included on that list of drugs, so the UN may use a different definition. This doesn't affect the importance or conclusions of this study, but the wording should be changed.*


We agree and are now following the WHO description of cannabis (<https://www.who.int/teams/mental-health-and-substance-use/alcohol-drugs-and-addictive-behaviours/drugs-psychoactive/cannabis>) by referring to it as the “most widely used illicit drug in the world”. (Line 44).

1. It would be informative to use your RNA-seq data to examine the expression of receptors for Δ^9 -THC such as CB1, CB2, and GPR55. CB1 might be the main one, but I am curious to see if others are present.

We have explored the protein expression of several cannabinoid receptors, including CB2, GPR18, GPR55 and TRPV1 (Bannister et al., 2019). These proteins, except TRPV1, were lowly expressed in mouse embryonic stem cells compared to the positive control (mouse brain extract, see Author response image 1). Furthermore, our experiment with Rimonabant showed that the proliferative effects of Δ^9 -THC are mediated through CB1.

Author response image 1.

Cannabinoid receptors and non-cannabinoid receptors protein expression in mouse embryonic stem cells.



1. Make sure to report exact p-values. You usually do this, but there are a few places where it says $p < 0.0001$. Also, report whether T-tests assumed equal variance (Student's) or unequal variance (Welch's). [In general, it's better to use unequal variance, unless there is good reason to assume equal variance.]


Prism, which was used for statistical analyses, only reports p-values to four decimal places. For all p-values that were $p < 0.0001$, the exact decimals were calculated in Excel using the “=T.DIST.2T(t, df)” function, where the Student's distribution and the number of degrees of freedom computed by Prism were inputted. Homoscedasticity was confirmed for all statistical analyses in Prism.

1. Figure 2A: An uncropped gel image should be provided as supplementary data. Additionally, show positive and negative controls (from cells known to either express CB1 or not express CB1)

The uncropped gel image is presented in Author response image 2. The antibody was validated on mouse brain extracts as a positive control as shown in Figure 1.

Author response image 2.

Uncropped gel corresponding to Fig. 2A where an anti-CB1 antibody was used.



1. Figure 6B: Please show a representative gating scheme for flow cytometry (including controls) as supplementary data. Also, was a live/dead stain used? What controls were used for compensation? These details should be reported.

The gating strategy is presented in Supplementary Figure 11. The Material and Methods section has also been expanded.

1. As far as I can tell, you only used female mESCs. It would be good to test the effects on male mESCs as well since these have some differences due to differences in X-linked gene expression (female mESCs have two active X chromosomes). I understand that you might not have a male BV:SC reporter line, so it would be acceptable to omit the mPGCLC experiments on male cells.

We have tested the 10nM-100μM dose range in the male R8 mESCs (Supplementary Figure 3). Similar results as with the female H18 cells were observed. Accordingly, PGCLCs induction was increased when R8 ESCs + EpiLCs were exposed to 100nM of Δ9-THC (Supplementary Figure 12). This is in line with Δ9-THC impact on fundamentally conserved metabolic pathways across species and sex, although it should be noted that one representative model of each sex is not sufficient to exclude sex-specific effects.

Reviewer #2 (Public Review):

In the study conducted by Verdikt et al, the authors employed mouse Embryonic Stem Cells (ESCs) and in vitro differentiation techniques to demonstrate that exposure to cannabis, specifically Δ9-tetrahydrocannabinol (Δ9-THC), could potentially influence early embryonic development. Δ9-THC was found to augment the proliferation of naïve mouse ESCs, but not formative Epiblast-like Cells (EpiLCs). This enhanced proliferation relies on binding to the CB1 receptor. Moreover, Δ9-THC exposure was noted to boost glycolytic

rates and anabolic capabilities in mESCs. The metabolic adaptations brought on by $\Delta 9$ -THC exposure persisted during differentiation into Primordial Germ Cell-Like Cells (PGCLCs), even when direct exposure ceased, and correlated with a shift in their transcriptional profile. This study provides the first comprehensive molecular assessment of the effects of $\Delta 9$ -THC exposure on mouse ESCs and their early derivatives. The manuscript underscores the potential ramifications of cannabis exposure on early embryonic development and pluripotent stem cells. However, it is important to note the limitations of this study: firstly, all experiments were conducted in vitro, and secondly, the study lacks analogous experiments in human models.

Reviewer #2 (Recommendations For The Authors):

1. EpiLCs, characterized as formative pluripotent stem cells rather than primed ones, are a transient population during ESC differentiation. The authors should consider using EpiSCs and/or formative-like PSCs (Yu et al., Cell Stem Cell, 2021; Kinoshita et al., Cell Stem Cell, 2021), and amend their references to EpiLCs as "formative".

Indeed, EpiLCs are a transient pluripotent stem cell population that is “functionally distinct from both naïve ESCs and EpiSCs” and “enriched in formative phase cells related to pre-streak epiblast” (Kinoshita et al., Cell Stem Cell, 2021). Here, we used the differentiation system developed by M. Saitou and colleagues to derive PGCLCs (Hayashi et al, 2011). Since EpiSCs are refractory to PGCLCs induction (Hayashi et al, 2011), we used the germline-competent EpiLCs and took advantage of a well-established differentiation system to derive mouse PGCLCs. Most authors, however, agree that in terms of epigenetic and metabolic profiles, mouse EpiLCs represent a primed pluripotent state. We have added that PGCs arise in vivo “from formative pluripotent cells in the epiblast” on lines 85-86.

1. Does the administration of $\Delta 9$ -THC, at concentrations from 10nM to 1uM, alter the cell cycle profiles of ESCs?

The proliferation of ESCs was associated with changes in the cell cycle, as presented in the new Supplementary Figure 2, which we discuss in lines 118-123.

1. Could $\Delta 9$ -THC treatment influence the differentiation dynamics from ESCs to EpiLCs?

No significant changes were observed in the pluripotency markers associated with ESCs and EpiLCs (Supplementary Figure 9). We have added this information in lines 277-279.

1. The authors should consider developing knockout models of cannabinoid receptors in ESCs and EpiLCs (or EpiSCs and formative-like PSCs) for control purposes.


This is an excellent suggestion. Due to time and resource constraints, however, we focused our mechanistic investigation of the role of CB1 on the use of rimonabant which revealed a reversal of $\Delta 9$ -THC-induced proliferation at 100nM.

1. Lines 134-136: "Importantly, SR141716 pre-treatment, while not affecting cell viability, led to a reduced cell count compared to the control, indicating a fundamental role for CB1 in promoting proliferation." Regarding Figure 2D, does the Rimonabant "+" in the "mock" group represent treatment with Rimonabant only? If that's the case, there appears to be no difference from the Rimonabant "-" mock. The authors should present results for Rimonabant-only treatment.

To be able to compare the effects +/- Rimonabant and as stated in the figure legend, each condition was normalized to its own control (mock with, or without Rimonabant). Author response image 3 is the unnormalized data showing the same effects of $\Delta 9$ -THC and Rimonabant on cell number.

Author response image 3.

Unnormalized data corresponding to the Figure 2D.



1. In Figure 3, both ESCs and EpiLCs show a significant decrease in oxygen consumption and glycolysis at a 10uM concentration. Do these conditions slow cell growth? BrdU incorporation experiments (Figure 1) seem to contradict this. With compromised bioenergetics at this concentration, the authors should discuss why cell growth appears unaffected.

Indeed, we believe that cell growth is progressively restricted upon increasing doses of $\Delta 9$ -THC (consider Supplementary Figure 2). In addition, oxygen consumption and glycolysis can be decoupled from cellular proliferation, especially considering the lower time ranges we are working with (44-48h).

1. Beyond $\Delta 9$ -THC exposure prior to PGCLCs induction, it would be also interesting to explore the effects of $\Delta 9$ -THC on PGCLCs during their differentiation.

We agree with the Reviewer. Our aim was to study whether exposure prior to differentiation could have an impact, and if so, what are the mediators of this impact. Full exposure during differentiation is another exposure paradigm that is relevant but would not have allowed us to show the metabolic memory of $\Delta 9$ -THC exposure. Future work, however, will be dedicated to analyzing the effect of continuous exposure through differentiation.

1. As PGC differentiation involves global epigenetic changes, it would be interesting to investigate how $\Delta 9$ -THC treatment at the ESCs/EpiLCs stage may influence PGCLCs' transcriptomes.

We also agree with the Reviewer. While this paper was not primarily focused on $\Delta 9$ -THC's epigenetic effects, we have explored the impact of $\Delta 9$ -THC on more than 100 epigenetic modifiers in our RNA-seq datasets. These results are shown in Supplementary Table 1 and Supplementary Figure 10 and discussed in lines 301-316.

1. Lines 407-408: *The authors should exercise caution when suggesting "potentially adverse consequences" based solely on moderate changes in PGCLCs transcriptomes.*

We agree and have modified the sentence as follows: "Our results thus show that exposure to $\Delta 9$ -THC prior to specification affects embryonic germ cells' transcriptome and metabolome. This in turn could have adverse consequences on cell-cell adhesion with an impact on PGC normal development in vivo."

1. *Investigating the possible impacts of $\Delta 9$ -THC exposure on cultured mouse blastocysts, implantation, post-implantation development, and fertility could yield intriguing findings.*

We thank the Reviewer for this comment. We have amended our discussion to include these points in the last paragraph.

1. *Given that naïve human PSCs and human PGCLCs differentiation protocols have been established, the authors should consider carrying out parallel experiments in human models.*

We have performed $\Delta 9$ -THC exposures in hESCs (Supplementary Figure 4 and Supplementary Figure 5), showing that $\Delta 9$ -THC alters the cell number and general metabolism of these cells. We present these results in light of the differences in metabolism between mouse and human embryonic stem cells on lines 135-141 and 185-188. Implications of these results are discussed in lines 474-486.

Reviewer #3 (Public Review):

Verdikt et al. focused on the influence of $\Delta 9$ -THC, the most abundant phytocannabinoid, on early embryonic processes. The authors chose an in vitro differentiation system as a model and compared the proliferation rate, metabolic status, and transcriptional level in ESCs, exposure to $\Delta 9$ -THC. They also evaluated the change of metabolism and transcriptome in PGCLCs derived from $\Delta 9$ -THC-exposed cells. All the methods in this paper do not involve the differentiation of ESCs to lineage-specific cells. So the results cannot demonstrate the impact of $\Delta 9$ -THC on preimplantation developmental stages. In brief, the authors want to explore the impact of $\Delta 9$ -THC on preimplantation developmental stages, but they only detected the change in ESCs and PGCLCs derived from ESCs, exposure to $\Delta 9$ -THC, which showed the molecular characterization of the impact of $\Delta 9$ -THC exposure on ESCs and PGCLCs.

Reviewer #3 (Recommendations For The Authors):

1. *To demonstrate the impact of $\Delta 9$ -THC on preimplantation developmental stages, ESCs are an appropriate system. They have the ability to differentiate three lineage-specific cells. The authors should perform differentiation experiments under $\Delta 9$ -THC-exposure, and detect the influence of $\Delta 9$ -THC on the differentiation capacity of ESCs, more than just differentiate to PGCLCs.*

We apologize for the lack of clarity in our introduction. We specifically looked at the developmental trajectory of PGCs because of the sensitivity of these cells to environmental insults and their potential contribution to transgenerational inheritance. We have expanded on these points in our introduction and discussion sections (lines 89-91 and 474-486). Because our data shows the relevance of $\Delta 9$ -THC-mediated metabolic rewiring in ESCs subsisting across differentiation, we agree that differentiation towards other systems (neuroprogenitors, for instance) would yield interesting data, albeit beyond the scope of the present study.

1. *Epigenetics are important to mammalian development. The authors only detect the change after $\Delta 9$ -THC-exposure on the transcriptome level. How about methylation landscape changes in the $\Delta 9$ -THC-exposure ESCs?*

We have explored the impact of $\Delta 9$ -THC on more than 100 epigenetic modifiers in our RNA-seq datasets. These results are shown in Supplementary Table 1 and Supplementary Figure 10, discussed in lines 301-316. While indeed the changes in DNA methylation profiles appear relevant in the context of $\Delta 9$ -THC exposure (because of Tet2 increased expression in EpiLCs), we highlight that other epigenetic marks (histone acetylation, methylation or ubiquitination) might be relevant for future studies.

1. *In the abstract, the authors claimed that "the results represent the first in-depth molecular characterization of the impact of $\Delta 9$ -THC exposure on preimplantation developmental stages." But they do not show whether the $\Delta 9$ -THC affects the fetus through the maternal-fetal interface.*

We have addressed the need for increased clarity and have modified the sentence as follows: "These results represent the first in-depth molecular characterization of the impact of $\Delta 9$ -THC exposure on early stages of the germline development."

1. *To explore the impact of cannabis on pregnant women, the human ESCs may be a more proper system, due to the different pluripotency between human ESCs and mouse ESCs.*

We have performed $\Delta 9$ -THC exposures in hESCs (Supplementary Figure 4 and Supplementary Figure 5). These preliminary results show that $\Delta 9$ -THC exposure negatively impacts the cell number and general metabolism of hESCs. With the existence of differentiation systems for hPGCLCs, future studies will need to assess whether $\Delta 9$ -THC-mediated metabolic remodelling is also carried through differentiation in human systems. We discuss these points in the last paragraph of our discussion section.

1. *All the experiments are performed in vitro, and the authors should validate their results in vivo, at least a $\Delta 9$ -THC-exposure pregnant mouse model.*

Our work is the first of its kind to show that exposure to a drug of abuse can alter the normal development of the embryonic germline. We agree with the Reviewer that to demonstrate transgenerational inheritance of the effects reported here, future experiments in an in vivo mouse model should be conducted. The metabolic remodeling observed upon cannabis exposure could also be directly studied in a human context, although these experiments would be beyond the scope of the present study. For instance, changes in glycolysis may be detected in pregnant women using cannabis, or directly measured in follicular fluid in a similar manner as done by Fuchs-Weizman and colleagues (Fuchs-Weizman et al., 2021). We hope that our work can provide the foundation to inform such in vivo studies.