

Broca's area, variation and taxic diversity in early *Homo* from Koobi Fora (Kenya)

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

September 4, 2023 (this version)

Reviewed preprint version 1

June 30, 2023

Posted to bioRxiv

June 7, 2023

Sent for peer review

May 10, 2023

Amélie Beaudet , Edwin de Jager

Laboratoire de Paléontologie, Évolution, Paléoécosystèmes et Paléoprimatologie (PALEVOPRIM), UMR 7262 CNRS & University of Poitiers, 6 Rue Michel Brunet, 86000 Poitiers, France • Department of Archaeology, University of Cambridge, Henry Wellcome Building, Fitzwilliam St, Cambridge CB2 1QH • School of Geography, Archaeology and Environmental Studies, University of the Witwatersrand, Private Bag 3, WITS 2050, South Africa

 https://en.wikipedia.org/wiki/Open_access

 <https://creativecommons.org/licenses/by/4.0/>

Abstract

Because brain tissues rarely fossilize, pinpointing when and how modern human cerebral traits emerged in the hominin lineage is particularly challenging. The fragmentary nature of the fossil material, coupled with the difficulty of characterizing such a complex organ, have been the source of long-standing debates. Prominent among them is the uncertainties around the derived or primitive state of the brain organization in the earliest representatives of the genus *Homo*, more particularly in key areas such as the Broca's area. By revisiting a particularly well-preserved fossil endocast from the Turkana basin (Kenya) attributed to early *Homo*, here we confirm that humans in Africa had a primitive organization of the Broca's area ca. 1.9 million years ago. Additionally, our description of KNM-ER 3732 adds further information about the variation pattern of the inferior frontal gyrus in fossil hominins, with implications for early *Homo* taxic diversity (i.e., one or two *Homo* species at Koobi Fora) and the nature of the mechanisms involved in the emergence of derived cerebral traits.

eLife assessment

This **important** study uses the brain endocast of a ~1.9-million-year-old hominin fossil from Kenya, attributed to genus *Homo*, to show that the organization of the Broca's area in members of early *Homo* was primitive. Specifically, the prefrontal sulcal pattern in this early *Homo* specimen more closely resembles that of chimpanzees than of modern humans. Because Broca's area is associated with speech function, the **compelling** evidence from this study is relevant for understanding the timing and trajectory of evolution of speech related traits in our genus. Coupled with its potential implications for taxonomic classification, this study will be of interest to paleoanthropologists, paleontologists, archaeologists, and neuroscientists.

The modern human brain is an exceptionally complex, highly specialized and extremely costly machinery. Because of the fragmentary nature of the hominin fossil record, assessing when and how changes in the brain of our ancestors happened, and inferring any related functional, behavioral and metabolic consequences, is particularly challenging (Zollikofer and Ponce de León, 2013). Nonetheless, reconstructing the chronological and taxonomical context of the emergence of derived cerebral traits is a prerequisite for disentangling underlying evolutionary processes. For instance, local reorganization of specific areas in an overall primitive hominin brain would support a mosaic-like evolutionary pattern (e.g., Holloway et al., 2004), and raise essential questions on the role of selection pressure (or absence of; rev. in Beaudet, 2021). Beyond the value of such information on the origins of the human brain, the assumption that *Homo* developed a uniquely complex brain organization (e.g., Tobias, 1987) requires further evidence.

Although brains rarely fossilize, it is possible to glean structural information about the evolutionary history of the hominin brain by studying sulcal imprints in fossil brain endocasts (Neubauer et al., 2014). In this regard, the Broca's area has been the focus of much interest in paleoneurology due to striking structural differences between extant human and chimpanzee brains and endocasts, and the implication of this area in articulated language (rev. in Beaudet, 2017). Ponce de León et al. (2021) thoroughly examined brain endocasts of *Homo* specimens in eastern Africa and Eurasia and demonstrated that the organization of the Broca's area in the earliest representatives of the genus before 1.5 Ma was primitive. Because the imprints of the Broca's cap in endocasts are not always readable, they used the coronal suture and surrounding sulcal imprints as a proxy to identify frontal lobe expansion (i.e., derived condition). Unfortunately, in some of the oldest and key specimens from Africa that could represent the >1.5 Ma condition (e.g., KNM-ER 1470) the interpretations remain inconclusive.

Testing the hypothesis of Ponce de León et al. (2021) of a primitive brain in the earliest representatives of the genus *Homo* before 1.5 Ma thus necessitates (i) an excellent preservation of very fine neuroanatomical details in fossil endocasts, and (ii) reliable information on their taxonomic identity (i.e., *Homo*) and stratigraphic context (i.e., before/after 1.5 Ma). The hominin specimen KNM-ER 3732 from Kenya, which fulfills these requirements, has the potential to shed new light on this conundrum. KNM-ER 3732 was discovered during the 1974-1975 field programme in the Koobi Fora Formation (area 115), east of Lake Turkana in Kenya (Leakey, 1976; Wood, 1991). KNM-ER 3732 consists of a calotte, left zygoma and a natural endocast. The expanded neurocranium and robust upper face supports an attribution to *Homo* sp. indet. (Wood, 1991). The specimen was lying below the KBS Tuff of the upper Burgi Member that is dated to 1.87 million years ago, which provides a minimum age (Feibel et al., 1983, 2009). Accordingly, if KNM-ER 3732 shows a derived cerebral condition, the hypothesis of >1.5 Ma *Homo* being associated to a primitive organization of the Broca's area is falsified. Here we provide a comparative study of the natural endocast of the pre-1.5 Ma *Homo* specimen KNM-ER 3732 to test the hypothesis of a late emergence of a modern Broca's area in the hominin lineage.

Results

KNM-ER 3732 offers a glimpse of early *Homo* prefrontal organization

The natural endocast of KNM-ER 3732, initially described by Holloway et al. (2004), preserves the dorsal part but misses the frontal pole, the occipital and temporal lobes, as well as the entire ventral surface (Fig. 1). Holloway et al. (2004) estimated the endocranial volume as about 750-800 cc but pointed out the low reliability of their estimate. Wood (1991: 132) identified an

“irregular bony excrescence on the surface of the right parietal at the level of lambda and 32 mm from the midline. It projects 6 mm from the surface of the bone, and may be an example of myositis ossificans associated with damage to the overlying temporal muscle”. Because of the location (right parietal) and nature (traumatic) of the outgrowth, it does not affect our area of study (i.e., left frontal bone) (Walczak et al., 2015 [↗](#)). The gyral and sulcal details are well-preserved, particularly in the prefrontal region. The superior, middle and inferior frontal sulci are visible on both hemispheres. On the left hemisphere, the Broca’s cap is prominent. A vertical groove, identified as the precentral sulcus, separates the Broca’s cap in half (**Fig. 2A** [↗](#)). On both hemispheres, the central sulcus intersects the inter-hemispheric scissure and the post-central sulcus seems to be connected to the lateral fissure. The intra-parietal sulcus can be found in the parieto-occipital region of both hemispheres. The examination of the internal surface of the braincase (**Fig. 2B** [↗](#)) confirms the pattern described on the natural endocast.

KNM-ER 3732 has a primitive prefrontal cortex

When compared to extant human and chimpanzee sulcal patterns, KNM-ER 3732 closely resembles the latter (**Fig. 2C-D** [↗](#)). In extant human brains and endocasts, the inferior frontal sulcus often transects the Broca’s cap, while the ascending ramus of the lateral fissure caudally borders the prominence (**Figure 2C-D** [↗](#); Connolly, 1950 [↗](#); de Jager et al., 2019 [↗](#), 2022 [↗](#)). In chimpanzees, the central sulcus is placed more rostrally, and the inferior portion of the precentral sulcus bisects the Broca’s cap that is bordered rostrally by the inferior frontal sulcus (**Figure 2E-F** [↗](#); Connolly, 1950 [↗](#); Falk et al., 2018 [↗](#)). The coronal suture runs in between the precentral and central sulci in KNM-ER 3732, which points towards the primitive configuration described in Ponce de León et al. (2021) [↗](#). The sulcal pattern seen in the prefrontal cortex of KNM-ER 3732, that was detected in both the natural endocast and the inner surface of the cranium (**Fig. 2A-B** [↗](#)), and more particularly the deeply marked precentral sulcus that incised the Broca’s cap, is not found in any of the early *Homo* specimens described in Ponce de León et al. (2021) [↗](#) but approximates the condition seen in the Dmanisi cranium D2282 (Fig. S1B in Ponce de León et al., 2021 [↗](#)). However, this pattern is seen in a contemporaneous non-*Homo* hominin specimen in South Africa, i.e., *Australopithecus sediba* (Falk, 2014 [↗](#)).

Discussion

Paleoneuroanatomy supports taxic diversity within early *Homo*

Overall, the present study not only demonstrates that Ponce de León et al.’s (2021) hypothesis of a primitive brain of early *Homo* cannot be rejected, but also adds information about the variation pattern of the inferior frontal gyrus. In particular, the diversity of the prefrontal sulcal patterns of hominin endocasts at Koobi for a revives the debate about the possible presence of two early *Homo* species in this locality (i.e., *Homo habilis sensu stricto* and *Homo rudolfensis*). In 1983, Falk (1983) [↗](#) published the description of the endocasts of two *Homo* specimens from Koobi Fora, KNM-ER 1470 and KNM-ER 1805. Her analysis supported the co-existence of two morphs, KNM-ER 1470 representing a more derived human-like sulcal pattern. Interestingly, cerebral evidence brought up by her analysis matched other studies that emphasized the more derived craniodental anatomy of specimens attributed to *Homo rudolfensis* as opposed to the more primitive (*Australopithecus*-like) traits identified within *Homo habilis sensu stricto* (Leakey et al., 2012 [↗](#)). While describing the external morphology of the neurocranium, Leakey (1976 [↗](#): 575) noted that KNM-ER 3732 was “strikingly similar to KNM-ER 1470”. This resemblance is not reflected in their cerebral organization since the present study rather suggests a primitive organization of the Broca’s cap in KNM-ER 3732. If KNM-ER 1470 had indeed a derived brain, taxic diversity as a source of variation cannot be discarded. If we go further down that route, the similarities between




Figure 1.

3D virtual rendering of the natural endocast of KNM-ER 3732 and identification of sulcal imprints. KNM-ER 3732 is shown in lateral right (**A**), lateral left (**B**), dorsal (**C**) and posterior (**D**) views. ar: ascending ramus of the lateral fissure; c: central sulcus; CO: coronal suture; fi: inferior frontal sulcus; fm: middle frontal sulcus; ip: intra-parietal sulcus; pc: pre-central sulcus; pt: post-central sulcus. Scale bar: 2 cm.




Figure 2.

Comparison of the sulcal patterns identified in the inferior frontal area of KNM-ER 3732. The natural endocranial cast (**A**) and braincase (**B**) of KNM-ER 3732 are compared to the virtual endocranial casts of extant human (**C**) and chimpanzee (**D**) individuals. Images not to scale. ar: ascending ramus of the lateral fissure; c: central sulcus; co: coronal suture; fi: inferior frontal sulcus; fm: middle frontal sulcus; ip: intra-parietal sulcus; pc: pre-central sulcus; pt: post-central sulcus.

KNM-ER 3732 and *Australopithecus sediba* revealed by our study could be an argument supporting the presence of *Australopithecus* in Koobi Fora or to the absence of a definite threshold between the two genera based on the morphoarchitecture of their endocasts (Wood and Collard, 1999 [↗](#)).

The evolutionary history of the human Broca's area unravelled

Beyond the taxonomic aspect, variation detected in Koobi Fora could provide information about underlying evolutionary mechanisms, and more specifically the process of fixation of an adaptive variant, i.e., a new organization of the Broca's area and the increase of neural interconnectivity in this region (Van Essen, 1987 [↗](#)). Such neurological changes might have had deep implication in the emergence of novel behaviors, such as articulated language (Beaudet, 2017 [↗](#), 2021 [↗](#)). Within this scenario, the study from Ponce de León et al. (2021) [↗](#) would suggest that this trait became stabilized by 1.5 million years ago. The identification of protracted brain growth as early as in *Australopithecus afarensis* (Gunz et al., 2020 [↗](#)), inducing longer exposure to social environment during brain maturation (rev. in Hublin et al., 2015 [↗](#)), could be consistent with a derived organization of the Broca's area being selected as a response to social environmental stimuli through developmental plasticity, culminating in this variant becoming dominant within *Homo*. In parallel, the possibility of allometric scaling and influence of brain size on sulcal patterns in early *Homo* has to be further explored.

Materials and Methods

Materials

KNM-ER 3732 is currently housed in the National Museums of Kenya in Nairobi (Kenya). We used brain and endocast atlases published in Connolly (1950) [↗](#), Falk et al. (2018) [↗](#) and de Jager et al. (2019 [↗](#), 2022 [↗](#); see also www.endomap.org (<http://www.endomap.org/>) [↗](#)) for comparing the pattern identified in KNM-ER 3732 to those described in extant humans and chimpanzees. To the best of our knowledge, these atlases are the most extensive atlases of extant human and chimpanzee brains/endocasts available to date and are widely used in the literature to explore variability in sulcal patterns. In **Figure 2** [↗](#), the extant human and chimpanzee conditions are illustrated by one extant human (adult female) and one extant chimpanzee (adult female) specimens from the Pretoria Bone Collection at the University of Pretoria (South Africa) and in the Royal Museum for Central Africa in Tervuren (Belgium), respectively (Beaudet et al., 2018 [↗](#)).

Digitization

Both the natural endocast and the cranium of KNM-ER 3732 were scanned using an Artec Space Spider 3D scanner and reconstructed with the software Artec Studio 16X. The 3D mesh can be viewed on MorphoSource (<https://www.morphosource.org/concern/media/000497752?locale=en>) [↗](#). The comparative specimens were imaged by microfocus X-ray tomography (Beaudet et al., 2018 [↗](#)) at the South African Nuclear Energy Corporation in Pelindaba (South Africa) and at the Centre for X-ray Tomography of Ghent University (UGCT) in Ghent (Belgium). Virtual endocasts were generated using Endex software (Subsol et al., 2010 [↗](#)).

Detection and identification of sulcal imprints

Sulcal imprints were automatically detected through a geometry-based method using curvature lines computed from the natural endocast and the inner surface of the cranium. Sulcal imprints, considered as variation points of the surface on a triangle mesh, were detected through a geometry-based method using curvature lines defined as salient subsets of the extrema of the principal curvatures on surfaces (Yoshizawa et al., 2008 [↗](#); Beaudet et al., 2016 [↗](#)). Vascular imprints and non-anatomical structures (e.g., fractures) were manually removed through a customized script written in MATLAB R2013a (Mathworks).

Acknowledgements

We thank Emmanuel Ndiema and the curatorial staff of the National Museums of Kenya for collection access. For scientific/technical discussion we are grateful to: G. Castelli (Cambridge), M. Mirazón-Lahr (Cambridge), R. Holloway (New York), F. Spoor (London). AB is funded by the National Research Foundation of South Africa (Research Development Grants for Y-Rated Researchers, grant number 129336), the University of Cambridge and the Centre National de la Recherche Scientifique. EdJ is funded by the University of Cambridge Harding Distinguished Postgraduate Scholars Programme. AB and EdJ are funded by the South Africa/France (PROTEA) Joint Research Programme (grant number 129923) and the McDonald Institute for Archaeological Research. We are grateful to the three reviewers and the editors for their comments and suggestions.

Competing Interest Statement

The authors declare no competing interest.

References

- Beaudet A. (2017) **The emergence of language in the hominin lineage: perspectives from fossil endocasts** *Frontiers in Human Neuroscience* **11**
- Beaudet A. (2021) **The enigmatic origins of the human brain** *Science* **372**:124–125
- Beaudet A, Dumoncel J, de Beer F, Duployer B, Durrleman S, Gilissen E, Hoffman J, Tenailleau C, Thackeray JF, Braga J. (2016) **Morphoarchitectural variation in South African fossil cercopithecoid endocasts** *Journal of Human Evolution* **101**:65–78
- Beaudet A, Carlson KJ, Clarke RJ, de Beer F, Dhaene J, Heaton J, Pickering TR, Stratford D. (2018) **Cranial vault thickness variation and inner structural organization in the StW 578 hominin cranium from Jacovec Cavern, South Africa** *Journal of Human Evolution* **121**:204–220
- Connolly CJ (1950) **External Morphology of the Primate Brain**. CC Thomas
- de Jager EJ, van Schoor AN, Hoffman JW, Oettlé AC, Fonta C, Mescam M, Risser L, Beaudet A. (2019) **Sulcal pattern variation in extant human endocasts** *Journal of Anatomy* **235**:803–810
- de Jager EJ, Risser L, Mescam M, Fonta C, Beaudet A. (2022) **Sulci 3D mapping from human cranial endocasts: A powerful tool to study hominin brain evolution** *Human Brain Mapping* **43**:4433–4443
- Falk D. (1983) **Cerebral cortices of East African early hominids** *Science* **221**:1072–1074
- Falk D. (2014) **Interpreting sulci on hominin endocasts: old hypotheses and new findings** *Frontiers in Human Neuroscience* **8**
- Falk D, Zollikofer CPE, Ponce de León M, Semendeferi K, Alatorre Warren JL, Hopkins WD (2018) **Identification of in vivo sulci on the external surface of eight adult chimpanzee brains: implications for interpreting early hominin endocasts** *Brain, Behavior and Evolution* **91**:45–58
- Feibel CS, Brown FH, McDougall I. (1983) **Stratigraphic context of fossil hominids from the Omo groups deposits: northern Turkana Basin, Kenya and Ethiopia** *American Journal of Physical Anthropology* **78**:595–622
- Feibel CS, Lepre CJ, Quinn RL (2006) **Stratigraphy, correlation, and age estimates for fossils from Area 123, Koobi Fora** *Journal of Human Evolution* **57**:112–122
- Gunz P, Neubauer S, Falk D, Tafforeau P, Le Cabec A, Smith TM, Kimbel WH, Spoor F, Alemseged Z. (2020) **Australopithecus afarensis endocasts suggest ape-like brain organization and prolonged brain growth** *Science Advances* **6**
- Holloway RL, Broadfield DC, Yuan MS (2004) **The Human Fossil Record: Brain Endocasts, the Paleoneurological Evidence**. Wiley-Liss
- Hublin JJ, Neubauer S, Gunz P. (2015) **Brain ontogeny and life history in Pleistocene hominins** *Philosophical Transactions of the Royal Society B: Biological Sciences* **2370**

- Leakey REF (1976) **New hominid fossils from the Koobi Fora formation in Northern Kenya** *Nature* **261**:574–576
- Leakey MG, Spoor F, Dean MC, Feibel CS, Antón SC, Kiarie C, Leakey LN (2012) **New fossils from Koobi Fora, northern Kenya, confirm taxonomic diversity in early Homo** *Nature* **488**:201–204
- Neubauer S. (2014) **Endocasts: Possibilities and limitations for the interpretation of human brain evolution** *Brain, Behavior and Evolution* **84**:117–134
- Ponce de León MS, Bienvenu T, Marom A, Engel S, Tafforeau P, Alatorre Warren JL, Lordkipanidze D, Kurniawan I, Murti DB, Suriyanto RA, Koesbardiati T, Zollikofer CPE (2021) **The primitive brain of early Homo** *Science* **372**:165–171
- Subsol G, Gesquière G, Braga J, Thackeray F. (2010) **3D automatic methods to segment “virtual” endocasts: state of the art and future directions** *American Journal of Physical Anthropology* **141**:226–227
- Tobias PV (1987) **The brain of Homo habilis: a new level of organization in cerebral evolution** *Journal of Human Evolution* **16**:741–761
- Van Essen DC (1987) **A tension-based theory of morphogenesis and compact wiring in the central nervous system** *Nature* **335**
- Walczak BE, Johnson CN, Howe BM (2015) **Myositis ossificans** *Journal of the American Academy of Orthopaedic Surgeons* **23**:612–622
- Wood B. (1991) **Koobi Fora Research Project**
- Wood B, Collard M. (1999) **The human genus** *Science* **284**:65–71
- Yoshizawa S, Belyaev A, Yokota H, Seidel HP (2008) **Fast, robust, and faithful methods for detecting crest lines on meshes** *Computer Aided Geometric Design* **25**:545–560
- Zollikofer CPE, Ponce de León MSP. (2013) **Pandora’s growing box: Inferring the evolution and development of hominin brains from endocasts** *Evolutionary Anthropology* **22**:20–33

Author information

Amélie Beaudet

Laboratoire de Paléontologie, Évolution, Paléoécosystèmes et Paléoprimatologie (PALEVOPRIM), UMR 7262 CNRS & University of Poitiers, 6 Rue Michel Brunet, 86000 Poitiers, France, Department of Archaeology, University of Cambridge, Henry Wellcome Building, Fitzwilliam St, Cambridge CB2 1QH, School of Geography, Archaeology and Environmental Studies, University of the Witwatersrand, Private Bag 3, WITS 2050, South Africa
For correspondence: beaudet.amelie@gmail.com
ORCID iD: [0000-0002-9363-5966](https://orcid.org/0000-0002-9363-5966)

Edwin de Jager

Laboratoire de Paléontologie, Évolution, Paléoécosystèmes et Paléoprimatologie (PALEVOPRIM), UMR 7262 CNRS & University of Poitiers, 6 Rue Michel Brunet, 86000 Poitiers, France

Editors

Reviewing Editor

Yonatan Sahle

University of Cape Town, South Africa

Senior Editor

George Perry

Pennsylvania State University, United States of America

Reviewer #1 (Public Review):

The cerebral cortex, or surface of the brain, is where humans do most of their conscious thinking. In humans, the grooves (sulci) and bumps (convolutions) have a particular pattern in a region of the frontal lobe called Broca's area, which is important for language. Specialists study features imprinted on the internal surfaces of braincases in early hominins by casting their interiors, which produces so-called endocasts. A major question about hominin brain evolution concerns when, where, and in which fossils a humanlike Broca's area first emerged, the answer to which may have implications for the emergence of language. The researchers used advanced imaging technology to study the endocast of a hominin (KNM-ER 3732) that lived about 1.9 million years ago (Ma) in Kenya to test a recently published hypothesis that Broca's remained primitive (apelike) prior to around 1.5 Ma. The results are consistent with the hypothesis and raise new questions about whether endocasts can be used to identify the genus and/or species of fossils.

Reviewer #2 (Public Review):

The authors present new data of endocranial surface details from the early Homo specimen KNM-ER 3732 and discuss the evolution of brain surface features that might be related to the evolution of language in the hominin lineage.

Comments and issues raised by the reviewers have been addressed adequately. I am sure that this contribution will revive discussion about these issues.

Reviewer #3 (Public Review):

The authors provide a detailed analysis of the sulcal and sutural imprints preserved on the natural endocast and associated cranial vault fragments of the KNM-ER3732 early Homo specimen. The analyses indicate a primitive ape-like organization of this specimen's frontal cortex. Given the geological age of around 1.9 million years, this is the earliest well-documented evidence of a primitive brain organization in African Homo.

The various points raised by the reviewers and the responses provided by the authors illustrate that paleoneurology is a research field where little consensus has been reached over the past century. This is due not only to the fragmentary preservation of most fossil endocasts, but also to the limitations of scientific inference in general, and paleoneurological inference in particular. Like any scientific hypothesis, a paleoneurological hypothesis cannot be proven, but at best be falsified, leaving a wide field of possible alternative hypotheses. Furthermore, endocranial morphology does not equate cerebral morphology. A classical

example: the endocranial Broca cap is not identical to the cortical Broca area. And last but not least, taxonomy cannot resolve questions of phylogeny.

Author Response

The following is the authors' response to the original reviews.

Reviewer #1 (Public Review):

The cerebral cortex, or surface of the brain, is where humans do most of their conscious thinking. In humans, the grooves (sulci) and bumps (convolutions) have a particular pattern in a region of the frontal lobe called Broca's area, which is important for language. Specialists study features imprinted on the internal surfaces of braincases in early hominins by casting their interiors, which produces so-called endocasts. A major question about hominin brain evolution concerns when, where, and in which fossils a humanlike Broca's area first emerged, the answer to which may have implications for the emergence of language. The researchers used advanced imaging technology to study the endocast of a hominin (KNM-ER 3732) that lived about 1.9 million years ago (Ma) in Kenya to test a recently published hypothesis that Broca's remained primitive (apelike) prior to around 1.5 Ma. The results are consistent with the hypothesis and raise new questions about whether endocasts can be used to identify the genus and/or species of fossils.

We would like to thank Rev. 1 for their comments on our paper.

Reviewer #2 (Public Review):

The authors tried to support the hypothesis that early Homo still had a primitive condition of Broca's cap (the region in fossil endocasts corresponding to Broca's area in the brain), being more similar to the condition in chimpanzees than in humans. The evidence from the described individual points to this direction but there are some flaws in the argumentation.

We are grateful to Rev. 2 for their comments, although we partially agree with some of them.

First, we would like to rectify the statement of Rev. 2 that we “tried to support the hypothesis that early Homo still had a primitive condition of Broca's cap”, indeed, our aim was to test this hypothesis and not to try to validate it.

First, only one human and one chimpanzee were used for comparison, although we know that patterns of brain convolutions (and in addition how they leave imprints in the endocranial bones) are very variable.

We understand the point raised by Rev. 2 about the variation of brain convolutions in humans and chimpanzees. We used atlases published by Connolly (1950), Falk et al. (2018) and de Jager et al. (2019, 2022) to analyse the endocast of KNM-ER 3732 and compare it to the extant human and chimpanzee cerebral conditions. However, in Figure 2, for the sake of clarity only two Homo and Pan specimens were used to illustrate the comparison (as it has been done in other published papers, e.g., Carlson et al., 2011; Science, Gunz et al., 2020 Sci Adv). In the revised version, we modified the manuscript to explain further our approach (line 156) “We used brain and endocast atlases published in Connolly (1950), Falk et al. (2018)

and de Jager et al. (2019, 2022; see also www.endomap.org (<http://www.endomap.org/>)) for comparing the pattern identified in KNM-ER 3732 to those described in extant humans and chimpanzees. To the best of our knowledge, these atlases are the most extensive atlases of extant human and chimpanzee brains/endocasts available to date and are widely used in the literature to explore variability in sulcal patterns. In Figure 2, the extant human and chimpanzee conditions are illustrated by one extant human (adult female) and one extant chimpanzee (adult female) specimens from the Pretoria Bone Collection at the University of Pretoria (South Africa) and in the Royal Museum for Central Africa in Tervuren (Belgium), respectively (Beaudet et al., 2018).”.

Second, the evidence from this fossil specimen adds to the evidence of previously describe individuals but still not yet fully prove the hypothesis.

We tempered our discussion by concluding that (line 116) “Overall, the present study not only demonstrates that Ponce de León et al.’s (2021) hypothesis of a primitive brain of early Homo cannot be rejected, but also adds information [...]”.

Third, there is a vicious circle in using primitive and derived features to define a fossil species and then using (the same or different) features to argue that one feature is primitive or derived in a given species. In this case, we expect members of early Homo to be derived compared to their predecessors of the genus Australopithecus and that's why it seems intriguing and/or surprising to argue that early Homo has primitive features. However, we should expect that there is some kind of continuum or mosaic in a time in which a genus "evolves into" another genus. This discussion requires far more discussions about the concepts we use, maybe less discussion about what is different between the two groups but more discussion about the evolutionary processes behind them.

We fully agree with Rev. 2 on this aspect. We believe that identifying these differences/similarities between fossil and extant hominids constitute the first step of a better understanding of the evolutionary mechanisms. Our work suggests indeed a certain continuity between genera and raises questions on the genus concept and how to interpret the specimens currently attributed to early Homo. In the revised version of the manuscript we included a reference to this possible scenario (line 134): “[...] or to the absence of a definite threshold between the two genera based on the morphoarchitecture of their endocasts (Wood and Collard, 1999).”.

Fourth, the data of convolutional imprints presented are rather subjective when identifying which impressions represent which brain convolutions. Not seeing an impression does not necessarily mean that the corresponding brain feature did not exist. Interestingly, the manuscript does not mention and discuss at all the frontoorbital sulcus. This is a sulcus that usually runs from the orbital surface of the frontal lobe up to divide the inferior frontal gyrus in chimpanzees, a condition totally different than in humans who do not have a frontoorbital sulcus. Could such a sulcus be identified, this would provide a far more convincing argument for a primitive condition in this specimen. In Australopithecus sediba, e.g., the condition in this region seems to be a mosaic in which some aspects of the morphology seem to be more modern while one of the sulcal impressions can well be interpreted as a short frontoorbital sulcus. For this specimen, by the way, I would come back to my third point above: some experts in the field might argue that this specimen could belong to Homo rather than Australopithecus...

We agree that the presence of a fronto-orbital sulcus would be more conclusive. However, this sulcus has not been identified in KNM-ER3732 and the region in which we would expect

to find it is not preserved. As demonstrated by Ponce de León et al. (2021), because of the topographic relationships between sulci (and cranial structures), it is possible to interpret imprints on endocasts and the evolutionary polarity of some traits even in the absence of landmarks such as the fronto-orbital sulcus. In *Australopithecus sediba* the main derived feature of the endocast corresponds to the ventrolateral bulge in the left inferior frontal gyrus, and not to the sulcal pattern itself (Carlson et al., 2011 Science). However, the discussion around the taxonomic status of this taxon confirms the urgent need for reconsidering specimens from that time period and clarifying the mosaic-like or concerted evolution of the derived Homo-like traits within our lineage. Regarding the subjective nature of this approach, we invite readers to examine the specimen on MorphoSource (<https://www.morphosource.org/concern/media/000497752?locale=en>) and to request access to the National Museums of Kenya to the physical or virtual specimen to falsify our hypothesis.

According to my arguments above, I think that this manuscript might revive interesting discussions about this topic but it is not likely to settle them because the data presented are not strong enough to fully support the hypothesis.

We would be more than happy to consider new/other specimens with similar chronological and geographical contexts and investigate further this hypothesis in the future.

Reviewer #3 (Public Review):

The authors provide a detailed analysis of the sulcal and sutural imprints preserved on the natural endocast and associated cranial vault fragments of the KNM-ER3732 early Homo specimen. The analyses indicate a primitive ape-like organization of this specimen's frontal cortex. Given the geological age of around 1.9 million years, this is the earliest well-documented evidence of a primitive brain organization in African Homo.

In the discussion, the authors re-assess one of the central questions regarding the evolution of early Homo: was there species diversity, and if yes, how can we ascertain it? The specimen KNM-ER1470 has assumed a central role in this debate because it purportedly shows a more advanced organization of the frontal cortex compared to other largely coeval specimens (Falk, 1983). However, as outlined in Ponce de León et al. 2021 (Supplementary Materials), the imprints on the ER1470 endocranium are unlikely to represent sulcal structures and are more likely to reflect taphonomic fracturing and distortion. Dean Falk, the author of the 1983 study, basically shares this view (personal communication). Overall, I agree with the authors that the hypothesis to be tested is the following: did early Homo populations with primitive versus derived frontal lobe organizations coexist in Africa, and did they represent distinct species?

I greatly appreciate that the authors make available the 3D surface data of this interesting endocast.

We are grateful to Rev. 3 for their comments and for contextualizing our finding. We would also like to point out that, although the 3D surface can be viewed on MorphoSource, permission from the National Museums of Kenya has to be requested for studying the specimen and getting access to the physical specimen and/or the 3D model.

Reviewer #1 (Recommendations For The Authors):

Holloway, Broadfield & Yuan (2004) estimate ER 3732 as having a cranial capacity of 750 cc, which is larger than chimps and australopiths and similar to ER 1470 (752 cc, same reference). (That for Dmanisi 2282 is somewhat smaller at around 650 cc.) Cranial capacities should be mentioned along with added discussion about possible allometric scaling of (increased) numbers of sulci with increasing brain size as well as possible shifts

in locations of sulci relative to cranial sutures in larger-brained (including due to ontogenetic maturation) in individuals/species. Could these variables (especially brain size) be relevant for your discussion/conclusions?

We thank Rev. 1 for their suggestion. We included the estimate by Holloway et al. (2004) (line 95): “Holloway et al. (2004) estimated the endocranial volume as about 750-800 cc but insisted on the low reliability of their estimate.”. Additionally, we raised the possibility of potential allometric effect (line 149): “In parallel, the possibility of allometric scaling and influence of brain size on sulcal patterns in early Homo has to be further explored.” for future discussion.

From the two figures, it appears that the authors produced a virtual endocast from the cranial remains of ER 3732 and compared its features with those seen on a virtual reproduction of the corresponding natural endocast. If so, this needs to be clarified in the text, not just the figures.

We thank Rev. 1 for their suggestions that were integrated.

Reviewer #3 (Recommendations For The Authors):

While the sulcal imprints on the left hemisphere can be interpreted unambiguously, the anatomical assignment of those on the right side may need to be reconsidered, as they are more ambiguous. For example, the postcentral sulcus (pt) almost touches the middle frontal sulcus, which is an unlikely natural configuration.

We agree that the configuration on the right hemisphere is intriguing, especially when compared to the extant human and chimpanzee atlases. As such, we decided to change the label for what we think could be the inferior frontal sulcus and leave a question mark instead.

I encourage the authors to include:

- *a posterior view in Figure 1, and mark the lambdoid suture, parts of which seem to be preserved especially on the left side. This will help the readership to better understand which parts of the endocranial morphology are preserved.*
- *a scale bar would be of great utility to appreciate the small size of this specimen. The distance from bregma to the Broca cap seems to be short, indicating an endocranial volume much smaller than the published estimate of 750 ccm. Perhaps the authors can provide a new estimate, which would provide further support for the arguments proposed in the discussion section, especially the question of any presence of Australopithecus at Koobi Fora.*

We included a posterior view of the specimen in Figure 1 and scale bar and modified the legend accordingly. Unfortunately, we were not able to identify with certainty the feature that could correspond to the lambdoid suture. We might see the impression where the parietal bone meets the occipital bone, but there is a risk of misidentification (which is an issue frequently raised in the literature, see for example Gunz et al. 2020 Sci Adv). Concerning the endocranial volume, in the revised version of the manuscript we included the estimate by Holloway et al. (2004). Because the specimen only preserves the superior part, we are reluctant in providing an estimate of the total volume. However, we agree that this would be an interesting feature to integrate in the interpretation of this specimen.

Minor points

- *This sentence needs to be clarified: «The superior temporal sulcus nearly intersects the lateral fissure on the right hemisphere».*
- *The terms «Broca's region» and «orbital cap» need some more context. Do the authors mean «Broca's cap» in either instance?*

We clarified/modified when needed, thank you very much.

We included minor corrections in addition to those recommended by the reviewers:

-Lines 50, 74, 142, 149: “Broca’s area” instead of “Broca’s cap”

-Line 73: “in the pre-1.5 Ma Homo specimen” instead of “in pre-1.5 Ma Homo specimen”

-Line 100: we specified “in human brains and endocasts”

-Line 120: “sulcal pattern” instead of “sulcal patterns”

-Line 144: “behaviors” (plural)