

GTPase activating protein DLC1 spatio-temporally regulates Rho signaling

Max Heydasch, Lucien Hinderling, Jakobus van Unen, Maciej Dobrzynski, Olivier Pertz 

Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland • Graduate School for Cellular and Biomedical Sciences, University of Bern, Switzerland

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

December 14, 2023 (this version)

Sent for peer review

July 10, 2023

Posted to bioRxiv

June 19, 2023

 https://en.wikipedia.org/wiki/Open_access Copyright information

Abstract

Tightly regulated spatio-temporal Rho GTPase activity patterns regulate morphogenetic processes such as cell migration. Emerging evidence suggests that binding of Guanine nucleotide exchange factors (GEFs) and GTPase activating proteins (GAPs) to the cytoskeleton or adhesions mediate feedback regulation to spatio-temporal Rho GTPase activation. To explore such feedback regulation, we study the Rho specific GAP Deleted in Liver Cancer 1 (DLC1) which binds to focal adhesions (FAs) through mechanosensitive interactions. Using a FRET biosensor, we show that DLC1 loss of function leads to global increase in Rho activity and contractility throughout the cell without affecting a striking lamellar RhoA activity band in fibroblasts. To interrogate the Rho GTPase signaling flux, we build a genetic circuit consisting of an optogenetic actuator to control Rho activity, and a Rho activity biosensor. In spreading cells at steady state, optogenetic manipulation of Rho activity reveals that DLC1 controls the rate of Rho activation rather than duration, both at FAs and at the plasma membrane (PM). Local and reversible optogenetic control of contractility shows that DLC1 associates/dissociates with FAs during their reinforcement/relaxation. This might provide positive feedback that locally increases the rate of Rho activation at FAs that experience local tension to facilitate FA disassembly. Our results indicate that DLC1 operates both at the PM and at FAs to regulate global Rho activity levels at steady state, or to amplify local Rho activity at FAs experiencing a strong mechanical input, presumably to induce robust FA disassembly. This provides new insights in the complexity of spatio-temporal Rho GTPase signaling.

eLife assessment

This study presents a **valuable** finding on how the GAP DLC1, a deactivator of the small GTPase RhoA, regulates RhoA activity globally as well as at Focal Adhesions. Using a new acute optogenetic system coupled to a RhoA activity biosensor, the authors present **solid** evidence that DLC1 amplifies local Rho activity at Focal Adhesions. Nevertheless, the proposed mechanism could be further supported by a deeper analysis of the data.

Introduction

Rho GTPases regulate the cytoskeletal dynamics that power morphogenetic processes at the cell- and tissue level (Etienne-Manneville and Hall, 2002). Their activity is tightly controlled in time and space by GEFs and GAPs, which respectively activate and inhibit Rho GTPases. In the GTP-loaded, active state, Rho GTPases bind downstream effectors that regulate cytoskeletal dynamics and other cellular processes. Based on classic experiments that include measuring population-averaged Rho GTPase activity levels, as well as overexpressing dominant positive/negative Rho GTPase mutants, it was proposed that Rac1 controls lamellipodial protrusion, Cdc42 regulates filopodium formation, and RhoA controls myosin-based contractility.

Recent studies in which spatio-temporal Rho GTPase activation dynamics were measured using fluorescence resonance energy transfer (FRET)-based biosensors, revealed a much higher signaling complexity. During cell migration, RhoA, Rac1 and Cdc42 are all activated in membrane protrusions at the leading edge (Itoh et al., 2002; Kraynov et al., 2000; Nalbant et al., 2004; Pertz et al., 2006). Further, the three Rho GTPases are activated in specific spatio-temporal sequences during the protrusion/retraction cycles that fluctuate on time and length scales of seconds and single micrometers (Machacek et al., 2009). A further level of complexity is that these Rho GTPase activity sequences vary depending on the cell system (Hu et al., 2022; Machacek et al., 2009; Martin et al., 2016). This indicates that complex spatio-temporal Rho GTPase signaling programs regulate processes such as cell migration. Similarly as for Ras GTPases (Schmick et al., 2015), it is now clear that these Rho GTPase signaling patterns most likely result from a precise spatio-temporal regulation of Rho GTPase signaling fluxes controlled by GEFs and GAPs (Bement and von Dassow, 2014; Fritz and Pertz, 2016; Pertz, 2010). In this signaling flux, specific GEFs will be locally activated to load Rho GTPases with GTP depending on different cellular inputs. These GTP-loaded GTPases will then subsequently diffuse in the plasma membrane by virtue of their C-terminal lipid moiety until they encounter a locally activated GAP which eventually deactivates them. The interplay of GEFs, GAPs and diffusion of Rho GTPases in the PM will ultimately lead to the formation of a Rho GTPase activity pattern. Modelling studies have suggested that sophisticated dynamic signaling behaviors can emerge from the interplay of GEFs, GAPs, and GTPases within a signaling network (Tsyganov et al., 2012). Steady-state images of Rho GTPase activity patterns do not provide information about the whole Rho GTPase flux mentioned above.

This signaling complexity is consistent with the large amount of GEFs and GAPs that are ubiquitously expressed in cells (Fusco et al., 2016, 2016; Moon and Zheng, 2003; Mosaddeghzadeh and Ahmadian, 2021; Müller et al., 2020; Rossman et al., 2005). Recently, a system-wide screen has revealed that many GEFs and GAPs localize to cytoskeletal structures as well as adhesion complexes such as focal adhesions (FAs) (Müller et al., 2020). This strongly suggests the existence of feedback mechanisms from the cytoskeleton and FAs to Rho GTPase signaling. In this study, it was shown that Rac1-specific GEFs bind to FAs at the front, while Rac1-specific GAPs bind to FAs and at the back of the lamella of fibroblasts. This precise, asymmetric positioning of GEFs and GAPs might then regulate the Rho GTPase flux that produces the wide gradient of Rac activity observed at the leading edge of fibroblasts (Itoh et al., 2002; Kraynov et al., 2000; Martin et al., 2016). Such feedback regulation might allow leading edge Rac1 activity to constantly integrate mechanosensitive inputs from FAs, allowing dynamic regulation of Rac1 activity required to fine tune cell migration. Another prominent example of such spatio-temporal feedback regulation are the excitable RhoA activity patterns observed in the cortex during cytokinesis in *Xenopus Laevis* (Bement and von Dassow, 2014), that have also been observed in mammalian cells (Graessl et al., 2017). Here, RhoA activity waves that control F-actin wave patterns are spatially regulated by a RhoGAP that is locally regulated by F-actin (Bement et al., 2015a). Thus, cytoskeletal feedback to Rho GTPase regulation might be crucial to generate dynamic Rho GTPase signaling patterns. Understanding such spatio-temporal feedback regulation is not accessible with classic genetic approaches in which long term perturbation of Rho GTPases,

their regulators and effectors rapidly result in a new mechanochemical state of the cell, that will not be informative about an initial signaling/mechanical state of interest (Isogai and Danuser, 2018). Further, it is currently unknown which specific features of spatio-temporal Rho GTPase signaling are regulated to produce signaling patterns. For example, does negative regulation by a GAP solely control Rho GTPase signal termination, or could it also modulate, less intuitive parameters such as the rate of activation? Tackling such questions requires new tools to acutely and transiently perturb the Rho GTPase flux to understand its spatio-temporal regulation, directly in single living cells. Note that this experimental paradigm was successful at analyzing the Mitogen activated protein kinase (MAPK network) (Blum et al., 2019; Dessauges et al., 2022; Ryu et al., 2015).

A promising candidate to investigate how mechanical feedback from FAs can spatio-temporally shape Rho GTPase activity is the RhoA-specific GAP Deleted in Liver Cancer 1 (DLC1). DLC1 has been shown to both bind the plasma membrane (PM) (Erlmann et al., 2009), as well as FAs (Haining et al., 2018; Kaushik et al., 2014). At the PM, interaction with phosphatidylinositol-4,5-bisphosphate (PI(4,5)P2) can regulate GAP activity (Erlmann et al., 2009). At the FA, DLC1 interacts with the r7-r8 domains of the FA protein talin in a mechanosensitive fashion (Gingras et al., 2008; Goult et al., 2021; Haining et al., 2018; Zhao et al., 2022), as well as with Focal adhesion kinase (Li et al., 2011). When talin is in a state of low mechanical strain, it binds to DLC1, and presumably leads to local RhoA inactivation. However, upon force application, the r7-r8 domains unfold leading to dissociation of DLC1 from talin, leading to loss of negative RhoA regulation. This suggests that DLC1 is regulated at FAs by mechanosensitive interactions, which can relay information about the mechanical state of the cell to control RhoA activity.

In this work, we explore the spatio-temporal regulation of RhoA activity by DLC1. We show that knocking out DLC1 in REF52 fibroblasts leads to increased amplitude in RhoA throughout the cell. This is accompanied by an increased contractility that augments FAs and stress fibers (SFs). To explore how DLC1 contributes to Rho GTPase activity fluxes, we built a genetic circuit consisting of an optogenetic actuator to activate Rho with light, and a spectrally compatible biosensor to measure Rho activity. Transient optogenetic recruitment of a Rho GEF domain at FAs or at the PM led to an increase in the rate of activation of Rho in *DLC1* KO versus WT cells at both subcellular localizations. However, we found that in both control and DLC1 null cells, RhoA activation was more efficient at FAs compared to the PM. Further, acute and local manipulation of contractility with the optogenetic actuator revealed that DLC1 dynamically associates/dissociates with FAs under acute mechanical tension/relaxation. Our results suggest a complex mechanism in which DLC1-dependent Rho regulation can occur both at the PM and at FAs, with a dependence on mechanosensitive signals on the latter.

Results

DLC1 deficient cells show increased formation of SFs and FAs

To investigate the function of DLC1 in cytoskeletal and adhesion dynamics, we knocked down (KD) DLC1 in rat REF52 fibroblasts via small interfering RNA (siRNA). We also created a knockout (KO) cell line via Crispr/Cas9, that has a frameshift deletion at position 841 of Exon 5 of the gene, resulting in an early in frame stop codon 100 bp downstream of the cut. Gene-editing was validated by sequencing the endogenous locus. We were unable to test expression levels using western blot because of the absence of commercially available antibodies specific for rat DLC1. We then also rescued the DLC1 KO cells with an mCherry-labeled DLC1 construct using stable transfection with a piggyBac transposase system. REF52 fibroblasts were seeded at subconfluent density onto fibronectin-coated glass bottom wells and allowed to spread for one hour before fixation, leading to an isotropic spreading state in control, wild-type (WT) cells. To assess the impact of DLC1 deficiency on the formation of cytoskeletal and adhesion structures, we stained for

F-actin (phalloidin) and paxillin (**Figure 1A** [↗](#)) or phospho-myosin light chain (pMLC) (**Figure 1B** [↗](#)). Consistent with an expected increase in RhoA activity, we observed a phenotype of increased contractility as documented by increased lamella size, SF content, number of FAs, and increased pMLC. This increased contractility led to the loss of the isotropic spreading observed in WT cells. Rescue with the DLC1 construct reverted the cells to a less contractile phenotype that displayed a lamella of identical size as WT cells, enabling the characteristic isotropic spreading observed in WT cells. Quantification of these images revealed both an increase in lamella size (**Figure 1C** [↗](#)) and FA area (**Figure 1D** [↗](#)) in DLC1 KD and KO cells versus WT and rescue cells (**Figure 1C** [↗](#)). As previously documented in fibroblasts (Kaushik et al., 2014 [↗](#)), these results indicate that DLC1 feeds into the regulation of contractility, SF and FA formation in fibroblasts. The less penetrant phenotype observed in KO versus KD suggests that the KO cells might be able to adapt to the long-term absence of DLC1, while KD cells had less than 48 hours to adjust to the new mechanical state. These experiments also validate the *DLC1* KO cells and *DLC1* KO rescued with mCherry-DLC1 that will be used in subsequent experiments.

DLC1 deficient cells display altered spreading dynamics, lack of polarization and efficient migration

We timelapsed REF52 WT and *DLC1* KO cells for multiple hours in presence of platelet derived growth factor (PDGF), a potent stimulator of fibroblast motility (Martin et al., 2014 [↗](#)), and evaluated their ability to migrate. We found that *DLC1* KO cells displayed a reduction in both their speed of migration and their ability to migrate directionally (**Figure S1A-C** [↗](#)). We then investigated cytoskeletal dynamics during spreading using REF52 fibroblasts stably expressing the F-actin marker LifeAct (Riedl et al., 2008 [↗](#)). Imaging early spreading (e.g. 20 minutes after plating) revealed no clear differences in edge dynamics in WT versus *DLC1* KO cells (**Figure 2A** [↗](#)). However, kymograph analysis clearly showed that the lamella that characteristically starts directly at the leading edge of the highly contractile REF52 cells (Martin et al., 2016 [↗](#), 2014 [↗](#)), was wider in *DLC1* KO versus WT cells. Later, 1 hour after plating, when the cells are still spreading, lamellipodial protrusion/retraction cycles of wider amplitude were observed in *DLC1* KO versus WT cells (**Figure 2B** [↗](#), Movie S1), with again a large increase in SF content (**Figure 2B** [↗](#), right panel). After initial spreading, REF52 fibroblasts break their symmetry and eventually display short episodes of polarized cell migration. Later, contractility builds up leading cells with spindle-like morphology. This process occurs on timescales of multiple hours. To capture these behaviors, we investigated cytoskeletal dynamics over the next 10 hours after reseeding at a slower timescale that precludes observation of lamellipodial dynamics. While WT cells efficiently broke their symmetry, and could display short stretches of polarized migration, *DLC1* KO cells displayed robust stress fibers that hampered symmetry breaking and polarization (**Figure 2C** [↗](#), Movie S2). *DLC1* KO cells therefore rapidly adopted a spindle-like shape. Quantification of the adoption of such a spindle shape phenotype showed that WT cells will assume the spindle-like shape much later than *DLC1* KO cells and often fail to do so entirely within the 10 hours of imaging (**Figure 2D** [↗](#)). Further highlighting a role for DLC1 in RhoA-mediated control of contractility, we found that DLC1 overexpression leads to strong reduction in the number of SFs (**Figure S2A** [↗](#)). Together, these results show that DLC1 loss of function leads to increased contractility at the onset of spreading, which later impedes polarization and directional cell migration.

DLC1 deficient cells display an increase in global RhoA activity, without any marked differences in RhoA activity pattern

To explore how DLC1 spatio-temporally controls RhoA activity, we used RhoA2G (Fritz et al., 2013 [↗](#)), a fluorescence resonance energy transfer (FRET)-based biosensor that reports on RhoA activity together with the Lifeact F-actin marker during spreading. As previously observed (Martin et al., 2014 [↗](#)), a RhoA activity band that correlates with the lamellar contractile actomyosin




Figure 1.

Cytoskeletal structures in WT and *DLC1* deficient cells

A,B. Representative immunofluorescence images of paxillin and F-actin (phalloidin) (**A**), and pMLC and F-actin (phalloidin) (**B**) immunostains are shown in inverted black and white (ibw) contrast as well as color composites (DAPI signal also included). Scale bar = 10μm.

C. Compared to WT cells (n=243) lamella size is significantly increased in *DLC1* KD cells (n=290, p-adjusted < 0.001) and *DLC1* KO cells (n=242, p-adjusted < 0.001), while *DLC1* rescue cells (n=57) show significantly lower lamella sizes than both, WT (p-adjusted = 0.0098), as well as KO cells (p-adjusted < 0.001).

D. *DLC1* KD cells (n=290) display a significant increase in the total area of focal adhesions per cell (p-adjusted < 0.001) compared to WT cells (n=243), while *DLC1* KO cells (n=242) do not show a significant difference (p-adjusted = 0.09). Rescue cells show a significant decrease in the total area of focal adhesions (p-adjusted < 0.001). ANOVA plus Tukey's honestly significant difference test.




Figure 2.

F-actin dynamics and cell morphodynamics in WT and *DLC1* KO cells

All time series are representative ibw contrast images of WT and *DLC1* KO cells expressing Lifeact-mCherry. Dotted lines mark the area used for the kymographs shown in the right panels.

A. Cells were imaged for 10 minutes in 10 second intervals during spreading immediately after replating. Dotted red lines display differences in lamella size in WT and *DLC1* KO cells. Time

B. One hour after reseeding, when cells are still isotropically spreading, *DLC1* KO cells display increased lamellipodia size and more prominent edge protrusion retraction cycles. High magnification insets are shown for time point 10 minutes and show the robust increase of contractility in *DLC1* KO cells.

C. Cells were imaged for 10 hours with 15 minutes intervals starting from spreading. WT cells break symmetry and display episodes of polarized motility. KO cells are much less dynamic and reach a contractile phenotype much faster, without being able to polarize.

D. Box plot showing at which time point cells exhibited the elongated phenotype described in (C). *DLC1* KO cells (n=42) exhibited this phenotype a mean of 90 minutes earlier than WT cells (n=22) p-adjusted = 0.002, Student's t-test. 17 additional WT cells failed to develop that phenotype entirely.

Scale bars = 20 μ m for all images.

network of the cell can be observed at the cell edge of WT REF52 fibroblast (**Figure 3A** [↗](#), Movie S3). In *DLC1* KO cells, we observed global elevation of RhoA activity throughout the cell. Quantification of both the average activity per cell and the local edge RhoA activity revealed increased RhoA activity (**Figure 3B** [↗](#)). Kymograph analysis revealed that the lamellar RhoA activity band remained of constant size during spreading (**Figure 3A** [↗](#)). Examining cells approximately 10 hours after spreading when they had fully spread, and WT cells had adopted a more contractile state, the *DLC1* KO cells still exhibited a global increase in RhoA activity (**Figure 3C,D** [↗](#)). This increase was however less strong than in spreading cells, most likely because of a higher state of contractility in WT well-spread versus spreading cells. We also evaluated the effect of overexpression of an mCherry-tagged *DLC1* on RhoA activity (**Figure S2B** [↗](#)). We observed that mild *DLC1* expression, that remains mostly bound to FAs and to the PM and does not compromise cell morphology, leads to a decrease in global RhoA activity. Strong *DLC1* expression that leads to a large cytosolic pool of *DLC1* further diminishes global RhoA activity but then compromises cell morphology. Together, these results suggest that *DLC1* signaling at the PM and at FAs controls global levels of RhoA activity within cells, rather than controlling the spatio-temporal RhoA activity pool observed at the lamella.

Design of an optogenetic actuator - Rho biosensor to interrogate the Rho GTPase flux

While FRET measurements as shown above provide insight about a steady-state RhoA activity pattern, they cannot probe in detail how a GEF/GAP/GTPase signaling network might affect the Rho GTPase flux which most likely emerges from the regulation of multiple GEFs and GAPs. For example, GAP-mediated negative regulation might control Rho GTPase signal duration, but might also be involved in adjusting the rate of activation. We reasoned that transient, acute optogenetic perturbations of the Rho GTPase flux, followed by measuring Rho activity dynamics might provide new insights about its regulation. To manipulate Rho activity in single living cells, we re-engineered an iLID-based optogenetic actuator based on a DH/PH domain of the GEF LARG (Oakes et al., 2017 [↗](#)) (**Figure 4A** [↗](#)). We used a high affinity nano SspB domain for efficient light-dependent recruitment of the LARG GEF domain to the iLID anchor. To better focus optogenetic activation, we also fused the iLID anchor to a stargazin membrane anchor that slows down its diffusion (Natwick and Collins, 2021 [↗](#)). Both the stargazin-iLID anchor and the SspB-LARG domains were separated by a P2A self-cleaving peptide, allowing for equimolar expression of both units from a single operon. We refer to this construct as optoLARG. To interrogate the Rho GTPase flux, we engineered a REF52 line that stably expressed the optogenetic construct as well as a rhotekin-based G protein binding domain (rGBD) that reports on Rho activation (Mahlandt et al., 2021 [↗](#)). The rGBD probe is labelled with a tandem dimeric Tomato (tdTomato) fluorophore that is spectrally orthogonal to optogenetic activation. We used a truncated CMV promoter to warrant low expression level of the rGBD probe to avoid dominant negative effects of the construct through inhibition of Rho signaling. We also expressed a spectrally orthogonal far-red Lifeact-miRFP construct that reports on F-actin dynamics. Because rGBD cannot distinguish between the activity of the RhoA, RhoB and RhoC isoforms, we refer to any rGBD measurement as Rho activity. This is in marked contrast with RhoA2G, the FRET biosensor we used above that is specific for RhoA (Fritz et al., 2013 [↗](#)), but is not spectrally compatible with optoLARG. Note that both biosensors are able to measure active Rho GTPase pools during robust morphogenetic events such as tail retraction (Fritz et al., 2013 [↗](#); Mahlandt et al., 2021 [↗](#)), while rGBD misses leading edge Rho GTPase activity pools widely observed using FRET probes (Machacek et al., 2009 [↗](#); Martin et al., 2016 [↗](#); Pertz et al., 2006 [↗](#)).

We then used a digital micromirror device (DMD) to spatially shine repetitive pulses of blue light delivered at different frequencies on 2 distinct regions of interest (ROIs) at the edge of fibroblasts. When light pulses were delivered at high frequency, we observed robust Rho activation at the edge, which correlated with assembly of contractile F-actin structures and potent edge retraction




Figure 3.

RhoA activation dynamics in WT and *DLC1* KO cells

Cells are stably expressing Lifeact-mCherry, and the RhoA2G FRET sensor. RhoA localization images show RhoA2G localization that is identical to RhoA. RhoA activity images display the computed FRET ratio. Images are color-coded according to the normalized scales shown below the panels.

A. Representative images of WT and *DLC1* KO cells during spreading. Kymographs for the violet lines are shown in the right panels. Note the RhoA activity band maintains constant width during spreading at the periphery in WT cells. Note increased global RhoA activity in *DLC1* KO cells, with maintenance of a similar RhoA band pattern at the cell edge.

B. Box plots of FRET ratio averaged over the whole cell (right panel) or a ROI placed at the cell edge (left panel). This shows that during spreading *DLC1* KO cells ($n = 149$) have an increased total RhoA activity compared to WT cells ($n = 114$, $p = 0.005$). In addition, the FRET ratio at just the cell edge is increased as well ($p = 0.011$).

C. Representative images of WT cells and *DLC1* KO cells that have transitioned in a contractile state 12 hours after plating. No difference in RhoA activity pattern can be observed between WT and *DLC1* KO cells, although the latter still display slightly higher global RhoA activity levels.

D. Box plots of FRET ratio averaged over the whole cell (right panel) or a ROI placed at the cell edge (left panel). This shows that contractile *DLC1* KO cells ($n = 103$) have an increased total RhoA activity ($p < 0.001$) and edge FRET ratio ($p < 0.001$) compared to WT cells ($n = 82$) compared to WT cells ($n = 82$). Students t-test.

(A,B) Scale bars = 20 μm .




Figure 4.

An optogenetic actuator- Rho biosensor circuit to probe Rho GTPase flux

A. Schematics of the optogenetic actuator - Rho biosensor to measure Rho GTPase flux. OptoLarg is based on an iLID system which does not interact with Ssb-LARG in the dark state. The iLID module is anchored to the plasma membrane by a Stargazin anchor that displays slow diffusion, allowing better focussing of optogenetic activation. Upon light exposure ssb-LARG is locally recruited to the plasma membrane, activating Rho. Rho activity is measured by a rGBD effector binding domain.

B. Time-series of REF52 cells locally stimulated with light first in a ROI at the cell top with a high stimulation frequency, and then with a ROI at the cell bottom with a low stimulation frequency. Light pulses have the same intensity. Thick and thin blue thunder symbols represent high and low optogenetic stimulation. Red dotted line is used for the kymograph shown in (C).

C. Kymographs of cells in (B). Light pulse stimulation regimes of top and bottom ROIs are shown in the upper box. Blue dotted boxes indicate the region and length of the stimulation.

Note how intense optogenetic stimulation in the top ROI leads to Rho activity, assembly of contractile F-actin structures, and robust edge retraction. Upon removal of the light input, the Rho activity and F-actin resume, and edge protrusion occurs again. Lower optogenetic stimulation in the bottom ROI leads to much lower Rho activity and F-actin structures, as well as lower edge retraction.

(B,C) Scale bars = 10 μm

(**Figure 4B** [↗](#), kymograph shown in 4C, Movie S4). Removal of the blue light pulses led to a subsequent decrease in Rho activity, and reversion of the edge to a protrusive state. Further, stimulating a ROI at the opposite edge of the cell with a lower light pulse frequency led to a lower Rho activation, and less robust edge retraction. These results indicate that depending on the strength of the light input, optoLARG can locally, reversibly, and quantitatively control Rho activity in single living cells on timescales of seconds, allowing us to acutely manipulate the Rho GTPase flux.

Optogenetic interrogation of the Rho GTPase flux in WT and *DLC1* KO cells

Because *DLC1* might both regulate the Rho GTPase flux both at the PM ([Erlmann et al., 2009](#) [↗](#)) and at FAs ([Haining et al., 2018](#) [↗](#)), we performed a series of experiments using optoLARG to transiently activate RhoA with the expectation of observing different perturbations in Rho activity fluxes in control versus *DLC1* KO cells. For that purpose, we produced WT and *DLC1* KO REF52 lines expressing optoLARG, the tdTomato-rGBD biosensor and mRFP-paxillin as a readout for focal adhesions. We first sought to locally activate RhoA by illuminating large ROIs at edges and evaluate any difference in Rho activity patterns using the spectrally compatible rGBD biosensor in WT versus *DLC1* KO cells. We however were not able to identify a threshold optogenetic light input that would induce different RhoA activity patterns and edge dynamics in WT versus the more contractile *DLC1* KO cells. We observed that the same light input could induce edge retraction in both WT and *DLC1* KO cells in a non-predictable manner. This might be due to the spatial heterogeneity of the mechanical states within cells, as well as small differences in optoLARG and rGBD expression levels in the stable cell lines. To address this caveat, we reasoned that we should activate Rho using optoLARG with a more subtle modality by illuminating only small ROIs and measuring small local fluctuations of Rho activity using rGBD without inducing large changes in mechanical states. Further, to dissect potential regulation of *DLC1* GAP activity by PI(4,5)P₂ at the PM ([Erlmann et al., 2009](#) [↗](#)) versus mechanosensitive interactions at the FA ([Haining et al., 2018](#) [↗](#)), we decided to make measurements on FAs but use ROIs outside of FAs (non-FAs ROIs) to probe the PM. For that purpose, we built an image analysis pipeline to first segment a cell in a field of view and to then segment FAs based on their fluorescent intensity, size and localization at the cell periphery using the paxillin-mRFP signal (**Figure 5A** [↗](#)). We then used these FA segmentations to illuminate small ROIs within FAs or in neighboring non-FA ROIs with one pulse of blue light using the DMD and evaluated rGBD fluorescence fluctuations at this location for a couple of minutes until Rho activity returned to baseline (**Figure 5B** [↗](#)). This optogenetic stimulation had no visible effect on edge retraction, or FA morphodynamics, as observed when stimulating larger ROIs. Quality control of our automated segmentation revealed high versus low paxillin signals in FA versus non-FA ROIs (**Figure 5C** [↗](#)). This experiment was performed on multiple cells, allowing to average out any experimental noise due to small differences in expression levels of optoLARG or rGBD. These data were then averaged for FA and non-FA ROIs. We observed that this pulsed optoLARG light input led to a local transient of Rho activity for approximately 2.5 minutes in WT cells when FAs or the PM were stimulated (**Figure 5D** [↗](#)). The identical optoLARG input led to an increased rate of Rho activation leading to higher amplitude when a FA versus a non-FA ROI was stimulated. Surprisingly, in *DLC1* KO cells, we observed increased Rho activation rate that also led to augmented Rho activity amplitude in comparison with WT cells in both FA and non-FA ROIs. This was also accompanied with faster deactivation kinetics of RhoA. These data suggest that *DLC1* limits the rate of Rho activation rather than signal duration to regulate RhoA activity in spread cells. This can occur both at the PM and at FAs. Further, FAs seem to provide a subcellular environment that favors GEF-mediated activation of Rho activity. The finding that *DLC1* KO also leads to faster deactivation suggests the existence of a negative feedback that points to complex modalities of spatio-temporal regulation that most likely involves additional GEFs/GAPs.




Figure 5.

Rho GTPase activation kinetics in WT versus DLC1 KO cells

A. Computer vision pipeline and experiment: 1) a cell is segmented using the rGBD-dTomato channel. 2) FAs and non-FAs are detected using the paxillin channel (see Material and Methods). ROIs on FAs (blue dots, shown in the bottom left panel) and ROIs on non-FA regions in between (orange dots, shown in the upper left panel) are selected for stimulation. 3) Image of the stimulation pattern (green channel) shows that the calibrated DMD can stimulate the regions with high spatial precision (image is overexposed to show the diffraction pattern of the mirrors).

B. The DMD is used to stimulate ROIs (FAs or non-FAs) with a pulse of blue light (blue line). rGBD signal fluctuations are then measured in the ROIs.

C. Distribution of paxillin-miRFP intensities in the FA and Non-FA ROIs normalized to the mean paxillin-miRFP intensity of the whole cell shows that our segmentation pipelines accurately identifies FA and Non-FA ROIs.

D. Normalized and averaged rGBD fluorescence fluctuations upon ROI optogenetic stimulation. For each stimulated ROIs, the fold change to the baseline (average activity from 0-150 seconds before optogenetic stimulation) is calculated. Median and 99% CI are shown. Regions on top of focal adhesions have a larger fold change in rGBD activity than regions between focal adhesions. DLC1-KO cells have a larger rGBD fold change in the initial time after stimulation (150-200 seconds), but then also fall back down to the baseline quicker. (KO-FA: N=3643, KO-NON-FA: N=3643, CTRL-FA: N = 2144, CTRL-NON-FA:2144. Mean number of regions per cell: ~16.90, Cells CTRL = 321, DLC1-KO=431. The same cell can appear multiple times in the experiment, but with a relaxation time in between and new stimulation regions).

E. The different dynamics described in (D) can be robustly observed in technical replicates of the experiment. In all replicates the rGBD recruitment is higher in FAs vs Non-FAs, and we see the trend of faster accumulation and faster return to baseline of rGBD in DLC1 KO vs CTRL cells.

DLC1 differently interact with FAs depending on their mechanical states

Our finding that we did not observe striking patterns of RhoA activity in the vicinity of FAs in spread cells challenges the idea that mechanical inputs regulate Rho activity in this specific regime of mechanical forces. We speculated that the DLC1/talin system in FAs is only mechanosensitive to stronger mechanical inputs. To evaluate FA-DLC1 interactions, we imaged fibroblasts rescued with mCherry-DLC1 and mRFP-paxillin as a marker for FAs. We observed that DLC1 binds to FAs at the onset of FA assembly, and that the local DLC1 density augmented with the paxillin density (**Figure S3A-C**). During FA disassembly, DLC1 density diminished concomitantly with decreasing paxillin density until the FA was fully disassembled (**Figure S3D-F**).

To characterize force-dependent interactions of FAs with DLC1 in regimes of strong mechanical perturbations, we engineered our mCherry-DLC1 rescue cells to stably express the optoLARG system, as well as the mRFP-paxillin construct. We then used the DMD to activate Rho-mediated contractility locally and transiently in small ROIs containing FAs with a standardized light input. The transient light pulse induced either of the two FA behaviors: light-mediated FA reinforcement as evidenced by a local increase in paxillin density, was followed by 1. FA relaxation (and decrease in paxillin density) upon light removal (**Figure 6A-C**, quantified in **Figure 6D**, Movie S5), or 2. FA rupture (**Figure S4A-C**). This was accompanied by a striking simultaneous decrease in DLC1 intensity at the FA during FA reinforcement, and a reassociation of DLC1 during FA relaxation upon light input removal. If the light input led to FA rupture, then the DLC1 signal gradually decreased along FA reinforcement until the FA ultimately ruptured in an all-or-nothing manner. FAs in the same cell that were not subjected to the light input displayed dynamics as in normal cells, as shown in **Figure 6** (**Figure S4D-F**). These results indicate that DLC1 dynamically dissociates and associates during acute FA reinforcement and relaxation in response to a strong mechanical input.

Discussion

A key question in Rho GTPase biology is how multiple GEFs and GAPs control spatio-temporal Rho GTPase signaling patterns to regulate cytoskeletal dynamics that power morphogenetic processes such as cell migration. The recent finding that many GEFs and GAPs themselves bind cytoskeletal and adhesion structures (Müller et al., 2020) strongly suggests direct feedback from the cytoskeleton to Rho GTPase signaling. Self-organizational properties of cytoskeletal and adhesions structures might therefore contribute to spatio-temporal control of Rho GTPase signaling. Because DLC1 has been proposed to bind to FAs in a mechanosensitive manner (Haining et al., 2018), we decided to study it as a prototypical example of an adhesion feedback to Rho GTPase signaling. Using novel technologies that allow for acute spatio-temporal perturbations, we provide some new insights about how Rho GTPases are spatio-temporally controlled by a RhoGAP that integrates mechanosensitive inputs.

We found that in spreading fibroblasts, *DLC1* KO globally augments contractility, as illustrated by a robust increase in SFs and FAs (**Figure 1**). Evaluation of F-actin dynamics immediately after spreading, when global contractility levels are still low, reveal that *DLC1* KO fibroblasts rapidly assemble a wider contractile lamella than WT cells, suggesting aberrant increase of myosin-based contractility already at an early stage of spreading (**Figure 2A**). One hour after cell plating, when cells are well spread but still retain an isotropic shape, we observed a global increase in contractility in *DLC1* KO versus WT cells (**Figure 3B**). On a timescale of multiple hours after spreading, when WT cells break symmetry, display polarized cell migration episodes, and finally adopt a contractile state leading to a spindle morphology, we find that *DLC1* KO cells transition directly to the contractile phenotype without being able to polarize (**Figure 2C,D**). Thus, DLC1 loss of function leads to a global increase of aberrant contractility.




Figure 6.

Optogenetic control of force-dependent DLC1 interactions with FAs

A. Color-coded fluorescence micrographs of REF52 fibroblast expressing mRFP-paxillin (left) and mCherry-DLC1 (right) and the optoLARG construct (not shown). The black boxes indicate the area used for close up images in (B). The white boxes indicate the ROI for optogenetic illumination. Selected FAs denoted by the pink and green arrowheads. The black dotted lines were used for the kymograph in B.

B. Kymographs showing two selected FAs, the grey box indicates the time at which optogenetic stimulation has been applied. Optogenetic stimulation is applied on select ROIs placed over FAs with 50 ms light pulses per frame (every 15 seconds) for a duration of 7.5 minutes.

C. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the respective arrowheads. Scale bars = 10 μ m.

D. Quantification of mcherry-DLC1 and mRFP-paxillin fluorescence signals during optoLARG mediated control of FA reinforcement and relaxation. Normalized mRFP-paxillin and mCherry-DLC1 from 2 FAs shown in panels A-C.

E. Model of Rho GTPase activity modulation by DLC1 at FAs and at the plasma membrane relevant to **Figure 5**. Left and right panels show schematics of Rho activation dynamics in response to optoLARG optogenetic input at FAs (left panel) and plasma membrane (right panels). Top and bottom panel show schematics for Rho activation dynamics in response to optoLARG optogenetic input in control (top) and DLC1 KO (bottom) panels.

F. Model of force dependent regulation of DLC1 at FAs relevant to this and **Figure S3**. Left panel, in absence of acute mechanical input, DLC1 increases with FA assembly and decreases with FA disassembly. Central panel, upon acute local increase of mechanical stress in response to application of an optoLARG optogenetic input, DLC1 unbinds from FA in a reinforcement regime, and rebinds FA in a relaxing regime when the optoLARG input is removed. Right panel, upon acute local increase of mechanical stress in response to application of an optoLARG optogenetic input some FAs rupture after DLC1 dissociation and FA rupture.

We had previously shown that REF52 fibroblasts display a wide band of RhoA activity that correlates with the lamellar contractile myosin network during spreading (Martin et al., 2016 [DOI](#), 2014 [DOI](#)). We found that *DLC1* KO leads to a global increase in RhoA activity throughout the cell, without however markedly modifying the spatial band of RhoA activity (**Figure 3A,B** [DOI](#)). Similar results were observed when cells were allowed to further polarize and assemble a robust actomyosin cytoskeleton (**Figure 3C,D** [DOI](#)). Further, we observed that mild *DLC1* expression, that can still mostly be titrated by FAs and the PM without spilling out into the cytosol, is already able to strongly downregulate global RhoA activity levels (**Figure S2B** [DOI](#)). We propose that *DLC1* regulates global levels of RhoA activity at steady state, rather than spatially patterned RhoA activity pools as observed in protrusive edges and filopodia (Fritz et al., 2013 [DOI](#); Machacek et al., 2009 [DOI](#); Pertz et al., 2006 [DOI](#)), in the lamella (Martin et al., 2016 [DOI](#), 2014 [DOI](#)) or at the cytokinetic furrow (Basant and Glotzer, 2018 [DOI](#); Bement et al., 2015b [DOI](#)). Note that the classic genetic perturbation paradigm cannot be used to study more subtle functions of *DLC1* at FAs because long-term loss of *DLC1* function results in cells being locked in a state of excessive contractility.

How does *DLC1* contribute to the Rho GTPase flux in cells? To answer this question, we used optoLARG to interrogate the Rho GTPase flux at the PM and at FAs, 2 sites at which *DLC1* can control Rho (Erlmann et al., 2009 [DOI](#); Haining et al., 2018 [DOI](#)). We 1st found that a transient optoLARG light input of identical amplitude at FAs led to increased Rho activity amplitude compared to stimulation in non-FA PM regions (schematized in **Figure 6E** [DOI](#)). This strongly suggest that the FA is a subcellular region which is highly permissive for GEF-mediated activation of Rho, which could involve a number of mechanisms including Src regulation of Rho release from RhoGDI (DerMardirossian et al., 2006 [DOI](#)). 2nd, surprisingly, we observed that *DLC1* KO cells displayed increased rates of Rho activation kinetics in both FA and non-FA regions in response to a transient optoLARG light input, also leading to increased Rho activity amplitude. This strongly suggests that *DLC1* regulates the rate of Rho activation rather than signal duration/termination. Given the promiscuous expression of GEFs and GAPs within a cell type (Moon and Zheng, 2003 [DOI](#); Rossman et al., 2005 [DOI](#)), other Rho GAP(s) might specifically control signal termination, which might by example be important to specify the lamellar pattern of RhoA activity in our REF52 cell system. The role of *DLC1* in controlling the rate of Rho activation remains to be understood. One can however speculate that *DLC1* provides negative feedback to control the rate of Rho activation. Such feedback structures are prevalent in different GTPase signaling networks (Tsyganov et al., 2012 [DOI](#); Wu and Lew, 2013 [DOI](#)), and can endow the latter with properties such as robustness to perturbations. 3rd, we observed that in addition to the increased rate of Rho activation in response to the optoLARG input, *DLC1* KO cells also display faster adaptation back to baseline Rho activity compared to control cells. Such non-linear behavior again suggests the existence of additional feedback regulation that might involve other GEFs/GAPs. An important feature of our optogenetic approach in which we perturb cells minimally without eliciting cytoskeletal responses is that we might be able to capture subtle dynamic processes occurring at steady state. Strong optoLARG input that leads to robust actomyosin contractility and edge retraction, as seen in **Figure 4** [DOI](#), might activate numerous mechanical feedback not present at steady-state, and thus not provide access to cellular regulation processes occurring in an unperturbed, normal state.

Together, our data strongly suggests that in spread fibroblasts at steady state, in absence of any acute mechanical input, *DLC1* can exert its GAP activity both at the PM and at FAs to control the rate of Rho activation. In these conditions, *DLC1* association with FAs follows FA assembly and disassembly (**Figure 6** [DOI](#)), strongly suggesting that mechanosensitive interactions with talin are not relevant in this context (Haining et al., 2018 [DOI](#)). Note that RhoA localization is homogeneously distributed at the PM, and any RhoA pool being regulated by *DLC1* at the FAs will immediately diffuse in the PM, explaining the absence of local pools of RhoA activity at FAs. This global regulation of RhoA activity levels by *DLC1* in fibroblasts is in marked contrast with neuronal growth cones, in which *DLC1* regulates a highly focused RhoA activity pattern that localizes to the tip of F-actin bundles in filopodia, most probably to regulate formin-dependent actin polymerization (Fusco et al., 2016 [DOI](#)). In this context, *DLC1* loss of function leads to an

enlargement of the RhoA activity pattern at filopodia, which results in longer and stabler filopodia, ultimately inducing more processive growth cone motility and neurite outgrowth. This shows that a specific RhoGAP, such as DLC1, can function in different ways in distinct cellular systems and mechanical contexts.

In the presence of acute mechanical perturbations, we observe distinct DLC1 dynamics at FAs that most likely involve documented mechanosensitive interactions with talin at FAs (Haining et al., 2018). Using optoLARG, we were able to control actomyosin contractility transiently and locally in a small ROI leading to FA reinforcement followed by relaxation (Figure 6). We found that in response to such a mechanical input, DLC1 rapidly dissociates from FAs that exhibit robust reinforcement (Figure 6, schematized in Figure 6F). The release of DLC1 in response to strong FA reinforcement correlates well with *in vitro* measurements that have shown that DLC1 dissociates from stretched talin due to unfolding of the r7-r8 domain (Haining et al., 2018). Our data also shows that FA relaxation can lead to rebinding of DLC1 to FAs (Figure 6), suggesting that the r7-r8 talin domain might refold in these conditions. Our results illustrate how optoLARG enables spatio-temporal, reversible control of contractility that might help unravel mechanosensitive interactions that are not accessible with classically used global chemical perturbations such as nocodazole-mediated microtubule depolymerization (Krendel et al., 2002), or treatment with lysophosphatidic acid (Pertz et al., 2006) leading to global RhoA activation and massive stimulation of contractility.

We propose that DLC1 provides two levels of regulation of Rho. In absence of any strong mechanical input, active DLC1 at FAs and at the PM throughout the cell seems to contribute to the regulation of steady-state global Rho activity in the cells by controlling its rate of activation. The significance of this mode of regulation is yet to be understood but most likely involves simultaneous regulation with additional Rho GEFs and GAPs. Distinct Rho GAPs might spatially control signal termination that shape the characteristic band of RhoA activity that coincides with the lamella. In response to strong mechanical stimuli leading to FA reinforcement and talin stretching, rapid unbinding of DLC1 from FAs might locally decrease local RhoGAP activity providing positive feedback that could rapidly increase the local rate of Rho activation, possibly strengthening contractility that might lead to FA disassembly as we have observed. Conversely, when the mechanical input vanishes, recruitment of DLC1 to relaxing FAs might rapidly switch off RhoA activity to return to a steady state of low contractility. We speculate that these non-linear signaling behaviors might allow to amplify local mechanical inputs to regulate all-or-nothing cytoskeletal processes such as tail retraction during cell migration. Such tail retraction events are characterized by short bursts of highly localized RhoA activity that are followed by rapid relaxation once the tail has been retracted (Martin et al., 2014; Pertz et al., 2006). Here, SF pulling on FAs at the back of a cell might allow local release of DLC1, rapid increase of local RhoA activity, leading to strong local contraction and tail retraction. Experimentally testing this hypothesis in the context of tail retraction is technically challenging because long term DLC1 perturbation leads to global contractility that precludes cell polarization, directional migration and thus tail retraction episodes. This illustrates an important limitation of the classic genetic perturbation toolkit that has been used to study Rho GTPase signaling.

Our results provide new insights into the complex network circuitry that regulates spatio-temporal Rho GTPase signaling and illustrate how we must tackle this problem. By allowing for precise and reversible perturbations, signaling optogenetics provide the opportunity to dissect the spatio-temporal Rho GTPase fluxes that control patterns of Rho GTPase activity at adequate time/length scales. In the future, a full understanding of Rho GTPase signaling fluxes and formation of signaling patterns will require more systematic approaches to consider the multiple GEFs/GAPs that are involved in shaping cell morphogenesis.

Materials and methods

Cell culture

Rat Embryo Fibroblast 52 (REF52) were cultured in Dubelcco's Modified Eagle Medium with 4.5 g/l glucose, 10% 4mM L-Glutamine, and 100 U/ml penicillin/streptomycin. Cells were grown at 37 °C and 5% CO₂.

SiRNA knockdown

4μl lipofectamine RNAi max was mixed with a total of 500μl of OPTIMEM medium and a pool of SiRNAs against the gene of interest at a final concentration of 60 nM (SiTools Biotech). The mix was added to a 6 well plate containing 100000 cells in a 2 ml medium volume medium.

CrispR/Cas9 Knockout generation

Two guide sequences targeting DNA within exon 5 of the DLC1 gene were selected using the CRISPOR tool for predicted high-specificity protospacer adjacent motif target sites in the rat genome³². Two complementary oligos each containing the DLC1 guide sequence and BbsI ligation adapters were synthesized (Microsynth) (sense sequence: 5'-CACCGAACCGAGAGAGCTACCCGG-3', antisense sequence: 5'-AAACCCGGGTAGCTCTCTCGGTTC-3'). The guide sequences were annealed and ligated into a pSpCas9(BB)-2A-GFP vector. REF52 cells were grown in six-well plates to 60% confluency and transfected with 1μg of the vector together with 1 μl of Lipofectamine 3000 and 5 μl of P3000 solution (ThermoFisher Scientific). Two days post transfection cells were detached, suspended in PBS +1% FBS and sorted into 96 well-plates using fluorescence-activated cell sorting. After expansion, individual clones were detached and suspended in the lysis buffer (10mM TRIS, 50mM KCL, 2.5mM MgCl₂, 0.45% Tween-20, 0.05% Gelatine, 0.12 mg/ml of proteinase K). The cell solution was frozen at -80 °C and subsequently kept at 63 °C for one hour. Proteinase K was inactivated by heating the solution to 95 °C for 15 minutes. PCR primers were designed to amplify a 500bp region around the CRISPR cut site (forward primer: 5'-AAGGAGTGTGTCTAACTCCACGCAGACCAG-3', reverse primer: 5'-CTCCTTAGGACTGTCGTGCTGTTTCTCT-3'). Genomic sequences were amplified by PCR and sequenced by Sanger sequencing.

DNA constructs

A "pB3.0" piggybac vector was created by adapting the pPBbsr2 (Matasci et al., 2011 [DOI](#)) to make it smaller in size. pPBbsr2 was digested with AscI and PacI, yielding 3922 bp and 2864 bp fragments. Full size pPBbsr2 was used as template for a PCR with forward primer 5'-TTAGCATTAAATTAAGCGCCGCGTTGCTGGCGTTTTTCC-3' and reverse primer 5'-GTGCCTTTACAACCTATGAGTAACCCGCGCGGACGATT-3', yielding a 1671bp insert that was ligated back into the receiving 3922bp part of the digested pPBbsr2 vector, creating "pB3.0-BLAST" (Blasticidine resistance). To introduce different resistance cassettes, we performed an overlap extension PCR with pB3.0-BLAST as template for PCR1 (primers A+B) and PCR2 (primers C+D) with the following primers: primer A: 5'-AAGGATGCCCAGAAGGTACCCCATTTGATGGGATCTGATCTGGG-3', primer B: 5'-GGAACTTTTTTGCTATTATGGTGCCATTCAGCTCTACGTAGCTACT-3', primer C 5'-CCTTTGAAAAACACGATAATACCACCGGTAAGTCGAGATGCATCGATGA-3', primer D: 5'-CTCCGCCTTTCTTGACGTGCGGTTCGAACCGCATTAGT-3'. Primers A and D, the product of PCR1 and PCR2 were used for fusing the two fragments by PCR. The resulting PCR product was ligated into the pB3.0-BLAST vector, digested with KpnI and PstI, by fusion cloning, yielding a pB3.0-noAB vector with a single AgeI site to introduce antibiotic resistance cassettes. To produce pB3.0-HYGRO and pB3.0-PURO, PCRs were performed on template vectors pHygro-PB and pPuro-PB, respectively. For pB3.0-HYGRO forward primer 5'-AACACGATAATACCACCATGAAAAAGCCTGAACCTACCGC-3' and reverse primer 5'-GCAGGCTCCCGTTTCTTATCGGCCATTTCAGCTCTA-3' were used. For pB3.0-PURO

forward primer 5'-AACACGATAATACCACCATGACCGAGTACAAGCCCACG-3' and reverse primer 5'-GCGTTCGGGCCACGGACTGGCCATTCAGCTCTA-3' were used. The respective inserts were inserted via fusion cloning in the pB3.0-noAB vector³³, digested with AgeI.

Construction of the pB3.0-optoLARG-mVenus-SspB-p2A-stargazin-mtq2-iLID optoLARG system was as follows. The cDNA for optoLARG-mVenus-SspB(nano)-p2A-iLID-CAAX was synthesized by custom gene synthesis (Genewiz, Azenta life sciences, US). The sequence contained the catalytical DH-PH domain of human RhoGEF12 (LARG, aa 766-1138), connected via a linker (GSGSGSGS) to full-length mVenus fused to SSPB(nano). This is followed by a p2A sequence (GSGATNFSLLKQAGDVEENPGP), the iLID module (Guntas et al., 2015 [DOI](#)) and a CAAX box (KRAS). The construct was cloned into pB3.0-BLAST containing a CAG promoter, yielding pB3.0-optoLARG-mVenus-SspB-p2A-iLID-CAAX. After publication of an improved plasma membrane anchor for the iLID-module (Natwick and Collins, 2021 [DOI](#)), we changed the part after the p2A sequence. The pB3.0-optoLARG-mVenus-SspB-p2A-iLID-CAAX vector was digested with BsrGI and AflII, and a custom gene synthesized stretch of cDNA containing SspB(nano)-p2A, full-length stargazin, mTurquoise2 and iLID was inserted. This yielded pB3.0-optoLARG-mVenus-SspB(nano)-p2A-stargazin-mtq2-iLID (exact sequences of the gene synthesis products are available on request).

An eGFP DLC1 plasmid containing the 1091aa long isoform originating from mouse (kindly provided by Monilola A. Olayioye) was used to produce mCherry-DLC1. The DLC1 gene was cut out of this plasmid with BamHI and, N-terminally fused to FKBP12-mCherry and inserted into pB3.0-BLAST containing a CAG-promotor, yielding pB3.0-FKBP12-mCherry-DLC1.

Spreading Assay

Glass bottom well plates (Celvis) were coated with 5 µg/ml of human plasma fibronectin purified protein (Merck) for one hour at room temperature. REF52 cells were seeded at a density of 7000 cells per well (using 24 well plates) in FluoroBrite DMEM medium containing 1% FBS, 0.1% BSA, 4mM L-Glutamine and 100 U/ml penicillin/streptomycin. Imaging was done with a Nikon Eclipse Ti-E inverted microscope with an automatic stage. Temperature was kept at 37°C with a temperature control system, humidity (100%) and CO₂ (~5%) with a gas mixer (Life Imaging Services). Focus drift was prevented by the equipped Perfect Focus System (Nikon). The microscope was controlled with Metamorph software (Universal Imaging).

Optogenetic Stimulation experiments

Glass bottom well plates (Celvis) were coated with 5 µg/ml of human plasma fibronectin purified protein (Merck) for one hour at room temperature. REF52 cells were seeded at 6000 cells/well into 12 well plates and allowed to attach for 12 hours. For both RhoA activity and mCherry-DLC1/miRFP-paxillin optogenetic stimulation experiments cells were incubated in FluoroBrite DMEM medium overnight. The microscope was controlled with NIS-Elements (Nikon) for observing DLC1 dynamics and with open-source micromanager software (Edelstein et al., 2014 [DOI](#)) for edge stimulation. Optogenetic stimulation was localized to specific regions using an Andor mosaic 3 DMD. For mCherry-DLC1/miRFP-paxillin experiments, ROIs were user-defined. For RhoA activity experiments, cells in different fields of view were selected by a user, cells were then segmented with a custom pixel-classifier based on a pre-trained VGG16 convolutional neural network, which was trained on a field of view of the experimental data. ROIs for stimulation masks were then automatically defined at the top and bottom of the cell for consecutive illumination with different light pulse regimes. Fields of views were imaged one after the other on multiple cells. For data extraction the stimulation mask was expanded by gaussian smoothing and then thresholded to cover a wider region, reducing artifacts caused by membrane dynamics. Image intensities were calculated for the overlap of this mask with the segmented cells and the full segmented cells respectively.

Immuno-histochemistry

One hour after seeding, cells were fixed for 10 minutes with 0.2% paraformaldehyde. A 0.1% Triton X solution was used for permeabilization, after which cells were incubated overnight at 4°C with the blocking buffer, containing 5% Bovine-Serum-Albumin and 0.05% Tween20 in PBS. After each step of the following protocol cells were washed three times with Phosphate-Buffered-Saline (PBS). Cells were incubated overnight with the primary antibody (1:250 dilution for paxillin (Abcam ab32084), 1:50 for pMLC (Cell Signaling Antibody #3671). Secondary antibody incubation (1:1000) lasted one hour. Phalloidin incubation (1:200 Phalloidin-Atto647N) was performed for 20 minutes and DAPI incubation for 10 hours. Cells were imaged immediately after the last step.

Image analysis

Lamella size, focal adhesion areas (**Figure 1** [↗](#)) and RhoA activity ratio measurements (**Figure 3** [↗](#)) were computed with CellProfiler in conjunction with a pixel classifier trained with Ilastik ([Berg et al., 2019](#) [↗](#)). Spreading dynamics (**Figure 2D** [↗](#)) were manually analyzed. FRET ratio images were computed and analyzed with custom python scripts. DLC1 and paxillin dynamics (**Figure S3D** [↗](#)) were calculated with manually drawn ROIs in Fiji ([Schindelin et al., 2012](#) [↗](#)).

Acknowledgements

This work was supported by the Swiss National Science Foundation IZSAZ3_173462 Argentinian-Swiss Joint Research Programme and by the Sinergia CRSII5_183550 grants to Olivier Pertz. We are grateful to the Microscopy Imaging Center of the University of Bern for support (<https://www.mic.unibe.ch> [↗](#)). We thank Kazuhiro Aoki (NIBB, Japan) for providing the pPBbsr2 vector, David L. Hacker (EPFL, Switzerland), for the pPuro-PB and pHygro-PB plasmids, Judith Trüb (UniBE, Switzerland) for the cloning of pB3.0-BLAST, Monilola A. Olayioye (University of Stuttgart, Germany) for the eGFP DLC1 plasmid, Dean E. Natwick and Sean R. Collins (University of California Davis) for sharing the Stargazin-iLID constructs. We are grateful to Miguel Vicente-Manzanares, Bernhard Wehrle-Haller and Daniel Riveline for constructive comments on the manuscript

Author contributions

O.P. supervised the research. O.P. and M.H. wrote the paper. M.H. designed and performed experiments. L.H. designed and performed optogenetic experiments. J.v.U. built the optoLARG/rGBD circuit. M.D. designed data analysis tools.

Additional Information

The authors declare no competing interests.

Supplementary Figure legends




Figure S1

Cell motility properties in WT versus *DLC1* KO cells

A. Example fields of view of tracked WT and *DLC1* KO cells. 20000 REF fibroblasts expressing a histone H2B-miRFP marker were seeded in a 24 well-plate well coated with fibronectin and stimulated with 20 ng/ml PDGF, imaged for 12 hours at 15 minutes interval. Tracking was performed using stardist³⁷ on the H2B image). *DLC1* KO cells display a large subpopulation of cells that remain extremely stationary.

B. Mean Square displacements (MSDs) for different time intervals, WT move more than *DLC1* KO cells for all possible lag times. Plot shows directionality (exponential bit between 0.25 and 1.5 hours). For times longer than 2 hours the curve is flat indicating that migration has characteristics of random walk at these timescales. Thick lines: mean, thin lines: standard deviation. Dotted grey line shows delta t chosen for figure C.

C. Distribution of velocities calculated as $\mu m/hour$ from Root-MSD. We again observe the bimodal distribution in the *DLC1* KO cell with a big subgroup of the cells move extremely slowly (5 $\mu m/hour$). Thick lines are mean and extrema. Two sided t-test: (statistic=25.1, p-value=1.6e-131)




Figure S2.

Effect of DLC1 overexpression on F-actin cytoskeleton and RhoA activity

A. Effect of DLC1 overexpression of F-actin. REF52 cells stably expressing Lifeact-mCherry were transfected with mCherry-DLC1 or mCherry plasmids, allowed to spread for 12 hours on fibronectin-coated coverslips and imaged. Images are shown in ibw contrast. Large mCherry-DLC1 pool in the cytosol levels documents its overexpression. Note how DLC1 overexpression leads to loss of contractile F-actin structures.

B. Effect of DLC1 overexpression of RhoA. Cells stably transfected with the RhoA2G biosensor were transfected with mCherry-paxillin or mCherry-DLC1. mCherry signals are shown in ibw contrast. RhoA2G FRET ratio, and expression levels are color-coded according to the scale. Note how low mCherry-DLC1 expression that remains associated with FA already lowers RhoA activity, while high mCherry-DLC1 expression leads to even lower RhoA activity and aberrant cell morphology.




Figure S3.

DLC1 dynamics at FAs in unperturbed cells

Panels **A-C** document FAs in an assembly state. This shows that the DLC1 signal augments concomitantly with paxillin signal during FA assembly. Panels **D-F** document FAs in a disassembly state. This shows that the DLC1 signal decreases concomitantly with paxillin signal during FA disassembly.

A,D. Color-coded fluorescence micrographs of REF52 fibroblast expressing miRFP-paxillin (top) and mCherry-DLC1 (bottom). The white boxes indicate the area used for close up images in (B,E).

B,E. Closeups of the ROIs (left panel) and kymographs (right panel) of selected FAs denoted by the red arrowheads. White dotted lines were used for the kymograph. In the kymograph, the grey box indicates the time at which optogenetic stimulation has been applied.

C,F. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the respective red arrowheads.

Scale bars = 10 μm .




Figure S4.

DLC1 dynamics in an optoLARG stimulated FA that undergoes reinforcement followed by disassembly, as well as FA behavior in absence of optoLARG stimulus

Panels **A-C** document FAs that when subjected to a nearby pulse of optogenetic Rho-mediated contractility display a behavior of FA reinforcement followed by disassembly. Optogenetic stimulation is applied on select ROIs placed over FAs with 50 ms light pulses per frame (every 15 seconds) for a duration of 8 minutes.

Panels **D-F** document the dynamics of DLC1 in a ROI not stimulated with light in the same cell as in shown in **Figure 6A-C**.

A. Color-coded fluorescence micrographs of REF52 fibroblast expressing miRFP-paxillin (top) and mCherry-DLC1 (bottom) and the optoLARG construct (not shown). The black boxes indicate the area used for close up images in (B). The white boxes indicate the ROI for optogenetic illumination.

B. Closeups of the ROIs (left panel) and kymographs (right panel) of selected FAs denoted by the arrowheads. The pink and green arrowheads indicate the FAs of interest in (B). The black arrowhead indicates the FA of interest in (C). The black dotted lines were used for the kymograph. In the kymograph, the grey box indicates the time at which optogenetic stimulation has been applied.

C. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the respective arrowheads. Scale bars = 10 μ m.

D. Color-coded fluorescence micrographs of REF52 fibroblast expressing miRFP-paxillin (top) and mCherry-DLC1 (bottom) and the optoLARG construct (not shown). The black boxes indicate the area used for close up images in (B,E). The white boxes indicate the ROI for optogenetic illumination.

E. Closeups of the ROIs (left panel) and kymographs (right panel) of selected FAs denoted by the arrowheads. The pink and green arrowheads indicate the FAs of interest in (B). The black arrowhead indicates the FA of interest in (D). The black dotted lines were used for the kymograph. In the kymograph, the grey box indicates the time at which optogenetic stimulation has been applied.

F. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the respective arrowheads.

Supplementary Movie legends

Movie S1: F-actin dynamics of WT and DLC1 KO REF52 cells during late spreading (relevant to Figure 2B [↗](#)).

Representative ibw contrast movie of WT and *DLC1* KO cells expressing Lifeact-mCherry during late spreading. Time: Minutes:seconds. Scale bar = 20 mm.

Movie S2: F-actin dynamics and edge dynamics of WT and DLC1 KO REF52 cells during acquisition of a polarized cell migration phenotype over a period of 15 hours (relevant to Figure 2C [↗](#)).

Representative ibw contrast movie of WT and *DLC1* KO cells expressing Lifeact-mCherry for 15. Edge dynamics were color-coded with respect to time. Time: Minutes:seconds. Scale bar = 20 mm.

Movie S3: F-actin and RhoA activity dynamics WT and DLC1 KO REF52 cells during late spreading (relevant to Figure 3A [↗](#)).

Representative movie of WT and *DLC1* KO cells expressing Lifeact-mCherry and the RhoA2G FRET biosensor during late spreading.

Left panels: F-actin channel in ibw contrast. Right panels: RhoA activity emission ratio color coded according to color scale.

Time: Minutes:seconds. Scale bar = 20 mm.

Movie S4: Characterisation of the optoLARG/rGBD genetic circuit (relati to Figure 5 [↗](#))

REF52 cells stably expressing the optoLARG/rGBD circuit as well as Lifeact-miRFP are locally stimulated with blue light of identical intensity, first in a ROI at the cell top with a high stimulation frequency, and then with a ROI at the cell bottom with a lower stimulation frequency. The stimulation frequencies are indicated by the appearance/disappearance frequency of the ROIs.

Left panel: rGBD signal, Right panel: F-actin lifeact-miRFP. Note the more robust edge contraction, as well as increased RhoA activity in response to high versus low light frequency stimulation.

Time: Minutes:seconds. Scale bar = 10 mm.

Movie S5: DLC1 dynamics at FAs during an optoLARG-induced contractility pulse (relevant to Figure 6A-C [↗](#), cell denoted by pink arrow)

REF52 KO cells rescued with mCherry-DLC1 and expressing mRFP-paxillin, and the optoLARG/rGBD circuit were imaged before, during and after a transient optogenetic stimulation in the whole field of view shown in the movie.

Time: Minutes:seconds. Scale bar = 5 mm.

References

- Basant A, Glotzer M (2018) **Spatiotemporal Regulation of RhoA during Cytokinesis** *Curr Biol* **28**:R570–R580
- Bement WM *et al.* (2015) **Activator-inhibitor coupling between Rho signalling and actin assembly makes the cell cortex an excitable medium** *Nat Cell Biol* **17**:1471–1483
- Bement WM *et al.* (2015) **Activator-inhibitor coupling between Rho signalling and actin assembly makes the cell cortex an excitable medium** *Nat Cell Biol* **17**:1471–1483
- Bement WM, von Dassow G. (2014) **Single cell pattern formation and transient cytoskeletal arrays** *Curr Opin Cell Biol* **26**:51–59
- Berg S *et al.* (2019) **. ilastik: interactive machine learning for (bio) image analysis** *Nat Methods* **16**:1–7
- Blum Y, Mikelson J, Dobrzyński M, Ryu H, Jacques M-A, Jeon NL, Khammash M, Pertz O (2019) **Temporal perturbation of ERK dynamics reveals network architecture of FGF2/MAPK signaling** *Mol Syst Biol* **15** <https://doi.org/10.15252/msb.20198947>
- DerMardirossian C, Rocklin G, Seo J-Y, Bokoch GM (2006) **Phosphorylation of RhoGDI by Src regulates Rho GTPase binding and cytosol-membrane cycling** *Mol Biol Cell* **17**:4760–4768 <https://doi.org/10.1091/mbc.e06-06-0533>
- Dessauges C, Mikelson J, Dobrzyński M, Jacques M, Frismantiene A, Gagliardi PA, Khammash M, Pertz O (2022) **Optogenetic actuator – ERK biosensor circuits identify MAPK network nodes that shape ERK dynamics** *Mol Syst Biol* **18** <https://doi.org/10.15252/msb.202110670>
- Edelstein AD, Tsuchida MA, Amodaj N, Pinkard H, Vale RD, Stuurman N (2014) **Advanced methods of microscope control using µManager software** *J Biol Methods* **1** <https://doi.org/10.14440/jbm.2014.36>
- Erlmann P, Schmid S, Horenkamp FA, Geyer M, Pomorski TG, Olayioye MA (2009) **DLC1 activation requires lipid interaction through a polybasic region preceding the RhoGAP domain** *Mol Biol Cell* **20**:4400–4411 <https://doi.org/10.1091/mbc.e09-03-0247>
- Etienne-Manneville S, Hall A (2002) **Rho GTPases in cell biology** *Nature* **420**:629–635
- Fritz RD *et al.* (2013) **A versatile toolkit to produce sensitive FRET biosensors to visualize signaling in time and space** *Sci Signal* **6** <https://doi.org/10.1126/scisignal.2004135>
- Fritz RD, Pertz O (2016) **The dynamics of spatio-temporal Rho GTPase signaling: formation of signaling patterns** *F1000Research* **5** <https://doi.org/10.12688/f1000research.7370.1>
- Fusco L, Lefort R, Smith K, Benmansour F, Gonzalez G, Barillari C, Rinn B, Fleuret F, Fua P, Pertz O (2016) **Computer vision profiling of neurite outgrowth dynamics reveals spatiotemporal modularity of Rho GTPase signaling** *J Cell Biol* **212**:91–111 <https://doi.org/10.1083/jcb.201506018>

Gingras AR *et al.* (2008) **The structure of the C-terminal actin-binding domain of talin** *EMBO J* **27**:458–469 <https://doi.org/10.1038/sj.emboj.7601965>

Goult BT, Brown NH, Schwartz MA (2021) **Talin in mechanotransduction and mechanomemory at a glance** *J Cell Sci* **134** <https://doi.org/10.1242/jcs.258749>

Graessl M *et al.* (2017) **An excitable Rho GTPase signaling network generates dynamic subcellular contraction patterns** *J Cell Biol* **17**:201706052–15

Guntas G, Hallett RA, Zimmerman SP, Williams T, Yumerefendi H, Bear JE, Kuhlman B (2015) **Engineering an improved light-induced dimer (iLID) for controlling the localization and activity of signaling proteins** *Proc Natl Acad Sci U S A* **112**:112–117

Haining AWM, Rahikainen R, Cortes E, Lachowski D, Rice A, von Essen M, Hytönen VP, del Río Hernández A (2018) **Mechanotransduction in talin through the interaction of the R8 domain with DLC1** *PLOS Biol* **16**:e2005599–20

Hu J, Gong X, Strömblad S (2022) **Local temporal Rac1-GTP nadirs and peaks restrict cell protrusions and retractions** *Sci Adv* **8** <https://doi.org/10.1126/sciadv.abl3667>

Isogai T, Danuser G (2018) **Discovery of functional interactions among actin regulators by analysis of image fluctuations in an unperturbed motile cell system** *Philos Trans R Soc B Biol Sci* **373** <https://doi.org/10.1098/rstb.2017.0110>

Itoh RE, Kurokawa K, Ohba Y, Yoshizaki H, Mochizuki N, Matsuda M (2002) **Activation of Rac and Cdc42 Video Imaged by Fluorescent Resonance Energy Transfer-Based Single-Molecule Probes in the Membrane of Living Cells** *Mol Cell Biol* **22**:6582–6591 <https://doi.org/10.1128/MCB.22.18.6582-6591.2002>

Kaushik S, Ravi A, Hameed FM, Low BC (2014) **Concerted modulation of paxillin dynamics at focal adhesions by deleted in liver cancer-1 and focal adhesion kinase during early cell spreading: DLC1 Regulates Paxillin Dynamics at Focal Adhesions via FAK** *Cytoskeleton* **71**:677–694 <https://doi.org/10.1002/cm.21201>

Kraynov VS, Chamberlain C, Bokoch GM, Schwartz MA, Slabaugh S, Hahn KM (2000) **Localized Rac activation dynamics visualized in living cells** *Science* **290**:333–337

Krendel M, Zenke FT, Bokoch GM (2002) **Nucleotide exchange factor GEF-H1 mediates cross-talk between microtubules and the actin cytoskeleton** *Nat Cell Biol* **4**:294–301 <https://doi.org/10.1038/ncb773>

Li G, Du X, Vass WC, Papageorge AG, Lowy DR, Qian X. (2011) **Full activity of the deleted in liver cancer 1 (DLC1) tumor suppressor depends on an LD-like motif that binds talin and focal adhesion kinase (FAK)** *Proc Natl Acad Sci* **108**:17129–17134 <https://doi.org/10.1073/pnas.1112122108>

Machacek M, Hodgson L, Welch C, Elliott H, Pertz O, Nalbant P, Abell A, Johnson GL, Hahn KM, Danuser G (2009) **Coordination of Rho GTPase activities during cell protrusion** *Nature* **461**:99–103 <https://doi.org/10.1038/nature08242>

Mahlandt EK, Arts JGG, van der Meer WJ, van der Linden FH, Tol S, van Buul JD, Gadella TWJ, Goedhart J. (2021) **Visualizing endogenous Rho activity with an improved localization-based, genetically encoded biosensor** *J Cell Sci* **134** <https://doi.org/10.1242/jcs.258823>

- Martin K, Reimann A, Fritz RD, Ryu H, Jeon NL, Pertz O (2016) **Spatio-temporal co-ordination of RhoA, Rac1 and Cdc42 activation during prototypical edge protrusion and retraction dynamics** *Sci Rep* **6** <https://doi.org/10.1038/srep21901>
- Martin K, Vilela M, Jeon NL, Danuser G, Pertz O (2014) **A growth factor-induced, spatially organizing cytoskeletal module enables rapid and persistent fibroblast migration** *Dev Cell* **30**:701–716 <https://doi.org/10.1016/j.devcel.2014.07.022>
- Matasci M, Baldi L, Hacker DL, Wurm FM (2011) **The PiggyBac transposon enhances the frequency of CHO stable cell line generation and yields recombinant lines with superior productivity and stability** *Biotechnol Bioeng* **108**:2141–2150 <https://doi.org/10.1002/bit.23167>
- Moon SY, Zheng Y (2003) **Rho GTPase-activating proteins in cell regulation** *Trends Cell Biol* **13**:13–22 [https://doi.org/10.1016/s0962-8924\(02\)00004-1](https://doi.org/10.1016/s0962-8924(02)00004-1)
- Mosaddeghzadeh N, Ahmadian MR (2021) **The RHO Family GTPases: Mechanisms of Regulation and Signaling** *Cells* **10** <https://doi.org/10.3390/cells10071831>
- Müller PM *et al.* (2020) **Systems analysis of RhoGEF and RhoGAP regulatory proteins reveals spatially organized RAC1 signalling from integrin adhesions** *Nat Cell Biol* **22**:498–511 <https://doi.org/10.1038/s41556-020-0488-x>
- Nalbant P, Hodgson L, Kraynov V, Touthkine A, Hahn KM (2004) **Activation of endogenous Cdc42 visualized in living cells** *Science* **305**:1615–1619
- Natwick DE, Collins SR (2021) **Optimized iLID Membrane Anchors for Local Optogenetic Protein Recruitment** *ACS Synth Biol* **10**:1009–1023 <https://doi.org/10.1021/acssynbio.0c00511>
- Oakes PW, Wagner E, Brand CA, Probst D, Linke M, Schwarz US, Glotzer M, Gardel ML (2017) **Optogenetic control of RhoA reveals zyxin-mediated elasticity of stress fibres** *Nat Commun* **8**:15817–12
- Pertz O (2010) **Spatio-temporal Rho GTPase signaling - where are we now?** *J Cell Sci* **123**:1841–1850 <https://doi.org/10.1242/jcs.064345>
- Pertz O, Hodgson L, Klemke RL, Hahn KM (2006) **Spatiotemporal dynamics of RhoA activity in migrating cells** *Nature* **440**:1069–1072 <https://doi.org/10.1038/nature04665>
- Riedl J *et al.* (2008) **Lifeact: a versatile marker to visualize F-actin** *Nat Methods* **5**:605–607 <https://doi.org/10.1038/nmeth.1220>
- Rossman KL, Der CJ, Sondek J. (2005) **GEF means go: turning on RHO GTPases with guanine nucleotide-exchange factors** *Nat Rev Mol Cell Biol* **6**:167–180
- Ryu H, Chung M, Dobrzyński M, Fey D, Blum Y, Lee SS, Peter M, Kholodenko BN, Jeon NL, Pertz O (2015) **Frequency modulation of ERK activation dynamics rewires cell fate** *Mol Syst Biol* **11** <https://doi.org/10.15252/msb.20156458>
- Schindelin J *et al.* (2012) **Fiji: an open-source platform for biological-image analysis** *Nat Methods* **9**:676–682 <https://doi.org/10.1038/nmeth.2019>
- Schmick M, Kraemer A, Bastiaens PIH (2015) **Ras moves to stay in place** *Trends Cell Biol* **25**:190–197

Tsyganov MA, Kolch W, Kholodenko BN (2012) **The topology design principles that determine the spatiotemporal dynamics of G-protein cascades** *Mol Biosyst* **8**:730–743

Wu C-F, Lew DJ (2013) **Beyond Symmetry Breaking: Competition and Negative Feedback in GTPase regulation** *Trends Cell Biol* **23**:476–483 <https://doi.org/10.1016/j.tcb.2013.05.003>

Zhao Y, Lykov N, Tzeng C (2022) **Talin-1 interaction network in cellular mechanotransduction (Review)** *Int J Mol Med* **49**:1–12 <https://doi.org/10.3892/ijmm.2022.5116>

Article and author information

Max Heydasch

Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland

Lucien Hinderling

Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland, Graduate School for Cellular and Biomedical Sciences, University of Bern, Switzerland
ORCID iD: [0000-0002-3956-9363](https://orcid.org/0000-0002-3956-9363)

Jakobus van Unen

Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland
ORCID iD: [0000-0002-7931-0989](https://orcid.org/0000-0002-7931-0989)

Maciej Dobrzynski

Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland
ORCID iD: [0000-0003-0757-4712](https://orcid.org/0000-0003-0757-4712)

Olivier Pertz

Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland
For correspondence: olivier.pertz@unibe.ch
ORCID iD: [0000-0001-8579-4919](https://orcid.org/0000-0001-8579-4919)

Copyright

© 2023, Heydasch et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Patricia Bassereau
Institut Curie, France

Senior Editor

Benoît Kornmann
University of Oxford, United Kingdom

Joint Public Review:

Summary:

The manuscript of Heydasch et al. addresses the spatiotemporal regulation of Rho GTPase signaling in living cells and its coupling to the mechanical state of the cell. They focus on a GAP of RhoA, the Rho-specific GAP Deleted in Liver Cancer 1 (DLC1). They first show that removing DLC1 either by a CRISPR KO or by downregulation using siRNA leads to increased contractility and globally elevated RhoA activity, as revealed by a FRET biosensor. This result was expected, since DLC1 is deactivating RhoA its absence should lead to increasing amounts of active RhoA. To go beyond global and steady levels of RhoA activity, the authors developed an acute optogenetic system to study transient RhoA activity dynamics in different genetic and subcellular contexts. In WT cells, they found that pulses of activation lead to an increased RhoA activity at focal adhesions (FA) compared to plasma membrane (PM), which suggests that FAs contain less RhoA GAPs, more RhoA, or that FAs involve positive feedback implying other GEFs for example. In DLC1 KO cells, they found that the RhoA response upon pulses of optogenetic activation was increased (higher peak) both at FA and PM, which could be expected since less GAP should increase the amount of active RhoA. But surprisingly, they observed a higher rate of RhoA deactivation in DLC1 KO cells, which is counterintuitive: less GAP should result in a slower rate of deactivation. Less GAP should also lead to a lower rate of observed RhoA activation (smaller koff) and delayed peak. From the data, it seems hard to conclude on these two expectations since the initial rates (slopes right after the activation) and times at peak appear similar in both WT and DLC1 KO cells. Further on, the authors study the dynamics of DLC1 on FAs depending on the mechanical state and nicely show a causal decrease of DLC1 enrichment at FA upon FA reinforcement, hereby probing a positive feedback where RhoA activation is further amplified as the force exerted at FA is increasing.

Strengths:

- Experiments are precise and well done.
- Technically, the work brings original and interesting data. The use of transient optogenetic activation within focal adhesions together with a biosensor of activity is new and elegant.
- The link between DLC1 and global contractility/RhoA activity is clear and convincing.
- The surprisingly higher rate of RhoA deactivation in DLC1 KO cells is convincing, as well as the differences in the dynamics of RhoA between focal adhesions and plasma membrane.
- The correlation between DLC1 enrichment and focal adhesion dynamics is very clear.

Weaknesses:

- There is no explanation for the higher rate of RhoA deactivation in DLC1 KO cells.
- For the optogenetic experiments, it is not clear if we are looking at the actual RhoA dynamics of the activity or at the dynamics of the optogenetic tool itself.
- There is no model to analyze transient RhoA responses, however, the quantitative nature of the data calls for it. Even a simple model with linear activation-deactivation kinetics fitted on the data would be of benefit for the conclusions on the observed rates and absolute amounts.