

Xanthomonas citri subsp. citri type III effector PthA4 directs the dynamical expression of a putative citrus carbohydrate-binding protein gene for canker formation

Reviewed Preprint

v2 • July 22, 2024

Revised by authors

Reviewed Preprint

v1 • March 11, 2024

Xinyu Chen, Huasong Zou , Tao Zhuo, Wei Rou, Wei Wu, Xiaojing Fan

Plant Protection College, Fujian Agriculture and Forestry University, Fuzhou, China • School of Life Sciences and Health, Huzhou College, Huzhou, China

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

Abstract

Xanthomonas citri subsp. *citri* (*Xcc*), the causal agent of citrus canker, elicits canker symptoms in citrus plants because of the transcriptional activator-like (TAL) effector PthA4, which activates the expression of the citrus susceptibility gene *CsLOB1*. This study reports the regulation of the putative carbohydrate-binding protein gene *Cs9g12620* by PthA4-mediated induction of *CsLOB1* during *Xcc* infection. We found that the transcription of *Cs9g12620* was induced by infection with *Xcc* in a PthA4-dependent manner. Even though it specifically bound to a putative TAL effector-binding element in the *Cs9g12620* promoter, PthA4 exerted a suppressive effect on the promoter activity. In contrast, *CsLOB1* bound to the *Cs9g12620* promoter to activate its expression. The silencing of *CsLOB1* significantly reduced the level of expression of *Cs9g12620*, which demonstrated that *Cs9g12620* was directly regulated by *CsLOB1*. Intriguingly, PthA4 interacted with *CsLOB1* and exerted feedback control that suppressed the induction of expression of *Cs9g12620* by *CsLOB1*. Transient overexpression and gene silencing revealed that *Cs9g12620* was required for the optimal development of canker symptoms. These results support the hypothesis that the expression of *Cs9g12620* is dynamically directed by PthA4 for canker formation through the PthA4-mediated induction of *CsLOB1*.

eLife assessment

This **valuable** study provides new insight into potential subtle dynamics in effector biology. The data presented generally support the claims, but in some cases, significant controls are missing and so the overall work is currently **incomplete**. If the limitations can be addressed, this work should be of broad relevance for biologists interested in molecular plant-microbe interactions.

<https://doi.org/10.7554/eLife.91684.2.sa2>

Introduction

Citrus canker is a severe bacterial disease that impacts the production of citrus because it affects most commercial cultivars of citrus in tropical and subtropical areas. The early symptoms on leaves, stems, and fruits are characterized by tiny, slightly raised blister-like lesions with water-soaked margins, which then develop into necrotic, raised lesions (Brunings et al., 2003). In addition to the severe defoliation and premature fruit dehiscence that usually occur on heavily infected trees, the blemished fruits also have significantly reduced commercial value. The citrus canker is associated with five different bacterial pathotypes, including three pathotypes of *Xanthomonas citri* subsp. *citri* (*Xcc*) (A, A*, and A^w) and two pathotypes of *X. fuscans* subsp. *aurantifolii* (B and C) (Gottwald et al., 2002; Sun et al., 2004). The *Xcc* A-type canker is the most prevalent and affects a wide range of hosts, including *Citrus* spp. and many closely related rutaceous plants (Graham et al., 2004).

Xcc PthA4 is a transcription activator-like (TAL) effector that is delivered into host plants by a type III secretion system (T3SS). The *Xcc* strains defective in the *pthA4* gene cannot *pthA4* alone induces weak canker phenotypes in citrus leaves (Duan et al., 2007). The heterologous expression of T3SS and *pthA4* homologous genes in *Escherichia coli* elicits similar canker symptoms in citrus but not in tobacco (*Nicotiana benthamiana*), soybean (*Glycine max*), or cotton (*Gossypium hirsutum*) (Kanamori et al., 1998). Variants of the PthA4 effector are widely present across all five bacterial subspecies that cause citrus canker disease (Swarup et al., 1992; Al-Saadi et al., 2007; Jalan et al., 2013). PthA4 acts as a major virulence determinant for the formation of cankers when translocated into the nuclei of its citrus host (Yang et al., 1995). The superhelical structure of this TAL effector recognizes host sequences through its repeat-variable di-residue (RVD), which specifically recognizes the effector-binding element (EBE) in the gene promoter of host plants, and thus, activates the expression of host target gene (Deng et al., 2012). Notably, PthA4 acts in a protein–protein interaction manner during infection with *Xcc* (Domingues et al., 2010). In addition to the proteins associated with DNA repair (Domingues et al., 2010; de Souza et al., 2012), PthA4 specifically interacts with citrus CsMAF1 and releases the suppressive effect on RNA Polymerase III to increase the transcription of tRNA and cell proliferation (Soprano et al., 2013). Thus, *Xcc* PthA4 functions through versatile mechanisms and modulates target genes not only at the transcriptional level but also at the protein level.

PthA4 activates the transcription of the *CsLOB1* susceptibility gene in host citrus plants owing to the TAL effector DNA-binding principles and specificity (Moscou et al., 2009; Boch et al., 2009; Hu et al., 2014; Li et al., 2014). Remarkably, gene editing of the coding region or promoter sequence of *CsLOB1* has been used to promote the resistance of citrus to infection with *Xcc* (Jia et al., 2016; Peng et al., 2017). *CsLOB1* belongs to the lateral organ boundary domain (LBD) gene family, which encodes a set of plant-specific transcription factors (TFs) that contain cysteine-rich DNA binding motifs (Husbands et al., 2007). Plant LBD genes are involved in the regulation of plant growth, as well as the production of anthocyanins and nitrogen metabolism, and also respond to both hormonal and environmental stimuli (Majer et al., 2010). There are as many as 36 LBD proteins in citrus plants. *CsLOB2* and *CsLOB3* show the same involvement in inducing the formation of pustules as *CsLOB1* when they are induced by *Xcc*, which expresses an artificial TAL effector (Zhang et al., 2017). In this case, many differentially expressed citrus genes have been identified, including those related to cell wall metabolism. However, the *CsLOB2* and *CsLOB3* promoters do not contain specific binding elements for PthA4; thus, neither can be induced by PthA4 during infection by *Xcc* (Zhang et al., 2017).

Since *CsLOB1* is a TF, elucidating its downstream target genes is particularly important to understand the molecular events associated with the development of pustule symptoms (Hu et al., 2014; Li et al., 2014). PthA4 and *CsLOB1* form a regulatory cascade that activate the citrus

genes during *Xcc* infection (Duan et al., 2018 [↗](#)). Our previous research showed that the ectopic expression of the TAL gene *avrXa7* in *Xcc* suppressed the development of cankers but induced a yellowing phenotype around the inoculation site (Sun et al., 2018 [↗](#)). Moreover, 138 upregulated citrus plant genes were identified to respond to wild-type (WT) *Xcc* infection but were relatively downregulated in leaves inoculated with *Xcc* 29-1 that expresses *avrXa7*. This study aimed to characterize the target genes of the citrus hosts that are directly regulated by the PthA4-mediated induction of CsLOB1 during *Xcc* infection, as well as its potential role in canker development.

Results

The expression of *Cs9g12620* depends on *pthA4* during *Xcc* infection

Our previous study on a transcriptome analysis reported that 138 genes were upregulated in citrus plants infected with the WT *Xcc* and downregulated in plants inoculated with *Xcc* that expressed *avrXa7*. Those genes were assumed to be related to the major *Xcc* virulence effector PthA4. PthA4 has 17.5 RVDs in its central domain that correspond to an EBE of 18 nucleotides of the target gene promoter. Based on the TAL effector-DNA binding code, PthA4 can bind an optimal EBE of the 19 bp oligonucleotide -TATAAACCCCAATACCCT-, in which the first nucleotide “T” at 0 position is strictly conserved (Figure 1A [↗](#), Figure 1—figure supplement 1 [↗](#)). For each of the 138 genes, we downloaded and analyzed the 2000 bp DNA sequence upstream of the translation start site from the public citrus genome database (<https://www.citrusgenomedb.org/> [↗](#)). Possible PthA4 EBEs were identified in the promoter region of the two genes *Cs9g12620* and *CsLOB1* (*Cs7g27640.1*) (Figure 1A [↗](#)).

The transcript levels of *Cs9g12620* and *CsLOB1* were studied to determine whether they were associated with PthA4. In view of the transcriptome analysis, their transcript levels were first studied in sweet orange (*Citrus sinensis*) plants inoculated with the WT *Xcc* 29-1 and *Xcc* 29-1 that expressed *avrXa7*. The inoculation of *Xcc* 29-1 that expressed *avrXa7* did not cause canker symptoms and displayed a yellow phenotype (Figure 1B [↗](#)). In comparison with the uninoculated control, the level of expression of *Cs9g12620* was induced remarkably by inoculation with the WT *Xcc* 29-1 but not *Xcc* 29-1 that expressed *avrXa7* (Figure 1C [↗](#)). Their transcript levels were then studied in *C. sinensis* plants inoculated with the *pthA4* Tn5 insertion mutant Mxac126-80, which was impaired in its ability to cause canker symptoms (Figure 1D [↗](#)). In comparison with the plants inoculated with the WT *Xcc* 29-1, *Cs9g12620* and *CsLOB1* were downregulated in the plants infected with Mxac126-80 (Figure 1E [↗](#)). These findings demonstrated that *Cs9g12620* was activated by *Xcc* infection in a PthA4-dependent manner.

Cs9g12620 shows different profiles of expression from its homolog *Cs9g12650* owing to the genetic diversity in promoter region

A BLAST search was performed in the citrus genome database using the *Cs9g12620* sequence. A highly conserved homolog *Cs9g12650* was found closely downstream of *Cs9g12620* in chromosome 9 of the *C. sinensis* chromosome (Figure 2A [↗](#)). Their amino acid sequences are 86% homologous. The existence of *Cs9g12620* and *Cs9g12650* in the *C. sinensis* chromosome was verified by PCR amplification with two primer sets anchored to the two genes, respectively (Figure 2B [↗](#)). Their transcription was then assessed by semi-quantitative reverse transcription PCR (RT-PCR) analysis. The results showed that *Cs9g12620* was expressed at a high transcript level, while no transcripts of *Cs9g12650* were found, even in the plants inoculated with the WT *Xcc* 29-1 (Figure 2C [↗](#)).

A sequence analysis of the *Cs9g12620* and *Cs9g12650* promoter regions revealed that are 71% homologous. A prediction of the promoter in Softberry (<http://linux1.softberry.com/berry.phtml> [↗](#)) identified the core structure region from the same location in both the *Cs9g12620* and *Cs9g12650* promoters (Figure 2—figure supplement 1 [↗](#)). It should be noted that the predicted PthA4 EBE




Figure 1.

The expression of *Cs9g12620* depends on PthA4 during *Xcc* infection. **(A)** Predicted PthA4 binding elements of the *Citrus sinensis* genes upregulated in response to *Xcc* infection. The optimal effector-binding element (EBE) that corresponded to the PthA4 repeat variable di-residues (RVDs) were predicted based on the TAL effector-DNA binding code. Putative PthA4 EBEs in promoters of *Cs9g12620* and *CsLOB1* are shown below the optimal EBE of PthA4. **(B)** The disease symptoms of *Xcc* 29-1/*avrXa7* on *C. sinensis* leaves. The cell suspension (10^8 CFU/mL) was infiltrated into plant leaves, and the phenotype was recorded at 5 dpi. All the experiments were repeated three times with similar results. **(C)** A qRT-PCR analysis of the expression of *Cs9g12620* and *CsLOB1* in plants inoculated with *Xcc* that expressed *avrXa7*. The level of expression of each gene in the uninoculated samples was set as 1, and the levels in other samples were calculated relative to those baseline values. Values are the mean results from three biological replicates and are the means $\pm$ SD. (ANOVA, $P < 0.01$). The experiment was repeated three times. **(D)** The phenotype elicited by the *pthA4* mutant Mx126-80. The analyses were performed as described in **B**. **(E)** qRT-PCR analysis of the transcript levels of *Cs9g12620* and *CsLOB1* in plants inoculated with *Xcc*29-1 and Mx126-80. The analyses were the same as those conducted in **C**. ANOVA, analysis of variance; dpi, days post-inoculation; qRT-PCR, real-time quantitative reverse transcription PCR; SD, standard deviation; *Xcc*, *Xanthomonas citri* subsp. *citri*. The online version of this article includes the following figure supplement(s) for **Figure 1**:


Figure 1—figure supplement 1.

The location of the PthA4 EBE in the promoter of *Cs9g12620* in *Citrus sinensis*. The promoter sequences were retrieved from the citrus genome database. The location of binding site of PthA4 EBE is indicated by an underline. Translation initiation site is indicated by an arrow.



Figure 2.

Cs9g12620 and *Cs9g12650* show different profiles of expression owing to the genetic variation in promoter. (A) The location of *Cs9g12620* and *Cs9g12650* in the *Citrus sinensis* chromosome. The location was revealed by a BLAST search of the Citrus Genome Database (<https://www.citrusgenomedb.org/>). (B) Verification of the existence of *Cs9g12620* and *Cs9g12650* in the *C. sinensis* genome by PCR amplification. The specific primers for either gene were used for qRT-PCR analysis, which are shown in Supplementary file 1b. M, DL2000 DNA marker. (C) Semi-quantitative RT-PCR detection of the transcription of *Cs9g12620* and *Cs9g12650*. *CsActin* was used as the internal control gene. A total of 2 µg of total RNA extracted from *C. sinensis* was used to synthesize the first single-stranded cDNA. The specific primers for each gene were the same as those of the qRT-PCR analysis and are shown in Supplementary file 1b. M, DL2000 DNA marker. (D) Luciferase assays of *Cs9g12620* and *Cs9g12650* promoter activity. The *Cs9g12620* and *Cs9g12650* promoter luciferase fusions were transiently expressed in the *Nicotiana benthamiana* leaves. Luciferase activity was measured with a CCD imaging system at 2 days post-agroinfiltration. The image on the right shows the quantification of luciferase signal using a microplate luminescence reader. Values are the means ± SD (n=3 biological replicates; Student's *t*-test, **p<0.01). (E) Luciferase assays of *Cs9g12620* and *Cs9g12620-M* with the deletion of 22 bp at the 3'-terminus. The *Cs9g12620* and *Cs9g12620-M* promoter luciferase fusions were transiently expressed in the *N. benthamiana* leaves. Luciferase activity was measured with a CCD imaging system at 2 days post-agroinfiltration. The luciferase signal was quantified as described in D. CCD, charge-coupling device; qRT-PCR, real-time quantitative reverse transcription PCR; RT-PCR, reverse transcription PCR; SD, standard deviation. The online version of this article includes the following figure supplement(s) for Figure 2:



Cs9g12620	AAGTCAACAT	TAGCACGAAC	TTGTACTTAT	TGGATTGATA	ATGAGAATGA
Cs9g12650	GATTTATTAT	GAGAACGA--	TTAAATATAT	TAT-CTGTTA	ATGTGAAAGA
Cs9g12620	ATGATTAGTA	CTAACATGTC	CGTACAAGTA	TGATTCA--T	ATGTAACA--
Cs9g12650	TTGTAAAGTA	TTGTTCTGCT	ATTATCGGCA	GCATGTGCCT	CTGTAATATG
Cs9g12620	--ACAGAGTC	TTGATTAAGT	TTTGTACTTA	GAAATATTGT	AGAACTAAGA
Cs9g12650	AAATAGAATT	TGTACTTTAT	AT-GTACTTC	AAAATGTTAG	CTGATAGAAT
Cs9g12620	CTA---TAGG	AGCAAATCAC	GGCAACC---	-----AAT	GAAGCTTCAT
Cs9g12650	TCAAAGTAGG	AGCAAATCAC	GGCAACCGGC	AACCAGTAAT	GAAACTTCAT
LB1					
Cs9g12620	TTGGGGCTGC	TCCAACACTT	GAATCAAGAC	TTTTGAGTGG	AAAAACAAGA
Cs9g12650	GTGGGG---C	TCCAACGCAT	TAATGAAGAA	ATTTGTGTGA	AAATACAAGA
Cs9g12620	ATTAATTGAA	ATTATGGGAA	ATGGGGAGAC	CAAGTGGCAC	ATCTCACGGC
Cs9g12650	CTT-----GAG	ATT-TGGGAA	ATGCTTGGAC	AAAGGGACAC	TTCTCGCGGC
LB2					
Cs9g12620	TAATTTGGTG	GCCCTATAAC	GTGGTT--CC	TTATCTTCTT	T--CTTCCTT
Cs9g12650	TAATTCGGTG	GCCCTATAAC	GTGGTTTTCT	TTTTCTTCTT	TTTCTTCCTT
Cs9g12620	TCTTTTTGTC	TCCCACTTTA	ATATATAAAA	AAATCATACA	CGTACATGCT
Cs9g12650	TGCTTTTGTC	TTCAAATTC	ATATA-AAAA	TCATTATTTG	CGTAC-TGAT
PthA4 EBE					
Cs9g12620	CCCGTTGAGC	GTTTTCATTC	TATAAATACT	GTATTCATCC	TAGTTGCTTC
Cs9g12650	CTTGTTGAAC	TAACTCATTC	TATAAATATC	CTATTCAGTC	TACTTGCTTC
Cs9g12620	ATCAATATAG	AAAAGATGCC	TTGGTTTCTC	TCTCTT	
Cs9g12650	TTCAATCATAG	AAAA-----	-----	-----	

Figure 2—figure supplement 1.

Alignment of the promoter regions of *Cs9g12620* and *Cs9g12650* in *Citrus sinensis*. The promoter sequences were retrieved from the citrus genome database. The putative binding sites of CsLOB1 (LB1 and LB2) are shadowed with blue. The location of binding site of PthA4 (EBE) is indicated by an underline. A green shadow shows the location of predicted promoter core structure. The lack of 22 bp at the 3' terminus is shown by a red box.

was located in the promoter core structure. A sequence alignment indicated that the last 22 nucleotides at the 3' terminus in *Cs9g12620* promoter had been lost genetically in the *Cs9g12650* promoter (**Figure 2—figure supplement 1** [↗](#)). Fragments of 463 bp in the *Cs9g12620* promoter and 460 bp in the *Cs9g12650* promoter that composed the core structure were fused with a luciferase reporter. The transient overexpression of the promoter luciferase fusion indicated that the *Cs9g12650* promoter was unable to drive the expression of luciferase in *N. benthamiana* (**Figure 2D** [↗](#)). This demonstrated that the *Cs9g12650* promoter region lacked promoter activity. To confirm the effect of 22 nucleotides on promoter activity, we subsequently created a *Cs9g12620* promoter mutant with a deletion of 22 nucleotides at the 3' terminus. The mutant completely lost its ability to drive the expression of luciferase, suggesting that the 22 nucleotides are essential for activity of the promoter (**Figure 2E** [↗](#)).

PthA4 suppresses *Cs9g12620* promoter activity independent of the binding action

The PthA4-dependent induction of *Cs9g12620* during *Xcc* infection prompted us to determine whether it is directly regulated by PthA4. The 463 bp *Cs9g12620* promoter was used in yeast one-hybrid (Y1H) assays. The EGY48 yeast co-transformed with pGBKT7-pthA4 and pG221 showed weak β -galactosidase activity, which demonstrated that PthA4 functions as a transcriptional activator. The co-transformation of pGBKT7-pthA4 and pG221-PC*Cs9g12620* resulted in deep blue, suggesting that PthA4 binds the *Cs9g12620* promoter. The enhanced β -galactosidase activity was quantified by the Miller method (**Figure 3A** [↗](#)). In an electrophoretic mobility shift assay (EMSA), the application of 0.117, 0.469, 1.875, 7.5, 30 and 120 μ g of GST-PthA4 proteins partially hindered the mobility of 25 ng *Cs9g12620* promoter DNA. The application of 7.5, 30 and 120 μ g GST-PthA4 proteins completely retarded the mobility of 25 ng promoter DNA (**Figure 3B** [↗](#)). The negative control glutathione-S transferase (GST) tag was unable to bind the *Cs9g12620* promoter DNA.




Figure 3.

PthA4 binds to and suppresses the *Cs9g12620* promoter. **(A)** A yeast one-hybrid assay shows that PthA4 interacts with the *Cs9g12620* promoter. pGBKT7-pthA4 and pG221-PCs9g12620 were co-transformed into the yeast strain EGY48 and screened on synthetic dropout SD/-Ura and SD/-Ura/-Trp media. The transformed cells were dissociated by repeated freeze-thaw treatment and then used for β -galactosidase assays on filters soaked with Z buffer that contained 20 μ g/mL X-gal. The pGBKT7 and pG221 vectors were used as negative controls by co-transformation with the corresponding constructs. The columns below show the quantitative assay of β -galactosidase by the Miller method. Values are the means $\pm$ SD (n=3 biological replicates). (ANOVA, $P < 0.01$). **(B)** An electrophoretic mobility shift assay (EMSA) shows that PthA4 binds to the *Cs9g12620* promoter. Purified GST-PthA4 was incubated with 25 ng of promoter DNA in gel shift binding buffer at 28°C for 30 min and then analyzed in 6% non-denaturing PAGE. A GST tag was used as a negative control. **(C)** The overexpression of PthA4 suppressed the *Cs9g12620* promoter. PthA4 was transiently co-expressed with a *Cs9g12620* promoter GUS fusion in *Nicotiana benthamiana*. The promoter-driven GUS activity was assayed at 2 days post-agroinfiltration. The upper images show histochemical staining of the *Cs9g12620* promoter-driven GUS activity. The bar chart below shows the qRT-PCR analysis of the transcript levels of *gusA* and *pthA4*. The level of expression of each gene in the samples that expressed the promoter alone was set as 1, and the levels of the other samples were calculated relative to that. Values are the mean results from three biological replicates and are the means $\pm$ SD (ANOVA, $P < 0.01$). **(D)** EMSA shows that PthA4 binds to the 19 bp predicted binding site (EBE) in the *Cs9g12620* promoter. The purified GST-PthA4 was incubated with 25 ng of synthetic 19 bp EBE DNA fragment in gel shift binding buffer at 28°C for 30 min and then analyzed in 6% non-denaturing PAGE. **(E)** Luciferase assays of the activity of *Cs9g12620* promoter with site-directed mutation in PthA4 EBE. The first nucleotide acid “T” in EBE was mutated into A, C, or G, which generated PCs9g12620-M_A, PCs9g12620-M_C, and PCs9g12620-M_G, respectively. The mutants were fused with luciferase and transiently overexpressed in *N. benthamiana* leaves. The luciferase activity was measured with a CCD imaging system at 2 days post-agroinfiltration. **(F)** PthA4 did not bind to PCs9g12620-M_A, PCs9g12620-M_C and PCs9g12620-M_G. The left image represents a yeast one-hybrid assay that did not show an interaction of PthA4 between PCs9g12620-M_A, PCs9g12620-M_C, and PCs9g12620-M_G. The experiments were conducted in the same manner as those in **A**. The right image represents an EMSA that shows that PthA4 does not bind to the promoter mutants. The analysis was the same as that described in **B**. **(G)** Luciferase assays showing the suppression of PthA4 on PCs9g12620-M_A, PCs9g12620-M_C, and PCs9g12620-M_G. The promoter luciferase fusions were co-expressed with PthA4 in *N. benthamiana*. Co-expression with the empty binary vector pHB was used as the control. The analysis was the same as that described in **E**. The image on the right shows the quantification of luciferase signal using a microplate luminescence reader. Values are the means $\pm$ SD (n=3 biological replicates). ANOVA, analysis of variance; CCD, charge-coupling device; GST, glutathione-S transferase; PAGE, polyacrylamide gel electrophoresis; qRT-PCR, real-time quantitative reverse transcription PCR; SD, standard deviation. * $p < 0.05$. ** $p < 0.01$ (Student’s *t*-test).

To examine whether the activity was affected by PthA4, the *Cs9g12620* promoter was cloned into the binary vector pCambia1381 fused to a *gusA* reporter gene. Co-transformation with *pthA4* did not induce the *Cs9g12620* promoter activity. Histochemical staining revealed that the β -glucuronidase (GUS) activity in the leaves that co-expressed *pthA4* suppressed the promoter activity (**Figure 3C**). For further verification, the levels of expression of *gusA* and *pthA4* were evaluated by real-time quantitative reverse transcription PCR (qRT-PCR). Compared with the expression of the promoter-GUS fusion alone or in combination with the empty vector pHB control, the transcript level of *gusA* was reduced when the *Cs9g12620* promoter-GUS fusion was co-transformed with *pthA4* (**Figure 3C**).

In the EMSA analysis that utilized a synthetic PthA4 EBE (-TCCCACTTTAATATATAAA-) as a probe, 6, 12 and 24 μ g of GST-PthA4 protein partially hindered the mobility of oligonucleotide (**Figure 3D**). This demonstrated that PthA4 truly bound to the predicted EBE located at the promoter core structure (**Figure 2—figure supplement 1**). The first base pair “T” at 0 position in the sequence was then mutated into ‘A’, ‘C’, and ‘G’ in the *Cs9g12620* promoter, which generated the point mutations PCs9g12620-M_A, PCs9g12620-M_C, and PCs9g12620-M_G, respectively. The activities of

PCs9g12620-M_A and PCs9g12620-M_C were reduced compared with the WT PCs9g12620. In contrast, the mutation “T” to “G” resulted in a complete loss of promoter activity (**Figure 3E**). The Y1H and EMSA assays demonstrated that PthA4 could not bind PCs9g12620-M_A, PCs9g12620-M_C, or PCs9g12620-M_G (**Figure 3F**). However, the co-transformation of PthA4 with PCs9g12620-M_A, PCs9g12620-M_C, and PCs9g12620-M_G retained the ability to suppress the promoter activity (**Figure 3G**).

CsLOB1 binds multiple sites to induce the Cs9g12620 promoter

CsLOB1 showed transcription activity since β -galactosidase activity was found when it was co-transformed with pG221 into yeast EGY48. The β -galactosidase activity was remarkably enhanced when CsLOB1 was co-transformed with pG221, which harbored PCs9g12620 (**Figure 4A**). The binding of CsLOB1 to PCs9g12620 was further verified by an EMSA analysis (**Figure 4B**). Most importantly, the Cs9g12620 promoter activity was significantly induced when CsLOB1 was co-transformed with the PCs9g12620-GUS fusion (**Figure 4C**).

Two putative CsLOB1-binding sites (-CGGC-) were found in the Cs9g12620 promoter and located at the -268 and -151 positions (designated LB1 and LB2, respectively) (**Figure 4D**, **Figure 2—figure supplement 1**). To examine the role of the two CsLOB1-binding sites, the two binding sites LB1 and LB2 were deleted to generate the mutants PCs9g12620-M_{LB1}, PCs9g12620-M_{LB2}, and Cs9g12620-M_{LB1/2}. Deletion of the LB1 or LB2 site did not block the binding of CsLOB1 to PCs9g12620 in yeast (**Figure 4D**). In the EMSA analysis, CsLOB1 retarded the mobility of PCs9g12620-M_{LB1}, PCs9g12620-M_{LB2}, and PCs9g12620-M_{LB1/2} DNA (**Figure 4E**). Furthermore, the activities of these three promoter mutants were induced by CsLOB1 (**Figure 4F**). These results demonstrated that CsLOB1 binds to multiple sites in the Cs9g12620 promoter to induce promoter activity. Citrus tristeza virus (CTV)-induced gene silencing was conducted in *C. sinensis* plants, which generated CsLOB1-silenced plants. The efficiency of infection was evaluated by analyses of the expression of P23 gene harbored in the CTV vector. In comparison with the empty CTV-infected plants, the transcript level of Cs9g12620 was significantly reduced in the CsLOB1 silenced plants (**Figure 4G**).

PthA4 dynamically regulates the expression of Cs9g12620 during Xcc infection

To examine the exact effect of PthA4 on the Cs9g12620 promoter during Xcc infection, the CsLOB1 promoter GUS fusion (PCsLOB1-GUS) was first co-transformed with PthA4. A comparison with PCsLOB1-GUS expressed alone showed that the expression of PthA4 induced the PCsLOB1 promoter activity (**Figure 5A**). The co-expression of CsLOB1 with PthA4 had no effect on the PthA4-induced CsLOB1 promoter activity. This demonstrated that CsLOB1 is directly regulated by PthA4. When PthA4 and CsLOB1 were co-expressed with the Cs9g12620 promoter, the expression of PthA4 strikingly suppressed the CsLOB1-induced Cs9g12620 promoter activity (**Figure 5B**). To verify the suppressive effect by PthA4, *A. tumefaciens* GV3101 that expressed PthA4-FLAG was arranged at diverse concentrations of OD₆₀₀ values of 0.05, 0.1, 0.2, and 0.4. The results showed that the efficiency of suppression correlated with the high level of expression of PthA4-FLAG (**Figure 5C**).




Figure 4.

The *Cs9g12620* promoter is directly activated by *CsLOB1*. **(A)** A yeast one-hybrid (Y1H) assay shows that *CsLOB1* interacts with the *Cs9g12620* promoter. pGBKT7-*CsLOB1* and pG221-*Cs9g12620* were co-transformed into yeast strain EGY48 and screened on synthetic dropout SD/-Ura and SD/-Ura/-Trp media. The transformed cells were dissociated by repeated freeze-thaw treatment and then used for β -galactosidase assays on filters soaked with Z buffer that contained 20 μ g/mL of X-gal. pGBKT7 and pG221 vectors were used as the negative control by co-transformation with corresponding constructs. The columns below show the quantitative assay of β -galactosidase by the Miller method. Values are the means $\pm$ SD (n=3 biological replicates) (ANOVA, $P < 0.01$). **(B)** An electrophoretic mobility shift assay (EMSA) shows that *CsLOB1* binds to the *Cs9g12620* promoter. Purified GST-*CsLOB1* was incubated with 25 ng of promoter DNA in gel shift binding buffer at 28°C for 30 min and was then analyzed by 6% non-denaturing PAGE. **(C)** *Cs9g12620* promoter activity is induced by *CsLOB1*. *CsLOB1* was transiently co-expressed with the *Cs9g12620* promoter GUS fusion in *Nicotiana benthamiana*. Promoter-driven GUS activity was assayed at 2 days post-agroinfiltration. The upper images show the histochemical staining of *Cs9g12620* promoter-driven GUS activity. The bar chart below shows the qRT-PCR analysis of transcript levels of *gusA* and *pthA4*. The level of expression of each gene in the samples that expressed the promoter alone was set as 1, and the levels of the other samples were calculated relative to that. Values are the mean results from three biological replicates and are the means $\pm$ SD (ANOVA, $P < 0.01$). **(D)** A Y1H assay shows that *CsLOB1* interacts with the *Cs9g12620* promoter *PCs9g12620* and *CsLOB1*-binding site deletion mutants *PCs9g12620-M_{LB1}*, *PCs9g12620-M_{LB2}*, and *PCs9g12620-M_{LB1/2}*. The upper diagram shows the putative *CsLOB1* binding sites and the corresponding deletion mutants. The bar chart below represents a Y1H assay that shows the interaction of *CsLOB1* with the promoter mutants. The analyses were same as those conducted in **A**. **(E)** An EMSA showed that *CsLOB1* binds to *Cs9g12620-M_{LB1}*, *PCs9g12620-M_{LB2}*, and *PCs9g12620-M_{LB1/2}*. Purified GST-*CsLOB1* was incubated with 25 ng of promoter DNA in gel shift binding buffer at 28°C for 30 min. The analyses were same as those in **B**. **(F)** Luciferase assays show the role of *CsLOB1* on *PCs9g12620-M_{LB1}*, *PCs9g12620-M_{LB2}*, and *PCs9g12620-M_{LB1/2}*. Promoter luciferase fusions were co-expressed with *CsLOB1* in *N. benthamiana*. Co-expression with the empty binary vector pHB was used as the control. Luciferase activity was measured with a CCD imaging system at 2 days post-agroinfiltration. The image on the right shows the quantification of luciferase signal using a microplate luminescence reader. Values are the means $\pm$ SD (n=3 biological replicates, ** $p < 0.01$ [Student's *t*-test]). **(G)** qRT-PCR analysis of the transcript level of *Cs9g12620* in *CsLOB1*-silenced plants. *CsLOB1* was silenced in *Citrus sinensis* using the citrus tristeza virus (CTV)-based gene silencing vector CTV33. The effective infection by CTV was evaluated by the expression of *P23* gene harbored in CTV33. The silencing efficiency of *CsLOB1* was verified by comparison with the levels of expression in the WT and empty CTV33-infected control plants. The reduced expression of *Cs9g12620* is shown by comparison with the level of expression in the empty CTV33-infected plants. The qRT-PCR was performed as described in **C**. ANOVA, analysis of variance; CCD, charge-coupled device; GUS, β -glucuronidase; PAGE, polyacrylamide gel electrophoresis; qRT-PCR, real-time quantitative reverse transcription PCR; SD, standard deviation; WT, wild type.




Figure 5.

CsLOB1-induced *Cs9g12620* promoter activity is suppressed by PthA4. **(A)** PthA4 induced the activity of *CsLOB1* promoter. PthA4 was transiently co-expressed with the *CsLOB1* promoter GUS fusion in *Nicotiana benthamiana*. GUS activity was assayed at 2 days post-agroinfiltration. The upper images show the histochemical staining of *CsLOB1* promoter (*PCsLOB1*)-driven GUS activity. The bar chart below shows the qRT-PCR analysis of the transcript levels of *gusA*, *CsLOB1*, and *pthA4*. The transcript level of each gene in the samples that expressed *PCsLOB1*-GUS was set as 1, and the levels of the other samples were calculated relative to that. Values are the mean results from three biological replicates and are the means $\pm$ SD (ANOVA, $P < 0.01$). **(B)** PthA4 suppressed the *CsLOB1*-induced activity of the *Cs9g12620* gene promoter. The suppression was evaluated by the co-expression of PthA4 with *CsLOB1* and *PCs9g12620*-GUS. Histochemical staining and a qRT-PCR analysis were performed as described in **A**. **(C)** PthA4 suppressed the *CsLOB1*-induced *PCs9g12620* activity in a dose-dependent manner. *CsLOB1* and *PCs9g12620*-LUC were co-expressed with PthA4-FLAG in *N. benthamiana*. The agrobacteria that expressed PthA4-FLAG were arranged to a cell concentration series of OD₆₀₀ values of 0.05, 0.1, 0.2, 0.4, and 0.6. The luciferase signal was quantified with a microplate luminescence reader. Values are the means $\pm$ SD ($n=3$ biological replicates, ANOVA, $P < 0.01$). The image on the bottom right shows the expression of PthA4-FLAG by immunoblotting with anti-FLAG. **(D)** A qRT-PCR analysis of the levels of expression of *pthA4*, *CsLOB1*, and *Cs9g12620* in *Citrus sinensis* leaves inoculated with *Xcc* 29-1. The cell suspension (10^8 CFU/mL) was infiltrated into the plant leaves. qRT-PCR was performed at 0, 5, and 10 days post-inoculation (dpi). The level of expression of each gene at 0 dpi was set as 1, and the level in other samples was calculated relative to those baseline values. Values are the mean results from three biological replicates and are the means $\pm$ SD. (ANOVA, $P < 0.01$). **(E)** The level of expression of *Cs9g12620* was dynamically related with that of PthA4 during *Xcc* infection. The disease symptoms caused by WT *Xcc* 049, TAL-free mutant 049E, and *Xcc* 049E/*pthA4* on *C. sinensis* leaves. The phenotype was recorded at 10 dpi. A qRT-PCR analysis was conducted to evaluate the levels of expression of *pthA4*, *CsLOB1*, and *Cs9g12620*. The experiment was performed as described in **D**. **(F)** Growth of wildtype *Xcc* 29-1 in *C. sinensis* plants. The *Xcc* 29-1 cells of 10^8 CFU/ml were infiltrated into citrus leaves. At 5 and 10 days post inoculation, bacteria were recovered from leaves and counted on Nutrient Agar (NA) plates. Error bars represent the standard deviation from three independent experiments (Student's *t*-test, **** $p < 0.0001$). **(G)** qRT-PCR analysis of *pectin esterase* (*orange1.1t02719*) and *expansin* (*cs9g15150*) genes expression levels in *C. sinensis* leaves inoculated with *Xcc* 29-1. The experiment was performed as described in **D**. ANOVA, analysis of variance; GUS, β -glucuronidase; qRT-PCR, real-time quantitative reverse transcription PCR; SD, standard deviation.

To study the regulatory pattern of *pthA4* on the expression of *Cs9g12620* during canker development, the levels of expression of *pthA4*, *CsLOB1*, and *Cs9g12620* were quantitatively assayed in plants inoculated with the WT *Xcc* 29-1. Even though the levels of expression of *pthA4*, *CsLOB1*, and *Cs9g12620* were enhanced at 5 and 10 days post-inoculation (dpi) compared with 0 dpi, their pattern of expression differed between 5 and 10 dpi. *pthA4* and *CsLOB1* showed the highest levels of expression at 10 dpi. In contrast, the level of expression of *Cs9g12620* at 10 dpi was lower than that at 5 dpi (**Figure 5D**). To verify this special pattern of expression, *Xcc* strain 049, its TAL-free mutant *Xcc* 049E, and *Xcc* 049E that expressed *pthA4* (*Xcc*049E/*pthA4*) were inoculated on *C. sinensis* plants. The TAL-free mutant 049E caused extremely weak canker symptoms, which were restored by expressing *pthA4* in *Xcc* 049E (**Figure 5E**). No transcript of *pthA4* was detected in the plants inoculated with *Xcc* 049E. In the plants inoculated with *Xcc* 049 or *Xcc* 049E/*pthA4*, the levels of expression of *pthA4* and *CsLOB1* were enhanced; moreover, the transcript level at 10 dpi was higher than that at 5 dpi (**Figure 5E**). In contrast, the level of expression of *Cs9g12620* at 10 dpi was distinctively lower than that at 5 dpi (**Figure 5E**). By comparison with 5 dpi, the cell number of *Xcc* 29-1 *in planta* was increased 1.5-fold at 10 dpi (**Figure 5F**). However, the expressions of *pectin esterase* and *expansin* genes associated with canker development were decreased at 10 dpi (**Figure 5G**). This supported the hypothesis that the overexpression of PthA4 exerts a feedback inhibition on the expression of *Cs9g12620*, which was consistent with the result that the transient overexpression of PthA4 suppressed the *Cs9g12620* promoter activation by *CsLOB1*. The dynamic regulation of *Cs9g12620* is closely related with canker development.

PthA4 interacts with CsLOB1

PthA4 exerted a role in suppressing the *Cs9g12620* promoter independently of its binding to EBE, which led us to determine whether PthA4 could physically interact with CsLOB1. In yeast two-hybrid (Y2H) assays, a positive interaction was observed in yeast strain AH109 co-transformed with AD-PthA4 and BD-CsLOB1 (**Figure 6A** [↗](#)). In a GST pull-down assay, GST-CsLOB1 successfully pulled down MBP-PthA4. In the control lane, the GST tag did not pull down MBP-PthA4 (**Figure 6B** [↗](#)). The interaction between PthA4 and CsLOB1 *in vivo* was further verified by split luciferase assays on *N. benthamiana* (**Figure 6C** [↗](#)).

Cs9g12620 is involved in the formation of canker symptoms

The *Cs9g12620*, as well as *CsLOB1* and *pthA4*, were transiently overexpressed in *C. sinensis*. The expression of *pthA4* induced a canker-like phenotype. The overexpression of *Cs9g12620* resulted in yellowing and hypertrophy of the leaves of *C. sinensis* plants (**Figure 7A** [↗](#)). The overexpression of *CsLOB1* did not cause a distinctive phenotype in *C. sinensis* plants. Thus, the overexpression of *Cs9g12620* elicited a canker-like symptom in *C. sinensis* that resembled that induced by *pthA4*, although the symptoms were weaker. Transmission electron microscopy (TEM) was conducted to investigate the changes in *C. sinensis* leaf structure. In the control leaves infiltrated with empty vector, large air spaces were found between the cells in spongy mesophyll. In comparison, more cells and less air space were found in the spongy mesophyll of the leaves that transiently expressed *Cs9g12620* and *pthA4* (**Figure 7B** [↗](#)).




Figure 6.

PthA4 interacts with CsLOB1. **(A)** A yeast two-hybrid assay shows that PhtA4 interacts with CsLOB1. pGADT7-PthA4 and pGBKT7-CsLOB1 were co-transformed in yeast cells and screened on synthetic dextrose media that lacked leucine and tryptophan (SD/-Leu-Trp). A single yeast colony was then selected for serial dilution and grown on SD/-Leu-Trp and SD/-Ade-His-Leu-Trp to examine the interaction. The yeast co-transformed with pGADT7-T and pGBKT7-53 served as a positive control. The yeast co-transformed with pGADT7-T and pGBKT7-lam served as a negative control. **(B)** GST pull-down assay of the interaction between PhtA4 and CsLOB1. The recombinant GST-CsLOB1 and MBP-PhtA4 proteins were used for GST pull-down assays. The GST protein served as a negative control. The pull-down of MBP-PhtA4 was verified by anti-MBP immunoblotting. The experiment was repeated three times with similar results. **(C)** A split luciferase assay for the interaction of PhtA4 and CsLOB1 *in vivo*. *Nicotiana benthamiana* leaves were co-infiltrated with agrobacteria that harbored 35S: PthA4-nLUC and 35S:cLUC-CsLOB1. The image on the right shows the expression of respective proteins. Images of chemiluminescence were obtained by treatment with 0.5 μ M luciferin 48 h post-infiltration. Similar results were obtained in three biological repeats. GUS, β -glucuronidase.




Figure 7.

Cs9g12620 is involved in the formation of canker symptoms. **(A)** The transient overexpression of Cs9g12620 results in hypertrophy in the *Citrus sinensis* leaves. The expression of *pthA4* was used as the positive control. The cultured *Agrobacterium* cells were adjusted to an OD₆₀₀ of 0.01 and infiltrated into young *C. sinensis* leaves. The phenotypes were recorded at 15 days post-agroinfiltration. The area of infiltration for each gene is indicated by a dotted line. **(B)** Transmission electron micrograph of *C. sinensis* leaf tissues. Transmission electron micrographs of the cross sections of citrus leaf tissue. The leaves were sampled at 15 days post-agroinfiltration. The changes in tissue structure induced by Cs9g12620 and *pthA4* are indicated by a box. The bars represent 200 μ m. **(C)** A qRT-PCR analysis of the silencing of Cs9g12620 in *C. sinensis*. Cs9g12620 was silenced using the citrus tristeza virus (CTV)-based gene silencing vector CTV33. The effective infection by CTV was evaluated by measuring the level of expression of the P23 gene harbored in CTV33. The silencing efficiency of Cs9g12620 was verified by comparison with the levels of expression in the WT and empty CTV33-infected control plants. Values are the mean results from three biological replicates. (Student's *t*-test, ***P* < 0.01). The experiment was repeated three times. **(D)** The canker symptoms on Cs9g12620-silenced citrus plants were caused by the WT Xcc 29-1. The Xcc 29-1 cells of 10⁷ CFU/mL were infiltrated into citrus leaves to analyze the development of canker symptoms. The canker symptoms were recorded at 6 days post-inoculation (dpi). The areas of infiltration are indicated by dotted lines. The inoculation of the plants infected by the empty vector CTV33 was used as the negative control. **(E)** Growth of Xcc 29-1 in Cs9g12620-silenced citrus plants. A volume of 10⁷ CFU/mL of Xcc 29-1 cells were infiltrated into citrus leaves. At 2, 4, and 6 dpi, bacteria were recovered from leaves and counted on Nutrient Agar (NA) plates. Error bars represent the standard deviation from three independent experiments (Student's *t*-test, **p* < 0.05, ***p* < 0.01, ****p* < 0.001). **(F)** A qRT-PCR analysis of Cs9g12620, *CsLOB1* and *pthA4* in Cs9g12620-silenced citrus plants inoculated with Xcc 29-1. The experiment was performed as described in C. **(G)** The canker symptoms on Cs9g12620-silenced citrus plants were caused by the WT Xcc 049, TAL-free mutant 049E, and 049E/*pthA4*. The experiment was performed as described in D. **(H)** Growth of WT Xcc 049, 049E, and 049E/*pthA4* in Cs9g12620-silenced citrus plants. At 6 dpi, bacteria were recovered from leaves and counted on nutrient rich (NA) plates. The experiment was performed as described in E. dpi, days post-inoculation; WT, wild type.

The contribution of Cs9g12620 to canker development was subsequently examined by citrus tristeza virus (CTV)-induced gene silencing in *C. sinensis* plants. The efficiency of gene silencing was evaluated in one-month-old leaves at 60 days post-stem inoculation. The infection by CTV was indicated by verification of the transcripts of CTV P23 gene. The transcript level of Cs9g12620 was then verified to be reduced by approximately 70% in the gene-silenced *C. sinensis* plants (Figure 7C). Inoculation of the WT Xcc 29-1 caused remarkably attenuated canker symptoms in the Cs9g12620-silenced plants. In comparison with the control plants that were infected with empty CTV, there were remarkably fewer symptoms on the Cs9g12620-silenced plants (Figure 7D). The Xcc 29-1 in the Cs9g12620-silenced plants grew more slowly than in the control plants. Significantly fewer cell numbers were isolated from the Cs9g12620-silenced plants than from the control plants at 2, 4, and 6 days post-inoculation (Figure 7E). The expression levels of Cs9g12620 and *CsLOB1* genes in Cs9g12620-silenced plants inoculated with Xcc 29-1 showed that the induced transcription levels of both were decreased, and the expression levels of *pthA4* were also significantly decreased, down-regulated by 38%, 15% and 16%, respectively (Figure 7F). The TAL-free 049E caused extremely weak canker symptom, and wildtype 049 and 049E/*pthA4* caused severe canker on control plants. On Cs9g12620-silenced plants, wildtype 049 and 049E/*pthA4* caused greatly reduced canker symptom, while the 049E/*pthA4* caused weak canker symptom alike to TAL-free mutant 049E (Figure 7G). Furthermore, the population of 049 and 049E/*pthA4* recovered from Cs9g12620-silenced plants was lower than that in control plants. By contrast, population of 049E was low in either Cs9g12620-silenced or control plants (Figure 7H). This demonstrated expression of *pthA4* in 049E did not restore bacterial virulence on Cs9g12620-silenced plants. The level of expression of Cs9g12620 is correlated with the formation of canker symptoms in citrus plants.

Discussion

The ectopic expression of *avrXa7* in *Xcc* 29-1 did not induce canker symptoms on citrus plants (Sun et al., 2018 [\[1\]](#)), which was consistent with a previous study that reported that the browning phenotype was elicited by an *Xcc* L2 strain that expressed *avrXa7* (Ishihara et al., 2003 [\[2\]](#)). This was hypothesized to be caused by two groups of differentially expressed genes, including (i) resistance genes induced by *AvrXa7*, and (ii) susceptibility genes induced by *PthA4* but suppressed by the expression of *AvrXa7*. A transcriptome analysis was utilized to identify the different relevant citrus genes and showed that *AvrXa7* specifically activated the aspartic protease gene *Cs7g06780.1* and linoleate 13S-lipoxygenase gene *orange1.1t03773.1* (Sun et al., 2018 [\[1\]](#)). In this study, we focused on identification of the genes induced by *PthA4* for canker formation. Based on the pattern of expression and analysis of promoter activity, *Cs9g12620* was proven to be dynamically regulated by the *PthA4*-mediated induction of *CsLOB1* for the development of canker symptoms. This supported the hypothesis that the *PthA4*-mediated induction of *CsLOB1* serves a critical role in the complicated regulatory network on gene expression during *Xcc* infection.

The original purpose of this study was to explore the genes that were directly targeted by *PthA4* through TAL effector-DNA recognition. Based on a sequence analysis, the two genes *Cs9g12620* and *CsLOB1* were predicted to contain *PthA4* EBEs. An analysis of the pattern of expression revealed that *Cs9g12620* and *CsLOB1* were induced by *Xcc* infection in a *PthA4*-dependent manner. Because the susceptibility gene *CsLOB1* has already been proven to be targeted by *Xcc* *PthA4*, the binding and regulatory role of *PthA4* on the *Cs9g12620* promoter was assayed experimentally. Y1H, EMSA, and promoter reporter fusion analyses were used and enabled the conclusion that *PthA4* bound to the *Cs9g12620* promoter, whereas it suppressed the promoter activity. Since two *CsLOB1* putative binding sites were predicted from the *Cs9g12620* promoter, we demonstrated that *CsLOB1* bound directly to the *Cs9g12620* promoter. Most importantly, *CsLOB1* could activate the *Cs9g12620* promoter. As such, *Cs9g12620* is actually a target of *CsLOB1*.

The ectopic expression of *CsLOB1* restores the mutant canker symptoms of *Xcc* *pthA4*, which demonstrates that *CsLOB1* is the key target of *PthA4* and thus, induces pustule symptoms (Duan et al., 2018 [\[3\]](#)). The transient overexpression of *CsLOB1* did not induce an obvious phenotype in the citrus plant leaves, which was consistent with the findings of previous research (Hu et al., 2014 [\[4\]](#)). Instead, the contribution of *CsLOB1* to canker symptoms has been verified by the transformation of ‘Duncan’ grapefruit (*Citrus x paradisi*) plants with a 35S:*CsLOB1*-glucocorticoid receptor construct, which suggested that the continuous expression of *CsLOB1* is essential for the development of canker symptoms (Duan et al., 2018 [\[3\]](#)). Duan et al. (2018) [\[3\]](#) reported that a number of citrus genes appear to be associated with *CsLOB1*, but only the *Cs2g20600* promoter probe interacted with *CsLOB1*. *Cs2g20600* encodes a zinc finger C3HC4-type RING finger protein with E3 ubiquitin ligase activity (Duan et al., 2018 [\[3\]](#)). In this study, we found that *Cs9g12620* was another target that is directly regulated by *CsLOB1*. The regulatory role of *CsLOB1* includes promoting host cell proliferation, regulating citrus cell wall remodeling, and affecting hormone signaling pathways, which is accompanied by the optimal growth of *Xcc* (Zou et al., 2021 [\[5\]](#)). Although it has been reported to specifically bind to -CGGC- in its target promoter region, *CsLOB1* retained the ability to bind the *Cs9g12620* promoter DNA when the two specific binding sites LB1 and LB2 were deleted. We hypothesized that there are other putative binding sites of *CsLOB1* in the *Cs9g12620* promoter, and multiple citrus genes are directly targeted by *CsLOB1* for the development of cankers during *Xcc* infection.

PthA4 acts as a sole disease TAL required by *Xcc* to cause symptoms of canker on citrus plants. It localizes to the nuclei following secretion into host cell by the type III secretion system. A transient overexpression analysis showed that *PthA4* exhibited a particular suppressive effect on the *CsLOB1*-activated *Cs9g12620* promoter activity. Since the *PthA4* binding site is critical for activity

of the *Cs9g12620* promoter, the binding of PthA4 suppressed the promoter activity. This prompted us to study whether PthA4 affects CsLOB1 alternatively. Y2H and pull-down assays showed that PthA4 interacted with CsLOB1 at the protein-protein level. This provided a new insight to study the dynamical regulatory role of PthA4 during *Xcc* infection since we found that PthA4 affects the expression of *Cs9g12620* in a dose-dependent manner. When PthA4 was expressed at a high level at 10 dpi, the level of expression of *Cs9g12620* was surprisingly lower. This suggested that there is a precise mechanism of the PthA4-CsLOB1-*Cs9g12620* regulatory cascade during infection with *Xcc*.

Although the amino acid sequence of *Cs9g12650* was 86.3% homologous with that of *Cs9g12620*, they differed in their promoter sequences. The promoter of *Cs9g12650* lacks 22 bp of nucleotides at its 3' terminus, which are critical for the activity of its promoter. The activity of *Cs9g12620* promoter was completely lost if the 22 bp sequence was deleted by mutation. Therefore, *Cs9g12650* is not transcribed in citrus plants. This supported the hypothesis that the null activity of *Cs9g12650* promoter was owing to genetic diversity in citrus plants, which leads to different patterns of expression between the two paralogs in citrus. The 22 bp sequence at the *Cs9g12620* promoter could be a potential site for gene editing to create citrus plants that are resistant to citrus canker. A similar regulation of expression has been found in *CsLOB1*; *CsLOB2* and *CsLOB3* show the same ability to induce pustule formation as *CsLOB1*, but their promoters do not contain the PthA4 EBE; therefore, neither can be induced by PthA4 (Zhang et al., 2017 [DOI](#)).

The *Cs9g12620* protein contains three distinctive conserved domains, including a signal peptide, a duplicated DUF642 region, and a bacterial exosortase protein domain. The duplicated DUF642 region has been shown to be a carbohydrate-binding domain. The exosortase system is involved in directing transport across the plasma membrane, with additional posttranslational modifications, such as glycosylation (Haft et al., 2012 [DOI](#)). This system is distributed across many biofilm-producing environmental bacteria and is closely related to the *N*-acyl amino acid synthases in Proteobacteria (Craig et al., 2011 [DOI](#)). It was hypothesized that *Cs9g12620* could possibly be involved in the transportation of carbohydrates in citrus cells. This was similar to the SWEET genes induced by the effectors that pathogenic bacteria secrete into plants, and they export intracellular sugar into the extracellular space, which can be utilized by pathogenic bacteria (Hu et al., 2014 [DOI](#)). The binding of carbohydrates and the exact role for canker formation merits further study.

In conclusion, the putative carbohydrate-binding protein gene *Cs9g12620* was identified as a target of CsLOB1, which was directly regulated by PthA4-mediated induction of CsLOB1. In addition to the enhanced pattern of expression in response to *Xcc* infection, *Cs9g12620* was dynamically feedback regulated by PthA4 in a dose-dependent manner. At the early infection stage, the presence of PthA4 is responsible for the induction of CsLOB1, which is necessary for the activation of *Cs9g12620*. At the late infection stage, the over-redundancy of PthA4 exerts a feedback suppression effect on the activation of *Cs9g12620* by CsLOB1 (**Figure 8** [DOI](#)). The findings provide strong evidence to understand the molecular complexity of PthA4-mediated induction of CsLOB1 during the development of citrus canker.

Materials and methods

Plant and bacterial materials

C. sinensis and *N. benthamiana* were grown in a greenhouse (26/24°C light/dark) under long-day conditions (16/8 h light/dark). The strains and plasmids used in this study are listed in Supplementary file 1a. *Xcc* strain 29-1 was collected from *C. sinensis* in Jiangxi Province, China (Ye et al., 2013 [DOI](#)). The *Xcc* 29-1/*avrXa7* strain is a *Xcc* 29-1 strain that expresses *avrXa7* from *Xanthomonas oryzae* pv. *oryzae*, the causal agent of rice bacterial blight (Sun et al., 2018 [DOI](#)). Mxac




Figure 8.

Proposed model of the expression of *Cs9g12620* directed by PthA4-mediated induction of CsLOB1. The regulation of putative carbohydrate-binding protein gene *Cs9g12620* was divided into two steps. At the early infection stage, the presence of PthA4 is responsible for the induction of CsLOB1, which is responsible for the induction of *Cs9g12620*. At the late infection stage, the over-redundancy of PthA4 exerts a feedback suppression effect on the activation of *Cs9g12620* by interacting with CsLOB1.

126-80 is a *pthA4* mutant with a Tn5 insertion at the 1789 bp position in the middle of RVD coding sequence (Song et al., 2015). *Xcc* strain 049 was collected from *C. sinensis* in Chongqing Province, China (Ye et al., 2013). *Xcc* 049E is a TAL-free strain of *Xcc* 049, and *Xcc* 049E/*pthA4* is *Xcc* 049E that expresses the *pthA4* gene (Ge et al., 2019). All the *Xcc* strains were cultured in nutrient broth (NB) or nutrient agar (NA, 1.5% agar) at 28°C (Song et al., 2015). *Agrobacterium tumefaciens* strain GV3101 was cultured in YEP medium (10 g/L yeast extract, 10 g/L peptone, and 5 g/L sodium chloride, pH 7.0) at 28°C. The yeast strains EGY48 for the Y1H assays and AH109 for the Y2H assays were cultured at 30L in YPD media (10Lg/L yeast extract, 20Lg/L peptone, and 20Lg/L glucose, pH 4.5–5.0).

Inoculation of *Xcc* in citrus host plants

The cultured *Xcc* cells were suspended in sterile distilled water to a final concentration of 10^8 CFU/mL ($OD_{600} = 0.3$). The bacterial suspension was inoculated onto fully expanded *C. sinensis* leaves with a needleless syringe (Wu et al., 2019). The canker disease symptoms were recorded at 5 or 10 dpi. To assess the bacterial growth *in planta*, 0.25 cm² leaf discs from the areas of infiltration were collected to count the cell numbers. The cell numbers were calculated as CFU per cm². All the experiments were repeated three times.

Real-time quantitative reverse transcription PCR (qRT-PCR)

qRT-PCR was performed using a CFX Connect Real Time PCR detection system (Bio-Rad, Shanghai, China) using iTaq Universal SYBR Green Supermix (Bio-Rad). The total RNA was isolated from *C. sinensis* and *N. benthamiana* leaves using a Plant RNA Kit (Omega, Shanghai, China). A volume of 2 µg of total RNA was reverse transcribed into single-stranded cDNA using AMV reverse transcriptase (TaKaRa, Dalian, China). The primer sequences used for qRT-PCR are listed in Supplementary file 1b. The *C. sinensis* *CsActin* and *N. benthamiana* *NbEF1 α* genes were used as internal controls to statistically analyze the levels of relative expression (Wu et al., 2019). All the experiments were repeated three times.

Luciferase assays

To study the interaction *in vivo*, the *PthA4* gene from *Xcc* was fused with nLUC at the N terminus. The *CsLOB1* gene from *C. sinensis* was fused with cLUC at the C terminus. cLUC-*CsLOB1* was co-expressed with nLUC-*PthA4* in 4-week-old *N. benthamiana* leaves. To analyze the promoter activity, the 463 bp *Cs9g12620* promoter region was directly introduced into vector pGWB435 by a Gateway cloning strategy to generate *PCs9g12620*-LUC. Site-directed mutation of the first nucleotide “T” in EBE of *PthA4* or deletion of *CsLOB1* binding sites LB1 and LB2 were created by overlapping PCR using the primers listed in Supplementary file 1b. The promoter mutants *M_A*, *M_C*, *M_G*, *M_{LB1}*, *M_{LB2}*, and *M_{LB1/2}* that were generated were fused with luciferase (LUC) in pGWB435. The activities of LUC were assayed after transient overexpression in *N. benthamiana*. For each assay at 2 dpi, the inoculated leaves were treated with 0.5 mM luciferin, kept in the dark for 1 min to quench the fluorescence, and then used to capture luciferase luminescence images by a cooled charge-coupled device (CCD) imaging apparatus (Roper Scientific, Trenton, NJ, USA). Each experiment had three replicates.

Yeast one-hybrid assays

The pGBKT7-*pthA4* and pGBKT7-*CsLOB1* constructs were created by inserting the *pthA4* and *CsLOB1* gene into the pGBKT7 vector, respectively (Supplementary file 1b). A 463 bp DNA fragment of the *Cs9g12620* gene promoter was cloned into the pG221 vector, which resulted in pG221-*PCs9g12620* (Supplementary file 1a and 1b). To examine the role of *PthA4* and *CsLOB1* binding sites in interaction, mutants *M_A*, *M_C*, *M_G*, *M_{LB1}*, *M_{LB2}*, and *M_{LB1/2}* were cloned in pG221 (Supplementary file 1b). Mutants *M_A*, *M_C*, and *M_G* were co-transformed with pGBKT7-*pthA4*, and mutants *M_{LB1}*, *M_{LB2}*, and *M_{LB1/2}* were co-transformed with pGBKT7-*CsLOB1* into yeast EGY48 using the standard LiAc-mediated yeast transformation (Ye et al., 2004). pG221 that harbored a

CYC1 core promoter was used for a transactivation assay in PthA4 and CsLOB1. The yeast transformants were screened on synthetic dropout SD/-Ura and SD/-Ura/-Trp media plates (Ye et al., 2004 [\[1\]](#)). The transformed cells were subjected to five freeze-thaw cycles using liquid nitrogen, spotted onto a sterile filter, and soaked with Z buffer that contained 20 µg/mL of X-gal. β-galactosidase activity was scored within 8 hours of incubation. The β-galactosidase activity was quantified using the standard Miller method (Marburg et al., 2016 [\[2\]](#)). Each experiment was performed three times.

Electrophoretic mobility shift assays

The binding of both PthA4 and CsLOB1 to the *Cs9g12620* promoter DNA (including M_{LB1} , M_{LB2} , and $M_{LB1/2}$) was detected by an Cy5-labeled probe electrophoretic mobility shift assay (EMSA) as described previously (Fan et al., 2020 [\[3\]](#)). The binding of PthA4 to the *Cs9g12620* promoter DNA (M_A , M_C , and M_G) was visualized by ethidium bromide staining as part of EMSA. The GST-pthA4 and GST-CsLOB1 constructs were created by separately inserting the *pthA4* and *CsLOB1* genes into the pET41a(+) vector (Supplementary file 1a). GST-PthA4, GST-CsLOB1, and GST tag were expressed separately in *E. coli* BL21 (DE3) by induction with 1.0 mM isopropyl-β-D-thiogalactopyranoside (IPTG). The purified GST-PthA4, GST-CsLOB1, and GST tag were subjected to ethidium bromide staining with concentration gradients by a two-fold dilution. The proteins were incubated with 25 ng of promoter DNA in gel shift binding buffer (50 ng/µL Poly (dI•dC), 0.05 M Tris-HCl, 1.0 mM EDTA, 0.15 M KCl, and 1.0 mM dithiothreitol). The GST-PthA4 protein, GST-CsLOB1 protein, and GST tag were diluted into series that consisted of 0.117, 0.469, 1.875, 7.5, 30, and 120 µg; 0.234, 0.938, 3.75, 7.5, 15, and 60 µg; and 1.953, 7.813, 3.75, and 15.625 µg; respectively. After incubation at 28°C for 30 min, the mixtures were subjected to a DNA mobility analysis by electrophoresis in a 6% non-denaturing acrylamide gel. The DNA bands were visualized by fluorescence imaging using a Typhoon Trio Variable Mode Imager (GE Healthcare, Chicago, IL, USA). Analyses of the binding to promoter mutants M_A , M_C , and M_G were conducted with the same strategy but using different GST-PthA4 protein series. Each experiment had three replicates.

GUS activity assays

The *Cs9g12620* promoter region was cloned into the binary vector pCambia1381 to fuse with the *gusA* gene (Supplementary file 1b). The coding sequences of *pthA4* and *CsLOB1* were individually cloned into the binary vector pHB to obtain pHB-pthA4 and pHB-CsLOB1, respectively. The promoter-GUS fusion was co-transformed with pHB-pthA4 or pHB-CsLOB1 to examine the promoter activity in *N. benthamiana*. β-Glucuronidase (GUS) activity was assayed by histochemical staining of the activity after 2 dpi (Hu et al., 2014 [\[4\]](#)). Moreover, the levels of expression of *gusA* in the leaf samples were evaluated by qRT-PCR to quantify the promoter activity (Sun et al., 2020 [\[5\]](#)). Each experiment had three replicates.

Yeast two-hybrid assays

The coding sequences of *pthA4* and *CsLOB1* were cloned individually into both the pGADT7 and pGBKT7 vectors. AD-PthA4/BD-CsLOB1 was then transformed into the yeast strain AH109 and screened on synthetic drop-out media that lacked leucine and tryptophan (SD/-Leu-Trp). Single colonies were cultured, serially diluted, and grown on SD/-Leu-Trp and SD/-Ade-Leu-Trp-His media to examine the possible interaction. Additionally, pGADT7-T and pGBKT7-53 were used as positive controls, while pGADT7-T and pGBKT7-lam served as negative controls. Each experiment had three replicates.

GST pull-down assay

PthA4 was cloned into pMAL-c4x to express the MBP-PthA4 fusion protein. *CsLOB1* was cloned into pET-41a(+) to express the GST-CsLOB1 fusion protein. MBP-PthA4 and GST-CsLOB1 were expressed separately in *E. coli* BL21 (DE3) by induction with 1.0 mM IPTG. After the proteins were

purified by Glutathione Sepharose 4 Fast Flow (GE Healthcare) and Amylose Resin (New England Biolabs, Ipswich, MA, USA) affinity chromatography, the GST pull-down assays were performed. Each experiment had three replicates.

Western blotting analysis

Western blotting was performed to detect the levels of expression of proteins in *N. benthamiana*. Leaf samples were harvested at 2 dpi. The leaves were ground in liquid nitrogen and extracted in extraction/washing buffer (Roche, Basel, Switzerland). The proteins that were extracted were eluted with 3×Laemmli loading buffer, resolved by 10% SDS-PAGE, and subjected to western blotting using anti-FLAG (Abmart, Shanghai, China), anti-nLUC (Abmart), or anti-cLUC polyclonal antibody (Abmart). Each experiment had three replicates.

Transient overexpression of the *Citrus sinensis* genes

The coding sequences of *Cs9g12620* were cloned into the binary vector pHB to obtain pHB-Cs9g12620. *A. tumefaciens* GV3101 cells were transformed with the resulting recombinant plasmids pHB-Cs9g12620, pHB-pthA4, and pHB-CsLOB1. For transient expression in *C. sinensis*, *A. tumefaciens* cells that harbored different constructs were cultured and grown overnight in YEP media with shaking. The cells were pelleted by centrifugation at 6,000 rpm for 10 min at 4°C, suspended in infiltration medium (10 mM MgCl₂, 10 mM MES, and 200 mM acetosyringone, pH 5.7) to an OD₆₀₀ of 0.01 and gently injected into the young leaves. The phenotypes were scored at 15 dpi. Each experiment had three replicates.

Transmission electron microscopy

To examine any changes in the leaf tissue, the agroinfiltrated *C. sinensis* leaves were sampled 15 dpi for observation by TEM. The samples were fixed using 3% glutaraldehyde in 0.1 M potassium phosphate buffer (pH 7.2) and incubated at room temperature for 4 h. The tissues were washed twice with the same buffer and post-fixed in 2% osmium tetroxide in 0.1 M potassium phosphate buffer at room temperature for 4 h (Etxeberria et al., 2009). The samples were then dehydrated in an ethanol series and embedded in Spurr's resin. Thin sections (80–100 nm thick) were cut with a diamond knife, collected on 200 mesh copper grids, stained with 2% uranyl acetate, and post-stained with lead citrate. Those sections were then examined under an electron microscope (SU8020; Hitachi, Ltd., Tokyo, Japan) in scanning transmission electron microscope mode. Each experiment had three replicates.

Silencing of *Cs9g12620* in the citrus plants

CTV33 is a stable expression vector based on the Citrus tristeza virus (El-Mohtar et al., 2014). A 488 bp DNA fragment of *Cs9g12620* and a 264 bp DNA fragment of *CsLOB1* were cloned separately into the CTV33 vector to generate the gene silencing vector. The CTV virion was prepared in *N. benthamiana* and inoculated in *C. sinensis* using the bark-flap method as previously described (El-Mohtar et al., 2014; Hajeri et al., 2014). Three citrus plants were inoculated for each CTV virion. The plants inoculated with the CTV33 empty vector were used as controls. The infection of CTV was evaluated by the level of expression of the *P23* gene harbored in the CTV33 vector. The RNA silencing efficiency of *Cs9g12620* and *CsLOB1* was evaluated by quantification of the transcript level in the leaves from newly emerging branches. The development of cankers was then assessed by infiltration with *Xcc* 29-1 inoculum of 10⁷ CFU/mL.

Bioinformatics analysis

Sequence similarity searches of the reference *C. sinensis* genome database (<https://www.citrusgenomedb.org/>) were performed using BLAST. Putative promoter sequences were predicted using the online promoter analysis program Neural Network Promoter Prediction (http://www.fruitfly.org/seq_tools/promoter.html). Putative PthA4 EBEs were predicted using DNAMAN

8.0 (Sun et al., 2018 [↗](#)). Putative signal peptides were predicted in PrediSi (<http://www.predisi.de/> [↗](#)) (Hiller et al., 2004 [↗](#)). The conserved domains were identified using the NCBI Conserved Domain Database.

Acknowledgements

We thank Gongyou Cheng and Lifang Zou in Shanghai Jiaotong University (Shanghai, China) for their gifts of Xcc strain 049, TAL-free mutant 049E, and 049E/pthA4. We also thank Dingzhong Tang and Haitao Cui for advice on using the cooled CCD imaging apparatus at the Plant Immune Center in Fujian Agriculture and Forestry University (Fuzhou, China). We would like to thank MogoEdit (<https://www.mogoeedit.com> [↗](#)) for its English editing during the preparation of this manuscript.

Additional information

Competing interests

The authors declare that they have no competing interests.

Funding

Funder	Grant reference number	Author
National Natural Science Foundation of China	31872919	Huasong Zou
National Natural Science Foundation of China	31801696	Xiaoqing Fan
National Natural Science Foundation of China	31701752	Tao Zhuo

References

- Al-Saadi A, Reddy JD, Duan Y, Brunings AM, Yuan Q, Gabriel DW (2007) **All Five Host-Range Variants of *Xanthomonas citri* Carry One *pthA* Homolog With 17.5 Repeats That Determines Pathogenicity on Citrus, but None Determine Host-Range Variation** *Molecular Plant-Microbe Interactions: MPMI* **20**:934–943 <https://doi.org/10.1094/MPMI-20-8-0934>
- Boch J, Scholze H, Schornack S, Landgraf A, Hahn S, Kay S, Lahaye T, Nickstadt A, Bonas U (2009) **Breaking the Code of DNA Binding Specificity of TAL-Type III Effectors** *Science* **326**:1509–432 <https://doi.org/10.4161/viru.1.5.12863>
- Brunings AM, Gabriel DW (2003) ***Xanthomonas citri*: breaking the surface** *Molecular Plant Pathology* **4**:141–157 <https://doi.org/10.1046/j.1364-3703.2003.00163.x>
- Craig JW, Cherry MA, Brady SF (2011) **Long-Chain N-Acyl Amino Acid Synthases Are Linked to the Putative PEP-CTERM/Exosortase Protein-Sorting System in Gram-Negative Bacteria** *Journal of Bacteriology* **193** <https://doi.org/10.1128/JB.05426-11>
- de Souza TA, Soprano AS, de Lira NP, Quaresma AJC, Pauletti BA, Leme AFP, Benedetti CE (2012) **The TAL Effector PthA4 Interacts with Nuclear Factors Involved in RNA-Dependent Processes Including a HMG Protein That Selectively Binds Poly(U) RNA** *Plos One* **7** <https://doi.org/10.1371/journal.pone.0032305>
- Deng D, Yan C, Pan X, Mahfouz M, Wang J, Zhu J, Shi Y, Yan N (2012) **Structural Basis for Sequence-Specific Recognition of DNA by TAL Effectors** *Science* **335**:720–723 <https://doi.org/10.1126/science.1215670>
- Domingues MN, Souza TAD, Cernadas RA, Oliveira MLP, Docena C, Farah CS, Benedetti CE (2010) **The *Xanthomonas citri* effector protein PthA interacts with citrus proteins involved in nuclear transport, protein folding and ubiquitination associated with DNA repair** *Molecular Plant Pathology* **11**:663–675 <https://doi.org/10.1111/j.1364-3703.2010.00636.x>
- Duan S, Jia H, Pang Z, Teper D, White F, Jones J, Zhou C, Wang N (2018) **Functional characterization of the citrus canker susceptibility gene *CsLOB1*** *Molecular Plant Pathology* **19**:1908–1916 <https://doi.org/10.1111/mpp.12667>
- Duan Y P, Castaneda AL, Zhao G, Erdos G, Gabriel DW (2007) **Expression of a Single, Host-Specific, Bacterial Pathogenicity Gene in Plant Cells Elicits Division Enlargement, and Cell Death.** *Molecular Plant-Microbe Interactions: MPMI* **12**:556–560 <https://doi.org/10.1094/MPMI.1999.12.6.556>
- El-Mohtar C, Dawson WO (2014) **Exploring the limits of vector construction based on Citrus tristeza virus** *Virology* **448**:274–283 <https://doi.org/10.1016/j.virol.2013.10.017>
- Etxeberria E, Gonzalez P, Achor D, Albrigo G (2009) **Anatomical distribution of abnormally high levels of starch in HLB-affected Valencia orange trees** *Physiological Molecular Plant Pathology* **74**:76–83 <https://doi.org/10.1016/j.pmpp.2009.09.004>

- Fan X *et al.* (2020) **The LysR-Type Transcriptional Regulator CrgA Negatively Regulates the Flagellar Master Regulator FlhDC in *Ralstonia solanacearum* GM1000** *Journal of Bacteriology* **203**:e00419–20 <https://doi.org/10.1128/JB.00419-20>
- Ge Z, Yan Y, Zhang C, Li S, Yang R, Li Y, Chen G, Zou L (2019) **Functional analysis of *gpd1* gene of *Xanthomonas citri* subsp. *citri* in pathogenesis on citrus** *Acta Phytopathologica Sinica* **49**:670–680 <https://doi.org/10.13926/j.cnki.apps.000392>
- Gottwald TR, Graham JH, Schubert TS (2002) **Citrus Canker: The Pathogen and Its Impact** *Plant Health Progress* **10**:15–48 <https://doi.org/10.1094/PHP-2002-0812-01-RV>
- Graham JH, Gottwald TR, Cubero J, Achor DS (2004) ***Xanthomonas axonopodis* pv. *citri*: factors affecting successful eradication of citrus canker** *Molecular Plant Pathology* **5**:1–15 <https://doi.org/10.1046/j.1364-3703.2004.00197.x>
- Haft DH, Payne SH, Selengut JD (2012) **Archaeosortases and Exosortases Are Widely Distributed Systems Linking Membrane Transit with Posttranslational Modification** *Journal of Bacteriology* **194**:36–48 <https://doi.org/10.1128/JB.06026-11>
- Hajeri S, Killiny N, El-Mohtar C, Dawson WO, Gowda S (2014) **Citrus tristeza virus-based RNAi in citrus plants induces gene silencing in *Diaphorina citri*, a phloem-sap sucking insect vector of citrus greening disease (Huanglongbing)** *Journal of Biotechnology* **176**:42–49 <https://doi.org/10.1016/j.jbiotec.2014.02.010>
- Hiller K, Grote A, Scheer M, Münch R, Jahn D (2004) **PrediSi: prediction of signal peptides and their cleavage positions** *Nucleic Acids Research* **32**:W375–379 <https://doi.org/10.1093/nar/gkh378>
- Hu Y, Zhang J, Jia H, Sosso D, Li T, Frommer WB, Yang B, White FF, Wang N, Jones JB (2014) **Lateral organ boundaries 1 is a disease susceptibility gene for citrus bacterial canker disease** *Proceedings of the National Academy of Sciences* **111**:E521–E529 <https://doi.org/10.1073/pnas.1313271111>
- Husbands A, Bell EM, Shuai B, Smith HMS, Springer PS (2007) **LATERAL ORGAN BOUNDARIES defines a new family of DNA-binding transcription factors and can interact with specific bHLH proteins** *Nucleic Acids Research* **35**:6663–6671 <https://doi.org/10.1093/nar/gkm775>
- Ishihara H, Ponciano G, Leach JE, Tsuyumu S (2003) **Functional analysis of the 3' end of *avrBs3/pthA* genes from two *Xanthomonas* species** *Physiological and Molecular Plant Pathology* **63**:329–338 <https://doi.org/10.1016/j.pmpp.2004.04.002>
- Jalan N, Kumar D, Andrade MO, Yu F, Jones JB, Graham JH, White FF, Setubal JC, Wang N (2013) **Comparative genomic and transcriptome analyses of pathotypes of *Xanthomonas citri* subsp. *citri* provide insights into mechanisms of bacterial virulence and host range** *BMC Genomics* **14** <https://doi.org/10.1186/1471-2164-14-551>
- Jia H, Orbovic V, Jones JB, Wang N (2016) **Modification of the PthA4 effector binding elements in Type I CsLOB1 promoter using Cas9/sgRNA to produce transgenic Duncan grapefruit alleviating XccΔpthA4:dCsLOB1.3 infection** *Plant Biotechnology Journal* **14**:1291–1301 <https://doi.org/10.1111/pbi.12495>
- Kanamori H, Tsuyumu S (1998) **Comparison of Nucleotide Sequences of Canker-forming and Non-canker-forming *pthA* Homologues in *Xanthomonas campestris* pv. *citri*** *Annals of the Phytopathological Society of Japan* **64**:462–470 <https://doi.org/10.3186/jjphytopath.64.462>

- Li Z, Zou L, Ye G, Xiong L, Ji Z, Zakria M, Hong N, Wang G, Chen G (2014) **A Potential Disease Susceptibility Gene CsLOB of Citrus Is Targeted by a Major Virulence Effector PthA of Xanthomonas citri subsp. citri** *Molecular Plant* **7**:912–915 <https://doi.org/10.1093/mp/sst176>
- Majer C, Hochholdinger F (2010) **Defining the boundaries: structure and function of LOB domain proteins** *Trends in Plant Science* **16**:47–52 <https://doi.org/10.1016/j.tplants.2010.09.009>
- Marburg S (2016) **The Burton and Miller Method: Unlocking Another Mystery of Its Coupling Parameter** *Journal of Computational Acoustics* **24**:155001601–155001620 <https://doi.org/10.1142/S0218396X15500162>
- Moscou MJ, Bogdanove AJ (2009) **A Simple Cipher Governs DNA Recognition by TAL Effectors** *Science* **326**:1501–1501 <https://doi.org/10.1126/science.1178817>
- Peng A, Chen S, Lei T, Xu L, He Y, Wu L, Yao L, Zou X (2017) **Engineering canker-resistant plants through CRISPR/Cas9-targeted editing of the susceptibility gene CsLOB1 promoter in citrus** *Plant Biotechnology Journal* **15**:1509–1519 <https://doi.org/10.1111/pbi.12733>
- Song X, Guo J, Ma WX, Ji Z, Zou L, Chen G, Zou H (2015) **Identification of seven novel virulence genes from Xanthomonas citri subsp. citri by Tn5-based random mutagenesis** *Journal of Microbiology* **53**:330–336 <https://doi.org/10.1007/s12275-015-4589-3>
- Soprano AS, Abe VY, Smetana JHC, Benedetti CE (2013) **Citrus MAF1, a Repressor of RNA Polymerase III, Binds the Xanthomonas citri Canker Elicitor PthA4 and Suppresses Citrus Canker Development** *Plant Physiology* **163**:232–242 <https://doi.org/10.1104/pp.113.224642>
- Sun D, Rou W, Zhou Y, Zhuo T, Fan X, Hu X, Zou H (2018) **Ectopic expression of the TAL effector AvrXa7 in Xanthomonas citri subsp. citri hinders citrus canker symptom formation by modulating transcriptional profile of citrus genes** *Biochemical and Biophysical Research Communications* **502**:479–485 <https://doi.org/10.1016/j.bbrc.2018.05.194>
- Sun T, Wu W, Wu H, Rou W, Zhou Y, Zhuo T, Fan X, Hu X, Zou H (2020) **Ralstonia solanacearum elicitor RipX Induces Defense Reaction by Suppressing the Mitochondrial atpA Gene in Host Plant** *International Journal of Molecular Sciences* **21**:2000–2013 <https://doi.org/10.3390/ijms21062000>
- Sun X *et al.* (2004) **Detection and Characterization of a New Strain of Citrus Canker Bacteria from Key/Mexican Lime and Alemow in South Florida** *Plant Disease* **88**:1179–1188 <https://doi.org/10.1094/PDIS.2004.88.11.1179>
- Swarup S, Yang Y, Kingsley MT, Gabriel DW (1992) **An Xanthomonas citri Pathogenicity Gene, pthA, Pleiotropically Encodes Gratuitous Avirulence on Nonhosts** *Molecular Plant-Microbe Interactions: MPMI* **5**:204–213 <https://doi.org/10.1094/MPMI-5-204>
- Wu W, Zhao Z, Luo X, Fan X, Zhuo T, Hu X, Liu J, Zou H (2019) **Response regulator VemR regulates the transcription of flagellar rod gene flgG by interacting with σ 54 factor RpoN2 in Xanthomonas citri subsp. citri** *Molecular plant pathology* **20**:372–381 <https://doi.org/10.1111/mpp.12762>

Yan Q, Wang N (2012) **High-Throughput Screening and Analysis of Genes of *Xanthomonas citri* subsp. *citri* Involved in Citrus Canker Symptom Development** *Molecular Plant-Microbe Interactions: MPMI* **25**:69–84 <https://doi.org/10.1094/MPMI-05-11-0121>

Yang Y, Gabriel DW (1995) **Xanthomonas avirulence/pathogenicity gene family encodes functional plant nuclear targeting signals** *Molecular Plant-Microbe Interactions: MPMI* **8**:627–631 <https://doi.org/10.1094/MPMI-8-0627>

Ye G, Hong N, Zou L, Zou H, Zakria M, Wang G, Chen G (2013) **Population-Based Genetic Diversity of Chinese Isolates of the Citrus Canker Pathogen *Xanthomonas citri* subsp *citri*** *Plant Disease* **97**:1187–1194 <https://doi.org/10.1094/PDIS-12-12-1201-RE>

Ye R, Yao Q, Xu Z, Xue H (2004) **Development of an efficient method for the isolation of factors involved in gene transcription during rice embryo development** *Plant Journal* **38**:348–357 <https://doi.org/10.1111/j.1365-313X.2004.02037.x>

Zhang J, Huguet LTapia JC, Hu Y, Jones J, Wang N, Liu S, White FF (2017) **Homologues of CsLOB1 in citrus function as disease susceptibility genes in citrus canker** *Molecular Plant Pathology* **18**:798–810 <https://doi.org/10.1111/mpp.12441>

Zou X, Du M, Liu Y, Wu L, Xu L, Long Q, Peng A, He Y, Andrade M, Chen S. (2021) **CsLOB1 regulates susceptibility to citrus canker through promoting cell proliferation in citrus** *The Plant Journal* **106**:1039–1057 <https://doi.org/10.1111/tpj.15217>

Editors

Reviewing Editor

Rebecca Bart

The Donald Danforth Plant Science Center, St Louis, United States of America

Senior Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Reviewer #1 (Public Review):

Previous Review:

The authors have identified the predicted EBE of PthA4 in the promoter of Cs9g12620, which is induced by Xcc. The authors identified a homolog of Cs9g12620, which has variations in the promoter region. The authors show PthA4 suppresses Cs9g12620 promoter activity independent of the binding action. The authors also show that CsLOB1 binds to the promoter of Cs9g12620. Interestingly, the authors show that PthA4 interacts with CsLOB1 at protein level. Finally, it shows that Cs9g12620 is important for canker symptoms. Overall, this study has reported some interesting discoveries and the writing is generally well done. However, the discoveries are affected by the reliability of the data and some flaws of the experimental designs.

Here are some major issues:

The authors have demonstrated that Cs9g12620 contains the EBE of PthA4 in the promoter region, to show that PthA4 controls Cs9g12620, the authors need to compare to the wild type Xcc and pthA4 mutant for Cs9g12620 expression. The data in Figure 1 is not enough.

The authors confirmed the interaction between PthA4 and the EBE in the promoter of Cs9g12620 using DNA electrophoretic mobility shift assay (EMSA). However, Fig. 2B is not convincing. The lane without GST-PthA4 also clearly showed mobility shift. For EMSA assay, the authors need also to include non-labeled probe as competitor to verify the specificity. The description of the EMSA in this paper suggests that it was not done properly. It is suggested the authors to redo this EMSA assay following the protocol: Electrophoretic mobility shift assay (EMSA) for detecting protein-nucleic acid interactions PMID: 17703195.

The authors also claimed that PthA4 suppresses the promote activity of Cs9g12620. The data is not convincing and also contradicts with their own data that overexpression of Cs9g12620 causes canker and silencing of it reduces canker considering PthA4 is required for canker development. The authors conducted the assays using transient expression of PthA4. It should be done with Xcc wild type, pthA4 mutant and negative control to inoculate citrus plants to check the expression of Cs9g12620.

Fig. 6 AB is not convincing. There are no apparent differences. The variations shown in B is common in different wild type samples. It is suggested that the authors to conduct transgenic instead of transient overexpression.

Gene silencing data needs more appropriate controls. Fig. D. seems to suggest canker symptoms actually happen for the RNAi treated. The authors need to make sure same amount of Xcc was used for both CTV empty vector and the RNAi. It is suggested a blink test is needed here.

Comments on revised version:

Point 1: Addressed well.

Point 2: The EMSA was reconducted with adding unlabeled DNA, however, the results are still not convincing. Firstly, in fig.3D lane 5, with the absence of unlabeled DNA, the shifted bound signal wasn't reduced significantly. Secondly, still in fig.3D lane 5, the free labeled DNA probe at the bottom of the gel didn't increase. Which together mean that the unlabeled DNA was unable to compete with the labeled DNA and the "bound" shifted bands might not be true positive.

Point 3: The authors didn't address the question clearly regarding the connection between the inhibition of Cs9g12620 promoter by PthA4 and the positive function of Cs9g12620 on citrus canker.

Point 4: The comment was not addressed. Fig.7A and B are not convincing. Firstly, no evidence shows the expression of transiently expressed genes. Secondly, hard to tell the difference in 7A. Thirdly, since CsLOB1 positively regulates Cs9g12620, why expressing of CsLOB1 is unable to cause phenotype, while expression of PthA4 does?

Point 5: addressed.

<https://doi.org/10.7554/eLife.91684.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1:

Point 1: The authors have demonstrated that Cs9g12620 contains the EBE of PthA4 in the promoter region, to show that PthA4 controls Cs9g12620, the authors need to compare

to the wild type Xcc and pthA4 mutant for Cs9g12620 expression. The data in Figure 1 is not enough.

The data in Figure 1 D and E show a pthA4 Tn5 insertion mutant Mxac126-80 and the expression level of Cs9g12620 in citrus inoculated with the pthA4 mutant.

Point 2: The authors confirmed the interaction between PthA4 and the EBE in the promoter of Cs9g12620 using DNA electrophoretic mobility shift assay (EMSA). However, Figure 2B is not convincing. The lane without GST-PthA4 also clearly showed a mobility shift. For the EMSA assay, the authors need also to include a non-labeled probe as a competitor to verify the specificity. The description of the EMSA in this paper suggests that it was not done properly. It is suggested the authors redo this EMSA assay following the protocol: Electrophoretic mobility shift assay (EMSA) for detecting protein-nucleic acid interactions PMID: 17703195.

Thank you very much for your comments. We have re-conducted the EMSA analysis based on your suggestion. The DNA probe was labeled with Cy5, included a non-labeled probe as a competitor. (Figure 3 B and D; Figure 4B and E)

Point 3: The authors also claimed that PthA4 suppresses the promote activity of Cs9g12620. The data is not convincing and also contradicts with their own data that overexpression of Cs9g12620 causes canker and silencing of it reduces canker considering PthA4 is required for canker development. The authors conducted the assays using transient expression of PthA4. It should be done with Xcc wild type, pthA4 mutant, and negative control to inoculate citrus plants to check the expression of Cs9g12620.

We have detected Cs9g12620 expression in silencing citrus plants inoculated wild type Xcc 29-1. (Figure 7F)

Point 4: Figure 6 AB is not convincing. There are no apparent differences. The variations shown in B are common in different wild-type samples. It is suggested that the authors conduct transgenic instead of transient overexpression.

It has been proven that transient expression of PthA4 leads to canker-like phenotype, suggesting that this experiment is effective. However, it will be more confident if conduct transgenic plant overexpressing pthA4 and Cs9g12620. We'll create the plants in our following research to confirm the phenotype.

Point 5: Gene silencing data needs more appropriate controls. Figure D seems to suggest canker symptoms actually happen for the RNAi treated. The authors need to make sure the same amount of Xcc was used for both CTV empty vector and the RNAi. It is suggested a blink test is needed here.

We used the same amount of Xcc to inoculate CTV empty vector and the RNAi. In either inoculation, the cultured Xcc cells were suspended in sterile distilled water to a final concentration of 108 CFU/mL (OD600 = 0.3).

Point 6: Figure 1. Please draw a figure to clearly show the location of the EBE in the promoter of Cs9g12620, including the transcription start site, and translational start site.

The EBE in Cs9g12620 promoter was indicated by underlined in Figure supplement 1. We did not sure about the translation start site, but the translation start site was labelled.

<https://doi.org/10.7554/eLife.91684.2.sa0>