

Identification of novel microcephaly-linked protein ABBA that mediates cortical progenitor cell division and corticogenesis through NEDD9-RhoA

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 1

April 9, 2024 (this version)

Sent for peer review

October 12, 2023

Posted to preprint server

September 14, 2023

Aurelie Carabalona ✉, **Henna Kallo**, **Maryanne Gonzalez**, **Liliia Andriichuk**, **Ellinoora Elomaa**, **Florence Molinari**, **Christiana Fragkou**, **Pekka Lappalainen**, **Marja W Wessels**, **Juha Saarikangas**, **Claudio Rivera** ✉

Aix Marseille Univ, INSERM, INMED, Marseille, France • Neuroscience Center, HiLIFE, University of Helsinki, Helsinki, Finland • Helsinki Institute of Life Science, HiLIFE • Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland • Aix Marseille Univ, Inserm, MMG, Marseille, France • Institute of Biotechnology, HiLIFE, University of Helsinki, Helsinki, Finland • Department of Clinical Genetics, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The cerebral cortex, which is responsible for higher cognitive functions, relies on the coordinated asymmetric division cycles of polarized radial glial progenitor cells for proper development. Defects in the mitotic process of neuronal stem cells have been linked to the underlying causes of microcephaly; however, the exact mechanisms involved are not fully understood. In this study, we present a new discovery regarding the role of the membrane-deforming cytoskeletal regulator protein called Abba (also known as MTSS1L/MTSS2) in cortical development. When Abba was absent in the developing brain, it led to a halt in radial glial cell proliferation, disorganized radial fibers, and abnormal migration of neuronal progenitors. During cell division, Abba localized to the cleavage furrow, where it recruited the scaffolding protein Nedd9, and positively influenced the activity of RhoA, a crucial regulator of cell division. Notably, we identified a variant of Abba (R671W) in a patient with microcephaly and intellectual disability, further highlighting its significance. The introduction of this mutant Abba protein in mice resulted in phenotypic similarities to the effects of Abba knockdown. Overall, these findings offer valuable mechanistic insights into the development of microcephaly and the cerebral cortex by identifying Abba as a novel regulator involved in ensuring the accurate progression of mitosis in neuronal progenitor cells.

eLife assessment

In this **valuable** study, the authors examine the role of the cytoskeletal regulatory protein Abba in governing the process of cell genesis in the developing cortex. This study provides insights into the mechanisms of microcephaly, a developmental malformation. The evidence supporting the study was felt to be **solid**, but the reviewers did note some technical weaknesses that limit the strength of some of the interpretations.

Introduction

Development of the central nervous system occurs by proliferation of neural progenitor cells, migration and differentiation of their progeny over substantial distance. These different steps require extensive membrane remodelling and cytoskeleton dynamics¹. Neuronal migration involves the coordinated extension and adhesion of the leading process along the radial glial scaffold with the forward translocation of the nucleus, which requires regulation of centrosome and microtubule dynamics by different proteins^{1–3}. However, little is known about the molecular mechanisms underlying membrane dynamics during neuronal migration and morphogenesis.

Cell membrane curvature is a micro-morphological change involved in many important cellular processes including endocytosis, exocytosis, and migration⁴. Recent studies demonstrated that members of an extended protein family, characterized by the presence of a membrane binding and deforming BAR (Bin–Amphiphysin–Rvs) domain function at the interface between the actin cytoskeleton and plasma membrane during the formation of membrane protrusions or invaginations^{5–7}. These proteins have been shown to be important for proper neuronal morphogenesis⁸, and can either generate positive membrane curvature to facilitate the formation of plasma membrane invaginations (e.g. BAR, N-BAR and F-BAR domain proteins), or induce negative membrane curvature to promote the formation of plasma membrane protrusions (I-BAR and IF-BAR domain proteins)^{9,10}. An important part of the membrane deforming mechanism include the interaction with phosphatidylinositol-4,5-bisphosphate PI(4,5)P₂-rich membranes through their BAR domain and also with actin cytoskeleton through Wasp Homology-2 domain (WH2).

Regulation of the actin cytoskeleton plays a crucial role in several developmentally important cellular processes including neuronal migration, axonal and dendritic extension and guidance, and dendritic spines formation¹¹. We previously discovered that MIM (Missing-In-Metastasis; Mtss1), a member of the I-BAR protein family, has an important role in bending the dendritic plasma membrane in order to form dendritic spines and sculpt their morphology¹². In a more recent study, a closely related protein Abba (Mtss1L) was found to be important for exercise induced increase synaptic potentiation in the hippocampus¹³. Although an important role of MIM and Abba in synapse formation and plasticity starts to emerge, information about the role of the early expression of this protein family during development is poor.

Mutations in BAR domain proteins are increasingly found in mental retardation disorders. For instance, altered expression and mutation of F-BAR family proteins, SrGAP2 & 3, are known to be associated to neurological disorders (mental retardation, the 3p syndrome, schizophrenia, early infantile epileptic encephalopathy and severe psychomotor delay^{14–18}). Furthermore, previous whole-exome sequencing of pre-screened multiplex consanguineous families identified *ABBA* as a likely candidate linked to neurogenic disorders¹⁹, suggesting that some BAR

containing proteins might play important functions during brain development. Despite Abba's potential role in neuronal development, very little is known about how this protein may lead to neurological pathology.

Regulation of the actin cytoskeleton and plasma membrane dynamics in glial cells of the developing brain has remained poorly understood. Interestingly, the I-BAR protein Abba is the only glia-enriched regulator that has been reported so far²⁰.

We sought to determine the consequences of altered expression of *Abba* in radial glial progenitor (RGP) cells on cell cycle progression and neurogenesis and to test for subsequent neuronal migration. To address these issues, we used *in utero* electroporation to silence *Abba* with shRNA in embryonic mouse brains. Downregulation of Abba's expression had a striking effect on RGP cells progression, resulting in a blocking of cytokinesis and a subsequent apoptosis. Biochemical analysis demonstrated that the cytokinesis block was due to an absence of interaction with the adhesion docking protein named Nedd9 (aka CAS-L or HEF1) that is well known to be a positive regulator of RhoA signaling in mitosis²¹. Using FRET based monitoring of RhoA activity, we also found a significant role of Abba on RhoA activity. Moreover, we observed a defect in neuronal migration associated with an increase in neuronal apoptosis. Importantly, we identified one patient harboring a missense variant in *ABBA* sequence and showing specific features such as: small head circumference, mental retardation, autism spectrum disorder (ASD) and craniofacial dysmorphism. We provide evidence that the (2011C>T(R671W)) patient mutation likely represents a dominant loss-of-function allele.

Overall, our data highlights the critical role of *Abba* in proper cortical development in neurogenesis and migration, and provides the first evidence that *ABBA* is a new causative gene for cortical malformation associated with ASD.

Materials and Methods

Plasmids construction

We utilized a single short hairpin RNA (shRNA) that targeted the coding sequence for Abba²⁰. It is worth noting that the rat and mouse Abba sequences share a high degree of similarity, with a 95% homology. The specific region targeted by the shRNA displayed 100% homology. Both Abba and Nedd9 shRNA were designed using the siRNA Wizard Software (InvivoGen ©). As a negative control, we employed a non-targeting shRNA.

These shRNAs were subcloned into an mU6pro vector²². Additionally, a pCAG-RFP vector was co-injected to facilitate the visualization of fluorescent cells. Abba mouse full length with silent mutations²⁰, ABBA human full length and human ABBA-R671W carrying vectors were made by GeneCust© (France) and subcloned into the pCAGIG-IRES-GFP vector (Addgene, Cambridge, MA, USA).

Bioinformatic methods

ABBA coding sequences were obtained from NCBI. The multiple sequence comparison was done by Log-Expectation (MUSCLE) method and visualized with DNASTAR MegAlign Pro. The impact of ABBA R671W on the structure and function of a human ABBA protein was predicted using PolyPhen-2 and HumVar model²³.

Human subjects

Families were identified through our clinical and research programs, personal communication, as well as the MatchMaker Exchange (MME) including GeneMatcher²⁴ (<http://www.genematcher.org>). Informed consent for publication and analysis of photos, imaging and clinical data was obtained from the patients' legal guardians. Brain magnetic resonance imaging (MRI) studies were performed on the subject and reviewed by the investigators in accordance with Department Ethical permission number (MEC-2012387) to conduct human exome studies.

RhoA activity assay

Fluorescence from Raichu-Rac1 was imaged using a confocal microscope (LSM 710/Axio; Carl Zeiss) controlled by ZEN 2011 software (Carl Zeiss) and equipped with water immersion objective (63×/1.0 NA; Carl Zeiss). Transfected cells were illuminated with an argon laser (Lasos; Lasertechnik GmbH) at 458 nm (CFP and FRET) and 514 nm (YFP) using 0.5–10% of full laser power. Emission was collected at 461–519 nm (CFP) and 519–621 nm (FRET and YFP). The pinhole was fully opened. Scanning was performed in XY mode using 6× digital zoom that resulted in a pixel XY size of 68 × 68 nm. FRET signal was calculated as reported elsewhere²³. In brief, using ImageJ 1.48v software images were background subtracted and FRET image generated using three images: MDonor (CFP excitation and CFP emission filters), MIndirectAcceptor (CFP excitation and YFP emission filters), and MDirectAcceptor (YFP excitation and YFP emission filters), using the following equation: $FRET = (MIndirectAcceptor - MDonor \times \beta - MDirectAcceptor \times (\gamma - \alpha\beta)) / (1 - \beta\delta)$. Coefficients α , β , δ , and γ were obtained by independent control experiments analyzing cells expressing CFP or YFP as described previously²⁵. Coefficients α , δ , and γ were determined in cells expressing only YFP (acceptor) and calculated according to equations: $\alpha = MDonor / MDirectAcceptor$; $\gamma = MIndirectAcceptor / MDirectAcceptor$; $\delta = MDonor / MIndirectAcceptor$. Coefficient β was determined in cells expressing only CFP (donor) and calculated according to the equation: $\beta = MIndirectAcceptor / MDonor$. After generation of FRET image, intensity of FRET signal was calculated using a mask obtained from a corresponding YFP image and covering neuronal cell bodies. The images were also analysed using Pix-Fret²⁶.

In Utero Electroporation

Animal experiments were performed in agreement with European directive 2010/63/UE and received approval N°: APAFIS#2797 from the French Ministry for Research and in agreement with the National Animal Experiment Board, Finland (license number ESAVI/18276/2018). Plasmids were transfected using intraventricular injection followed by *in utero* electroporation^{27,28}. Timed pregnant C57BL6N mice (Janvier Labs; E14; E0 was defined as the day of confirmation of sperm-positive vaginal plug) received buprenorphine (Buprecare, 0.03 mg/kg) and were anesthetized with sevoflurane/isoflurane (4%–4.5% induction, 2% anaesthesia maintenance) 30 minutes later. For pain management, bupivacaine (2 mg/kg) was administered via a subcutaneous injection at the site of the future incision. Uterine horns were exposed, the lateral ventricle of each embryo was injected using pulled glass capillaries and a microinjector (Picospritzer II; General Valve Corporation, Fairfield, NJ, USA) with Fast Green (2 mg/ml; Sigma, St Louis, MO, USA) combined with the different DNA constructs at 1.5 µg/µl or with 1.5 µg/µl shRNA constructs together with 0.5 µg/µl pCAGGS-RFP. Plasmids were further electroporated by discharging a 4000 mF capacitor charged to 50V with a BTX ECM 830 electroporator (BTX Harvard Apparatus, Holliston, MA, USA). The voltage was discharged in five electrical pulses at 950ms intervals via 5mm tweezer-type electrodes laterally pinching the head of each embryo through the uterus. The mice were followed-up post-surgically and given buprenorphine (Buprecare 0.03 mg/kg) if needed. The number of animals has been calculated on the basis of the requirement for adequate numbers of brain slices and sections for sufficient imaging to provide statistically significant data on the effects of RNAi, small molecules, and other reagents used in the proposed analysis, plus controls.

Organotypic cultures and Live Imaging

To assess the impact of altered ABBA expression in the RGP cells on neurogenesis and cell cycle progression, we utilized a combination of *in utero* electroporation and time-lapse imaging of organotypic cultures. The organotypic cultures were prepared as described²⁸ with the following exceptions. After surrounding the brains (E17) with 4.5% UltraPure Low Melting Point Agarose (Invitrogen, 16520-100), 350 µm thick slices were cut with Vibratome 1000 Plus Sectioning System (No. 054018). Before placing the slices on a coated (83 µg/ml Laminin, Sigma L2020; 8,3 µg/ml Poly-L-Lysine, Sigma P4707) cell culture insert (Millicell, PICMORG50), sections were carefully cleaned from agarose to ensure adequate oxygenation. The inserts with slice cultures were transferred into a 6-well plate containing 1 ml of warmed slice culture medium. For live-imaging we used Andor Dragonfly 550 high speed Spinning Disc confocal equipped with Nikon Perfect Focus System, motorized XY stage, and environmental chamber. The system uses Fusion 2.0 software (version 2.1.0.48). The image acquisition was executed using Andor iXon 888 U3 EMCCD camera, 20x magnification with voxel size 0.65 x 0.65 x 0.9524 micron³ (channel/excitation: RFP/561 nm, GFP/488 nm). Time lapse-imaging of organotypic cultures was started at E17+DIV0 with intervals of 10 minutes and total duration of 15 hours. The data was collected from 6 pups (2 scramble, 4 shRNA), 17 slices (5 scramble, 12 shRNA) and 24 areas (9 scramble, 15 shRNA; some slices were recorded from two different areas).

Immunostaining of Brain Slices

Mice brains were fixed (E17) in Antigen Fix (Diapath), Brain slices were sectioned coronally (80µm) on a vibratome (Leica microsystems). Brain slices were washed with PBS (Phosphate-buffered saline pH 7.4) and stained in PBS 0.3% Triton X-100 supplemented with 5% of donkey serum. Primary antibodies were incubated overnight at 4°C, sections were then washed with PBS and incubated in secondary antibodies for 2 hours at room temperature. Antibodies used in this study were: Abba (homemade), Vimentin (Millipore, MAB3400), Ki67 (Millipore, AB9260), phospho-histone H3 (Abcam, ab14955), Pax6 (biolegend, PRB-278P).

Cells culture and transfection

Rat C6 glioma cells were cultured at 37°C under a humidified atmosphere with 5% CO₂ with a complete medium DMEM supplemented with 10% foetal bovine serum (FBS, Sigma) and 100 units/mL antibiotics/antimycotics. Cells were transfected using the Neon® Transfection System (Life Technologies) according to the manufacturer's protocol. Briefly, cells were trypsinized and counted with the cell counter SCEPTER (Millipore). Electroporation was performed with 500 000 cells and a total amount of 5 µg of DNA containing a reporter plasmid encoding RFP (1:5) and shRNAs or mismatch constructs, with the following configuration: 1860 V, 1 pulse, 20 ms. Cells were then cultured on a 6 wells plate during 2 days before RNA extraction.

Immunocytochemistry

HEK293T cells were fixed in Antigen Fix (Diapath), for 15 min and blocked at room temperature for 1 h with 5% normal goat serum, 0.3% Triton X-100 in PBS and incubated overnight at 4°C with a set of antibodies against Abba (homemade), Phalloidin (Thermo Fisher, A12379), Nedd9 (Abcam, ab18056), Hoechst (Thermo Fisher, 62249). Cells were then washed with PBS and incubated in secondary antibodies for 2 hours at room temperature.

RT-PCR and qPCR

Total RNA was isolated from C6 using RNeasy Plus Mini kit and cDNA was synthesized using the Quantitect Reverse Transcription kit, according to the manufacturer's protocol (QIAGEN). Quantitative PCR (qPCR) was performed on a Light cycler 480 using SYBR-Green chemistry (Roche) and specific primers for Abba (F: CGAGACTCGCTGCAGTATTCC and R: CCATTCACAGAGTAGCAGTCG,

113 bp), Nedd9 (Qiagen, QT01610385) and CyclophilinA (Qiagen, QT00177394) as a control probe. qPCR was performed with 5 μ L of diluted cDNA template, specific primers (0.6 μ M) and SYBR Green I Master Mix (7.5 μ L) at a final volume of 15 μ L. Each reaction was performed at an annealing temperature of 60°C and for 50 cycles. Reactions were performed in duplicate and melting-curve analysis was performed to assess the specificity of each amplification. A standard curve was performed for each gene with a control cDNA diluted at different concentrations. Relative expression was assessed with the calculated concentration in respect to the standard. All experiments were performed in triplicate.

Microscopy and Image Analysis

All images were collected with an IX80 laser scanning confocal microscope (LSM 710/Axio; Carl Zeiss) controlled by ZEN 2011 software (Carl Zeiss). Cells & Brain sections were imaged using a 60x 1.42 N.A. oil objective or a 10x 0.40 N.A. air objective. All images were analyzed using ImageJ (NIH, Bethesda, MD). To quantify the total number of C6 cells alive and in mitosis on 12mm diameter coverslips 2 & 3 days after Abba shRNA & control transfections, we have arbitrarily designed 4 identical regions comprised of 9 squares using ZEN software acquisition system and count manually the total number of alive and mitotic cells in each condition. All cell quantifications in brain slices were initially normalized to electroporation efficacy.

Statistical data

Statistical analyses were performed with Prism (GraphPad Software, La Jolla, CA, USA). A two-sample Student's t-test was used to compare means of two independent groups if the distribution of the data was normal. If the values come from a Gaussian distribution (D'agostino-Pearson omnibus normality test) the parametric unpaired t-test with Welch's correction was used. But, when the normality test failed, the non-parametric Mann-Whitney test was used. Significance was accepted at the level of $p < 0.05$.

No statistical methods were used to predetermine sample sizes, but our sample sizes are similar to those generally employed in the field. No randomization was used to collect all the data, but they were quantified blindly.

Yeast two-hybrid screen

Yeast two-hybrid screening was performed by Hybrigenics Services (Paris, France). The Abba coding sequence containing amino acids (aa) 1-715 was cloned into pB27 (N-LexA-bait-C fusion) and pB66 (N-GAL4-bait-C fusion) and sequence verified. These constructs were used as baits to screen the mouse embryo brain RP2 prey library. Total of 91 million (pB27) and 31 (pB66) million interactions were analyzed and the detected interactions were assigned with statistical confidence score, the Predicted Biological Score (PBS) (www.hybrigenics-services.com .

Co-immunoprecipitation

C6 cells or cortical tissue were washed with ice cold PBS and lysed using lysis buffer (20 mM Tris-HCl [pH 8.0], 137 mM NaCl, 10% glycerol, 1% Triton X-100, 2 mM EDTA) with 50 μ g/ml PMSF (Roche) and a protease inhibitor cocktail (Roche). Cell lysate was agitated for 30 min at 4°C and centrifuged at 12000 rpm for 20 min at 4°C. The supernatant was collected and protein concentrations were determined with a Pierce™ BCA Protein Assay kit, according to the manufacturer's protocol (Thermo Scientific).

Before Co-IP, lysates were precleared with washed Pierce Protein A/G Magnetic Beads (Thermo Scientific) overnight in gentle shaking at 4°C, to remove nonspecific binding. Beads used for Co-IP were washed and pre-blocked with 3% BSA in PBS overnight in gentle shaking at 4 °C. Washing steps were done using a wash buffer (1% BSA, 150 mM NaCl, 0.1% Tween-20 in TBS).

For Co-IP, 1000 µg of protein from precleared lysate was incubated with 20 µl of specific primary antibodies (Abba: homemade; Nedd9: Abcam, ab18056) overnight in gentle shaking at 4°C. Negative controls were incubated without primary antibodies under the same conditions. Immunocomplexes were pulled down by incubating the antigen-antibody mixture with pre-blocked beads for 1 h in gentle shaking at room temperature. Beads were washed 3 times with wash buffer and once with purified water before elution. Bound proteins were eluted from the beads by incubating them in 1 x Laemmli buffer containing 150 mM NaCl for 10 min at room temperature.

Original C6 cell lysate and immunoprecipitated proteins were resolved by SDS-PAGE (4-20% Mini-PROTEAN TGX Gel; Bio-Rad) and transferred using Trans-Blot Turbo Mini PVDF Transfer Packs and Transfer System (Bio-Rad). Membranes were blocked with TBS containing 0.1% Tween-20 and 5% milk and blotted with primary antibodies overnight in shaking at 4°C. HRP-conjugated secondary antibodies used in this study were Rabbit anti-mouse IgG and Goat anti-rabbit IgG (Invitrogen). Protein bands were visualized using the ChemiDoc™ MP imaging system (Bio-Rad).

Flow-cytometry

To measure cell cycle length, 4×10⁵ cells were seeded in 6-well plates, transfected and cultured for 24h. Plasmid transfection was performed with jetPEI by following the Batch HTS protocol suggested in the data sheet of the product. Briefly, transfection complexes, which consist of both DNA and jetPEI reagent diluted in NaCl, is mixed with the suspension of cells obtained after trypsinization and distributed in the wells for culture over 24h. Then, cell cycle synchronization was induced by 24h serum starvation and re-started using fetal calf serum supplemented DMEM. The cells were detached with trypsin+EDTA and fixed with 3.7% formaldehyde after 24 hours in culture. 5×10⁵ fixed cells from each stop point were stained in 200ul of Hoechst 33342 (1:2000 dilution) for 30min, washed with PBS 1X and diluted in flow buffer. Acquisition of 100 000 total events was performed in NovoCyte Quanteon 4025 and the analysis performed in (FlowJo v10.8 or NovoExpress 1.6.1) software. Graphs and statistical analysis were performed in GraphPad Prism 5.

Results

Abba knockdown disrupted radial glia morphology

Previous studies of the expression pattern of *Abba* in the developing brain has disclosed a conspicuous expression in radial glia. In order to investigate the function of *Abba* in the developing brain, we used a mouse *in utero* RNAi approach to knockdown *Abba* at E14. For this purpose, we opted for shRNA knockdown using a previously validated targeting sequence²⁰. Revalidation of the knockdown efficiency of this target with three different *Abba*-shRNA (*Abba*-shRNA1-3) in rat C6 glioma cells showed a 70– 80% reduction of *Abba* mRNA and protein levels with *Abba*-shRNA3 (Supplemental Figure 1, *Abba*-shRNA3).

To determine whether knockdown of *Abba* expression alters neuronal migration, *Abba*-shRNA3 combined with a red fluorescent protein (RFP) construct was introduced into neural progenitor cells of mouse neocortex by *in utero* electroporation at E14 and evaluated at E16-18 (**Figure 1A**). Electroporation of E14 targets the progenitors that will migrate and populate cortical layers II/III neurons. In E17 brain sections, we observed that some neurons previously electroporated at 3 days earlier with the scrambled construct reached the cortical plate as expected (**Figure 1B**). However, *in utero* expression of *Abba*-shRNA3 induced a significant arrest of cells within the ventricular zone (VZ) (**Figure 1A-C**; scramble: 24.3 ± 6.4%; n=9, *Abba*-shRNA3: 63.7 ± 2.9%; p<0.0001; n=8). Surprisingly, we also observed a striking decrease in the total number of RFP positive cells in brain sections electroporated with *Abba*-shRNA3 at E17 and E18 (**Figure 1D-E**);

scramble: $969,2 \pm 109,0$ cells/mm²; n=9, Abba-shRNA3: $311,3 \pm 28,61$ cells/mm²; p<0.0001; n=8). These results indicate an important role of Abba in the proliferation and migration of cortical neuronal progenitors.

We also evaluated the impact of *Abba* knockdown on radial glial morphology by imaging vimentin-stained *Abba* knockdown cells at 3 days post-electroporation. All E17 *Abba* knockdown brains, but not in age-matched controls, displayed disruption of radial glial fibers in the VZ/SVZ (**Figure 1D** and **F**). Control vimentin-positive cells displayed the characteristic features and polarized morphology of radial glia with processes extending from the VZ to the pial surface. In contrast, in *Abba* knockdown brains, vimentin-positive cells exhibited misoriented apical processes, detached end feet in the ventricular surface (**Figure 1D**) and decreased number of radial glial progenitor cells in the VZ (**Figure 1F**; scramble: $126,5 \pm 13,4$ cells, n=8; Abba-shRNA3: $65,1 \pm 26,2$ cells, n=7; p=0.0033). All together these data show that the loss of Abba can disrupt radial glial cells morphology thus affecting their progenitor function.

These data suggest that *Abba* is required for radial glial organization, neuronal migration, regulation of neuronal proliferation and cell survival. Importantly, these results are in accord with the reported association of *ABBA* with developmental neurological diseases including intellectual disability and microcephaly¹⁹.

ABBA is required for cell cycle progression of radial glia progenitors

Development of the cerebral cortex occurs through a series of stages, beginning with RGP. These stem cells exhibit an unusual form of cell-cycle-dependent nuclear oscillation between the apical and the basal regions of the ventricular zone, known as interkinetic nuclear migration (INM). RGP cells undergo mitosis only when they reach the SVZ^{30,31}. It has been shown that inhibition of INM disrupts RGP cell cycle progression^{32–34}.

To test the effect of altered radial glial cell morphology on cell cycle progression, we have quantified the nuclear distance from the VS for RGP cells after electroporation of scrambled or Abba RNAi (**Figure 2A**). We observed an accumulation of Abba-shRNA3 RGP cells close to the ventricular surface (0-10 mm; scramble: $30,2 \pm 3,2\%$, n=7; Abba-shRNA3: $38,3 \pm 2,3\%$, n=7; p=0,000156) which can correspond to mitotic cell, but also further away (>30 mm; scramble $12,0 \pm 3,3\%$, n=7; Abba-shRNA3: $21,9 \pm 2,4\%$, n=7; p=0,000023) which support our previous observations on cells detaching from the VS (**Figure 1D** and **F**). Furthermore, we observed that Abba-shRNA3 decrease the percentage of the RGP cells marker Pax6 (**Figure 2B** and **C**; scramble: $31,9 \pm 0,6\%$, n=8; Abba-shRNA3: $21,1 \pm 0,6\%$, n=5; p<0.0001). We have then tested for cell cycle effects, and found a decrease on the percentage of Abba shRNA3-expressing RGP cells positive for the cell cycle marker Ki67 (**Figure 2D** and **E**; scramble: $46,5 \pm 0,8\%$, n=7; Abba-shRNA3: $23,7 \pm 1,369\%$, n=5; p= <0.0001) and also for the late G2/M phases marker phospho-histone 3 (PH3, **Figure 2 F** and **G**; scramble: $4,5 \pm 0,1764\%$, n=9; Abba-shRNA3: $1,264 \pm 0,5191\%$, n=5; p= <0.0001). More detailed examination of the impact of Abba-shRNA3 on cell cycle progression using flow cytometry showed an accumulation in S phase. Interestingly, these effect was rescued by overexpression of Abba-FL (Supplemental fig 2 G1: scramble: $45,2 \pm 1,4 \%$, Abba-shRNA3: $37,4 \pm 1,0 \%$, Abba-shRNA3+Abba-FL: $42,8 \pm 1,3 \%$; S: scramble: $34,3 \pm 2,8 \%$, Abba-shRNA3: $48,2 \pm 2,7 \%$, Abba-shRNA3+Abba-FL: $38,9 \pm 3,4 \%$; G2: scramble: $17,1 \pm 2,1 \%$, Abba-shRNA3: $14,4 \pm 2,7\%$, Abba-shRNA3+Abba-FL: $15,9 \pm 1,9 \%$; n=10; p<0.001). These results indicate that Abba plays a major role in coordinating cell cycle progression in radial glia.




Figure 1.

***In utero* knockdown of *Abba* expression is linked to radial glial disruption and alters neuronal migration.**

A. Representative coronal sections of E16-E18 mice brains electroporated at E14 with either scramble or Abba-shRNA3 and immunostained for vimentin (in blue). In Abba-shRNA3 brain sections, vimentin staining revealed a marked disruption of radial glial apical fibers. **B.** Representative neocortical coronal sections showing migration of transfected cells 3 days after electroporation at E14 with either scramble or Abba-shRNA. **C.** Quantification of RFP-positive cell distribution in the cortex at E17 (VZ/SVZ: scramble $24.33 \pm 2.13\%$, Abba-shRNA3 $63.71 \pm 1.03\%$; IZ: scramble $39.80 \pm 2.22\%$, Abba-shRNA3 $29.71 \pm 0.85\%$; CP: scramble $35.86 \pm 1.73\%$, Abba-shRNA3 $6.58 \pm 0.53\%$; $n=9$ scramble and 8 Abba-shRNA3; (VZ/SVZ: $p < 0.0001$; IZ: $p = 0.0002$; CP: $p < 0.0001$) showing a significant increase of Abba knockdown cells in SVZ/lower IZ. **D.** Higher magnifications of regions indicated by a square in C. **E.** Total number of RFP positive cells in mouse neocortex at E17 (scramble $969.2 \pm 109.0\%$; Abba-shRNA3 $311.3 \pm 28.61\%$; $n=9$ scramble and 8 Abba-shRNA3) showing a striking reduction of RFP positive cells in Abba knockdown electroporated brains. Error bars represent mean $\pm$ s.d. $^{**}P < 0.002$, $^{***}P < 0.001$, Scale bars $100\mu\text{m}$ (A). **F.** Quantification of radial glial progenitor cells at the VZ (scramble $126.50 \pm 13.36\%$, Abba-shRNA3 $65.14 \pm 9.91\%$). mean $\pm$ SD. VZ: ventricular Zone, IZ: Intermediate Zone, CP: Cortical Plate. Error bars represent mean $\pm$ s.d. $^{**}P < 0.002$. Scale bars represent $100\mu\text{m}$ (A), $50\mu\text{m}$ (B).




Figure 2.

Abba down-regulation induced inhibition of basal nuclear migration in RGP cells blocks cell cycle progression.

A. Quantification of the nuclear distance from the VS for RGP cells after electroporation of scrambled or Abba-shRNA3. Nuclear distribution of RFP RGP cells was significantly altered after electroporation of Abba-shRNA3 (0-10: scramble, 30.20 ± 1.21%; Abba-shRNA3, 38.34 ± 0.89%; 10-20: scramble, 34.35 ± 0.69%; Abba-shRNA3, 22.80 ± 1.03%; 20-30: scramble, 23.42 ± 1.00%; Abba-shRNA3, 17.00 ± 1.10%; >30: scramble, 12.04 ± 1.27%; Abba-shRNA3, 21.85 ± 0.74%; (0-10: P = 0.000156; 10-20: P = <0.000001; 20-30: P = 0.001030; >30: P = 0.000023). **B-C.** E14 mice brains were subjected to *in utero* electroporation with either scramble or Abba-shRNA3. Brains were then fixed at E17 and stained with the RGP cell marker Pax6. We observe an important decrease of Pax6 RGP cells expressing Abba-shRNA3 (scramble, 31.87 ± 0.57%; Abba-shRNA3, 21.08 ± 0.63%; n = 8 scramble; 5 Abba-shRNA3). **D-G.** E14 mice brains were subjected to *in utero* electroporation with either scramble or Abba-shRNA3. Brains were then fixed at E17 and stained with the cell cycle marker Ki67 (D) and PH3 (F). We observe a striking decrease in the percent of cycling (E: Ki67; scramble, 46.54 ± 0.79%, n = 7; Abba-shRNA3, 23.71 ± 1.37%, n = 5 Abba-shRNA3) and but not in mitotic (G: PH3; scramble, 4.50 ± 0.17%; Abba-shRNA3, 4.25 ± 0.65%) cells with low expression of Abba. **G.** Flow cytometry analysis of cells electroporated with Abba-shRNA3 show accumulation in S-phase and recovery after additional expression of Abba-FL (**G1**: scramble 45.2 ± 1.4 %, Abba-shRNA3 37.4 ± 1.0 %, Abba-shRNA3+Abba-FL 42.8 ± 1.3 %; **S**: scramble 34.3 ± 2.8 %, Abba-shRNA3 48.2 ± 2.7 %, Abba-shRNA3+Abba-FL 38.9 ± 3.4; **G2**: scramble 17.1 ± 2.1 %, Abba-shRNA3 14.4 ± 2.7 %, Abba-shRNA3+Abba-FL 15.9 ± 1.9 %; n = 10). Error bars represent mean ± s.d. **P < 0.002, ***P < 0.001. Scale bar represents 20µm.

Abba localizes to the cleavage furrow and is important for cytokinesis

To further characterize the role of Abba in mitotic progression, we synchronized C6 rat glioma cell line and stained with Abba antibody. We found that Abba localized to the putative apical membrane initiation site and cytokinetic bridge, proximal to the contractile actin ring during late telophase and cytokinesis (**Figure 3A** and **B** respectively). The actomyosin contractile ring is formed at the plasma membrane in response to accumulation of phosphoinositide 4,5 biphosphate, PI(4,5)P₂, to generate constricting force to separate the two daughter cells³⁵. Abba is known to interact with PI(4,5)P₂ via its membrane deforming I-BAR domain²⁰ that has the capacity to deform and recognise the membrane curvature.

To determine the impact of the loss of Abba on cytokinesis, we quantified the number of C6 cells in cytokinesis 3 days after transfection with Abba-shRNA3 or scramble shRNA. Importantly we found a two-and-a-half-fold increase in the percentage of Abba-shRNA3-expressing cells in cytokinesis (**Figure 3B** and **D**; scramble 0,8285 ± 0,1161%, n=12; Abba-shRNA3 2,078 ± 0,2682%, n=12; p=0,0358), suggesting a prominent role of Abba in cytokinesis. To validate this *in vivo*, we performed live imaging in cortical organotypic slices from animals that were *in utero* electroporated with either Abba-shRNA3 or scramble vectors. Analysis of live imaging recording showed a strong mitotic block at the ventricular surface (**Figure 3C**). We then examined the impact of cytokinesis block on cell survival and found a lower total number of cells 4 days after transfection with Abba-shRNA3 compared with control scramble (**Figure 3E**; scramble: 380,3 ± 48,16%, n=12; Abba-shRNA3: 155,8 ± 22,32%, n=12; p<0.0001). Taken together, these data provide evidence that Abba is required for the completion of mitosis, and its absence results in a reduced number of cell progeny.

Abba recruits Nedd9 to the cytokinetic bridge and is required for RhoA activation

To investigate the mechanism by which Abba operates in mitosis, we performed a yeast-two-hybrid (Y2H) screen in E10.5-E12.5 mouse brain embryo library using full length mouse Abba as a bait. The resulting analysis identified three high (PBS: B) / good (PBS: C) confidence interactors: the E3 ubiquitin ligase Beta-Trcp2 (PBS: B), scaffolding protein Nedd9 (PBS: C), the transcription factor Otx2 (PBS: C) (**Figure 4A**; Supplemental table 2). Among these candidate proteins, Nedd9 (aka HEF-1 or Cas-L) caught our interest as it was previously reported in breast cancer cells to localize to the cleavage furrow during cytokinesis where it activates RhoA, as well as abnormal expression of Nedd9 results in cytokinesis defect^{21,36,37} (**Figure 4B**).

We first tested whether Abba and Nedd9 physically interact, as suggested by the Y2H screen. To this end, we performed immunoprecipitation experiments from embryonic day 18 cortical tissue as well as C6 cells using anti-Nedd9 as well as anti-Abba antibodies. The precipitated Nedd9-Abba complex was separated on SDS-page and followed by western blot with anti-Abba and anti-Nedd9 antibodies respectively. Indeed, Abba precipitated with Nedd9 in both directions, but not using control beads from brain cortex homogenates (**Figure 4C**) and C6 cells (Supplemental figure 3). Nedd9 stability is not affected by the availability of Abba as western blots show no changes in Nedd9 expression levels in Abba-shRNA3 expressing cells (data not shown). Interestingly however, when we analyzed the localization of Nedd9, we found that Abba was required for targeting Nedd9 to the cleavage furrow/cytokinetic bridge (**Figure 4D**). These data suggest that Abba recruits Nedd9 to the cleavage furrow during RGP cells division, which fits to its role in sensing curved PI(4,5)P₂ rich membranes. Nedd9 plays a role in RGP cytokinesis *in vivo*^{21,36,37}. To knockdown Nedd9, we used shRNA targeting Nedd9 which reduced mRNA levels by 75% in C6 cells (Supplemental Figure 3A). We used a previously characterised Nedd9-shRNA to determine




Figure 3.

Abba down-regulation blocks cytokinesis

A. Cellular distribution of Abba in C6 cells during cell division. Upper panel shows two example immunofluorescent microscopy images of Abba (red). Left most figure shows the projection images at the cross line of the image. Mid and lower panels show the distribution of Abba at different cell division stages in relation to β -tubulin (green) aggregation **B.** Immunofluorescence microscopy images from C6 cells 72 hours after transfection with Abba-shRNA3 shows an absence of Abba expression at cytokinesis. **C.** Brain slices prepared from in utero electroporated animals with either Abba-shRNA or scramble were cultured at E17 and subjected to live imaging for a duration of 15–20 hours. The length of the time lapse was adjusted as necessary to capture significant events during interkinetic nuclear migration (INM). In the case of the scramble group, an RGP cell underwent a single mitotic event at the ventricular surface of the brain slice, observed at timepoints 1:50 and 2:20. At timepoint 3:00, two cells can be observed (indicated by arrows), along with the presence of basal fibers (indicated by arrowheads). Conversely, in the Abba knockdown group, the nucleus of the RGP (indicated by an arrow) initially underwent apical INM and subsequently divided at 8:30. Remarkably, the resulting daughter cells remained at the ventricular surface for at least 7 hours, suggesting that the absence of Abba hinders cells from exiting the mitotic state. **D.** Abba-shRNA3-expressing cells show increase in cytokinesis (scramble $0,8285 \pm 0,1161\%$, $n=12$; Abba-shRNA3 $2,078 \pm 0,2682\%$, $n=12$) **E.** Quantification of the total number of RFP positive cells in the neocortex illustrating the impact of cytokinesis block on cell survival. We observed a lower total number of cells 4 days after transfection with Abba-shRNA3 compared with control scramble (scramble $380,3 \pm 48,16\%$; $n=12$, Abba-shRNA3 $155,8 \pm 22,32\%$; $p < 0.0001$; $n=12$). Error bars represent mean $\pm$ s.d. * $P < 0.03$, ** $P < 0.002$, *** $P < 0.001$. Scale bars $50\mu\text{m}$ (A), $20\mu\text{m}$ (D).




Figure 4.

Role of Abba-Nedd9 complex in RGP cells division

A. Good and high confidence interactions detected in yeast two-hybrid screen performed in mouse brain embryo library using full length mouse Abba as a bait. **B.** Schematic representation of the Abba and Nedd9 proteins. **C.** E18 cortical homogenates were subjected to a direct pulldown assay with endogenous Abba and Nedd9. Nedd9 co-immunoprecipitated with Abba, but not in control pulldowns showing the specificity of the interaction. **D.** immunocytochemistry on C6 cells 72 hours after transfection with Abba-shRNA3 shows that lack of Abba did not decrease Nedd9 expression (green) but rather its localization. Scale bars 20µm. **E.** Representative coronal sections of E17 mice brains transfected at E14 with Nedd9-shRNA and immunostained for vimentin (in blue). Vimentin staining revealed a marked disruption of radial glial apical fibers. **F.** Quantification of RFP-positive cell distribution in the cortex at E17 (VZ/SVZ: scramble 25 ± 2.30%, Nedd9-shRNA 46.92±2.78%; IZ: scramble 38.94±2.32%, Nedd9-shRNA 37.37±2.96%; CP: scramble 36.06±1.95%; Nedd9-shRNA 15.71±2.25%; n=8 scramble and 7 Nedd9-shRNA; (VZ/SVZ: $p<0.0001$; IZ: $p=0.0002$; CP: $p<0.0001$) showing a significant increase of Nedd9 knockdown cells in SVZ/lower IZ. Scale bars 100µm (E), 50µm (D).

whether knockdown of Nedd9 expression alters RGP cells morphology and function. To this end, Nedd9-shRNA combined with a red fluorescent protein (RFP) construct was introduced into neural progenitor cells of mouse neocortex by *in utero* electroporation at E14 and stained with the radial glial cell marker, vimentin. Interestingly, similarly to Abba knockdown, Nedd9 downregulation resulted in disruption of INM as a larger portion of progenitors accumulated close to the ventricular surface (**Figure 4E-F**; scramble: $25 \pm 2.30\%$ n=8, Nedd9-shRNA: $46.92 \pm 2.78\%$; n=7; $p < 0.0001$). Using the late G2/M phases marker, PH3, we observed similar results as with Abba shRNA and point mutation no difference in number of PH3 positive cells (Supplemental figure 3C; scramble: $4.50 \pm 0.18\%$, n=9; Nedd9-shRNA: $5.07 \pm 0.35\%$, n=6, $P=0.1135$). The role of Nedd9 in cell cycle progression is consistent with previous published results^{21,36,37} and provide evidence that both Abba and Nedd9 function to ensure correct mitotic completion of RGP cells.

Finally, to determine whether Abba is important for Nedd9 activity and how Abba and Nedd9 converge to regulate the cell-cycle, we assessed the activity of RhoA, a downstream signalling molecule of Nedd9³⁸ which is important for cytokinetic actomyosin ring assembly and dynamics³⁸⁻⁴⁰. To determine RhoA activity we used a GFP-based FRET construct for RhoA designated “Ras and interacting protein chimeric unit” Raichu-RhoA. In order to test the efficacy of the assay we used Calpeptin, a RhoA activator which, as expected, increased RhoA activity (% change compared to Raichu-RhoA: Calpeptin 132.2 ± 3.2 n=33; $p=0.002$) confirming the functionality of our assay (**Figure 5**). Importantly, when Abba was silenced by shRNA there was a significant 31,5% decrease in Raichu-RhoA activity as compared to the control shRNA treated cells (**Figure 5B**; scramble: $103.1.03 \pm 5.3$; n=102, Abba-shRNA3: 68.5 ± 2.3 ; n=46; $p < 0.0001$). In addition, this effect was rescued by the co-expression of shRNA insensitive plasmid carrying the full-length sequence of Abba (Abba-shRNA + Abba-FL 139.8 ± 19.6 n=26; $p < 0.0001$ compared to Abba-shRNA3). Abba-FL did not change RhoA activity significantly compare to scramble expressing cells (Abba-FL 105.2 ± 16.0 ; n=37). These data provide support for the function of Abba in spatial assembly of cytokinetic signaling through the Nedd9-RhoA axis.

Identification of *ABBA* missense variant R671W in an individual with neurodevelopmental syndrome

Mutations in genes effecting mitotic progression of neuronal progenitors have been linked to neurodevelopmental disorders⁴¹. Importantly, we identified a patient carrying a novel heterozygous missense variant in *ABBA* coding *MTSS1/MTSS2* gene. This mutation (2011C>T(R671W)) affects residues that are conserved from Zebrafish to Human and reside in exon 15 (**Figure 6A** and **B**). Consistent with a *de novo* origin of the variant, it was not found in the parents of the affected individual or in normal controls. This patient presents some characteristic craniofacial dysmorphism as mild deep-set eyes, thickened helices, synophrys, narrow forehead, bitemporal narrowing, small bilateral epicanthal folds, upslanting palpebral fissures, medial flaring of eyebrows, flat midface, V-shaped palate, and dysmorphic ears. Brain magnetic resonance imaging sequences showed the presence of a microcephalic brain associated with a smaller head linked to mild IQ. The patient also suffered from Attention-Deficit/Hyperactivity Disorder (ADHD) (**Figure 6C**, Supplemental Table 1). Moreover, a recent publications found six additional patients carrying the same *ABBA* variant that shared similar brain malformation^{42,43}. This variant replaces an arginine to tryptophan that has a large aromatic group that could lead to destabilization of a tertiary structure. Indeed, bioinformatic prediction using AlphaFold predicts (pLDDT) with high 84.4 confidence score that R671 is located in an alpha helical fold close to the C-terminal end of the protein⁴⁴. Furthermore, analysis using Poly-Phen-2 structure and sequence-based algorithm to predict the impact of R671W substitution on the structure and function of *ABBA*, we found a high probability that the mutation is damaging for the protein function²³ (Naïve bayes probability score 0.952) (**Figure 6D**).




Figure 5.

Abba downregulation decreases RhoA activity.

A. Confocal images were captured of C6 cells 72 hours post-transfection with the Raichu-RhoA vector for monitoring RhoA activity. Six experimental groups were examined: transfected with scramble control, Abba-shRNA3, Abba-fl, rescue of Abba-shRNA3 with Abba-fl and expression of the mutant Abba-R671W. In addition to testing the sensitivity of the assay to RhoA activation by calpeptin. **B.** Quantification of FRET density (Raichu-RhoA activity) showed a decreased of Raichu-RhoA activity in C6 cells transfected with Abba-shRNA3 as compared to transfected with the scramble (scramble: $103,1.03 \pm 5.3$; $n=102$, Abba-shRNA3: $68,5 \pm 2,3$; $n=46$; $p<0.0001$, Abba-shRNA + ABBA-FL: $139,8 \pm 19,6$ $n=26$, Abba-FL: $105,2 \pm 16,0$; $n=37$, Abba-R671W: $142 \pm 17,9$ $n=15$). Error bars represent mean $\pm$ s.d. ** $P < 0.002$, *** $P < 0.001$. Scale bar 25 μ m (**A**).

Expression of R671W Human variant results in defects in mitosis and neuronal migration

To test the effects of the human missense variant in mouse brain development, we introduced the human *ABBA* cDNA encoding either wild type or patient mutation R671W into developing mouse brains by *in utero* electroporation at E14 followed by imaging and quantification of targeted cell at E17. Interestingly, whereas wild type *ABBA* (*ABBA*-FL) expression showed no defects in radial glial morphology or neuronal migration, the R671W mutant expression resulted in disorganized radial glial fibers and accumulation of multipolar neurons in the SVZ/lower IZ with very few bipolar neurons (**Figure 6E**). When analyzing the distribution of electroporated neuron progenitors, we observed clear accumulation in the VZ/SVZ (**Figure 6F**; *ABBA*-FL $18.67 \pm 1.44\%$; $n=8$, *ABBA*-R671W $39.77 \pm 2.73\%$ $n=6$; $P=0.0001$). We also found a significant increase in RGP nuclei at the VS of the brain (**Figure 6G**, C *ABBA*-FL $32.56 \pm 2.58\%$; $n=8$, *ABBA*-R671W $50.04 \pm 2.04\%$; $n=6$; $p=0.000001$). Interestingly, no difference in accumulation of PH3+ cells was found (Supplemental Figure 3 B and C; *Abba*-human-WT $4.50 \pm 0.17\%$; $n=9$, *Abba*-human-mutation $4.71 \pm 1.98\%$; $p=0.76$; $n=5$), indicative of a dysfunction in neuronal maturation but a less prominent impact on cell division. In line with these results the effect of *ABBA*-R671W on RhoA activation was not significantly different to scramble expressing cells (142 ± 17.9 $n=15$). The data presented indicate that the abnormal expression of the R671W mutation, rather than the normal protein, leads to abnormal development of the developing cortex. These findings offer evidence suggesting a potential connection between the missense mutation in *ABBA* discovered in the patient and abnormal brain development. However, it is unlikely that the mechanism behind this phenotype is a result of *ABBA*-R671W acting in a dominant-negative manner on cell cycle progression.

Discussion

Abba was initially discovered as a novel regulator of actin and plasma membrane dynamics, specifically in radial glial C6 cells. However, its role in brain radial glial progenitor (RGP) cells in live organisms has remained unclear. In this study, we observed that decreased expression of *Abba* in radial glia within the cortical ventricular zone leads to impaired radial migration and disrupted organization of radial glia. Notably, knock-down of *Abba* specifically hindered cytokinesis in RGP cells, resulting in abnormal cell morphology, mitosis, and reduced numbers of glial and neuronal cells. The underlying mechanism behind this effect appears to involve impaired signaling through the actin regulatory protein RhoA, likely through its interaction with Nedd9 during the early and late stages of telophase. In accordance with these findings, we identified a missense variant in a patient with microcephaly and intellectual disability exhibiting similar clinical characteristics. Furthermore, overexpression of the mutant protein in cortical progenitors of mice resulted in phenotypes resembling, although not identical to, the down-regulation of *Abba* in the cortex. The results presented in this study highlight the critical importance of *Abba* in cortical development and contribute to elucidating the potential mechanism underlying abnormal brain development associated with human variants of *ABBA*.

Radial glial exiting from cytokinesis requires *Abba*

BAR proteins can interact with PI(4,5)P₂-rich membranes through their BAR domain, and several studies indicate that PI(4,5)P₂ play a central role in cytokinesis. Indeed, it has been reported that in mammalian cells PI(4,5)P₂ accumulates at the cleavage furrow and recruits membrane proteins required for stability of the furrow because of a role in adhesion between the contractile ring and the plasma membrane. Moreover, overexpression of proteins that bind to PI(4,5)P₂ perturbs cytokinesis completion, by interfering with adhesion of the plasma membrane to the contractile ring at the furrow, but not ingression of the cleavage furrow^{45,46}. Immunocytochemistry of

Abba shows a strong accumulation at the cleavage furrow. In addition, in absence of Abba, we observed an accumulation of cells blocked in cytokinesis (**Figure 3**). These observations and previous findings showing that Abba can interact with PI(4,5)P₂-rich membranes through its BAR domain as well as with the actin cytoskeleton through Wasp Homology-2 domain (WH2), lead us to suspect that Abba could link the actin contractile ring to the plasma membrane and participate in the mechanism for the invagination of the nuclear membrane during cytokinesis.

Abba interacts with Nedd9 to regulate RhoA activity

Here we have found that Abba interacts directly with Nedd9 and down regulation of Abba leads to significant changes in RhoA activation. Nedd9 is a focal adhesion protein involved in mitotic entry and cleavage furrow ingression⁴⁷. Similar to the results found in this study, previous results have shown that cells with abnormal expression of Nedd9 remain in G1 subsequently leading to cell apoptosis³⁷. This is consistent with the view that Nedd9 triggers cells to enter mitosis. Nedd9 is known to be a positive regulator of RhoA signaling in mitosis. Indeed, activation of RhoA is required for cell rounding at early stages of mitosis and cleavage furrow ingression, and deactivation of RhoA is essential for cell abscission²¹. Moreover, it has been shown that Nedd9 overexpression blocks cells at abscission by maintaining levels of RhoA activation, and decreased Nedd9 expression inhibits cleavage furrow ingression by non-activated RhoA pathway. In light of these results, we hypothesized that during cytokinesis at the cleavage furrow the membrane composition changes and is enriched in PI(4,5)P₂ leading to Abba recruitment and its interaction with Nedd9 which is essential to activate the RhoA pathway cleavage furrow ingression.

Role of ABBA in neuronal migration and microcephaly

The proper formation of the cerebral cortex relies on the sequential generation of cortical layers, which is highly dependent on the normal production, survival, and migration of neuronal progenitors. Abnormalities in these processes are associated with pathological conditions such as microcephaly, leading to impaired cognitive development. Interestingly, both downregulation of *Abba* and overexpression of the *ABBA* point mutation construct result in a similar accumulation of neuronal progenitors in the ventricular zone (VZ) and subventricular zone (SVZ). This may be attributed to abnormal proliferation of neuronal progenitors caused by disrupted interaction with critical proteins involved in cell cycle progression. This disruption aligns with the hypothesized impact on Nedd9 interaction and reduced RhoA activity observed in conditions of low Abba expression.

One possible explanation for these results could be the disruption of the predicted local tertiary structure. In this scenario, heterozygous expression of the human R671W variant would exert a dominant negative effect on *ABBA*'s role in brain development, leading to microcephaly and cognitive delay. This notion is supported by recent work disclosing additional patients carrying the R671W variant⁴². In the same study the significant neurological phenotypes were observed in a *Drosophila* model where the ortholog of human MTSS2 and MTSS1 was deleted. However, from a clinical genetics' standpoint, it is unlikely to find patients with the recurrent R671W mutation without any homozygous or compound heterozygous loss-of-function mutations elsewhere in the *ABBA* gene. This could also suggest a gain-of-function effect of the R671W mutation. Supporting this notion, overexpressing *ABBA*-R671W in cells expressing the wild-type *Abba* in this study did not result in a dominant-negative decrease in RhoA activation, nor did it affect the expression of PH3 *in vivo*. These findings make it plausible to suggest that a mechanism responsible for the phenotype associated with overexpression of the human variant may primarily involve post-cell division processes, such as cell migration.

Although previous research has indicated that *Abba* is not expressed in mature neurons²⁰, more recent studies have shown transient expression of *Abba* in neurons during conditions of increased neuronal plasticity¹³. While our study primarily focused on Abba's role in radial glial cell proliferation, the data also indicate an additional role in the migration of immature neurons. This

is supported by the observed accumulation of Abba-shRNA3 electroporated cells in the ventricular zone. Additionally, we observed a significant decrease in process length in migrating cortical neurons with downregulated Abba and expressing ABBA-R671W mutant but interestingly dendritic processes were increased in length in neurons overexpressing *ABBA* (Supplemental Figure 4). Given that disorganization of radial glial progenitor cells does not lead to neuronal cell death, these findings suggest that Abba may be involved in additional molecular mechanisms in differentiated neurons. Further experiments are needed to determine whether this phenotype is intrinsic to Abba's role in migrating neurons or if it is influenced by the disruption of guiding radial glia.

In conclusion, this study provides compelling evidence for the significant role of *Abba* in brain development. We demonstrate that Abba's involvement in the cell division of radial glial progenitors is a key component of this mechanism. Furthermore, these findings have translational relevance as they contribute to a better understanding of the putative mechanisms underlying human variants associated with microcephaly and developmental disabilities.

Data Availability

All data produced in the present study are available upon reasonable request to the authors

Acknowledgements

We thank Drs Lauren Briere and Gabrielle Lemire for discussion about the data. We thank Biomedicum Imaging Unit (Helsinki, Finland) for their technical assistance with the time-lapse imaging of organotypic cultures. This project was supported by Eranet Neuron III program project ACROBAT, ANR project GABGANG and Academy of Finland grant 341361 to C.R., and by Fondation pour la Recherche Medicale "aide au retour en France" to A.C. Academy of Finland (317038, 319907) and Sigrid Juselius foundation to J.S. and Finnish Cultural Foundation to E.E.

Author Contributions

A.C conceived the project. A.C. designed and performed most of the experiments, analyzed and interpreted all the data, and prepared figures. A.C. and C.R. wrote the original draft of the manuscript. J.S. designed experiments, interpreted data, edited the manuscript. E.E. performed and analyzed experiments and edited the manuscript. H.K. performed organotypic slices culture and live imaging experiments. All authors read and approved the final manuscript.

Conflict of interest statement

None declared

References

1. Ayala R., Shu T., Tsai L.-H. (2007) **Trekking across the brain: the journey of neuronal migration** *Cell* **128**:29–43

2. Higginbotham H. R., Gleeson J. G. (2007) **The centrosome in neuronal development** *Trends Neurosci* **30**:276–283
3. Lambert de Rouvroit C., Goffinet A. M. (2001) **Neuronal migration** *Mech. Dev.* **105**:47–56
4. Liu S., Xiong X., Zhao X., Yang X., Wang H. (2015) **F-BAR family proteins, emerging regulators for cell membrane dynamic changes—from structure to human diseases** *J. Hematol. Oncol. J Hematol Oncol* **8**
5. Doherty G. J., McMahon H. T. (2008) **Mediation, Modulation, and Consequences of Membrane-Cytoskeleton Interactions** *Annu. Rev. Biophys* **37**:65–95
6. Frost A., Unger V. M., De Camilli P. (2009) **The BAR Domain Superfamily: Membrane-Molding Macromolecules** *Cell* **137**:191–196
7. Takano K., Takano K., Toyooka K., Suetsugu S. (2008) **EFC/F-BAR proteins and the N-WASP-WIP complex induce membrane curvature-dependent actin polymerization** *EMBO J* **27**:2817–2828
8. Qualmann B., Koch D., Kessels M. M. (2011) **Let's go bananas: revisiting the endocytic BAR code** *EMBO J* **30**:3501–3515
9. Guerrier S., et al. (2009) **The F-BAR domain of srGAP2 induces membrane protrusions required for neuronal migration and morphogenesis** *Cell* **138**:990–1004
10. Mattila P. K., et al. (2007) **Missing-in-metastasis and IRSp53 deform PI(4,5)P2-rich membranes by an inverse BAR domain-like mechanism** *J. Cell Biol* **176**:953–964
11. Spillane M., et al. (2011) **The actin nucleating Arp2/3 complex contributes to the formation of axonal filopodia and branches through the regulation of actin patch precursors to filopodia** *Dev. Neurobiol* **71**:747–758
12. Saarikangas J., et al. (2015) **MIM-Induced Membrane Bending Promotes Dendritic Spine Initiation** *Dev. Cell* **33**:644–659
13. Chatzi C., et al. (2019) **Exercise-induced enhancement of synaptic function triggered by the inverse BAR protein, Mtss1L** *eLife* **8**
14. Wilson N. K. A., et al. (2011) **A Novel Microduplication in the Neurodevelopmental Gene SRGAP3 That Segregates with Psychotic Illness in the Family of a COS Proband** *Case Rep. Genet* **2011**
15. Carlson B. R., et al. (2011) **WRP/srGAP3 facilitates the initiation of spine development by an inverse F-BAR domain, and its loss impairs long-term memory** *J. Neurosci. Off. J. Soc. Neurosci* **31**:2447–2460
16. Endris V., et al. (2002) **The novel Rho-GTPase activating gene MEGAP/ srGAP3 has a putative role in severe mental retardation** *Proc. Natl. Acad. Sci. U. S. A* **99**:11754–11759
17. Waltereit R., et al. (2012) **Srgap3^{-/-} mice present a neurodevelopmental disorder with schizophrenia-related intermediate phenotypes** *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol* **26**:4418–4428

18. Saitsu H., et al. (2012) **Early infantile epileptic encephalopathy associated with the disrupted gene encoding Slit-Robo Rho GTPase activating protein 2 (SRGAP2)** *Am. J. Med. Genet. A* **158**:199–205
19. Alazami A. M., et al. (2015) **Accelerating Novel Candidate Gene Discovery in Neurogenetic Disorders via Whole-Exome Sequencing of Prescreened Multiplex Consanguineous Families** *Cell Rep* **10**:148–161
20. Saarikangas J., et al. (2008) **ABBA regulates plasma-membrane and actin dynamics to promote radial glia extension** *J. Cell Sci* **121**:1444–1454
21. Dadke D., Jarnik M., Pugacheva E. N., Singh M. K., Golemis E. A. (2006) **Deregulation of HEF1 impairs M-phase progression by disrupting the RhoA activation cycle** *Mol. Biol. Cell* **17**:1204–1217
22. Yu J.-Y., DeRuiter S. L., Turner D. L. (2002) **RNA interference by expression of short-interfering RNAs and hairpin RNAs in mammalian cells** *Proc. Natl. Acad. Sci. U. S. A* **99**:6047–6052
23. Adzhubei I. A., et al. (2010) **A method and server for predicting damaging missense mutations** *Nat. Methods* **7**:248–249
24. Sobreira N., Schiettecatte F., Valle D., Hamosh A. (2015) **GeneMatcher: a matching tool for connecting investigators with an interest in the same gene** *Hum. Mutat* **36**:928–930
25. van Rheenen J., Langeslag M., Jalink K. (2004) **Correcting confocal acquisition to optimize imaging of fluorescence resonance energy transfer by sensitized emission** *Biophys. J* **86**:2517–2529
26. Feige JN., Sage D., Wahli W., Desvergne B., Gelman L. (2005) **PixFRET, an ImageJ plug-in for FRET calculation that can accommodate variations in spectral bleed-throughs** *Microsc Res Tech.* **2517** **68**:51–8
27. Saito T., Nakatsuji N. (2001) **Efficient gene transfer into the embryonic mouse brain using in vivo electroporation** *Dev. Biol* **240**:237–246
28. Tabata H., Nakajima K. (2001) **Efficient in utero gene transfer system to the developing mouse brain using electroporation: visualization of neuronal migration in the developing cortex** *Neuroscience* **103**:865–872
29. Wiegreffe C., Feldmann S., Gaessler S., Britsch S. (2017) **Time-lapse Confocal Imaging of Migrating Neurons in Organotypic Slice Culture of Embryonic Mouse Brain Using In Utero Electroporation** *J. Vis. Exp. JoVE* <https://doi.org/10.3791/55886>
30. Kosodo Y. (2012) **Interkinetic nuclear migration: beyond a hallmark of neurogenesis** *Cell. Mol. Life Sci* **69**:2727–2738
31. Spear P. C., Erickson C. A. (2012) **Apical movement during interkinetic nuclear migration is a two-step process** *Dev. Biol* **370**:33–41
32. Tsai J.-W., Lian W.-N., Kemal S., Kriegstein A. R., Vallee R. B. (2010) **Kinesin 3 and cytoplasmic dynein mediate interkinetic nuclear migration in neural stem cells** *Nat. Neurosci* **13**:1463–1471

33. Hu D. J.-K., et al. (2013) **Dynein recruitment to nuclear pores activates apical nuclear migration and mitotic entry in brain progenitor cells** *Cell* **154**:1300–1313
34. Carabalona A., Hu D. J.-K., Vallee R. B. (2016) **KIF1A inhibition immortalizes brain stem cells but blocks BDNF-mediated neuronal migration** *Nat. Neurosci* **19**:253–262
35. Miller A. L. (2011) **The contractile ring** *Curr. Biol. CB* **21**:R976–978
36. Zhang S., Wu L. (2015) **Roles of neural precursor cell expressed, developmentally downregulated 9 in tumor-associated cellular processes (Review)** *Mol. Med. Rep* **12**:6415–6421
37. Zhong J., et al. (2012) **NEDD9 stabilizes focal adhesions, increases binding to the extra-cellular matrix and differentially effects 2D versus 3D cell migration** *PloS One* **7**
38. Singh M., Cowell L., Seo S., O'Neill G., Golemis E. (2007) **Molecular basis for HEF1/NEDD9/Cas-L action as a multifunctional co-ordinator of invasion, apoptosis and cell cycle** *Cell Biochem. Biophys* **48**:54–72
39. Jarin Z., et al. (2019) **Unusual Organization of I-BAR Proteins on Tubular and Vesicular Membranes** *Biophys. J* **117**:553–562
40. Glotzer M. (2001) **Animal cell cytokinesis** *Annu. Rev. Cell Dev. Biol.* **17**:351–386
41. Klingler E., Francis F., Jabaudon D., Cappello S. (2021) **Mapping the molecular and cellular complexity of cortical malformations** *Science* **371**
42. Huang Y., et al. (2022) **The recurrent de novo c.2011C>T missense variant in MTSS2 causes syndromic intellectual disability** *Am J Hum Genet.* **109**
43. Corona-Rivera, et al. (2023) **MTSS2-related neurodevelopmental disorder: Further delineation of the phenotype** *Eur. J. Med. Genet* **66**
44. Jumper J., et al. (2021) **Highly accurate protein structure prediction with AlphaFold** *Nature* **596**:583–589
45. Field S. J., et al. (2005) **PtdIns(4,5)P2 functions at the cleavage furrow during cytokinesis** *Curr. Biol. CB* **15**:1407–1412
46. Normand G., King R. W. (2010) **Understanding cytokinesis failure** *Adv. Exp. Med. Biol.* **676**:27–55
47. Zhang S., Wu L. (2015) **Roles of neural precursor cell expressed, developmentally downregulated 9 in tumor-associated cellular processes (Review)** *Mol. Med. Rep* **12**:6415–6421

Article and author information

Aurelie Carabalona

Aix Marseille Univ, INSERM, INMED, Marseille, France

For correspondence: aurelie.carabalona@inserm.fr

ORCID iD: [0000-0003-2509-7198](https://orcid.org/0000-0003-2509-7198)

Henna Kallo

Neuroscience Center, HiLIFE, University of Helsinki, Helsinki, Finland

Maryanne Gonzalez

Neuroscience Center, HiLIFE, University of Helsinki, Helsinki, Finland

Liliia Andriichuk

Neuroscience Center, HiLIFE, University of Helsinki, Helsinki, Finland

Ellinoora Elomaa

Helsinki Institute of Life Science, HiLIFE, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

Florence Molinari

Aix Marseille Univ, Inserm, MMG, Marseille, France

Christiana Fragkou

Aix Marseille Univ, INSERM, INMED, Marseille, France

Pekka Lappalainen

Institute of Biotechnology, HiLIFE, University of Helsinki, Helsinki, Finland

Marja W Wessels

Department of Clinical Genetics, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

Juha Saarikangas

Neuroscience Center, HiLIFE, University of Helsinki, Helsinki, Finland, Helsinki Institute of Life Science, HiLIFE, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

Claudio Rivera

Aix Marseille Univ, INSERM, INMED, Marseille, France, Neuroscience Center, HiLIFE, University of Helsinki, Helsinki, Finland

For correspondence: claudio.rivera@helsinki.fi

Copyright

© 2024, Carabalona et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Samuel Pleasure

University of California, San Francisco, San Francisco, United States of America

Senior Editor

Sacha Nelson

Brandeis University, Waltham, United States of America

Reviewer #1 (Public Review):

The manuscript investigates the role of the membrane-deforming cytoskeletal regulator protein Abba in cortical development and its potential implications for microcephaly. It is a valuable contribution to the understanding of Abba's role in cortical development. The strengths and weaknesses identified in the manuscript are outlined below:

Clinical Relevance:

The authors identified a patient with microcephaly and a patient with an intellectual disability harboring a mutation in the Abba variant (R671W) adding a clinically relevant dimension to the study.

Mechanistic Insights:

The study offers valuable mechanistic insights into the development of microcephaly by elucidating the role of Abba in radial glial cell proliferation, radial fiber organization, and the migration of neuronal progenitors. The identification of Abba's involvement in the cleavage furrow during cell division, along with its interaction with Nedd9 and positive influence on RhoA activity, adds depth to our understanding of the molecular processes governing cortical development. Though the reported results establish the novel interaction between Abba and Nedd9, the authors have not addressed whether the mutant protein loses this interaction and whether that results in the observed effects.

In Vivo Validation:

The overexpression of mutant Abba protein (R671W) resulting in phenotypic similarities to Abba knockdown effects supports the significance of Abba in cortical development.

<https://doi.org/10.7554/eLife.92748.1.sa1>

Reviewer #2 (Public Review):

Summary:

Carabalona and colleagues investigated the role of the membrane-deforming cytoskeletal regulator protein Abba (MTSS1L/MTSS2) in cortical development to better understand the mechanisms of abnormal neural stem cell mitosis. The authors used short hairpin RNA targeting Abba20 with a fluorescent reporter coupled with in-utero electroporation of E14 mice to show changes to neural progenitors. They performed flow cytometry for in-depth cell cycle analysis of Abba-shRNA impact on neural progenitors and determined an accumulation in the S phase. Using culture rat glioma cells and live imaging from cortical organotypic slides from mice in utero electroporated with Abba-shRNA, the authors found Abba played a prominent role in cytokinesis. They then used a yeast-two-hybrid screen to identify three high-confidence interactors: Beta-Trcp2, Nedd9, and Otx2. They used immunoprecipitation experiments from E18 cortical tissue coupled with C6 cells to show Abba's requirement for Nedd9 localization to the cleavage furrow/cytokinetic bridge. The authors performed a shRNA knockdown of Nedd9 by in-utero electroporation of E14 mice and observed similar results as with the Abba-shRNA. They tested a human variant of Abba using in-utero electroporation of cDNA and found disorganized radial glial fibers and misplaced, multipolar neurons, but lacked the impact of cell division seen in the shRNA-Abba model.

Strengths:

A fundamental question in biology about the mechanics of neural stem cell division.

Directly connecting effects in Abba protein to downstream regulation of RhoA via Nedd9.

Incorporation of human mutation in ABBA gene.

Use of novel technologies in neurodevelopment and imaging.

Weaknesses:

Unexplored components of the pathway (such as what neurogenic populations are impacted by Abba mutation) and unleveraged aspects of their data (such as the live imaging) limit the scope of their findings and leave significant questions about the effect of ABBA on radial glia development.

(1) The claim of disorganized radial glial fibers lacks quantifications.

On page 11, the authors claim that knockdown of Abba leads to changes in radial glial morphology observed with vimentin staining. Here they claim misoriented apical processes, detached end feet, and decreased number of RGP cells in the VZ. However, they do not provide quantification of process orientation to better support their first claim.

Measurements of radial glia fiber morphology (directionality, length) and angle of division would be metrics that can be applied to data. Some of these analyses could be done in their time-lapse microscopy images, such as to quantify the number of cell divisions during their period of analysis (though that is short-15 hours).

(2) It is unclear where the effect is:

-In RG or neuroblasts? Is it in cell cleavage that results in the accumulation of cells at VZ (as sometimes indicated by their data like in Figure 2A or 4D)? Interrogation of cell death (such as by cleaved caspase 3) would also help. Given their time-lapse, can they identify what is happening to the RG fiber? The authors describe a change in "migration" but do not show evidence for this for either progenitor or neuroblast populations. Given they have nice time-lapse imaging data, could they visualize progenitor versus young neuron migration? Analysis of neuroblasts (such as with doublecortin expression in the tissue) would also help understand any issues in migration (of neurons v stem cells).

-At cleavage furrow? In abscission? There is high-resolution data that highlights the cleavage furrow as the location of interest (Figure 3A), however, there is also data (Figure 3B) to suggest Abba is expressed elsewhere as well and there is an overall soma decrease. More detail of the localization of Abba during the division process would be helpful for example, could cleavage furrow proteins, such as Aurora B, co-localization (and potentially co-IP) help delineate subpopulations of Abba protein? Furthermore, the FRET imaging is a unique way to connect their mutation with function - could they measure/quantify differences at furrow compared to the rest of soma to further corroborate that the Abba-associated RhoA effect was furrow-enriched?

-The data highlights nicely that a furrow doesn't clearly form when ABBA expression and subsequent RhoA activity are decreased (in Figure 3 or 5A). Does this lead to cells that can't divide because of poor abscission, especially since "rounding" still occurs? Or abnormal progenitors (with loss of fiber or inability to support neuroblast migration)? Or abnormal progression of progenitors to neuroblasts?

(3) Limited to a singular time point of mouse cortical development

On page 13, the authors outline the results of their Y2H screen with the identification of three high-confidence interactors. Notably, they used an E10.5-E12.5 mouse brain embryo library rather than one that includes E14, the age of their in-utero electroporation mice. Many of the authors' claims focus on in-utero electroporation of shRNA-Abba of E14 mice that are then evaluated at E16-18. Justification for the focus on this age range should be included to support that their findings can then be applied to all mouse corticogenesis.

(4) Detail of the effect of the human variant of the ABBA mutation in mice is lacking.

Their identification of the R671W mutation is interesting and the IUE model warrants more characterization, as they did with their original KD experiments.

-Could they show that Abba protein levels are decreased (in either cell lines or electroporated tissue)?

-While time-lapse morphology might not have been performed, more analysis on cell division phenotype (such as plane of division and radial glia morphology) would be helpful.

<https://doi.org/10.7554/eLife.92748.1.sa0>