

Tumor Purity-Related Genes for Predicting the Prognosis and Drug Sensitivity of DLBCL Patients

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 1

January 31, 2024 (this version)

Posted to preprint server

October 5, 2023

Sent for peer review

September 27, 2023

Zhenbang Ye, Ning Huang, Yongliang Fu, Rongle Tian, Wenting Huang 

Department of Pathology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China; • Department of Hepatobiliary Surgery, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100021, China • Department of Pathology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital & Shenzhen Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Shenzhen, 518116, China

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Background

Diffuse large B-cell lymphoma (DLBCL) is the predominant type of malignant B-cell lymphoma. Although various treatments have been developed, the limited efficacy calls for more and further exploration of its characteristics.

Methods

Datasets from Gene Expression Omnibus (GEO) database were used for identifying the tumor purity of DLBCL. Survival analysis was employed for analyzing the prognosis of DLBCL patients. Immunohistochemistry was conducted to detect the important factor that influenced the prognosis. Drug sensitive prediction was performed to evaluate the value of the constructed model.

Results

VCAN, CD3G and C1QB were identified as three key genes that impacted the outcome of DLBCL patients both in GEO datasets and samples from our center. Among them, VCAN and CD3G+ T cells were correlated with favorable prognosis, and C1QB was correlated with worse prognosis. The ratio of CD68+ macrophages and CD8+ T cells was associated with better prognosis. In addition, CD3G+ T cells ratio was significantly correlated with CD68+ macrophages, CD4+ T cells and CD8+ T cells ratio, indicating it could play an important role in the anti-tumor immunity in DLBCL. The riskScore model constructed based on the RNASeq data of VCAN, C1QB and CD3G work well in predicting the prognosis and drug sensitivity.

Conclusion

VCAN, CD3G and C1QB were three key genes that influenced the tumor purity of DLBCL, and could also exert certain impact on drug sensitivity and prognosis of DLBCL patients.

eLife assessment

The findings in this study are **useful** and may have practical implications for predicting DLBCL risk subject to further validating the bioinformatics outcomes. We found the approach and data analysis **solid**. However, some concerns regarding the drug sensitivity prediction and the links between the selected genes for the risk scores have been raised that need to be addressed by further functional works.

Introduction

The latest refined classification by the World Health Organization (WHO) categorizes large B-cell lymphoma as a heterogeneous group of B-cell lymphomas[1]. Diffuse large B-cell lymphoma (DLBCL) is the most prevalent type among them, accounting for around 30% of all non-Hodgkin lymphomas. DLBCL can be classified into three subtypes based on its immunohistochemical expression patterns: germinal center B-cell-like (GCB), activated B-cell-like (ABC), and unclassified[2]. After undergoing R-CHOP chemotherapy, about 60% of patients achieve long-term remission; however, approximately 30% of patients experience relapse, resulting in poor prognosis and a considerable number of deaths from refractory lymphoma[3]. Consequently, to explore the characteristics of DLBCL in detail is urgently needed for developing more effective therapy.

Solid tumor tissue comprises tumor cells and the surrounding stroma, which encompasses diverse types of matrix cells, immune cells, endothelial cells[4], etc. The tumor microenvironment (TME) is a complex and dynamic system that consists of the extracellular matrix and a variety of cellular components. Recent studies have unveiled multiple subgroups of immune cells within the microenvironment of DLBCL, including T cells, B cells, NK cells, monocytes/macrophages, dendritic cells, as well as the distribution of stromal cell components like fibroblasts and endothelial cells[5, 6]. Despite the relatively limited composition of the TME in DLBCL, its role in tumor proliferation and evasion of the immune system should not be disregarded. The interaction between tumors and the microenvironment is a vital factor that impacts the development and prognosis of B-cell lymphoma[7]. Nevertheless, the existing research on the influence of the TME on the prognosis of DLBCL patients is limited and lacks a consensus.

Moreover, the comprehensive investigation of non-immune cell components in the TME is still lacking. Previous research on stroma in DLBCL has predominantly indicated that a higher quantity of extracellular matrix is associated with a more favorable prognosis, while increased vascular density is associated with poorer prognosis[8]. Furthermore, higher stromal scores have been associated with an improved prognosis in DLBCL patients[9]. Additionally, a fibrotic tumor microenvironment has been correlated with a better prognosis after DLBCL chemotherapy and immunotherapy[10]. These research findings stem from computational analysis of stromal and immune scoring in gene databases and have not been experimentally validated as of yet.

Tumor purity quantifies the relative ratio of tumor cells to the surrounding stromal components in solid tumors, elucidating the dynamics between tumor cells and their microenvironment[11]. It can partly reflect the characteristics of TME, namely, a higher tumor purity indicates a lower abundance of stromal components in TME. Tumor purity is associated with patient prognosis, and the strength of this association varies across different tumor types[12-14]. Therefore, when investigating the influence of TME on the prognosis of DLBCL, it is crucial to analyze not only the immune cell components but also the significance of non-immune cell components.

This study utilized bioinformatic analysis to establish the relationship between immune and stromal components and the prognostic outcomes of DLBCL patients. We developed a novel immunohistochemical panel to assess prognostic outcomes and treatment sensitivity by detecting the expression of VCAN, CD3G, C1QB, CD68, CD4 and CD8 in both the TME and tumor cells of 190 DLBCL patients. We then explored their relationship with DLBCL clinicopathological features as well as overall survival (OS).

Materials and Methods

Data collection and tumor purity-related genes (TPGs) selection

The RNA-Sequence and clinical data of GSE53786 and GSE32918 datasets were download from Gene Expression Omnibus (GEO) database. The first gene symbols of GSE53786 datasets were retained when one probe detected multiple genes. Average expression value of genes in each dataset were calculated and used when one gene was detected by multiple probes. Tumor purity was assessed by ESTIMATE (Estimation of Stromal and Immune cells in Malignant Tumor tissues using Expression data) algorithm and the then its correlation with genes expression was analyzed. The genes with $|r| \geq 0.5$ and p value < 0.05 was defined as the TPGs.

TPGs function analysis

Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses were executed to analyze the biological processes, cellular components, molecular functions and pathways related to the TPGs. The statistical significance was considered as $p.adjust < 0.05$. The protein-protein interactions (PPI) analysis was utilized to investigate the interaction among TPGs, and those with interactive confidence greater than 0.90 on the STRING platform (version 11.5) were selected to establish an interaction network with Cytoscape software (version 3.8.2).

Prognostic model

The prognostic model was constructed with “survival” package in R (version 4.1.3). The genes enrolled in this model was selected among the prognostic and PPI hub TPGs by function “step” in “survival” package, which can optimize the model. The prognostic model was represented by
$$\text{riskScore} = \sum_{i=1}^n \text{gene expression}_i \times \text{coef}_i$$
.

Clinical specimens and follow-up

190 patients from Cancer hospital Chinese Academy of Medical Sciences, the CHCAMS cohort, were enrolled in this study (Supplementary table 1). All patients received surgery or biopsy during September, 2010 and September, 2020, and then standard follow-ups were carried out until March, 2023. The overall survival (OS) was defined as the interval between the operation and death or the last follow-up. The specimens from the CHCAMS cohort were used for immunohistochemistry assay. The study was designed according to the Declaration of Helsinki and approved by the institutional ethics committee of Cancer Hospital Chinese Academy of Medical Sciences. Informed consent was taken from all the patients.

Ihc

Paraffin embedded DLBCL tissues of CHCAMS Cohort were used for immunohistochemistry (IHC). After de-paraffinization and hydration, heat-induced method was performed for antigen retrieval. Primary antibody of VCAN (AB177480, 1:100, Abcam, USA), CD3G (AB134096, 1:1000, Abcam, USA), C1QB (AB92508, 1:50, Abcam, USA), CD68 (303565, 1:1000, Abcam, USA), CD4 (ZM-0418, ZSGB-BIO, China), CD8 (ZA-0508, ZSGB-BIO, China) was incubated at 4°C overnight. Sections were washed with TBS-T buffer, and then incubated with secondary antibody, and finally stained with DAB. The quantitative analysis of the slices was conducted by QuPath-0.4.3. VCAN and C1QB were assessed by H-score, and CD3G, CD68, CD4, CD8 were assessed as the ratio of the corresponding positive cells among all cells.

Drug sensitivity prediction

Drug sensitivity prediction was conducted utilizing “oncoPredict” packages in R 4.1.3. The drug sensitivity data was collected from Genomics of Drug Sensitivity in Cancer (GDSC). The drugs that was analyzed in this study was selected according to clinical practice or clinical trials searched in Pubmed.

Statistical analysis

Data in this study was shown in the form of mean $\pm$ SEM. Correlation between two variates was determined with Spearman analysis. Kaplan–Meier (K–M) curve and Log rank test were used for survival analysis. The cut-offs of survival analysis were provided by X-tile. The independent risk factor analysis was performed with Cox regression analysis. Receiver operating characteristic (ROC) curve was used for test the efficacy of prognostic model. The clinicopathological characteristics difference analysis was conducted with χ^2 test, Fisher’s exact test or Wilcoxon rank sum test. The drug sensitivity scores were compared with Wilcoxon rank sum test. In this study, $p < 0.05$ were considered statistically significant.

Results

Tumor purity related genes were correlated with extracellular matrix organization and immune response

Based on GSE53786 dataset, we first assessed the tumor purity of DLBCL, which ranged from 17.2% to 67.4% (**Figure 1A** [↗](#)). In order to screen out the TPGs, we then analyzed the correlation between genes expression and tumor purity. According to the thresholds mentioned above, 642 genes were identified as TPGs, among which 31 genes were positively correlated with tumor purity, while 611 genes were negatively correlated with it (**Figure 1B** [↗](#)). In addition, tumor purity did have influence on the prognosis of DLBCL patients, which showed that patients with high tumor purity had lower OS rate than those with low tumor purity (**Figure 1C** [↗](#), $p = 0.025$).

Next, we performed GO and KEGG enrichment analyses to explore the functions and signaling pathways in which these TPGs were involved. It turned out that the TPGs were mainly associated with extracellular matrix organization and immune response (**Figure 1C** [↗](#), **1D** [↗](#)). Not only did the enrichment results confirm that these genes were reliable to be related with the tumor purity, but it also laid solid foundations for the sequent analyses.

A prognostic model was constructed with three TPGs

With the 642 TPGs, we exerted PPI analysis to investigate their interaction and the hub genes (**Figure 2A** [↗](#)). The TPGs who had five or more interactive genes were shown in **Figure 2B** [↗](#), and defined as hub genes. Then, we performed univariate Cox regression analysis to figure out the TPGs that were associated with the prognosis of DLBCL patients, and 103 genes were identified




Figure 1

TPGs were screened out with GSE53786 dataset.

(A) The range (17.2 –67.4%) of tumor purity of samples in GSE53786. (B) The heatmap showing genes defined as TPGs. (C) The K-M curve showed high tumor purity was correlated with poor prognosis in DLBCL patients in GSE53678 dataset (Patients were divided into two groups according to the best-cutoff provided by “survminer” package in R). (D) GO analysis of TPGs. (E) KEGG analysis of TPGs. TPGs, tumor purity-related genes; K-M, Kaplan-Meier; DLBCL, diffuse large B cell lymphoma; GO, gene ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes.

(**Figure 2C** [↗](#)). Interestingly, most of the TPGs were correlated with good outcome (with $HR < 1$), and only six genes were associated with poor outcome (with $HR > 1$). Through conducting intersection analysis, we found nine genes (LUM, VCAN, YAP1, COL5A2, SDC2, TWIST1, CD3G, C1QB and C3) were intersection genes, indicating that they had an active effect in modulating the tumor purity, as well as influencing the prognosis of DLBCL patients.

After ascertaining the key genes, we tried to construct a prognostic model with them. The model was constructed by Cox regression, and the three selected genes (VCAN, CD3G, C1QB) and their parameters like coefficient, HR and 95%CI of HR, were shown in **Figure 3A** [↗](#). It showed that VCAN and CD3G were correlated with good prognosis and C1QB was correlated with poor prognosis. All patients were divided into high and low-risk group according to the median value of riskScore (**Figure 3B** [↗](#)). As expected, the high-risk group has worse prognosis than the low-risk group (**Figure 3C** [↗](#)). In addition, the three genes were differentially expressed between high and low-risk group, with VCAN and CD3G showing high expression level in low-risk group, and C1QB showing high expression level in high-risk group, which was consistent with the coefficient (**Figure 3D** [↗](#)). To appraise the efficacy of these prognostic model, we conducted survival analysis and ROC analysis. High-risk group had lower OS rate than low-risk group (**Figure 3E** [↗](#), $p < 0.001$), and the areas under curve (AUC) for 1-year, 3-year and 5-year ROC were 0.73, 0.77 and 0.77 respectively (**Figure 3F** [↗](#)). Just similar to tumor purity, the high riskScore indicated bad outcome, which was consistent with the positive correlation between tumor purity and riskScore (**Figure 3G** [↗](#)). This TPGs signature prognostic model manifested satisfying prognostic efficacy.

When we applied this model to GSE32918 dataset, it still did excellently and the results were in accordance with that in GSE53786 dataset (**Figure 4A** [↗](#), **4B** [↗](#); **Supplementary Figure 2** [↗](#)). Next, we analyzed the relationship between riskScore and some clinicopathological characteristics provided in GSE53786 dataset. The results showed that high-risk group had more ABC type DLBCL, while low-risk group had more the GCB type DLBCL (**Figure 4C** [↗](#)). Besides, high-risk group displayed higher lactic dehydrogenase (LDH) ratio (**Figure 4F** [↗](#)). However, the Eastern Cooperative Oncology Group (ECOG) performance and stage was not associated with the riskScore (**Figure 4D** [↗](#), **4E** [↗](#)). Still, the high-risk group has more Stage III and Stage IV patients, but less Stage I patients than low-risk group. Finally, we employed the univariate and multivariate analysis to explore whether riskScore was an independent prognostic factor for DLBCL patients. As expected, the riskScore was associated with the poor prognosis (**Figure 4G** [↗](#), $p < 0.001$, $HR = 1.545$, 95%CI 1.284–1.861) and was an independent prognostic factor (**Figure 4H** [↗](#), $p = 0.002$, $HR = 1.474$, 95%CI 1.156–1.879).

The prognostic value of VCAN, CD3G and C1QB were validated by IHC assay

With the purpose of the further validation of the prognostic value of VCAN, CD3G and C1QB, we detected the expression of these genes in CHCAMS cohort by IHC. For VCAN, the patients were divided into high and low group according to the cut-off of the H-score (275.42) provided X-tile. The survival analysis showed that patients with high expression of VCAN had higher OS rate (**Figure 5A** [↗](#), $p = 0.003$). For CD3G, previous study revealed that it was a component of T cell receptor complex, for which it could be regarded as a marker of T cells. Therefore, we assessed the expression level of CD3G by counting the CD3G+ T cells ratio, and divided patients by the cut-off (2.5%). The survival analysis revealed that patients with high CD3G+ T cells infiltration showed favorable prognosis (**Figure 5B** [↗](#), $p < 0.001$). For C1QB, the patients in high expression group (cut-off = 82.41) showed adverse prognosis (**Figure 5C** [↗](#), $p = 0.015$). Although the detection of protein level was not convenient to build a prognostic model for the difference of assessment methods and the lake of coefficient, these results were in accordance with those of GEO datasets, which successfully proved the prognostic value of VCAN, CD3G and C1QB.




Figure 2

The key gene candidates used for constructing prognostic model were selected.

(A) The PPI network of TPGs (orange nodes representing genes positively correlated with tumor purity, and green nodes representing genes negatively correlated with tumor purity). (B) The barplot showing hub genes with five or more interactive genes. (C) The forest plot showing prognostic TPGs of DLBCL patients in GSE53786. (D) The venn plot showing intersection genes of PPI hub gene and prognostic TPGs. PPI, protein-protein interaction.




Figure 3

TPGs signature prognostic model was constructed.

(A) Three genes enrolled in the prognostic model. (B) The patients in GSE53786 dataset were divided into high and low-risk group according to the median riskScore based on the prognostic model. (C) High-risk group had worse prognosis than low-risk group. (D) The heatmap showing expression discrepancy of the three genes. (E) Survival analysis revealed that high-risk group had poor prognosis in GSE53786 dataset. (F) The ROC curve showed that the prognostic model performed well in predicting 1-year, 3-year and 5-year prognosis in GSE53786 dataset. (G) Tumor purity was positively correlated with riskScore in GSE53786 dataset. ROC, receiver operating characteristic.




Figure 4

The riskScore of three TPGs signature prognostic model was an independent prognostic factor in DLBCL patients.

(A) Survival analysis results in GSE32918 dataset was consistent to that of GSE53786 dataset. (B) The prognostic model also did well in GSE32918 dataset. (C) High-risk group in GSE53786 dataset contained more ABC type DLBCL, while low-risk group contained more GCB type DLBCL. (D) The ECOG performance of two groups (GSE53786 dataset) showed no statistical difference. (E) More patients in high-risk group were at Stage III or Stage IV, and less patients were at Stage I, compared with low-risk group, although no statistical significance was shown (GSE53786 dataset). (F) High-risk group had higher LDH ratio (GSE53786 dataset). (G) The riskScore was associated with poor prognosis of DLBCL patients in GSE53786 dataset. (H) The riskScore was an independent prognostic factor for DLBCL patient. * $p < 0.05$, ** $p < 0.01$, ns, not significant. ABC, activated B cell; GCB, germinal center B cell; ECOG, Eastern Cooperative Oncology Group; LDH, lactic dehydrogenase.




Figure 5

The analysis of CHCAMS cohort.

(A) The representative image of VCAN staining of high and low expression groups and the survival analysis based on VCAN expression. (B) The representative image of CD3G staining of high and low CD3G+ T cells ratio groups and the survival analysis based on CD3G+ T cells ratio. (C) The representative image of C1QB staining of high and low expression group and the survival analysis based on C1QB expression. (D) The representative image of CD68 staining of high and low CD68+ macrophages ratio groups and the survival analysis based on CD68+ macrophages ratio. (E) The representative image of CD8 staining of high and low CD8+ T cells ratio groups and the survival analysis based on CD8+ T cells ratio. (F) The representative image of CD4 staining of high and low CD4+ T cells ratio groups and the survival analysis based on CD4+ T cells ratio. (G) The correlation between VCAN, CD3G+ T cells ratio, C1QB and CD68+ macrophages, CD8+ T cells and CD4+ T cells ratio. "x" means no statistical significance. (H) The CD3G+ T cells infiltration varied from colon to testis originating DLBCL in male. (I) The VCAN expression level was different between intra- and extra-lymph node DLBCL.

Given that these genes could potentially influence the tumor purity of DLBCL, we then analyzed the relationship between them and CD68+ macrophages, CD4+ T cells and CD8+ T cells. As was shown in **Supplementary figure 3A** [\[15\]](#), CD68+ macrophages [(17.75±1.05) %] account for more ratio than CD4+ T cells [(0.68±0.20) %] and CD8+ T cells [(6.69±0.56) %] ($p < 0.001$, Kruskal-Wallis Test and Dunn's Test). In the survival analysis of these three types of immune cells, we found that CD68+ macrophages, CD8+ T cells and CD4+ T cells were associated with better prognosis (**Figure 5D-F** [\[15\]](#), $p = 0.029$, $p = 0.002$, $p = 0.053$). And XCELL and QUANTISEQ algorithm revealed that M2 macrophages accounted for more proportion than M1 macrophages in GSE53786 and GSE32918 (**Supplementary figure 3L-O** [\[15\]](#)). Besides, the ratio of CD3G+ T cells was positively correlated with that of CD68+ macrophages, CD8+ T cells and CD4+ T cells, C1QB expression level was positively correlated with CD8+ T cells, and VCAN expression level was positively correlated with CD8+ T cells ratio (**Figure 5G** [\[15\]](#)).

In addition to the above analyses, we also explored the relationship between these three genes and the proliferation and location of DLBCL. It turned out that CD3G+ T cells ratio was higher in DLBCL originated from groin and testis, and VCAN featured higher expression in lymph node originated DLBCL (**Figure 5H-I** [\[15\]](#), $p < 0.05$, $p < 0.01$, **Supplementary figure 3B-K** [\[15\]](#)).

The TPGs signature model could also predict the drug sensitivity of DLBCL patients

In order to learn about the ability of the previously mentioned model to predict drug sensitivity, we performed the prediction with “oncoPredict” package in R. Fifteen drugs (**Supplementary table 2**) included in the GDSC and used in clinical practice or under clinical trials (searched on Pubmed) were enrolled in this prediction analysis.

As is shown in **Figure 6A** [\[15\]](#) (prediction of GSE53786), patients in high-risk group could be sensitive to Carmustine, Cytarabine, Oxaliplatin, Vincristine, Vorinostat, and Bortezomib, but no drug could work better in low-risk group. And in GSE32918 (**Figure 6B** [\[15\]](#)), Carmustine, Cytarabine, Oxaliplatin, Vorinostat, Afuresertib, Bortezomib, Ibrutinib and Tamoxifen could work better in high-risk group, and Vincristine (sensitivity score: low-risk vs high-risk = 0.219 ± 0.026 vs 0.223 ± 0.031) could work better in low-risk group. The discrepancy between the prediction in two datasets might be due to the samples and sequencing platforms. However, the intersection analysis of the drugs to which the high-risk patients in both datasets could be sensitive revealed that Carmustine, Cytarabine, Oxaliplatin, Vorinostat and Bortezomib could be reliable candidates for treating high-risk patients based on the three TPGs signature prognostic model (**Supplementary table 2**).

Discussion

In this study, bioinformatics techniques were employed to identify three genes (VCAN, CD3G, C1QB) that exhibit associations with prognosis in both immune and stromal environments, thereby revealing their relationship with the prognosis of DLBCL patients. The findings indicate that higher expression of VCAN, increased infiltration of CD3G+ T cells, and decreased expression of C1QB are correlated with favorable prognostic outcomes. Conversely, a lower infiltration of CD68+ macrophages and lower infiltration of CD8+ T cells are associated with poorer prognosis. Furthermore, we investigated the relationship between risk genes related to tumor purity and treatment sensitivity and established a list of possible drugs that might be helpful for enhancing outcomes.

Previous studies have extensively investigated the VCAN gene in relation to tumorigenesis and metastasis [\[15\]](#). VCAN, also known as versican, is a crucial component of extracellular matrix [\[16\]](#), and exists in several isoforms [\[17\]](#). Research has shown that VCAN plays a




Figure 6

Drug sensitivity prediction revealed therapeutic candidates for high-risk group.

(A) Drug sensitivity prediction results with statistical significance in GSE53786 dataset. (B) Drug sensitivity prediction results with statistical significance in GSE53786 dataset.

multifaceted role in TME depending on the cell type expressing it. When expressed by myeloid cells, VCAN induces an anti-inflammatory and immunosuppressive microenvironment. Conversely, its expression by stromal cells typically leads to a pro-inflammatory response[18]. In gastric cancer, high expression of VCAN has been associated with increased infiltration of fibroblasts, significant enrichment of stromal-associated signaling pathways and poor prognosis[19]. In hepatocellular carcinoma, VCAN exhibits a strong association with immune checkpoint gene expression[20]. Despite these findings in other tumor types, the role of VCAN in DLBCL has not been explored yet. Our study reveals that high expression of VCAN is actually associated with a more favorable prognosis. This suggests that VCAN may have different functions in different tumor types. One possible mechanism through which VCAN influences prognosis is that VCAN overexpression in DLBCL may also impact tumor cell proliferation. A study has shown that overexpression of VCAN V1 has an inhibitory effect on cell proliferation, partly due to its promotion of activation-induced cell death in lymphoid cell lines[17]. Hence, the high expression of VCAN in DLBCL could impact not only the TME but also tumor cell proliferation, suggesting a potential mechanism for the observed preferable prognosis.

C1q is synthesized in the tumor microenvironment and functions as an extracellular matrix protein, and C1QB is a component of C1q[21]. Previous studies have provided insights into the diverse roles of C1q in cancer progression. However, the majority of these results, as observed in non-small cell lung carcinoma and gastric cancer, indicate that high C1q expression in TME is associated with a poor prognosis [22-24]. Additionally, C1QB has been found to exert an impact on the TME and is positively associated with infiltration levels of CD8+ T cell, as well as with M1 and M2 macrophages in osteosarcoma[25]. Moreover, C1QB expression shows a positive correlation with predictive biomarkers for immunotherapy, such as PD-L1 expression and CD8+ T cell infiltration[24]. Furthermore, in malignant melanoma, C1QB promotes proliferation, migration and invasion, while inhibiting cell apoptosis[26], and the high-expression group exhibits significant enrichment of genes related to immune and apoptosis[21]. In our study, we found that high expression of C1QB in DLBCL was associated with a worse prognosis and positively correlated with CD8+ T cells infiltration. Based on these findings, we propose that C1QB in DLBCL might share similarities with its functions in other tumor types, particularly regarding the promotion of recruitment and subsequent deactivation of CD8+ T cells within the TME through the induction of immune checkpoint effects. These results shed light on the intricate role of C1QB in TME and its potential significance as a prognostic marker in DLBCL.

CD3G is a member of the TCR/CD3 complex primarily expressed in lymphocytes subgroups. It plays a crucial role in initiating the activation of T cells[27]. It is also involved in coupling antigen recognition[28]. It is reported to associate with long-term OS and good prognosis in breast invasive carcinoma[29] as well as in head and neck squamous cell carcinoma[30]. However, its role in DLBCL has not been fully explored. In our study, we revealed that high infiltration of CD3G+ T cells is correlated with good prognosis. The infiltration of CD3G+ T cells was found to be positively related to the infiltration of CD8+, CD4+ and CD68+ cells. This indicates that CD3G+ T cells in DLBCL may enhance the tumor antigen recognition process and stimulate the infiltration of immune cells, leading to an increased abundance of immune cell infiltration in the TME. The presence of CD3G+ T cells in the TME may contribute to a favorable prognosis by facilitating the activation of immune responses against tumor cells.

Macrophages play a crucial role in TME, and CD68 is a surface marker specific to macrophages. Macrophages can be roughly classified into two types based on their functional features: M1 or M2. M1 macrophages exert anti-tumor effects, whereas M2 macrophages promote tumor growth and progression in TME[31]. A previous study found that low infiltration of CD68+ macrophages was associated with an inferior prognosis[32]. Similarly, our study has yielded similar results, revealing a noteworthy correlation between a high proportion of CD68+ macrophages in the TME and improved prognosis. Additionally, by analyzing the datasets, we observed a higher proportion

of M1 macrophages infiltration compared to M2 macrophages. This suggests that, within our DLBCL cohort, these macrophages may also exhibit the M1 phenotype and consequently play a protective role against tumor progression.

CD8 is widely recognized as a marker of CD8⁺ T cells, also known as cytotoxic T cells[33]. These cells are crucial for the immune response against tumors. However, in DLBCL, CD8⁺ T cells exhibits elevated levels of inhibitory molecules on their surface, such as PD-1, PD-L1, TIM3. High expression of TIM3, an inhibitory immune checkpoint receptor, on CD8⁺ T cells has been associated with tumor progression and poor outcomes[34, 35]. These inhibitory molecules may impair the function of CD8⁺ T cells and hinder their anti-tumor activity. Surprisingly, our study demonstrates a correlation between the infiltration of CD8⁺ T cells and favorable prognosis in DLBCL. Here, we propose a hypothesis that in our study, the observed high expression of VCAN might create a suppressive environment for PD-1⁺ CD8⁺ T cells[18, 36]. Intriguingly, our study revealed a statistically significant correlation between VCAN expression, C1QB expression and CD8⁺ T cell infiltration. VCAN has the potential to modulate immune infiltration by reducing the immunosuppressive phenotype of immune cells[37], thus enabling a more efficient anti-tumor response. This aspect is still worth of consideration.

Taken together, our findings underscore the significant roles of VCAN, CD3G, C1QB, which influence both the TME and the behavior of tumor cells. The interaction between each component and the TME is rather complicated. To fully comprehend the underlying mechanisms and identify potential therapeutic targets in DLBCL, further investigation is required.

However, this study still has several limitations that should be addressed. Firstly, the patients included in this study were from a single center, which may introduce biases into the results. Although we made efforts to minimize these biases, it is inevitable that some may persist. Secondly, we hypothesized that VCAN, CD3G and C1QB could serve as continuous prognostic parameters, thereby eliminating the need for a cut-off. However, the methodology used in this study, which utilized IHC staining to assess the protein expression levels, may have potential limitations. While IHC is a widely used technique, additional validation is needed to confirm the prognostic value of VCAN, CD3G and C1QB in DLBCL. Furthermore, due to the potential variability in interpreting IHC results across different centers, a standardized coefficient and formula have not been established to calculate the final prognostic index for patients with DLBCL. Developing a standardized approach would be beneficial in ensuring consistent and accurate interpretation of IHC results. To address these limitations and expand upon our findings, future studies should strive to incorporate a diverse range of patients from multiple centers. Additionally, it is crucial to employ rigorous experimental techniques to authenticate the prognostic significance of VCAN, CD3G, and C1QB in DLBCL.

Data Availability

The GEO datasets can be obtained from <https://www.ncbi.nlm.nih.gov/gds/?term=>. Due to the nature of this study, the data of CHCAMS cohort can be accessed from Dr. Wenting Huang upon reasonable request.

<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi>

<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE53786>

Author contributions

Conceptualization: Wenting Huang, Ning Huang, Zhenbang Ye, Data Curation: Zhenbang Ye, Ning Huang, Formal Analysis: Zhenbang Ye, Ning Huang, Investigation: Zhenbang Ye, Methodology: Zhenbang Ye, Ning Huang, Wenting Huang, Resources: Wenting Huang, Yongliang Fu, Supervision: Wenting Huang, Visualization: Zhenbang Ye, Ning Huang, Validation: Ning Huang, Writing-Original Draft: Zhenbang Ye, Ning Huang, Writing-Review & Editing: Wenting Huang, Yongliang Fu, Rongle Tian

Funding


This work is supported by Shenzhen High-level Hospital Construction Fund. The funders had no role in study design, data collection and analysis, interpretation of data, or preparation of the manuscript.

Acknowledgements

Not applicable.


Conflicts of interest

The authors made no disclosures.



Supplementary Figure 1


Flow chart and study design of this research.



Supplementary Figure 2

The correlation between tumor purity and prognosis and riskScore in GSE32918 dataset.

(A) High ESTIMATE-Score group had better prognosis. (B) ESTIMATE-Score was negatively associated with riskScore. Note: ESTIMATE-Score manifests the tumor purity, with high ESTIMATE-Score indicating low tumor purity and high immune and stromal components. Because the GSE32918 datasets were sequenced with illumina platform, the ESTIMATE algorithm only provided ESTIMATE Score to represent the tumor purity.



Supplementary Figure 3

The immunoenvironment analysis and clinicopathological analysis of CHCAMS cohort and GEO datasets.

(B) The ratio of CD68+ macrophages, CD8+ T cells and CD4+ T cells in CHCAMS cohort. (B–D) The correlation between VCAN, CD3G+ T cells, C1QB and types of DLBCL in CHCAMS cohort. (E–K) The correlation between VCAN, CD3G+ T cells, C1QB and DCBCL origination in CHCAMS cohort. “Location to LN” means the relative location of the tumor to lymph node, that is, intra-lymph node or extra-lymph node. (L–M) The predicted M1 and M2 macrophages proportion in GSE53786 dataset. (N–O) The predicted M1 and M2 macrophages proportion in GSE532918 dataset. $**p < 0.01$, $***p < 0.001$.

References

1. Alaggio R., et al. (2022) **The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Lymphoid Neoplasms** *Leukemia* **36**:1720–1748
2. Sehn L.H., Salles G. (2021) **Diffuse Large B-Cell Lymphoma** *N Engl J Med* **384**:842–858
3. Autio M., et al. (2021) **Immune cell constitution in the tumor microenvironment predicts the outcome in diffuse large B-cell lymphoma** *Haematologica* **106**:718–729
4. Joyce J.A., Pollard J.W. (2009) **Microenvironmental regulation of metastasis** *Nat Rev Cancer* **9**:239–52
5. Steen C.B., et al. (2021) **The landscape of tumor cell states and ecosystems in diffuse large B cell lymphoma** *Cancer Cell* **39**:1422–1437
6. Ciavarella S., et al. (2018) **Dissection of DLBCL microenvironment provides a gene expression-based predictor of survival applicable to formalin-fixed paraffin-embedded tissue** *Ann Oncol* **29**:2363–2370
7. Ennishi D., et al. (2020) **Toward a New Molecular Taxonomy of Diffuse Large B-cell Lymphoma** *Cancer Discov* **10**:1267–1281
8. Miyawaki K., et al. (2022) **A germinal center-associated microenvironmental signature reflects malignant phenotype and outcome of DLBCL** *Blood Adv* **6**:2388–2402
9. Schmitz R., et al. (2018) **Genetics and Pathogenesis of Diffuse Large B-Cell Lymphoma** *N Engl J Med* **378**:1396–1407
10. Lou X., et al. (2022) **CCL8 as a promising prognostic factor in diffuse large B-cell lymphoma via M2 macrophage interactions: A bioinformatic analysis of the tumor microenvironment** *Front Immunol* **13**
11. Gong Z., Zhang J., Guo W. (2020) **Tumor purity as a prognosis and immunotherapy relevant feature in gastric cancer** *Cancer Med* **9**:9052–9063
12. Lou S., et al. (2021) **Comprehensive Characterization of Tumor Purity and Its Clinical Implications in Gastric Cancer** *Front Cell Dev Biol* **9**
13. Zhao Y., et al. (2023) **Tumor purity-associated genes influence hepatocellular carcinoma prognosis and tumor microenvironment** *Front Oncol* **13**
14. Zhang C., et al. (2017) **Tumor Purity as an Underlying Key Factor in Glioma** *Clin Cancer Res* **23**:6279–6291
15. Baghy K., et al. (2016) **Proteoglycans in liver cancer** *World J Gastroenterol* **22**:379–93
16. Wight T.N (2017) **Provisional matrix: A role for versican and hyaluronan** *Matrix Biol* **60**:38–56
17. Fujii K., et al. (2015) **Versican upregulation in Sézary cells alters growth, motility and resistance to chemotherapy** *Leukemia* **29**:2024–32

18. Wight T.N., et al. (2020) **Versican-A Critical Extracellular Matrix Regulator of Immunity and Inflammation** *Front Immunol* **11**
19. Song J., et al. (2022) **Versican enrichment predicts poor prognosis and response to adjuvant therapy and immunotherapy in gastric cancer** *Front Immunol* **13**
20. Wang M.Q., et al. (2022) **VCAN, expressed highly in hepatitis B virus-induced hepatocellular carcinoma, is a potential biomarker for immune checkpoint inhibitors** *World J Gastrointest Oncol* **14**:1933–1948
21. Yang H., et al. (2022) **Prognostic and immune-related value of complement C1Q (C1QA, C1QB, and C1QC) in skin cutaneous melanoma** *Front Genet* **13**
22. Li Z., et al. (2023) **Differentiation-related genes in tumor-associated macrophages as potential prognostic biomarkers in non-small cell lung cancer** *Front Immunol* **14**
23. Mangogna A., et al. (2019) **Prognostic Implications of the Complement Protein C1q in Gliomas** *Front Immunol* **10**
24. Jiang J., et al. (2020) **Identification of TYROBP and C1QB as Two Novel Key Genes With Prognostic Value in Gastric Cancer by Network Analysis** *Front Oncol* **10**
25. Chen L.H., et al. (2021) **Complement C1q (C1qA, C1qB, and C1qC) May Be a Potential Prognostic Factor and an Index of Tumor Microenvironment Remodeling in Osteosarcoma** *Front Oncol* **11**
26. Zheng Y., et al. (2021) **IRF4-activated TEX41 promotes the malignant behaviors of melanoma cells by targeting miR-103a-3p/C1QB axis** *BMC Cancer* **21**
27. Wang M., Windgassen D., Papoutsakis E.T. (2008) **Comparative analysis of transcriptional profiling of CD3+, CD4+ and CD8+ T cells identifies novel immune response players in T-cell activation** *BMC Genomics* **9**
28. Chen Z., et al. (2021) **A Machine Learning Model to Predict the Triple Negative Breast Cancer Immune Subtype** *Front Immunol* **12**
29. Wang Q., Li P., Wu W. (2019) **A systematic analysis of immune genes and overall survival in cancer patients** *BMC Cancer* **19**
30. Wang J., et al. (2021) **Establishment and validation of immune microenvironmental gene signatures for predicting prognosis in patients with head and neck squamous cell carcinoma** *Int Immunopharmacol* **97**
31. Zhang Y., et al. (2021) **GM-CSF enhanced the effect of CHOP and R-CHOP on inhibiting diffuse large B-cell lymphoma progression via influencing the macrophage polarization** *Cancer Cell Int* **21**
32. Croci G.A., et al. (2021) **SPARC-positive macrophages are the superior prognostic factor in the microenvironment of diffuse large B-cell lymphoma and independent of MYC rearrangement and double-/triple-hit status** *Ann Oncol* **32**:1400–1409
33. Farhood B., Najafi M., Mortezaee K. (2019) **CD8(+) cytotoxic T lymphocytes in cancer immunotherapy: A review** *J Cell Physiol* **234**:8509–8521

34. Roussel M., et al. (2021) **Functional characterization of PD1+TIM3+ tumor-infiltrating T cells in DLBCL and effects of PD1 or TIM3 blockade** *Blood Adv* **5**:1816–1829
35. Xu-Monette Z.Y., et al. (2019) **Immune Profiling and Quantitative Analysis Decipher the Clinical Role of Immune-Checkpoint Expression in the Tumor Immune Microenvironment of DLBCL** *Cancer Immunol Res* **7**:644–657
36. Hirani P., et al. (2021) **Targeting Versican as a Potential Immunotherapeutic Strategy in the Treatment of Cancer** *Front Oncol* **11**
37. Huang X.Y., et al. (2021) **Bioinformatics analysis of the prognosis and biological significance of VCAN in gastric cancer** *Immun Inflamm Dis* **9**:547–559

Article and author information

Zhenbang Ye

Department of Pathology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China;
ORCID iD: [0000-0002-3169-5023](https://orcid.org/0000-0002-3169-5023)

Ning Huang

Department of Hepatobiliary Surgery, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100021, China

Yongliang Fu

Department of Pathology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China;

Rongle Tian

Department of Pathology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China;

Wenting Huang

Department of Pathology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China;; Department of Pathology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital & Shenzhen Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Shenzhen, 518116, China
For correspondence: huangwt@cicams.ac.cn

Copyright

© 2024, Ye et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Mohammad Karimi

King's College London, London, United Kingdom

Senior Editor

Tony Ng

King's College London, London, United Kingdom

Reviewer #1 (Public Review):

Summary:

Ye et al. identified a novel tumour microenvironment (TME) signature that can help to prognosticate DLBCL. They first interrogated a publicly available dataset to identify tumour purity-related genes (TPGs) and found these TPGs were associated with extracellular matrix organisation and immune response. Protein-protein interaction analysis identified hub genes that were associated with prognosis, and 3 genes (VCAN, CD3G, C1QB) were selected to construct a prognosis model. The authors attempted to validate the findings on immunohistochemistry (IHC) and showed prognostication using an IHC assay. Finally, they showed a possible prediction of drug sensitivity using the novel signature in DLBCL.

Strengths:

This study investigated both immune and non-immune TME related to tumour purity. Tumour purity has not been thoroughly investigated in DLBCL. Hence, the prognostic significance of tumour purity demonstrated in this paper brought into light another potential area of research in DLBCL. Similarly, the investigation into non-immune TME was novel and thought-provoking, as most research in DLBCL TME has mostly been in the immune microenvironment.

The bioinformatics approach in identifying the key TPGs was well conducted, such as the GO and KEGG enrichment analysis which supported the role of these TPGs in the modulation of the microenvironment. The findings were also validated in another dataset, which increased the confidence in this model. However, it was not clear to me why the authors chose VCAN, CD3G, and C1QB out of the 9 intersection genes that they found. It would perhaps be useful to show the statistical justification in the Supplementary Results section.

The possible translation of these findings into clinical practice by immunohistochemistry (IHC) was a useful tool to make the findings applicable in the clinical setting. However, as stated by the authors, the real-life clinical application of these findings may be more challenging as these antigens seemed to be expressed in a continuum, rather than in a discrete manner. For example, in Figure 5A, even the low VCAN status still demonstrated strong cytoplasmic staining. Similarly, in Figure 5C, it seemed to be difficult to differentiate strong from background staining. This means pre-analytical variables may affect the staining and standardisation among different laboratories may be difficult to achieve without external controls.

Weaknesses:

Though the rationale behind choosing the TPG genes and its correlation with non-immune TME was clear, the justification for investigating CD68+ macrophages, CD4+ T cells, and CD8+ T cells was not as strong. This was done in a subsection that was supposed to investigate the prognostic values of IHC staining in VCAN, CD3G, and C1QB. Hence, the analysis of the immune compartment of the TME was rather superficial. For example, it would be insufficient to correlate CD4+ and CD8T+ T cells without understanding their deeper

phenotypes such as regulatory vs memory or exhausted vs activated. An attempt was made to subtype the macrophages by bioinformatics approach but it was not further investigated with IHC.

Similarly, the investigation into drug sensitivity was only done in-silico. This investigation was adequate for hypothesis generation. However, it was not enough to substantiate the claim that TPGs can be used to predict drug sensitivity. This claim requires functional in-vitro experiments to validate the bioinformatics approach, or even correlation with clinical data when the identified drugs were used in DLBCL, for example in the ReMODL-B cohort that used bortezomib.

<https://doi.org/10.7554/eLife.92841.1.sa1>

Reviewer #2 (Public Review):

In this study, Zhenbang Ye and colleagues investigate the links between microenvironment signatures, gene expression profiles, and prognosis in diffuse large B-cell lymphoma (DLBCL). They show that increased tumor purity (ie, a higher proportion of tumor cells relative to surrounding stromal components) is associated with a worse prognosis. They then show that three genes associated with tumor purity (VCAN, CD3G, and C1QB) correlate with patterns of immune cell infiltration and can be used to create a risk-scoring system that predicts prognosis, which can be replicated by immunohistochemistry (IHC), and response to some therapies.

1. The two strengths of the study are its relatively large sample size (n = 190) and the strong prognostic significance of the risk-scoring system. It is worth noting that the validation of this scoring with IHC, a simple technique already routinely used for the diagnosis and classification of DLBCL, increases the potential for clinical translation. However, the correlative nature of the study limits the conclusions that can be drawn in regard to links between the risk scoring system, the tumor microenvironment, and the biology of DLBCL.

2. The tumor microenvironment has been extensively studied in DLBCL and a prognostic implication has already been established (for instance, Steen et al., Cancer Cell, 2021). In addition, associations have already been established in non-Hodgkin lymphoma between prognosis and expression of C1QB (Rapier-Sharman et al., Journal of Bioinformatics and Systems Biology, 2022), VCAN (S. Hu et al., Blood, 2013), and CD3G (Chen et al., Medical Oncology, 2022). Nevertheless, one of the strengths and novelty aspects of the study is the combination of these 3 genes into a risk score that is also valid by immunohistochemistry (IHC), which substantially facilitates a potential clinical translation.

3. Figures 1A-B: tumor purity is calculated using the ESTIMATE (Estimation of Stromal and Immune cells in Malignant Tumor tissues using Expression data) algorithm (Yoshihara et al., Nature Communications, 2013). The ESTIMATE algorithm is based on two gene signatures ("stromal" and "immune"). It is therefore expected that tumor purity measured by the ESTIMATE algorithm will correlate with the expression of multiple genes. Importantly, C1QB is included in the stromal signature of the ESTIMATE algorithm meaning that, by definition, it will be correlated with tumor purity in that setting.

4. Figure 2A: as established in Figure 1C, high tumor purity is associated with worse prognosis. Later in the manuscript, it is also shown that C1QB expression is associated with a worse prognosis. However, Figure 2A shows that C1QB is associated with decreased tumor purity. It therefore makes it less likely that the prognostic role of C1QB expression is related to its impact on tumor purity. The prognostic impact could be related to different patterns of immune cell infiltration, as shown later. However, the evidence presented in the study is correlative and natural and not sufficient to draw this conclusion.

5. Figure 3G: although there is a strong prognostic implication of the risk score on prognosis, the correlation between the risk score and tumor purity is significant but not very strong ($R = 0.376$). It is therefore likely that other important biological factors explain the correlation between the risk score and prognosis.

6. Figure 6: the drug sensitivity analysis includes a wide range of established and investigational drugs with varied mechanisms of action. Although the difference in sensitivity between tumors with low and high-risk scores shows statistical significance for certain drugs, the absolute difference appears small in most cases and is of unclear biological significance. In addition, even though the risk score is statistically related to drug sensitivity, there is no direct evidence that the differences in drug sensitivity are directly related to tumor purity.

<https://doi.org/10.7554/eLife.92841.1.sa0>