

The evolution of transposable elements in *Brachypodium distachyon* is governed by purifying selection, while neutral and adaptive processes play a minor role

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2
February 27, 2024 (this version)

Reviewed preprint version 1
December 5, 2023

Posted to preprint server
October 13, 2023

Sent for peer review
October 13, 2023

Robert Horvath , Nikolaos Minadakis, Yann Bourgeois, Anne C. Roulin 

Department of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland • DIADE, University of Montpellier, CIRAD, IRD, Montpellier, France • University House, Winston Churchill Ave, Portsmouth PO1 2UP, United Kingdom

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Understanding how plants adapt to changing environments and the potential contribution of transposable elements (TEs) to this process is a key question in evolutionary genomics. While TEs have recently been put forward as active players in the context of adaptation, few studies have thoroughly investigated their precise role in plant evolution. Here we used the wild Mediterranean grass *Brachypodium distachyon* as a model species to identify and quantify the forces acting on TEs during the adaptation of this species to various conditions, across its entire geographic range. Using sequencing data from more than 320 natural *B. distachyon* accessions and a suite of population genomics approaches, we reveal that putatively adaptive TE polymorphisms are rare in wild *B. distachyon* populations. After accounting for changes in past TE activity, we show that only a small proportion of TE polymorphisms evolved neutrally (< 10%), while the vast majority of them are under moderate purifying selection regardless of their distance to genes. TE polymorphisms should not be ignored when conducting evolutionary studies, as they can be linked to adaptation. However, our study clearly shows that while they have a large potential to cause phenotypic variation in *B. distachyon*, they are not favored during evolution and adaptation over other types of mutations (such as point mutations) in this species.

eLife assessment

This **valuable** study seeks to disentangle the different selective forces shaping the evolutionary dynamics of transposable elements (TEs) in the wild grass *Brachypodium distachyon*. Using haplotype-length metrics, and genetic and environmental differentiation tests, the authors present **convincing** evidence that positive selection on TE polymorphisms is rare and that the distribution of TE ages points to purifying selection being the main force acting on TE evolution in this species. This study will be relevant for anyone interested in the role of TEs in evolution and adaptation.

Introduction

Transposable elements (TEs) are an intrinsic part of eukaryotic genomes and their evolution [1–12]. In addition to modulating genome size, the ability of TEs to create genetic diversity through insertion and excision events can lead to new phenotypes on which selection can act. TEs can alter phenotypes through various mechanisms, including the functional disruption of genes [1, 2], large-scale changes in the regulatory apparatus [3, 4], alteration of epigenetic landscapes [5, 6], ectopic recombination and structural rearrangements [7, 8]. In plants, the dynamics of TE loss and proliferation play a major role in genome evolution [e.g., 9–12]. TEs therefore constitute potentially important drivers of plant evolution, both in nature and during domestication [13].

Beyond their influence on genome structure, and given that their transpositional activity can be stress-inducible [for review 14], TEs are often regarded as more likely than classical point mutations to produce the diversity needed for individuals to respond quickly to challenging environments [15–17]. For instance, punctual TE polymorphisms can lead to gains of fitness and evolve under positive selection [2, 18–23]. TE polymorphisms can even induce more extreme changes in gene expression than single nucleotide polymorphisms (SNPs) in plants [24, 25].

Despite such evidence, whether TE polymorphisms are major contributors to adaptation to changing environments is still debated. Indeed, TE transposition can be disruptive, and purifying selection has been shown to play an important role in TE evolution [e.g., 26, 27]. Based on simulations, it has been suggested that the persistence of TE polymorphisms within a genome without an uncontrolled accumulation, can only be achieved if weak purifying selection is the main force governing TE evolution [28–31]. The uncertainty surrounding this important question in evolutionary genomics results from the limited number of studies that comprehensively tested the extent to which selection shapes TE allele frequencies, both in plants [23, 32] and animals [26, 33–36] and characterized the distribution of fitness effects of new TE insertions. To clarify this question, we used the plant model system *Brachypodium distachyon* [37] to disentangle the effects of purifying and positive selection on TE polymorphisms in natural populations.

B. distachyon is a wild annual grass endemic to the Mediterranean basin and Middle East. Recent genetic studies based on more than 320 natural accessions spanning from Spain to Iraq (hereafter referred to as the *B. distachyon* diversity panel) revealed that *B. distachyon* accessions cluster into three main genetic lineages (the A, B and C genetic lineages), which further divide into five main genetic clades that display little evidence for historical gene flow (Fig. 1A; [38, 39]). Niche modeling analyses suggest that the species moved southward during the last glacial period and recolonized Europe and the Middle East within the last five thousand years [39]. Consequently, while some *B. distachyon* genetic clades currently occur in the same broad geographical areas (Fig. 1A), natural accessions are adapted to a mosaic of habitats [38, 39]. These past and more recent shifts in the species distribution led to clear footprints of positive selection in the genome [39, 40] and make *B. distachyon* an ideal study system to investigate the contribution of TEs to the adaptation of plants in the context of environmental changes.

In *B. distachyon*, TEs are exhaustively annotated and account for approximately 30% of the genome [37]. Recent TE activity has been reported for many families, but despite past independent bottlenecks and expansions experienced by the different genetic clades, no lineage-specific TE family activity has been observed [32]. Rather, TE activity tends to be homogeneous throughout the species range and across genetic clades, indicating a high level of conservation of the TE regulatory apparatus [32]. While purifying selection shapes the accumulation patterns of




Fig 1.

Distribution of the studied accessions and TE polymorphism frequencies. Panel A: Map showing the geographical distribution of the accessions used in the current study. The phylogenetic tree illustrates the phylogeny between the five genetic clades. This panel was made based on the data and results published by Stritt et al. [38] and Minadakis et al. [39]. Panel B: Observed (blue) and simulated (gray) X^tX values of TE polymorphisms in *B. distachyon*. Dotted lines show the 2.5% and 97.5% quantiles of the simulated X^tX values. Panel C-G: Folded site frequency spectrum of TE polymorphisms and synonymous SNPs in all clades. C: A_East; D: A_Italia; E: B_West; F: B_East; G: C.

TEs in this species [32], some TE polymorphisms have been observed in the vicinity of genes [32], potentially affecting gene expression [41]. These early studies, based on a relatively small number of accessions originating exclusively from Spain and Turkey, suggested that TE polymorphisms could contribute to functional divergence and local adaptation in *B. distachyon* [32].

To test this hypothesis, we used the *B. distachyon* diversity panel to identify TE polymorphisms in a large set of 326 natural accessions spanning the whole species distribution. We combined a set of population genomic analyses to assess the proportion of TE polymorphisms associated with positive or purifying selection as well as neutral evolution. We also quantified the strength of purifying selection through forward simulations. Altogether, our work provides the first quantitative estimate, to our knowledge, of the adaptive, neutral, and disruptive potential of TEs, while accounting for changes in TE activity, in a plant harboring a relatively small genome. Altogether, our result advocate against an extended role of TEs in recent adaptation.

Results

Genetic variation in *Brachypodium distachyon*

Using the *B. distachyon* diversity panel (Fig. 1A), we identified 97,660 TE polymorphisms in our *B. distachyon* dataset, of which 9,172 were retrotransposons, 52,249 were DNA-transposons and 36,239 were unclassified. We also identified 9 million SNPs across the 326 samples, including 182,801 synonymous SNPs. A Principal Component Analysis (PCA) performed either with SNPs or TE polymorphisms reflects the previously described population structure of *B. distachyon* [38, 39], with the first two components of the PCA splitting the data according to the demographic structure (Additional file 1: Fig. S1). Investigating the genetic variation caused by retrotransposons and DNA-transposons revealed that the observed diversity in retrotransposons strongly correlated with the demographic structure (Mantel test; $r = 0.79$, p value = 0.001), while the observed diversity in DNA-transposons only had a weaker correlation (Mantel test; $r = 0.36$, p value = 0.001) with the demographic structure (Additional file 1: Fig. S2).

From the initial TE and SNP dataset, we could estimate the time of origin in generations (age) of 50,891 TE polymorphisms and 108,855 synonymous SNPs based on pairwise differences in identity by descent (IBD) regions around the focal mutation (see Materials and methods). The results of the age estimate analysis were checked by contrasting the observed correlation between allele age and frequency of synonymous SNPs to the theoretical predictions of Kimura and Ohta [42] for neutrally evolving mutations. We found that, the observed correlation matched expectations (Additional file 1: Fig. S3), with older alleles found on average at higher frequencies than younger ones. Furthermore, most TE polymorphisms in our dataset were young and only a few were very old (Additional file 1: Fig. S4).

The overall contribution of TEs to clade differentiation and adaptation is limited

To examine the overall contribution of TEs to evolution and adaptation in *B. distachyon*, we first identified regions of the genomes that were likely affected by recent selective sweeps. The fast increase in the frequency of a beneficial allele is expected to lead to a longer than average haplotype around the mutation under positive selection. Such events (known as selective sweeps) can be identified by computing the integrated haplotype score (iHS) around focal mutations [43]. We therefore computed iHS along the genome for the four derived genetic clades. Regions of the genomes with significantly higher iHS than average are expected to harbor mutations that were under positive selection during evolution and adaptation. We hypothesized that if TEs

constitute an important part of the genetic makeup that led to adaptation in a given genetic clade, then they should be more frequently fixed or at higher frequencies in regions with high iHS than in the corresponding regions that did not experience recent selective sweeps in other clades.

First, we tested if more TE polymorphisms were fixed in a specific region of the genome if a genetic clade had a high iHS, and presumably experienced a selective sweep, than in other genetic clades. An analysis of covariance (ANCOVA) revealed that the number of fixed TE polymorphisms per clade did not significantly differ between high iHS regions and the same regions in other clades (**Table 1** [↗](#)). These results indicate that there is no correlation between the overall number of fixed TE polymorphisms per clade in a region and recent selective sweeps. However, the number of fixed TEs in genomic regions along the genome was significantly affected by the total number of TEs in the region, the TE superfamily, the TE age, the genetic clade and the overall genetic features of the region (e.g., recombination rate, see Materials and methods) but not by the iHS itself (**Table 1** [↗](#)). Similarly, we tested if the allele frequency of TE polymorphisms was significantly higher in a specific region of the genome if a genetic clade had a high iHS than in other genetic clades. A second ANCOVA revealed that the allele frequency of TE polymorphisms was significantly influenced by the TE superfamily, TE age, clade and overall genetic features of the region but not by the iHS (**Table 2** [↗](#)). These results indicate that TEs in high iHS regions did not experience a significant increase in their frequency and that TEs in high iHS regions are experiencing the same selective constraints as other TEs. Similar results were obtained when investigating the number of fixed TE polymorphisms (Additional file 2: Table S1) and the allele frequency of TE polymorphisms (Additional file 2: Table S2) in high iHS regions using a subset of our dataset with an expected lower false negative TE call rate, that only included samples with a genome-wide mapping coverage of at least 20x (see Discussion and Materials and Methods for more details).

A complementary approach to explore the impact of positive selection on TEs consists in investigating their genetic differentiation among populations. Using the five genetic clades as focal populations, we computed X^tX values, a standardized measure of genetic differentiation corrected for the neutral covariance structure across populations [[44](#) [↗](#), [45](#) [↗](#)], for each TE polymorphism. Mutations affected by positive selection are expected to be over-differentiated between clades and display significantly higher X^tX values than other mutations [[45](#) [↗](#)]. In contrast, a low X^tX value implies that the mutation is less differentiated than other mutations and potentially evolves under balancing selection, whereas purifying selection and a neutral evolution are not expected to impact the differentiation of a mutation among populations [[44](#) [↗](#)]. We contrasted the observed X^tX values computed for each TE polymorphism to a simulated pseudo-observed dataset (simulated observations under the demographic model inferred from the covariance matrix of the SNP dataset, for more details see [[45](#) [↗](#)]) and found that only a small fraction of the TE polymorphisms (0.06%) displayed X^tX values higher than the 97.5% quantile of the simulated values (**Fig. 1B** [↗](#)). This indicates that only a few TE polymorphisms are over-differentiated among genetic clades and might have been affected by positive selection. However, a relatively larger portion of the TE polymorphisms (4.3%) displayed X^tX values smaller than the 2.5% quantile of the simulated values (**Fig. 1B** [↗](#)), indicating that balancing selection might also shape TE frequency in *B. distachyon*.

To further examine the contribution of TEs to adaptation, we tested whether and how many TE polymorphisms were significantly associated with environmental factors. If the presence of a TE provides an advantage in a certain environment and contributes to adaptation, we expected a correlation between the environment and the presence/absence of this TE. In this context, we performed genome-environment association analyses (GEA) using all TEs and SNPs identified across the 326 samples and 32 environmental factors associated with precipitation, solar radiation, temperature, elevation and aridity (see in Materials and methods for the full list). The GEA revealed that only nine of the 97,660 TE polymorphisms were significantly associated with some environmental factors (Additional file 2: Table S3), confirming that TEs only had a limited

Variable	Sum of squares	degrees of freedom	<i>F</i> value	<i>P</i> value
Total number of TEs in the region	28969.6	1	35405.64	< 0.001
TE superfamily	887.5	14	77.48	< 0.001
Clade	587	3	239.13	< 0.001
Genomic region	136.7	80	2.09	< 0.001
TE age	45.5	2	27.81	< 0.001
High iHS	0	1	0.03	0.869

Table 1

ANCOVA predicting the number of fixed TE polymorphisms per clade in candidate regions under positive selection.

Variable	Sum of squares	degrees of freedom	<i>F</i> value	<i>P</i> value
TE superfamily	453.2	14	247.3	< 0.001
Clade	17.7	3	45.18	< 0.001
Genomic region	147	80	14	< 0.001
TE age	2	2	7.7	< 0.001
High iHS	0.1	1	0.79	0.374

Table 2

ANCOVA predicting the allele frequency of TE polymorphisms per clade in candidate regions under positive selection.

contribution to adaptation in *B. distachyon*. Importantly, two of these nine TEs were found in a gene, and three were in the vicinity of genes (less than 2 kilobase (kb) away, Additional file 2: Table S3).

Purifying selection dominates the evolution of TE polymorphisms in *B. distachyon*

To further characterize the forces governing the evolution of TE polymorphisms in *B. distachyon*, we examined the genome-wide frequency distribution of TEs. We first computed the folded site frequency spectrum (SFS) and found that the folded SFS of TE polymorphisms was shifted toward a higher proportion of rare minor alleles compared to neutral sites in all genetic clades (**Fig. 1C-G**). Splitting the TE data into DNA-transposons and retrotransposons resulted in similar folded SFS and shifts in both TE classes (Additional file 1: Fig. S5 and S6).

These shifts could be the result of purifying selection as the analyses presented above indicate that positive selection has a negligible effect on TE polymorphism frequencies in *B. distachyon*. However, in contrast to SNPs, TEs do not evolve in a clock-like manner, as their transposition rate is known to vary between generations [46, 47]. Changes in transposition rate and purifying selection can lead to similar shifts in the SFS but can be disentangled using age-adjusted SFS [48]. In brief, if TE polymorphisms are evolving neutrally, they are expected to accumulate on average at the same rate in a population as neutral SNPs of the same age. Hence, Δ frequency, the difference between the average frequency of TE polymorphisms and neutral sites in a specific age bin, will remain close to 0 regardless of the polymorphisms' age. In contrast, if TE polymorphisms evolve under purifying selection, they will tend to occur at lower frequencies than neutral SNPs of the same age, as selection will prevent them from accumulating in the population. Consequently, the Δ frequency will reach negative values for older TE polymorphisms [48].

Because this model does not allow for back mutations, as typically observed for DNA-transposons that can excise from the genome, we primarily investigated the age-adjusted SFS of retrotransposons in the four derived clades. This analysis revealed that retrotransposons are indeed prevented by natural selection from randomly accumulating, as older retrotransposons are significantly less frequent than neutral SNPs of the same age (**Fig. 2**; one-sided Wilcoxon test, Bonferroni corrected p value < 0.01).

As previous studies showed that the distance between TE polymorphisms and the next gene can impact the strength of selection affecting TEs [6, 49, 50], we further split our retrotransposon polymorphisms into three categories based on their distance to the next gene: retrotransposons (i) in and up to 1 kb away from genes, (ii) between 1 kb and 5 kb away and (iii) more than 5 kb away. The age-adjusted SFS of all three categories displayed the same pattern as that observed for the whole retrotransposon polymorphism dataset: older retrotransposon polymorphisms were significantly less frequent than neutral sites of the same age regardless of their distance to genes (one-sided Wilcoxon test, Bonferroni corrected p value < 0.01), indicating that retrotransposons more than 5 kb away from genes are also affected by purifying selection (**Fig. 2**).

Retrotransposon polymorphisms tended to be more deleterious than SNPs predicted to have a high impact on fitness. Indeed, the age-adjusted SFS of retrotransposons resulted in a larger deviation of Δ frequency from 0 than for non-synonymous SNPs and high effect SNPs (**Fig. 2**). In addition, Δ frequency in the oldest (last) age bin was significantly more negative than in all other age bins in the A_East, B_East and B_west clades (one-sided Wilcoxon test, Bonferroni corrected p value < 0.01). In the A_Italia clades the oldest age bin was not significantly different from the second oldest age bin (two-sided Wilcoxon test, Bonferroni corrected p value N.S.). While older non-synonymous SNPs and high effect SNPs were generally less frequent than neutrally evolving SNPs at the same age, the negative Δ frequency trend was reversed for the oldest non-synonymous SNPs and high




Fig. 2

Age-adjusted SFS of retrotransposons. The top row shows the age-adjusted SFS of all retrotransposons (colored), non-synonymous SNPs (light gray) and high effect SNPs (dark gray) in the four derived clades. The bottom row shows the age-adjusted SFS of retrotransposons based on their distance to the next gene in the four derived clades. The X axes show the age range of the mutations in each bin, and the age range of each bin was chosen so that each bin represents the same number of retrotransposon observations in the top row. The different columns show the four derived clades: A: A_East; B: A_Italia; C: B_West; D: B_East. Boxplots are based on 100 estimations of Δ frequency. Significant deviations of Δ frequency estimates from 0 in the age-adjusted SFS of retrotransposons are shown with asterisks (one-side Wilcoxon tests, Bonferroni corrected p value < 0.01 ; ***).

effect SNPs (**Fig. 2**). In all clades, Δ frequency in the oldest age bin was significantly higher than at least the lowest Δ frequency observed in the other age bins for non-synonymous SNPs, as well as high effect SNPs (one-sided Wilcoxon test, Bonferroni corrected p value < 0.01). This might be because not all predicted non-synonymous SNPs and high effect SNPs might result in fitness effects. Those SNPs can therefore evolve neutrally or nearly neutrally and persist as polymorphic SNPs much longer in a population than those affecting fitness negatively. Hence, even the oldest retrotransposon polymorphisms seem to be mostly non-neutral and are affected by purifying selection.

To assess whether similar forces may drive retrotransposon and DNA-transposon evolution, we repeated the analysis for DNA-transposons. The age-adjusted SFS of DNA-transposons revealed very similar patterns, with Δ frequency showing significant deviations from 0 in older age bins (one-sided Wilcoxon test, Bonferroni corrected p value < 0.01), but DNA-transposon polymorphisms seemed less deleterious than non-synonymous SNPs and high effect SNPs (Additional file 1: Fig. S7).

Forward simulations allow us to quantify the strength of purifying selection

To evaluate to what extent the proportion of neutrally evolving mutations in the focal group of mutations affects the shape of the age-adjusted SFS, we ran forward simulation with mutations under multiple selective constraints, and we tested what ratio of neutral to selected mutations can lead to an age-adjusted SFS similar to that observed for retrotransposons in *B. distachyon*. Specifically, we investigated the conditions under which we observed a Δ frequency in the oldest age bin significantly smaller than Δ frequency in all other age bins. Our simulations revealed that the shape of the age-adjusted SFS of retrotransposons could only be reproduced if less than 10% of the mutations were neutrally evolving for most of the selective constraint investigated (Additional file 1: Fig. S8 and Additional file 2: Table S4).

Finally, we used the results from our simulations to narrow down the selection strength affecting retrotransposons in *B. distachyon* by investigating the age of the oldest retrotransposons in our dataset. The main difference between the age-adjusted SFS of mutations evolving under weak and strong purifying selection is that the oldest mutations are much older in the simulation with weak purifying selection than in the simulation with strong purifying selection. This age difference arises because mutations under strong purifying selection are removed from the population more effectively and, therefore, cannot persist as long in the population. Examining the age of the last retrotransposon bins in the age-adjusted SFS revealed that the ages of the oldest retrotransposons were the most similar to the expected ages of the oldest mutations in our simulations, with a scaled selection coefficient (S) of -5 and -8 (**Fig. 3**), indicating that retrotransposons in *B. distachyon* are under moderate purifying selection. In simulations with a nearly neutral selection coefficient ($S = -1$), the simulated mutations were much older than the oldest observed retrotransposons (**Fig. 3**). Conversely, in simulations with a strong purifying selection coefficient ($S < -10$), they were much younger than the oldest observed retrotransposons (**Fig. 3**).

Discussion

B. distachyon is a widely used model species in evolutionary genomics, molecular ecology, developmental biology and crop functional genomics [for review 51, 52] with past and ongoing TE movements in its genome [32]. In this study, we used a diversity panel containing next-generation sequencing data from over 320 individuals sampled across the whole geographical range of *B. distachyon* to examine the role of TEs during evolution and adaptation. We investigated the frequency with which positive selection led to an increase in the frequency and fixation of TEs and quantified the strength of purifying selection on TE polymorphisms. Accounting for




Fig. 3

Relative age difference ((mutation age in simulations - observed mutation age)/maximum absolute age difference) between simulated and observed data in the last bin of the age-adjusted SFS. A: 25% quantile; B: 50% quantile; C: 75% quantile.

population structure and fluctuant transposition rates, we demonstrate that TEs are rarely part of the genetic makeup that was positively selected during environmental adaptation in *B. distachyon*. Furthermore, we show that the majority of TE polymorphisms found in the natural population of this model species are under weak to moderate purifying selection, with only a small minority of TE polymorphisms evolving neutrally.

Rare instances of positive selection on TEs

By combining complementary approaches, we were able to demonstrate that TEs are rarely the target of positive selection in *B. distachyon*. We first probed for footprints of positive selection on TE polymorphisms using the five genetic clades as focal populations. In conducting this analysis, we did not find TE polymorphisms to be at high frequencies or fixed at higher rates than expected, in regions of the genome presumably harboring selective sweeps in at least one of the genetic clades (high iHS regions). This suggested that TEs were rarely the target of positive selection, which we confirmed with a genome-wide scan for overly differentiated TE polymorphisms using X^tX analysis. Indeed, this approach revealed that only a very small proportion of TE polymorphisms are more differentiated than expected under a neutral scenario.

Importantly, the X^tX analysis also revealed that a non-negligible fraction of the TE polymorphisms is less differentiated than expected and are shared among genetic clusters. This could be the result of selection favoring the same TE polymorphisms in different accessions to adapt to similar environmental constraints across genetic clades. To test this scenario, we performed GEA with 32 environmental factors, and found only nine TE polymorphisms significantly associated with any of these, and representing a very small proportion ($< 0.01\%$) of all the TE polymorphisms we identified. Interestingly though, these nine TE polymorphisms were associated with environmental variables pertaining to precipitation, temperature and altitude, which are known to drive adaptation in *B. distachyon* [39]. Some insertions were found within or in close proximity of genes, making these polymorphisms very good candidates for future functional validation.

Single TE insertions can have a drastic impact on phenotypic variation and be affected by positive selection [for review, see 16, 53, 54]. For instance, TEs have increased in frequency through positive selection in humans [23] or during range expansion in *Arabidopsis* [25] and *D. melanogaster* [20, 22]. Evidently, *B. distachyon* exhibits a different pattern, as causal mutations for adaptation in this grass species are rarely TEs. Only a few studies have thoroughly quantified the extent to which positive selection influences the evolution of TEs [25, 26, 28, 30, 31]. But two of these drew similar conclusions to us, in the green anole *Anolis carolinensis* [30] and in the invasive species *Drosophila Suzukii* [31]. In addition, a large number of candidate genes for adaptation were identified with a similar approach focusing on SNPs [39], indicating that population structure or demographic events are not limiting factors for the methods we used. Altogether, these observations call for a closer investigation of which forces, e.g., purifying selection or neutral evolution, are important in shaping TE allele frequency in natural populations.

Moderate purifying selection is the dominant force during TE evolution

Our results suggest that purifying selection is an important factor limiting the ability of TE polymorphisms to fix and increase their frequency in *B. distachyon*. This finding that purifying selection is the main force shaping the landscape of TE polymorphisms in *B. distachyon* is in line with similar observations made for example in maize [55], *Arabidopsis thaliana* [56, 57] and *Drosophila simulans* [58]. Indeed, one of the significant explanatory variables in our ANCOVA models was the genetic clade, a proxy for the effective population size (N_e), which affects the efficiency with which selection can fix beneficial mutations and purge deleterious ones. In *B. distachyon*, the number of fixed TE polymorphisms per clade and the frequency of TE

polymorphisms were negatively correlated with N_e , indicating that the accumulation of TEs is significantly lower in genetic clades with a larger N_e , potentially because of a greater efficacy of purifying selection.

It is widely accepted that most new TE insertions have a deleterious or no effect on the fitness of the host [22, 26, 28, 30, 33–36, 58]. To properly quantify the effect of purifying selection on TE evolution in *B. distachyon*, we used age-adjusted SFS analyses to evaluate the selective constraint experienced by TE polymorphisms while accounting for previously reported changes in their activity [32]. While this method can only be applied to retrotransposons (because the model does not allow back mutations), it provided a first clue on the importance of purifying selection on TE evolution and revealed that overall, retrotransposons evolved under purifying selection in all four derived genetic clades. Indeed, the Δ frequency was significantly smaller than 0, especially for older retrotransposons, meaning that old retrotransposons are less common than neutrally evolving SNPs at the same age. This further demonstrates that even after accounting for the different genetic clades and using a large sample size, retrotransposons evolve under purifying selection in *B. distachyon*. One caveat of the approach used in this study is that TE calling pipelines based on short-reads tend to have higher false positive and false negative call rates than SNP calling pipelines, which is also the case for the TEPID TE calling pipeline used here [57, 59]. A high false negative TE calling rate however might bias our TE frequency estimates toward lower frequencies, which could drive the observed patterns in the age-adjusted SFS. To assess if the false negative TE calling rate in our study substantially affected our results, we re-run the age-adjusted SFS on a subset of our dataset only including samples with a genome-wide mapping coverage of at least 20x, as higher mapping coverages are expected to reduce the false negative call rate [27, 59]. Using the TE allele frequencies estimated based on this subset of our data to estimate Δ frequency revealed similar results of the age-adjusted SFS based on the whole dataset (Additional file 1: Fig. S9), indicating that our observation of retrotransposons evolving under purifying selection is not solely driven by a high false negative TE calling rate.

We also revealed that only a minority of retrotransposons evolved neutrally, as the observed shape of the Δ frequency curve could only be reproduced in our simulation if the proportion of neutrally evolving mutations in our focal mutations was below 10%. This estimate gives a first glimpse into the distribution of fitness effects of new TE insertions, a fundamental parameter in genetics that describes the way in which new TE insertions can contribute to evolution and adaptation [60]. Here, we show for the first time that new TE insertions have a less than 10% chance to insert into the genome of *B. distachyon* in a way that will allow them to evolve neutrally, advocating for a large potential of TEs to create, through their movement, new phenotypic variation on which selection can act on. PCAs based on TE polymorphisms allowed us to recover the population structure of *B. distachyon*, implying that demographic history and hence neutral processes may indeed partially explain the differences in the TE distribution we observed between genetic clades, as shown in *Arabidopsis thaliana* and *Arabidopsis lyrata* [26, 61], *Drosophila melanogaster* [62], humans [63] and the green anole (*Anolis carolinensis*; [30]). However, and in line with our simulations, the first two axes of the PCA explain less than 7% of the variance, indicating that neutrally evolving TEs contribute only mildly to overall TE diversity in our system.

Because TEs can cause phenotypic variation through new insertions [1–8], it is not surprising that most new insertions interfere with the function of the genome, especially in a species with a small genome, such as *B. distachyon* (272 Mb) [37]. The proportion of neutrally evolving TE polymorphisms is expected to be very small in genes, as insertions in genic regions are likely to result in loss-of-function [1, 2]. Similarly, TE insertions in close proximity to genes are expected to be highly disruptive, as regulatory elements such as *cis*-regulatory elements are predominantly located in the proximity of genes. In *A. thaliana*, for instance, TEs located in the vicinity of genes (less than 2 kb) globally result in downregulation [64]. Although only specific families alter gene expression in *B. distachyon* [41], the observed Δ frequency for retrotransposon polymorphisms in genes and in their 1 kb surroundings matched our

expectations. The fact that TE polymorphisms located more than 5 kb away from genes are also evolving under purifying selection was more surprising. That said, little is known about the distance between *cis*-regulatory sequences and genes in *B. distachyon*. In plants, TEs are believed to affect gene expression in *trans* through the production of small-interfering RNA [65–69]. Hence, the fact that only a small proportion of TEs can accumulate neutrally indicates that, in a gene-dense genome such as that of *B. distachyon* (42.5% of the genome are genes) [69], TE insertions in any genomic compartment may result in some *cis*- or *trans*-regulatory effects visible to selection. Finally, we tested whether TE polymorphisms located more than 5 kb away from genes are evolving under purifying selection could be due to mapping or other artefacts by comparing the shape of the age-adjusted SFS of retrotransposons and SNPs more than 5 kb away from genes. However, the age-adjusted SFS of SNPs 5 kb away from genes differs from the one of retrotransposons (Additional file 1: Fig. S10), indicating that the shape of the age-adjusted SFS of retrotransposons more than 5 kb away from genes is not likely to be the result of artefacts in regions of the genome far away from genes.

To further ascertain the strength of purifying selection, we used forward simulation and showed that simulations assuming a moderately weak selection pressure ($S = -5$ or $S = -8$) against TE polymorphisms best fitted our observed data. In theory, no TE polymorphisms under strong purifying selection should be present in a natural population, as such mutations are expected to be quickly lost, especially in a predominantly selfing species where most loci are expected to be homozygous. Therefore, it is not surprising that TE polymorphisms which persist in *B. distachyon* are under weak to moderate selection, as also shown, for example, for the L1 retrotransposons in humans [27] or the BS retrotransposon family in *Drosophila melanogaster* [62].

While some of the parameters we chose for our simulations, such as the dominance or selfing rate, can affect the efficiency of TE purging, it is unlikely that discrepancies in the true and assumed values for these parameters would have led to drastically different results. For example, we assumed codominance for all mutations, which might not hold true for each TE polymorphism. However, because of the high selfing rate observed in *B. distachyon* [38], heterozygous loci are expected to be rare, and dominance is unlikely to have a strong impact on our observations. Similarly, with a higher selfing rate, deleterious TE polymorphisms should be removed more efficiently by purifying selection. To check whether a lower selfing rate could allow a higher proportion of TE polymorphisms to evolve neutrally, we reran the simulations assuming fully outcrossing individuals. This also resulted in simulation with weak to moderate selection strength on TE polymorphisms best fitting the observed data, further strengthening our results (Additional file 1: Fig. S11).

While the analyses of positive selection and GEA were based on both DNA-transposons and retrotransposons, we only used retrotransposons to assess the strength of selection on TE polymorphisms, as the age-adjusted SFS was developed with the assumption of no back mutations [48]. Yet, DNA-transposons do not solely transpose through cut and paste mechanisms as they would otherwise not be so abundant in Eukaryotic genomes. DNA-transposons can also create extra copies of themselves by transposing during chromosome replication or repair from a position that has already been replicated, or repaired [70]. We therefore repeated the age-adjusted SFS analyses using DNA-transposons to evaluate whether DNA-transposons were affected by similar selective constraints. The folded SFS of DNA-transposons and retrotransposons display similar shifts toward high proportions of rare alleles and Δ frequency deviations from 0 in the age-adjusted SFS of DNA-transposons and retrotransposons are comparable. Hence, we argue that the conclusion drawn for retrotransposons also holds for DNA-transposons, and that purifying selection affect TEs broadly. To further examine our conclusion on purifying selection, we investigated the selective regime affecting different retrotransposons and DNA-transposons superfamilies. Thereby, we generated age-adjusted SFS for the four most common TE superfamilies Copia, Ty3 (also known under the name Gypsy, but we will avoid using this name because of its problematic nature see [71]), Helitron and MITE and found similar deviations of

the Δ frequency from 0 in the four investigated TE superfamilies (Additional file 1: Fig. S12–S15). These results indicate that our conclusion on the broad effect of purifying selection is not driven by a single TE superfamily but is at least common among the four most numerous TE superfamilies.

Conclusion

Adaptation to different environmental conditions is a complex process that involves various mutation types. Here, we show that the vast majority of TE polymorphisms are under purifying selection in the small genome of *B. distachyon*. Conversely, only a very small proportion of TEs seem to have contributed to adaptation. The observed lack of neutrally evolving TE polymorphisms in *B. distachyon* advocates for a large potential of TE polymorphisms to contribute to the genetic diversity and phenotypic variation on which selection can act and highlights the need to consider TE polymorphisms during evolutionary studies. Finally, our work shows that the ability of TEs to cause phenotypic variation does not necessarily translate into being favored during evolution and adaptation over other mutations with more subtle effects, such as SNPs.

Materials and methods

Whole-genome resequencing data

In this study, we analyzed a total of 326 publicly available whole-genome sequencing data from *Brachypodium distachyon* accessions sampled around the Mediterranean Basin (Fig. 1A; Additional file 2: Table S5). Our *B. distachyon* dataset consisted of 47 samples published by Gordon et al. [8], 57 samples published by Skalska et al. [72], 65 samples published by Gordon et al. [73], 86 samples published by Stritt et al. [38] and 71 samples published by Minadakis et al. [39], covering all five genetic clades previously described in this species [38, 39]. Each sample was assigned to a genetic clade based on previously published results [39].

Data processing

Raw reads were trimmed using Trimmomatic 0.36 [74] and mapped to the *B. distachyon* reference genome version 3.0 [37] using bowtie2 [75] and yaha [76], and TE polymorphisms were identified using the TEPID pipeline [59] and the recently updated TE annotation by Stritt et al. [77] and Wyler et al. [69]. TE polymorphisms include both TE insertion polymorphisms (TIPs; insertions absent from the reference genome but present in at least one natural accession) and TE absence polymorphisms (TAPs; insertions present in the reference genome but absent from at least one natural accession). The class, superfamily and family of each TE call were assigned based on the TEPID results and the TE annotation from the reference genome. TIPs that were less than 100 base pairs (bp) apart in different samples and assigned to the same TE family were merged.

Single nucleotide polymorphisms (SNPs) were called using GATK v.4.0.2.1 [78] using HaplotypeCaller [79] following Minadakis et al. [39]. The SNP calls were hard filtered using the following conditions: QD < 5.0; FS > 20.0; SOR > 3.0; MQ < 50.0; MQRankSum < 2.5; MQRankSum > -2.5; ReadPosRankSum < 2.0; ReadPosRankSum > -2.0. Because *B. distachyon* displays a high selfing rate [38], most genetic variants are expected to be homozygous within an individual. Hence, all TE calls were treated as homozygous, and heterozygous SNP calls were removed from our dataset to reduce false variant calls. Additionally, all sites with multiallelic TE and SNP calls were removed. SNPs were classified as synonymous, non-synonymous and of high fitness effect using SnpEff [80]. SNPs and TE polymorphisms were merged into a single vcf file using custom scripts provided in github (see section Availability of data and materials).

To estimate the age of each SNP and TE polymorphism, the SNPs and TEs found in the A_East, A_Italia, B_East and B_West clades were polarized using the C clade, which was identified as the most ancestral *B. distachyon* clade [38]. GEVA was run separately for each clade using the genetic map produced by Huo et al. [82] and a mutation rate of 7×10^{-9} substitutions/generation. The theoretical prediction of the correlation between allele age and allele frequency of neutrally evolving mutations based on N_e [42] was compared to the observed correlation between allele age and frequency of synonymous SNPs to check the sanity of the age estimates.

The observed SNP and TE diversity was first examined using a principal component analysis (PCA), and correlations between TE diversity and genetic clades were tested with a mantel test using the ade4 package version 1.7-22 [83] in R version 4.1.2 [84]. The folded site frequency spectrum (SFS) was computed for TE polymorphisms and SNPs using the minor allele frequency in R version 4.1.2 [84]. Finally, the map of the geographical distribution of the used accessions was done in R using the rnatuarearth package 0.3.3 [85].

Mapping coverage is known to influence false discovery rate [27, 59]. To investigate the impact of false positive and false negative TE calls on our results, we down sampled the TE dataset to only include TEs that have been called in samples that had at least an average mapping coverage of 20x. The allele frequencies of TEs present in our high coverage dataset was recalculated only considering samples with at least an average mapping coverage of 20x. This second TE dataset was then used to check if using a dataset with a higher mapping coverage and presumably a lower false TE calling rate impacted our results.

Analyses of positive selection

Regions of the genome affected by positive selection were identified using the integrated haplotype score (iHS), a measure of the amount of extended haplotype homozygosity along the ancestral allele relative to the derived allele for a given polymorphic site [86]. iHS was calculated using the SNP dataset, and regions displaying longer haplotypes and hence high iHS were identified in R using the rehh package [87, 88]. The threshold to distinguish between regions of high iHS and other regions was selected such that less than 5% of the *B. distachyon* genome was classified as high iHS regions in each clade (Additional file 2: Table S6). Candidate regions under positive selection were defined as all regions that were found to have high iHS in each clade separately.

A first ANCOVA was used to model the number of fixed TE polymorphisms in each clade found in the candidate region under positive selection based on the following genetic features: total number of TEs, TE superfamily, TE age (split into three categories: young: age < 10,000 generations; intermediate: age between 10,000 generations and 60,000 generations; old: age > 60,000 generations), clade, genomic region (a unique ID for each candidate region under positive selection) and iHS classification of the regions in each clade (high or average). A second ANCOVA was used to model the allele frequency of TE polymorphisms found in the candidate region under positive selection based on the following genetic features: TE superfamily, TE age, clade, genomic region and iHS classification of the regions in each clade. The TE superfamily was included to account for different evolutionary behaviors of TEs from different superfamilies. Age accounted for differences in the fixation rate and frequency distribution between young and old TEs. The clade was included to account for clade-specific differences such as differences in N_e . Finally, a unique ID for each candidate region under positive selection was included to account for region-specific differences such as differences in the recombination rate and GC content. In the end, regions that were found to have a high iHS in some clades were compared to the same regions in the other clades. All ANCOVAs were run in R using the car package [89].

The standardized allele frequency of a mutation across populations (X^tX) values [44] were computed for the combined TE and SNP dataset using Baypass version 2.3 [45, 90]. The X^tX values were used to identify over- and under differentiated TE polymorphisms between clades. A pseudo-observed dataset (POD) of 100,000 SNPs was simulated under the demographic model inferred from the covariance matrix of the SNP dataset. The POD was then used to determine the 97.5% (over-differentiated polymorphisms) and 2.5% (under differentiated polymorphisms) quantiles.

Genome-environment association analyses

We identified TE polymorphisms significantly associated with environmental factors using genome-environment association analyses (GEA) following Minadakis et al. [39]. GEAs were run with GEMMA 0.98.5 [91] using the combined TE and SNP vcf file against the following 32 environmental factors extracted by Minadakis et al. [39]: altitude, aridity from March to June, aridity from November to February, annual mean temperature, mean temperature of warmest quarter, mean temperature of coldest quarter, annual precipitation, precipitation of wettest month, precipitation of driest month, precipitation seasonality, precipitation of wettest quarter, precipitation of driest quarter, precipitation of warmest quarter, precipitation of coldest quarter, mean diurnal Range, isothermality, temperature seasonality, maximum temperature of warmest month, minimum temperature of coldest month, temperature annual range, mean temperature of wettest quarter, mean temperature of driest quarter, precipitation from March to June, precipitation from November to February, solar radiation from March to June, solar radiation from November to February, mean temperature between March and June, mean temperature between November and February, maximum temperature between March and June, maximum temperature between November and February, minimum temperature between March and June and minimum temperature between November and February. We applied a False Discovery Rate (FDR, [92]) threshold of 5% to control for false positive rates.

Age-adjusted frequency spectra and analyses of purifying selection

Footprints of purifying selection on TE polymorphisms were first evaluated using folded SFS. An age-adjusted site frequency spectrum (age-adjusted SFS) approach was used to further investigate the impact of purifying selection on retrotransposons while accounting for nonconstant transposition rates. Briefly, the age-adjusted SFS is a summary statistic that describes the difference between the average frequency of TEs at a specific age and the average frequency of neutral sites of the same age [48]. Therefore, the TE dataset was sorted by age and split into equally large bins with respect to the number of observations in each age bin. Neutral sites were then randomly down-sampled to match the number of observations in the TE dataset and its age distribution [48].

The difference between the average TE and neutral site frequency, or Δ frequency, was computed for each age bin [48]. This method allows for an unbiased comparison between the allele frequencies of TEs and neutral sites, and is robust to transposition rate changes and demographic changes [48]. However, the theory behind this method was developed assuming no back mutations and is therefore best suited for retrotransposons, as DNA-transposons can exit an insertion site [48]. We used the synonymous SNPs identified with SnpEff as the neutrally evolving sites. However, because estimating the population wide frequency of TEs is more challenging than estimating SNP frequencies, putative biases in frequency estimates need to be assessed before performing age-adjusted SFS analyses. To do so, the SNP dataset was resampled so that the SNP dataset used in the age-adjusted SFS had a frequency distribution that matched the observed TE frequency distribution. The age-adjusted SFS of retrotransposons was contrasted against the age-adjusted SFS of non-synonymous, as well as against high fitness effect SNPs. Therefore, 10,000 non-synonymous and high fitness effect SNPs were randomly selected for each clade to reach approximately the same number of retrotransposon polymorphisms, non-

synonymous and high fitness effect SNPs for final comparisons. To estimate the variation in Δ frequency estimates, all age-adjusted SFS were computed 100 times. All Wilcoxon tests and Bonferroni p value corrections were done in R version 4.1.2 [84].

Forward simulation

We used SLiM 4.0.1 [93, 94] to run forward simulations and assess the proportion of neutrally evolving retrotransposons and the average selection strength affecting them. The simulations were designed to reflect the population size and demographic history of *B. distachyon*. The simulated genomic fragment was 1 megabase (Mb) long and included neutral (synonymous) mutations as well as focal mutations that evolved under different selective constraints. The focal mutations were a mix of neutrally evolving mutations and mutations evolving under a constant selection pressure. Therefore, the ratio (r) of focal mutations that evolved neutrally was either 0%, 5%, 10%, 25% or 50%. The scaled selection coefficient (S , defined as $N_e s$, with s the strength of selection and N_e the effective population size) affecting the remaining focal mutations was set at the beginning of the simulation to be either -1, -5, -8, -10, -12, -15, -20 or -50 to cover effectively neutral ($0 > S \Delta -1$), intermediate ($-1 > S \Delta -10$) and strongly deleterious ($-10 > S$) selective constraints. The selfing rate was set to 70%, as *B. distachyon* is a highly selfing species with occasional outcrossing [38, 39]. In addition, a high recombination rate was chosen to minimize the effects of linked selection in the small genomic fragment simulated. Simulations for each combination of these two parameters were run 20 times to assess the variation in the resulting age-adjusted SFS. The shape of the resulting age-adjusted SFS was used to narrow down the ratio of neutrally evolving TE polymorphisms. Similarly, the age distribution of the mutations in the oldest bin of the age-adjusted SFS was used to narrow down the strength of selection affecting TE polymorphisms.

Acknowledgements

We would like to thank Jeffrey Ross-Ibarra and Mitra Menon as well as Fabrizio Menardo, Michael Thieme, Wenbo Xu, Jigisha, Lars Kaderli and Serafin Schefer for all the discussions and their comments on this project. We thank Emmanuelle Botté (<https://manuscribe.com.au>) for professional editing of the manuscript.

Declarations

Consent for publication

All authors have read and approved the submission of this manuscript.

Availability of data and materials

The datasets supporting the conclusions of this article are publicly available on the European Nucleotide Archive (<https://www.ebi.ac.uk/ena/browser/home>) and National Center for Biotechnology Information (<https://www.ncbi.nlm.nih.gov/sra/>), and the archive numbers of the accessions used are listed in the Additional file 1: Table S5. The scripts generated are available on GitHub https://github.com/Roberthorv/TE_in_Brachypodium/tree/main.

Funding

This project was funded by the Swiss National Science Foundation (project 31003A_182785) and the Research Priority Program Evolution in Action from the University of Zürich. Data analyzed in this paper were generated in collaboration with the Genetic Diversity Center (GDC), ETH Zürich.

Authors' contributions

A.R. and R.H. conceived the study. R.H. carried out the study, did the TE and SNP calling, performed the age-adjusted SFS analyses, conducted the X^tX analyses and ran the forward simulations. N.M. performed the GEA. Y.B. ran the iHS analyses. A.R. acquired the fundings. R.H. wrote the manuscript and A.R. revised it. All authors discussed the results and commented on the manuscript.

References

1. Bhattacharyya MK, Smith AM, Ellis TH, Hedley C, Martin C (1990) **The wrinkled-seed character of pea described by Mendel is caused by a transposon-like insertion in a gene encoding starch-branching enzyme** *Cell* **60**:115–22
2. Hof AE, Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, et al. (2016) **The industrial melanism mutation in British peppered moths is a transposable element** *Nature* **534**:102–105
3. Feschotte C (2008) **Transposable elements and the evolution of regulatory networks** *Nat. Rev. Genet* **9**:397–405
4. Qiu Y, Köhler C (2020) **Mobility connects: transposable elements wire new transcriptional networks by transferring transcription factor binding motifs** *Biochem. Soc. Trans* **48**:1005–1017
5. Slotkin RK, Martienssen R (2007) **Transposable elements and the epigenetic regulation of the genome** *Nat. Rev. Genet* **8**:272–85
6. Hollister JD, Gaut BS (2009) **Epigenetic silencing of transposable elements: a trade-off between reduced transposition and deleterious effects on neighboring gene expression** *Genome Res* **19**:1419–1428
7. Xiao H, Jiang N, Schaffner E, Stockinger EJ, van der Knaap E (2008) **A retrotransposon-mediated gene duplication underlies morphological variation of tomato fruit** *Science* **319**
8. Gordon SP, Contreras-Moreira B, Woods DP, Des Marais DL, Burgess D, Shu S, et al. (2017) **Extensive gene content variation in the *Brachypodium distachyon* pan-genome correlates with population structure** *Nat. Commun* **8**
9. Bennetzen JL, Kellogg EA (1997) **Do plants have a one-way ticket to genomic obesity?** *Plant Cell* **9**:1509–1514
10. Vitte C, Panaud O (2003) **Formation of solo-LTRs through unequal homologous recombination counterbalances amplifications of LTR retrotransposons in rice *Oryza sativa* L** *Mol. Biol. Evol* **20**:528–540
11. Piégu B, Guyot R, Picault N, Roulin A, Sanyal A, Kim H, et al. (2006) **Doubling genome size without polyploidization: dynamics of retrotransposition-driven genomic expansions in *Oryza australiensis*, a wild relative of rice** *Genome Res* **16**:1262–1269
12. Wendel JF, Jackson SA, Meyers BC, Wing RA (2016) **Evolution of plant genome architecture** *Genome Biol* **17**:1–14
13. Lisch D (2013) **How important are transposons for plant evolution?** *Nat. Rev. Genet* **14**:49–61
14. Negi P, Rai AN, Suprasanna P (2016) **Moving through the stressed genome: emerging regulatory roles for transposons in plant stress response** *Front. Plant Sci* **7**:1–20

15. Rey O, Danchin E, Mirouze M, Loot C, Blanchet S (2016) **Adaptation to Global Change: A Transposable Element-Epigenetics Perspective** *Trends Ecol. Evol* **31**:514–526
16. Dubin MJ, Mittelsten Scheid O, Becker C (2018) **Transposons: a blessing curse** *Curr. Opin. Plant Biol* **42**:23–29
17. Quadrana L, Etcheverry M, Gilly A, Caillieux E, Madoui MA, Guy J, et al. (2019) **Transposition favors the generation of large effect mutations that may facilitate rapid adaption** *Nat. Commun* **10**:1–10
18. González J, Karasov TL, Messer PW, Petrov DA (2010) **Genome-wide patterns of adaptation to temperate environments associated with transposable elements in Drosophila** *PLoS Genet* **6**
19. Studer A, Zhao Q, Ross-Ibarra J, Doebley J (2011) **Identification of a functional transposon insertion in the maize domestication gene tb1** *Nat. Genet* **43**:1160–1163
20. Barrón MG, Fiston-Lavier AS, Petrov DA, González J (2014) **Population genomics of transposable elements in Drosophila** *Annu. Rev. Genet* **48**:561–581
21. Rishishwar L, Wang L, Wang J, Yi SV, Lachance J, Jordan IK (2018) **Evidence for positive selection on recent human transposable element insertions** *Gene* **675**:69–79
22. Niu XM, Xu YG, Li ZW, Bian YT, Hou XH, Chen JF, et al. (2019) **Transposable elements drive rapid phenotypic variation in Capsella rubella** *Proc. Natl. Acad. Sci. USA* **116**:6908–6913
23. Jiang J, Xu YC, Zhang ZQ, Chen JF, Niu XM, Hou XH, et al. (2022) **Forces driving transposable element load variation during Arabidopsis range expansion** *bioRxiv* :1–12
24. Uzunović J, Josephs EB, Stinchcombe JR, Wright SI (2019) **Transposable elements are important contributors to standing variation in gene expression in Capsella grandiflora** *Mol. Biol. Evol* **36**:1734–1745
25. Castanera R, Morales-Díaz N, Gupta S, Purugganan M, Casacuberta JM (2023) **Transposons are a major contributor to gene expression variability under selection in rice populationse** *Life* **12**
26. Bourgeois Y, Ruggiero RP, Hariyani I, Boissinot S (2020) **Disentangling the determinants of transposable elements dynamics in vertebrate genomes using empirical evidences and simulations** *PLoS Genet* **16**
27. Stritt C, Gordon SP, Wicker T, Vogel JP, Roulin AC (2018) **Recent activity in expanding populations and purifying selection have shaped transposable element landscapes across natural accessions of the mediterranean grass Brachypodium distachyon** *Genome Biol. Evol* **10**:304–318
28. Charlesworth B (1991) **Transposable elements in natural populations with a mixture of selected and neutral insertion sites** *Genet. Res* **57**:127–134
29. Charlesworth B, Langley CH, Sniegowski PD (1997) **Transposable element distributions in Drosophila** *Genetics* **147**:1993–1995
30. Charlesworth B, Charlesworth D (1983) **The population dynamics of transposable elements** *Genet. Res* **42**:1–27

31. Charlesworth B (1996) **Background selection and patterns of genetic diversity in *Drosophila melanogaster*** *Genet. Res* **68**:131–149
32. Lockton S, Ross-Ibarra J, Gaut BS (2008) **Demography and weak selection drive patterns of transposable element diversity in natural populations of *Arabidopsis lyrata*** *Proc. Natl. Acad. Sci. U S A* **105**:13965–13970
33. Boissinot S, Davis J, Entezam A, Petrov D, Furano AV (2006) **Fitness cost of LINE-1 (L1) activity in humans** *Proc. Natl. Acad. Sci. U S A* **103**:9590–9594
34. Blumenstiel JP, Chen X, He M, Bergman CM (2014) **An age-of-allele test of neutrality for transposable element insertions** *Genetics* **196**:523–38
35. Rech GE, Bogaerts-Márquez M, Barrón MG, Merenciano M, Villanueva-Cañas JL, Horváth V, et al. (2019) **Stress response, behavior, and development are shaped by transposable element-induced mutations in *Drosophila*** *PLoS Genet* **15**
36. Mérel V, Gibert P, Buch I, Rodriguez Rada V, Estoup A, Gautier M, et al. (2021) **The Worldwide Invasion of *Drosophila suzukii* Is Accompanied by a Large Increase of Transposable Element Load and a Small Number of Putatively Adaptive Insertions** *Mol. Biol. Evol* **38**:4252–4267
37. International Brachypodium Initiative (2010) **Genome sequencing and analysis of the model grass *Brachypodium distachyon*** *Nature* **463**:763–768
38. Stritt C, Gimmi EL, Wyler M, Bakali AH, Skalska A, Hasterok R, et al. (2022) **Migration without interbreeding: Evolutionary history of a highly selfing Mediterranean grass inferred from whole genomes** *Mol. Ecol* **31**:70–85
39. Minadakis N, Williams H, Horvath R, Caković D, Stritt C, Thieme M, et al. (2023) **The demographic history of the wild crop relative *Brachypodium distachyon* is shaped by distinct past and present ecological niches** *Peer Community J* **3**:1–19
40. Bourgeois Y, Stritt C, Walser JC, Gordon SP, Vogel JP, Roulin AC (2018) **Genome-wide scans of selection highlight the impact of biotic and abiotic constraints in natural populations of the model grass *Brachypodium distachyon*** *Plant J* **96**:438–451
41. Wyler M, Stritt C, Walser JC, Baroux C, Roulin AC (2020) **Impact of transposable elements on methylation and gene expression across natural accessions of *Brachypodium distachyon*** *Genome Biol. Evol* **12**:1994–2001
42. Kimura M, Ohta T (1973) **The age of a neutral mutant persisting in a finite population** *Genetics* **75**:199–212
43. Voight BF, Kudaravalli S, Wen X, Pritchard JK (2006) **A map of recent positive selection in the human genome** *PLoS Biol* **4**
44. Günther T, Coop G (2013) **Robust identification of local adaptation from allele frequencies** *Genetics* **195**:205–20
45. Olazcuaga L, Loiseau A, Parrinello H, Paris M, Fraimout A, Guedot C, et al. (2020) **A whole-genome scan for association with invasion success in the fruit fly *Drosophila suzukii* using contrasts of allele frequencies corrected for population structure** *Mol. Biol. Evol* **37**:2369–2385

46. Garcia Guerreiro MP (2012) **What makes transposable elements move in the *Drosophila* genome** *Heredity* **108**:461–468
47. Belyayev A (2014) **Bursts of transposable elements as an evolutionary driving force** *J. Evol. Biol* **27**:2573–2584
48. Horvath R, Menon M, Stitzer M, Ross-Ibarra J (2022) **Controlling for variable transposition rate with an age-adjusted site frequency spectrum** *Genome Biol. Evol* **14**
49. Wright SI, Agrawal N, Bureau TE (2003) **Effects of recombination rate and gene density on transposable element distributions in *Arabidopsis thaliana*** *Genome Res* **13**:1897–1903
50. Horvath R, Slotte T (2017) **The Role of small RNA-based epigenetic silencing for purifying selection on transposable elements in *Capsella grandiflora*** *Genome Biol. and Evol* **9**:2911–2920
51. Raissig MT, Woods DP (2022) **The wild grass *Brachypodium distachyon* as a developmental model system** *Curr. Top. Dev. Biol* **147**:33–71
52. Hasterok R, Catalan P, Hazen SP, Roulin AC, Vogel JP, Wang K, et al. (2022) ***Brachypodium*: 20 years as a grass biology model system; the way forward?** *Trends Plant Sci* **27**:1002–1016
53. Casacuberta E, González J (2013) **The impact of transposable elements in environmental adaptation** *Mol. Ecol* **22**:1503–1517
54. Bourgeois Y, Boissinot S (2019) **On the population dynamics of junk: a review on the population genomics of transposable elements** *Genes* **10**:1–23
55. Stitzer MC, Khaipho-Burch MB, Hudson AI, Song B, Valdez-Franco JA, Ramstein G, et al. (2023) **Transposable element abundance subtly contributes to lower fitness in maize**
56. Quadrana L, Silveira AB, Mayhew GF, LeBlanc C, Martienssen RA, Jeddeloh JA, et al. (2016) **The *Arabidopsis thaliana* mobilome and its impact at the species level** *eLife* **5**
57. Baduel P, Leduque B, Ignace A, Gy I, Gil J, Loudet O, et al. (2021) **Genetic and environmental modulation of transposition shapes the evolutionary potential of *Arabidopsis thaliana*** *Genome Biol* **22**:1–26
58. Langmüller AM, Nolte V, Dolezal M, Schlötterer C (2023) **The genomic distribution of transposable elements is driven by spatially variable purifying selection** *Nucleic Acids Res* :1–11
59. Stuart T, Eichten SR, Cahn J, Karpievitch Y, Borevitz JO, Lister R (2016) **Population scale mapping of transposable element diversity reveals links to gene regulation and epigenomic variation** *eLife* **5**
60. Eyre-Walker A, Woolfit M, Phelps T (2006) **The distribution of fitness effects of new deleterious amino acid mutations in humans** *Genetics* **173**:891–900
61. Lockton S, Gaut BS (2010) **The evolution of transposable elements in natural populations of self-fertilizing *Arabidopsis thaliana* and its outcrossing relative *Arabidopsis lyrata*** *BMC Evol. Biol* **10**:1–11

62. González J, Macpherson JM, Messer PW, Petrov DA (2009) **Inferring the strength of selection in drosophila under complex demographic models** *Mol. Biol. Evol* **26**:513–526
63. Xue AT, Ruggiero RP, Hickerson MJ, Boissinot S (2018) **Differential effect of selection against LINE retrotransposons among vertebrates inferred from whole-genome data and demographic modeling** *Genome Biol. Evol* **10**:1265–1281
64. Wang X, Weigel D, Smith LM (2013) **Transposon variants and their effects on gene expression in Arabidopsis** *PLoS Genet* **9**
65. McCue AD, Nuthikattu S, Slotkin RK (2013) **Genome-wide identification of genes regulated in trans by transposable element small interfering RNAs** *RNA Biol* **10**:1379–1395
66. McCue AD, Nuthikattu S, Reeder SH, Slotkin RK (2012) **Gene expression and stress response mediated by the epigenetic regulation of a transposable element small RNA** *PLoS Genet* **8**
67. McCue AD, Slotkin RK (2012) **Transposable element small RNAs as regulators of gene expression** *Trends Genet* **28**:616–623
68. Cho J (2018) **Transposon-derived non-coding RNAs and their function in plants** *Front. Plant Sci* **9**:1–6
69. Wyler M, Keller B, Roulin AC (2022) **Potential impact of TE-derived sRNA on gene regulation in the grass Brachypodium distachyon**
70. Wicker T, Sabot F, Hua-Van A., Bennetzen JL, Capy P, Chalhoub B, et al. (2007) **A unified classification system for eukaryotic transposable elements** *Nat. Rev. Genet* **8**:973–982
71. Wei K, Aldaimalani R, Mai D, Zinshteyn D, Satyaki PRV, Blumenstiel J, et al. (2022) **Rethinking the “gypsy” retrotransposon: A roadmap for community-driven reconsideration of problematic gene names** *OSF Preprints* :1–13 <https://doi.org/10.31219/osf.io/fma57>
72. Skalska A, Stritt C, Wyler M, Williams HW, Vickers M, Han J, et al. (2020) **Genetic and methylome variation in turkish Brachypodium distachyon accessions differentiate two geographically distinct subpopulations** *Int. J. Mol. Sci* **21**
73. Gordon SP, Contreras-Moreira B, Levy JJ, Djamei A, Czedik-Eysenberg A, Tartaglio VS, et al. (2020) **Gradual polyploid genome evolution revealed by pan-genomic analysis of Brachypodium hybridum and its diploid progenitors** *Nat. Commun* **11**
74. Bolger AM, Lohse M, Usadel B (2014) **Trimmomatic: a flexible trimmer for Illumina sequence data** *Bioinformatics* **30**:2114–2120
75. Langmead B, Salzberg S (2012) **Fast gapped-read alignment with Bowtie2** *Nat. Methods* **9**:357–359
76. Faust GG, Hall IM (2012) **YAHA: fast and flexible long-read alignment with optimal breakpoint detection** *Bioinformatics* **28**:2417–2424
77. Stritt C, Wyler M, Gimmi EL, Pippel M, Roulin AC (2020) **Diversity, dynamics and effects of long terminal repeat retrotransposons in the model grass Brachypodium distachyon** *New Phytol* **227**:1736–1748

78. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. (2010) **The genome analysis toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data** *Genome Res* **20**:1297–1303
79. Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van der Auwera GA, et al. (2017) **Scaling accurate genetic variant discovery to tens of thousands of samples**
80. Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, et al. (2012) **A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3** *Fly* **6**:80–92
81. Albers PK, McVean G (2020) **Dating genomic variants and shared ancestry in population-scale sequencing data** *PLoS Biol* **18**
82. Huo N, Garvin DF, You FM, McMahon S, Luo MC, Gu YQ, et al. (2011) **Comparison of a high-density genetic linkage map to genome features in the model grass *Brachypodium distachyon*** *Theor. Appl. Genet* **123**:455–464
83. Dray S, Dufour AB (2007) **The ade4 Package: Implementing the duality diagram for ecologists** *J. Stat. Softw* **22**:1–20
84. R Core Team (2021) **R: A language and environment for statistical computing**
85. Massicotte P, South A (2023) **rnaturalearth: World map data from natural earth**
86. Sabeti PC, Reich DE, Higgins JM, Levine HZP, Richter DJ, Schaffner SF, et al. (2002) **Detecting recent positive selection in the human genome from haplotype structure** *Nature* **419**:832–837
87. Gautier M, Klassmann A (2017) **Vitalis R. rehh 2.0: a reimplementa-tion of the R package rehh to detect positive selection from haplotype structure** *Mol. Ecol. Resour* **17**:78–90
88. Gautier M (2012) **Vitalis R. rehh: An R package to detect footprints of selection in genome-wide SNP data from haplotype structure** *Bioinformatics* **28**:1176–1177
89. Fox J, Weisberg S (2019) **An R companion to applied regression**
90. Gautier M (2015) **Genome-wide scan for adaptive divergence and association with population-specific covariates** *Genetics* **201**:1555–1579
91. Zhou X, Stephens M (2012) **Genome-wide efficient mixed-model analysis for association studies** *Nat. Genet* **44**:821–824
92. Benjamini Y, Hochberg Y (1995) **Controlling the false discovery rate: a practical and powerful approach to multiple testing** *J. R. Stat. Soc* **57**:289–300
93. Haller BC, Messer PW (2019) **Evolutionary modeling in SLiM 3 for beginners** *Mol. Biol. Evol* **36**:1101–1109
94. Haller BC, Messer PW (2019) **SLiM 3: forward genetic simulations beyond the Wright-Fisher smodel** *Mol. Biol. Evol* **36**:632–637

Article and author information

Robert Horvath

Department of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland

For correspondence: robert.horvath@bluewin.ch

ORCID iD: [0000-0002-3221-8835](https://orcid.org/0000-0002-3221-8835)

Nikolaos Minadakis

Department of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland

ORCID iD: [0000-0003-0643-3299](https://orcid.org/0000-0003-0643-3299)

Yann Bourgeois

DIADÉ, University of Montpellier, CIRAD, IRD, Montpellier, France, University House, Winston Churchill Ave, Portsmouth PO1 2UP, United Kingdom

Anne C. Roulin

Department of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland

For correspondence: anne.roulin@botinst.uzh.ch

ORCID iD: [0000-0002-6668-3321](https://orcid.org/0000-0002-6668-3321)

Copyright

© 2023, Horvath et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Senior Editor

Detlef Weigel

Max Planck Institute for Biology Tübingen, Tübingen, Germany

Reviewer #1 (Public Review):

The study presented in this manuscript presents very convincing evidence that purifying selection is the main force shaping the landscape of TE polymorphisms in *B. distachyon*, with only a few putatively adaptive variants detected, even though most conclusions are based on the 10% of polymorphisms contributed by retrotransposons. That first conclusion is not novel, however, as it had already been clearly established in natural *A. thaliana* strains (Baduel et al. *Genome Biol* 2021) and in experimental *D. simulans* lines (Langmüller et al. *NAR* 2023). In contrast to the conclusions reached in *A. thaliana*, however, Horvath et al. report here a seemingly deleterious effect of TE insertions even very far away from genes (>5kb), a striking observation for a genome of relatively similar size. However, SNPs within these regions show similar allele frequency deviations, suggesting this effect may be due to

mapping quality artefacts in gene poor regions of the genome. An additional caveat of this study is the lack of orthogonal benchmarking of the TE polymorphisms calls by a pipeline known for a high rate of false positives (see detailed Private Recommendations #1). The authors note that their conclusions are still valid using only the highest covered samples (>20x), yet this coverage threshold is relatively low and higher coverage would mostly reduce the rate of false negatives.

Nonetheless, this set of observations makes an important addition to the knowledge of TE dynamics in the wild and questions our understanding of the main molecular mechanisms through which TEs can impact fitness.

<https://doi.org/10.7554/eLife.93284.2.sa1>

Reviewer #2 (Public Review):

Transposable elements are known to have a strong potential to generate diversity and impact gene regulation, and they are thought to play an important role in plant adaptation to changing environments. Nevertheless, very few studies have performed genome-wide analyses to understand the global effect of selection on TEs in natural populations. Horvath et al., used available whole-genome re-sequencing data from a representative panel of *B. distachyon* accessions to detect TE insertion polymorphisms (TIPs) and estimate their time of origin. Using a thorough combination of population genomics approaches, the authors demonstrate that only a small amount of the TE polymorphisms are targeted by positive selection or potentially involved in adaptation. By comparing the age-adjusted population frequencies of TE polymorphisms and neutral SNPs, the authors found that retrotransposons are affected by purifying selection independently of their distance to genes. Finally, using forward simulations they were able to quantify the strength of selection acting on TE polymorphisms, finding that retrotransposons are mainly under moderate purifying selection, with only a minority of the insertions evolving neutrally.

Horvath et al., use a convincing set of strategies and their conclusions are well supported by the data. I think that incorporating polymorphism's age to the analysis of purifying selection is an interesting way to reduce the possible bias introduced by the fact that SNPs and TEs polymorphisms do not occur at the same pace. The fact that TE polymorphisms far from genes are also under purifying selection is an interesting result that reinforces the idea that trans-regulatory effect of TE insertions might not be a rare phenomenon, a matter that may be demonstrated in future studies.

<https://doi.org/10.7554/eLife.93284.2.sa0>

Author Response

The following is the authors' response to the original reviews.

eLife assessment

*This valuable study seeks to disentangle the different selective forces shaping the evolutionary dynamics of transposable elements (TEs) in the wild grass *Brachypodium distachyon*. Using haplotype-length metrics, and genetic and environmental differentiation tests, the authors present in large parts convincing evidence that positive selection on TE polymorphisms is rare, and that the distribution of TE ages points to purifying selection being the main force acting on TE evolution in this species. A caveat of this study, as of other studies that seek to assess TE insertion polymorphisms with short reads, is that the rates of false negatives and false positives are difficult to estimate,*

which may have major effects on the interpretation. This study will be relevant for anyone interested in the role of TEs in evolution and adaptation.

Thank you for considering our manuscript for publication in eLife. We appreciate the constructive comments and suggestions of the reviewers. We have addressed the raised issues by the reviewers. Below, we provide a more detailed response to each of the reviewer comments.

Public Reviews:

Reviewer #1:

*The study presented in this manuscript presents very convincing evidence that purifying selection is the main force shaping the landscape of TE polymorphisms in *B. distachyon*, with only a few putatively adaptive variants detected, even though most conclusions are based on the 10% of polymorphisms contributed by retrotransposons. That first conclusion is not novel, however, as it had already been clearly established in natural *A. thaliana* strains (Baduel et al. *Genome Biol* 2021) and in experimental *D. simulans* lines (Langmüller et al. *NAR* 2023), two studies that the authors do not mention, or improperly mention. In contrast to the conclusions reached in *A. thaliana*, however, Horvath et al. report here a seemingly deleterious effect of TE insertions even very far away from genes (>5kb), a striking observation for a genome of relatively similar size. If confirmed, as a caveat of this study is the lack of benchmarking of the TE polymorphisms calls by a pipeline known for a high rate of false positives (see detailed Private Recommendations #1), this set of observations would make an important addition to the knowledge of TE dynamics in the wild and questioning our understanding of the main molecular mechanisms through which TEs can impact fitness.*

Thank you for your positive evaluation of our paper. We have now adjusted the manuscript to include the mentioned studies (Line 330-333) and to address the issue of false positive and false negative calls. The detailed responses to all the raised points are below.

Reviewer #2:

Summary:

*Transposable elements are known to have a strong potential to generate diversity and impact gene regulation, and they are thought to play an important role in plant adaptation to changing environments. Nevertheless, very few studies have performed genome-wide analyses to understand the global effect of selection on TEs in natural populations. Horvath et al. used available whole-genome re-sequencing data from a representative panel of *B. distachyon* accessions to detect TE insertion polymorphisms (TIPs) and estimate their time of origin. Using a thorough combination of population genomics approaches, the authors demonstrate that only a small amount of the TE polymorphisms are targeted by positive selection or potentially involved in adaptation. By comparing the age-adjusted population frequencies of TE polymorphisms and neutral SNPs, the authors found that retrotransposons are affected by purifying selection independently of their distance to genes. Finally, using forward simulations they were able to quantify the strength of selection acting on TE polymorphisms, finding that retrotransposons are mainly under moderate purifying selection, with only a minority of the insertions evolving neutrally.*

Strengths:

Horvath et al., use a convincing set of strategies, and their conclusions are well supported by the data. I think that incorporating polymorphism's age into the analysis of

purifying selection is an interesting way to reduce the possible bias introduced by the fact that SNPs and TEs polymorphisms do not occur at the same pace. The fact that TE polymorphisms far from genes are also under purifying selection is an interesting result that reinforces the idea that the trans-regulatory effect of TE insertions might not be a rare phenomenon, a matter that may be demonstrated in future studies.

Weaknesses:

TEs from different classes and orders strongly differ in multiple features such as size, the potential impact of close genes upon insertion, insertion/elimination ratio (ie, MITE/TIR excision, solo-LTR formation), or insertion preference. Given such diversity, it is expected that their survival rates on the genome and the strength of selection acting on them could be different. The authors differentiate DNA transposons and retrotransposons in some of the analyses, the specificities of the most abundant plant TE types (ie, LTR/Gypsy, LTR/Copia, MITE DNA transposons) are not considered.

The authors used a short-read-based approach to detect TIPs and TAPs. It is known that detecting TE polymorphisms is challenging and can lead to false negatives, depending on the method used and the sequencing coverage. The methodology used here (TEPID) has been previously applied to other species, but it is unclear if the sensitivity of the TIP/TAP caller is equivalent to that of the SNP caller and how these potential differences may affect the results.

Thank you for your positive evaluation of our paper. We have now adjusted the manuscript and the discussion to include the mentioned points on the different TE superfamilies and the reliability of the TE calls. The detailed responses to all the raised points are below.

Private Recommendations:

Reviewer #1:

(1) TE polymorphisms (presence and absence variants) were called from short-read sequencing data using a pipeline (TEPID, Stuart et al. eLife 2016) that is known to have a low specificity as well as a low sensitivity in its detection of presence variants (Baduel et al. MIMB 2021). An assessment of the rate of false positives and false negatives in the data presented in this study and how it varies across TE superfamilies is therefore of crucial importance as it may bias all downstream analyses, especially if it impacts the identification of polymorphisms contributed by retrotransposons, as these are the basis of most conclusions of the manuscript. Nonetheless, the fact that the PCA of the polymorphisms contributed by DNA transposons is less able to distinguish genetic clades than with those contributed by retrotransposons, suggests the issue of false positives is most preeminent for DNA transposons. However, high rates of false positives may explain why no significant increase in TE frequency is detected within selective sweep regions, a result that runs against the expectation of hitch-hiking of neutral or weakly deleterious polymorphisms which the authors claim is the category of many TE polymorphisms. Furthermore, given that the reference genome belongs to the B_{east} clade, and the TEPID is better at calling absence than presence it may bias analyses in this clade (where clade-specific insertions will take the form of absence in other clades which are well detected) compared to other clades (where clade-specific insertions will be presence polymorphisms and may be missed). A benchmark of TE polymorphism calls could be done by de novo assembling one genome from each clade or by cross-checking at least the presence variant calls from TEPID with those made with another of the many TE calling pipelines available.

We agree with this issue raised by both reviewers regarding the effects of false negative and false positive TE calls. We also think that some reasonable follow-ups should be done to check

the potential impact of the false negative and false positive TE calls on the presented results, without turning the manuscript in a method comparison paper as this is not the main goal of this study. Therefore, we generated a subsample of our dataset that included only accession with an average genome wide mapping coverages of at least 20x, as the false negative TE call rate is correlated with the mapping coverage and a high mapping coverage is expected to lead to a reduction in the false negative TE call rates. We then used this subsample to check if our results would change if our dataset had a lower false negative TE call rate. However, reducing the rate of false negative calls through the use of only higher coverage samples did not change our results and interpretations.

Re-running the ANCOVA analyses revealed similar results regarding the accumulation of TEs in selective sweep regions. This was added to the main text Line 143-148: “Similar results were obtained when investigating the number of fixed TE polymorphisms (Additional file 2: Table S1) and the allele frequency of TE polymorphisms (Additional file 2: Table S2) in high iHS regions using a subset of our dataset with an expected lower false negative TE call rate, that only included samples with a genome-wide mapping coverage of at least 20x (see Discussion and Materials and Methods for more details).” and in Additional file 2: Table S1 and S2.

Further, we re-ran the age-adjusted SFS based on this subset of our dataset and found that the results and conclusions from the age-adjusted SFS were not only driven by false negative TE calls. This was also included in the text Line 338-349: “One caveat of the approach used in this study is that TE calling pipelines based on short-reads tend to have higher false positive and false negative call rates than SNP calling pipelines, which is also the case for the TEPID TE calling pipeline used here [57, 59]. A high false negative TE calling rate however might bias our TE frequency estimates toward lower frequencies, which could drive the observed patterns in the age-adjusted SFS. To assess if the false negative TE calling rate in our study substantially affected our results, we re-run the age-adjusted SFS on a subset of our dataset only including samples with a genome-wide mapping coverage of at least 20x, as higher mapping coverages are expected to reduce the false negative call rate [27, 59]. Using the TE allele frequencies estimated based on this subset of our data to estimate π frequency revealed similar results of the age-adjusted SFS based on the whole dataset (Additional file 1: Fig. S9), indicating that our observation of retrotransposons evolving under purifying selection is not solely driven by a high false negative TE calling rate.” and in Additional file 1: Fig. S9.

The details of this analyses have been added to the materials and methods Line 493-498: “Mapping coverage is known to influence false discovery rate [27, 59]. To investigate the impact of false positive and false negative TE calls on our results, we down sampled the TE dataset to only include TEs that have been called in samples that had at least an average mapping coverage of 20x. The allele frequencies of TEs present in our high coverage dataset was recalculated only considering samples with at least an average mapping coverage of 20x. This second TE dataset was then used to check if using a dataset with a higher mapping coverage and presumably a lower false TE calling rate impacted our results.”

(2) If confirmed, the observation that retrotransposons located more than 5kb away from genes appear to be also affected by purifying selection (L209) is indeed surprising. The authors should add a comparison with SNPs at the same distance from genes to strengthen the claim and make sure it is not the result of mapping artifacts, such as alignment quality dropping far away from genes.

We added a comparison of the age-adjusted SFS of SNPs and retrotransposons more than 5 kb away from genes to evaluate if the observed shape of the age-adjusted SFS of retrotransposons more than 5 kb away from genes were due to artefacts. The results are included on line 383-389: “Finally, we tested whether TE polymorphisms located more than 5 kb away from genes are evolving under purifying selection could be due to mapping or other

artefacts by comparing the shape of the age-adjusted SFS of retrotransposons and SNPs more than 5 kb away from genes. However, the age-adjusted SFS of SNPs 5 kb away from genes differs from the one of retrotransposons (Additional file 1: Fig. S10), indicating that the shape of the age-adjusted SFS of retrotransposons more than 5 kb away from genes is not likely to be the result of artefacts in regions of the genome far away from genes.” and Additional file 1: Fig. S10.

(3) The authors' claim that most TE polymorphisms are under weak to moderate purifying selection (L273) relies on the comparison of the age of polymorphisms in the oldest age bin with forward simulations. However, the conclusions from these comparisons cannot be extrapolated to the fitness effects of all TE polymorphisms as variants in the oldest age bin are de facto a biased sample of the variants of a category, a point the authors highlight.

We adjusted the mentioned paragraph to better highlight this point. Line 390-397: “To further ascertain the strength of purifying selection, we used forward simulation and showed that simulations assuming a moderately weak selection pressure ($S = -5$ or $S = -8$) against TE polymorphisms best fitted our observed data. In theory, no TE polymorphisms under strong purifying selection should be present in a natural population, as such mutations are expected to be quickly lost, especially in a predominantly selfing species where most loci are expected to be homozygous. Therefore, it is not surprising that TE polymorphisms which persist in *B. distachyon* are under weak to moderate selection, as also shown, for example, for the L1 retrotransposons in humans [27] or the BS retrotransposon family in *Drosophila melanogaster* [62].”

L220-228 for high-effect SNPs. Indeed, the most deleterious TE polymorphisms would be purged very quickly and never contribute to variants in the oldest age bin. Unless new arguments can be made to support this claim, this conclusion should be rephrased to claim instead that even the oldest TE polymorphisms are still mostly non-neutral and under weak to moderate purifying.

This has been adjusted. Line 231-232: “. Hence, even the oldest retrotransposon polymorphisms seem to be mostly non-neutral and are affected by purifying selection.”

L214: replace smaller with more negative for clarity.

Done.

L233: Given the discussion L220-228, the oldest age bin seems to be biased in its composition and thus not useful for comparisons. The sentence should therefore be rephrased to reflect that DNA transposon polymorphisms appear to be actually less deleterious than high-effect SNPs in S9A and B based on the penultimate age bin.

This has been fixed.

Reviewer #2:

- I wonder if false negative detection could artificially increase the evidence for purifying selection by increasing the amount of low-frequency variants. This could be easily checked if long-read data or genome assembly is available for any of the samples in the collection, by comparing the TIP/TAP prediction with the actual sequence.*

We agree with this point from the reviewers that false negative calls can lead to misinterpretations of the observed low-frequencies of the TEs. (But see response to the first comment of reviewer #1). Unfortunately, long-read data from the sample used here are not available to estimate false negative call rates. However, to check if the observed results are mainly driven by high false negative rates, we re-run the age-adjusted SFS based on samples with at least 20x mapping coverage, which should result in the reduction the false negative TE calling rate. The results and conclusions from this second analyses were included in the text Line 338-349: “One caveat of the approach used in this study is that TE calling pipelines based on short-reads tend to have higher false positive and false negative call rates than SNP calling pipelines, which is also the case for the TEPID TE calling pipeline used here [57, 59]. A high false negative TE calling rate however might bias our TE frequency estimates toward lower frequencies, which could drive the observed patterns in the age-adjusted SFS. To assess if the false negative TE calling rate in our study substantially affected our results, we re-run the age-adjusted SFS on a subset of our dataset only including samples with a genome-wide mapping coverage of at least 20x, as higher mapping coverages are expected to reduce the false negative call rate [27, 59]. Using the TE allele frequencies estimated based on this subset of our data to estimate π frequency revealed similar results of the age-adjusted SFS based on the whole dataset (Additional file 1: Fig. S9), indicating that our observation of retrotransposons evolving under purifying selection is not solely driven by a high false negative TE calling rate.” and in Additional file 1: Fig. S9.

- *Supplementary Figure S1. DNA transposons are much worse at separating the samples in comparison to LTR-retrotransposons. Doesn't this suggest that these two classes have very different dynamics in the population and maybe different intensities of the selection forces acting on them? Could this profile be explained as DNA transposons being older and likely more fixed in all the clades, whereas retrotransposons are more recent and more specific to some populations? Another possibility might be that some B. distachyon DNA transposons had an unusually high excision rate. In any case, in my opinion, this reinforces the need to study the different TE orders in more detail.*

Indeed, different TE orders and superfamilies can have different excision rates, age distributions and be under different selective regimes. To investigate the possibility that different TE orders are affected by very different selective regimes, we split our TE dataset into the four different TE types: Copia, Ty3, Helitron and MITE. We then re-run the age-adjusted SFS analyses and added our results to the text Line 422-430: “To further examine our conclusion on purifying selection, we investigated the selective regime affecting different retrotransposons and DNA-transposons superfamilies. Thereby, we generated age-adjusted SFS for the four most common TE superfamilies Copia, Ty3 (also known under the name Gypsy, but we will avoid using this name because of its problematic nature see [71]), Helitron and MITE and found similar deviations of the π frequency from 0 in the four investigated TE superfamilies (Additional file 1: Fig. S12–S15). These results indicate that our conclusion on the broad effect of purifying selection is not driven by a single TE superfamily but is at least common among the four most numerous TE superfamilies.” and in Additional file 1: Fig. S12–S15.

- *Line 112: “most TE polymorphisms in our dataset were young and only a few were very old”. Does this change substantially among TE orders/superfamilies?*

Indeed, there are some differences in the age distribution of the TEs depending on the superfamilies, However, the differences are no substantial as the age bins in the age-adjusted SFS of the different TE superfamilies are fairly similar. See Additional file 1: Fig. S12–S15.

- *Figure 2. Is difficult to read, especially lower panels. I think the grey border of the boxplots makes visualization difficult.*

The gray borders have been removed.