

Association with TFIIC limits MYCN localization in hubs of active promoters and chromatin accumulation of non-phosphorylated RNA Polymerase II

Reviewed Preprint

v2 • July 17, 2024

Revised by authors

Reviewed Preprint

v1 • February 7, 2024

Raphael Vidal, Eoin Leen, Steffi Herold, Mareike Müller, Daniel Fleischhauer, Christina Schüle-Völk, Dimitrios Papadopoulos, Isabelle Röschert, Leonie Uhl, Carsten P Ade, Peter Gallant, Richard Bayliss, Martin Eilers , Gabriele Büchel 

Theodor Boveri Institute, Department of Biochemistry and Molecular Biology, Biocenter, University of Würzburg; Am Hubland, 97074 Würzburg, Germany. • Comprehensive Cancer Center Mainfranken; Josef-Schneider-Str. 6, 97080 Würzburg, Germany. • Astbury Centre for Structural Molecular Biology, Faculty of Biological Sciences, University of Leeds; Leeds LS2 9JT, United Kingdom. • Mildred Scheel Early Career Center, University Hospital Würzburg; Josef-Schneider-Str. 6, 97080 Würzburg, Germany. • Theodor Boveri Institute, Core Unit High-Content Microscopy, Biocenter, University of Würzburg; Am Hubland, 97074 Würzburg, Germany.

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Summary

MYC family oncoproteins regulate the expression of a large number of genes and broadly stimulate elongation by RNA polymerase II. While the factors that control the chromatin association of MYC proteins are well understood, much less is known about how interacting proteins mediate MYC's effects on transcription. Here we show that TFIIC, an architectural protein complex that controls the three-dimensional chromatin organization at its target sites, binds directly to the amino-terminal transcriptional regulatory domain of MYCN. Surprisingly, TFIIC has no discernible role in MYCN-dependent gene expression and transcription elongation. Instead, MYCN and TFIIC preferentially bind to promoters with paused RNAPII and globally limit the accumulation of non-phosphorylated RNAPII at promoters. Consistent with its ubiquitous role in transcription, MYCN broadly participates in hubs of active promoters. Depletion of TFIIC further increases MYCN localization to these hubs. This increase correlates with a failure of the nuclear exosome and BRCA1, both of which are involved in nascent RNA degradation, to localize to active promoters. Our data suggest that MYCN and TFIIC exert a censoring function in early transcription that limits promoter accumulation of inactive RNAPII and facilitates promoter-proximal degradation of nascent RNA.

eLife assessment

This study presents the **valuable** finding that TFIIC interacts with MYCN to regulate RNA polymerase II dynamics by dissecting its impact on 3D chromatin architecture. Authors provide **convincing** evidence that MYCN and TFIIC show long-range chromatin contacts, and that the expression of each protein limits the function of the other. The notion emerges that TFIIC helps MYCN to maintain output at promoters while decreasing less productive associations at larger more extensively connected chromatin hubs. The paper is of interest to molecular biologists working on MYCN-dependent regulation of gene expression.

<https://doi.org/10.7554/eLife.94407.2.sa4>

Introduction

The MYC family of proto-oncogenes is at the epicenter of cellular regulatory networks that govern cell growth, proliferation, and differentiation (Dang, 2012, Kress, Sabo et al., 2015). The three MYC paralogs (MYC, MYCN, and MYCL) are central players in normal development and tissue homeostasis, and when dysregulated, fuel many of the processes that are hallmarks of cancer (Dhanasekaran, Deutzmann et al., 2022, Hanahan, 2022). Among them, MYCN has attracted attention for its causal role in the development of neuroblastoma and other childhood tumors (Rickman, Schulte et al., 2018).

Both MYC and MYCN bind to virtually all active promoters and profoundly alter the dynamics of RNA polymerase II (RNAPII) transcription, with an increase in pause release and elongation being most apparent (Herold, Kalb et al., 2019, Walz, Lorenzin et al., 2014). One consequence of these changes are alterations in expression of a broad range of target genes (Dhanasekaran, Hansen et al., 2023). Unrelated to those changes in gene expression, MYC and MYCN also control RNAPII function to limit the accumulation of R-loops, to facilitate promoter-proximal double-strand break repair and to co-ordinate transcription elongation with DNA replication (Papadopoulos, Uhl et al., 2023). Which of these effects are critical for the oncogenic functions of MYC is an open question and consequently the partner proteins via which MYC proteins alter RNAPII dynamics are under intense investigation (Baluapuri, Hofstetter et al., 2019, Baluapuri, Wolf et al., 2020, Buchel, Carstensen et al., 2017, Das, Lewis et al., 2022, Heidelberger, Voigt et al., 2018, Kalkat, Resetca et al., 2018, Lourenco, Resetca et al., 2021, Oksuz, Henninger et al., 2023). Direct interactions of MYC and MYCN with MAX, WDR5 and MIZ1 control the localization of MYC on chromatin (Blackwood & Eisenman, 1991, Blackwood, Lüscher et al., 1992, Thomas, Wang et al., 2015, Vo, Wolf et al., 2016, Walz et al., 2014). MYC-dependent effects on RNAPII elongation involves the transfer of elongation factors SPT5 and PAF1c from MYC onto RNAPII (Baluapuri et al., 2019, Endres, Solvie et al., 2021, Jaenicke, von Eyss et al., 2016). Both MYC and MYCN also interact with and activate topoisomerases I and II, suggesting that MYC/N-dependent pause release also involves the relieve of torsional stress that builds up during early transcription (Das, Kuzin et al., 2022).

Intriguingly, both MYC and MYCN form prominent complexes with TFIIC and mapping experiments for MYCN show that the amino terminal transcription-regulatory domain is required for the interaction (Buchel et al., 2017, Heidelberger et al., 2018). TFIIC was first identified as a general transcription factor for RNA polymerase III (RNAPIII) (Orioli, Pascali et al., 2012). Surprisingly, TFIIC also binds to thousands of genomic sites that are not shared with RNAPIII and are called extra TFIIC (ETC) sites (Noma, Cam et al., 2006). Many ETC sites localize to RNAPII-

transcribed promoters and such sites are often juxtaposed to MYCN binding sites (Buchel et al., 2017 [DOI](#), Mogtaderi, Wang et al., 2010 [DOI](#), Noma et al., 2006 [DOI](#), Oler, Alla et al., 2010 [DOI](#)). TFIIC is an architectural protein that can affect the three-dimensional chromatin organization at its binding sites (Noma et al., 2006 [DOI](#), Van Bortle & Corces, 2013 [DOI](#)). Functionally TFIIC can act as insulator that blocks the spread of chromatin states (Raab, Chiu et al., 2012 [DOI](#)), affect cohesin loading (Buchel et al., 2017 [DOI](#)), and bind at the borders of topologically associating domains (TADs) (Van Bortle, Nichols et al., 2014 [DOI](#)).

Intriguingly, TFIIC compartmentalizes RNAPII promoters and gene expression. Specifically, the effects of TFIIC on the expression of E2F-dependent and neuronal genes correlate with its effects on the three-dimensional chromatin architecture. In the case of neuronal genes, TFIIC prevents the localization of activity-dependent genes to sites of active transcription before stimulation (Crepaldi, Policarpi et al., 2013 [DOI](#), Ferrari, de Llobet Cudalon et al., 2020, Policarpi, Crepaldi et al., 2017 [DOI](#)). Similarly, TFIIC together with the activity-dependent neuroprotector homeobox protein (ADNP) controls the three-dimensional architecture of cell cycle genes (Ferrari et al., 2020 [DOI](#)). Our previous experiments had suggested that TFIIC may have a role in MYCN-dependent control of RNAPII function and we here show that MYCN and TFIIC together have an unexpected censoring function that excludes non-functional RNAPII from hubs of active promoters.

Results

TFIIC interacts directly with the amino-terminus of MYCN

TFIIC is present in MYCN immunoprecipitates and the amino terminal region of recombinant MYCN binds to TFIIC in pull-down experiments performed with cell lysates (Buchel et al., 2017 [DOI](#)). To test whether MYCN and TFIIC interact directly, we expressed the six subunits of TFIIC together with a FLAG-tagged MYCN construct comprising amino acids 2-137 using a recombinant baculovirus to infect insect cells. Performing pulldown experiments with cell lysates and immobilized FLAG-tagged MYCN, we recovered all six subunits of the TFIIC complex in the eluate along with the FLAG-tagged MYCN construct, demonstrating that TFIIC can bind to the amino terminal region of MYCN (**Figure 1A** [DOI](#)). TFIIC is composed of two subcomplexes, designated TauA (1A) and TauB (1B), each comprising three of the six subunits. We first attempted co-expression of FLAG-MYCN with individual subcomplexes, but the 1B complex was too unstable to be isolated, and so we focused on the stable 1A complex. To test whether MYCN binds directly to 1A, we purified MYCN and the 1A complex separately (**Figure S1A** [DOI](#), B) and then mixed the purified preparations (**Figure 1B-D** [DOI](#)). In gel filtration experiments, the isolated MYCN protein eluted with a molecular weight of around 50 kDa, whereas a fraction of MYCN eluted with a much larger molecular weight together with the 1A complex when both were mixed (**Figure 1B-D** [DOI](#)). Native-mass spectrometry (Tamara, den Boer et al., 2022 [DOI](#)) showed the molecular weight of this complex to be 204.7 kDa, very close to the sum of the molecular weights of a 1:1:1:1 complex (**Figure S1C** [DOI](#), D). We concluded that the MYCN amino terminal region and the 1A subcomplex of TFIIC form a stable complex with each other in solution.

As described in the introduction, TFIIC is both a general transcription factor for RNAPIII and has been implicated in regulation of RNAPII-dependent transcription. To test whether TFIIC is required for the proliferation of neuroblastoma cells, we individually depleted three of its subunits, TFIIC2, TFIIC3 and TFIIC5, by stable expression of doxycycline-inducible shRNAs. Controls confirmed that each shRNA efficiently depleted its target protein (**Figure S1E** [DOI](#)). To test both for a general requirement of each subunit in neuroblastoma cell proliferation and for MYCN-specific effects, we expressed each shRNA in SH-EP-MYCN-ER cell. SH-EP cells express endogenous MYC and are engineered to stably express a MYCN-ER chimera (Herold et al., 2019 [DOI](#)). Activation of the MYCN-ER by addition of 4-hydroxytamoxifen (4-OHT) suppresses the expression of endogenous MYC (**Figure S1F** [DOI](#)), in effect causing a switch from MYC to MYCN expression. Under control




Figure 1.

TFIIC directly interacts with MYCN.

A. Immunoblots showing levels of FLAG-tagged MYCN (amino acids 2-137) and the six subunits of the TFIIC complex after a pulldown assay using anti-FLAG affinity columns. Multiple columns labelled “Wash” represent the sequential washings ($n = 2$).

B. Size exclusion chromatography graph of MYCN (aa 1-137)/1A (black trace) or MYCN alone (red trace). The blue box marks the fractions used for panel C and D ($n = 2$).

C. Coomassie staining of fractions of the MYCN (aa 1-137)/1A complex (fractions marked with blue box in panel B).

D. Immunoblot of fractions of the MYCN (aa 1-137)/1A complex (fractions marked with blue box in panel B).

E. Growth curve (measured as % confluence) of SH-EP-MYCN-ER cells expressing Dox-inducible shRNA targeting *TFIIC2*, *TFIIC3* or *TFIIC5* under the indicated conditions. Data show mean $\pm$ standard deviation (SD) ($n = 3$).

conditions, depletion of TFIIC2 or TFIIC3 attenuated proliferation, whereas depletion of the small TFIIC5 subunit had little effect. Addition of 4-OHT had little effect by itself, but greatly enhanced the inhibitory effect of depletion of each subunit (**Figure 1E**), suggesting that TFIIC has both general and MYCN-specific roles in neuroblastoma cell proliferation.

TFIIC limits promoter binding of non-phosphorylated RNAPII in MYCN-expressing cells

The proximity of TFIIC binding sites to MYCN binding sites at many promoters transcribed by RNAPII prompted us to explore the role of TFIIC in MYCN-driven transcription (Buchel et al., 2017). Using an antibody that detects total RNAPII, we had previously shown that activation of MYCN in SH-EP-MYCN-ER cells impacts RNAPII in two ways: First, MYCN promotes pause release and MYCN's effect on elongation correlates with its effects on gene expression (Herold et al., 2019, Rahl, Lin et al., 2010, Walz et al., 2014). At the same time, activation of MYCN uniformly reduces RNAPII occupancy at all active promoters; this correlates with a reduced accumulation of promoter-proximal R-loops, suggesting that MYCN limits the accumulation of RNAPII at promoters that is impaired in productive elongation and/or splicing (Herold et al., 2019).

Previous experiments using antibodies directed against total RNAPII had yielded results that varied among different antibodies (Buchel et al., 2017) and RV, unpublished). To directly compare the effects of MYCN and TFIIC on promoter-bound RNAPII with those on elongating RNAPII in a side-by-side manner, we conducted parallel ChIP-seq analyses using antibodies that specifically recognize non-phosphorylated and Ser2-phosphorylated (pSer2) RNAPII, respectively. Visual inspection of multiple individual genes (**Figure 2A**) and global analyses (**Figure 2B,C** and **Figure S2A**,B) showed that activation of MYCN caused a global decrease in promoter association of non-phosphorylated RNAPII. A control ChIP-seq experiment established that addition of 4-OHT had no such effect in SH-EP cells that do not express a MYCN-ER chimera, demonstrating that the reduction is due to activation of MYCN (**Figure S2C**). At the same time, MYCN promoted an increase in transcription elongation as best documented by pSer2-RNAPII occupancy at the transcription end site (TES) (**Figures 2A**, C). This increase was most pronounced on MYCN-activated genes. Comparison with RNA-sequencing (RNA-seq) data showed that it correlated closely with MYCN-dependent changes in gene expression (**Figure 2D**). Depletion of TFIIC5 abrogated the MYCN-dependent decrease in chromatin association of non-phosphorylated RNAPII but had no effect on MYCN-dependent changes in transcription elongation (**Figure 2A-C**). In many experimental systems, MYC and MYCN effects on elongation parallel closely with the corresponding changes in gene expression. Consistent with these observations, RNA-seq showed that neither depletion of TFIIC3 nor of TFIIC5 had significant effects on either basal gene expression or MYCN-dependent changes in steady-state mRNA levels (**Figures 2E** and **Figure S2D**). Finally, we aimed to understand whether the localization of TFIIC and MYCN at promoters is linked to the dynamics of RNAPII at promoters and performed ChIP-seq of TFIIC5 in SH-EP-MYCN-ER cells in the presence of 5,6-Dichlorobenzimidazole-1- β -D-ribofuranoside (DRB), which stabilizes paused RNAPII and prevents pause-release of RNAPII (Mancebo, Lee et al., 1997). Inspection of individual genes (**Figure S2E**) and average density plots of all active promoters (**Figure 2F**) showed that activation of MYCN globally enhanced TFIIC5 association with regions surrounding active transcription start sites, consistent with previous observations (Buchel et al., 2017). Addition of DRB increased TFIIC5 association with active promoters and the combination of MYCN activation and DRB had a strong additive effect, arguing that MYCN preferentially recruits TFIIC5 to promoters with paused RNAPII.




Figure 2.

MYCN and TFIIC antagonize accumulation of non-phosphorylated RNAPII.

A. Browser tracks for non-phosphorylated RNAPII (top) and RNAPII pSer2 (bottom) ChIP-Rx at the indicated gene loci. SH-EP-MYCN-ER cells were treated with Dox (1 μ g/ml, 48 h) and/or 4-OHT, respectively. EtOH was used as control.

B. Average density plot of ChIP-Rx signal for non-phosphorylated RNAPII. Data show mean (line) $\pm$ standard error of the mean (SEM indicated by the shade) of different gene sets based on an RNA-seq of SH-EP-MYCN-ER cells $\pm$ 4-OHT. The y-axis shows the number of spike-in normalized reads and it is centered to the TSS $\pm$ 2 kb. N = number of genes in the gene set defined in the methods (n = 2).

C. Density plot of ChIP-Rx signal for RNAPII pSer2 as described for panel B. The signal is centered to the TES $\pm$ 2 kb (n = 2).

D. Average bin dot plot showing fold change for RNAPII pSer2 ChIP-Rx reads over TES $\pm$ 2 kb and RNA-seq of SH-EP-MYCN-ER for the same genes $\pm$ MYCN + TFIIC5 (blue) or + MYCN $\pm$ TFIIC5 (red). The plot shows 20 bins representing a total of 13,239 and 12,330 genes for $\pm$ MYCN + TFIIC5 and + MYCN $\pm$ TFIIC5 datasets, respectively (n = 3 for RNA-seq, n = 2 pSer2 RNAPII ChIP-Rx).

E. Average bin dot plot for RNA-seq of SH-EP-MYCN-ER showing log2 mRNA expression normalized by control per bin. Cells were treated with 1 μ g/ml Dox ("– TFIIC5", 48 h) and/or 4-OHT (" + MYCN", 4 h) or EtOH as control. Expression was normalized by its control. Each bin represents 150 genes of a total of 14,085 genes. Dotted line marks the relative expression at 0 (n = 3).

F. Density plot of ChIP-Rx signal for TFIIC5. Data show mean (line) $\pm$ SEM (shade) for 14,722 genes. The signal is centered to the TSS $\pm$ 2 kb (n = 2).

MYCN takes part in three-dimensional networks of active promoters

The effects of TFIIC on the expression of E2F-dependent and neuronal genes correlate with its effects on the three-dimensional chromatin architecture (see Introduction). This raised the possibility that complex formation of TFIIC similarly affects three-dimensional chromatin interactions of MYCN. To test this hypothesis, we used phosphorylated linker HiChIP (pLHiChIP), a modification of the HiChIP protocol, which identifies pairs of DNA loci that are brought into close spatial proximity to a specific protein (**Figure S3A** [↗](#)) (Mumbach, Rubin et al., 2016). We initially performed pLHiChIP for MYCN in SH-EP-MYCN-ER neuroblastoma cells (**Figure 3A** [↗](#)). ChIP experiments measuring the occupancy of multiple promoters bound by both MYCN and MYC showed that the chromatin association of MYCN is much greater than that of MYC, allowing analysis of MYCN function on chromatin with little interference from MYC (**Figure S3B** [↗](#)). Visual inspection of the MYCN pLHiChIP showed that thousands of pairs of MYCN binding sites are in close spatial proximity with each other (**Figure 3A** [↗](#)). Appropriate quality controls established the validity of these results: For example, the Hi-C module of pLHiChIP protocol yielded a high percentage of valid interactions pairs (**Figure S3C** [↗](#), D). Relative to the Hi-C input, the MYCN HiChIP is strongly enriched for interactions that connected two MYCN-binding sites with each other, confirming the specificity of the signal (**Figure S3E** [↗](#)).

The pLHiChIP data analysis revealed a total of 4,591 distinct binary interactions involving MYCN (**Figure 3B** [↗](#)). To facilitate a comprehensive functional analysis of MYCN anchor sites, we performed ChIP-seq for RNAPII from SH-EP-MYCN-ER cells and integrated the HiChIP data with these data (**Figure 3A** [↗](#)), along with other relevant annotations, such as SINE elements. Remarkably, our analysis revealed that 1,277 out of the 4,591 MYCN anchors (28%) were positioned on RNAPII promoters (**Figure 3B** [↗](#)). Furthermore, 864 out of the 4,591 MYCN interactions (19%) exhibited connections between promoters and either exonic or intronic sequences, consistent with previous observations showing that MYCN binding sites are often located within transcribed regions (Buchel et al., 2017 [↗](#)). Additionally, MYCN interactions were observed between promoters and SINE repetitive elements (647/4,591; 14%) including interactions with tRNAs (**Figure S3F** [↗](#)). These connections are likely shared with TFIIC (see below). We also observed interactions between promoters or gene bodies and enhancers, accounting for 453 out of the 4,591 interactions (10%).

We used the RNAPII and published MYCN (Herold et al., 2019 [↗](#)) ChIP-seq data as well as sequencing datasets of mRNA (Buchel et al., 2017 [↗](#)) and nascent (4sU-labeled) (Papadopoulos, Solvie et al., 2022 [↗](#)) RNA to identify the specific properties of promoters within three-dimensional MYCN interactions. This showed that significantly more MYCN and RNAPII bound to promoters that participated in such interactions and that these promoters were more active than MYCN-bound promoters without such interactions, although the latter difference was small (**Figure 3C** [↗](#)). We next used the binary interactions as a starting point to reconstruct interaction networks. This showed that MYCN participated in three-dimensional promoter hubs of different sizes (**Figure 3D** [↗](#)), with the largest connecting up to 34 promoters and 10 enhancers (**Figure 3E** [↗](#)). This number of promoters is consistent with published estimates (Palacio & Taatjes, 2022 [↗](#)). Functional annotation of the promoters that are contained in these large hubs showed that they are highly enriched in genes encoding ribosomal proteins and ribosome biogenesis genes transcribed by RNAPII (**Figure S3G** [↗](#)). We concluded from the data that MYCN participates in three-dimensional hubs of highly active promoters; we will use the term “promoter hubs” (Lim & Levine, 2021 [↗](#)) for these structures.




Figure 3.

MYCN is part of three-dimensional promoter hubs.

A. Top: Representative browser track of MYCN three-dimensional chromatin interactions. Height shows the number of paired end tags (PETs) indicating the interaction intensity and the width of the line shows the start and end positions of each anchor. Middle and bottom: Browser tracks showing the number of reads of MYCN and total RNAPII ChIP-Rx, respectively. Unless stated, all experiments were performed in SH-EP-MYCN-ER cells treated with 4-OHT (200 nM, 4 h). The ruler at the bottom shows the genomic coordinates. (n = 3 independent biological replicates for MYCN pLHiChIP; n = 2 for RNAPII ChIP-Rx).

B. Bar chart listing functional annotations of all binary MYCN interactions (N = 4,591; N indicates total number).

C. Boxplots showing relative binding of the indicated proteins (RNAPII, MYCN, TFIIIC5) to promoter regions or expression levels of the corresponding genes (mRNA by RNA-seq; 4sU by 4sU-seq). Red boxes: Genes bound by MYCN and part of MYCN-hubs; Blue boxes: Genes bound by MYCN that are not part of MYCN-hubs. Each pair was normalized to the median of the corresponding “blue” gene set. p-values were obtained by pairwise comparisons using Student’s *t*-test. (n = 2 for TFIIIC5 and RNAPII ChIP-Rx).

D. Boxplot showing the number of promoters in each cluster, with each red dot representing one cluster.

E. Network reconstruction of the three biggest clusters based on MYCN pLHiChIP interactions. Each anchor is represented by a node (“triangle”) and the lines show interactions between the anchors. The colors are indicating the different functional annotation.

Binding to TFIIC antagonizes MYCN localization in promoter hubs

To determine whether TFIIC participates in three-dimensional chromatin structures, we performed pLHiChIP using a previously validated antibody against TFIIC5 (Buchel et al., 2017). In these experiments, thousands of pairs of TFIIC binding sites yielded a robust signal, demonstrating that they are in close spatial proximity with each other (Figure 4A). Specifically, these analyses identified a total of 3,499 binary non-contiguous interactions for TFIIC. Comparison to RNAPII ChIP-seq data showed that 31% of these interactions connected a promoter or the gene body of an RNAPII-transcribed gene to a SINE repetitive element (Figure 4B). Importantly, 1,107 out of 4,591 MYCN interactions shared one (1,075) or both (32) anchors with TFIIC-containing interactions arguing that both proteins participate in common hubs (Figure 4C). This is consistent with our previous data, which showed that many promoters contain a TFIIC-bound site close to a MYCN-bound E-box element (Buchel et al., 2017). A significant percentage of the interactions that are shared between MYCN and TFIIC contained an E-box in one anchor and A- or B-boxes, which are bound by TFIIC, in the other anchor sequence, arguing that MYCN and TFIIC can participate in the same interactions (Figure S4A). Most interactions that are shared by MYCN and TFIIC connected promoters to other promoters, to gene bodies or to SINE elements. Conversely, TFIIC5 anchors without overlapping MYCN loops (“TFIIC5 only”) did not show an enrichment for promoter interactions (Figure 4D). We concluded that both MYCN and TFIIC5 are present in promoter hubs. To determine how TFIIC affects MYCN involvement in promoter hubs, we used SH-EP-MYCN-ER neuroblastoma cells that express a Dox-inducible shRNA targeting *TFIIC5* (Figure S1E) and performed spike-in phosphorylated linker HiChIP (spLHiChIP), a spike-in variation of pLHiChIP that allows quantitative comparisons between different samples. These experiments showed that depletion of TFIIC5 strongly increased the number of chromatin interactions of MYCN, arguing that association with TFIIC antagonizes MYCN participation in promoter hubs (Figure 4E, F and Figure S4B). HiChIP experiments performed for TFIIC5 showed that expression of MYCN moderately and blockade of pause release by DRB strongly decreased the frequency of three-dimensional interactions of TFIIC5 (Figures 4G, H). Stratification showed that interactions of TFIIC5 with promoters, gene bodies and SINE elements were all decreased upon incubation of cells with DRB (Figure S4C), arguing that MYCN/TFIIC5-bound genes are not part of three-dimensional promoter hubs.

TFIIC is required for promoter association of factors involved in nascent RNA degradation

Depletion of TFIIC led to the accumulation of non-phosphorylated RNAPII and an enhanced presence of MYCN in promoter hubs, suggesting that lack of TFIIC might lead to enhanced or uncontrolled functionality of RNAPII. To ascertain whether TFIIC is required for functionality of elongating RNAPII, we first performed an rMATS analysis, which analyzes changes in splicing (Shen, Park et al., 2014b; Wang, Pan et al., 2017). This showed that depletion of TFIIC3 caused significant increases in intron retention and exon skipping, evidence of aberrant splicing (Figure S5A, B). However, most of these alterations occurred in downstream exons, and were not observed for TFIIC5. Furthermore, expression of MYCN weakened this effect for TFIIC3 (Figure S5A, B), arguing that it does account for the requirement for TFIIC function in MYCN-expressing cells.

To explore promoter-proximal events, we made use of the fact that many of the factors that determine premature termination and degradation of nascent RNA have recently been elucidated and initially used PLA assays to survey a series of factors involved in premature termination (Rodriguez-Molina, West et al., 2023). Appropriate controls using immunofluorescence and PLA assays with single antibodies established the specificity of each assay (Figure S5C). Consistent with the ChIP-seq data, activation of MYCN enhanced the proximity between TFIIC5 and total RNAPII and depletion of TFIIC5 reduced the signal, confirming the specificity of the assay (Figure




Figure 4.

TFIIC antagonizes MYCN participation in promoter hubs.

A. Representative example of pLHiChIP track for MYCN (red) and TFIIC5 (blue) interactions (conventions as in [Figure 3A](#)) (n = 2).

B. Bar chart listing the total number of functional annotations for all TFIIC5 binary interactions (N = 3,499).

C. Venn diagram showing the number of interactions shared between MYCN and TFIIC5. The diagram at the left shows the types of overlaps between connections.

D. Bar chart listing the interactions functional annotations for MYCN anchors not overlapping with TFIIC5 anchors (“MYCN only”) as well as TFIIC5 anchors without overlapping MYCN anchors (“TFIIC5 only”) and their joint anchors.

E. Representative example of MYCN spLHiChIP track for MYCN interactions in the presence (blue) or absence (red) of TFIIC5 (n = 2).

F. Bar graph showing the fold change of all MYCN spLHiChIP interactions comparing “+ TFIIC5” and “- TFIIC5” in SH-EP-MYCN-ER cells expressing a Dox-inducible shRNA targeting *TFIIC5*. n1,2 indicates two independent biological replicates.

G. Representative example of TFIIC5 spLHiChIP track without (blue) or with (red) induction of MYCN for SH-EP-MYCN-ER cells (conventions as in [Figure 3A](#)).

H. Bar graph showing the number of TFIIC5 interactions normalized by the relative binding of TFIIC5 ChIP-Rx signals for the same coordinates. Coordinates defined as TSS ± 2 kb of 14,722 genes.

SSD [\(Figure 5A\)](#)). Activation of MYCN and depletion of TFIIC5 had only weak effects on the proximity of RNAPII with NELFE, which associates with pausing RNAPII. Similarly, depletion of TFIIC5 either alone or in combination with MYCN activation enhanced the proximity of RNAPII with PP2A, which is recruited to promoters via its interaction with the integrator termination complex ([Cossa, Parua et al., 2021](#) [\(Figure 5A\)](#), [Vervoort, Welsh et al., 2021](#) [\(Figure 5A\)](#)), and with PNUTS, a targeting subunit of PP1 that is globally involved in termination (**Figure 5A** [\(Figure 5A\)](#)) ([Cortazar, Sheridan et al., 2019](#) [\(Figure 5A\)](#), [Estell, Davidson et al., 2023](#), [Landsverk, Sandquist et al., 2020](#) [\(Figure 5A\)](#)).

Key factors in degradation of aberrant nascent RNAs include XRN2, a 5'-3' RNA exonuclease, and the exosome, a 3'-5' exonuclease RNA complex ([Cortazar, Erickson et al., 2022](#) [\(Figure 5B\)](#), [Gerlach, Garland et al., 2022](#) [\(Figure 5B\)](#), [Noe Gonzalez, Blears et al., 2021](#) [\(Figure 5B\)](#)). We have previously shown that MYCN can recruit the nuclear exosome to its target promoters ([Papadopoulos et al., 2022](#) [\(Figure 5B\)](#)). MYCN can also recruit BRCA1, which in turn stimulates binding of the mRNA decapping enzyme DCP1 that initiates RNA degradation ([Herold et al., 2019](#) [\(Figure 5B\)](#)). No significant effects were observed in the proximity between RNAPII with the XRN2 exonuclease (**Figure 5A** [\(Figure 5A\)](#)). In contrast, CUT&RUN experiments using antibodies directed EXOSC5, a core structural subunit of the exosome ([Kilchert, Wittmann et al., 2016](#) [\(Figure 5B\)](#)), showed that depletion of TFIIC5 strongly decreased the association of EXOSC5 with regions around the TSS, both in control and in MYCN-expressing cells (**Figure 5B** [\(Figure 5B\)](#)). To test whether MYCN-dependent recruitment of BRCA1 depends on TFIIC, we performed ChIP experiments at multiple MYCN-bound promoters (**Figure 5C** [\(Figure 5C\)](#) and **Figure S5E** [\(Figure S5E\)](#)). Consistent with our previous observations, only low amount of BRCA1 were found at core promoter in SH-EP cells and that induction of MYCN recruited BRCA1 to all tested promoters ([Herold et al., 2019](#) [\(Figure 5C\)](#)). Depletion of TFIIC5 had variable effects on the presence of BRCA1 by itself, but abrogated BRCA1 recruitment by MYCN at all promoters tested (**Figure 5C** [\(Figure 5C\)](#) and **Figure S5E** [\(Figure S5E\)](#)), demonstrating that TFIIC5 is required for MYCN-dependent recruitment of BRCA1 to multiple target sites. Collectively, the data show that TFIIC is required for association of the nuclear exosome and of BRCA1 with active promoters and that these effects are enhanced in cells expressing MYCN, arguing that they can account for the enhanced dependence of MYCN expressing cells on TFIIC. A model summarizing our data is shown in **Figure 6** [\(Figure 6\)](#).

Discussion

One of the most puzzling aspects of MYC biology is the apparent discrepancy between the global association of MYC and MYCN oncoproteins with open chromatin, which includes virtually all active promoters, particularly in tumor cells, and their much more restricted and often weak effects on the expression of downstream target genes. Most MYC/N binding sites on chromatin do not appear to be functional with respect to gene expression and transcriptional elongation ([Kress et al., 2015](#) [\(Figure 6\)](#), [Sabo, Kress et al., 2014](#) [\(Figure 6\)](#), [Walz et al., 2014](#) [\(Figure 6\)](#)). This raises the possibility that such binding sites are indeed non-functional ([Pellanda, Dalsass et al., 2021](#) [\(Figure 6\)](#)). Alternatively, MYCN has functions in transcription that go beyond gene regulation. In support of the later model, we and others have uncovered critical roles of MYC and MYCN in maintaining the genomic stability of tumor cells, for example, by coordinating transcription with DNA replication and by enabling promoter-proximal double-strand break repair ([Balupuri et al., 2020](#) [\(Figure 6\)](#), [Papadopoulos et al., 2023](#) [\(Figure 6\)](#)). This raises the question of which direct interaction partners of MYC and MYCN mediate these effects and what the underlying mechanisms are. Here we show that TFIIC binds directly to the amino-terminus of MYCN, and that a MYCN/TFIIC complex globally limits the accumulation of non-phosphorylated RNAPII at promoters.

Several mechanisms can account for these observations: First, the amino-terminus of MYC proteins is a transcription-activating domain that can interact with cyclin T1 and CDK8 ([Eberhardy & Farnham, 2002](#) [\(Figure 6\)](#)) and potentially other co-activators, hence TFIIC may block MYCN interaction with co-activators that recruit RNAPII to promoters. Since MYCN/TFIIC also restricts RNAPII accumulation at promoters that are not activated by MYCN, this is unlikely to be the sole mode of




Figure 5.

TFIIIC is required for promoter association of the exosome and of BRCA1.

A. Boxplots showing the number of proximity ligation assay (PLA) signals between RNAPII and NELFE, PP2A, PNUTS or XRN2. SH-EP-MYCN-ER cells were treated with 1 μ g/ml Dox ("– TFIIIC5", 48 h) and/or 4-OHT (" + MYCN"). EtOH was used as control. For clarity purposes, 500 cells pooled from different replicates were plotted. p-values were calculated comparing the PLA signal of all cells using unpaired Wilcoxon rank sum test. The grey dotted line indicates the median in the control condition (n = 3).

B. Density plot of CUT&RUN for EXOSC5 binding (N = 14,704 genes) in SH-EP-MYCN-ER cells expressing a Dox-inducible shRNA targeting *TFIIIC5* treated with 4-OHT. Data show mean $\pm$ SEM (shade).

C. BRCA1 ChIP in SH-EP-MYCN-ER cells expressing a Dox-inducible shRNA targeting *TFIIIC5* treated with 4-OHT (4 h). Shown is the mean of technical triplicates of one representative experiment with identical results (n = 2).




Figure 6.

Model.

Model summarizing our findings. We propose that complex formation with the TFIIC complex antagonize the localization of MYCN in promoter hubs and that this enables access of the nuclear exosome and BRCA1 to promoters with paused or stalled RNAPII. Both the exosome and BRCA1 have been implicated in fostering the degradation of nascent RNA at promoters. The precise mechanisms by which MYCN and TFIIC limit accumulation of non-phosphorylated RNAPII at promoters remain to be determined.

action. Second, MYCN and TFIIC binding sites are often downstream of the transcription start sites and the three-dimensional structures formed by MYCN and TFIIC may sterically restrict elongation and favour termination of non-phosphorylated RNAPII. Third, TFIIC is an architectural protein complex and re-localization of promoters may be central to the mechanism of MYCN/TFIIC action. Active promoters coalesce at a discrete number of sites in the eucaryotic nucleus (Lim & Levine, 2021 [↗](#), Palacio & Taatjes, 2022 [↗](#)). Many of the factors involved in basal transcription undergo liquid-liquid phase separation, arguing that transcription initiation takes place in condensate-like structures established by transcription-regulatory domains and by RNAPII itself (Hnisz, Shrinivas et al., 2017 [↗](#), Lewis, Das et al., 2023 [↗](#), Lu, Yu et al., 2018 [↗](#)). Consistent with their ubiquitous role in transcription, MYC proteins form condensates by themselves and enter condensates formed by the RNAPII-associated mediator complex (Boija, Klein et al., 2018 [↗](#), Solvie, Baluapuri et al., 2022, Yang, Chung et al., 2022 [↗](#)). As described in the introduction, TFIIC can restrict the localization of genes to sites of active transcription. We show here that MYCN/TFIIC complexes, in contrast to MYCN, are not part of hubs of active promoters, leading us to propose that TFIIC antagonizes MYCN participation in promoter hubs and limits the accumulation of paused or stalled RNAPII at promoters (**Figure 6** [↗](#)). Since TFIIC competes with the Aurora-A kinase for binding of MYCN and Aurora-A in turn promotes elongation by RNAPII, we propose that the competition between both complexes enables transcriptional elongation on promoters that are activated by MYCN (Buchel et al., 2017 [↗](#), Roeschert, Poon et al., 2021).

As described above, depletion of TFIIC has no discernible role in elongation and changes in polyA⁺ mRNA levels, raising the question of what the function of the association might be. The fate of nascent RNA is controlled by several interrelated processes leading to splicing and polyadenylation of the full-length transcript on the one hand, and mis-splicing, premature termination and RNA degradation on the other and MYCN has been implicated in both processes (Noe Gonzalez et al., 2021, Papadopoulos et al., 2022 [↗](#), Schmid & Jensen, 2019 [↗](#)). Depletion of TFIIC strongly decreased the association of the nuclear exosome, an RNA exonuclease complex that degrades multiple forms of aberrant nascent RNA (Schmid & Jensen, 2019 [↗](#)), and of BRCA1, which in turn recruits the mRNA decapping enzyme DCP1 at promoters (Herold et al., 2019 [↗](#)). The association of both the exosome and BRCA1 with promoters were enhanced in MYCN expressing cells, arguing that they are likely to reflect - at least in part - the activity of the TFIIC/MYCN complex. At the same time, TFIIC3 depletion caused splicing errors, which occurred predominantly in downstream exons and were independent of MYCN, arguing that the effect on splicing may reflect an activity of TFIIC that is independent of its function at promoters. We propose that the TFIIC/MYCN complex formation exerts a censoring and quality control function for RNAPII at promoters that have not received a full complement of activating signals and hypothesize that this contributes to focus the transcription machinery and metabolic resources on the genes that drive the growth of MYCN-driven tumors.

Acknowledgements

We thank Vanessa Luzak and Yordan Sbirkov for initial experiments on chromatin architecture. This work was funded by grants from German Cancer Aid (Mildred Scheel Early Career Center, #70113303; G.B.), German Cancer Aid (#70113870; M.E.), German Research Foundation (DFG) (INST 93/1023-1-FUGG and EI222/21-1; M.E.), Alex's Lemonade Stand Foundation (Crazy 8 Initiative; M.E. and G.B.), Wilhelm Sander-Stiftung (#2023.002.1; S.H.) and Medical Research Council (MR/V029975/1; E.L. and R.B.).

Author Contributions

R.V., E.L., S.H., M.M., D.F., I.R., D.P. and G.B. performed experiments; C.S.-V. analyzed PLA assays; R.V., L.U. and P.G. analyzed sequencing data; C.P.A. generated high-throughput data; G.B., R.B., and M.E. devised and supervised experiments and G.B. and M.E. wrote the paper.

Declaration of Interests

M.E. is a founder and shareholder of Tucana Biosciences.

Corresponding Authors

Correspondence should be addressed to Gabriele Büchel (gabriele.buechel@uni-wuerzburg.de) and Martin Eilers (martin.eilers@uni-wuerzburg.de).

Data and materials availability

The following published datasets were used: MYCN ChIP-seq and RNA-seq: GSE111905; 4sU-seq: GSE164569. Reviewers can access data from this work under GEO accession GSE223058.

Methods

Cell culture

Cell line derived from human neuroblastoma (SH-EP; CVCL_RR78) was verified by STR profiling and grown in RPMI-1640 (Thermo Fisher Scientific). Murine neuroblastoma cells (NHO2A) were grown in RPMI-1640. Murine NIH-3T3 (CVCL_0594) cells were grown in DMEM (Thermo Fisher Scientific). Media were supplemented with 10% fetal calf serum (Capricorn Scientific GmbH) and penicillin/streptomycin (Sigma-Aldrich). All cells were routinely tested for mycoplasma contamination. Inhibitors were used in the following concentrations: MLN8237: 1 μ M, 4 h; DRB: 100 μ M, 2 h. Growth curves were obtained using the Incucyte® Live-Cell Analysis System.

Transfection and lentiviral infection

For lentivirus production, HEK293TN (CVCL_UL49) cells were transfected using PEI (Polyethyleneimine, Sigma-Aldrich). Lentiviruses expressing a shRNA targeting *GTFIIC5* (targeting sequence #1: AAGCGCAGCACCTACAACACTACA, #2: TTGATAAATCTTGGCATCTGGG) were produced by transfection of pINDUCER11 (Sequence #1) and pLT3GEPIR (Sequence #2) plasmid together with the packaging plasmid psPAX.2 and the envelope plasmid pMD2.G into HEK293TN cells. Virus-containing supernatant was harvested 24 h and 48 h after transfection. SH-EP-MYCN-ER cells were infected with lentiviral supernatants in the presence of 4 μ g/ml polybrene (Sigma-Aldrich) for 24 h. Cells were sorted for GFP and RFP expression. shRNA expression was stimulated by addition of doxycycline (1 μ g/ml) for 12 h and cells were FACS-sorted for RFP-GFP double-positive cells. Lentiviruses expressing a shRNA targeting *GTFIIC2* or *GTFIIC3* (targeting sequence *GTFIIC2*: TGAAGCAGAAGAATGGTCTGGA, *GTFIIC3*: TTCATCATTTTCTTGTTTCAC) were produced by

transfection of and pLT3GEPiR plasmid together with the packaging plasmid psPAX.2 and the envelope plasmid pMD2.G into HEK293TN cells. After harvesting supernatant and infection SH-EP-MYCN-ER cells were selected using puromycin and sorted for GFP expression.

For experiments, cells were harvested 48 h after induction with doxycycline (1 µg/ml) or ethanol as control. For induction of the MYCN chimera cells were treated with 4-OHT (200 nM, 4 h) as indicated.

Constructs

All constructs were cloned from human cDNA, or subcloned from plasmids containing specific human cDNA sequences. All TFIIC subunit containing complexes (TFIIC/3xFLAG-MYCN; 1A/3xFLAG-MYCN; 1A) were expressed using the MultiBac system. For 1A/3xFLAG-MYCN, TFIIC3, TFIIC5, Strep-TFIIC6, and 3xFLAG-MYCN aa 2-137, were cloned into pACEBAC1, pIDC, pIDK, and pIDS respectively. These were sequentially assembled into a single vector using Cre-Lox recombination. For the TFIIC/3xFLAG MYCN complex, 10xHis-TFIIC1, TFIIC3, TFIIC5, TFIIC6, and 3xFLAG MYCN aa 2-137 were all cloned into pACEBAC1. A single pACEBAC1 vector containing all of these inserts was generated by repetitive sub-cloning of the subunits. The acceptor pACEBAC1 was opened using I-CeuI and SpeI. The region of interest was removed from the donor pACEBAC1 clone using I-CeuI and AvrII. The region of interest was then ligated into the acceptor pACEBAC1. An internal AvrII site had to be silently removed from TFIIC3 by site directed mutagenesis prior to sub-cloning. The pACEBAC1 with these five members was then used in successive Cre-Lox recombination reactions with pIDK TFIIC4 and then pIDC TFIIC2 in order to generate a construct with all seven members of the complex. For the 1A complex 6xhis-TEV-TFIIC3 was cloned into pACEBAC1. This was Cre-loxed with pIDK Strep-TFIIC6, and pIDC TFIIC5. MultiBac constructs were validated by restriction enzyme digest and sequencing. Post assembly, constructs containing multiple subunits were validated by analytical PCR and restriction enzyme digest.

Protein expression and purification

For *Sf9* expression, bacmid DNA was generated by transformation of expression vectors into *E. coli* DH10MultiBac cells. Purified bacmid DNA (1.6 – 8.3 µg) was transfected into a monolayer of *Sf9* cells, in a T25 flask, using X-tremeGENE™ HP DNA Transfection Reagent (Roche). After 7 days at 28 °C the P1 virus stock was collected. 1 ml of P1 virus was used to infect 300 ml of *Sf9* cells at a density of 2×10^6 cells/ml. The infected cells were grown for 72 h at 120 rpm and 27 °C prior to harvesting by centrifugation at 1,000 xg for 10 min. The supernatant was used to infect large scale cultures of *Sf9* cells. 100 ml of virus stock was used per litre of *Sf9* cells at a density of 1.5×10^6 cells/ml. These were harvested after 48 h. *E.coli* expression was performed using BL21(DE3)RIL cells. 10 ml from overnight cultures were used to seed each Litre of LB media (supplemented with 50 µg/ml kanamycin and 35 µg/ml chloramphenicol). The cells were grown at 37 °C and 220 rpm until mid log phase (~ 0.6 OD₆₀₀) whereupon expression was induced using 0.6 mM final isopropyl β-D-1-thiogalactopyranoside (IPTG). Cells were grown overnight at 20 °C, and were harvested by centrifugation at 6,000 xg for 15 minutes. Cell pellets were stored at -80 °C prior to purification.

For the purification of TFIIC/3xFLAG-MYCN and 1A/3xFLAG-MYCN complexes *Sf9* pellets were resuspended in a lysis buffer (25 mM Tris pH8, 200 mM NaCl, EDTA free cOmplete PI (Roche), 10 µg/ml DNaseI) then lysed by sonication. Lysate was clarified at 40,000 xg for 20 min. Clarified lysate was applied to 2 ml bed volume Anti-FLAG M2 Affinity Gel (Sigma-Aldrich). The slurry was rotated in the cold room for two hours prior to application to a gravity flow column. Flow was collected and resin washed as follows. For the TFIIC/3xFLAG-MYCN complex the resin was washed with 40 ml of wash buffer (25 mM Tris pH8, 200 mM NaCl) in four 10 ml batches. For the 1A/3xFLAG-MYCN complex the resin was washed with two 10 ml batches of wash buffer, followed by 10 ml of wash buffer with a final NaCl concentration of 350 mM, followed by 10 ml of wash buffer with a final NaCl concentration of 400 mM. The resin was finally washed with 10 ml of

wash buffer. In both cases the protein was eluted by addition of 17.5 ml wash buffer spiked with a final concentration of 0.25 mg/ml 3xFLAG peptide. For the 1A/3XFLAG-MYCN complex the eluate fractions were spiked with 2 mM final β -mercaptoethanol and then concentrated to less than 1 ml with a Vivaspin 6 10 kDa MWCO centrifugal concentrator. Size Exclusion Chromatography (SEC) was performed using a Superose 6 10/300 GL column attached to an ÄKTA pure FPLC system (Cytiva). The SEC buffer (25 mM Tris pH8, 200 mM NaCl, 3 mM DTT) was flowed over the column at 0.3 ml/min and fractions collected every 0.25 ml over the peaks. In order to do an MYCN alone SEC run, the MYCN peak fractions from several SEC runs of the complex were combined and re-concentrated down to 300 μ l. This was then run as previous. For all purified proteins, protein concentration was determined by absorbance at 280 nm. Unless otherwise stated proteins were aliquoted, flash frozen in liquid N₂ and stored at -80 °C prior to use.

For the purification of 1A alone, *Sf9* pellets were resuspended in a base buffer (25 mM Tris pH 7.4, 2.7 mM KCl, 287 mM NaCl, 2 mM β -mercaptoethanol) supplemented with 0.1 mg/ml DNase I, and 1 EDTA free cOmplete PI (Roche) tablet per 30 ml of buffer. Cells were lysed by sonication, then the lysate clarified at 40,000 xg for 25 min. Clarified lysate was applied to 2.5 ml bed volume His-select Cobalt Affinity Gel (Sigma-Aldrich). The slurry was rotated in the cold room for an hour prior to application to a gravity flow column. The resin was washed sequentially as follows: 50 ml base buffer, 10 ml ATP buffer (25 mM Tris pH 7.4, 137 mM NaCl, 2.7 mM KCl, 5 mM MgCl₂, 5 mM ATP), 50 ml base buffer with 5 mM imidazole, 50 ml base buffer with 10 mM imidazole. Protein was eluted sequentially by addition of 25 ml of base buffer with 150 mM imidazole followed by 10 ml of the same buffer with 500 mM imidazole. Eluted fractions were pooled and 0.7 mg of his-tagged TEV NIa was added. The mixture was dialysed overnight, using 10 kDa MWCO Snakeskin dialysis tubing, against 4 l of buffer (25 mM Tris pH 7.4, 137 mM NaCl, 2.7 mM KCl, 2 mM β -mercaptoethanol). Post-dialysis the TEV NIa processed complex was rebound to Cobalt resin as per previous. The flow fraction was collected and concentrated. SEC was performed as per the 1A/3XFLAG-MYCN complex. 1A was concentrated, as per pre-SEC, down to 16.8 μ M and used for 1A/MYCN complex reconstruction immediately.

Protein purification was assessed by a combination of Coomassie stained SDS-PAGE gels and immunoblotting (TFIIIC90: A301-239A, AB_890667; TFIIC5: A301-242A, AB_890669; TFIIC102: A301-238A, AB_890671, Bethyl Laboratories; TFIIC110: sc-81406, AB_2115237; MYCN: sc-53993, AB_831602, Santa Cruz Biotechnologies; TFIIC35: NBP2-31851, AB_2891101; TFIIC1: NBP2-14077, AB_2891102, Novus Biologicals). The SEC elution position of molecular weight standards were established by using Bio-Rad gel filtration standards (cat #1511901). 5 mg/ml BSA was additionally used as the 66 kDa marker. Dextran blue was used to establish the void position of the column.

Aurora-A aa 122-403 D274N C290A C393A was purified as previously described (Bayliss, Sardon et al., 2003 [\[4\]](#)). The final buffer was 20 mM Hepes pH 7.5, 150 mM NaCl, 5 mM MgCl₂, 10% glycerol, 1 mM β -mercaptoethanol. *E. coli* expressed MYCN aa 1-137 pellets were initially disrupted using base buffer (pH 7.5, 100 mM KH₂PO₄, 10 mM Tris, 300 mM NaCl, 2 mM β -mercaptoethanol) which was supplemented with 30 mg of lysozyme, 0.3 μ g DNaseI, and 0.9 μ M MnCl₂ per 30 ml of base buffer. This was sonicated 15 seconds on 15 seconds off for 150 seconds at 40 % amplitude. The lysate was initially clarified by centrifugation for 30 min at 30,000 xg. As this construct of MYCN expresses into inclusion bodies, the protein had to be recovered from the pellet using urea. All urea solutions were prepared immediately prior to use. The pellet was disrupted by vigorous pipetting after addition of 30 ml of 1 M urea in base buffer followed by sonication as previous. This solution was clarified by centrifugation for 20 min at 30,000 xg. This protocol of disruption of the pellet, following addition of urea containing base buffers, followed by clarification was repeated sequentially for solutions containing 2 M and 4 M urea. 6xHis-tagged MYCN aa 1-137 was found in the soluble fraction of the 4 M urea solution. This was diluted 1/4 with 25 mM Tris pH 7.4, 2.7 mM KCl, 137 mM NaCl, 2 mM β -mercaptoethanol prior to application of the protein to His-Select Cobalt affinity gel (Sigma-Aldrich). The slurry was rotated for 60 min prior to application to a gravity flow column. The resin was washed sequentially as follows with imidazole containing

buffer (25 mM Tris pH 7.4, 2.7 mM KCl, 137 mM NaCl, 2 mM β -mercaptoethanol): 50 ml 0 mM imidazole, 50 ml 5 mM imidazole, 25 ml 10 mM imidazole, 25 ml 150 mM imidazole, and 10 ml 500 mM imidazole. Pure fractions were pooled, along with 0.7 mg of his-tagged TEV N1a, and were dialysed overnight using 3.5 kDa MWCO snakeskin dialysis tubing against 4 l of buffer (25 mM Tris pH 7.4, 2.7 mM KCl, 137 mM NaCl, 2 mM β -mercaptoethanol). Post TEV N1a processing, the protein was re-applied to His-Select Cobalt affinity gel (Sigma-Aldrich). The flow through fractions were pooled and concentrated (Vivaspin Turbo 3,000 MWCO) prior to SEC. SEC was performed using an ÄKTA prime and 16/600 S75 column. Pure fractions were pooled and concentrated as per pre SEC. Protein concentration was determined using absorbance at 280 nm.

1A/3xFLAG-MYCN native mass spectrometry

The 1A/3xFLAG-MYCN complex was desalted into 200 mM ammonium acetate using 75 μ l Zeba spin desalting columns (ThermoFisher Scientific). Samples were analysed by nanoelectrospray ionisation MS using a quadrupole-orbitrap MS (Q-Exactive UHMR, ThermoFisher Scientific) using gold/palladium coated nanospray tips prepared in-house. The MS was operated in positive ion mode using a capillary voltage of 1.3 kV, capillary temperature of 250 °C and S-lens RF of 200 V. In source trapping was used with a desolvation voltage of -100 V for 4 μ s. Extended trapping was not used. The quadrupole mass range was 2000-15000 m/z. Nitrogen gas was used in the HCD cell with a trap gas pressure setting of 5. Orbitrap resolution was 12500, detector m/z optimisation was low. 5 microscans were averaged. Mass calibration was performed by a separate injection of sodium iodide at a concentration of 2 μ g/ μ l. Data processing was performed using QualBrowser 4.2.28.14 and deconvoluted using UniDec (Marty, Baldwin et al., 2015 [DOI](#)).

1A/3xFLAG-MYCN intact mass spectrometry

Protein desalting and mass analysis was performed by LC-MS using an M-class ACQUITY UPLC (Waters UK, Manchester, UK) interfaced to a Xevo QToF G2-XS mass spectrometer (Waters UK, Manchester, UK). Samples were diluted to 1 μ M using 0.1% TFA. 1 μ l of the 1 μ M sample was run on an Acquity UPLC Protein BEH C4 column (300 Å, 1.7 μ m, 2.1 mm $\times$ 100 mm, Waters UK) with an Acquity UPLC Protein BEH VanGuard Pre-Column (300 Å, 1.7 μ m, 2.1 mm $\times$ 5 mm, Waters UK). Buffer A was 0.1% formic acid in water, and buffer B was 0.1% formic acid in AcN (v/v basis). System flow rate was kept constant at 50 μ l/min. Protein sample was loaded on to the trap column in 20% acetonitrile/0.1% formic acid and washed for 5 min. Following valve switching, the bound protein was eluted by a gradient of 20-95% solvent B in A over 10 min. The column was subsequently washed with 95 % solvent B in A for 5 min before re-equilibration at 20% solvent B in A ready for the next injection. The mass spectrometer was calibrated using a separate injection of glu-fibrinopeptide. Data were processed using MassLynx 4.2.

Protein identification/Peptide LC-MS analysis

Gel bands were excised and chopped into small pieces ($\sim$ 1 mm³), covered with 30% ethanol in a 1.5 ml microcentrifuge tube and heated to 56 °C for 30 min with shaking. The supernatant was removed, replaced with fresh ethanol solution and was again heated to 56 °C for 30 min with shaking. This was repeated until all Coomassie stain was removed from the gel. The gel slices were then dehydrated by covering with 100% acetonitrile and left for five minutes before the supernatant was discarded and replaced with a fresh aliquot of acetonitrile. Proteins were reduced by adding 100 μ l 20 mM DTT solution before incubation at 57 °C for 1 h with shaking. The supernatant was removed and once the gel pieces were at room temperature proteins were alkylated by adding 100 μ l 55 mM iodoacetic acid. The samples were then incubated at room temperature in the dark for 30 min with shaking. After removing the supernatant, the gel slices were then covered with 100% acetonitrile and left for five minutes. The acetonitrile was removed, and the gel pieces were left to dry in a laminar flow hood for 60 min. Once dry, the gel slices were cooled on ice then they were covered with ice-cold protease solution and left on ice for 20 min to rehydrate. Trypsin solution (20 ng/ μ l in 25 mM ammonium bicarbonate) was added to the

unknown bands, chymotrypsin solution (25 ng/μl in 25 mM ammonium bicarbonate). Excess protease solution was removed, and the gel slices were covered with a minimal amount of 25 mM ammonium bicarbonate. After briefly vortexing and centrifuging, the gel slices were incubated at 37 °C while shaking for 18 h. The resulting digest was vortexed and centrifuged. The supernatant was recovered and added to a 1.5 ml tube containing 5 μl acetonitrile/ water/ formic acid (60/35/5; v/v). 50 μl acetonitrile/ water/ formic acid (60/35/5; v/v) was added to the gel slices and vortexed for an additional 10 min. The supernatant was pooled with the previous wash and one additional wash of the gel slices was performed. The pool of three washes was dried by vacuum centrifugation. The peptides were reconstituted in 20 μl 0.1% aqueous trifluoroacetic acid prior to analysis. 4 μl sample were injected onto an in house-packed 20 cm capillary column (inner diameter 75 μm, 3.5 μm Kromasil C18 media). An EasyLC nano liquid chromatography system was used to apply a gradient of 4–40% ACN in 0.1% formic acid over 45 min at a flow rate of 250 nl/min. Total acquisition time was 60 min including column wash and re-equilibration. Separated peptides were eluted directly from the column and sprayed into an Orbitrap Velos Mass Spectrometer (ThermoFisher Scientific, Hemel Hempstead, UK) using an electrospray capillary voltage of 2.7 kV. Precursor ion scans were acquired in the Orbitrap with resolution of 60,000. Up to 20 ions per precursor scan were selected for fragmentation in the ion-trap. Dynamic exclusion of 30 s was used. Peptide MS/MS data were processed with PEAKS Studio X+ (Bioinformatic Solutions Inc, Waterloo, Ontario, Canada) and searched against the Uniprot databases for *H. sapiens* and *S. frugiperda* proteins (release 2021_02). Carbamidomethylation was selected as a fixed modification, variable modifications were set for oxidation of methionine and deamidation of glutamine and asparagine. MS mass tolerance was 15 ppm, and fragment ion mass tolerance was 0.3 Da. The peptide false discovery rate was set to 1%.

Immunoblots

Whole cell extracts were prepared using RIPA buffer (50 mM HEPES pH 7.9, 140 mM NaCl, 1 mM EDTA; 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS) containing protease and phosphatase inhibitor cocktails (Sigma-Aldrich). Lysates were cleared by centrifugation and protein concentrations were determined by Bradford.

Protein samples were separated on Bis-Tris gels and transferred to a PVDF membrane (Millipore). For immunoblots showing multiple proteins with similar molecular weight, one representative loading control is shown. Vinculin (VCL) or GAPDH were used as loading control. Antibodies used in this study: MYCN (B8.4.B): sc-53993, AB_831602, Santa-Cruz Biotechnology; TFIIC5: A301-242A, AB_890669, Bethyl Laboratories; MYC (Y69): ab32072, AB_731658, Abcam; VCL (h-VIN1): V9131, AB_477629, Sigma-Aldrich; GAPDH: 2118, AB_561053, Cell Signaling; TFIIC2: sc-81406, AB_2115237, Santa-Cruz; TFIIC3: sc-393235, Santa-Cruz.

Proximity Ligation Assay (PLA)

SH-EP-MYCN-ER cells were plated in 384-well plates (PerkinElmer), treated with Dox and/or 4-OHT and fixed with methanol for 20 min. After blocking for 30 min with 5% BSA in PBS, cells were incubated over night at 4 °C with primary antibodies: Total RNA Polymerase II (F12): sc-55492, (N20): sc-899, Santa-Cruz Biotechnology; TFIIC5: A301-242A, Bethyl Laboratories; NELFE: ABE48, Merck; PP2A: 2038, Cell Signaling; PNUTS: A300-439-1, Bethyl Laboratories; XRN2: A301-103A, Bethyl Laboratories. PLA was performed using Duolink® In Situ Kit (Sigma-Aldrich) according to the manufacturer's protocol. Nuclei were counterstained using Hoechst 33342 (Sigma-Aldrich). Pictures were taken with Z-stacks of four planes at 0.5 μm distance at 40-fold magnification in Operetta® CLS High-Content Imaging System. Analysis was performed in Harmony® High Content Imaging and Analysis Software. Thirty image fields per well were acquired with a total of at least 500 cells per sample.

High-throughput sequencing

ChIP and ChIP-seq was performed as described previously (Roeschert et al., 2021 [DOI](#)). For each ChIP or ChIP-Rx seq experiment, 5×10^7 cells per immunoprecipitation condition were fixed for 5 min at room temperature with formaldehyde (final concentration, 1%). Fixation was stopped by adding 125 mM glycine for 5 min. Cells were harvested in ice-cold PBS containing protease and phosphatase inhibitors (Sigma-Aldrich). As exogenous control (spike-in), murine NHO2A or NIH3T3 cells were added at a 1:10 cell ratio during cell lysis. Cell lysis was carried out for 20 min in lysis buffer I (5 mM PIPES pH 8.0, 85 mM KCl, 0.5% NP-40) and nuclei were collected by centrifugation (1,500 rpm, 10 min, 4 °C). Crosslinked chromatin was prepared in lysis buffer II (10 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 1% sodium deoxycholate, 0.1% SDS) and fragmented by using the Covaris Focused Ultrasonicator M220 for 50 min per ml lysate. For each IP reaction, 15 μ l (for ChIP) or 100 μ l (for ChIP-seq) Dynabeads Protein A and Protein G (Thermo Fisher Scientific) were pre-incubated overnight with rotation in the presence of 5 mg/ml BSA and 3 μ g (for ChIP) or 10–15 μ g (for ChIP-seq) antibody (MYCN (B8.4.B): sc-53993, AB_831602; TFIIIC5: A301-242A, AB_890669; RNAPII pSer2: ab5095, AB_304749; RNAPII (8WG16): sc-56767, AB_785522; BRCA1: A300-000A, AB_67367, Bethyl Laboratories). Chromatin was added to the beads, and IP was performed for a minimum of 6 h at 4 °C with rotation. Beads were washed three times each with washing buffer I (20 mM Tris pH 8.1, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS), washing buffer II (20 mM Tris pH 8.1, 500 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS), washing buffer III (10 mM Tris pH 8.1, 250 mM LiCl, 1 mM EDTA, 1% NP-40, 1% sodium deoxycholate; including a 5 min incubation with rotation) and TE buffer. Chromatin was eluted twice by incubating with 200 μ l elution buffer (100 mM NaHCO₃, 1% SDS in TE) for 15 min with rotation. Input samples and eluted samples were de-crosslinked overnight. Protein and RNA were digested with proteinase K and RNase A, respectively. DNA was isolated by phenol-chloroform extraction followed by an ethanol precipitation and analyzed by qPCR using StepOnePlus Real-Time PCR System (Thermo Fisher Scientific) and SYBR Green Master Mix (Thermo Fisher Scientific) or sequencing on the Illumina Next-Seq 2000.

After DNA extraction occupancy of different proteins were assessed by RT-PCR. Primers were used for *TFAP4* (forward: CCGGGCGCTGTTACTA; reverse: CAGGACACGGAGAACTACAG), *POLG* (forward: CTCTCAAGGAGCAGGTGGA; reverse: TCATAACCTCCCTTCGACCG), *NPM1* (forward: TTCACCGGAAGCATGG; reverse: CACGCGAGGTAAGTCTACG), Intergenic region (forward: TTTTCTCACATTGCCCCCTGT; reverse: TCAATGCTGTACCAGGCAAA), *NCL* (forward: CTACCACCCTCATCTGAATCC; reverse: TTGTCTCGCTGGGAAAGG), *NME1* (forward: GGGGTGGAGAGAAGAAAGCA; reverse: TGGGAGTAGGCAGTCATTCT), *PLD6* (forward: GCTGTGGGTCCCGATTA; reverse: CCTCCAGAGTCAGAGCCA), *TAF4B* (forward: AAGGTCGTCGCTCACAC; reverse: GCGTGGCTATATAAACATGGCT), *RPL22* (forward: CCGTAGCTTCTCTCTGCTC; reverse: ACCTCTGGGCTTCCTGTCT), *CCND2* (forward: GCCAGCTGCTGTTCTCCTTA; reverse: CCCCTCTCTTTCAATCTC). Shown analysis of RT-PCR show mean of technical triplicates as well as an overlay of each data point to indicate the distribution of the data.

For ChIP- or ChIP-Rx-seq, DNA was quantified using the Quant-iT PicoGreen dsDNA assay (Thermo Fisher Scientific). DNA library preparation was done using the NEBnext Ultra II DNA Library Prep Kit (New England Biolabs) following manufacturer's instructions. Quality of the library was assessed on the Fragment Analyzer (Agilent) using the NGS Fragment High Sensitivity Analysis Kit (1–6,000 bp; Agilent). Finally, libraries were subjected to cluster generation and base calling for 75 cycles on Illumina NextSeq 500 platform.

CUT&RUN followed by sequencing was performed as described in Meers et al. (Meers, Tenenbaum et al., 2019 [DOI](#)). 5×10^5 cells were harvested using Accutase (Sigma-Aldrich). Cells were washed in wash buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 0.5 mM Spermidine). After washing, cells were coupled to ConA-coated magnetic beads (Polysciences Europe), permeabilized and subsequently

incubated with the primary antibody (EXOSC5: NBP2-14952, 1:100, Novus Biologicals) in antibody binding buffer (wash buffer, 0.05% digitonin, 2 mM EDTA) over night at 4 °C. Cells were washed in Dig-Wash buffer (Wash buffer, 0.05% digitonin) and MNase (700 ng/ml) was added to each sample for 1 h at 4 °C. After washing with Dig-wash buffer and Low-Salt Rinse buffer (20 mM HEPES pH 7.5, 0.5 mM Spermidine, 0.05% digitonin) incubation buffer (3.5 mM HEPES pH 7.5, 10 mM CaCl₂, 0.05% digitonin) was added for 30 min at 4 °C. STOP buffer (170 mM NaCl, 20 mM EGTA, 0.05% digitonin, 20 µg/ml RNase A, 25 µg/ml Glycogen) was added to stop the MNase digestion. DNA fragments were released at 37 °C for 30 min. Decrosslinking was performed for 1 h at 50 °C after adding 0.01% SDS and 10 mg/ml Proteinase K followed by Phenol Chloroform extraction. Precipitated DNA was resuspended in 30 µl 0.1x TE buffer. For library preparation NEBnext Ultra II DNA Library Prep Kit (New England Biolabs) was used. Pre-PCR samples were purified using AmpureXP beads with a ratio of 1.75x. Then the eluted material was used with 16 PCR cycles. The libraries were cleaned using Agentcourt AMPure XP Beads (ratio 0.8x, Beckman Coulter), quality, quantity and fragment size assessed on the Fragment Analyzer (Agilent) using the NGS Fragment High Sensitivity Analysis Kit (1-6,000 bp; Agilent). Finally, libraries were subjected to cluster generation and base calling for 75 cycles on Illumina NextSeq 500 platform.

Phosphorylated linker Hi-C (pLHi-C) was based on Hi-C (Dixon, Selvaraj et al., 2012) and *in situ* Hi-C (Rao, Huntley et al., 2014). The main differences consisted in replacing the digestion enzyme and the Klenow fill-in step to increase resolution and overall protocol efficiency. 1x10⁶ cells per condition were fixed with formaldehyde (1% final concentration) for 10 min at room temperature (RT). Reaction was stopped by adding 125 mM glycine for 15 min at RT. Cells were harvested with ice-cold PBS supplemented with protease and phosphatase inhibitors (Sigma-Aldrich). Nuclei isolation was performed by adding buffer I (10 mM Tris-HCl pH 8, 10 mM NaCl, 0.5% NP-40, 1x protease inhibitors; Sigma-Aldrich) coupled with douncer homogenization. The resulting solution was centrifuged (2,500 xg, 5 min, 4 °C), the nuclei pellet was washed with 1x NEBuffer 3.1 (NEB) and incubated in buffer IIa (0.2% SDS, 1x NEBuffer 3.1; NEB) for 10 min at 50 °C. Buffer IIb (2% Triton X-100, 1.5 x NEBuffer 3.1, 1x protease inhibitors;) was added to a final volume of 943 µl. Sample was incubated for 15 min at 37 °C. 850 U DpnII (NEB) and 35 µl rSAP (NEB) were added and sample was incubated for at least 8 h at 37 °C. Modified DNA oligos, 5'-GATCCCCAAATCT-3' and 5'-GATCAGAT[BtdT]TGGG-3' with 5' end phosphate (Sigma-Aldrich), were annealed (81 µl of each Oligo 100 µM, 1x T4 ligase buffer; NEB) at 98 °C for 5 min, followed by 20 min at RT. After digestion, sample was washed with 1x T4 ligase buffer (NEB) (2,500 xg, 5 min, 4 °C). Nuclei pellet was resuspended in buffer IIIa (0.3% SDS, 1x T4 ligase buffer) and incubated for 15 min at 60 °C. Buffer IIIb (3% Triton X-100, 1x T4 ligase buffer, 1x protease inhibitors; NEB, Sigma-Aldrich) was added to a final volume of 1 ml and sample was incubated for 15 min at 37 °C. The nuclei pellet was resuspended in buffer IV (160 µl annealed oligos, 1.2x T4 ligase buffer, 6,000 U T4 ligase, 1x protease inhibitors; NEB) to a final volume of 800 µl after washing it with 1x T4 ligase buffer (NEB) (2,500 xg, 5 min, 4 °C). Sample was incubated overnight at 16 °C. rSAP in the digestion followed by phosphorylated linker in ligation was introduced to decrease the likelihood of self-ligation, increasing the final yield. Nuclei pellet was isolated by centrifugation (2,500 xg, 5 min, 4 °C) and mixed with buffer V (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.1% SDS, 1x protease inhibitors). Chromatin fragmentation was performed using the Covaris Focused Ultrasonicator M220 for 30 min per ml lysate. Samples were de-crosslinked overnight. Protein and RNA were digested with proteinase K and RNase A, respectively. DNA was isolated by ChIP DNA Clean & Concentrator (Zymo Research) according to the manufacturer's instructions. Biotin pull-down was performed with MyOne Streptavidin C1 beads (Thermo Fisher Scientific) according to manufacturer's instructions. Library preparation was performed on beads using NEBNext ChIP-Sequencing Library Prep Master Mix Set for Illumina (NEB). The manufacturer's instructions were followed, apart from introducing a washing step between each module of the protocol with buffer VI (3x, 5 mM Tris-HCl pH 8, 0.5 mM EDTA, 1 M NaCl, 0.05% Tween) followed by one with 1x TE buffer. Quality of the library was assessed on the Fragment Analyzer (Agilent) using the NGS Fragment High Sensitivity Analysis Kit (1-6,000 bp; Agilent). Libraries were subjected to cluster generation and base calling for 150 cycles paired end on Illumina NextSeq 500 platform.

Spike-in phosphorylated linker HiChIP (spLHiChIP) and phosphorylated linker HiChIP (pLHiChIP) were based on pLHi-C, ChIP-Rx, and the original HiChIP protocol (Mumbach et al., 2016 [DOI](#)). spLHiChIP and pLHiChIP benefit from the same modules of pLHi-C fixation, digestion, ligation, chromatin fragmentation, and library preparation. The difference consists in the ChIP module, which follows the exact same steps from IP of target protein to de-crosslinked overnight stated in the ChIP workflow. The ChIP module allocated between the chromatin fragmentation and library preparation in pLHi-C workflow defines the spLHiChIP and pLHiChIP workflows. spLHiChIP differs from pLHiChIP by adding murine NHO2A cells as an exogenous control (spike-in) at a 1:20 cell ratio during nuclei isolation. This allows normalization by murine paired end valid tags in a similar fashion as murine reads for ChIP-Rx. 10×10^6 cells per condition were fixed with formaldehyde (1% final concentration) for 10 min at room temperature (RT). Reaction was stopped by adding 125 mM glycine for 15 min at RT. Cells were harvested with ice-cold PBS supplemented with protease and phosphatase inhibitors (Sigma-Aldrich). Nuclei isolation was performed by adding buffer I (10 mM Tris-HCl pH 8, 10 mM NaCl, 0.5% NP-40, 1x protease inhibitors; Sigma-Aldrich) coupled with douncer homogenization. The resulting solution was centrifuged (2,500 xg, 5 min, 4 °C), the nuclei pellet was washed with 1x NEBuffer 3.1 (NEB) and incubated in buffer IIa (0.2% SDS, 1x NEBuffer 3.1; NEB) for 10 min at 50 °C. Buffer IIb (2% Triton X-100, 1.5 x NEBuffer 3.1, 1x protease inhibitors;) was added to a final volume of 943 μ l. Sample was incubated for 15 min at 37 °C. 850 U DpnII (NEB) and 35 μ l rSAP (NEB) were added and sample was incubated for at least 8 h at 37 °C. Modified DNA oligos, 5'-GATCCCCAAATCT-3' and 5'-GATCAGAT[BtndT]TGGG-3' with 5' end phosphate (Sigma-Aldrich), were annealed (81 μ l of each Oligo 100 μ M, 1x T4 ligase buffer; NEB) at 98 °C for 5 min, followed by 20 min at RT. After digestion, sample was washed with 1x T4 ligase buffer (NEB) (2,500 xg, 5 min, 4 °C). Nuclei pellet was resuspended in buffer IIIa (0.3% SDS, 1x T4 ligase buffer) and incubated for 15 min at 60 °C. Buffer IIIb (3% Triton X-100, 1x T4 ligase buffer, 1x protease inhibitors; NEB, Sigma-Aldrich) was added to a final volume of 1 ml and sample was incubated for 15 min at 37 °C. The nuclei pellet was resuspended in buffer IV (160 μ l annealed oligos, 1.2x T4 ligase buffer, 6,000 U T4 ligase, 1x protease inhibitors; NEB) to a final volume of 800 μ l after washing it with 1x T4 ligase buffer (NEB) (2,500 xg, 5 min, 4 °C). Sample was incubated overnight at 16 °C. Nuclei pellet was isolated by centrifugation (2,500 xg, 5 min, 4 °C) and mixed with buffer V (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.1% SDS, 1x protease inhibitors). Chromatin fragmentation was performed using the Covaris Focused Ultrasonicator M220 for 30 min per ml lysate. For each IP reaction, 100 μ l Dynabeads Protein A and Protein G (Thermo Fisher Scientific) were pre-incubated overnight with rotation in the presence of 5 mg/ml BSA and 10 μ g antibody (MYCN (B8.4.B): sc-53993, AB_831602; TFIIIC5: A301-242A, AB_890669). Chromatin was added to the beads, and IP was performed for a minimum of 6 h at 4 °C with rotation. Beads were washed three times each with washing buffer I (20 mM Tris pH 8.1, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS), washing buffer II (20 mM Tris pH 8.1, 500 mM NaCl, 2 mM EDTA, 1% Triton X-100, 0.1% SDS), washing buffer III (10 mM Tris pH 8.1, 250 mM LiCl, 1 mM EDTA, 1% NP-40, 1% sodium deoxycholate; including a 5 min incubation with rotation) and TE buffer. Chromatin was eluted twice by incubating with 200 μ l elution buffer (100 mM NaHCO₃, 1% SDS in TE) for 15 min with rotation. Input samples and eluted samples were de-crosslinked overnight. Protein and RNA were digested with proteinase K and RNase A, respectively. DNA was isolated by phenol-chloroform extraction followed by an ethanol precipitation. DNA library preparation was done using the NEBnext Ultra II DNA Library Prep Kit (New England Biolabs) following the manufacturer's instructions. Quality of the library was assessed on the Fragment Analyzer (Agilent) using the NGS Fragment High Sensitivity Analysis Kit (1-6,000 bp; Agilent). Finally, libraries were subjected to cluster generation and base calling on Illumina NextSeq 500 platform.

RNA sequencing was performed by extracting RNA with RNeasy Mini Kit (Qiagen) according to manufacturer's instructions. On-column DNase I digestion was performed followed by mRNA isolation with the NEBNext Poly (A) mRNA Magnetic Isolation Kit (NEB). Library preparation was done with the Ultra II Directional RNA Library Prep for Illumina following the manufacturer's manual. Libraries were size selected using SPRIselect Beads (Beckman Coulter) after amplification with 9 PCR cycles. Library quantification and size determination was performed with the

Fragment Analyzer (Agilent) using the NGS Fragment High Sensitivity Analysis Kit (1-6,000 bp; Agilent). Libraries were subjected to cluster generation and base calling for 100 cycles paired end on Illumina NextSeq 2000 platform.

Bioinformatics analysis and statistics

All libraries were subjected to NextSeq 500 or NextSeq 2000 (Illumina) sequencing following the manufacturer's guidelines. Base call data was converted and demultiplexed to FASTQ files by bcl2fastq Conversion Software v1.1.0 (Illumina). Quality control of each dataset was done using FastQC tool.

ChIP-seq and CUT&RUN sequencing reads were aligned to human (hg19/ GRCh37 assembly) using Bowtie2 v2.3.5.1. with default parameters (Langmead & Salzberg, 2012 [↗](#)). Normalization by aligned read number was performed on the input and on all correspondent IP samples. For each sample, the number of spike-in normalized reads was calculated by dividing the number of reads mapping exclusively to hg19 by the number of reads mapping exclusively to mm10 and multiplying this number with the smallest number of reads mapping to mm10 of all samples. After normalization, files were converted to bedGraph with "bedtools genomecov" v2.26 (Quinlan & Hall, 2010 [↗](#)). These files were used to plot genome browser track examples via the package plotgardener v1.012 in R v4.1.1.

MACS2 v2.1.2 (Zhang, Liu et al., 2008 [↗](#)) was used for peak calling with the following ChIP-seq datasets & parameters, with inputs serving as control: MYCN (GSM3044606, GSM3044608; dup = 5, $q = 1 \times 10^{-3}$), TFIIC5 in SH-EP-MYCN-ER (this work; dup = 5, $q = 10^{-3}$), and RAD21 (this work; dup = 5, $p = 10^{-2}$). H3K4me1, H3K4me3, and H3K27ac peaks were called with SICER v1.1 (Xu, Grullon et al., 2014 [↗](#)), using input as control and the parameters: redundancy threshold = 1, window size = 200 bp, fragment size = 75 bp, effective genome fraction = 0.74, gap size = 600, FDR = 0.001. Peak calling results were refined and confirmed by visual inspection.

RNA sequencing reads were aligned to human (hg19/ GRCh37 assembly) using STAR aligner v2.7.9a with the default parameters and gene quantification (--quantMode GeneCounts) (Dobin, Davis et al., 2013 [↗](#)). The output table with the gene counts was loaded into R v4.1.1. and processed with DESeq2 package accordingly to their manual (Love, Huber et al., 2014 [↗](#)). Genes with or more than 15 counts in at least 3 samples or more were selected, and all samples were normalized by size factors. The fold average for 3 independent biological replicates was calculated on treatment to control as showed in the correspondent figure and legend. The result was averaged in 100 or 150 bins for a total of 14,085 genes and displayed as dot plot. Alternative splicing events were identified using rMATS (Shen et al., 2014b [↗](#)).

pLHi-C, pLHiChIP and spLHiChIP sequencing reads were processed in two steps. In the first step they were all processed in the same way, HiC-Pro v 2.11.4 (Servant, Varoquaux et al., 2015) trims the reads for the linker sequence, aligns it to hg19 and filters it by DpnII restriction fragments of the same genome assembly. The default parameters were used for the alignment steps, except for the length of the seed (-L) parameter that was set to 25 for the global options and 15 for the local one, and mismatch (-N) that was set to 1. Read trimming was performed by inserting combinations of the linker sequence (GATCAGATTGGGGATC, GATCCCCAAATCTGATC) in the ligation site parameter. Reads considered as duplicates were accepted and multi-mapped reads were only removed for pLHiChIP and spLHiChIP. Quality control for all steps until all valid pairs (paired end tags; PETs) output was also performed thru HiC-Pro. We compared HiC-Pro's standard quality controls example (IMR90 replicate 1 sample; GSM862724), according to Servant et al. 2015 [↗](#) (Servant et al., 2015 [↗](#)), to the base of our methods variation, pLHi-C.

In the second step, spLHiChIP samples undergo the same workflow described for the first step replacing hg19 for mm10 assembly. PETs from mm10 were removed from hg19 PETs and a spike-in normalization was applied to hg19 PETs in the same way explained for ChIP-Rx hg19 aligned

reads. DNA loop calling was performed on pLHiChIP and pLHiChIP PETs using hicchipper v0.7.7 (Lareau & Aryee, 2018 [\[link\]](#)). ChIP-seq peak calling sites were used as pre-determined peaks. All the default parameters with a loop maximum distance (`--max-dist`) of 5,000,000 were used for all proteins. We compared the DNA loops calling quality control for Oct4 HiChIP (GSM2238510) from the original method description (Mumbach et al., 2016 [\[link\]](#)) to our method variation for MYCN and TFIIC5. An extra quality control to show enrichment and specificity was performed by plotting MYCN pLHiChIP, its input pLHi-C and its ChIP-seq with pLHiChIP and pLHi-C normalized to the same number of PETs. DNA loops calling results were refined and confirmed by visual inspection. The mango output files from the loop calling were then loaded in R, filter by a q-value equal or smaller than 0.01 and used in all further plots and analyses. All examples showing DNA loops, ChIP-seq and annotation tracks in different genomic regions/locus were plotted using plotgardener package in R environment.

The different genome functional annotations were retrieved from UCSC table browser (Karolchik, Hinrichs et al., 2004 [\[link\]](#)). Promoters, gene bodies and transcription end sites (TES) come from Reference Sequence (RefSeq) collection of genes for hg19 assembly and were adapted via R language. For RNAPII targets, promoters were defined as TSS $\pm$ 0.5 kb (Buchel et al., 2017 [\[link\]](#)), gene bodies as the sum of exons and introns, TES as TES $\pm$ 0.3 kb. Enhancers were defined as in Walz et al. (2014) [\[link\]](#) (Walz et al., 2014 [\[link\]](#)): H3K3me1 and H3K27ac overlapping sites without H3K4me3. Overlapping and discontinuous regions were obtained by using peak calling outputs as inputs in intersectBed (BEDTools) with standard parameters.

Overlapping between different pLHiChIP was calculated by “bedtools pairtopair” function (BEDTools) using the options to prevent self-hits and ignore strands. The type of overlap was set to report when both anchors of the different pLHiChIP overlaps (`-type either`). This provided outputs of loops with and without overlapping that could be used for other analyses. For discrimination between the different types of overlaps, the type both was removed from the type either and the two types were then reported. Overlaps were called joint sites and adapted as Venn diagrams by centering the coordinates and numbers to MYCN anchors or MYCN-TFIIC5 anchors. TADs overlap corresponds to common sites in at least eight out of nine cell lines reported in (GSE35156) (Rao et al., 2014 [\[link\]](#)).

The GenomicInteractions v1.28 (Harmston, Ing-Simmons et al., 2015 [\[link\]](#)) package in R language was used to calculate and report the overlaps between the different pLHiChIP anchors and their functional annotations. Overlaps on different functional annotations were reported in the following order of priority: enhancer, promoter, TES, gene body, tRNA, rRNA, miRNA, snRNA, snoRNA, SINE, LINE, LTR, and satellite. The bar graphs reporting the functional annotation of each anchor are proportional to the total number of loops of each pLHiChIP, numbers were rounded and plotted by GenomicInteractions and ggplot2 v3.3.5 packages in R environment. Motif analysis was performed by using the FIMO algorithm (Grant, Bailey et al., 2011 [\[link\]](#)) of the MEME Suite software toolkit v5.3.3 (Bailey, Johnson et al., 2015 [\[link\]](#)). The function getfasta (BEDTools) extracted the hg19 FASTA sequence from each pLHiChIP anchor to be used independently as input for motif search. The background model was generated by the function fasta-get-markov (MEME Suite) using hg19 FASTA with standard parameters. Pre-defined motifs for MYCN (E-box) (MA0104.4: JASPAR 2018 (Sandelin, Alkema et al., 2004 [\[link\]](#))), B-box, and A-box were scanned using the options “`-thresh 1.0 --text`” with the default parameters (Chan, Lin et al., 2021 [\[link\]](#), Pavesi, Conterio et al., 1994 [\[link\]](#)). Resulting hits were then filtered by $p\text{-value} \leq 1 \times 10^{-4}$. In R environment, duplicated hits of the same motif in the same anchor were filtered by the one with the lowest p-value. Different motifs in the same anchor window were reported according to the decreasing priority: E-box, B-box, and A-box. The overlap between anchors and motifs was computed by GenomicInteractions package and the corresponding heatmap was plotted in R.

To visualize the influence of the MYCN interaction on gene expression (**Fig 1C**), MYCN bound genes (as defined by MACS2 on MYCN ChIP-seq data, above) were subdivided into genes that are bound by MYCN without loops (“no MYCN loops”) and genes bound by MYCN forming loops (above, “MYCN loops”). For both classes, RNA-sequencing reads between TSS & TSS+2 kb were counted using “bedtools intersect” on spike-normalized ChIP-seq data for RNAPII (A10), MYCN (GSM3044606), TFIIC5; processed transcript data were obtained from Büchel *et al.* 2017 (RNA-sequencing, GSE111905 (Büchel *et al.*, 2017)) and Papadopoulos *et al.* 2021 (4sU-sequencing, GSE164569 (Papadopoulos *et al.*, 2022)), respectively. Plots were generated with “boxplot” and statistical significance calculated using Students’ t-test (unpaired, two-sided, unequal variance), both in R v3.6.3.

Over-representation analysis for Molecular Signatures Database (MsigDB (Subramanian, Tamayo *et al.*, 2005)) were calculated by clusterProfiler package (Wu, Hu *et al.*, 2021) in R environment. Different functional classes of loops were retrieved from the overlaps and discontinuous loops by the GenomicInteractions package. Gene names were converted to HGNC symbol name using AnnotationDbi package and sorted by the number of PETs. Duplicated gene names were merged and the median of their number of PETs was used. A hypergeometric test was calculated on the sorted list for MSigDB collections H (hallmark gene sets) or C5 (ontology gene sets) using a 0.05 q-value threshold.

MYCN spLHiChIP treated sample was normalized by its control counterpart in each biological replicate before plotted as graph bar in R environment. The ratio of TFIIC5 binding (ChIP-Rx) per number of interaction (spLHiChIP) was calculated in R environment v4.1.1. The number of ChIP-Rx reads was calculated to TSS ± 2 kb for all genes using bedtools intersect (BEDtools) with default parameters. The number of spLHiChIP interactions was calculated for the same region. The ratio was calculated on reads to number of interactions per each sample and plotted as bar graph.

Networks were reconstituted by using the R package igraph v1.2.11 and Cytoscape v3.9 (Brockmann, Poon *et al.*, 2013). The different classes of loops were retrieved via the GenomicInteractions package and converted to the simple interaction file format and to a node table. These files were used in Cytoscape for visualization, using the Edge-weighted Spring-Embedded Layout based on the number of PETs for each loop. The resulting MYCN network reports the nodes as functional annotations; explained before. In parallel, the sif file was used as input for the igraph package. This package was used to define the membership of each anchor to the different clusters and to quantify the number of nodes per cluster. The statistical significance comparing the number of nodes per cluster between two groups were calculated via Wilcoxon rank sum test.

Density plots for the different regions were performed using ngs.plot v2.41.3 (Shen, Shao *et al.*, 2014a). Default parameters were used for samples normalized by reads number. For spike-in normalized samples, we used a custom variation of ngs.plot that allows the spike-in normalization by skipping the internal normalization step. All densities plots and metaplots were based on gene expression for SH-EP-MYCN-ER cells ± MYCN induction (4-OHT treatment) – evaluated by RNA-sequencing (GSE111905; Herold *et al.*, 2019). Based on this, all expressed genes gene set consists of 14,704 genes, down- and up-regulated gene sets are represented by 613 and 921 genes, respectively. MYCN-down and -up regulated genes are defined by negative and positive fold change upon ±4-OHT treatment of genes with q-values less than 0.05 and expression filter by counts per million. In R v4.1.1., Wilcoxon rank sum test (unpaired, two-sided) was used to compare the different conditions on the plotted region per ChIP.

PLA’s p-values were calculated by comparing the signal of the pool of the cells in all replicates using Wilcoxon rank sum test (unpaired, two-sided) in R v4.1.1.

Materials, data and code availability

All plasmids generated are available upon request.

The following published datasets were used: MYCN ChIP-seq and RNA-seq: GSE111905; 4sU-seq: GSE164569. Reviewers can access data from this work under GEO accession GSE223058.

Supplementary Figures




Figure S1

Characterization of MYCN/TFIIIC complexes.

- A.** SDS-PAGE gel of recombinant purified MYCN (aa 1-137) expressed in *E. coli* cells.
- B.** SDS-PAGE gel of recombinant purified 1A subcomplex which was expressed in *Spodoptera frugiperda* (Sf9) cells.
- C.** Deconvoluted spectra of native mass spectrometry for the 1A/3x FLAG-MYCN complex. Masses are shown in red, identities of complexes are marked in blue.
- D.** Box plot of the 30 possible 1A/3x FLAG-MYCN complex molecular weights based on the masses observed using intact mass spectrometry. The red line indicates the mass observed for the complex by native mass spectrometry (204,763 Da).
- E.** Immunoblot showing levels of TFIIIC2, TFIIIC3 or TFIIIC5 and MYC in SH-EP-MYCN-ER cells expressing doxycycline (Dox)-inducible shRNAs targeting the different *TFIIIC* subunits (n = 3). Where indicated cells were treated with Dox (1 µg/ml, 48 h) and/or 4-OHT (200 nM, 4 h), respectively. EtOH was used as control. In the following panels addition of 4-OHT is indicated by "+ MYCN" and Dox treatment by "- TFIIIC".
- F.** Immunoblot of MYC and MYCN in SH-EP-MYCN-ER cells after induction of MYCN with 4-OHT (200 nM, 4 h). VCL was used as a loading control.




Figure S2.

Effects of MYCN and TFIIC on RNA polymerase II.

A. Metagenes plot of ChIP-Rx signal for non-phosphorylated RNAPII. Data show mean (line) $\pm$ standard error of the mean (SEM indicated by the shade) of different gene sets based on an RNA-seq of SH-EP-MYCN-ER cells $\pm$ 4-OHT.

B. Metagenes plot of ChIP-Rx signal for RNAPII pSer2. Data are presented as described in A.

C. Average density plot of ChIP-Rx signal for non-phosphorylated RNAPII in SH-EP cells treated with 4-OHT. Data show mean (line) $\pm$ SEM (indicated by the shade) of different gene sets based on an RNA-seq of SH-EP-MYCN-ER cells $\pm$ 4-OHT. The signal is centered to the TSS $\pm$ 2 kb (n = 2).

D. Average bin dot plot for RNA-seq of SH-EP-MYCN-ER showing mRNA expression normalized by control per bin. Cells were co-treated with 1 μ g/ml Dox (" - TFIIC3", 48 h) and/or 4-OHT (" + MYCN", 4 h) as EtOH as control. Expression was normalized by its control. Each bin represents 100 genes of a total of 12,091 genes. Dotted line marks the relative expression at 1 (n = 3).

E. Browser tracks for TFIIC5 ChIP-Rx at the indicated gene loci. SH-EP-MYCN-ER cells were treated with DRB and/or 4-OHT, respectively. EtOH was used as control.




Figure S3

Characterization of HiChIP Methods.

- A.** Diagram showing the workflow of pLHiChIP. The workflow corresponds to the original HiChIP protocol. For a detailed description, see Methods.
- B.** MYCN and MYC ChIP in SH-EP-MYCN-ER cells before and after activation of MYCN with 4-OHT (200 nM, 24 h). Shown is the mean and points of technical triplicates normalized to an intergenic region ($n = 2$).
- C.** (Top) Diagram illustrating the difference between valid and invalid pairs after mapping of sPLHiChIP data. Valid pairs harbor non-contiguous elements of chromosomal DNA, whereas invalid pairs harbor contiguous stretches of chromosomal DNA. (Bottom) Bar graph showing percentage of valid and invalid reads in our dataset (right) compared to a published HiC protocol (left).
- D.** Table showing quality controls for MYCN and TFIIC5 sPLHiChIP compared to the original Oct4 HiChIP.
- E.** Representative example of pLHiChIP track for MYCN (red) and a pLHiC track (blue) showing the strongly increased PETs number of the MYCN interactions after MYCN-immunoprecipitation relative to input (pLHi-C). Data are superimposed with a browser view of a MYCN ChIP-seq.
- F.** Table showing interactions involving *tRNA* genes in MYCN pLHiChIP.
- G.** Table showing the top three terms, their p-values and the FDR from enrichment analysis for MSigDB C5 collection of all MYCN network.




Figure S4

Three-dimensional interactions of MYCN and TFIIIC.

A. Heatmap showing analysis of sequence motifs characteristic for interacting boxes on both anchors of the interactions.

Color reflects the number of interactions.

B. Table summarizing numbers of MYCN interactions in the presence and absence of TFIIIC5 in SH-EP-MYCN-ER cells and the TFIIIC5-dependent change. Data merged of 2 independent biological replicates.

C. Table displaying total numbers of TFIIIC5 interactions in SH-EP-MYCN-ER ± MYCN + DRB.




Figure S5

Effects of TFIIC on splicing and termination factors.

A. Volcano Plot depicting changes in intron retention upon TFIIC3 knock-down in the absence (left) and presence (right) of MYCN. Triangle reflects the effect of the splicing error after TFIIC knock-down.

B. Volcano Plot depicting changes in exon skipping upon TFIIC3 knock-down in the absence (left) and presence (right) of MYCN. Triangle reflects the effect of the splicing error after TFIIC knock-down.

C. Controls for proximity ligation assays (PLAs) shown in [Figure 5A](#). Immunofluorescence show specificity of the antibody. Single antibodies PLAs were performed in parallel to each PLA. Mean of signal in the nucleus and cytoplasm was calculated and compared to mean of the signal per nucleus of the PLA.

D. Boxplots showing the number of proximity ligation assay (PLA) signals between RNAPII and TFIIC5. SH-EP-MYCN-ER cells were treated with 1 μ g/ml Dox ("– TFIIC5", 48 h) and/or 4-OHT ("+ MYCN"). EtOH was used as control. For clarity purposes, 500 cells pooled from different replicates were plotted. p-values were calculated comparing the PLA signal of all cells using unpaired Wilcoxon rank sum test. The grey dotted line indicates the median in the control condition (n = 3).

E. BRCA1 ChIP in SH-EP-MYCN-ER cells expressing a Dox-inducible shRNA targeting *TFIIC5* treated with 4-OHT. Data show fold change of BRCA1 binding after induction of MYCN in the presence (blue) or absence (red) of TFIIC5 (n = 2).

References

- Bailey TL, Johnson J, Grant CE, Noble WS (2015) **The MEME Suite** *Nucleic Acids Res* **43**:W39–49
- Baluapuri A *et al.* (2019) **MYC Recruits SPT5 to RNA Polymerase II to Promote Processive Transcription Elongation** *Mol Cell* **74**:674–687
- Baluapuri A, Wolf E, Eilers M (2020) **Target gene-independent functions of MYC oncoproteins** *Nature reviews* **21**:255–267
- Bayliss R, Sardon T, Vernos I, Conti E (2003) **Structural basis of Aurora-A activation by TPX2 at the mitotic spindle** *Mol Cell* **12**:851–62
- Blackwood EM, Eisenman RN (1991) **Max: A helix-loop-helix zipper protein that forms a sequence-specific DNA-binding complex with myc** *Science* **251**:1211–1217
- Blackwood EM, Lüscher B, Eisenman RN (1992) **Myc and Max associate in vivo** *Genes Dev* **6**:71–80
- Boija A *et al.* (2018) **Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains** *Cell* **175**:1842–1855
- Brockmann M *et al.* (2013) **Small molecule inhibitors of aurora-a induce proteasomal degradation of N-myc in childhood neuroblastoma** *Cancer cell* **24**:75–89
- Buchel G *et al.* (2017) **Association with Aurora-A Controls N-MYC-Dependent Promoter Escape and Pause Release of RNA Polymerase II during the Cell Cycle** *Cell reports* **21**:3483–3497
- Chan PP, Lin BY, Mak AJ, Lowe TM (2021) **tRNAscan-SE 2.0: improved detection and functional classification of transfer RNA genes** *Nucleic Acids Res* **49**:9077–9096
- Cortazar MA, Erickson B, Fong N, Pradhan SJ, Ntini E, Bentley DL (2022) **Xrn2 substrate mapping identifies torpedo loading sites and extensive premature termination of RNA pol II transcription** *Genes & development* **36**:1062–1078
- Cortazar MA, Sheridan RM, Erickson B, Fong N, Glover-Cutter K, Brannan K, Bentley DL (2019) **Control of RNA Pol II Speed by PNUTS-PP1 and Spt5 Dephosphorylation Facilitates Termination by a “Sitting Duck Torpedo” Mechanism** *Mol Cell* **76**:896–908
- Cossa G, Parua PK, Eilers M, Fisher RP (2021) **Protein phosphatases in the RNAPII transcription cycle: erasers, sculptors, gatekeepers, and potential drug targets** *Genes Dev* **35**:658–676
- Crepaldi L, Policarpi C, Coatti A, Sherlock WT, Jongbloets BC, Down TA, Riccio A (2013) **Binding of TFIIC to sine elements controls the relocation of activity-dependent neuronal genes to transcription factories** *PLoS Genet* **9**
- Dang CV (2012) **MYC on the path to cancer** *Cell* **149**:22–35

- Das SK *et al.* (2022) **MYC assembles and stimulates topoisomerases 1 and 2 in a “topoisome”** *Mol Cell* **82**:140–158
- Das SK, Lewis BA, Levens D (2022) **MYC: a complex problem** *Trends Cell Biol*
- Dhanasekaran R, Deutzmann A, Mahauad-Fernandez WD, Hansen AS, Gouw AM, Felsher DW (2022) **The MYC oncogene - the grand orchestrator of cancer growth and immune evasion** *Nature reviews Clinical oncology* **19**
- Dhanasekaran R *et al.* (2023) **MYC Overexpression Drives Immune Evasion in Hepatocellular Carcinoma That Is Reversible through Restoration of Proinflammatory Macrophages** *Cancer research* **83**:626–640
- Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B (2012) **Topological domains in mammalian genomes identified by analysis of chromatin interactions** *Nature* **485**:376–80
- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) **STAR: ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21
- Eberhardy SR, Farnham PJ (2002) **Myc recruits P-TEFb to mediate the final step in the transcriptional activation of the cad promoter** *The Journal of biological chemistry* **277**
- Endres T *et al.* (2021) **Ubiquitylation of MYC couples transcription elongation with double-strand break repair at active promoters** *Mol Cell* **81**:830–844
- Estell C, Davidson L, Eaton JD, Kimura H, Gold VAM, West S (2023) **A restrictor complex of ZC3H4 WDR82, and ARS2 integrates with PNUTS to control unproductive transcription.** *Mol Cell* **83**
- Ferrari R *et al.* (2020) **TFIIIC Binding to Alu Elements Controls Gene Expression via Chromatin Looping and Histone Acetylation** *Mol Cell* **77**:475–487
- Gerlach P, Garland W, Lingaraju M, Salerno-Kochan A, Bonneau F, Basquin J, Jensen TH, Conti E (2022) **Structure and regulation of the nuclear exosome targeting complex guides RNA substrates to the exosome** *Mol Cell* **82**:2505–2518
- Grant CE, Bailey TL, Noble WS (2011) **FIMO: scanning for occurrences of a given motif** *Bioinformatics* **27**:1017–8
- Hanahan D (2022) **Hallmarks of Cancer: New Dimensions** *Cancer Discov* **12**:31–46
- Harmston N, Ing-Simmons E, Perry M, Baresic A, Lenhard B (2015) **GenomicInteractions: An R/Bioconductor package for manipulating and investigating chromatin interaction data** *BMC Genomics* **16**
- Heidelberger JB, Voigt A, Borisova ME, Petrosino G, Ruf S, Wagner SA, Beli P (2018) **Proteomic profiling of VCP substrates links VCP to K6-linked ubiquitylation and c-Myc function** *EMBO Rep*
- Herold S *et al.* (2019) **Recruitment of BRCA1 limits MYCN-driven accumulation of stalled RNA polymerase** *Nature* **567**:545–549
- Hnisz D, Shrinivas K, Young RA, Chakraborty AK, Sharp PA (2017) **A Phase Separation Model for Transcriptional Control** *Cell* **169**:13–23

- Jaenicke LA, von Eyss B, Carstensen A, Wolf E, Xu W, Greifenberg AK, Geyer M, Eilers M, Popov N (2016) **Ubiquitin-Dependent Turnover of MYC Antagonizes MYC/PAF1C Complex Accumulation to Drive Transcriptional Elongation** *Mol Cell* **61**:54–67
- Kalkat M *et al.* (2018) **MYC Protein Interactome Profiling Reveals Functionally Distinct Regions that Cooperate to Drive Tumorigenesis** *Mol Cell* **72**:836–848
- Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, Kent WJ (2004) **The UCSC Table Browser data retrieval tool** *Nucleic Acids Res* **32**:D493–6
- Kilchert C, Wittmann S, Vasiljeva L (2016) **The regulation and functions of the nuclear RNA exosome complex** *Nature reviews* **17**:227–39
- Kress TR, Sabo A, Amati B (2015) **MYC: connecting selective transcriptional control to global RNA production** *Nat Rev Cancer* **15**:593–607
- Landsverk HB, Sandquist LE, Bay LTE, Steurer B, Campsteijn C, Landsverk OJB, Marteiijn JA, Petermann E, Trinkle-Mulcahy L, Syljuasen RG (2020) **WDR82/PNUTS-PP1 Prevents Transcription-Replication Conflicts by Promoting RNA Polymerase II Degradation on Chromatin** *Cell reports* **33**
- Langmead B, Salzberg SL (2012) **Fast gapped-read alignment with Bowtie 2** *Nature methods* **9**:357–9
- Lareau CA, Aryee MJ (2018) **hichipper: a preprocessing pipeline for calling DNA loops from HiChIP data** *Nature methods* **15**:155–156
- Lewis BA, Das SK, Jha RK, Levens D (2023) **Self-assembly of promoter DNA and RNA Pol II machinery into transcriptionally active biomolecular condensates** *Sci Adv* **9**
- Lim B, Levine MS (2021) **Enhancer-promoter communication: hubs or loops?** *Curr Opin Genet Dev* **67**:5–9
- Lourenco C *et al.* (2021) **MYC protein interactors in gene transcription and cancer** *Nat Rev Cancer*
- Love MI, Huber W, Anders S (2014) **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2** *Genome Biol* **15**
- Lu H, Yu D, Hansen AS, Ganguly S, Liu R, Heckert A, Darzacq X, Zhou Q (2018) **Phase-separation mechanism for C-terminal hyperphosphorylation of RNA polymerase II** *Nature* **558**:318–323
- Mancebo HS *et al.* (1997) **P-TEFb kinase is required for HIV Tat transcriptional activation in vivo and in vitro** *Genes & development* **11**:2633–44
- Marty MT, Baldwin AJ, Marklund EG, Hochberg GK, Benesch JL, Robinson CV (2015) **Bayesian deconvolution of mass and ion mobility spectra: from binary interactions to polydisperse ensembles** *Anal Chem* **87**:4370–6
- Meers MP, Tenenbaum D, Henikoff S (2019) **Peak calling by Sparse Enrichment Analysis for CUT&RUN chromatin profiling** *Epigenetics & chromatin* **12**

- Moqtaderi Z, Wang J, Raha D, White RJ, Snyder M, Weng Z, Struhl K (2010) **Genomic binding profiles of functionally distinct RNA polymerase III transcription complexes in human cells** *Nat Struct Mol Biol* **17**:635–40
- Mumbach MR, Rubin AJ, Flynn RA, Dai C, Khavari PA, Greenleaf WJ, Chang HY (2016) **HiChIP: efficient and sensitive analysis of protein-directed genome architecture** *Nat Methods* **13**:919–922
- Gonzalez M, Noe, Blears D, Svejstrup JQ (2021) **Causes and consequences of RNA polymerase II stalling during transcript elongation** *Nature reviews* **22**:3–21
- Noma K, Cam HP, Maraia RJ, Grewal SI (2006) **A role for TFIIC transcription factor complex in genome organization** *Cell* **125**:859–72
- Oksuz O *et al.* (2023) **Transcription factors interact with RNA to regulate genes** *Mol Cell* **83**:2449–2463
- Oler AJ *et al.* (2010) **Human RNA polymerase III transcriptomes and relationships to Pol II promoter chromatin and enhancer-binding factors** *Nat Struct Mol Biol* **17**:620–8
- Orioli A, Pascali C, Pagano A, Teichmann M, Dieci G (2012) **RNA polymerase III transcription control elements: themes and variations** *Gene* **493**:185–94
- Palacio M, Taatjes DJ (2022) **Merging Established Mechanisms with New Insights: Condensates, Hubs, and the Regulation of RNA Polymerase II Transcription** *J Mol Biol* **434**
- Papadopoulos D *et al.* (2022) **MYCN recruits the nuclear exosome complex to RNA polymerase II to prevent transcription-replication conflicts** *Mol Cell* **82**:159–176
- Papadopoulos D, Uhl L, Ha SA, Eilers M (2023) **Beyond gene expression: how MYC relieves transcription stress** *Trends Cancer*
- Pavesi A, Conterio F, Bolchi A, Dieci G, Ottonello S (1994) **Identification of new eukaryotic tRNA genes in genomic DNA databases by a multistep weight matrix analysis of transcriptional control regions** *Nucleic Acids Res* **22**:1247–56
- Pellanda P *et al.* (2021) **Integrated requirement of non-specific and sequence-specific DNA binding in Myc-driven transcription** *The EMBO journal* **40**
- Policarpi C, Crepaldi L, Brookes E, Nitarska J, French SM, Coatti A, Riccio A (2017) **Enhancer SINEs Link Pol III to Pol II Transcription in Neurons** *Cell reports* **21**:2879–2894
- Quinlan AR, Hall IM (2010) **BEDTools: a flexible suite of utilities for comparing genomic features** *Bioinformatics* **26**
- Raab JR, Chiu J, Zhu J, Katzman S, Kurukuti S, Wade PA, Haussler D, Kamakaka RT (2012) **Human tRNA genes function as chromatin insulators** *The EMBO journal* **31**:330–50
- Rahl PB, Lin CY, Seila AC, Flynn RA, McCuine S, Burge CB, Sharp PA, Young RA (2010) **c-Myc regulates transcriptional pause release** *Cell* **141**
- Rao SS *et al.* (2014) **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping** *Cell* **159**:1665–80

- Rickman DS, Schulte JH, Eilers M (2018) **The Expanding World of N-MYC-Driven Tumors** *Cancer Discov* **8**:150–163
- Rodriguez-Molina JB, West S, Passmore LA (2023) **Knowing when to stop: Transcription termination on protein-coding genes by eukaryotic RNAPII** *Mol Cell* **83**:404–415
- Roeschert I *et al.* (2021) **Combined inhibition of Aurora-A and ATR kinase results in regression of MYCN-amplified neuroblastoma** *Nat Cancer* **2**:312–326
- Sabo A *et al.* (2014) **Selective transcriptional regulation by Myc in cellular growth control and lymphomagenesis** *Nature* **511**:488–492
- Sandelin A, Alkema W, Engstrom P, Wasserman WW, Lenhard B (2004) **JASPAR: an open-access database for eukaryotic transcription factor binding profiles** *Nucleic Acids Res* **32**:D91–4
- Schmid M, Jensen TH (2019) **The Nuclear RNA Exosome and Its Cofactors** *Adv Exp Med Biol* **1203**:113–132
- Servant N, Varoquaux N, Lajoie BR, Viara E, Chen CJ, Vert JP, Heard E, Dekker J, Barillot E (2015) **HiC-Pro: an optimized and flexible pipeline for Hi-C data processing** *Genome Biol* **16**
- Shen L, Shao N, Liu X, Nestler E (2014) **ngs.plot: Quick mining and visualization of next-generation sequencing data by integrating genomic databases** *BMC Genomics* **15**
- Shen S, Park JW, Lu ZX, Lin L, Henry MD, Wu YN, Zhou Q, Xing Y (2014) **rMATS: robust and flexible detection of differential alternative splicing from replicate RNA-Seq data** *Proceedings of the National Academy of Sciences of the United States of America* **111**:E5593–601
- Solvie D *et al.* (2022) **MYC multimers shield stalled replication forks from RNA polymerase** *Nature*
- Subramanian A *et al.* (2005) **Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles** *Proceedings of the National Academy of Sciences of the United States of America* **102**:15545–50
- Tamara S, den Boer MA, Heck AJR (2022) **High-Resolution Native Mass Spectrometry** *Chemical reviews* **122**:7269–7326
- Thomas LR *et al.* (2015) **Interaction with WDR5 promotes target gene recognition and tumorigenesis by MYC** *Mol Cell* **58**:440–52
- Van Bortle K, Corces VG (2013) **The role of chromatin insulators in nuclear architecture and genome function** *Curr Opin Genet Dev* **23**:212–8
- Van Bortle K, Nichols MH, Li L, Ong CT, Takenaka N, Qin ZS, Corces VG (2014) **Insulator function and topological domain border strength scale with architectural protein occupancy** *Genome Biol* **15**
- Vervoort SJ *et al.* (2021) **The PP2A-Integrator-CDK9 axis fine-tunes transcription and can be targeted therapeutically in cancer** *Cell* **184**:3143–3162
- Vo BT *et al.* (2016) **The Interaction of Myc with Miz1 Defines Medulloblastoma Subgroup Identity** *Cancer cell* **29**:5–16

Walz S *et al.* (2014) **Activation and repression by oncogenic MYC shape tumour-specific gene expression profiles** *Nature* **511**:483–7

Wang J, Pan Y, Shen S, Lin L, Xing Y (2017) **rMATS-DVR: rMATS discovery of differential variants in RNA** *Bioinformatics* **33**:2216–2217

Wu T *et al.* (2021) **clusterProfiler 4.0: A universal enrichment tool for interpreting omics data** *Innovation (N Y)* **2**

Xu S, Grullon S, Ge K, Peng W (2014) **Spatial clustering for identification of ChIP-enriched regions (SICER) to map regions of histone methylation patterns in embryonic stem cells** *Methods Mol Biol* **1150**:97–111

Yang J, Chung C-I, Koach J, Liu H, Zhao Q, Yang X, Shen Y, Weiss WA, Shu X (2022) **Phase separation of Myc differentially regulates gene transcription** *Biorxiv*

Zhang Y *et al.* (2008) **Model-based analysis of ChIP-Seq (MACS)** *Genome Biol* **9**

Editors

Reviewing Editor

Andrés Aguilera

CABIMER, Universidad de Sevilla, Seville, Spain

Senior Editor

Tony Ng

King's College London, London, United Kingdom

Reviewer #1 (Public Review):

Summary:

In this manuscript entitled "Association with TFIIC limits MYCN accumulation in hubs of active promoters and chromatin accumulation of non-phosphorylated RNA polymerase II" the authors examine how the cohesin complex component (and RNA pol III associated factor) TFIIC interacts with MYCN and controls transcription. They confirm that TFIIC co-purifies with MYCN, dependent on its amino terminus, as shown in previous work. The authors also find that TFIIC and MYCN are both found in promoter hubs and suggest that TFIIC inhibits MYCN association with these hubs. Finally, the authors indicate that TFIIC/MYCN alter exosome function, and BRCA1 dependent effects, at MYCN regulated loci.

In the revised manuscript the authors have adequately addressed or responded to our questions and comments. The exception concerns point #2 in our initial review:

(2) The authors indicate in Figure 2 that TF3C has essentially no effect on MYCN- dependent gene expression and/or transcription elongation. Yet a previous study (PMID: 29262328) associated with several of the same authors concluded that TF3C positively affects transcription elongation. The authors to not attempt to reconcile these disparate results and the point still needs to be clarified.

Authors' Response

We agree that the data in this manuscript do not support the role on transcription elongation. This point was also raised by Reviewer 3. Comparing our new results to the data published

previously we can summarize that the data sets in the two studies show three key results: First, the traveling ratio of RNAPII changes upon induction of MYCN. Second, RNAPII decreases at the transcription start side and third, it increases towards the end side.

We agree that in the previous study we linked the traveling ratio directly to elongation. However performing ChIP-seq with different RNAPII antibodies showed us that for example RNAPII (N20), which is unfortunately discontinued, gives different results compared to RNAPII (A10). Combining our new results using the RNAPII (8WG16) antibody shows that the traveling ratio is not only reflecting transcription elongation but also includes that the RNAPII is kicked-off chromatin at the start side.

Reviewer revised response:

The explanation for the change in interpretation of the previous study (Buchel, et. al., 2017) in light of the differing results using different RNA pol2 antibodies used in the present study seems reasonable. However the final manuscript may well result in some confusion in the literature in regards to TF3C and elongation. This is because, while the authors refer to the earlier paper frequently, they do not directly discuss the re-interpretation of the elongation conclusion of the earlier paper. It seems likely that a reader of the present paper will find this issue confusing when trying to reconcile the results of the two papers.

<https://doi.org/10.7554/eLife.94407.2.sa3>

Reviewer #2 (Public Review):

This manuscript reports several interesting observations that invite follow-up. The notion that hubs, and perhaps condensates that may (or may not embrace them) are functionally and physiologically important is an open issue at this time. The authors note that TFIIC helps to prune extraneous connections from hubs, but do not comment that the connections that are maintained are also reinforced. At the same time only modest changes in gene expression associated with expanded or decreased connections and changes in bound proteins. One interesting possibility might be that standard methods for assessing expression miss changes global or background transcription. It seems that the TFIIC-MYCN-ER connection has features that would help to suppress such background. The results invite a more global consideration of TFIIC than as primarily RNAPIII/small RNA transcription factor and of MYCN as an E-box dependent transcription factor. The results use state of the art methods to develop interesting new ideas that have the potential to instruct further studies that may reveal new mechanisms of action for TFIIC and MYCN.

The work is however subject to a couple of caveats. First, the authors should be more cautious when drawing firm conclusions about the dynamics and kinetics of transcription from the static snapshots obtained from most genomic methods. For example, please take a look at Figure 1F of "Transcription elongation defects link oncogenic SF3B1 mutations to targetable alterations in chromatin landscape" by Buddu et al, <https://doi.org/10.1016/j.molcel.2024.02.032>. Here, an increase in RNAPser2P is seen in gene bodies and a bit at the TES- superficially inviting the conclusion that expression is increased (a similar erroneous conclusion has been claimed in other genomic studies), but the increase is in fact, not due to increased transcription, rather to impaired elongation- this conclusion required performing TT-Seq which allowed inferences to be made about elongation rates. Acknowledging this qualification would help advise the reader.

The authors also need to discuss directly what differences between the MYC predominant SH-EP cells and the MYCN-predominant SH-EP-MYCNER+tamoxifen are qualitative versus quantitative. MYCNER indeed associates much more with chromatin than did MYC, but there seems to be a lot more MYCER than there was MYC prior to the addition of tamoxifen. (The true control for this would be to prepare SH-EP-MYCNER cells expressed from the same

promoter as was MYCNER. Some discussion of qualitative versus quantitative differences should be acknowledged.

Strengths:

Use of a variety of methods to assess the genomic response to increased MYCN in the presence or absence of TFIIC. Clearly establishes in vitro and in vivo the TFIIC-MYCN complex

Weaknesses:

Dynamic inferences are made without kinetic experiments.

<https://doi.org/10.7554/eLife.94407.2.sa2>

Reviewer #3 (Public Review):

Summary:

Vidal et al. investigated how TFIIC may mediate MYCN effects on transcription. The work builds upon previous reports from the same group where they describe MYCN interactors in neuroblastoma cells (Buchel et al, 2017), which include TFIIC, and their different roles in MYCN-dependent control of RNA polymerase II function (Herold et al, 2019) (Roeschert et al, 2021) (Papadopoulos et al, 2022). Using baculovirus expression systems, they confirm that MYCN-TFIIC interaction is direct, and likely relevant for neuroblastoma cell proliferation. However, transcriptomics analyses led them to conclude that TFIIC is largely dispensable for MYCN-dependent gene expression. Instead, they propose that TFIIC limits MYCN-mediated promoter-promoter 3D chromatin contacts, which would in turn facilitate the recruitment of the nascent RNA degradation machinery and restrict the accumulation of non-phosphorylated RNA polymerase II at promoters. How this mechanism may impact on MYCN-driven neuroblastoma cell biology remains to be elucidated.

Strengths:

This study presents a nice variety of genomic datasets addressing the specific role of TFIIC in MYCN-dependent functions. In particular, the technically challenging HiChIP sequencing experiments performed under various conditions provide very useful information about the interplay between MYCN and TFIIC in the regulation of 3D chromatin contacts. The authors show that MYCN and TFIIC participate both in unique and overlapping long-range chromatin contacts and that the expression of each of these proteins limits the function of the other. Together, their results suggest a dynamic and interconnected relationship between MYCN and TFIIC in regulating 3D chromatin contacts.

Weaknesses:

(1) Mechanistic questions regarding the specific role of TFIIC in regulating MYCN function remain unsolved. Why is it important to restrict MYCN association to promoter hubs? Do the authors find any TFIIC-dependent phenotype that is restricted or particularly enhanced at these locations? Both the effects on the accumulation of non-phosphorylated RNA pol II and the recruitment of the nascent RNA degradation machinery seem to be global.

(2) Two specific points regarding RNA pol II ChIPseq results remain unclear:

-It is unfortunate that although both RNAPII (N20) and RNAPII (A10) antibodies were raised against the N-terminal domain, they give different results according to the authors. Caution should be taken, as it may imply that some previous results could be explained by epitope masking.

-I am sorry if I missed something crucial, but to my understanding, the disparities regarding the ChIPseq results obtained using the 8WG16 antibody are not fully resolved. In Figure S7C from their previous publication (Buchel et al, 2017) the authors concluded that "Intriguingly, ChIP sequencing showed that activation of N-MYC had no significant effect on chromatin association of hypo-phosphorylated Pol II". Is this not a similar experiment, using the same antibody and experimental conditions as in Figure 2 from the current manuscript? They now conclude that "activation of MYCN caused a global decrease in promoter association of non-phosphorylated RNAPII".

(3) Conducting ChIP-qPCR experiments for all nascent RNA degradation factors to be compared would have enabled a more direct and comprehensive comparison.

<https://doi.org/10.7554/eLife.94407.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public Review):

(1) In Figure 1, the authors show that TF3C binds to the amino terminus of MYCN (Myc box I region), as shown previously. The data in Figure 1 B-D support, but do not rigorously confirm a 'direct' interaction because it has not been ruled out that accessory proteins mediating the association may be present in the mixture.

In Figure 1B-D we have purified MYCN and the TFIIC/TauA complex separately and then mixed the purified preparations, demonstrating that the purified proteins interact. We have additionally performed mass spectrometry, which shows that the TauA/MYCN complex is formed without further accessory proteins, as the molecular weight would be higher. Based on the Coomassie stained SDS-PAGE gels, there is no plausible contaminating band in the purified complex that could be mediating the interaction between MYCN and TauA, either in the purified complex (Figure 1C), or in the purified protein used to reconstitute the complex (Figure S1A & S1B).

(2) The authors indicate in Figure 2 that TF3C has essentially no effect on MYCNdependent gene expression and/or transcription elongation. Yet a previous study (PMID: 29262328) associated with several of the same authors concluded that TF3C positively affects transcription elongation. The authors make no attempt to reconcile these disparate results and need to clarify this point.

We agree that the data in this manuscript do not support the role on transcription elongation. This point was also raised by Reviewer 3. Comparing our new results to the data published previously we can summarize that the data sets in the two studies show three key results: First, the traveling ratio of RNAPII changes upon induction of MYCN. Second, RNAPII decreases at the transcription start side and third, it increases towards the end side.

We agree that in the previous study we linked the traveling ratio directly to elongation. However performing ChIP-seq with different RNAPII antibodies showed us that for example RNAPII (N20), which is unfortunately discontinued, gives different results compared to RNAPII (A10). Combining our new results using the RNAPII (8WG16) antibody shows that the traveling ratio is not only reflecting transcription elongation but also includes that the RNAPII is kicked-off chromatin at the start side.

(3) Figures 2B and C show that unphosphorylated pol2 is TSS-centered, and Ser2-P pol2 occupation is centered beyond the TES. From this data, however, the reader can't tell how much of the phospho-Ser2- pol2 is centered on the TSS. The authors should include overall plots over TSS and TES, and also perhaps the gene-body to allow a better comparison for TSS and TES plotted for both antibodies over the collected gene sets.

We focused on the TSS for unphosphorylated RNAPII and the TES for pSer2-RNAPII, as these are the regions with specific enrichment of the respective antibodies. As requested for comparison, we now include metagenes showing TSS, gene-body, and TES for both antibodies as new Figure S2A and B. Additionally, we included density plots for unphosphorylated RNAPII at the TES as well as for pSer2-RNAPII at the TSS as a Figure for the Reviewers (Figure 1).

(4) The authors see more TF3C at promoters in cells with MYCN (Figure 2F). What are the levels of TF3C in the absence and presence of MYCN?

As shown in the immunoblot in Figure S1E, TF3C5 levels do not change upon induction of MYCN. We therefore think that MYCN helps to recruit TFIIC5 to RNAPII promoter sites. This is also in accordance to what we previously reported 1.

(5) The finding that TF3C is increased at TSS (Figure 2F) doesn't necessarily indicate that 1) MYCN is recruiting TF3C there, and 2) that this is due to the phosphorylation status of pol2. It could mean many other things. The logic of conflating these 3 points based on the data shown is questionable.

We showed previously that knock-down of MYCN affects TFIIC5 binding, showing that MYCN is required for binding of TFIIC5 at promoter sites 1.

Additionally, we included data with DRB treated cells (Figure 2F), which prevents RNAPII loading by preventing downstream de novo elongation. Those data show that TFIIC5 binding at the TSS is massively increased upon induction of MYCN and additionally upon treatment with DRB. Conversely, we observed that the major effect of TFIIC knock-down was at the nonphosphorylated RNAPII at the TSS on MYCN induction (Figure 2B). Therefore, we would argue that our assumption fits well to the data presented in the manuscript.

(6) Figure 3A doesn't add much to the paper, as it is overplotted and no relationship is clear, except that Pol2 and MYCN occupy many of the same sites. Perhaps a less complex or different type of plot would allow the interactions to be better visible.

We agree with the comment and since in another comment we were asked to show the same window for all shown Hi-ChIP data plots, we changed Figure 3A.

(7) That depletion of TF3C leads to increased promoter hubs may or may not have anything to do with its association with MYCN (Figure 4E). This could be a direct consequence of its known structural function in cohesin complexes, and the MYCN changes as a secondary consequence of this (also see point 4, above).

As shown in Büchel et al. (2017) 1 MYCN is needed to recruit RAD21 and depletion of RAD21 has no impact on the recruitment of MYCN. Since RAD21 is part of the cohesin complex we would exclude that the MYCN changes are a secondary consequence.

(8) Depletion of TF3C5 results in a loss of EXOSC5 (exosome) at TSS in the presence and absence of MYCN (Figure 5B). As TF3C5 is a cohesin, could this simply be a consequence

of genomic structure changes?

We agree that the discovered changes in EXOSC5 can be due to depletion of TFIIC5. TFIIC has been shown to recruit cohesin 1 and condensin complexes 2, as well as inducing chromatin architectural changes 3. However, MYCN is needed to recruit TFIIC and depletion of TFIIC had no impact on MYCN recruitment 1. Furthermore, MYCN has been shown to recruit exosome 4. Therefore, we would argue that either MYCN can directly play a role or thru chromatin architectural changes.

(9) The authors suggest that RNA dynamics are affected by changes in exosome function (RNA degradation, etc). What effect, if any does TF3C depletion have on the overall gene expression profile?

We show in the manuscript that TFIIC depletion in unperturbed cells has no effect on the global gene expression profile in the time frame analyzed (Figure 2E and S2B).

Reviewer #2 (Public Review):

(1) Dynamic inferences are made without kinetic experiments.

While we agree that we did not collect kinetic data to study the dynamics of RNA polymerase we would argue that the integration of our different data sets make it possible to draw conclusions about dynamic interferences. The transcription cycle and its sequential steps have been well described. In this sense, we use the non-phosphorylated RNAPII data that is situated between RNAPII recruitment and initiation and RNAPII-pSer2 that shows pause-release to elongation to draw conclusions on the dynamic. Likewise, we also made use of our previous published datasets.

Reviewer #2 (Recommendations For The Authors):

(1) A number of changes are reported in hub size, expression, etc. upon treatment with tamoxifen to activate MCN-ER. But MYC is already present in the SHEP cells, so why doesn't MYC support these same phenomena? It would seem that either the ability to cooperate with TFIIC to clear non-productive polymerase complexes from promoters is particular to MYCN, or else it reflects a quantitative increase in total MYC proteins due to the entry of MYCN-ER into the nucleus with tamoxifen. The authors should address or discuss this issue.

It could be that protein levels are the limiting factor between MYC and MYCN observed effects in this system. This interpretation would be in accordance with the results of Lorenzin et al. 5, which reported that different levels of MYC had different targets based on the affinity to Eboxes and protein level. A similar profile of MYC levels compared to function was also reported regarding SPT5 6. Those high protein levels mimic what is found in certain tumors in contrast to physiological levels. In this sense, the observed differences can also be between physiological and oncological levels of MYC proteins.

On the other hand, it has been described both a core MYC- and an isoform specific-signature of target genes. MYCN is described to be involved in gene expression during the S-phase of the cell cycle 7. This suggests that there are differences between MYC and MYCN other than gene sets. The interaction with TFIIC appears to be one of these differences. We have found multiple TFIIC subunits as part of the MYCN interactome, but the interaction of TFIIC with MYC is weaker and we are uncertain how relevant it is 7,8. We show here that depletion of different subunits of the TFIIC complex show a MYCN-dependent growth defect (Figure 1 E). Similarly, nuclear exosome is a MYCN-specific dependence 4, and we show here that MYCNdependent recruitment of the exosome requires TFIIC5. We take this as an indication

that there is an intrinsic difference between MYC and MYCN and that MYCN engages TFIIC for this pathway.

(2) Reciprocal to TFIIC recruitment to MYCN- rRNA, and other RNAPIII genes. Does this happen targets would be MYCN association with tRNA genes, 5S, and if so, is this association TFIIC dependent? What happens to the expression of these genes?

We did observe MYCN in interactions involving tRNA and other RNAPIII sites, such as SINE elements and tRNAs (Figure 4B, 4D, S3F, and S4B). There was no relevant number of 5S rRNA involved in interactions – either because the difficulty to properly map these repetitive regions or due to biology. In any case, none of those regions appeared to be specifically dependent on TFIIC as the overall number of interactions increased in TFIIC depletion regardless of the genomic annotation (Figure S4B). Regarding the expression of RNAPIII genes, we are constrained by technical limitations of poly(A) enrichment RNA-seq to globally analyze it in an unbiased way. However, we addressed this point for tRNAs expression in an earlier work 1 and found that tRNA levels do not change upon TFIIC depletion. We think this is because tRNAs are stable transcripts and RNAPIII recycling can occur in a TFIICindependent manner 9. Conversely, we reported no significant expression changes in RNAPII genes upon TFIIC depletion in this work.

(3) The authors show that TFIIC depletion does not alter the RNA-expression profile; how do they account for this? Can they comment on "background" transcription that it would seem should be suppressed by TFIIC-dependent removal of various hypofunctional polymerases?

Since TFIIC is important for the removal of non-functional RNAPII we would not expect changes to the gene expression profile upon depletion of TFIIC in the time frame analyzed. Monitoring the elongating form of RNAPII by measuring pSer2 indeed shows us that transcription elongation is not affected.

(4) Global changes in expression are difficult to assess with DESEQ2. This hypenormalizing algorithm is not really suited to distinguish differential, but universal upregulation from some targets being truly upregulated while others are downregulated. The authors should comment.

The authors acknowledge that DESEQ2 relies on the conjecture that genewise estimates of dispersion are generally unchanged among samples. We address this comment in two different ways. We include those in the Figure for the Reviewers (Figure 2). The first was to sequence samples deeper to avoid any bias created by random effect of lower coverage, the range of total reads increased from 6.8-9.3 to 16.5-20.7 million reads. The second was to compare the fold average bin dot plot for RNA-seq of SH-EP-MYCN-ER showing mRNA expression normalized by control per bin using the DESEQ2 (Figure 2A) normalization to TMM in edgeR (Figure 2B) and to quantile normalization (Figure 2C). No major differences were found from the original data or using the different methods, but we updated the Figure 2E in the manuscript to include the deeper sequencing dataset, we also adjusted it to show -/+ MYCN and transformed to log2 to make it more intuitive. Overall, it enhances our original understanding that gene expression remains largely unaffected by TFIIC5 knockdown.

(5) On page 7, the authors claim that MYCN-ER increased Ser-2 can reflect MYCN-stimulated transcription elongation. In fact, without kinetic studies, this is not fully supported. Accumulation of Ser-2 RNAPII along a gene can reflect increased initiation of full-speed RNAPs or a pile-up of RNAPs slowing down. This should be resolved or qualified.

While we agree that we did not collect kinetic data to study the dynamics of RNA polymerase we would argue that the integration of our different data sets make it possible to draw conclusions about dynamic interferences. We showed on the one side that pSer-2 accumulates on the TES and on the other side the induction of MYCN-ER up-regulates gene expression which proves productive transcription elongation.

(6) pLHiChIP needs to be better described, the Mumbach reference is not sufficient.

We have reformulated the pLHiChIP in the method section and hope that this will provide now a better description of the method.

(7) Can the authors recheck all the labels in Figure 2D-I believe there is an error involving + or - MYCN.

We carefully rechecked all the labels in Figure 2 and it was correct as it was. We understand the confusion that may have created comparing Figure 2D and Figure 2E. To avoid confusion, we updated Figure 2E to show the same direction of Figure 2D. We also log2 transformed the y-axis of Figure 2E to foster a more intuitive reading.

(8) Why are there different scales for the regions of chromosome 17 shown in Figures 3 and 4? It would be easier to compare if the examples were all shown at the same scale (about 2 MB is shown in another Figure).

We now show the same region of chromosome 17 in Figure 3 and 4.

Reviewer #3 (Public Review):

(1) The connection between the three major findings presented in this study regarding the role of TFIIC in the regulation of MYCN function remains unclear. Specifically, how the TFIICdependent restriction of MYCN localization to promoter hubs enhances the association of factors involved in nascent RNA degradation to prevent the accumulation of inactive RNA polymerase II at promoters is not apparent. As they are currently presented, these findings appear as independent observations. Cross-comparison of the different datasets obtained may provide some insight into addressing this question.

We previously observed that TFIIC does not affect MYCN recruitment, while MYCN affects TFIIC binding 1. Moreover, our group reported that MYCN recruits exosome 4 and BRCA1 to promoter-proximal regions 10 to clear out non-functional RNAPII. We are currently reporting that MYCN-TFIIC complexes exclude non-functional RNAPII. However, MYCN-active promoter hubs have more RNAPII and more transcription than MYCN-active promoter outside hubs. Furthermore, TFIIC binding occurs upstream of BRCA1 and exosome recruitments as depletion of TFIIC leads to recruitment decrease of both factors. Therefore, we argue that TFIIC is required for the proper function of those MYCN-active promoter hubs.

(2) Another concern involves the disparities in RNA polymerase II ChIP-seq results between this study and earlier ones conducted by the same group. In Figure 2, the authors demonstrate that activation of MYCN results in a reduction of non-phosphorylated RNA polymerase II across all expressed genes. This discovery contradicts prior findings obtained using the same methodology, where it was concluded that the expression of MYCN had no significant effect on the chromatin association of hypo-phosphorylated RNA polymerase II (Buchel et al, 2017). In this regard, the choice of the 8WG16 antibody raises concern, as fluctuations in the signal may be attributed to changes in the phosphorylation levels of the Cterminal domain. It remains unclear why the authors decided against using antibodies targeting the N-terminal domain of RNA

polymerase II, which are unaffected by phosphorylation and consistently demonstrated a significant signal reduction upon MYCN activation in their previous studies (Buchel et al, 2017) (Herold et al, 2019). Similarly, the authors previously proposed that depletion of TFIIC5 abrogates the MYCN-dependent increase of Ser2phosphorylated RNA polymerase II (Buchel et al, 2017), whereas they now show that it has no obvious impact. These aspects need clarification.

We politely disagree that our discoveries are contradicting each other. Comparing our new results to the data published previously we can summarize that the data sets in the two studies show three key results: First, the traveling ratio of RNAPII changes upon induction of MYCN. Second, RNAPII decreases at the transcription start side and third, it increases towards the end side.

We agree that in the previous study we linked the traveling ratio directly to elongation. However performing ChIP-seq with different RNAPII antibodies showed us that for example RNAPII (N20), which is unfortunately discontinued, gives different results compared to RNAPII (A10). Combining our new results using the RNAPII (8WG16) antibody shows that the traveling ratio is not only reflecting transcription elongation but also includes that the RNAPII is kicked-off chromatin at the start side.

In the previous study we only performed manual ChIP experiments for RNAPII (8WG16) and pSer2. Now we did a global analysis which is more meaningful and is also reflected in the RNA sequencing data.

(3) Finally, the varied techniques employed to explore the role of TFIIC in MYCNdependent recruitment of nascent RNA degradation factors make it challenging to draw definitive conclusions about which factor is affected and which one is not. While conducting ChIPseq experiments for all factors may be beyond the scope of this manuscript, incorporating proximity ligation assays (PLA) or ChIP-qPCR assays with each factor would have enabled a more direct and comprehensive comparison.

We understand the criticism that we are comparing different assays. We have performed PLAs with different antibodies. Since the controls of the PLAs were not sufficient for us, we refrain from using them. ChIP-qPCR experiments are much more challenging to do side by side compared to PLAs, which is why we decided against looking at all factors with this method.

Recommendations For The Authors:

Reviewer #3 (Recommendations For The Authors):

(1) Figure 2: Why did the authors choose the 8WG16 antibody? Does TFIIC5 depletion suppress the MYCN-dependent reduction of total RNA polymerase II binding to promoters that they consistently showed in previous studies? Given that phosphorylation of the CTD impacts 8WG16 recognition, including Ser5-phosphorylated RNA polymerase II ChIPseq experiments might clarify this issue.

We used the RNAPII (8WG16) antibody to exactly map non-phosphorylated RNAPII which shows us the binding of non-functional RNAPII.

(2) Figures 3 and 4: As it stands, the manuscript does not convincingly establish a functional connection between the results in Figures 2, 3, and 4 or elucidate potential mechanisms. Are changes in RNA polymerase II levels upon MYCN activation more pronounced at promoters located at MYCN hubs? Do changes in MYCN-enriched chromatin contacts upon TFIIC5 depletion somehow correlate with alterations in RNA

polymerase II levels? Performing similar cross-comparisons as in Figure 3C may help address this issue. Furthermore, it is not clear how the authors concluded that MYCN/TFIIC5-bound genes are not part of these so-called promoter hubs.

In Figure 3C we show that RNAPII levels are more pronounced upon MYCN activation at promoters located at MYCN hubs. Additionally, we show non-phosphorylated ChIP-seq on TSS and RNAPII-pSer2 ChIP-seq on TES density plots for promoters with MYCN interactions in the Figure for the Reviewers (Figure 3). We found no other difference than binding compared to the overall global analysis for all expressed genes showed in Figure 2B and Figure 2C. This goes on the same direction of the high expression observed of those genes in MYCN interactions observed in Figure 3C.

The changes observed in Figures 2B and 2C are global and do include the promoters with MYCN interactions. At the same time, it is required a higher number of replicates to statistically distinguish the MYCN interaction differences between TFIIC5 presence and depletion. We acknowledge this limitation, and we therefore restrain any attempt towards this end. We base our conclusions on the other parts of the manuscript and on our previous studies that show that MYCN recruits TFIIC, BRCA1, and the exosome to promoter proximal regions 1,4,10.

(3) Figure 5: According to the PLA results, activation of MYCN could enhance RNA polymerase II-NELFE interaction in a TFIIC5-dependent manner. Considering the raised issues regarding the use of the 8WG16 antibody, this result might be of relevance.

Nevertheless, PLA does not seem to be the optimal technique to address these questions, and I would rather suggest performing ChIP-qPCR experiments for all the factors to be compared. Finally, do the authors conclude that the TFIIC5 effect on MYCN-dependent changes in RNA polymerase II depends upon the recruitment of EXOSC5 and BRCA1? If so, it would be interesting to determine whether depletion of these factors phenocopies the effects observed with TFIIC5.

We understand the criticism that we are comparing different assays. We have performed PLAs with different antibodies. Since the controls of the PLAs were not sufficient for us, we refrain from using them.

(4) In Figure S2 the labels should be EtOH, 4-OHT, and Input.


We changed this accordingly.

(5) On page 7, the sentence "We have shown previously that TFIIC5 depletion does not cause significant changes in expression of multiple tRNA genes that are transcribed by RNAPIII (Buchel et al., 2017)" appears to lack a connection.

We agree with the reviewer and we deleted this sentence from the manuscript.


Author response image 1.

(A) Density plot of ChIP-Rx signal for non-phosphorylated RNAPII. Data show mean (line) $\pm$ standard error of the mean (SEM indicated by the shade) of different gene sets based on an RNA-seq of SH-EP-MYCN-ER cells $\pm$ 4-OHT. The y-axis shows the number of spike-in normalized reads and it is centered to the TES $\pm$ 2 kb. N = number of genes in the gene set defined in the methods. (B) Density plot of ChIP-Rx signal for RNAPII pSer2 as described for panel A. The signal is centered to the TSS $\pm$ 2 kb.




Author response image 2.

Bin dot plot for RNA-seq of SH-EP-MYCN-ER showing mRNA expression normalized by control per bin comparing the fold average using DESEQ2 (A), normalization to TMM in edgeR (B) and to quantile normalization (C).



Author response image 3.

Average density plot of ChIP-Rx signal for non-phosphorylated RNAPII (A) or RNAPII pSer2 (B) at promoters with MYCN interactions.



References

- (1) Büchel, G., Carstensen, A., Mak, K.-Y., Roeschert, I., Leen, E., Sumara, O., Hofstetter, J., Herold, S., Kalb, J., and Baluapuri, A. (2017). Association with Aurora-A controls NMYC-dependent promoter escape and pause release of RNA polymerase II during the cell cycle. *Cell reports* 21, 3483-3497.
- (2) Yuen, K.C., Slaughter, B.D., and Gerton, J.L. (2017). Condensin II is anchored by TFIIC and H3K4me3 in the mammalian genome and supports the expression of active dense gene clusters. *Sci Adv* 3, e1700191. [10.1126/sciadv.1700191](https://doi.org/10.1126/sciadv.1700191).
- (3) Ferrari, R., de Llobet Cucalon, L.I., Di Vona, C., Le Dilly, F., Vidal, E., Lioutas, A., Oliete, J.Q., Jochem, L., Cutts, E., Dieci, G., et al. (2020). TFIIC Binding to Alu Elements Controls Gene

Expression via Chromatin Looping and Histone Acetylation. *Mol Cell* 77, 475-487 e411. 10.1016/j.molcel.2019.10.020.

(4) Papadopoulos, D., Solvie, D., Baluapuri, A., Endres, T., Ha, S.A., Herold, S., Kalb, J., Giansanti, C., Schulein-Volk, C., Ade, C.P., et al. (2021). MYCN recruits the nuclear exosome complex to RNA polymerase II to prevent transcription-replication conflicts. *Mol Cell*. 10.1016/j.molcel.2021.11.002.

(5) Lorenzin, F., Benary, U., Baluapuri, A., Walz, S., Jung, L.A., von Eyss, B., Kisker, C., Wolf, J., Eilers, M., and Wolf, E. (2016). Different promoter affinities account for specificity in MYC-dependent gene regulation. *Elife* 5. 10.7554/eLife.15161.

(6) Baluapuri, A., Hofstetter, J., Dudvarski Stankovic, N., Endres, T., Bhandare, P., Vos, S.M., Adhikari, B., Schwarz, J.D., Narain, A., Vogt, M., et al. (2019). MYC Recruits SPT5 to RNA Polymerase II to Promote Processive Transcription Elongation. *Mol Cell* 74, 674-687 e611. 10.1016/j.molcel.2019.02.031.

(7) Baluapuri, A., Wolf, E., and Eilers, M. (2020). Target gene-independent functions of MYC oncoproteins. *Nat Rev Mol Cell Biol*. 10.1038/s41580-020-0215-2.

(8) Koch, H.B., Zhang, R., Verdoodt, B., Bailey, A., Zhang, C.D., Yates, J.R., 3rd, Menssen, A., and Hermeking, H. (2007). Large-scale identification of c-MYC-associated proteins using a combined TAP/MudPIT approach. *Cell Cycle* 6, 205-217. 10.4161/cc.6.2.3742.

(9) Ferrari, R., Rivetti, C., Acker, J., and Dieci, G. (2004). Distinct roles of transcription factors TFIIB and TFIIC in RNA polymerase III transcription reinitiation. *Proc Natl Acad Sci U S A* 101, 13442-13447. 10.1073/pnas.0403851101.

(10) Herold, S., Kalb, J., Büchel, G., Ade, C.P., Baluapuri, A., Xu, J., Koster, J., Solvie, D., Carstensen, A., and Klotz, C. (2019). Recruitment of BRCA1 limits MYCN-driven accumulation of stalled RNA polymerase. *Nature* 567, 545-549.

<https://doi.org/10.7554/eLife.94407.2.sa0>