

# RAS-p110 $\alpha$ signalling in macrophages is required for effective inflammatory response and resolution of inflammation

## Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

## Reviewed preprint version 1

April 12, 2024 (this version)

## Posted to preprint server

January 3, 2024

## Sent for peer review

December 12, 2023

**Alejandro Rosell, Agata A. Krygowska, Marta Alcón Pérez, Mathieu-Benoit Voisin, Juan de Paz, Héctor Sanz-Fraile, Vinothini Rajeeve, Alberto Berral-González, Cristina Cuesta, Ottilie Swinyard, Enrique Gabandé-Rodríguez, Julian Downward, Jordi Alcaraz, Javier De Las Rivas, Pedro Cutillas, Esther Castellano** 

Tumour-Stroma Signalling Lab., Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain • Centre for Cancer and Inflammation, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom • Centre for Microvascular Research, William Harvey Research Institute, Queen Mary University of London, Charterhouse Square, London, EC1M 6BQ, United Kingdom • Unit of Biophysics and Bioengineering, Department of Biomedicine, School of Medicine and Health Sciences, Universitat de Barcelona, Barcelona, Spain • Centre for Cancer Genomics and Computational Biology, Cell Signalling and Proteomics Laboratory, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom • Bioinformatics and Functional Genomics, Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain • Oncogene Biology Laboratory, Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK • Institute for Bioengineering of Catalonia (IBEC), The Barcelona Institute for Science and Technology (BIST), Barcelona 08028, Spain

 [https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access)

 Copyright information

## Abstract

Macrophages are crucial in the body's inflammatory response, with tightly regulated functions for optimal immune system performance. Our study reveals that the RAS-p110 $\alpha$  signalling pathway, known for its involvement in various biological processes and tumorigenesis, regulates two vital aspects of the inflammatory response in macrophages: the initial monocyte movement and later-stage lysosomal function. Disrupting this pathway, either in a mouse model or through drug intervention, hampers the inflammatory response, leading to delayed resolution and the development of more severe acute inflammatory reactions in live models. This discovery uncovers a previously unknown role of the p110 $\alpha$  isoform in immune regulation within macrophages, offering insight into the complex mechanisms governing their function during inflammation. With emerging potential to activate p110 $\alpha$  using small molecules, targeting the RAS-p110 $\alpha$  pathway could be a promising approach for treating chronic inflammation. This therapeutic prospect holds significant promise for easing inflammatory disorders and improving the quality of life for affected patients.

## eLife assessment

This **useful** study investigates the impact of disrupting the interaction of RAS with the PI3K subunit p110 $\alpha$  in macrophage function in vitro and inflammatory responses in vivo. **Solid** data overall supports a role for RAS-p110 $\alpha$  signalling in regulating macrophage activity and so inflammation, however for many of the readouts presented the magnitude of the phenotype is not particularly pronounced. Further analysis would be required to substantiate the claims that RAS-p110 $\alpha$  signalling plays a key role in macrophage function. Of note, the molecular mechanisms of how exactly p110 $\alpha$  regulating the functions in macrophage have not yet been established.

## Introduction

Phosphatidylinositol 3-kinases (PI3K) are a family of lipid kinases that phosphorylate phosphatidylinositides (PtdIns) at the 3'-hydroxyl group<sup>1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. Upon activation, PI3K phosphorylates phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP<sub>3</sub>). PIP<sub>3</sub> serves as a second messenger that recruits proteins containing pleckstrin homology (PH) domains, such as Akt (also known as protein kinase B)<sup>2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. This activation of PI3K regulate various cellular functions, including cell proliferation, growth, survival, motility, inflammation, and metabolism, among others<sup>1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. In macrophages, the activation of PI3K-Akt signalling is crucial to restrict inflammation and to promote anti-inflammatory responses in Toll Like Receptors (TLR)-induced macrophages, contributing to macrophage polarization<sup>5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>.

PI3Ks are heterodimeric lipid kinases composed of catalytic and adaptor/regulatory subunits that can be categorized into three classes based on their structures and substrate specificities<sup>3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. Class I catalytic isoforms, including p110 $\alpha$ , p110 $\beta$ , p110 $\gamma$ , and p110 $\delta$ , play essential roles in integrating signals from growth factors, cytokines, and other environmental cues. While p110 $\alpha$  and p110 $\beta$  are ubiquitously expressed, p110 $\delta$  and p110 $\gamma$  are largely restricted to the myeloid and lymphoid lineages<sup>7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. While p110 $\alpha$  is primarily associated with cell growth regulation and survival in epithelial cells, it must be considered that this isoform is also expressed in immune cells, including macrophages. The role played by p110 $\alpha$  in macrophages is not well understood, although some studies have suggested that it might regulate the survival and the regulation of the phagocytic activity of macrophages<sup>10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. In the context of cancer, impairment of RAS binding to p110 $\alpha$  in macrophages results in reduced recruitment of macrophages to the tumour site<sup>11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. Additionally, this disruption leads to a change in macrophage polarization, favouring a more proinflammatory M1 state<sup>12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. These findings suggest that p110 $\alpha$  plays a crucial role in regulating macrophage-dependent functions. However, despite these insights, the precise impact of p110 $\alpha$  on macrophage function and the underlying molecular mechanisms influencing the inflammatory response are not yet fully understood.

Inflammation is a complex and tightly regulated series of events triggered by various stimuli such as pathogens, harmful mechanical and chemical agents, and autoimmune reactions. The inflammatory response primarily occurs in vascularized connective tissues, involving a dynamic interplay of plasma components, circulating cells, blood vessels, and cellular and extracellular factors<sup>13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>. During inflammation, mediators released by recruited leukocytes orchestrate a response that aims to facilitate tissue repair and protect the body against harmful stimuli<sup>16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100</sup>.

Macrophages play a vital role in the inflammatory response by performing functions such as antigen presentation, phagocytosis, and immunomodulation<sup>16</sup>,<sup>17</sup>. Their role begins with the active recruitment of monocytes from the bloodstream to the site of infections<sup>19</sup>, where they differentiate into macrophages and recognize microbes and cellular debris through specific mechanisms. Macrophages subsequently actively participate in phagocytosis, a vital process involving the internalization and elimination of pathogens. Microbial destruction predominantly takes place within lysosomes and phagolysosomes<sup>20</sup>–<sup>23</sup>. At later stages of inflammation, macrophages contribute to the resolution of inflammation, thus preventing progression from acute to persistent inflammation that would cause additional tissue damage.

In this study we have used a combination of cell biology techniques and animal models to better understand the role of RAS-dependent activation of p110α at the different stages of the inflammatory response. Our findings show that RAS-p110α signalling plays a key role in the initial stages of inflammation, facilitating the extravasation of monocytes from the bloodstream by promoting the necessary cytoskeletal changes. Subsequently, RAS-p110α has a crucial role in lysosomal acidification and activation of cathepsins, which are indispensable for efficient degradation of lysosomal cargo; when these functions are impaired, prolonged acute inflammatory responses and delayed resolution steps are observed. These results significantly enhance our understanding of the complex mechanisms governing the immune response to inflammation, emphasizing the pivotal role played by RAS-p110α signalling in orchestrating proper monocyte extravasation and maintaining optimal lysosomal function.

## Results

### Disruption of RAS-p110α causes prolonged and more acute responses to inflammatory stress *in vivo*

Previous data suggested that, in a tumoral setting, somatic disruption of RAS-p110α prevents macrophage recruitment to tumours<sup>11</sup>,<sup>12</sup> and favours polarization of macrophages to a proinflammatory phenotype<sup>12</sup>, suggesting a possible role for p110α in macrophage function. To determine whether RAS-dependent activation of p110α participates in innate or adaptive immune responses to inflammation in macrophages, we used an established mouse model designed for the tamoxifen-inducible disruption of RAS binding to p110α<sup>11</sup>,<sup>24</sup> (**Supplementary Fig. S1A**). This mouse model introduces two-point mutations, T208D and K227A, in the RAS Binding Domain (RBD) of the endogenous *Pik3ca* allele (*Pik3ca*<sup>RBD</sup>), enabling the selective disruption of the RAS-p110α interaction<sup>24</sup>. Wild-type (*Pik3ca*<sup>WT</sup>) and *Pik3ca*<sup>RBD</sup> mice were bred with mice containing a floxed *Pik3ca* allele<sup>25</sup>, along with a strain containing a conditional Cre recombinase (Cre-ERT2) allele targeted to the ubiquitously expressed Rosa26 locus. The resulting *Pik3ca*<sup>WT/Flox</sup> and *Pik3ca*<sup>RBD/Flox</sup> mice displayed no discernible phenotype and exhibited behavior consistent with *Pik3ca*<sup>WT</sup> mice<sup>11</sup>,<sup>24</sup>. Activation of Cre-recombinase by tamoxifen led to the excision of the floxed *Pik3ca* allele, resulting in mice expressing either one *Pik3ca*<sup>WT</sup> allele (*Pik3ca*<sup>WT/-</sup>) or one *Pik3ca*<sup>RBD</sup> allele (*Pik3ca*<sup>RBD/-</sup>)<sup>11</sup>. This inducible genetic manipulation strategy allows us to selectively disrupt the RAS-p110α interaction in a controlled and temporally regulated manner, providing a valuable tool for dissecting the contributions of this pathway to immune responses in the context of inflammation.

Bone marrow derived macrophages (BMDMs) were generated from tibias and femurs of *Pik3ca*<sup>WT/Flox</sup> and *Pik3ca*<sup>RBD/Flox</sup> mice, ensuring efficient removal of the floxed allele (**Supplementary Fig. S1B**). Subsequently, these BMDMs were induced towards a proinflammatory state through stimulation with LPS and IFN-γ.


To assess the impact of RAS-p110 $\alpha$  binding deficiency on inflammatory intracellular signalling, we first examined Akt activation. The results revealed a notable decrease in Akt activation levels under inflammatory conditions in BMDMs lacking RAS-p110 $\alpha$  binding, while no discernible change was observed in ERK activation, another well-known RAS effector (**Fig. 1A**). Interestingly, the decrease in Akt activation was accompanied by a decrease in the activation of NF- $\kappa$ B (**Fig. 1B**), a pivotal transcription factor that regulates the expression of pro-inflammatory genes<sup>26</sup>. This suggests a specific impairment in the Akt signaling pathway in the absence of RAS-p110 $\alpha$  interaction during proinflammatory stimulation.

Consequently, our next objective was to investigate whether in vivo disruption of RAS-p110 $\alpha$  might lead to altered inflammatory responses. To address this, we administered tamoxifen to 10–12-week-old *Pik3ca*<sup>WT/flox</sup> and *Pik3ca*<sup>RBD/flox</sup> mice with tamoxifen and, after a two-week interval, conducted a paw swelling assay. In this assay, zymosan (10  $\mu$ g/ $\mu$ l) or PBS was injected into the hind paws of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> mice with paw thickness measured at regular intervals over a 5-day period. The results revealed a significant increase in paw inflammation in *Pik3ca*<sup>RBD/-</sup> mice compared to *Pik3ca*<sup>WT/-</sup> mice, evident from the earlier time points measured and persisting throughout the experiment (**Fig. 1C**). Paws from the *Pik3ca*<sup>WT/flox</sup> and *Pik3ca*<sup>RBD/flox</sup> mice, which had not received tamoxifen, exhibited comparable levels of inflammation (**Supplementary Fig. S1C**). This reaffirms that the absence of p110 $\alpha$  triggers a significant alteration in the inflammatory response and confirms that *Pik3ca*<sup>WT/flox</sup> and *Pik3ca*<sup>RBD/flox</sup> mice do not show any phenotype, as previously described<sup>11, 24</sup>. Additionally, analysis of the blood sedimentation rate, a reliable indicator of systemic inflammation levels<sup>27</sup>, showed a lower basal sedimentation ratio in *Pik3ca*<sup>RBD/-</sup> mice under PBS-treated conditions (**Supplementary Fig. S1D**). Following zymosan injection, both genotypes exhibited an increase in sedimentation rate, but the rise was more pronounced in *Pik3ca*<sup>RBD/-</sup> mice, indicating a heightened systemic inflammatory response. These findings underscore that disruption of RAS-p110 $\alpha$  interaction results in an exacerbated inflammatory state, reflected in both localized paw inflammation and systemic inflammatory mediator levels.

To delve deeper into the inflammatory response induced by zymosan, we collected paw samples at two- and five-days post-injection and conducted Haematoxylin & Eosin (H&E) studies. This approach enabled a comprehensive analysis, allowing us not only to scrutinize the early stages of inflammation but also to monitor the subsequent resolution phase of the inflammatory process. Paws obtained from mice injected with zymosan for 2 days displayed extensive areas of damaged connective tissue in both *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> paws (**Fig 1D and E**). Notably, the inflamed region in *Pik3ca*<sup>RBD/-</sup> mice was larger compared to control samples (**Fig. 1D and E**). After 5 days of zymosan injection, there was a significant reduction in the inflamed area, although it remained comparatively larger in the paws from *Pik3ca*<sup>RBD/-</sup> mice (**Fig. 1D and E**). This observation suggests a prolonged and heightened inflammatory response in mice lacking RAS-p110 $\alpha$  interaction, emphasizing the role of this interaction in the regulation of inflammatory processes.

The cellular composition within the inflammatory abscess offers crucial insights into the severity and progression of the disease. Our observations revealed features indicative of an acute inflammatory response, including an inflammatory abscess with elevated numbers of polymorphonuclear cells (**Fig. 1F**, blue arrows), primarily neutrophils, and macrophages (**Fig. 1F**, green arrows) displaying altered cell shape and increased cell death, accompanied by fibrin deposition (**Fig. 1F**, yellow arrows). Notably, *Pik3ca*<sup>RBD/-</sup> mice exhibited lower numbers of macrophages, a larger necrotic area with increased chromatin remnants, and reduced fibrin content compared to the paws from *Pik3ca*<sup>WT/-</sup> mice. By day 5, paws from *Pik3ca*<sup>WT/-</sup> mice showed a significant increase in the number of macrophages (**Fig. 1F**, green arrows), a decrease in polymorphonuclear cells, and a nearly complete resolution of the necrotic area, indicating the initiation of the resolution phase. Moreover, there were abundant activated fibroblasts (**Fig. 1F**, orange arrows), suggesting active production of new connective tissue. In contrast, paws from *Pik3ca*<sup>RBD/-</sup> mice exhibited a delayed healing process characterized by a larger central area of





**Figure 1.**

### ***Pik3ca*<sup>RBD/-</sup> mice have impaired responses to inflammatory insults.**

**(A)** Western blotting showing activation of Akt and ERK in *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs activated towards a pro-inflammatory state with LPS and IFN- $\gamma$  at the indicated time points; **(B)** Western blotting showing activation of NF- $\kappa$ B (p65) in *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs activated towards a pro-inflammatory state with LPS and IFN- $\gamma$  at the indicated time points; **(C)** *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> mice were injected with zymosan or PBS in the back-hind paws and inflammation was measured and plotted over time, *Pik3ca*<sup>WT/-</sup> PBS n = 6; *Pik3ca*<sup>RBD/-</sup> PBS n = 5; *Pik3ca*<sup>WT/-</sup> Zymosan n = 4; *Pik3ca*<sup>RBD/-</sup> Zymosan n = 4; Error bars indicate mean  $\pm$  SEM. Significance using 2-way ANOVA test: \*\*p < 0.01 **(D and E)** Representative H&E images of the inflamed area of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> paws injected with zymosan at the indicated times; **(F)** Representative images of cellularity present in the inflamed abscess of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> paws injected with zymosan at the indicated times. Blue arrows indicate apoptotic polymorphonuclear cells, yellow arrows indicate fibrin deposition, green arrows indicate macrophages and orange arrows indicate activated fibroblasts; **(G)** Representative images of macrophages (CD68 positive cells) present in the inflamed abscess of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> paws injected with zymosan; **(H)** Quantification of macrophages (CD68 positive cells) present in the inflamed abscess of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> paws injected with zymosan; **(I)** Representative H&E images of the inflamed area and cellularity of *Pik3ca*<sup>WT/WT</sup> paws and *Pik3ca*<sup>WT/WT</sup> paws treated with BYL-719 injected with zymosan. Blue arrows indicate apoptotic polymorphonuclear cells, yellow arrows indicate fibrin deposition, green arrows indicate macrophages; **(J)** Representative images and quantification of macrophages (CD68 positive cells) present in the inflamed abscess of *Pik3ca*<sup>WT/WT</sup> paws and *Pik3ca*<sup>WT/WT</sup> paws treated with BYL-719 injected with zymosan; **(K)** Quantification of macrophages (CD68 positive cells) present in the inflamed abscess of *Pik3ca*<sup>WT/WT</sup> paws and *Pik3ca*<sup>WT/WT</sup> paws treated with BYL-719 injected with zymosan. Error bars indicate mean  $\pm$  SEM. Significance using Student's t test: \*\*p < 0.01.

polymorphonuclear cells, abundant fibrin deposits, reduced numbers of infiltrating macrophages, and limited fibroblast activity (**Fig. 1F**). These findings collectively indicate an imbalance in the inflammatory response and a slower progression towards resolution in the absence of RAS-p110 $\alpha$  interaction, emphasizing the pivotal role of this interaction in orchestrating an effective and timely resolution of inflammation.

To evaluate the presence of macrophages within the inflammatory lesion, we performed specific immunohistochemical (IHC) analysis using the macrophage-specific marker CD68 in paws from day 5, where more cellular preservation and initiation of the healing process had been observed. A notable decrease in the number of CD68-positive cells was observed in the inflamed abscess region of *Pik3ca*<sup>RBD/-</sup> mice (**Fig. 1G and H**).


To further confirm the involvement of p110 $\alpha$  signalling in the acute inflammatory response, *Pik3ca*<sup>WT/WT</sup> mice were subjected to daily treatment with BYL719 (Alpelisib), a specific inhibitor of p110 $\alpha$  isoform. After an initial 48-hour treatment period, the mice received injections of zymosan or PBS into their back-hind paws and were sacrificed 2 days later for analysis. Similar to what we had observed in *Pik3ca*<sup>RBD/-</sup> mice, the inflamed area in BYL719-treated mice exhibited a larger extension than the inflamed area from non-treated mice (**Fig. 1I** and **Supplementary Fig. S1E**). The central necrotic region was significantly larger in the BYL719-treated mice, and it contained large amount of apoptotic polymorphonuclear cells (**Fig. 1I**, blue arrows) and lower number of macrophages (**Fig. 1I**, green arrows), resembling the observations from *Pik3ca*<sup>RBD/-</sup> mice. Immunostaining for CD68 revealed a reduction in the number of macrophages present in the inflammatory abscess upon inhibition of p110 $\alpha$  signalling with BYL719 treatment (**Fig. 1J and K**).

In summary, our findings highlight that both genetic disruption of RAS-p110 $\alpha$  interaction and pharmacological inhibition of p110 $\alpha$  contribute to an expanded inflamed area and central necrotic region, concurrently reducing macrophage infiltration in a zymosan-induced inflammation model. These results collectively underscore the pivotal role of the p110 $\alpha$  isoform of PI3K in orchestrating the resolution of inflammatory responses, emphasizing its significance in modulating the dynamics and outcomes of inflammatory processes.

## Disruption of RAS binding to p110 $\alpha$ impairs the number of inflammatory monocytes in blood and spleen

Given the decrease in macrophages observed in the inflammatory abscess, we next set out to determine whether disruption of RAS binding to p110 $\alpha$  had an effect on the number of monocytes circulating in the blood of adult mice. *Pik3ca*<sup>WT/flox</sup> and *Pik3ca*<sup>RBD/flox</sup> mice were treated with tamoxifen at 12–14 weeks of age and 4 weeks later, blood was collected by cardiac puncture and immune populations were analysed by flow cytometry. We found a decrease in the number of circulating classical (or inflammatory) monocytes (Ly6C<sup>Hi</sup>/Ly6G<sup>-</sup>/CD11b<sup>+</sup> cells) (**Fig. 2A**) in *Pik3ca*<sup>RBD/-</sup> mice and no changes were observed in non-classical (or non-inflammatory) monocytes (Ly6C<sup>Lo</sup>/Ly6G<sup>-</sup>/CD11b<sup>+</sup> cells) (**Fig. 2B**). Together with the decrease in inflammatory monocytes, we observed an increase in the number of neutrophils (Ly6C<sup>-</sup>/Ly6G<sup>+</sup>/CD11b<sup>+</sup> cells) (**Supplementary Fig. S2A**). We did not detect differences in the numbers of T-cells (CD3<sup>+</sup>, CD8<sup>+</sup> or CD4<sup>+</sup>) (**Supplementary Fig. S2B**) or B-cells (CD19<sup>+</sup>) (**Supplementary Fig. S2C**) in the blood after the disruption of RAS binding to p110 $\alpha$ . We also analysed these same cell populations in *Pik3ca*<sup>WT/flox</sup> and *Pik3ca*<sup>RBD/flox</sup> mice and no differences were found (data not shown).

Since splenic monocytes resemble their blood counterparts<sup>28</sup>, we next aimed at determining whether splenic monocyte population would also be altered after disruption of RAS p110 $\alpha$  interaction. As observed in circulating blood, the number of classical monocytes (Ly6C<sup>Hi</sup>/Ly6G<sup>-</sup>/CD11b<sup>+</sup>) in the spleen decreased after disruption of RAS-p110 $\alpha$  interaction (**Fig. 2C**) and no differences were found in non-classical activated monocytes (Ly6C<sup>Lo</sup>/Ly6G<sup>-</sup>/CD11b<sup>+</sup>)



**Figure 2.**

### Disruption of RAS-p110 $\alpha$ interaction decreases the number of inflammatory monocytes in blood and spleen.

Twelve-week-old mice were treated with tamoxifen and after four weeks, flow cytometry analysis was performed to determine (A) inflammatory monocytes in circulating blood; (B) alternatively activated monocytes in circulating blood; (C) inflammatory monocytes in spleen; (D) classically activated monocytes in spleen; (E) macrophages in spleen; (F) macrophages in spleen's white pulp; (G) macrophages in spleen's red pulp. Data are presented as percentage of positive cells for the indicated markers. Black dots represent data from *Pik3ca*<sup>WT</sup> mice and red dots represent data from *Pik3ca*<sup>RBD</sup> mice. Each dot represents an individual mouse. Error bars indicate mean  $\pm$  SEM. Significance using Student's t test: \* p < 0.05; \*\*p < 0.01.

(Fig. 2D). We also checked the levels of resident macrophages present in the spleen and results showed that spleens from the *Pik3ca*<sup>RBD/-</sup> mice had a decrease in the number of differentiated macrophages (F4/80<sup>+</sup>/CD11b<sup>+</sup>/CD45<sup>+</sup>) (Fig. 2E). Analysis of macrophages in the white and red pulp of the spleen indicated that *Pik3ca*<sup>RBD/-</sup> mice had a significant decrease in the macrophage population in the former region (Fig. 2F and 2G). No differences were found in the number of granulocytes (Supplementary Fig. S2D), B-cells (Supplementary Fig. S2E) or T-cells (Supplementary Fig. S2F) present in the spleen between *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> mice. There were no differences in spleen size from *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> mice either (Supplementary Fig. S2G).


Given the decrease in the number of inflammatory monocytes and macrophages observed in blood and spleens of *Pik3ca*<sup>RBD/-</sup> mice, we wondered if disruption of RAS activation of p110α could lead to a decrease in myeloid bone marrow precursors. Myeloid lineage descends from a common myeloid progenitor (CMP) in bone marrow and traverse into blood as mature cells<sup>29</sup>. CMP differentiates into the Granulocyte-Macrophage Progenitor (GMP) and the Megakaryocyte-Erythroid Progenitor (MEP). The CMP can generate all types of myeloid colonies, whereas the GMP or the MEP produces only granulocyte macrophage (GM) or megakaryocyte erythrocyte (MegE) lineage cells, respectively (Supplementary Fig. S2H). No significant differences were noted in the quantity of CMP progenitors within the bone marrow *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> mice (Supplementary Fig. S2I-L). This observation suggests that the variations observed in blood and spleen parameters are not attributable to impairments in progenitor fate.

Finally, bone marrow precursors from *Pik3ca*<sup>RBD/flox</sup> and *Pik3ca*<sup>WT/flox</sup> mice were differentiated into macrophages *in vitro*, and the number of bone marrow-derived macrophages (BMDM) was determined by flow cytometry analysis. No differences were found in the number of BMDMs obtained from *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> mice (Supplementary Fig. S2M), suggesting that the disruption of RAS-p110α signalling do not interfere with the ability of bone marrow precursors to differentiate to macrophages.

## Disruption of RAS binding to p110α impairs monocyte transendothelial extravasation in response to inflammatory cues

The decrease in classical monocytes observed in the inflammatory abscess of paws from *Pik3ca*<sup>RBD/-</sup> mice may indicate a decreased ability to mount an effective immune response. We carried out trans-well assays since they are widely used to quantify transendothelial migration. Fibroblasts were seeded in the lower chamber to provide a continuous supply of Ccl2<sup>30</sup>, as this cytokine is well known for its ability to drive chemotaxis of myeloid cells under inflammatory conditions<sup>31</sup>. *Pik3ca*<sup>RBD/-</sup> or *Pik3ca*<sup>WT/-</sup> BMDMs were seeded in the upper chamber of the transwell. Additionally, we stimulated *Pik3ca*<sup>RBD/-</sup> or *Pik3ca*<sup>WT/-</sup> BMDMs with lipopolysaccharide (LPS) and Interferon gamma (IFN-γ) to mimic/recapitulate the proinflammatory phenotype typically associated with bacterial infection that causes macrophage activation<sup>32</sup>. Results showed that in both unstimulated and LPS+IFN-γ stimulated BMDMs, a lower number of *Pik3ca*<sup>RBD/-</sup> BMDMs were able to go through the transwell pore membranes compared to *Pik3ca*<sup>WT/-</sup> BMDMs (Fig. 3A). As expected, when BMDMs were stimulated towards a proinflammatory phenotype, a decrease in their migratory ability is observed<sup>33</sup>.

Extravasation entails the migration of monocytes through the endothelium<sup>34, 35</sup>, so we conducted random migration assays of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs growing in matrigel coated plates under unstimulated conditions or activated towards an inflammatory phenotype by addition of LPS and IFN-γ to the media. Analysis of the data revealed that, under pro-inflammatory conditions, *Pik3ca*<sup>RBD/-</sup> BMDMs displayed a decrease in migration speed (Fig. 3B). Migration was also evaluated in *Pik3ca*<sup>WT/WT</sup> BMDMs treated with BYL-719. Results confirmed a



**Figure 3.**

### Disruption of RAS-p110 $\alpha$ interaction impairs transendothelial extravasation to sites of inflammation.

(A) Graph indicating the quantification of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs passing through 8  $\mu$ m membrane pore transwells for 24 hours. Membrane attached macrophages on the lower part of the transwell were stained with crystal violet and quantified. Each dot represents an independent experiment. Error bars indicate mean  $\pm$  SEM; (B) Random migration of *Pik3ca*<sup>WT/-</sup>, *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/WT</sup> BMDMs treated with BYL719 (500 ng/ml) was analysed by time-lapse video microscopy and cell tracing in the presence or absence of LPS (100 ng/ml) and IFN- $\gamma$  (20 ng/ml); (C) Schematic representation of myeloid chimera generation strategy; (D) Graph showing level of bone marrow reconstitution with *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> myeloid lineage; (E) Graph quantifying the number of neutrophils in the blood of chimera mice treated with anti-GR1 (25 $\mu$ g/mouse/day); (F-H) Intra vital imaging quantification of the number of: (F) rolling monocytes per 5 minutes; (G) adherent monocytes per 500  $\mu$ m of vessel segment; (H) extravasated monocytes per mm<sup>2</sup> of tissue. Each dot represents an individual mouse. (I) Chimera *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> mice were injected with zymosan or PBS in the back-hind paws and inflammation was measured and plotted over time, *Pik3ca*<sup>WT/-</sup> PBS n = 5; *Pik3ca*<sup>RBD/-</sup> PBS n = 5; *Pik3ca*<sup>WT/-</sup> Zymosan n = 5; *Pik3ca*<sup>RBD/-</sup> Zymosan n = 5.

Statistical significance was obtained using Mann-Whitney test (A-H) or 2-way ANOVA test (I): n.s. non-significant; \*p < 0.05; \*\*p < 0.01; \*\*\*p < 0.001; \*\*\*\*p < 0.0001.



decrease in migration speed of macrophages upon treatment with BYL-719 (**Fig. 3B**) indicating that inhibition of p110 $\alpha$ , either genetically or pharmacologically, reduces the ability of macrophages to migrate under inflammatory conditions.

To determine if disruption of RAS binding to p110 $\alpha$  impairs monocyte ability to extravasate through the endothelium *in vivo* in response to an inflammatory stress, we analysed monocyte extravasation through the mesenteric vein in response to intraperitoneal Ccl2 injection. It is well established that loss of p110 $\alpha$  function leads to significant impairment in endothelial and lymphatic system<sup>12, 36</sup>, so in order to determine if monocytes from *Pik3ca*<sup>RBD/-</sup> mice presented an alteration in extravasation, we generated chimeric mice in which only the bone marrow were defective in RAS binding to p110 $\alpha$  (**Fig. 3C**). For this, bone marrow from *Pik3ca*<sup>RBD/-</sup> or *Pik3ca*<sup>WT/-</sup> mice was injected through the tail vein of irradiated LysM-GFP donor mice<sup>37</sup>. Engraftment of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> bone marrow could be followed by disappearance of the eGFP signal in donor mice (**Fig. 3D** and **Supplementary Fig. S3A**). Neutrophils (but no monocytes) were depleted using an anti-GR1 antibody (**Fig. 3E** and **Supplementary Fig. S3B**) to avoid interference with monocyte extravasation. Intraperitoneal injection of Ccl2 was performed in the peritoneum of the chimera mice to induce extravasation of monocytes through the mesenteric vein and rolling, adhesion and extravasation was measured by intravital microscopy. Flow cytometry analysis demonstrated no differences in Ccr2 expression between *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> monocytes (**Supplementary Fig. S3C**). We observed that blocking RAS-p110 $\alpha$  interaction did not induce a decrease in the number of rolling monocytes (**Fig. 3F**) or in the number of monocytes that adhere to the endothelium (**Fig. 3G**). However, when extravasation was measured, data showed that disruption of RAS binding to p110 $\alpha$  caused a significant decrease in the number of monocytes that were capable of extravasating through the endothelium of the mesenteric vein (**Fig. 3H**).


Additionally, in order to determine whether the differences observed in the inflammatory response of *Pik3ca*<sup>RBD/-</sup> mice were due to lack of monocyte extravasation, we repeated the zymosan paw swelling assay in the chimera mice. As previously, we observed that disruption of RAS-p110 $\alpha$  only in the immune system led to higher inflammatory response and a delay in the initiation of the resolution phase (**Fig. 3I**).

## Disruption of RAS-p110 $\alpha$ activation in macrophages induces changes in cytoskeleton reorganization

During transendothelial migration, leukocytes undergo cytoskeletal rearrangements that allow them to squeeze through the tight spaces between endothelial cells and enter the underlying tissue<sup>38, 39</sup>. Therefore, we next sought to determine whether the differences observed in *Pik3ca*<sup>RBD/-</sup> BMDMs extravasation and migration were attributable to altered cytoskeletal dynamics. To do so, we first aimed at examining cell shape and spread in *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs in pro-inflammatory conditions after LPS+IFN- $\gamma$  treatment. Treatment with LPS+IFN $\gamma$  induced an increase in cell spread in both *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs when compared to their respective unstimulated counterparts (**Fig. 4A and 4B**). However, cell spread in LPS+IFN $\gamma$  activated *Pik3ca*<sup>RBD/-</sup> BMDMs was significantly decreased compared to that observed *Pik3ca*<sup>WT/-</sup> BMDMs (**Fig. 4A and 4B**). Additionally, *Pik3ca*<sup>RBD/-</sup> BMDMs were more elongated and did not acquire the typical rounded shape known to be induced in macrophages after treatment with LPS+IFN- $\gamma$ <sup>40</sup> (**Fig. 4A and 4C**). We also analysed cell height and results showed that *Pik3ca*<sup>RBD/-</sup> BMDMs are higher both in unstimulated conditions and after LPS+IFN- $\gamma$  stimulation (**Fig. 3D**).

Actin is a bona fide regulator of cell shape<sup>41</sup>, so we next aimed at exploring the actin cytoskeleton in *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs. Actin fractionation assays were carried out in unstimulated or LPS+IFN $\gamma$  activated *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs. Our results revealed that disruption of RAS-p110 $\alpha$  binding caused a decrease in the F-actin pool in unstimulated BMDMs





**Figure 4.**

### Disruption of RAS-p110 $\alpha$ signalling impairs BMDM ability to remodel their cytoskeleton in response to inflammatory cues.

(A) Representative images of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs unstimulated or activated with LPS+ IFN- $\gamma$ ; (B) violin plot quantifying spread area. IF images of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs co-stained with phalloidin and DAPI were used to analyse spread area. Results are represented as a violin plot. Three independent biological replicates were analysed ( $n \geq 250$  total cells); (C) Quantification of cell circularity of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs unstimulated or activated with LPS+ IFN- $\gamma$ . Quantification was performed using the same images from panel (A); (D) Representative 3D projections of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs unstimulated or activated with LPS+IFN- $\gamma$  and violin plot showing quantification of the corresponding cell height. 3D projections and cell height were analysed using same images used in panel (A). (E) Violin plot representing F-actin pool in *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs unstimulated or activated with LPS+ IFN- $\gamma$ . Three independent biological replicates were analysed ( $n \geq 250$  total cells); (F) Violin plot representing G-actin pool (measured by DNase staining) in *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs unstimulated or activated with LPS+ IFN- $\gamma$ . Three independent biological replicates were analysed ( $n > 250$  total cells); (G) *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs were activated with LPS+ IFN- $\gamma$  and western blots were performed to determine expression levels of the indicated proteins.

Statistical significance was obtained using Mann-Whitney test: n.s. non-significant; \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ ; \*\*\*\* $p < 0.0001$ .

(Fig. 4E [↗](#)), with no changes observed in the G-actin pool when compared to matched controls (Fig. 4F [↗](#)). Analysis of actin dynamics after activation with LPS+IFN $\gamma$  showed that, in proinflammatory conditions, the F-actin pool is increased, as expected<sup>42</sup> (Fig. 4E [↗](#)). However, actin polymerization was significantly active in *Pik3ca*<sup>RBD/-</sup> BMDMs, leading to a striking increase in F-actin when compared to that observed in controls (Fig. 4E [↗](#)). In parallel, a decrease in the G-actin pool in *Pik3ca*<sup>RBD/-</sup> BMDMs was observed, suggesting an increase in the stabilization of F-actin after disruption of RAS binding to p110 $\alpha$  (Fig. 4F [↗](#)).

We performed Western blot analysis to assess the expression and activation of proteins involved in actin polymerization dynamics. Interestingly, we found elevated levels of serine 3 phosphorylated Cofilin in *Pik3ca*<sup>RBD/-</sup> BMDMs both in unstimulated and in proinflammatory conditions (Fig. 4G [↗](#)). Phosphorylation of Cofilin at Ser3 hampers its ability to depolymerize F-actin, resulting in the stabilization of actin and affecting actin filament dynamics. The increase in phosphorylated Cofilin, thus, would explain the increase in actin stabilization observed in *Pik3ca*<sup>RBD/-</sup> BMDMs. Additionally, we also observed an upregulation in the phosphorylation of LIM domain kinase (LIMK), a known kinase responsible for phosphorylating Cofilin at Ser3 (Fig. 4G [↗](#)), suggesting that the heightened Ser3 phosphorylation of Cofilin may be LIMK-mediated.

We next sought to investigate whether the enhanced F-actin stabilization corresponded to increased cellular rigidity. Consequently, we measured the elastic modulus of *Pik3ca*<sup>RBD/-</sup> BMDMs, revealing a significant increase in cell stiffness following RAS-p110 $\alpha$  disruption under both basal and proinflammatory conditions. To validate that the heightened cell stiffness was attributed to p110 $\alpha$ , we treated *Pik3ca*<sup>WT/WT</sup> BMDMs with BYL-719 and observed an amplified elastic modulus in these cells when stimulated with LPS+IFN $\gamma$ , confirming that the inhibition of p110 $\alpha$  indeed results in augmented cellular rigidity.

In summary, our findings suggest that the impeded extravasation of RAS-p110 $\alpha$  deficient monocytes in response to inflammatory cues is associated with disturbances in cytoskeletal dynamics, resulting in heightened cellular stiffness and diminished deformability.

## Disruption of RAS-p110 $\alpha$ activation impacts the secretome of macrophages

We next wondered whether the absence of RAS binding to p110 $\alpha$  affects the ability of BMDMs to acquire a pro-inflammatory state characterized by increased expression of markers such as CD80, CD86, and MHCII<sup>43, 44</sup>. To assess this, *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs were stimulated with LPS+IFN $\gamma$ , followed by flow cytometry analysis. Expression levels of CD80 (Supplementary Fig. S4A [↗](#)), CD86 (Supplementary Fig. S4B [↗](#)), and MCHII (Supplementary Fig. S4C [↗](#)) were examined. No significant differences were observed in the expression levels of any of these markers between the two genotypes under study.

During the inflammatory response, macrophages release a diverse array of bioactive molecules that have a profound impact on immune regulation, tissue repair, and disease progression. Thus, we analysed the secretome of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs both under steady state condition and during phagocytosis. We utilized apoptotic LKR10 cells, a murine lung cancer cell line, as the substrate in our phagocytosis assay. LKR10 cells were exposed to cisplatin for 16 h and apoptosis was confirmed by Western Blot (Supplementary Fig. S4D [↗](#)). Our goal was to create a more physiologically relevant experimental setting that more closely mimics the complex nature of the inflammatory response. For the secretome analysis, macrophages were incubated with or without apoptotic cells for 16 h. Culture supernatants were collected, clarified, and subjected to label-free quantitative proteomics analysis.

A total of 127 peptides corresponding to 105 proteins (**Supplementary Table 1** [↗](#)) showed differential expression between *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDM secretomes at steady state conditions. Additionally, 359 peptides corresponding to 210 proteins (**Supplementary Table 2** [↗](#)) were present at significantly different levels in the secretomes of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDM during phagocytosis of apoptotic cells. Next, we compared these peptides with the list of secreted proteins available at The Human Protein Atlas, and removed those that do not correspond to secreted proteins. After this step, 18 proteins were found to be differentially secreted by *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDM in steady state conditions (**Fig. 5A** [↗](#)), and 38 by *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDM during phagocytosis (**Fig. 5B** [↗](#) and **Supplementary Fig. S4E and S4F** [↗](#)). Surprisingly, most proteins were secreted at lower levels in *Pik3ca*<sup>RBD/-</sup> BMDMs, independently of the conditions under study. 12 proteins were differentially secreted in both experimental conditions under study (**Fig. 5C** [↗](#)).


Differentially secreted proteins are involved in two main biological processes: complement activation (C1qa, C1qb and C1qc connected through physical interaction and C3, C9 and Cfb through coexpression) and lysosome function, mainly cathepsins (Ctsd, Ctsb, Ctsz, Cst3, Psap, Anxa1 and Gsn). Together, the complement cascade and lysosome function work in concert to provide an effective defence against pathogens and promote overall maintenance of cellular homeostasis<sup>22</sup> [↗](#),<sup>45</sup> [↗](#)<sup>47</sup> [↗](#) and data from the secretome analysis of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs suggests that RAS activation of p110α may play a crucial role in the regulation of both response pathways.

## Disruption of RAS-p110α signalling leads to altered lysosomal function

We next aimed to functionally validate the proteomics data suggesting that RAS-p110α activation regulates lysosomal function in macrophages. First, we performed immunofluorescence analysis of lysosomal-associated membrane protein 1 (LAMP1) to assess lysosomal biogenesis and function in *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs. Analysis of LAMP1 in *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs showed a decrease in LAMP1 expression in steady state conditions (**Fig. 6A** [↗](#)) suggesting a decrease in the number of lysosomes after disruption of RAS-p110α interaction. However, *Pik3ca*<sup>RBD/-</sup> showed increased levels of Lamp1 expression when subjected to phagocytosis of apoptotic cells, suggesting an increase in the number of phagolysosomes present in these cells.

We hypothesised that the increase in Lamp1 expression in BMDMs lacking RAS-p110α interaction during phagocytosis could be attributed to aberrant lysosomal function and phagolysosome retention. Thus, we next evaluated lysosomal activity by using the lysosomotropic dye lysotracker red coupled with epifluorescence analysis, since it is specifically taken up by acidic organelles. As such, its accumulation is proportional to the number of acidic vesicles. Data analysis showed that disruption of RAS-PI3K in BMDMs leads to a significant decrease in lysotracker uptake, both in unstimulated conditions and also after activation with LPS+IFN-γ (**Fig. 6B** [↗](#)). Similar results were obtained with control BMDMs treated with BYL719, the p110α specific inhibitor. Data analysis showed a decrease in the uptake of lysotracker after p110α inhibition (**Fig. 6B** [↗](#)), further suggesting that loss of p110α function in BMDMs results in altered lysosomal pH. To confirm that lysosomal pH of *Pik3ca*<sup>RBD/-</sup> BMDMs was less acidic, we next stained *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs with Green lysosensor, a pH-sensitive dye that exhibit a pH-dependent increase in fluorescent intensity upon lysosomal acidification. As shown in **Fig. 6C** [↗](#), *Pik3ca*<sup>RBD/-</sup> BMDMs presented attenuated fluorescent intensity both in unstimulated and activated when compared with that in the control group, indicating that their lysosomal content is less acidic than in *Pik3ca*<sup>WT/-</sup> BMDMs.


Considering that *Pik3ca*<sup>RBD/-</sup> lysosomes were less acidic, we next investigated the expression level and activation of some of the cathepsins identified in the secretome analysis (**Fig. 5** [↗](#)) by western blotting. Cathepsins play a critical role in lysosomal protein degradation. They are initially



**Figure 5.**

### Secretome analysis of *Pik3ca*<sup>RBD/-</sup> BMDMs suggested a defect in complement activation and lysosomal function.

(**A** and **B**) Volcano plots of secretome analysis from *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs in non-stimulated conditions (**A**) or phagocytosing apoptotic cells (**B**). The x-axis shows the log<sub>2</sub> FC of each identified protein and the y-axis the corresponding -log<sub>10</sub> P value. Statistically significant peptides with FC > 2 and p-value < 0.05 are in blue; peptides that do not pass this threshold are in grey; peptides with FC > 2 but p-value ≥ 0.05 are in green; (**C**) Venn diagram showing the overlap of *Pik3ca*<sup>RBD/-</sup> vs *Pik3ca*<sup>WT/-</sup> BMDMs differentially expressed proteins at rest and during phagocytosis of apoptotic cells. Proteins displayed in the blue square represents peptides differentially expressed in resting BMDMs, the brown square represents peptides differentially expressed during phagocytosis, and the peptides differentially expressed in both conditions are displayed in purple; (**D** and **E**) Network analysis of significantly expressed proteins identified in the secretome analysis of *Pik3ca*<sup>RBD/-</sup> vs *Pik3ca*<sup>WT/-</sup> BMDMs in steady state conditions (**D**) or during phagocytosis of apoptotic cells (**E**). The nodes represent individual proteins, and the edges represent known interactions between proteins, either co-expression (purple) or physical interaction (thick pink). The size of each node reflects the significance of differential expression. Proteins in green have been implicated in lysosomal function, while proteins in orange are members of the complement cascade.



**Figure 6.**

### Disruption of RAS-p110α activation in BMDMs leads to abnormal lysosomal function

(A) Representative IF images and quantification analysis of lamp intensity in *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs in steady state conditions and during phagocytosis of apoptotic LKR10 cells. Three independent biological replicates were analysed ( $n \geq 250$  total cells). Error bars indicate mean  $\pm$  SEM; (B) Quantification analysis of lysosomal function in *Pik3ca*<sup>WT/-</sup>, *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/WT</sup> BMDMs treated with BYL719 BMDMs unstimulated or activated with LPS + IFN- $\gamma$  using Lysotracker staining. Three independent biological replicates were analysed ( $n \geq 250$  total cells); (C) Violin plot displaying lysosome pH acidity in unstimulated and LPS+IFN- $\gamma$  stimulated *Pik3ca*<sup>WT/-</sup> (black), *Pik3ca*<sup>RBD/-</sup> (red) and *Pik3ca*<sup>WT/WT</sup> BMDMs treated with BYL719 (Blue) BMDMs, as determined by Lysosensor staining. Three independent biological replicates were analysed ( $n \geq 250$  total cells); (D) *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs were activated with LPS (100 ng/ml) and IFN- $\gamma$  (20 ng/ml) and expression and activation of Cathepsin B and D were blotted; (E) Quantification of phagocytosed apoptotic cells in *Pik3ca*<sup>WT/-</sup>, *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/WT</sup> BMDMs treated with BYL719 BMDMs. Apoptotic cells were labelled with a red tracker and allowed to be phagocytosed by control, *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/WT</sup> BMDMs treated with BYL719 for 16 hours, measuring the amount of cell tracker that was internalized at different time points. Each dot of the graph indicates a different independent experiment. Error bars indicate mean  $\pm$  SEM; (F) Graph displaying the ability of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs to degrade GFP-labelled (pH sensitive) and red tracker-labelled (pH insensitive) apoptotic cells. Three independent biological replicates were analysed. Error bars indicate mean  $\pm$  SEM; (G) Representative image and quantification of Cathepsin D expression in the inflammatory abscess of *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> paws injected with zymosan; (H) Representative image and quantification of Cathepsin D expression in the inflammatory abscess of paws from control and *Pik3ca*<sup>WT/WT</sup> mice treated with BYL719 injected with zymosan.

Statistical significance was obtained using Mann-Whitney test: \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ ; \*\*\*\* $p < 0.0001$ .

synthesized as inactive precursors and are activated through proteolysis in the lysosome at a low pH<sup>48</sup>. We found a reduction in the expression and activation of cathepsin D and cathepsin B in *Pik3ca*<sup>RBD/-</sup> BMDMs upon stimulation with LPS+IFN- $\gamma$  (Figure 6D). This observation suggests that the impaired lysosomal pH observed in *Pik3ca*<sup>RBD/-</sup> BMDMs could potentially account for the observed decrease in cathepsin activity.

Acidification of lysosomes and cathepsin activation are critical steps for activation of the resolute stage of the inflammatory response, so we next evaluated the ability of *Pik3ca*<sup>RBD/-</sup> BMDM's lysosomal compartment to degrade internalised particles. We set up a phagocytosis assay in which *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs were labelled with a green tracker and were allowed to engulf apoptotic cells that were labelled with Cell tracker Red CMTPX. We expected that, if disruption of RAS-p110 $\alpha$  interaction impairs lysosomal degradation of engulfed material, after a certain period of time *Pik3ca*<sup>RBD/-</sup> BMDMs presented an accumulation of the red tracker from non-digested apoptotic cells. Flow cytometry analysis showed that *Pik3ca*<sup>RBD/-</sup> BMDMs accumulate significantly higher levels of red tracker signal from apoptotic cells than *Pik3ca*<sup>WT/-</sup> BMDMs over time, suggesting that disruption of RAS-p110 $\alpha$  in BMDMs causes an accumulation of non-digested apoptotic cells (Fig. 6E). Furthermore, inhibiting p110 $\alpha$  with BYL719 also resulted in a significant accumulation of apoptotic cell debris within BMDMs over time, confirming the involvement of p110 $\alpha$  signalling in the digestion of lysosomal content.

To further confirm the impairment of phagocytosis in *Pik3ca*<sup>RBD/-</sup> BMDMs to degrade phagocytosed particles, we set up another phagocytosis assay in which *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs would phagocytose apoptotic cells that were transduced with GFP (which is pH sensitive<sup>49</sup>) and also labelled with Celltracker Red CMTPX (pH insensitive). *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> BMDMs were allowed to engulf these apoptotic cells for 16 h and after this time, apoptotic cells were eliminated by washes and BMDMs were analysed by flow cytometry at different time points to measure GFP signal. Our results showed that GFP signal is lost significantly faster in control BMDMs than in *Pik3ca*<sup>RBD/-</sup> BMDMs (Fig. 6F). As expected, we did not observe any decay in the red tracker signal. Collectively, these data evidence that *Pik3ca*<sup>RBD/-</sup> BMDMs exhibit a significant impairment in removing engulfed particles, that can be attributed to the absence of lysosome acidification.

We have shown a delayed clearance of apoptotic cells in *Pik3ca*<sup>RBD/-</sup> mice after zymosan injection (Fig. 1). This, together with previous data took us to analyse the levels of Cathepsin D in the inflamed abscess from the paws of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> mice. Results showed a significant decrease in the levels of Cathepsin D in the inflamed area of *Pik3ca*<sup>RBD/-</sup> mice (Fig. 6G) and BYL719-treated mice (Fig. 6H).

In summary, our findings underscore the crucial role of RAS-p110 $\alpha$  signalling axis in maintaining the balance of the inflammatory response and promoting timely resolution, providing further evidence for the significant involvement of RAS-p110 $\alpha$  signalling in the response to inflammatory stimuli.

## Discussion

In this study, we provide compelling evidence that disruption of RAS-p110 $\alpha$  signalling or chemical inhibition of p110 $\alpha$  impairs response to inflammatory stresses due to defects in both monocyte extravasation during the early stages of the inflammatory response and decreased lysosomal function during the later stages.

Monocyte extravasation during the inflammatory response is a critical step that allows immune cells to reach the site of infection or injury<sup>19</sup>. Impairment on this process causes slower or inadequate immune response, as macrophages are crucial for detecting and engulfing pathogens



or cellular debris at the site of inflammation, as well as delayed resolution of inflammation, resulting in prolonged inflammation and potential tissue damage<sup>50, 51</sup>. Our data shows that RAS binding to p110 $\alpha$  is involved in macrophage extravasation by modulating actin dynamics through LIMK-cofilin signalling, two well-known regulators of actin polymerization that facilitate the disassembly of actin filaments<sup>52</sup>. During monocyte extravasation, actin filaments form actin-rich protrusions that are essential for monocyte migration across the endothelium<sup>39, 53</sup>. Our data show that, during extravasation, RAS-p110 $\alpha$  signalling regulates actin depolymerization so monocytes are able to squeeze through endothelial cells<sup>39, 41</sup>.

The disruption of RAS-p110 $\alpha$  in macrophages emerges as a pivotal determinant in cellular mechanics, elucidating a cascade of events resulting in increased F-actin levels. The increase in cytoskeletal components, particularly the actin-modulating proteins LIMK and cofilin and the structural protein F-actin, causes a notable increase in cell stiffness and a concurrent decrease in cell deformability. The orchestration of these changes underscores the intricate balance in cytoskeletal dynamics. Notably, the regulatory role of Rho-GTPases comes into focus as potential mediators of this phenomenon. LIMK phosphorylation can occur through PI3K inhibition<sup>54</sup>, as well as through downstream effectors of phosphoinositide 3-kinase (PI3K), such as phosphorylated p21-activated kinase (PAK)<sup>55</sup> or Rho kinase (ROCK)<sup>54, 56</sup>, which are regulated by Rho GTPases. Rho-GTPases are well-known mediators of actin cytoskeletal rearrangements<sup>57</sup>, and their dysregulation is known to impact cell mechanics<sup>58, 59</sup>. The observed increase in cofilin and F-actin levels, and the subsequent alterations in cell stiffness and deformability, may thus be governed by the modulation of Rho-GTPase activity. This intricate interplay provides a compelling avenue for further investigation into the molecular mechanisms through which RAS-p110 $\alpha$  disruption influences Rho-GTPases, ultimately shaping the biomechanical properties of macrophages and may open avenues for therapeutic interventions targeting cytoskeletal dynamics in macrophages.

Monocyte extravasation shares many features with monocyte egression from the bone marrow. During monocyte egress, monocytes also undergo changes in cytoskeleton dynamics to detach from the sinusoidal endothelial cells and to migrate through the endothelial fenestrae and basement membrane into the bone marrow sinusoids<sup>60</sup>. Thus, the decrease in the number of classical monocytes in the *Pik3ca*<sup>RBD/-</sup> mice in blood may be due, at least in part, to a defect in the extravasation process. This result may also explain, at least partially, the lack of macrophage recruitment to lung tumours found in previous studies using the RBD mouse model<sup>11, 12</sup>.

Our findings also revealed a crucial role of RAS-p110 $\alpha$  activation in the acute phase of the inflammatory response by regulating effective lysosomal degradation of phagocytosed and engulfed material. Phagocytosis constitutes a vital step in the inflammatory response, whereby phagosomes bind to lysosomes to form phagolysosomes in which pathogens are eradicated to facilitate an appropriate host response. This response encompasses antigen presentation to engage T-cell responses, secretion of inflammatory mediators that guide the adaptive immune response, and initiation of tissue repair mechanisms<sup>22, 47</sup>. Our data provide evidence that the activation of RAS-p110 $\alpha$  signalling pathway is involved in the critical process of lysosomal acidification, which is essential for the efficient degradation of internalized particles and the activation of proteolytic enzymes, ultimately resulting in the formation of fully functional lysosomes. Consequently, lack of lysosome acidification impairs the expression and activation of important proteases such as cathepsin B and cathepsin D. The proper functioning of lysosomes is essential for mounting a robust response to inflammatory stress. Lysosomes play a central role in the breakdown of pathogens and dead cells by providing the necessary degradative enzymes and maintaining an acidic environment that facilitates the degradation of engulfed particles<sup>20, 21</sup>. When lysosomal function is compromised in macrophages, the degradation of phagocytosed material becomes impaired, leading to the accumulation of toxic debris. This accumulation subsequently triggers inflammation and causes damage to the surrounding tissues, leading to chronic inflammation and appearance of lysosomal storage diseases (LSD)<sup>61, 62</sup>.

Our data showed that the disruption of Ras-p110 $\alpha$  in macrophages presents a multifaceted impact in inflammatory response, notably manifesting as a dual reduction in NF- $\kappa$ B activation and cathepsin expression coupled with diminished lysosomal function. NF- $\kappa$ B, a pivotal regulator of inflammatory responses<sup>63, 64</sup>, is intricately linked to lysosomal dynamics, with lysosomes playing a crucial role in modulating NF- $\kappa$ B signalling through the degradation of relevant molecules<sup>65, 66</sup>. Additionally, the observed decrease in NF- $\kappa$ B activation aligns with the decline in lysosomal function, which has previously been suggested in previous reports<sup>67, 68</sup>. This bidirectional influence, where perturbations in lysosomal function coincide with alterations in NF- $\kappa$ B activity, underscores the intricate and context-dependent nature of this regulatory network, suggesting a plausible mechanistic interdependence. Our data suggest that disruption in Ras-p110 $\alpha$  signalling cascade could trigger downstream effects impacting both NF- $\kappa$ B activity and lysosomal integrity. This intricate nexus between Ras-p110 $\alpha$  signalling, NF- $\kappa$ B activation, cathepsin expression, and lysosomal function underscores the complexity of cellular regulatory networks. Further analysis of the molecular pathways connecting these observations holds the potential to reveal novel insights into the coordinated regulation of inflammatory and lysosomal processes in macrophages. Thus, further investigations are necessary to decipher the causal relationships and unravel the broader implications of Ras-p110 $\alpha$  disruption in shaping these intricate cellular responses during the inflammatory response.

The possible functional link between the increased actin polymerization and lysosomal dysfunction observed in *Pik3ca*<sup>RBD/-</sup> mice remains an unanswered question. Lysosome acidification and actin polymerization are tightly interconnected processes that have pivotal roles in numerous cellular functions<sup>69–71</sup>. Studies have demonstrated that lysosome acidification can influence actin polymerization dynamics through the activation of specific actin-binding proteins and the modulation of actin-regulatory proteins<sup>69, 72</sup>. Conversely, disruption in lysosome acidification, such as impaired proton pump activity or lysosomal storage disorders, have been associated with changes in actin polymerization and the organization of the cytoskeleton. Notably, it has been reported that Cofilin plays a crucial role in assembling the macromolecular components of the acidification machinery in nascent endosomes by depolymerizing F-actin<sup>73</sup>. Therefore, the intricate relationship between lysosome acidification and actin polymerization suggests a potential reciprocal influence, where perturbations in one process could impact the other. Further investigation is required to determine the precise regulatory mechanisms by which p110 $\alpha$  influences both lysosome acidification and actin polymerization, whether it occurs in a linear manner or through separate pathways.

In summary, our findings offer valuable insights into the underlying mechanisms governing the immune response to inflammation, emphasizing the critical role of RAS-p110 $\alpha$  signalling in the context of inflammation. Given the relevance of p110 $\alpha$  signalling in the resolution of inflammation and the persistent dysregulated inflammation seen in diseases such as rheumatoid arthritis, inflammatory bowel disease, psoriasis, and systemic lupus erythematosus, among others, exploring the potential of p110 $\alpha$  as a therapeutic target in those indications is intriguing. The identification of a p110 $\alpha$  small molecule activator<sup>74</sup> provides a tool to evaluate the therapeutic potential of transiently activating p110 $\alpha$  in these diseases by testing if enhancing p110 $\alpha$  function promotes the resolution of inflammation and potentially alleviate the symptoms and progression of these inflammatory conditions.

## Additional declarations

The authors declare no competing interests.

## Authors' disclosures

The authors declare no conflict of interest.

## Authors' contributions

**Conception and design:** E. Castellano. **Development of methodology:** A. Rosell, M. Alcón Pérez, M.B. Voisin, J. de Paz, V. Rajeeve, C. Cuesta, O. Swinyard, E. Gabandé-Rodriguez, H. Fraile. **Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.):** A. Rosell, M. Alcón Pérez, A. Krywoska, M.B. Voisin, J. Downward, P. Cutillas, E. Castellano. **Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis):** A. Rosell, M. Alcón Pérez, A. Krywoska, M.B. Voisin, V. Rajeeve, A. Berral-Gonzalez, J. De Las Rivas, P. Cutillas, E. Castellano. **Writing, review, and/or revision of the manuscript:** A. Rosell, M. Alcón Pérez, M.B. Voisin, J. De Las Rivas, P. Cutillas, E. Castellano.

## Acknowledgements

This work was supported by grants from the Spanish Ministry of Science and Innovation (RTI2018-099161-A-I00), Programa JAE-Intro ICU from CSIC (JAEICU-21- IBMCC-6), JCyL (CSI185-20), Marie Curie Initial Training Network on Tumour Infiltrating Myeloid Cell Compartment (PF7 MCA-ITN317445) and CRUK-Barts Cancer Centre Development Fund. This research was co-financed by FEDER funds. The CIC is supported by the Programa de Apoyo a Planes Estratégicos de Investigación de Estructuras de Investigación de Excelencia of Castilla y León autonomous government (CLC-2017-01) and AECC Excellence program Stop Ras Cancers (EPAEC222641CICS).


## Supplementary data

## Methods

### Animal studies

For removal of *Pik3ca*-floxed allele, *Pik3ca*<sup>RBD/lox</sup> and *Pik3ca*<sup>WT/flox</sup> mice<sup>64</sup> were given 3.2 mg tamoxifen (Sigma) dissolved in 80 µl of corn oil by oral gavage once per day during 3 consecutive days. Efficiency of tamoxifen treatment was routinely performed by genotyping for the presence of the floxed allele.


For **paw oedema studies** *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>WT/-</sup> mice<sup>64</sup> were divided into groups of four two weeks before oedema induction. Before inducing the paw oedema, the mice were anesthetized with 4% isoflurane. To induce the oedema, mice received ipsilateral i.pl. injection (30 µl) of either zymosan (10 µg/µl, Sigma Aldrich) or PBS into the back-hind paw. Injection of 0.1mg/kg Buprenorphine (NOAH, Vetergesic) was given for pain prevention. Paw thickness was measured using a calliper every hour during the first 6 hours after injection and then at 8, 10 hours and twice per day afterwards. Buprenorphine was injected twice per day during the length of the experiment. Mice were kept, managed, and sacrificed in the NUCLEUS animal facility of the



Supplementary Figure S1.

### *Pik3ca*<sup>RBD/-</sup> mice have more extended inflamed tissue after zymosan injection


(A) Schematic representation of the mouse model used in which the interaction of p110α with RAS was disrupted by the introduction of two-point mutations, T208D and K227A, into the endogenous *Pik3ca* gene (*Pik3ca*<sup>RBD</sup>). Wild-type (*Pik3ca*<sup>WT</sup>) and *Pik3ca*<sup>RBD</sup> mice were bred with mice containing a floxed *Pik3ca* allele and a mouse carrying a conditional Cre recombinase (Cre-ERT2) allele targeted to the ubiquitously expressed Rosa26 locus; (B) Representative PCR showing efficiency of floxed allele removal after treatment with 4-hydroxytamoxifen for 48 hours; (C) *Pik3ca*<sup>WT/Flox</sup> and *Pik3ca*<sup>RBD/Flox</sup> mice were injected with zymosan or PBS in the back-hind paws and inflammation was measured and plotted over time, *Pik3ca*<sup>WT/Flox</sup> n = 5; *Pik3ca*<sup>RBD/Flox</sup> PBS n = 5; *Pik3ca*<sup>WT/Flox</sup> Zymosan n = 5; *Pik3ca*<sup>RBD/Flox</sup> Zymosan n = 5; Error bars indicate mean ± SEM. Significance using 2-way ANOVA test: n.s. non-significant (D) Blood from the same mice was obtained by cardiac puncture prior culling and blood sedimentation was measured. The clear layer formed at the top was used for data quantification. Black dots represent data from *Pik3ca*<sup>WT/-</sup> mice and red dots represent data from *Pik3ca*<sup>RBD/-</sup> mice. Each dot represents an individual mouse. Error bars indicate mean ± SEM. Significance using Student's t test: \* p < 0.05; \*\*p < 0.01. *Pik3ca*<sup>WT/-</sup> PBS n = 6; *Pik3ca*<sup>RBD/-</sup> PBS n = 5; *Pik3ca*<sup>WT/-</sup> Zymosan n = 4; *Pik3ca*<sup>RBD/-</sup> Zymosan n = 4; (E) Representative images showing zymosan-induced inflammatory area in *Pik3ca*<sup>WT/WT</sup> mice and *Pik3ca*<sup>WT/WT</sup> mice treated with BYL-719.



## Supplementary Figure S2.

### Disruption of RAS-p110α interaction does not affect the number of B-cells or T-cells in blood or spleen.


Twelve-week-old mice were treated with tamoxifen and after four weeks, flow cytometry analysis was performed to determine abundance of different immune populations. **(A)** Granulocytes in circulating blood; **(B)** CD3, CD4+ and CD8+ T-cells in circulating blood; **(C)** B-cells in circulating blood; **(D)** Granulocytes in spleen; **(E)** B-cells in spleen; **(F)** T-cells in spleen; **(G)** Spleen weight compared to full body; **(H)** Schematic representation of myeloid precursors differentiation (CMP-common myeloid progenitor; GMP-Granulocyte-monocyte progenitor; MEP-Megakaryocyte-erythroid progenitor); **(I)** Graph showing percentage of bone marrow cells progenitor cells (have not acquire any lineage marker, Lin<sup>+</sup>); **(J)** Graph showing percentage of CMP cells within progenitor cells; **(K)** Graph showing percentage of MEP cells within progenitor cells; **(L)** Graph showing percentage of GMP cells within progenitor cells; **(M)** Graph depicting % of differentiated macrophages obtained in culture from bone marrow precursors in the presence of m-CSF. Black dots represent data from *Pik3ca*<sup>WT/-</sup> mice and red dots represent data from *Pik3ca*<sup>RBD/-</sup> mice. Each dot represents an individual mouse. Error bars indicate mean ± SEM. Significance using Student's t test: n.s. non-significant.



Supplementary Figure S3.

### *Pik3ca*<sup>RBD/-</sup> monocytes do not extravasate through the endothelium

(A) Graph showing levels of neutrophil reconstitution in chimera's bone marrow. Blood from chimera mice was collected and GFP in granulocytes was analysed by flow cytometry to determine engraftment percentage. (B) Graph quantifying the number of monocytes in the blood of chimera mice treated with anti-GR1 (25 µg/mouse/day) for 3 days. (C) Graph showing levels of expression of CCR2 in *Pik3ca*<sup>WT/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs detected by flow cytometry. Error bars indicate mean ± SEM. Significance using Mann-Whitney test: n.s. no-significant



Supplementary Figure S4.

### Disruption of RAS-p110α interaction alters the secretome of BMDMs

*Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs were stimulated with LPS+IFN-γ and expression of (A) CD80; (B) CD86; and (C) MHCII were analysed by flow cytometry (*Pik3ca*<sup>RBD/-</sup> n = 5; *Pik3ca*<sup>WT/-</sup> n = 3). Error bars indicate mean ± SEM. Significance using Mann-Whitney test: n.s. no-significant. (D) WB showing induction of apoptosis in LKR10 cell line treated with cisplatin for 16 hours; (E) Dendrogram showing hierarchical clustering of peptides found in the secretome analysis of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs in steady state conditions; (F) Dendrogram showing hierarchical clustering of peptides found in the secretome analysis of *Pik3ca*<sup>RBD/-</sup> and *Pik3ca*<sup>RBD/-</sup> BMDMs phagocytosing apoptotic cells.



|    | A      | B                 | C         | D         | E         | F           | G           | H                      | I                                                                                                 |
|----|--------|-------------------|-----------|-----------|-----------|-------------|-------------|------------------------|---------------------------------------------------------------------------------------------------|
|    | db_ID  | UniProt_EntryName | Gene_Name | log2(FC)  | AveSignal | t-statistic | p-value     | Peptide                | Protein_Description                                                                               |
| 1  | p00979 | 1433E_MOUSE       | Ywhae     | -2.62309  | 74.03284  | -2.39545    | 0.025186147 | EAAENSLVAYK            | 14-3-3 protein epsilon OS=Mus musculus GN=Ywhae PE=1 SV=1                                         |
| 2  | p00568 | 1433F_MOUSE       | Ywhah     | -3.80235  | 87.71693  | -2.81089    | 0.012094651 | AVTELENPLSNEDR         | 14-3-3 protein eta OS=Mus musculus GN=Ywhah PE=1 SV=2                                             |
| 3  | p04399 | 1433S_MOUSE       | Sfn       | -4.61501  | 105.35784 | -2.21317    | 0.041899479 | NLLSVAYK               | 14-3-3 protein sigma OS=Mus musculus GN=Sfn PE=1 SV=2                                             |
| 4  | p02233 | ACTB1_MOUSE       | Actb12    | -4.77874  | 49.54558  | -2.68071    | 0.015875482 | HQGVVVMGMGQK           | Beta-actin-like protein 2 OS=Mus musculus GN=Actb2 PE=1 SV=1                                      |
| 5  | p00304 | AKR1A1_MOUSE      | Akr1a1    | -4.58145  | 88.03299  | -3.51125    | 0.003186888 | ALEVLVAK               | Alcohol dehydrogenase [NADP+] OS=Mus musculus GN=Akr1a1 PE=1 SV=3                                 |
| 6  | p04738 | ALDOA_MOUSE       | Aldoa     | -5.52762  | 85.98573  | -2.79612    | 0.013657994 | QLLTADDR               | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                               |
| 7  | p02412 | ALDOA_MOUSE       | Aldoa     | -6.47337  | 168.83214 | -2.34654    | 0.033239605 | IGEHTPSALAMENANVLAR    | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                               |
| 8  | p01918 | ALDOA_MOUSE       | Aldoa     | -4.06699  | 89.00265  | -2.26262    | 0.039080506 | GILAADESTGSIK          | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                               |
| 9  | p00505 | ALDOA_MOUSE       | Aldoa     | -4.37786  | 92.11914  | -2.17614    | 0.047341789 | ADDGRPFPPQVVK          | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                               |
| 10 | p00369 | ALDR_MOUSE        | Akr1b1    | -3.25918  | 50.91124  | -2.53517    | 0.029889815 | SPPGGVTEAVK            | Aldose reductase OS=Mus musculus GN=Akr1b1 PE=1 SV=3                                              |
| 11 | p03031 | AN32B_MOUSE       | Anp32b    | -6.63248  | 156.82890 | -2.66971    | 0.045628004 | LAELFSLTHLNLGNNLK      | Acidic leucine-rich nuclear phosphoprotein 32 family member B OS=Mus musculus GN=Anp32b PE=1 SV=1 |
| 12 | p03304 | APOE_MOUSE        | ApoE      | -2.90888  | 87.23308  | -2.39517    | 0.031311937 | LGADMEDLR              | Apolipoprotein E OS=Mus musculus GN=ApoE PE=1 SV=2                                                |
| 13 | p02551 | ARF4_MOUSE        | Arf4      | -5.45667  | 120.56874 | -2.55781    | 0.031159731 | ILMVGLDAAAG            | ADP-ribosylation factor 4 OS=Mus musculus GN=Arf4 PE=1 SV=2                                       |
| 14 | p04971 | ARP2_MOUSE        | Arp2      | -5.96573  | 40.95109  | -3.21875    | 0.006802071 | RLDIAGIR               | Actin-related protein 2 OS=Mus musculus GN=Arp2 PE=1 SV=1                                         |
| 15 | p06167 | ARPC2_MOUSE       | Actr2     | -3.92057  | 56.19169  | -2.42426    | 0.030857962 | VGNEIK                 | Actin-related protein 2 OS=Mus musculus GN=Arp2 PE=1 SV=1                                         |
| 16 | p02747 | ARPC4_MOUSE       | Arp4      | -4.41613  | 111.62260 | -2.45985    | 0.031937251 | IVAEELFK               | Actin-related protein 2/3 complex subunit 4 OS=Mus musculus GN=Arp4 PE=1 SV=3                     |
| 17 | p02534 | BGH3_MOUSE        | Tgfb1     | -5.50174  | 91.78027  | -2.93345    | 0.009339267 | LGDEPALR               | Transforming growth factor-beta-induced protein lg-h3 OS=Mus musculus GN=Tgfb1 PE=1 SV=1          |
| 18 | p00061 | BGH3_MOUSE        | Tgfb1     | -5.14440  | 72.21607  | -2.38970    | 0.048884543 | ADHATHNGVVLHLDK        | Transforming growth factor-beta-induced protein lg-h3 OS=Mus musculus GN=Tgfb1 PE=1 SV=1          |
| 19 | p02283 | BLVRB_MOUSE       | Blvrb     | -6.23169  | 101.75248 | -3.97429    | 0.001885854 | AIIFGATGR              | Flavin reductase (NADPH) OS=Mus musculus GN=Blvrb PE=1 SV=3                                       |
| 20 | p04893 | C1QC_MOUSE        | C1qc      | -3.76890  | 72.61010  | -2.14293    | 0.045378462 | QVSSGGVLLR             | Complement C1q subcomponent subunit C OS=Mus musculus GN=C1qc PE=1 SV=2                           |
| 21 | p00699 | CALM1_MOUSE       | Calm1     | -3.28594  | 101.49865 | -2.36434    | 0.031162013 | DGNGYISAAELR           | Calmodulin OS=Mus musculus GN=Calm1 PE=1 SV=2                                                     |
| 22 | p05860 | CATD_MOUSE        | Ctsd      | -3.72920  | 107.46557 | -2.15842    | 0.044756496 | ITTEPSELLK             | Cathepsin D OS=Mus musculus GN=Ctsd PE=1 SV=1                                                     |
| 23 | p03549 | CAZAI_MOUSE       | Capza1    | -4.90584  | 133.31643 | -2.64031    | 0.018655110 | LLLNNDNLLR             | F-actin-capping protein subunit alpha-1 OS=Mus musculus GN=Capza1 PE=1 SV=4                       |
| 24 | p02009 | CD14_MOUSE        | Cd14      | -4.02664  | 68.61897  | -2.40057    | 0.037599594 | GNPFLDSEHSEK           | Monocyte differentiation antigen CD14 OS=Mus musculus GN=Cd14 PE=1 SV=1                           |
| 25 | p02687 | CLH1_MOUSE        | Cltc      | -6.27808  | 156.39427 | -2.34448    | 0.047619848 | ISGETIFVTAPHEATAGIGVNR | Clathrin heavy chain 1 OS=Mus musculus GN=Cltc PE=1 SV=3                                          |
| 26 | p05695 | CMMPK2_MOUSE      | Cmpk2     | -7.08929  | 97.66407  | -2.47981    | 0.035384012 | TKEEALEANNVFR          | UMP-CMP kinase 2, mitochondrial OS=Mus musculus GN=Cmpk2 PE=1 SV=2                                |
| 27 | p02363 | DESM_MOUSE        | Des       | -6.63594  | 148.71331 | -3.43881    | 0.027812451 | IESLNEIEAFLKK          | Desmin OS=Mus musculus GN=Des PE=1 SV=3                                                           |
| 28 | p05053 | DPLY2_MOUSE       | Dpysl2    | -4.37311  | 62.87575  | -2.50284    | 0.024491460 | SAAEVIAQAR             | Dihydropyrimidinase-related protein 2 OS=Mus musculus GN=Dpysl2 PE=1 SV=1                         |
| 29 | p04320 | DYL2_MOUSE        | Dynl2     | -4.24626  | 68.19918  | -3.27247    | 0.032328410 | NFSQSYTHETK            | Dynein light chain 2, cytoplasmic OS=Mus musculus GN=Dynl2 PE=1 SV=2                              |
| 30 | p00729 | ECM1_MOUSE        | Ecm1      | -5.19102  | 146.05218 | -4.09735    | 0.006697195 | DILTDLRS               | Extracellular matrix protein 1 OS=Mus musculus GN=Ecm1 PE=1 SV=2                                  |
| 31 | p02417 | EF1A2_MOUSE       | Eef1a2    | -4.06820  | 90.30546  | -2.38601    | 0.029853339 | IGIGITVPVGR            | Elongation factor 1-alpha 2 OS=Mus musculus GN=Eef1a2 PE=1 SV=1                                   |
| 32 | p05782 | EF1B_MOUSE        | Eef1b     | -3.66103  | 170.14153 | -2.27517    | 0.032170909 | TPAGLQVLNDYLADK        | Elongation factor 1-beta OS=Mus musculus GN=Eef1b PE=1 SV=5                                       |
| 33 | p06134 | EF2_MOUSE         | Eef2      | -5.91556  | 101.58858 | -2.97047    | 0.008634240 | VFGSVYGLTK             | Elongation factor 2 OS=Mus musculus GN=Eef2 PE=1 SV=2                                             |
| 34 | p05280 | EF2_MOUSE         | Eef2      | -3.19095  | 68.66118  | -2.69037    | 0.031652774 | MGPNYTPGK              | Elongation factor 2 OS=Mus musculus GN=Eef2 PE=1 SV=2                                             |
| 35 | p06367 | EF2_MOUSE         | Eef2      | -3.23794  | 105.16843 | -2.23066    | 0.036816051 | VNFTVDQIR              | Elongation factor 2 OS=Mus musculus GN=Eef2 PE=1 SV=2                                             |
| 36 | p00460 | EF2_MOUSE         | Eef2      | -4.55647  | 126.75721 | -2.22125    | 0.044921497 | ARPPFDGLAEDIDKGEVSAR   | Elongation factor 2 OS=Mus musculus GN=Eef2 PE=1 SV=2                                             |
| 37 | p06113 | ESD2_MOUSE        | Esd       | -4.60197  | 89.45241  | -2.62372    | 0.021188125 | VFEHSSVELK             | S-formylglutathione hydrolase OS=Mus musculus GN=Esd PE=1 SV=1                                    |
| 38 | p03613 | FABP5_MOUSE       | Fabp5     | -5.53684  | 107.62544 | -3.93606    | 0.005897717 | LMESHGFEEYMK           | Fatty acid-binding protein, epidermal OS=Mus musculus GN=Fabp5 PE=1 SV=3                          |
| 39 | p00611 | FABP5_MOUSE       | Fabp5     | -6.93746  | 90.48486  | -2.87799    | 0.020949532 | LMESHGFEEYMK           | Fatty acid-binding protein, epidermal OS=Mus musculus GN=Fabp5 PE=1 SV=3                          |
| 40 | p05529 | FLNA_MOUSE        | Flna      | -5.26445  | 167.11916 | -2.85913    | 0.024889758 | VYTPPMAPGSYLISIK       | Filamin A OS=Mus musculus GN=Flna PE=1 SV=5                                                       |
| 41 | p06706 | FLNA_MOUSE        | Flna      | -4.14179  | 127.55676 | -2.60133    | 0.029026511 | YIPVQGGPVGNNVYGGDHPFK  | Filamin A OS=Mus musculus GN=Flna PE=1 SV=5                                                       |
| 42 | p01763 | G3P_MOUSE         | Gapdh     | -3.89247  | 76.45923  | -2.44953    | 0.024854423 | GAAGNIPASTGAARK        | Glycerinaldehyde-3-phosphate dehydrogenase OS=Mus musculus GN=Gapdh PE=1 SV=2                     |
| 43 | p02183 | GPI_MOUSE         | Gpi       | -9.24929  | 61.22198  | -6.73154    | 0.000150963 | HVVALSTYAK             | Glucose-6-phosphate isomerase OS=Mus musculus GN=Gpi PE=1 SV=4                                    |
| 44 | p02266 | GELS_MOUSE        | Gsn       | -4.76822  | 69.46049  | -3.45218    | 0.019022185 | HVYNEVYVQR             | Gelsolin OS=Mus musculus GN=Gsn PE=1 SV=3                                                         |
| 45 | p02970 | GLRX1_MOUSE       | Glrx      | -7.67625  | 134.09504 | -3.57431    | 0.007454774 | KTOEILSQLPFK           | Glutaredoxin-1 OS=Mus musculus GN=Glrx PE=1 SV=3                                                  |
| 46 | p02969 | GLRX1_MOUSE       | Glrx      | -6.62854  | 138.50807 | -3.66024    | 0.022916283 | KTOEILSQLPFK           | Glutaredoxin-1 OS=Mus musculus GN=Glrx PE=1 SV=3                                                  |
| 47 | p05810 | GLRX1_MOUSE       | Glrx      | -6.95820  | 150.98028 | -3.17830    | 0.025553613 | KTOEILSQLPFK           | Glutaredoxin-1 OS=Mus musculus GN=Glrx PE=1 SV=3                                                  |
| 48 | p00559 | GPNMB_MOUSE       | Gpnmb     | -2.58240  | 183.93820 | -2.13527    | 0.048901657 | DVYVITDQIPVFTMSQK      | Transmembrane glycoprotein NMB OS=Mus musculus GN=Gpnmb PE=1 SV=2                                 |
| 49 | p01636 | GRIK1_MOUSE       | Grik1     | -7.35968  | 88.48448  | -2.79172    | 0.021296807 | FNNMLK                 | Glutamate receptor ionotropic, kainate 1 OS=Mus musculus GN=Grik1 PE=2 SV=2                       |
| 50 | p01581 | HA13_MOUSE        | H2-D1     | -4.33720  | 150.12532 | -3.55960    | 0.012404526 | FISVGYVDNTEFVR         | H-2 class I histocompatibility antigen, D-K alpha chain OS=Mus musculus GN=H2-D1 PE=1 SV=1        |
| 51 | p01580 | HA1K_MOUSE        | H2-K1     | -4.33720  | 150.12532 | -3.55960    | 0.012404526 | FISVGYVDNTEFVR         | H-2 class I histocompatibility antigen, K-K alpha chain OS=Mus musculus GN=H2-K1 PE=1 SV=1        |
| 52 | p03379 | HB2_MOUSE         | Hbb-b2    | -4.66941  | 85.58305  | -2.92283    | 0.014035564 | LHVDPENFR              | Hemoglobin subunit beta-2 OS=Mus musculus GN=Hbb-b2 PE=1 SV=2                                     |
| 53 | p06735 | HMOX1_MOUSE       | Hmox1     | -3.00732  | 114.70633 | -2.29491    | 0.048081881 | YLGDLGGQVQLK           | Heme oxygenase 1 OS=Mus musculus GN=Hmox1 PE=1 SV=1                                               |
| 54 | p01346 | HOMEZ_MOUSE       | Homez     | -4.54273  | 86.84165  | -2.31735    | 0.031281112 | EQLAIK                 |                                                                                                   |
| 55 | p06185 | IF5A2_MOUSE       | Eif5a2    | -10.52357 | 156.26635 | -3.29315    | 0.017115649 | VHLVGIDIFTGK           | Eukaryotic translation initiation factor 5A-2 OS=Mus musculus GN=Eif5a2 PE=1 SV=3                 |
| 56 | p06186 | IF5A2_MOUSE       | Eif5a2    | -6.39888  | 156.09916 | -2.49305    | 0.020732725 | VHLVGIDIFTGK           | Eukaryotic translation initiation factor 5A-2 OS=Mus musculus GN=Eif5a2 PE=1 SV=3                 |
| 57 | p05312 | IPO5_MOUSE        | Ipo5      | -4.42108  | 123.46736 | -3.64426    | 0.006827466 | SLVEADTPVK             | Importin-5 OS=Mus musculus GN=Ipo5 PE=1 SV=3                                                      |
| 58 | p03023 | IQAAT_MOUSE       | Iqgap1    | -4.81313  | 133.82886 | -3.12844    | 0.020998024 | LAAYANNAIAQK           | Ras GTPase-activating like protein IQGAP1 OS=Mus musculus GN=Iqgap1 PE=1 SV=2                     |

**Supplementary Table 1.**

**Differentially expressed peptides identified in *Pik3ca*<sup>RBD/-</sup> vs *Pik3ca*<sup>WT/-</sup> secretome of BMDMs under steady state conditions**

|     | A      | B            | C        | D        | E         | F        | G           | H                       | I                                                                                                  |
|-----|--------|--------------|----------|----------|-----------|----------|-------------|-------------------------|----------------------------------------------------------------------------------------------------|
| 60  | p01911 | ITIH3_MOUSE  | Itih3    | -5,80756 | 72.67106  | -4,14492 | 0.009499008 | GHSVFKPSLDQGR           | Inter-alpha-trypsin inhibitor heavy chain H3 OS=Mus musculus GN=Itih3 PE=1 SV=3                    |
| 61  | p01483 | ITIH3_MOUSE  | Itih3    | -4,75206 | 43.98185  | -2,28199 | 0.048834983 | FAHNVVTTR               | Inter-alpha-trypsin inhibitor heavy chain H3 OS=Mus musculus GN=Itih3 PE=1 SV=3                    |
| 62  | p06507 | K1C27_MOUSE  | Krt27    | -3,99871 | 57.78918  | -2,28197 | 0.040171056 | VTMQLNDR                | Keratin, type I cytoskeletal 27 OS=Mus musculus GN=Krt27 PE=1 SV=1                                 |
| 63  | p03892 | KAP9_MOUSE   | Prkar1a  | -4,08671 | 156.63972 | -2,57432 | 0.042972739 | LTVDADLEPVQFEDGQK       | cAMP-dependent protein kinase type I-alpha regulatory subunit OS=Mus musculus GN=Prkar1a PE=1 SV=3 |
| 64  | p01975 | KATL2_MOUSE  | Katn2    | -6,58928 | 115.75448 | -3,02652 | 0.007188948 | GLLYGPPPTGK             | Katanin p60 ATPase-containing subunit A-like 2 OS=Mus musculus GN=Katn2 PE=2 SV=2                  |
| 65  | p01641 | KCD12_MOUSE  | Kctd12   | -6,46669 | 166.10145 | -2,36965 | 0.035645082 | FNLEQAFDK               | BTB/POZ domain-containing protein KCTD12 OS=Mus musculus GN=Kctd12 PE=1 SV=1                       |
| 66  | p01821 | KPYM_MOUSE   | Pkm      | -5,18495 | 88.35175  | -3,05771 | 0.007575392 | GDYPLEAVR               | Pyruvate kinase PKM OS=Mus musculus GN=Pkm PE=1 SV=4                                               |
| 67  | p03381 | KSYK_MOUSE   | Syk      | -5,64892 | 91.99532  | -2,91747 | 0.012112354 | LIATTAHEK               | Tyrosine-protein kinase SYK OS=Mus musculus GN=Syk PE=1 SV=2                                       |
| 68  | p01875 | LASP1_MOUSE  | Lasp1    | -3,19614 | 120.21599 | -2,38764 | 0.033019550 | GFSVVAADPELQR           | LIM and SH3 domain protein 1 OS=Mus musculus GN=Lasp1 PE=1 SV=1                                    |
| 69  | p04813 | LASP1_MOUSE  | Lasp1    | -3,17596 | 40.90733  | -2,25497 | 0.041733101 | COSELOSQVR              | LIM and SH3 domain protein 1 OS=Mus musculus GN=Lasp1 PE=1 SV=1                                    |
| 70  | p05045 | LDHA_MOUSE   | Ldha     | -7,48129 | 63.71028  | -2,13969 | 0.046440818 | RHPHSTMIK               | L-lactate dehydrogenase A chain OS=Mus musculus GN=Ldha PE=1 SV=3                                  |
| 71  | p03928 | LDHC_MOUSE   | Ldhc     | -5,88676 | 88.57748  | -2,95233 | 0.008973032 | LVITAGAR                | L-lactate dehydrogenase C chain OS=Mus musculus GN=Ldhc PE=1 SV=2                                  |
| 72  | p05998 | LEO3_MOUSE   | Lgals3   | -7,70489 | 86.05659  | -3,33027 | 0.008984627 | VAVNDHALQYNHR           | Galectin-3 OS=Mus musculus GN=Lgals3 PE=1 SV=3                                                     |
| 73  | p05997 | LEO3_MOUSE   | Lgals3   | -4,86224 | 85.03049  | -2,94024 | 0.009205941 | VAVNDHALQYNHR           | Galectin-3 OS=Mus musculus GN=Lgals3 PE=1 SV=3                                                     |
| 74  | p05538 | LMNA_MOUSE   | Lmna     | -4,15277 | 101.22930 | -2,31311 | 0.032844347 | TALNSTGEEVAMR           | Prolamin-A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                                     |
| 75  | p00433 | MOES_MOUSE   | Men      | -4,32043 | 48.19669  | -2,69292 | 0.013670496 | QMVGQEDLEK              | Moesin OS=Mus musculus GN=Men PE=1 SV=3                                                            |
| 76  | p05813 | MOES_MOUSE   | Men      | -5,18351 | 143.28100 | -2,70895 | 0.043560856 | TQEQLASEMAELTR          | Moesin OS=Mus musculus GN=Men PE=1 SV=3                                                            |
| 77  | p01144 | MYL3_MOUSE   | My3      | -3,81919 | 73.81483  | -3,51726 | 0.004315833 | EGNGTVMGAELR            | Myosin light chain 3 OS=Mus musculus GN=My3 PE=1 SV=4                                              |
| 78  | p01142 | MYL6_MOUSE   | My6      | -3,81919 | 73.81483  | -3,51726 | 0.004315833 | EGNGTVMGAELR            | Myosin light polypeptide 6 OS=Mus musculus GN=My6 PE=1 SV=3                                        |
| 79  | p00350 | NDKA_MOUSE   | Nme1     | -3,00862 | 84.88688  | -2,18166 | 0.039674556 | YMLGETNPADSKPTTR        | Nucleoside diphosphate kinase A OS=Mus musculus GN=Nme1 PE=1 SV=1                                  |
| 80  | p05608 | NDKB_MOUSE   | Nme2     | -4,42424 | 94.98301  | -2,20593 | 0.039336023 | FTAIKPDGVQR             | Nucleoside diphosphate kinase B OS=Mus musculus GN=Nme2 PE=1 SV=1                                  |
| 81  | p01835 | OTUB1_MOUSE  | Otub1    | -4,26562 | 71.24322  | -3,01089 | 0.009433455 | GEGGTTNPHVFPEGEPEK      | Ubiquitin thioesterase OTUB1 OS=Mus musculus GN=Otub1 PE=1 SV=2                                    |
| 82  | p05816 | PARK7_MOUSE  | Par7     | -3,57628 | 172.05670 | -2,44281 | 0.045250543 | TGQPYDVVVLPGNGLGAQNLS   | Protein deglycase DJ-1 OS=Mus musculus GN=Par7 PE=1 SV=1                                           |
| 83  | p02289 | PCBP1_MOUSE  | Pcbp1    | -4,21269 | 48.09057  | -2,17281 | 0.047653607 | IANPVEGSSGR             | Poly(rC)-binding protein 1 OS=Mus musculus GN=Pcbp1 PE=1 SV=1                                      |
| 84  | p05329 | PESP1_MOUSE  | Pebp1    | -5,33878 | 159.43485 | -2,61691 | 0.040513442 | VLTTPTVMNRPSSISWGLDPQK  | Phosphatylethanolamine-binding protein 1 OS=Mus musculus GN=Pebp1 PE=1 SV=3                        |
| 85  | p04009 | PER1_MOUSE   | Prph     | -8,82554 | 131.99956 | -3,02913 | 0.006662090 | MALDIEATYR              | Peripherin OS=Mus musculus GN=Prph PE=1 SV=2                                                       |
| 86  | p01696 | PGAM1_MOUSE  | Pgam1    | -3,86373 | 123.97364 | -2,63326 | 0.023486664 | FSGUYDADLSAPGHEEAK      | Phosphoglycerate mutase 1 OS=Mus musculus GN=Pgam1 PE=1 SV=3                                       |
| 87  | p04123 | PIPNA_MOUSE  | Pipna    | -8,46627 | 99.96031  | -2,72852 | 0.042581907 | MLAPEGALNHEK            | Phosphatidylinositol transfer protein alpha isoform OS=Mus musculus GN=Pipna PE=1 SV=2             |
| 88  | p01679 | PLD4_MOUSE   | Pld4     | -5,82449 | 115.05170 | -2,25504 | 0.048119538 | EQGLLR                  | Phospholipase D4 OS=Mus musculus GN=Pld4 PE=1 SV=1                                                 |
| 89  | p03894 | PLEC_MOUSE   | Plnc     | -5,18071 | 66.79403  | -2,31556 | 0.043431178 | LTVDENVR                | Plectin OS=Mus musculus GN=Plnc PE=1 SV=3                                                          |
| 90  | p02196 | PLF4_MOUSE   | Pf4      | -8,87763 | 84.65049  | -3,25665 | 0.008776991 | FLISLEVIK               | Platelet factor 4 OS=Mus musculus GN=Pf4 PE=1 SV=1                                                 |
| 91  | p06817 | PLSL_MOUSE   | Lcp1     | -4,72870 | 172.89474 | -2,57042 | 0.030525937 | YTLNLEDIGGGQK           | Plastin-2 OS=Mus musculus GN=Lcp1 PE=1 SV=4                                                        |
| 92  | p00594 | PLSL_MOUSE   | Lcp1     | -4,53563 | 119.98343 | -2,28069 | 0.046076964 | AYYHLEQVAPK             | Plastin-2 OS=Mus musculus GN=Lcp1 PE=1 SV=4                                                        |
| 93  | p04114 | PLST_MOUSE   | Pls3     | -3,00524 | 139.27803 | -2,20917 | 0.039484235 | MINLSVPDTR              | Plastin-3 OS=Mus musculus GN=Pls3 PE=1 SV=3                                                        |
| 94  | p05516 | PLTP_MOUSE   | Pltp     | -7,49544 | 101.22057 | -3,70357 | 0.007901860 | YTSAAALDLVK             | Phospholipid transfer protein OS=Mus musculus GN=Pltp PE=1 SV=1                                    |
| 95  | p02128 | PLTP_MOUSE   | Pltp     | -5,23056 | 178.64594 | -2,98934 | 0.017695170 | GVOIPLPEGINFVR          | Phospholipid transfer protein OS=Mus musculus GN=Pltp PE=1 SV=1                                    |
| 96  | p06462 | PP1A_MOUSE   | Ppia     | -4,11789 | 143.72076 | -2,21186 | 0.036802790 | VSFELFADKVPK            | Peptidyl-prolyl cis-trans isomerase A OS=Mus musculus GN=Ppia PE=1 SV=2                            |
| 97  | p05303 | PRDX2_MOUSE  | Prdx2    | -7,86175 | 83.03399  | -2,76504 | 0.016201871 | SLSQNYGLVK              | Peroxisome oxidin-2 OS=Mus musculus GN=Prdx2 PE=1 SV=3                                             |
| 98  | p03465 | PRDX3_MOUSE  | Prdx3    | -3,75675 | 107.39118 | -2,38908 | 0.039603084 | LIADPTGAFK              | Peroxisome oxidin-3, mitochondrial OS=Mus musculus GN=Prdx3 PE=1 SV=2                              |
| 99  | p05451 | PSA3_MOUSE   | Psm3     | -3,00883 | 69.70984  | -2,34945 | 0.047254359 | AVENSSTAIGIR            | Proteasome subunit alpha type-3 OS=Mus musculus GN=Psm3 PE=1 SV=3                                  |
| 100 | p00403 | PSME2_MOUSE  | Psm2     | -4,11663 | 126.61836 | -2,43162 | 0.031850182 | ALPIDPPDPFPPKDDMETKDEK  | Proteasome activator complex subunit 2 OS=Mus musculus GN=Psm2 PE=1 SV=4                           |
| 101 | p03937 | RAN_MOUSE    | Ran      | -4,31902 | 74.71141  | -2,41820 | 0.028004649 | LVLVGGDGTGK             | GTP-binding nuclear protein Ran OS=Mus musculus GN=Ran PE=1 SV=3                                   |
| 102 | p00739 | RIN_MOUSE    | Rnh1     | -2,85060 | 163.64283 | -2,57552 | 0.037337382 | DISSAVGANPALTELSLR      | Ribonuclease inhibitor OS=Mus musculus GN=Rnh1 PE=1 SV=1                                           |
| 103 | p03275 | ROA3_MOUSE   | Hrmpa3   | -4,21423 | 179.76796 | -2,27442 | 0.046568411 | LFQGLGFATTTDDSLR        | Heterogeneous nuclear ribonucleoprotein A3 OS=Mus musculus GN=Hrmpa3 PE=1 SV=1                     |
| 104 | p01257 | S100A4_MOUSE | S100a4   | -4,96465 | 118.33344 | -2,46407 | 0.025556167 | ELPFSFLK                | Protein S100-A4 OS=Mus musculus GN=S100a4 PE=1 SV=1                                                |
| 105 | p00850 | SAA3_MOUSE   | Saa3     | -5,15245 | 71.79133  | -2,89574 | 0.020391896 | DPNHFRAPGLPK            | Serum amyloid A-3 protein OS=Mus musculus GN=Saa3 PE=1 SV=1                                        |
| 106 | p02567 | SELB_MOUSE   | Eefsec   | -8,86181 | 119.45437 | -3,58640 | 0.024420020 | ILTPTLK                 | Selenocysteine-specific elongation factor OS=Mus musculus GN=Eefsec PE=1 SV=2                      |
| 107 | p00754 | SGP1_MOUSE   | Sgp1     | -5,81057 | 170.32554 | -3,09284 | 0.017938629 | DLALEKGK                | Sphingosine 1-phosphate lyase 1 OS=Mus musculus GN=SGP1 PE=1 SV=1                                  |
| 108 | p02261 | SODC_MOUSE   | Sod1     | -5,92187 | 54.46354  | -3,18209 | 0.004332977 | HYGLDGNVATGAK           | Superoxide dismutase [Cu-Zn] OS=Mus musculus GN=Sod1 PE=1 SV=2                                     |
| 109 | p02097 | SPB6_MOUSE   | Serpinb6 | -4,42458 | 126.55736 | -2,57216 | 0.019873408 | GTTASQAQAALADK          | Serpin B6 OS=Mus musculus GN=Serpinb6 PE=1 SV=1                                                    |
| 110 | p05404 | STBL5_MOUSE  | Stxbp5l  | -7,12546 | 61.58937  | -5,08652 | 0.004124063 | SRSSSSISIDK             | Syntaxin-binding protein 5-like OS=Mus musculus GN=Stxbp5l PE=1 SV=1                               |
| 111 | p02562 | TALD3_MOUSE  | Talpid3  | -9,60934 | 167.90716 | -3,65754 | 0.006617559 | ILRGVONNKK              | Protein TALPID3 OS=Mus musculus GN=Talpid3 PE=1 SV=1                                               |
| 112 | p04723 | TBA8_MOUSE   | Tuba8    | -4,29694 | 126.22057 | -2,22659 | 0.036598884 | QLHPQEILITGK            | Tubulin alpha-8 chain OS=Mus musculus GN=Tuba8 PE=1 SV=1                                           |
| 113 | p00760 | TBD4A_MOUSE  | Tubba4a  | -5,76779 | 145.12171 | -2,76191 | 0.025010880 | YLTAAVFR                | Tubulin beta-4A chain OS=Mus musculus GN=Tubba4a PE=1 SV=3                                         |
| 114 | p01922 | TERA_MOUSE   | Vcp      | -6,58926 | 115.75448 | -3,02652 | 0.007188948 | GLLYGPPPTGK             | Transitional endoplasmic reticulum ATPase OS=Mus musculus GN=Vcp PE=1 SV=4                         |
| 115 | p06588 | TERA_MOUSE   | Vcp      | -5,00155 | 107.91094 | -2,71525 | 0.016063354 | WALSQSNPSALR            | Transitional endoplasmic reticulum ATPase OS=Mus musculus GN=Vcp PE=1 SV=4                         |
| 116 | p03355 | TKT_MOUSE    | Tkt      | -4,71784 | 101.44240 | -2,42580 | 0.026794412 | LGQSDPAPLQHGVQDIYQK     | Transketolase OS=Mus musculus GN=Tkt PE=1 SV=1                                                     |
| 117 | p02446 | TKT_MOUSE    | Tkt      | -4,54961 | 70.71485  | -2,22514 | 0.040937220 | IALLGDTR                | Transketolase OS=Mus musculus GN=Tkt PE=1 SV=1                                                     |
| 118 | p05228 | TKT_MOUSE    | Tkt      | 3,44586  | 185.68523 | 2,56959  | 0.043423762 | SKDDGVTVYAGVYTHLEALAAES | Transketolase OS=Mus musculus GN=Tkt PE=1 SV=1                                                     |

|     | A      | B           | C      | D        | E         | F        | G           | H                  | I                                                                                   |
|-----|--------|-------------|--------|----------|-----------|----------|-------------|--------------------|-------------------------------------------------------------------------------------|
| 119 | p06144 | TLN1_MOUSE  | Tln1   | -3,40156 | 118.31330 | -2,38927 | 0.041010289 | VGGDPVAVWLK        | Talin-1 OS=Mus musculus GN=Tln1 PE=1 SV=2                                           |
| 120 | p01411 | TLN2_MOUSE  | Tln2   | -5,07328 | 58.30177  | -2,61716 | 0.035162598 | EVANSTANLVK        | Talin-2 OS=Mus musculus GN=Tln2 PE=1 SV=3                                           |
| 121 | p03339 | TPD52_MOUSE | Tpd52  | -3,57197 | 131.58990 | -2,19041 | 0.042843111 | LGSISSQFEK         | Tumor protein D52 OS=Mus musculus GN=Tpd52 PE=1 SV=2                                |
| 122 | p05892 | TSP1_MOUSE  | Thbs1  | -5,99335 | 135.09165 | -2,76600 | 0.028389362 | TIYVITLQDSIR       | Thrombospondin-1 OS=Mus musculus GN=Thbs1 PE=1 SV=1                                 |
| 123 | p04203 | UBC_MOUSE   | Ubc    | -6,32829 | 87.73204  | -3,07201 | 0.005604908 | MOIVFK             | Polyubiquitin-C OS=Mus musculus GN=Ubc PE=1 SV=2                                    |
| 124 | p06302 | VAT1_MOUSE  | Vat1   | -5,22864 | 91.91971  | -2,90915 | 0.010874302 | VLLVPGPEK          | Synaptic vesicle membrane protein VAT-1 homolog OS=Mus musculus GN=Vat1 PE=1 SV=3   |
| 125 | p00702 | VIME_MOUSE  | Vim    | -2,42628 | 66.98323  | -2,43263 | 0.033509489 | DDGVINETSQHHDDLE   | Vimentin OS=Mus musculus GN=Vim PE=1 SV=3                                           |
| 126 | p01187 | VP26A_MOUSE | Vps26a | -3,95940 | 105.01774 | -3,35981 | 0.015779092 | EITGIGPSTTTETETIAK | Vacuolar protein sorting-associated protein 26A OS=Mus musculus GN=Vps26a PE=1 SV=1 |
| 127 | p02298 | WDR1_MOUSE  | Wdr1   | -3,87118 | 47.63178  | -2,62349 | 0.034843743 | IAVVGEGR           | WD repeat-containing protein 1 OS=Mus musculus GN=Wdr1 PE=1 SV=3                    |
| 128 | p06892 | WDR1_MOUSE  | Wdr1   | -4,49047 | 129.10988 | -2,13972 | 0.044338552 | VYSILASTLK         | WD repeat-containing protein 1 OS=Mus musculus GN=Wdr1 PE=1 SV=3                    |

Supplementary Table 1. (continued)

|    | A      | B                 | C         | D         | E         | F           | G           | H                    | I                                                                                        |
|----|--------|-------------------|-----------|-----------|-----------|-------------|-------------|----------------------|------------------------------------------------------------------------------------------|
|    | db_ID  | UniProt_EntryName | Gene_Name | log2(FC)  | AveSignal | t-statistic | p.value     | Peptide              | Protein_Description                                                                      |
| 1  | p06759 | 1433B_MOUSE       | Ywhab     | -1.78402  | 52.58793  | -2.21177    | 0.036401704 | YLSEVASENKK          | 14-3-3 protein beta/alpha OS=Mus musculus GN=Ywhab PE=1 SV=3                             |
| 2  | p06718 | 1433E_MOUSE       | Ywhae     | -4.94857  | 91.21487  | -4.07894    | 0.000717391 | YLAEFATGNDK          | 14-3-3 protein epsilon OS=Mus musculus GN=Ywhae PE=1 SV=1                                |
| 3  | p06979 | 1433E_MOUSE       | Ywhae     | -2.87466  | 74.03284  | -3.17608    | 0.004233231 | EAAENSLVAYK          | 14-3-3 protein epsilon OS=Mus musculus GN=Ywhae PE=1 SV=1                                |
| 4  | p06648 | 1433E_MOUSE       | Ywhae     | -2.67412  | 74.70144  | -2.60467    | 0.016225713 | YDEHVESMK            | 14-3-3 protein epsilon OS=Mus musculus GN=Ywhae PE=1 SV=1                                |
| 5  | p06204 | 1433E_MOUSE       | Ywhae     | -3.02200  | 47.49620  | -2.21727    | 0.036384658 | HLPAANTGESK          | 14-3-3 protein epsilon OS=Mus musculus GN=Ywhae PE=1 SV=1                                |
| 6  | p04476 | 1433G_MOUSE       | Ywhah     | -2.53209  | 72.12524  | -3.04458    | 0.005973975 | NSVVEASEAYK          | 14-3-3 protein eta OS=Mus musculus GN=Ywhah PE=1 SV=2                                    |
| 7  | p04513 | 1433G_MOUSE       | Ywhah     | -3.74557  | 90.64788  | -3.28468    | 0.003265352 | NVTELEPLSNEER        | 14-3-3 protein gamma OS=Mus musculus GN=Ywhah PE=1 SV=2                                  |
| 8  | p04399 | 1433H_MOUSE       | Sfn       | -4.41175  | 105.35784 | -3.58949    | 0.002482170 | NLLSVAYK             | 14-3-3 protein sigma OS=Mus musculus GN=Sfn PE=1 SV=2                                    |
| 9  | p01613 | 1433Z_MOUSE       | Ywhaz     | -3.29595  | 100.16061 | -2.60678    | 0.016936840 | FLIPNASGPESK         | 14-3-3 protein zeta/delta OS=Mus musculus GN=Ywhaz PE=1 SV=1                             |
| 10 | p05512 | 1433Z_MOUSE       | Ywhaz     | -1.65449  | 61.18041  | -2.34382    | 0.027386666 | SVTEGQALSNEER        | 14-3-3 protein zeta/delta OS=Mus musculus GN=Ywhaz PE=1 SV=1                             |
| 11 | p06719 | 1433Z_MOUSE       | Ywhaz     | -2.02942  | 63.96062  | -2.14600    | 0.044414766 | YLAEVAGDDK           | 14-3-3 protein zeta/delta OS=Mus musculus GN=Ywhaz PE=1 SV=1                             |
| 12 | p06721 | 1433Z_MOUSE       | Ywhaz     | -2.20935  | 51.27537  | -2.09924    | 0.046114546 | YLAEVAGDDKK          | 14-3-3 protein zeta/delta OS=Mus musculus GN=Ywhaz PE=1 SV=1                             |
| 13 | p00191 | 6PGD_MOUSE        | Pgd       | -4.88403  | 92.08933  | -3.75973    | 0.001162702 | AGQAVDDFIEK          | 6-phosphogluconate dehydrogenase, decarboxylating OS=Mus musculus GN=Pgd PE=1 SV=3       |
| 14 | p00229 | 6PGD_MOUSE        | Pgd       | -5.12285  | 100.86449 | -3.98866    | 0.006167639 | AFIOAJAAK            | 6-phosphogluconate dehydrogenase, decarboxylating OS=Mus musculus GN=Pgd PE=1 SV=3       |
| 15 | p02174 | 6PGD_MOUSE        | Pgd       | -6.33119  | 114.46576 | -4.49578    | 0.011737784 | HMLPANIQAQK          | 6-phosphogluconate dehydrogenase, decarboxylating OS=Mus musculus GN=Pgd PE=1 SV=3       |
| 16 | p06006 | 6PGD_MOUSE        | Pgd       | -3.34516  | 88.21334  | -2.36582    | 0.027753957 | VDDFLAEAK            | 6-phosphogluconate dehydrogenase, decarboxylating OS=Mus musculus GN=Pgd PE=1 SV=3       |
| 17 | p04259 | A2M_MOUSE         | A2m       | -4.51169  | 114.77785 | -2.21407    | 0.040075526 | MVSGFIPLKPTVK        | Alpha-2-macroglobulin-P OS=Mus musculus GN=A2m PE=2 SV=2                                 |
| 18 | p02233 | ACTBL_MOUSE       | Actb12    | -3.07554  | 49.54558  | -2.44146    | 0.025957403 | HQVMVMGMGQK          | Beta-actin-like protein 2 OS=Mus musculus GN=Actb12 PE=1 SV=1                            |
| 19 | p02234 | ACTBL_MOUSE       | Actb12    | -2.96352  | 48.97585  | -2.25737    | 0.036749916 | HQVMVMGMGQK          | Beta-actin-like protein 2 OS=Mus musculus GN=Actb12 PE=1 SV=1                            |
| 20 | p04623 | ACTG_MOUSE        | Actg1     | -2.53173  | 65.83073  | -2.98046    | 0.006359747 | QEYDESGPSIVHR        | Actin, cytoplasmic 2 OS=Mus musculus GN=Actg1 PE=1 SV=1                                  |
| 21 | p02156 | ACTG_MOUSE        | Actg1     | -2.55417  | 73.13892  | -2.92636    | 0.007226119 | GYSFITTAER           | Actin, cytoplasmic 2 OS=Mus musculus GN=Actg1 PE=1 SV=1                                  |
| 22 | p04622 | ACTG_MOUSE        | Actg1     | -2.42095  | 66.05454  | -2.75318    | 0.011358391 | QEYDESGPSIVHR        | Actin, cytoplasmic 2 OS=Mus musculus GN=Actg1 PE=1 SV=1                                  |
| 23 | p04621 | ACTG_MOUSE        | Actg1     | -1.61693  | 84.42025  | -2.26244    | 0.032664348 | QEYDESGPSIVHR        | Actin, cytoplasmic 2 OS=Mus musculus GN=Actg1 PE=1 SV=1                                  |
| 24 | p03045 | ACTN1_MOUSE       | Actn1     | -4.51630  | 109.89988 | -3.04442    | 0.008266194 | LAILGHNEYSK          | Alpha-actinin-1 OS=Mus musculus GN=Actn1 PE=1 SV=1                                       |
| 25 | p04124 | ACTN1_MOUSE       | Actn1     | -2.89387  | 119.56354 | -2.59523    | 0.023616488 | MLDAEDVGTARPEK       | Alpha-actinin-1 OS=Mus musculus GN=Actn1 PE=1 SV=1                                       |
| 26 | p07182 | ACTS_MOUSE        | Acta1     | -2.43244  | 69.38335  | -2.63015    | 0.014706279 | EITALAPSTMK          | Actin, alpha skeletal muscle OS=Mus musculus GN=Acta1 PE=1 SV=1                          |
| 27 | p01183 | ACTS_MOUSE        | Acta1     | -2.93148  | 91.16393  | -2.61054    | 0.015094627 | EITALAPSTMK          | Actin, alpha skeletal muscle OS=Mus musculus GN=Acta1 PE=1 SV=1                          |
| 28 | p06778 | ACTS_MOUSE        | Acta1     | -3.29018  | 137.29575 | -2.17646    | 0.048761100 | YPIEHGIITNWDDMEK     | Actin, alpha skeletal muscle OS=Mus musculus GN=Acta1 PE=1 SV=1                          |
| 29 | p01467 | ANX_MOUSE         | Ina       | -2.76624  | 108.04476 | -2.76189    | 0.012305562 | EYQDLLNVK            | Alpha-internexin OS=Mus musculus GN=Ina PE=1 SV=3                                        |
| 30 | p05345 | AKR1A1_MOUSE      | Akr1a1    | -4.78376  | 96.64329  | -2.61485    | 0.016233541 | SPAQILLR             | Alcohol dehydrogenase [NADP+] OS=Mus musculus GN=Akr1a1 PE=1 SV=3                        |
| 31 | p02193 | AKR1A1_MOUSE      | Akr1a1    | -3.24957  | 49.49514  | -2.56713    | 0.018016635 | HPHDEVPALR           | Alcohol dehydrogenase [NADP+] OS=Mus musculus GN=Akr1a1 PE=1 SV=3                        |
| 32 | p01964 | AKR1A1_MOUSE      | Akr1a1    | -2.69917  | 122.98782 | -2.38362    | 0.027242061 | GLEVTAYSLGSSDR       | Alcohol dehydrogenase [NADP+] OS=Mus musculus GN=Akr1a1 PE=1 SV=3                        |
| 33 | p05344 | AKCL2_MOUSE       | Akr1e2    | -4.78376  | 96.64329  | -2.61485    | 0.016233541 | SPAQILLR             | 1,5-anhydro-D-fructose reductase OS=Mus musculus GN=Akr1e2 PE=1 SV=1                     |
| 34 | p02212 | AL3A2_MOUSE       | Aldh3a2   | -3.95411  | 109.19023 | -2.67348    | 0.020453126 | HLTPVTLEGGK          | Fatty aldehyde dehydrogenase OS=Mus musculus GN=Aldh3a2 PE=1 SV=2                        |
| 35 | p00056 | ALDOA_MOUSE       | Aldoa     | -4.53586  | 92.43587  | -3.05445    | 0.005473093 | ADGRPPFPOVVK         | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                      |
| 36 | p01918 | ALDOA_MOUSE       | Aldoa     | -4.63298  | 89.00265  | -3.21479    | 0.005842858 | GILADESTGSIK         | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                      |
| 37 | p04531 | ALDOA_MOUSE       | Aldoa     | -4.12109  | 67.95854  | -3.02015    | 0.007081122 | PHPHYALTPEQK         | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                      |
| 38 | p01273 | ALDOA_MOUSE       | Aldoa     | -2.20783  | 43.71517  | -2.21530    | 0.036128748 | ELSDIAHR             | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                      |
| 39 | p02412 | ALDOA_MOUSE       | Aldoa     | -3.63271  | 168.83214 | -2.25840    | 0.039398288 | IGEHPTSALAIMENANVLAR | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                      |
| 40 | p03769 | ALDOA_MOUSE       | Aldoa     | -1.48660  | 52.86032  | -2.15367    | 0.041158377 | LQSIGTENTEENRR       | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                      |
| 41 | p03770 | ALDOA_MOUSE       | Aldoa     | -1.88103  | 46.47046  | -2.14400    | 0.042431578 | LQSIGTENTEENRR       | Fructose-bisphosphate aldolase A OS=Mus musculus GN=Aldoa PE=1 SV=2                      |
| 42 | p06243 | ALDOB_MOUSE       | Aldob     | -2.74732  | 59.33084  | -2.21846    | 0.035886521 | VLAAYVK              | Annexin A1 OS=Mus musculus GN=Anxa1 PE=1 SV=2                                            |
| 43 | p00276 | ANXA1_MOUSE       | Anxa1     | -6.04788  | 103.19305 | -4.79025    | 0.001030391 | ALDLELK              | Annexin A1 OS=Mus musculus GN=Anxa1 PE=1 SV=2                                            |
| 44 | p05784 | ANXA1_MOUSE       | Anxa1     | -2.57564  | 93.05821  | -2.96344    | 0.007720844 | TPAQFDADLR           | Annexin A1 OS=Mus musculus GN=Anxa1 PE=1 SV=2                                            |
| 45 | p04309 | APOE_MOUSE        | ApoE      | -22.19466 | 72.81929  | -15.05079   | 0.000000002 | NEVHTMLQGSTEIR       | Apolipoprotein E OS=Mus musculus GN=ApoE PE=1 SV=2                                       |
| 46 | p03714 | APOE_MOUSE        | ApoE      | -4.92855  | 96.51014  | -3.63721    | 0.001904059 | LQAEFQAR             | Apolipoprotein E OS=Mus musculus GN=ApoE PE=1 SV=2                                       |
| 47 | p04310 | APOE_MOUSE        | ApoE      | -3.65804  | 96.46549  | -2.83217    | 0.010342481 | NEVHTMLQGSTEIR       | Apolipoprotein E OS=Mus musculus GN=ApoE PE=1 SV=2                                       |
| 48 | p05547 | APOE_MOUSE        | ApoE      | -2.63960  | 71.12006  | -2.73495    | 0.012813105 | TANLGAGAAQPLR        | Apolipoprotein E OS=Mus musculus GN=ApoE PE=1 SV=2                                       |
| 49 | p03352 | APOE_MOUSE        | ApoE      | -2.80457  | 88.85990  | -2.43673    | 0.023870062 | LGPLVEQGR            | Apolipoprotein E OS=Mus musculus GN=ApoE PE=1 SV=2                                       |
| 50 | p01218 | APOE_MOUSE        | ApoE      | -1.75194  | 101.65972 | -2.11121    | 0.048995593 | ELEEQLGPVETTR        | Apolipoprotein E OS=Mus musculus GN=ApoE PE=1 SV=2                                       |
| 51 | p04475 | ARCB1_MOUSE       | Arcp1b    | -3.84947  | 94.09629  | -3.56111    | 0.001973724 | NSVSGISVLGGK         | Actin-related protein 2/3 complex subunit 1B OS=Mus musculus GN=Arcp1b PE=1 SV=4         |
| 52 | p03495 | ARCB1_MOUSE       | Arcp1b    | -2.63241  | 65.50715  | -2.33971    | 0.031119686 | ASSEGGAATGAGLDSLHK   | Actin-related protein 2/3 complex subunit 1B OS=Mus musculus GN=Arcp1b PE=1 SV=4         |
| 53 | p02747 | ARPC4_MOUSE       | Arcp4     | -3.81589  | 111.62260 | -2.46146    | 0.031846265 | IVAEFFLK             | Actin-related protein 2/3 complex subunit 4 OS=Mus musculus GN=Arcp4 PE=1 SV=3           |
| 54 | p06284 | ARPC5_MOUSE       | Arcp5     | -2.98346  | 95.22647  | -2.13587    | 0.046020588 | ALAAGGVGSIVR         | Actin-related protein 2/3 complex subunit 5 OS=Mus musculus GN=Arcp5 PE=1 SV=3           |
| 55 | p06072 | B2M_MOUSE         | B2m       | -2.69647  | 94.21887  | -2.78021    | 0.011259704 | REMSDADFSGK          | Beta-2-microglobulin OS=Mus musculus GN=B2m PE=1 SV=2                                    |
| 56 | p04950 | B4GALT5_MOUSE     | B4galt5   | -5.12767  | 137.63393 | -2.96098    | 0.032574566 | RELLRLRPR            | Beta-1,4-galactosyltransferase 5 OS=Mus musculus GN=B4galt5 PE=2 SV=2                    |
| 57 | p06784 | BGH3_MOUSE        | Tgfb1     | -6.07846  | 106.93092 | -3.97960    | 0.002674289 | DLNLNHLK             | Transforming growth factor-beta-induced protein ig-h3 OS=Mus musculus GN=Tgfb1 PE=1 SV=1 |

Supplementary Table 2.

Differentially expressed peptides identified in *Pik3ca*<sup>RBD/-</sup> vs *Pik3ca*<sup>WT/-</sup> secretome of BMDMs phagocytosing apoptotic cells



|     | A      | B            | C       | D         | E         | F        | G           | H                 | I                                                                                        |
|-----|--------|--------------|---------|-----------|-----------|----------|-------------|-------------------|------------------------------------------------------------------------------------------|
| 59  | p02534 | BGH3_MOUSE   | Tgfb1   | -3.22916  | 91.78027  | -2.92801 | 0.009447494 | ILGDPEARL         | Transforming growth factor-beta-induced protein ig-h3 OS=Mus musculus GN=Tgfb1 PE=1 SV=1 |
| 60  | p05299 | BGH3_MOUSE   | Tgfb1   | -3.15890  | 55.25223  | -2.47681 | 0.043070585 | SLQGDGVVSSK       | Transforming growth factor-beta-induced protein ig-h3 OS=Mus musculus GN=Tgfb1 PE=1 SV=1 |
| 61  | p03725 | BLVRB_MOUSE  | Blvrb   | -3.56425  | 56.68616  | -2.76974 | 0.015161193 | LQDVTDDHIR        | Flavin reductase (NADPH) OS=Mus musculus GN=Blvrb PE=1 SV=3                              |
| 62  | p02283 | BLVRB_MOUSE  | Blvrb   | -6.43583  | 101.75248 | -2.77262 | 0.017038042 | IAIFGATGR         | Flavin reductase (NADPH) OS=Mus musculus GN=Blvrb PE=1 SV=3                              |
| 63  | p03702 | BLVRB_MOUSE  | Blvrb   | -2.80611  | 93.79123  | -2.38096 | 0.025988024 | LPSEGQPGRAVHVGDVR | Flavin reductase (NADPH) OS=Mus musculus GN=Blvrb PE=1 SV=3                              |
| 64  | p03726 | BLVRB_MOUSE  | Blvrb   | -2.50905  | 56.08433  | -2.27151 | 0.031618446 | LQDVTDDHIR        | Flavin reductase (NADPH) OS=Mus musculus GN=Blvrb PE=1 SV=3                              |
| 65  | p06402 | C170B_MOUSE  | Cep170b | -2.34826  | 64.89792  | -2.26848 | 0.047038634 | VPEALX            | Complement C1q subcomponent subunit A OS=Mus musculus GN=C1qa PE=1 SV=2                  |
| 66  | p04945 | C1QA_MOUSE   | C1qa    | -2.96963  | 71.15993  | -2.40305 | 0.030844289 | RGDEVIEKDPK       | Complement C1q subcomponent subunit A OS=Mus musculus GN=C1qa PE=1 SV=2                  |
| 67  | p04781 | C1QA_MOUSE   | C1qa    | -3.25673  | 161.21739 | -2.27788 | 0.036938255 | QNPMTLGNVIFDK     | Complement C1q subcomponent subunit A OS=Mus musculus GN=C1qa PE=1 SV=2                  |
| 68  | p06225 | C1QB_MOUSE   | C1qb    | -1.79175  | 60.91096  | -2.34380 | 0.029073670 | NTNANENVEPR       | Complement C1q subcomponent subunit B OS=Mus musculus GN=C1qb PE=1 SV=2                  |
| 69  | p04893 | C1QC_MOUSE   | C1qc    | -3.14080  | 72.81010  | -2.41369 | 0.026180105 | QVSGGVGLLR        | Complement C1q subcomponent subunit C OS=Mus musculus GN=C1cq PE=1 SV=2                  |
| 70  | p00699 | CALM_MOUSE   | Calml1  | -2.42614  | 101.49865 | -2.70739 | 0.014102471 | DNGYSAXELR        | Calmodulin OS=Mus musculus GN=Calml1 PE=1 SV=2                                           |
| 71  | p02831 | CAP1_MOUSE   | Cap1    | -9.41797  | 104.08797 | -3.03607 | 0.029298392 | KEPALLELEK        | Adenylyl cyclase-associated protein 1 OS=Mus musculus GN=Cap1 PE=1 SV=4                  |
| 72  | p06413 | CAP2_MOUSE   | Cap2    | -2.72076  | 77.34416  | -2.79951 | 0.010784241 | VPTISINK          | Adenylyl cyclase-associated protein 2 OS=Mus musculus GN=Cap2 PE=1 SV=1                  |
| 73  | p00753 | CAPG_MOUSE   | Capg    | -5.15897  | 103.73530 | -2.48752 | 0.000451191 | DLALAIR           | Macrophage-capping protein OS=Mus musculus GN=Capg PE=1 SV=2                             |
| 74  | p02168 | CATB_MOUSE   | Ctcb    | -3.02404  | 51.66206  | -2.34417 | 0.029051040 | HEADGMMGGHAIR     | Cathepsin B OS=Mus musculus GN=Ctcb PE=1 SV=2                                            |
| 75  | p06480 | CATD_MOUSE   | Ctcd    | -2.60598  | 117.14097 | -2.11768 | 0.046385096 | VSLPTTYLTK        | Cathepsin D OS=Mus musculus GN=Ctcd PE=1 SV=1                                            |
| 76  | p01809 | CATZ_MOUSE   | Ctaz    | -2.92733  | 106.42594 | -2.99538 | 0.007810845 | GDOLALLGR         | Cathepsin Z OS=Mus musculus GN=Ctaz PE=1 SV=1                                            |
| 77  | p04509 | CATZ_MOUSE   | Ctaz    | -2.14611  | 73.56134  | -2.49845 | 0.020907716 | NVNGVNYASVTR      | Cathepsin Z OS=Mus musculus GN=Ctaz PE=1 SV=1                                            |
| 78  | p05934 | CATZ_MOUSE   | Ctaz    | -5.34704  | 108.29607 | -2.50599 | 0.037085145 | TYPRRHEYLSPADLPK  | Cathepsin Z OS=Mus musculus GN=Ctaz PE=1 SV=1                                            |
| 79  | p06146 | CATZ_MOUSE   | Ctaz    | -1.63263  | 57.37968  | -2.07839 | 0.048150555 | VDGYGSLSGR        | Cathepsin Z OS=Mus musculus GN=Ctaz PE=1 SV=1                                            |
| 80  | p03549 | CAZA1_MOUSE  | Capza1  | -4.00091  | 133.31643 | -2.96840 | 0.009866219 | LLINDNLLLR        | F-actin-capping protein subunit alpha-1 OS=Mus musculus GN=Capza1 PE=1 SV=4              |
| 81  | p02516 | CBPC4_MOUSE  | Aghl1   | -3.44246  | 62.57933  | -3.80858 | 0.011705640 | ILEDIR            | Monocyte differentiation antigen CD14 OS=Mus musculus GN=Cd14 PE=1 SV=1                  |
| 82  | p02009 | CD14_MOUSE   | Cd14    | -4.80809  | 69.61987  | -3.77054 | 0.003748351 | GNPFLDSESHSEK     | Complement factor B OS=Mus musculus GN=Cfb PE=1 SV=2                                     |
| 83  | p05410 | CFAB_MOUSE   | Cfb     | -2.41188  | 86.96799  | -2.93192 | 0.008002775 | SDDAVVTEK         | Complement factor B OS=Mus musculus GN=Cfb PE=1 SV=2                                     |
| 84  | p04641 | CFAB_MOUSE   | Cfb     | -2.52250  | 79.67581  | -2.88783 | 0.008842270 | STGWSLDLQTR       | Complement factor B OS=Mus musculus GN=Cfb PE=1 SV=2                                     |
| 85  | p00309 | CFAB_MOUSE   | Cfb     | -2.96764  | 69.32049  | -2.54414 | 0.019381761 | ALFVSECKK         | Complement factor B OS=Mus musculus GN=Cfb PE=1 SV=2                                     |
| 86  | p03658 | CFAB_MOUSE   | Cfb     | -4.77847  | 87.78475  | -2.25765 | 0.034814188 | LQNGYSAEIR        | Complement factor B OS=Mus musculus GN=Cfb PE=1 SV=2                                     |
| 87  | p02766 | CLH1_MOUSE   | Clhc    | -4.71991  | 10.08561  | -3.10002 | 0.021760986 | VLDSNVSEHR        | Claudin heavy chain 1 OS=Mus musculus GN=Clhc PE=1 SV=3                                  |
| 88  | p04426 | CLH1_MOUSE   | Clhc    | -2.74538  | 130.32308 | -2.46641 | 0.026309636 | NLAGAEELFAR       | Claudin heavy chain 1 OS=Mus musculus GN=Clhc PE=1 SV=3                                  |
| 89  | p02135 | CLIC1_MOUSE  | Clic1   | -4.64918  | 105.06739 | -4.05110 | 0.010392484 | GVTFTVTVDTK       | Chloride intracellular channel protein 1 OS=Mus musculus GN=Clic1 PE=1 SV=3              |
| 90  | p00325 | CLUS_MOUSE   | Ciu     | -4.62606  | 79.69258  | -4.18690 | 0.004285371 | IDSLESDR          | Clusterin OS=Mus musculus GN=Ciu PE=1 SV=1                                               |
| 91  | p04689 | CLUS_MOUSE   | Ciu     | -3.27112  | 154.32708 | -2.22512 | 0.048141725 | ASQDITLFDOR       | Clusterin OS=Mus musculus GN=Ciu PE=1 SV=1                                               |
| 92  | p05868 | CMKP2_MOUSE  | Cmpk2   | -4.76052  | 103.11605 | -4.28402 | 0.001326950 | TLTLOSVESELK      | UMP-CMP kinase 2, mitochondrial OS=Mus musculus GN=Cmpk2 PE=1 SV=2                       |
| 93  | p06340 | CO3_MOUSE    | C3      | -3.18535  | 49.73706  | -2.38820 | 0.029723905 | VLVVTGSGNR        | Complement C3 OS=Mus musculus GN=C3 PE=1 SV=3                                            |
| 94  | p02430 | CO3_MOUSE    | C3      | -2.88025  | 48.14074  | -2.42886 | 0.032010158 | IGQGEVEVK         | Complement C3 OS=Mus musculus GN=C3 PE=1 SV=3                                            |
| 95  | p05877 | CO3_MOUSE    | C3      | -3.45744  | 73.38719  | -2.42309 | 0.032348183 | TVAMHTLDPK        | Complement C3 OS=Mus musculus GN=C3 PE=1 SV=3                                            |
| 96  | p01335 | CO3_MOUSE    | C3      | -3.03029  | 86.71632  | -2.23161 | 0.034094404 | EQALILK           | Complement C3 OS=Mus musculus GN=C3 PE=1 SV=3                                            |
| 97  | p06638 | COF2_MOUSE   | Cf2     | -2.04667  | 97.70002  | -2.21212 | 0.036374469 | YALVDATYETK       | Cofilin-2 OS=Mus musculus GN=Cf2 PE=1 SV=1                                               |
| 98  | p06126 | COF2_MOUSE   | Cf2     | -2.65252  | 38.98716  | -2.19459 | 0.039655488 | VFNDDMK           | Cofilin-2 OS=Mus musculus GN=Cf2 PE=1 SV=1                                               |
| 99  | p06159 | COR1C_MOUSE  | Coro1c  | -9.33812  | 100.82697 | -8.26275 | 0.002306740 | GVGVAWHPTAR       | Coronin-1C OS=Mus musculus GN=Coro1c PE=1 SV=2                                           |
| 100 | p01216 | COTL1_MOUSE  | Cotl1   | -3.93733  | 106.13154 | -2.81318 | 0.013197038 | ELEEDFIR          | Coadactin-like protein OS=Mus musculus GN=Cotl1 PE=1 SV=3                                |
| 101 | p05887 | CXCL16_MOUSE | Cxcl16  | -3.30823  | 106.16795 | -2.24563 | 0.048900502 | TSSGTGIPQGLTDHIR  | C-X-C motif chemokine 16 OS=Mus musculus GN=Cxcl16 PE=1 SV=2                             |
| 102 | p03636 | CYC_MOUSE    | Cysa    | -3.18116  | 66.43035  | -2.54457 | 0.030821767 | TGQAGGSYTDANK     | Cytochrome c, somatic OS=Mus musculus GN=Cysa PE=1 SV=2                                  |
| 103 | p04691 | CYTB_MOUSE   | Catb    | -4.31383  | 131.46925 | -2.58969 | 0.017573186 | QIVAGTNLFK        | Cystatin-B OS=Mus musculus GN=Catb PE=1 SV=1                                             |
| 104 | p04139 | CYTC_MOUSE   | Cat3    | -2.15597  | 76.21185  | -2.50011 | 0.020832971 | MLGAPEADANEEGVRR  | Cystatin-C OS=Mus musculus GN=Cat3 PE=1 SV=2                                             |
| 105 | p04135 | CYTC_MOUSE   | Cat3    | -1.62012  | 74.45510  | -2.39501 | 0.026598772 | MLGAPEADANEEGVRR  | Cystatin-C OS=Mus musculus GN=Cat3 PE=1 SV=2                                             |
| 106 | p06069 | DESM_MOUSE   | Des     | -2.46983  | 78.87907  | -2.81348 | 0.009435647 | VELQELNDR         | Desmin OS=Mus musculus GN=Des PE=1 SV=3                                                  |
| 107 | p05053 | DPS11_MOUSE  | Dpsyl1  | -62.87375 | 164.52708 | -2.45452 | 0.028272750 | VASAVASAVAR       | Dihydropyrimidine-related protein 2 OS=Mus musculus GN=Dpsyl2 PE=1 SV=2                  |
| 108 | p06128 | DPSYL2_MOUSE | Dpsyl2  | -2.56425  | 109.40214 | -2.56425 | 0.033883083 | VNLYPYR           | Dihydropyrimidine-related protein 2 OS=Mus musculus GN=Dpsyl2 PE=1 SV=2                  |
| 109 | p02552 | DX39B_MOUSE  | Ddx39b  | -2.61300  | 48.76373  | -2.77324 | 0.013651111 | ILNDVQDR          | Spliceosome RNA helicase Ddx39b OS=Mus musculus GN=Ddx39b PE=1 SV=1                      |
| 110 | p06844 | EF1A1_MOUSE  | Eef1a1  | -4.12524  | 107.77769 | -3.65806 | 0.002362711 | YYVTIDAPGHR       | Elongation factor 1-alpha OS=Mus musculus GN=Eef1a1 PE=1 SV=3                            |
| 111 | p06845 | EF1A1_MOUSE  | Eef1a1  | -2.96104  | 107.14322 | -2.43344 | 0.022441406 | YYVTIDAPGHR       | Elongation factor 1-alpha OS=Mus musculus GN=Eef1a1 PE=1 SV=3                            |
| 112 | p05628 | EF1A1_MOUSE  | Eef1a1  | -4.39136  | 44.36988  | -2.41229 | 0.034733185 | YEENIK            | Elongation factor 1-alpha OS=Mus musculus GN=Eef1a1 PE=1 SV=3                            |
| 113 | p02417 | EF1A2_MOUSE  | Eef1a2  | -2.29403  | 90.30546  | -2.45421 | 0.01618801  | IGGIGTVPSVGR      | Elongation factor 1-alpha OS=Mus musculus GN=Eef1a2 PE=1 SV=1                            |
| 114 | p04727 | EF1A2_MOUSE  | Eef1a2  | -4.94584  | 70.68335  | -2.84723 | 0.014845479 | QLIVGVNK          | Elongation factor 1-alpha OS=Mus musculus GN=Eef1a2 PE=1 SV=1                            |
| 115 | p03694 | EF1A2_MOUSE  | Eef1a2  | -2.85549  | 96.43270  | -2.28750 | 0.032716810 | LPLDDVYK          | Elongation factor 1-alpha OS=Mus musculus GN=Eef1a2 PE=1 SV=1                            |
| 116 | p00315 | EF1G_MOUSE   | Eef1g   | -3.09868  | 62.65829  | -2.53090 | 0.023173962 | ALIAAQYGAQVR      | Elongation factor 1-gamma OS=Mus musculus GN=Eef1g PE=1 SV=3                             |

|     | A      | B           | C       | D        | E         | F        | G             | H                      | I                                                                              |
|-----|--------|-------------|---------|----------|-----------|----------|---------------|------------------------|--------------------------------------------------------------------------------|
| 117 | p06134 | EF2_MOUSE   | Eef2    | -5.47746 | 101.58858 | -3.86808 | 0.001250482   | VFSGVVSTGLK            | Elongation factor 2 OS=Mus musculus GN=Eef2 PE=1 SV=2                          |
| 118 | p02820 | EF2_MOUSE   | Eef2    | -6.73345 | 77.97263  | -2.68055 | 0.021596989   | KEDLYLKPIOR            | Elongation factor 2 OS=Mus musculus GN=Eef2 PE=1 SV=2                          |
| 119 | p06367 | EF2_MOUSE   | Eef2    | -2.18443 | 105.16843 | -2.15770 | 0.04267630605 | VNFYTVYDVR             | Elongation factor 2 OS=Mus musculus GN=Eef2 PE=1 SV=2                          |
| 120 | p03798 | EFHD2_MOUSE | Eefhd2  | -3.81234 | 92.12205  | -3.22893 | 0.024174872   | LEISIDVTEGVK           | EF-hand domain-containing protein D2 OS=Mus musculus GN=Eefhd2 PE=1 SV=1       |
| 121 | p03618 | ENO4_MOUSE  | Eno1    | -3.67675 | 95.59045  | -3.06763 | 0.005303620   | LMIEDMGTEK             | Alpha-enolase OS=Mus musculus GN=Eno1 PE=1 SV=3                                |
| 122 | p02405 | ENO4_MOUSE  | Eno1    | -3.29383 | 67.25741  | -2.50531 | 0.019154702   | IGAEVYHNLLK            | Alpha-enolase OS=Mus musculus GN=Eno1 PE=1 SV=3                                |
| 123 | p01016 | ENO4_MOUSE  | Eno1    | -7.04595 | 99.74000  | -2.64930 | 0.021383514   | EALLELLK               | Alpha-enolase OS=Mus musculus GN=Eno1 PE=1 SV=3                                |
| 124 | p02133 | ENO4_MOUSE  | Eno1    | -2.37495 | 51.13751  | -2.11197 | 0.049046352   | GVSGVAEHNNK            | Alpha-enolase OS=Mus musculus GN=Eno1 PE=1 SV=3                                |
| 125 | p06653 | ENO4_MOUSE  | Eno2    | -3.69828 | 96.10529  | -3.82381 | 0.000877856   | YDLDFK                 | Gamma-enolase OS=Mus musculus GN=Eno2 PE=1 SV=2                                |
| 126 | p01238 | ENPL_MOUSE  | Hsp90b1 | -2.82702 | 92.99985  | -2.58492 | 0.018227678   | ELISNASDALDK           | Endoplasmic reticulum protein OS=Mus musculus GN=Hsp90b1 PE=1 SV=2             |
| 127 | p05559 | EPCR_MOUSE  | Procr   | -2.14024 | 70.76069  | -2.41957 | 0.023001308   | TAAVVWSQGEPSK          | Endothelial protein C receptor OS=Mus musculus GN=Procr PE=1 SV=3              |
| 128 | p01997 | EPPL_MOUSE  | Eppk1   | -3.76146 | 111.72495 | -2.38976 | 0.026381458   | GLVGLLEK               | Epilplakin OS=Mus musculus GN=Eppk1 PE=1 SV=2                                  |
| 129 | p02129 | FKBP1_MOUSE | Fkbp1a  | -1.81219 | 74.00769  | -2.30332 | 0.030102003   | GVGVETIPQDGR           | Peptidyl-prolyl cis-trans isomerase FKBP1A OS=Mus musculus GN=Fkbp1a PE=1 SV=2 |
| 130 | p01715 | FLNA_MOUSE  | Flna    | -3.85949 | 102.03047 | -2.92616 | 0.003046247   | GTAGTGLALAVEGPEAK      | Filamin-A OS=Mus musculus GN=Flna PE=1 SV=5                                    |
| 131 | p06490 | FLNA_MOUSE  | Flna    | -2.27785 | 77.38693  | -2.80382 | 0.010383572   | VTADGGLPEFSGNANK       | Filamin-A OS=Mus musculus GN=Flna PE=1 SV=5                                    |
| 132 | p01041 | FLNA_MOUSE  | Flna    | -2.45944 | 76.28744  | -2.69098 | 0.012549004   | AFPGGLGGNAGSPAR        | Filamin-A OS=Mus musculus GN=Flna PE=1 SV=5                                    |
| 133 | p01002 | FLNA_MOUSE  | Flna    | -4.01208 | 102.97692 | -2.74278 | 0.015982742   | EAGAGGLAAVEGSPK        | Filamin-A OS=Mus musculus GN=Flna PE=1 SV=5                                    |
| 134 | p03901 | FLNA_MOUSE  | Flna    | -2.61403 | 102.97280 | -2.60386 | 0.017858541   | LTYSLSQESGLK           | Filamin-A OS=Mus musculus GN=Flna PE=1 SV=5                                    |
| 135 | p06076 | FLNA_MOUSE  | Flna    | -2.23963 | 79.75902  | -2.54620 | 0.018451428   | VTGGLCADNSVGR          | Filamin-A OS=Mus musculus GN=Flna PE=1 SV=5                                    |
| 136 | p01640 | FLNA_MOUSE  | Flna    | -4.32562 | 153.83900 | -2.85374 | 0.023204894   | FNEEHPDPSFVPVAVSPSGDAR | Filamin-A OS=Mus musculus GN=Flna PE=1 SV=5                                    |
| 137 | p06016 | FLNB_MOUSE  | Flnb    | -3.11789 | 122.93197 | -2.20846 | 0.049645153   | GDGELLAVQITDGEKPR      | Filamin-B OS=Mus musculus GN=Flnb PE=1 SV=3                                    |
| 138 | p01763 | G3P_MOUSE   | Gapdh   | -3.07285 | 76.45923  | -2.99211 | 0.007866543   | GAAGNIPASTGAAK         | Glyceraldehyde 3-phosphate dehydrogenase OS=Mus musculus GN=Gapdh PE=1 SV=2    |
| 139 | p06215 | G3P_MOUSE   | Gapdh   | -4.02775 | 68.42285  | -2.95854 | 0.049800033   | VPENLKK                | Glyceraldehyde 3-phosphate dehydrogenase OS=Mus musculus GN=Gapdh PE=1 SV=2    |
| 140 | p03993 | GDI1_MOUSE  | Gdi1    | -3.27229 | 61.33179  | -2.96724 | 0.009144431   | LYSELNKK               | Rab GDP-dissociation inhibitor alpha OS=Mus musculus GN=Gdi1 PE=1 SV=3         |
| 141 | p05571 | GDIR1_MOUSE | Arhgdia | -4.47737 | 91.12964  | -3.37305 | 0.003416564   | TDYMWGSGPR             | Rho GDP-dissociation inhibitor 1 OS=Mus musculus GN=Arhgdia PE=1 SV=3          |
| 142 | p00901 | GELS_MOUSE  | Gsn     | -3.12865 | 55.81121  | -2.57406 | 0.019795971   | DSQEEKTEALTSK          | Gelsolin OS=Mus musculus GN=Gsn PE=1 SV=3                                      |
| 143 | p02970 | GLRX1_MOUSE | Glrx    | -5.33550 | 134.09504 | -2.40165 | 0.04383241    | KTQEILSQLPKF           | Glutaredoxin-1 OS=Mus musculus GN=Glrx PE=1 SV=3                               |
| 144 | p05877 | GNPMB_MOUSE | Gnpmb   | -2.83813 | 68.89721  | -2.64041 | 0.015348080   | AIYPSIK                | Transmembrane glycoprotein NMB OS=Mus musculus GN=Gnpmb PE=1 SV=2              |
| 145 | p04449 | BIP_MOUSE   | Hspa5   | -3.45163 | 104.34677 | -3.45651 | 0.003983405   | NGLTSGPENTVFDK         | 78 kDa glucose-regulated protein OS=Mus musculus GN=Hspa5 PE=1 SV=3            |
| 146 | p02725 | HSPA5_MOUSE | Hspa5   | -3.33011 | 39.74239  | -2.77912 | 0.003821963   | ITTDNDNR               | 78 kDa glucose-regulated protein OS=Mus musculus GN=Hspa5 PE=1 SV=3            |
| 147 | p00665 | GSTM1_MOUSE | Gstm1   | -3.94078 | 89.49407  | -3.16945 | 0.006408767   | ADIVENQVMDTR           | Glutathione S-transferase Mu 1 OS=Mus musculus GN=Gstm1 PE=1 SV=2              |
| 148 | p06698 | GSTM1_MOUSE | Gstm1   | -3.70752 | 111.45089 | -2.41917 | 0.042416683   | VIATPFSK               | Glutathione S-transferase Mu 1 OS=Mus musculus GN=Gstm1 PE=1 SV=2              |
| 149 | p02052 | GSTO1_MOUSE | Gsto1   | -1.99845 | 72.08230  | -3.26205 | 0.031303150   | GSAPPQVPVEGQIR         | Glutathione S-transferase omega-1 OS=Mus musculus GN=Gsto1 PE=1 SV=2           |
| 150 | p04155 | GSTP2_MOUSE | Gstp2   | -4.48819 | 106.33990 | -3.26708 | 0.014116523   | MLLADGGGSWK            | Glutathione S-transferase P 2 OS=Mus musculus GN=Gstp2 PE=1 SV=2               |
| 151 | p04072 | H12_MOUSE   | H1-2    | -5.56501 | 100.43318 | -2.28444 | 0.041579553   | ASGPVSELTK             | Histone H1.2 OS=Mus musculus GN=Hist1h12 PE=1 SV=2                             |
| 152 | p02063 | H1T_MOUSE   | H1-6    | -3.70967 | 71.88766  | -3.13548 | 0.014204953   | ALAAAGYDVEK            | Hist                                                                           |

|     | A      | B            | C        | D           | E         | F        | G           | H                 | I                                                                                   |
|-----|--------|--------------|----------|-------------|-----------|----------|-------------|-------------------|-------------------------------------------------------------------------------------|
| 175 | p03408 | INAR2_MOUSE  | Iftar2   | -4.07357    | 150.80767 | -3.44434 | 0.019180408 | LILSWELNKK        | Interferon alpha/beta receptor 2 OS=Mus musculus GN=Ifnar2 PE=1 SV=2                |
| 176 | p05312 | IPCS_MOUSE   | Ipcr5    | -123.467186 | -17.1615  | -0.17615 | 0.019180408 | LILSWELNKK        | Imprin-5 OS=Mus musculus GN=Ipcr5 PE=1 SV=3                                         |
| 177 | p05337 | IQGAP1_MOUSE | Iqgap1   | -3.93438    | 156.54796 | -2.26395 | 0.038980613 | SNOQLENDLMDIK     | Ras GTPase-activating-like protein IQGAP1 OS=Mus musculus GN=Iqgap1 PE=1 SV=2       |
| 178 | p06294 | IREB2_MOUSE  | Ireb2    | -6.12356    | 93.07646  | -4.14877 | 0.006532466 | VLEEAAYR          | Iron-responsive element-binding protein 2 OS=Mus musculus GN=Ireb2 PE=1 SV=2        |
| 179 | p00293 | KC10_MOUSE   | Krt10    | -2.06228    | 77.74877  | -2.40683 | 0.029562353 | ALEESNYELEKG      | Keratin, type I cytoskeletal 10 OS=Mus musculus GN=Krt10 PE=1 SV=4                  |
| 180 | p03215 | KC10_MOUSE   | Krt10    | -2.21032    | 88.93281  | -2.30532 | 0.037142148 | LENEIQTYR         | Keratin, type I cytoskeletal 10 OS=Mus musculus GN=Krt10 PE=1 SV=3                  |
| 181 | p02734 | KC10_MOUSE   | Krt13    | -3.19166    | 69.08462  | -2.62278 | 0.023033006 | ITMKNLNDNR        | Keratin, type I cytoskeletal 13 OS=Mus musculus GN=Krt13 PE=1 SV=2                  |
| 182 | p03084 | KC18_MOUSE   | Krt15    | -2.49317    | 56.34026  | -2.45981 | 0.020134573 | LASVLDK           | Keratin, type I cytoskeletal 15 OS=Mus musculus GN=Krt15 PE=1 SV=2                  |
| 183 | p04850 | KC18_MOUSE   | Krt18    | -5.40285    | 72.87465  | -3.50665 | 0.005782506 | QSVSDIHLRL        | Keratin, type I cytoskeletal 18 OS=Mus musculus GN=Krt18 PE=1 SV=5                  |
| 184 | p03154 | KC18_MOUSE   | Krt18    | -2.79152    | 81.76535  | -2.87725 | 0.017605640 | LEAEIATYR         | Keratin, type I cytoskeletal 18 OS=Mus musculus GN=Krt18 PE=1 SV=5                  |
| 185 | p02452 | KC18_MOUSE   | Krt18    | -3.44246    | 62.57933  | -3.60858 | 0.011705640 | IEDLR             | Keratin, type I cytoskeletal 18 OS=Mus musculus GN=Krt18 PE=1 SV=5                  |
| 186 | p03006 | KC18_MOUSE   | Krt18    | -3.57119    | 40.36469  | -2.50563 | 0.027628225 | KVVDQDNTIR        | Keratin, type I cytoskeletal 18 OS=Mus musculus GN=Krt18 PE=1 SV=5                  |
| 187 | p04849 | KC18_MOUSE   | Krt18    | -5.04432    | 71.92173  | -2.44660 | 0.030993176 | QSVSDIHLRL        | Keratin, type I cytoskeletal 18 OS=Mus musculus GN=Krt18 PE=1 SV=5                  |
| 188 | p01395 | KC18_MOUSE   | Krt18    | -2.13667    | 49.46444  | -2.31494 | 0.037794967 | ETMQDLNDR         | Keratin, type I cytoskeletal 18 OS=Mus musculus GN=Krt18 PE=1 SV=5                  |
| 189 | p00298 | KC19_MOUSE   | Krt19    | -2.88935    | 75.72268  | -3.07166 | 0.007820664 | ALEQANGELEVK      | Keratin, type I cytoskeletal 19 OS=Mus musculus GN=Krt19 PE=1 SV=1                  |
| 190 | p02770 | KC19_MOUSE   | Krt19    | -2.95454    | 92.13499  | -2.55108 | 0.021458785 | IVLOIDNAR         | Keratin, type I cytoskeletal 19 OS=Mus musculus GN=Krt19 PE=1 SV=1                  |
| 191 | p05280 | KC19_MOUSE   | Krt19    | -6.80407    | 107.19553 | -3.66237 | 0.022876547 | SLLGEAEHYNNLPPTK  | Keratin, type I cytoskeletal 19 OS=Mus musculus GN=Krt19 PE=1 SV=1                  |
| 192 | p05770 | KC21_MOUSE   | Krt1     | -3.31559    | 94.59955  | -3.06302 | 0.006439817 | TNAENEFVTK        | Keratin, type II cytoskeletal 1 OS=Mus musculus GN=Krt1 PE=1 SV=4                   |
| 193 | p04375 | KC25_MOUSE   | Krt5     | -2.33376    | 42.68853  | -2.51187 | 0.024059801 | KNYDEINKR         | Keratin, type II cytoskeletal 5 OS=Mus musculus GN=Krt5 PE=1 SV=1                   |
| 194 | p01606 | KC27_MOUSE   | Krt73    | -2.99541    | 90.32561  | -2.75794 | 0.014746608 | FLEQGNQVLQTK      | Keratin, type II cytoskeletal 73 OS=Mus musculus GN=Krt73 PE=1 SV=1                 |
| 195 | p00147 | KC28_MOUSE   | Krt8     | -2.97741    | 124.01242 | -3.07610 | 0.008040143 | ASLEAAIADEAQR     | Keratin, type II cytoskeletal 8 OS=Mus musculus GN=Krt8 PE=1 SV=4                   |
| 196 | p00083 | KC28_MOUSE   | Krt8     | -2.84441    | 76.52934  | -2.71838 | 0.016783488 | AEAEITVYQIK       | Keratin, type II cytoskeletal 8 OS=Mus musculus GN=Krt8 PE=1 SV=4                   |
| 197 | p05579 | KC28_MOUSE   | Krt8     | -3.94800    | 42.71343  | -2.37362 | 0.042069295 | TEISEMRN          | Keratin, type II cytoskeletal 8 OS=Mus musculus GN=Krt8 PE=1 SV=4                   |
| 198 | p01975 | KATL2_MOUSE  | Katnal2  | -3.11865    | 115.75448 | -2.22528 | 0.040925818 | GLLLYPPGTGK       | Katanin p60 ATPase-containing subunit A-like 2 OS=Mus musculus GN=Katnal2 PE=2 SV=2 |
| 199 | p01821 | KPYM_MOUSE   | Pkm      | -5.48083    | 88.35175  | -6.3712  | 0.002254349 | GDPYLEAVR         | Pyruvate kinase PKM OS=Mus musculus GN=Pkm PE=1 SV=4                                |
| 200 | p02727 | KPYM_MOUSE   | Pkm      | -2.71690    | 97.51797  | -2.73134 | 0.011672215 | ITLDNAYMEK        | Pyruvate kinase PKM OS=Mus musculus GN=Pkm PE=1 SV=4                                |
| 201 | p03118 | KPYM_MOUSE   | Pkm      | -2.44775    | 100.75845 | -2.59838 | 0.015519393 | LDDSPATIR         | Pyruvate kinase PKM OS=Mus musculus GN=Pkm PE=1 SV=4                                |
| 202 | p04040 | KPYM_MOUSE   | Pkm      | -2.98742    | 73.66449  | -2.55517 | 0.017118718 | QPIAVNTIR         | Pyruvate kinase PKM OS=Mus musculus GN=Pkm PE=1 SV=4                                |
| 203 | p02726 | KPYM_MOUSE   | Pkm      | -1.89146    | 79.60893  | -2.32585 | 0.035725816 | ITLDNAYMEK        | Pyruvate kinase PKM OS=Mus musculus GN=Pkm PE=1 SV=4                                |
| 204 | p03015 | KRT35_MOUSE  | Krt35    | -2.71782    | 62.52846  | -2.80882 | 0.014043205 | LAADDFR           | Pyruvate kinase PKM OS=Mus musculus GN=Pkm PE=1 SV=4                                |
| 205 | p03381 | KSKY_MOUSE   | Syk      | -4.04813    | 91.99532  | -2.94023 | 0.011596367 | LIATTAHEK         | Tyrosine-protein kinase SYK OS=Mus musculus GN=Syk PE=1 SV=2                        |
| 206 | p01875 | LASP1_MOUSE  | Lasp1    | -3.36569    | 120.21599 | -2.16514 | 0.049778892 | GFSVADTPELQR      | LIM and SH3 domain protein 1 OS=Mus musculus GN=Lasp1 PE=1 SV=1                     |
| 207 | p04898 | LDHA_MOUSE   | Ldha     | -2.45981    | 102.48687 | -2.42491 | 0.043032060 | QVYVLEAEVK        | L-lactate dehydrogenase A chain OS=Mus musculus GN=Ldha PE=1 SV=3                   |
| 208 | p05295 | LDHA_MOUSE   | Ldha     | -3.03393    | 97.69394  | -2.83653 | 0.028636216 | PELQETDADKEQWK    | L-lactate dehydrogenase A chain OS=Mus musculus GN=Ldha PE=1 SV=3                   |
| 209 | p05604 | LDHA_MOUSE   | Ldha     | -1.93738    | 62.59679  | -2.18936 | 0.038173853 | VLTLPPEAR         | L-lactate dehydrogenase A chain OS=Mus musculus GN=Ldha PE=1 SV=3                   |
| 210 | p03928 | LdHC_MOUSE   | Ldhc     | -3.53970    | 88.57748  | -2.49566 | 0.023204069 | LVITAGAR          | L-lactate dehydrogenase C chain OS=Mus musculus GN=Ldhc PE=1 SV=2                   |
| 211 | p03670 | LEG1_MOUSE   | Lgals1   | -2.46707    | 52.35011  | -2.24855 | 0.033652300 | LPDGHEFK          | Galectin-1 OS=Mus musculus GN=Lgals1 PE=1 SV=3                                      |
| 212 | p02007 | LEG3_MOUSE   | Lgals3   | -6.69476    | 92.98716  | -3.44358 | 0.002744794 | GNDAVFHFNPR       | Galectin-3 OS=Mus musculus GN=Lgals3 PE=1 SV=3                                      |
| 213 | p05994 | LEG3_MOUSE   | Lgals3   | -5.58633    | 86.05059  | -2.40310 | 0.040091053 | VAVYVALLLYNNHR    | Galectin-3 OS=Mus musculus GN=Lgals3 PE=1 SV=3                                      |
| 214 | p02662 | LEG3_MOUSE   | Lgals3   | -3.02703    | 110.62700 | -2.16099 | 0.040529677 | IOVLAEDHFK        | Galectin-3 OS=Mus musculus GN=Lgals3 PE=1 SV=3                                      |
| 215 | p05765 | LG3BP_MOUSE  | Lgals3bp | -4.97194    | 158.83830 | -3.66966 | 0.008248956 | TMQALEFTVPVPELAK  | Galectin-3-binding protein OS=Mus musculus GN=Lgals3bp PE=1 SV=1                    |
| 216 | p04957 | LG3BP_MOUSE  | Lgals3bp | -4.44566    | 74.26951  | -3.10808 | 0.008429848 | RIEVSMSVVK        | Galectin-3-binding protein OS=Mus musculus GN=Lgals3bp PE=1 SV=1                    |
| 217 | p04487 | LMNA_MOUSE   | Lmna     | -5.04272    | 89.42719  | -4.05920 | 0.001045830 | NSNLVGAHHEELQOSR  | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 218 | p04779 | LMNA_MOUSE   | Lmna     | -91.20368   | 58.02003  | -2.29984 | 0.030513796 | SLETEAGLR         | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 219 | p05538 | LMNA_MOUSE   | Lmna     | -3.00940    | 101.22930 | -2.76402 | 0.012847167 | TALNISTGEEVAMR    | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 220 | p00502 | LMNA_MOUSE   | Lmna     | -2.11498    | 70.72597  | -2.68123 | 0.014028212 | AAYEALGDAR        | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 221 | p04370 | LMNA_MOUSE   | Lmna     | -3.13524    | 76.83885  | -2.67009 | 0.018419516 | NIYSEELR          | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 222 | p01058 | LMNA_MOUSE   | Lmna     | -1.90549    | 64.21477  | -2.41548 | 0.026034548 | EDQLQELNR         | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 223 | p05262 | LMNA_MOUSE   | Lmna     | -2.16373    | 58.52003  | -2.29984 | 0.030513796 | SLETEAGLR         | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 224 | p04214 | LMNA_MOUSE   | Lmna     | -3.30505    | 166.03390 | -2.37432 | 0.045488448 | MOQLDEYQELLDIK    | Prelamin A/C OS=Mus musculus GN=Lmna PE=1 SV=2                                      |
| 225 | p01826 | MARCS_MOUSE  | Marcks   | -3.38875    | 46.75072  | -2.67646 | 0.023494877 | GEATAPRGEAAVSSPSK | Myristoylated alanine-rich C-kinase substrate OS=Mus musculus GN=Marcks PE=1 SV=2   |
| 226 | p03370 | MdHC_MOUSE   | Mdh1     | -4.00895    | 60.25191  | -4.22297 | 0.042168125 | LQVADDDVK         | Malate dehydrogenase, cytoplasmic OS=Mus musculus GN=Mdh1 PE=1 SV=3                 |
| 227 | p06391 | MdHM_MOUSE   | Mdh2     | -2.86262    | 79.00935  | -2.33476 | 0.028704936 | VNVPVIGGHAGK      | Malate dehydrogenase, mitochondrial OS=Mus musculus GN=Mdh2 PE=1 SV=3               |
| 228 | p02635 | MdHM_MOUSE   | Mdh2     | -2.13121    | 47.77924  | -2.28891 | 0.044676470 | IGEAETGVK         | Malate dehydrogenase, mitochondrial OS=Mus musculus GN=Mdh2 PE=1 SV=3               |
| 229 | p02431 | MdG_MOUSE    | Magohb   | -4.97216    | 89.84459  | -2.54687 | 0.044603277 | ISGLDYNQSK        | Protein mago nashi homolog 2 OS=Mus musculus GN=Magohb PE=2 SV=1                    |
| 230 | p00354 | MOES_MOUSE   | Men      | -2.97461    | 75.11937  | -2.72808 | 0.011571034 | ATSELANAR         | Moesin OS=Mus musculus GN=Men PE=1 SV=3                                             |
| 231 | p00434 | MOES_MOUSE   | Men      | -2.52802    | 64.86943  | -2.58000 | 0.016181599 | AMQVQEDLEK        | Moesin OS=Mus musculus GN=Men PE=1 SV=3                                             |
| 232 | p02703 | MOES_MOUSE   | Men      | -3.11494    | 66.06252  | -2.46731 | 0.023958814 | ISQLEMAR          | Moesin OS=Mus musculus GN=Men PE=1 SV=3                                             |

|     | A      | B           | C      | D         | E         | F        | G           | H                  | I                                                                             |
|-----|--------|-------------|--------|-----------|-----------|----------|-------------|--------------------|-------------------------------------------------------------------------------|
| 233 | p01144 | MYL3_MOUSE  | Myi3   | -3.141623 | 73.81483  | -2.72613 | 0.018563162 | EGNGTVMGAELR       | Myosin light chain 3 OS=Mus musculus GN=Myi3 PE=1 SV=4                        |
| 234 | p02263 | MYL6_MOUSE  | Myi6   | -7.33130  | 68.94790  | -3.03312 | 0.009707444 | HVLVTLEAK          | Myosin light polypeptide 6 OS=Mus musculus GN=Myi6 PE=1 SV=3                  |
| 235 | p01142 | MYL6_MOUSE  | Myi6   | -3.41623  | 73.81483  | -2.72613 | 0.018563162 | EGNGTVMGAELR       | Myosin light polypeptide 6 OS=Mus musculus GN=Myi6 PE=1 SV=3                  |
| 236 | p01996 | NDKA_MOUSE  | Nme1   | -3.85147  | 111.70147 | -2.40498 | 0.025542291 | GLVGEIK            | Nucleoside diphosphate kinase A OS=Mus musculus GN=Nme1 PE=1 SV=1             |
| 237 | p05609 | NDKB_MOUSE  | Nme2   | -3.49510  | 94.27497  | -2.63184 | 0.015639747 | FTIAIKPDGVQR       | Nucleoside diphosphate kinase B OS=Mus musculus GN=Nme2 PE=1 SV=1             |
| 238 | p04354 | NDKB_MOUSE  | Nme2   | -2.46029  | 55.46242  | -2.29505 | 0.031246264 | NIHSD3SVESAER      | Nucleoside diphosphate kinase B OS=Mus musculus GN=Nme2 PE=1 SV=1             |
| 239 | p00219 | NEXN_MOUSE  | Nexn   | -6.04788  | 103.19305 | -4.79025 | 0.010303931 | ADLEIK             | Nucleophosmin OS=Mus musculus GN=Nxm1 PE=1 SV=1                               |
| 240 | p01575 | NPM_MOUSE   | Npm1   | -3.94926  | 68.82489  | -2.82891 | 0.040160257 | NIYNYIK            | Nucleobindin-1 OS=Mus musculus GN=Nucb1 PE=1 SV=2                             |
| 241 | p03389 | NUCB1_MOUSE | Nucb1  | -3.10240  | 67.50553  | -3.13941 | 0.007360156 | VNVPVQAQALK        | Nucleobindin-1 OS=Mus musculus GN=Nucb1 PE=1 SV=2                             |
| 242 | p06732 | NUCB1_MOUSE | Nucb1  | -2.61106  | 76.14803  | -2.94645 | 0.008020054 | YLESGEOR           | Nucleobindin-1 OS=Mus musculus GN=Nucb1 PE=1 SV=2                             |
| 243 | p04940 | NUCB1_MOUSE | Nucb1  | -3.51237  | 62.90359  | -2.73339 | 0.012497532 | RFEELAAR           | Nucleobindin-1 OS=Mus musculus GN=Nucb1 PE=1 SV=2                             |
| 244 | p03836 | NUCB1_MOUSE | Nucb1  | -1.88344  | 54.13689  | -2.40157 | 0.030931497 | LSQETALQR          | Nucleobindin-1 OS=Mus musculus GN=Nucb1 PE=1 SV=2                             |
| 245 | p01992 | NUF1_MOUSE  | Nuf1   | -1.98954  | 70.36965  | -2.14534 | 0.047744551 | QSELETNFEETLK      | Nucleolin OS=Mus musculus GN=Nuf1 PE=1 SV=2                                   |
| 246 | p01136 | PA2G4_MOUSE | Pa2g4  | -6.06403  | 101.01561 | -3.21817 | 0.044603277 | ISGLDYNQSK         | Polliferation-associated protein 2G4 OS=Mus musculus GN=Pa2g4 PE=1 SV=3       |
| 247 | p05188 | PABP1_MOUSE | Pabpc1 | -5.07333  | 108.66393 | -3.16036 | 0.016340193 | SGVGNFIK           | Polyadenylation-binding protein 1 OS=Mus musculus GN=Pabpc1 PE=1 SV=2         |
| 248 | p05973 | PATFA_MOUSE | Pfatf  | -2.93649  | 72.24840  | -2.94325 | 0.008077737 | VAIDNTAK           | Platelet-activating factor acetylhydrolase OS=Mus musculus GN=Patf2 PE=2 SV=2 |
| 249 | p03996 | PATFA_MOUSE | Pfatf2 | -2.07246  | 64.15170  | -2.49180 | 0.021193903 | LYYPQVQDQR         | Platelet-activating factor acetylhydrolase OS=Mus musculus GN=Patf2 PE=2 SV=2 |
| 250 | p04345 | PATFA_MOUSE | Pfatf2 | -4.10205  | 131.05141 | -2.48486 | 0.039520634 | ALSADLDEGDPK       | Platelet-activating factor acetylhydrolase OS=Mus musculus GN=Patf2 PE=2 SV=2 |
| 251 | p01484 | PDI3_MOUSE  | Pdia3  | -3.91612  | 104.94463 | -3.97623 | 0.004237727 | FHTNLSLVK          | Protein disulfide-isomerase A3 OS=Mus musculus GN=Pdia3 PE=1 SV=2             |
| 252 | p01743 | PDI3_MOUSE  | Pdia3  | -3.91800  | 105.40147 | -3.21017 | 0.012707784 | FVMOEFESR          | Protein disulfide-isomerase A3 OS=Mus musculus GN=Pdia3 PE=1 SV=2             |
| 253 | p03066 | PDI3_MOUSE  | Pdia3  | -2.41746  | 63.82095  | -2.40951 | 0.030465842 | LAPEYEAATR         | Protein disulfide-isomerase A3 OS=Mus musculus GN=Pdia3 PE=1 SV=2             |
| 254 | p03677 | PDLT_MOUSE  | Pdlit  | -6.06177  | 104.33049 | -3.26381 | 0.032584403 | LPENPLIK           | Phosphoglycerate mutase 1 OS=Mus musculus GN=Pgam1 PE=1 SV=3                  |
| 255 | p01696 | PGLM1_MOUSE | Pgam1  | -3.91421  | 123.97364 | -2.45188 | 0.032389524 | FSGWYDADLSPAGHEEAK | Phosphoglycerate mutase 2 OS=Mus musculus GN=Pgam2 PE=1 SV=3                  |
| 256 | p03284 | PGLM2_MOUSE | Pgam2  | -7.41988  | 66.24074  | -4.98291 | 0.008297388 | VLUAHQSLR          | Phosphoglycerate kinase 1 OS=Mus musculus GN=Pgk1 PE=1 SV=4                   |
| 257 | p02645 | PGK1_MOUSE  | Pgk1   | -3.09753  | 132.85474 | -2.65243 | 0.016836891 | IQULNNLDK          | Phospholipase D4 OS=Mus musculus GN=Pld4 PE=1 SV=1                            |
| 258 | p01679 | PLD4_MOUSE  | Pld4   | -4.42425  | 111.65170 | -3.11118 | 0.011216170 | QFQLLR             | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 259 | p00418 | PLEC_MOUSE  | Plec   | -2.79042  | 75.87133  | -2.95303 | 0.009950221 | AOAEALQELQR        | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 260 | p02027 | PLEC_MOUSE  | Plec   | -4.32107  | 52.16458  | -3.04633 | 0.010236396 | AGVGPPELHEK        | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 261 | p01751 | PLEC_MOUSE  | Plec   | -4.49707  | 67.19625  | -2.95008 | 0.014739266 | QLQLAQEAQK         | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 262 | p06798 | PLEC_MOUSE  | Plec   | -3.54863  | 146.32893 | -2.82711 | 0.015407534 | YSSELTLSQYIK       | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 263 | p01884 | PLEC_MOUSE  | Plec   | -2.37928  | 71.90537  | -2.60463 | 0.017999072 | GGELYVTDTAR        | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 264 | p04977 | PLEC_MOUSE  | Plec   | -3.14376  | 34.41156  | -2.68950 | 0.018704187 | REEQAALHK          | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 265 | p04546 | PLEC_MOUSE  | Plec   | -2.66130  | 72.20927  | -2.53792 | 0.019197023 | QAEAEALR           | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 266 | p06231 | PLEC_MOUSE  | Plec   | -3.61889  | 136.72738 | -2.65143 | 0.021300032 | VYQSLGSAVAGGLK     | Plectin OS=Mus musculus GN=Plec PE=1 SV=3                                     |
| 267 | p04704 | PLEC_MOUSE  | Plec   | -2.07962  | 68.05262  | -2.50993 | 0.022801937 | QLAEEDLAQR         | Plectin OS=Mus mus                                                            |



|     | A      | B           | C        | D        | E         | F        | G           | H                  | I                                                                                             |
|-----|--------|-------------|----------|----------|-----------|----------|-------------|--------------------|-----------------------------------------------------------------------------------------------|
| 291 | p04721 | RADI_MOUSE  | Rdx      | -5.62475 | 102.05090 | -2.63084 | 0.022121476 | QLFDQVVK           | Radixin OS=Mus musculus GN=Rdx PE=1 SV=3                                                      |
| 292 | p00297 | RADI_MOUSE  | Rdx      | -2.30883 | 61.63237  | -2.33344 | 0.028013351 | ALELEOER           | Radixin OS=Mus musculus GN=Rdx PE=1 SV=3                                                      |
| 293 | p03937 | RAN_MOUSE   | Ran      | -3.13699 | 74.71141  | -2.64505 | 0.017739645 | LVLVGDGGTGK        | GTP-binding nuclear protein Ran OS=Mus musculus GN=Ran PE=1 SV=3                              |
| 294 | p06558 | RBI1B_MOUSE | Rab11b   | -5.38848 | 89.44255  | -2.66471 | 0.032839410 | VVLIGDSGVGK        | Ras-related protein Rab-11B OS=Mus musculus GN=Rab11b PE=1 SV=3                               |
| 295 | p03662 | RBBP7_MOUSE | Rbbp7    | -2.97984 | 130.23626 | -2.71435 | 0.035721399 | LNVWDLGK           | Histone-binding protein RBBP7 OS=Mus musculus GN=Rbbp7 PE=1 SV=1                              |
| 296 | p05337 | RENK_MOUSE  | Atg6ap2  | -3.82593 | 67.37101  | -3.00498 | 0.015090397 | SFGSVVFR           | Renin receptor OS=Mus musculus GN=Atg6ap2 PE=1 SV=2                                           |
| 297 | p01970 | RIF1_MOUSE  | Rif1     | -3.02394 | 62.15434  | -2.09968 | 0.046072559 | GLGQLIR            | Telomere-associated protein RIF1 OS=Mus musculus GN=Rif1 PE=1 SV=2                            |
| 298 | p02596 | RL7L_MOUSE  | Rpl7l1   | -3.02394 | 62.15434  | -2.09968 | 0.046072559 | INQLIR             |                                                                                               |
| 299 | p05837 | ROA3_MOUSE  | Rplp0    | -5.60565 | 175.94860 | -2.73916 | 0.029518949 | TSFFQALGITTK       | 60S acidic ribosomal protein P0 OS=Mus musculus GN=Rplp0 PE=1 SV=3                            |
| 300 | p03515 | RN180_MOUSE | Rnf180   | -3.72269 | 159.57688 | -2.22638 | 0.039102962 | LLFKASDPK          | E3 ubiquitin-protein ligase RNF180 OS=Mus musculus GN=Rnf180 PE=1 SV=2                        |
| 301 | p02368 | ROA3_MOUSE  | Hmnpa3   | -2.63936 | 103.62954 | -2.22785 | 0.039801830 | ITETVEIEDR         | Heterogeneous nuclear ribonucleoprotein A3 OS=Mus musculus GN=Hmnpa3 PE=1 SV=1                |
| 302 | p06311 | RS19_MOUSE  | Rps19    | -5.98115 | 113.28451 | -5.62832 | 0.001453984 | VLOALEGLK          | 40S ribosomal protein S19 OS=Mus musculus GN=Rps19 PE=1 SV=3                                  |
| 303 | p04083 | RS21_MOUSE  | Rps21    | -2.69923 | 76.20512  | -3.18641 | 0.004466875 | MGESDDSLR          | 40S ribosomal protein S21 OS=Mus musculus GN=Rps21 PE=1 SV=1                                  |
| 304 | p00111 | RS3_MOUSE   | Rps3     | -4.13065 | 115.24356 | -2.90724 | 0.045687123 | AELENEFLTR         | 40S ribosomal protein S3 OS=Mus musculus GN=Rps3 PE=1 SV=1                                    |
| 305 | p01473 | RSSA_MOUSE  | Rpsa     | -1.86185 | 74.36178  | -2.12508 | 0.044607934 | FAAATGATPIAGR      | 40S ribosomal protein SA OS=Mus musculus GN=Rpsa PE=1 SV=4                                    |
| 306 | p03662 | S10A6_MOUSE | S100a6   | -2.33270 | 66.23138  | -2.37313 | 0.026041751 | LMDDLDR            | Protein S100-A6 OS=Mus musculus GN=S100a6 PE=1 SV=3                                           |
| 307 | p00077 | SAA3_MOUSE  | Saa3     | -3.42563 | 90.54078  | -3.40337 | 0.002695239 | ADGFANWGR          | Serum amyloid A-3 protein OS=Mus musculus GN=Saa3 PE=1 SV=1                                   |
| 308 | p00850 | SAA3_MOUSE  | Saa3     | -4.31196 | 71.79133  | -2.67241 | 0.028692264 | DPNHFRPAGLPK       | Serum amyloid A-3 protein OS=Mus musculus GN=Saa3 PE=1 SV=1                                   |
| 309 | p02612 | SAP_MOUSE   | Psap     | 22.47654 | 65.88784  | 16.89663 | 0.000016609 | IPVEDMAR           | Prosaposin OS=Mus musculus GN=Psap PE=1 SV=2                                                  |
| 310 | p02613 | SAP_MOUSE   | Psap     | -2.50759 | 77.20331  | -2.53785 | 0.019199741 | IPVEDMAR           | Prosaposin OS=Mus musculus GN=Psap PE=1 SV=2                                                  |
| 311 | p03544 | SAP_MOUSE   | Psap     | -3.10218 | 65.82346  | -2.28594 | 0.036360064 | LLLGTEK            | Prosaposin OS=Mus musculus GN=Psap PE=1 SV=2                                                  |
| 312 | p06370 | SEPT8_MOUSE | Septin8  | -4.89958 | 137.86507 | -3.08575 | 0.022155101 | VNIPIRAK           | Septin-8 OS=Mus musculus GN=Sept8 PE=1 SV=1                                                   |
| 313 | p06581 | SH3L1_MOUSE | Sh3bgr1  | -2.02620 | 67.99412  | -2.12721 | 0.043504711 | VYIASAGSGTAIK      | SH3 domain-binding glutamic acid-rich-like protein OS=Mus musculus GN=Sh3bgr1 PE=1 SV=1       |
| 314 | p04428 | SHPS1_MOUSE | Sirpa    | -4.27550 | 93.75550  | -3.69742 | 0.002180445 | NNMDFSIR           | Tyrosine-protein phosphatase non-receptor type substrate 1 OS=Mus musculus GN=Sirpa PE=1 SV=1 |
| 315 | p00347 | SNX2_MOUSE  | Snx2     | -4.99249 | 91.90760  | -3.63922 | 0.004646325 | ALNSOLAEVEEK       | Sorting nexin-2 OS=Mus musculus GN=Snx2 PE=1 SV=2                                             |
| 316 | p00707 | SODC_MOUSE  | Sod1     | -1.63271 | 76.31442  | -2.26812 | 0.035261862 | DGVANVSIEDR        | Superoxide dismutase [Cu-Zn] OS=Mus musculus GN=Sod1 PE=1 SV=2                                |
| 317 | p02261 | SODC_MOUSE  | Sod1     | -2.40190 | 54.46354  | -2.17744 | 0.040524640 | HVGLDGNTAGK        | Superoxide dismutase [Cu-Zn] OS=Mus musculus GN=Sod1 PE=1 SV=2                                |
| 318 | p02097 | SPB6_MOUSE  | Serpinb6 | -2.66762 | 126.55736 | -2.48367 | 0.023825213 | GTTASQMAQALADK     | Serpin B6 OS=Mus musculus GN=Serpinb6 PE=1 SV=1                                               |
| 319 | p04888 | SYNE3_MOUSE | Syne3    | -3.02394 | 62.15434  | -2.09968 | 0.046072559 | QVQLLR             |                                                                                               |
| 320 | p04326 | TAGL2_MOUSE | Tagln2   | -1.81520 | 59.65370  | -2.35964 | 0.026455861 | NFSDNQLQEGK        | Transgelin-2 OS=Mus musculus GN=Tagln2 PE=1 SV=4                                              |
| 321 | p00949 | TBA3_MOUSE  | Tuba3a   | -3.94385 | 94.57577  | -3.12024 | 0.008215264 | DVNAAIATIK         | Tubulin alpha-3 chain OS=Mus musculus GN=Tuba3a PE=1 SV=1                                     |
| 322 | p01062 | TBA4A_MOUSE | Tuba4a   | -3.20382 | 54.37042  | -2.59373 | 0.019686734 | EDMAALEK           | Tubulin alpha-4A chain OS=Mus musculus GN=Tuba4a PE=1 SV=1                                    |
| 323 | p01660 | TBB1_MOUSE  | Tubb1    | -2.89800 | 106.17752 | -2.92352 | 0.008156524 | FGPOLNADLR         | Tubulin beta-1 chain OS=Mus musculus GN=Tubb1 PE=1 SV=1                                       |
| 324 | p02713 | TBB5_MOUSE  | Tubb5    | -2.47240 | 79.55134  | -2.23850 | 0.043520586 | ISVYNEATGKG        | Tubulin beta-5 chain OS=Mus musculus GN=Tubb5 PE=1 SV=1                                       |
| 325 | p03095 | TCPB_MOUSE  | Cct2     | -6.32664 | 104.92853 | -3.78341 | 0.005343633 | LAVEAVLR           | T-complex protein 1 subunit beta OS=Mus musculus GN=Cct2 PE=1 SV=4                            |
| 326 | p03940 | TCPO_MOUSE  | Cct8     | -4.11424 | 104.83273 | -2.66547 | 0.022183348 | LVPGGGATEIELAK     | T-complex protein 1 subunit theta OS=Mus musculus GN=Cct8 PE=1 SV=3                           |
| 327 | p05943 | TCPO_MOUSE  | Cct8     | -5.04706 | 78.98807  | -2.62336 | 0.049619420 | VADIALHYANK        | T-complex protein 1 subunit theta OS=Mus musculus GN=Cct8 PE=1 SV=3                           |
| 328 | p02847 | TERA_MOUSE  | Vcp      | -7.77575 | 90.44112  | -4.36114 | 0.001465500 | KGDFILVR           | Transitional endoplasmic reticulum ATPase OS=Mus musculus GN=Vcp PE=1 SV=4                    |
| 329 | p01922 | TERA_MOUSE  | Vcp      | -3.11865 | 115.75448 | -2.22528 | 0.040929818 | GILLYPGPGTGK       | Transitional endoplasmic reticulum ATPase OS=Mus musculus GN=Vcp PE=1 SV=4                    |
| 330 | p06588 | TERA_MOUSE  | Vcp      | -2.88294 | 107.91094 | -2.20104 | 0.043963154 | WALSSQNPALSR       | Transitional endoplasmic reticulum ATPase OS=Mus musculus GN=Vcp PE=1 SV=4                    |
| 331 | p03409 | TITIN_MOUSE | Ttn      | -3.10218 | 65.46006  | -2.36080 | 0.032332643 | LILTEGK            | Titin OS=Mus musculus GN=Ttn PE=1 SV=1                                                        |
| 332 | p02446 | TKT_MOUSE   | Tkt      | -4.12963 | 70.71485  | -3.39265 | 0.003755586 | IHALGGDTK          | Transketolase OS=Mus musculus GN=Tkt PE=1 SV=1                                                |
| 333 | p03098 | TKT_MOUSE   | Tkt      | -2.57655 | 64.00280  | -2.43306 | 0.023611150 | LAVSQVPR           | Transketolase OS=Mus musculus GN=Tkt PE=1 SV=1                                                |
| 334 | p02507 | TKT_MOUSE   | Tkt      | -3.04038 | 139.90204 | -2.53749 | 0.027838819 | ILATTPQEDAPSDIANIR | Transketolase OS=Mus musculus GN=Tkt PE=1 SV=1                                                |
| 335 | p02875 | TKT_MOUSE   | Tkt      | -2.91125 | 57.24536  | -2.24925 | 0.039061330 | KISSDLGHVPVK       | Transketolase OS=Mus musculus GN=Tkt PE=1 SV=1                                                |
| 336 | p03464 | TLN1_MOUSE  | Tln1     | -4.40194 | 162.29866 | -3.25066 | 0.008865765 | LLAALLEDGNGRPLQAQK | Talin-1 OS=Mus musculus GN=Tln1 PE=1 SV=2                                                     |
| 337 | p02259 | TPIS_MOUSE  | Tpi1     | -4.26355 | 92.92296  | -2.65759 | 0.017292973 | HVFGESEDLGKQ       | Triosephosphate isomerase OS=Mus musculus GN=Tpi1 PE=1 SV=4                                   |
| 338 | p04963 | TPM2_MOUSE  | Tpm2     | -6.46774 | 105.22339 | -4.03476 | 0.003698954 | ROLVEEELDR         | Tropomyosin beta chain OS=Mus musculus GN=Tpm2 PE=1 SV=1                                      |
| 339 | p00011 | TPM3_MOUSE  | Tpm3     | -2.08266 | 73.42371  | -3.02699 | 0.005842977 | ADADEAEVASLNR      | Tropomyosin alpha-3 chain OS=Mus musculus GN=Tpm3 PE=1 SV=3                                   |
| 340 | p05604 | TR168_MOUSE | Trim68   | -4.12385 | 71.69103  | -3.17194 | 0.004129146 | TFADVDR            |                                                                                               |
| 341 | p01249 | U2AF2_MOUSE | U2af2    | -3.43475 | 145.98405 | -2.96557 | 0.013008369 | ELLTSGPLK          | Splicing factor U2AF 65 kDa subunit OS=Mus musculus GN=U2af2 PE=1 SV=3                        |
| 342 | p06207 | UBE2V_MOUSE | Ube2v2   | -4.65531 | 92.39200  | -3.75170 | 0.002461951 | VILQELR            | Ubiquitin-conjugating enzyme E2 variant 2 OS=Mus musculus GN=Ube2v2 PE=1 SV=4                 |
| 343 | p00121 | UBA1_MOUSE  | Uba1     | -2.19778 | 65.48175  | -2.28899 | 0.034570313 | AENYDISPADR        | Ubiquitin-like modifier-activating enzyme 1 OS=Mus musculus GN=Uba1 PE=1 SV=1                 |
| 344 | p06745 | UBC_MOUSE   | Ubc      | -2.61702 | 66.71835  | -3.21420 | 0.003604753 | TLSDYNIQK          | Polyubiquitin-C OS=Mus musculus GN=Ubc PE=1 SV=2                                              |
| 345 | p01377 | UBC_MOUSE   | Ubc      | -4.87984 | 100.13700 | -2.67058 | 0.014361942 | ESTLHLVLR          | Polyubiquitin-C OS=Mus musculus GN=Ubc PE=1 SV=2                                              |
| 346 | p04203 | UBC_MOUSE   | Ubc      | -3.28498 | 67.79204  | -2.13473 | 0.044244418 | MQIFVK             | Polyubiquitin-C OS=Mus musculus GN=Ubc PE=1 SV=2                                              |
| 347 | p05772 | UBE2N_MOUSE | Ube2n    | -1.85689 | 56.10898  | -2.48118 | 0.020539135 | TNEAQAIETAR        | Ubiquitin-conjugating enzyme E2 N OS=Mus musculus GN=Ube2n PE=1 SV=1                          |
| 348 | p06302 | VAT1_MOUSE  | Vat1     | -5.30929 | 91.91971  | -4.15433 | 0.000863248 | VLLVGPGEK          | Synaptic vesicle membrane protein VAT-1 homolog OS=Mus musculus GN=Vat1 PE=1 SV=3             |

|     | A      | B           | C      | D        | E         | F        | G           | H               | I                                                                                 |
|-----|--------|-------------|--------|----------|-----------|----------|-------------|-----------------|-----------------------------------------------------------------------------------|
| 349 | p01489 | VIME_MOUSE  | Vim    | -2.94706 | 67.65907  | -2.93252 | 0.007120883 | FANYIDK         | Vimentin OS=Mus musculus GN=Vim PE=1 SV=3                                         |
| 350 | p03720 | VIME_MOUSE  | Vim    | -2.60550 | 67.74121  | -2.83425 | 0.008986034 | LDEIQNMK        | Vimentin OS=Mus musculus GN=Vim PE=1 SV=3                                         |
| 351 | p01073 | VIME_MOUSE  | Vim    | -3.86348 | 89.00676  | -2.89556 | 0.010118630 | EAEASTQSFR      | Vimentin OS=Mus musculus GN=Vim PE=1 SV=3                                         |
| 352 | p01477 | VIME_MOUSE  | Vim    | -2.31600 | 73.58086  | -2.69319 | 0.012485930 | FADLSEANR       | Vimentin OS=Mus musculus GN=Vim PE=1 SV=3                                         |
| 353 | p00702 | VIME_MOUSE  | Vim    | -2.99191 | 66.98323  | -2.91792 | 0.014158672 | DGVOINETSQHDDLE | Vimentin OS=Mus musculus GN=Vim PE=1 SV=3                                         |
| 354 | p03729 | VIME_MOUSE  | Vim    | -2.05100 | 66.82180  | -2.10815 | 0.047286817 | LQEEMLQR        | Vimentin OS=Mus musculus GN=Vim PE=1 SV=3                                         |
| 355 | p00269 | VINC_MOUSE  | Vim    | -4.14077 | 101.28537 | -2.61990 | 0.022570675 | ALASQLQDSLK     | Vinculin OS=Mus musculus GN=Vim PE=1 SV=4                                         |
| 356 | p04237 | VINC_MOUSE  | Vcl    | -2.62025 | 107.39300 | -2.40182 | 0.033623725 | MTGLVDEADTK     | Vinculin OS=Mus musculus GN=Vcl PE=1 SV=4                                         |
| 357 | p05612 | VPS18_MOUSE | Vps18  | -4.60381 | 70.69222  | -2.94684 | 0.012349315 | TLFSSPR         |                                                                                   |
| 358 | p02572 | VPS35_MOUSE | Vps35  | -4.10564 | 106.93690 | -3.76292 | 0.009774472 | ILVGTNLVR       | Vacuolar protein sorting-associated protein 35 OS=Mus musculus GN=Vps35 PE=1 SV=1 |
| 359 | p02298 | WDR1_MOUSE  | Wdr1   | -2.92870 | 47.63178  | -2.79666 | 0.027196253 | IAVVGEGR        | WD repeat-containing protein 1 OS=Mus musculus GN=Wdr1 PE=1 SV=3                  |
| 360 | p04343 | ZN329_MOUSE | Znf329 | -3.02394 | 62.15434  | -2.09968 | 0.046072559 | NIAGLIR         |                                                                                   |

Supplementary Table 2. (continued)



University of Salamanca according to current European (2007/526/CE) and Spanish (RD 1201/2005 and RD53/2013) legislation. All experiments were approved by the Bioethics Committee of the Cancer Research Center.

## Isolation, culture and treatments of BMDM

Bone marrow cells from tibias and femurs of 12–14-week-old *Pik3ca*<sup>RBD/Lox</sup> mice and *Pik3ca*<sup>WT/Lox</sup> littermates were cultured with DMEM supplemented with 10% FBS, 100 units/ml penicillin, 100 µg/mL streptomycin, 2 mM L-Glutamine and 20 ng/mL M-CSF for 7 days. 4-hydroxytamoxifen (Sigma Aldrich) (100 nM) was added to culture media on day 3 to eliminate *Pik3ca*-Lox allele. The differentiated BMDM were then detached using cell dissociation buffer (C5914-100, Merck) and cultured in DMEM supplemented with 10% FBS, 100 µg/mL streptomycin, 2 mM L-Glutamine, 20 ng/mL M-CSF for unstimulated BMDMs. For macrophage polarization towards an inflammatory phenotype 20 ng/ml IFN $\gamma$  (Peprotech) and 100 ng/ml LPS (Sigma Aldrich) was added to culture media.

## Generation of chimeric animals

Chimeric mice exhibiting WT or RBD deficient leukocytes were generated by lethal irradiation with 5.5 Gy twice, 4 h apart of LysM GFP recipient animals (mice exhibiting endogenously GFP fluorescent monocytes and neutrophils) followed by an injection of bone marrow cells ( $1.5 \times 10^6$  cells/recipient i.v.) from C57BL/6 WT or RBD donor mice. Chimerism was then assessed 4 weeks later by flow cytometry from blood samples (reconstitution of  $99.8 \pm 0.2$  % and  $99.8 \pm 0.1$  % for WT and RBD deficient donor cells, respectively; n =5 mice per group).

## Neutrophil depletion

Neutrophil depletion of chimeric mice was induced by intraperitoneal injection of anti-GR1 25µg/mouse/day for 3 days). Numbers of blood circulating monocytes and neutrophils were quantified by flow cytometry pre- and post-depletion. Neutrophils were found to be reduced by 99.5%, whilst this anti GR1-depleting protocol had no effect on blood monocyte proportion (n=5 mice/group).

## Brightfield intravital confocal microscopy

Mesenteric inflammation was induced following intraperitoneal injection of mouse recombinant CCL2 (500ng/mouse in 500uL of PBS). Six hours later, anesthetized chimeric mice (150 mg/kg ketamine, 7.5 mg/kg xylazine, i.p.) were placed in supine position on a heating pad (37 °C) for maintenance of body temperature. The mesenteric vascular bed was exteriorized, placed on a purpose-built stage of an upright brightfield microscope (Zeiss Axioskop). Mesenteries were superfused with warmed (37 °C) Tyrode's solution (Sigma). After a 5-min equilibration period, analysis of leukocyte-endothelium interactions was made in at least 9 (and up to 16) randomly selected segments (100 µm in length) of post-capillary venules (20–40 µm in diameter) for each mouse. Leukocyte rolling was quantified by counting the number of rolling cells passing a fixed transversal line in the middle of the vessel segment for 5 min. Leukocyte adhesion (stationary position of the cell for 30 s or longer) was quantified along a 100 µm vessel length and data were normalized as the number of cells per 500µm vessel segments. Leukocyte extravasation response was quantified within 50 µm on either side of the 100 µm vessel segment in the perivenular tissue; and data were normalized as the number of extravasated leukocytes per mm<sup>2</sup> of extravascular tissue. At the end of the analysis period, mice were humanely killed by cervical dislocation.

## BMDM elastic modulus

The elastic modulus of BMDMs was quantitatively assessed utilizing Atomic Force Microscopy (AFM). Specifically, cells were cultured on cover glass slides, which were subsequently positioned in a specialized BIO-AFM setup integrated with an inverted optical microscope (Nikon TE2000).

The BIO-AFM was equipped with a V-shaped, four-sided pyramidal silicon nitride tip (Bruker AFM Probes) to facilitate accurate force measurements. To avoid localized variations and ensure data representativeness, no more than three cells were selected from a single field of view for mechanical characterization, and these cells were never contiguous. Subsequently, the stiffness of each individual cell was characterized through measurements obtained at three perinuclear points. For each point, force-displacement (F vs. z) curves were captured (10  $\mu\text{m}$  amplitude at a speed of 5  $\mu\text{m/s}$ ). The determination of the elastic modulus was performed based on the analysis of force-displacement curves. A total of five curves were recorded for each perinuclear point at an indentation depth of 300 nm. Data were analysed by fitting the curves to the Hertz model as previously described<sup>75</sup>,<sup>76</sup>. Finally, the elastic modulus for each cellular population under distinct experimental conditions was calculated based on a minimum of fifteen measurements per independent cell, with each condition comprising at least ten independent cells ( $n \geq 10$ ).

## Immunofluorescence

BMDM were fixed using 4% paraformaldehyde, permeabilized with 0.1% Triton X-100 and blocked for 1 h with 3% BSA in PBS before incubation with the primary antibodies, used at a 1:100 dilution: Lamp1 (#553792, BD Pharmingen), Deoxyribonuclease I-Alexa Fluor™ 488 Conjugate (#D12371, Invitrogen, 1:2000). To stain actin cytoskeleton, Alexa Fluor™ 647 Phalloidin (Invitrogen, 1:10 000) was directly added to the primary antibody mixture. Alexa Fluor 488- or Alexa Fluor 555-conjugated secondary antibodies (Invitrogen) were used to detect the indicated proteins at a 1:1000 dilution. Cells were counterstained with DAPI on the mounting solution (ProLong Gold Antifade Reagent with DAPI, Invitrogen). Images were taken using a Zeiss LSM510 confocal microscope or Leica DM6 B THUNDER Imager 3D Tissue.

## Transwell migration assay

Transwell migration assays were carried out using the 6.5 mm Transwell® with 8.0  $\mu\text{m}$  Pore Polyester Membrane Insert (Corning).  $9 \times 10^4$  MEFs from wild type mice were used as a chemo-attractant to encourage macrophage migration.  $8 \times 10^5$  BMDM were seeded in the transwell. Transwells were performed following the manufacturer instructions.

## Random migration assay

For random migration assays, BMDMs were seeded in 24-well plates coated with matrigel (0.5mg/ml) and labelled using CellTracker™ Red CMTPX Dye 1  $\mu\text{M}$  (ThermoFisher) for 30 minutes. 24 hours later LPS + IFN- $\gamma$  was added when necessary. Triplicates of each condition and genotype were prepared. Time-lapse imaging was carried out for 24 h. One image was taken every 10 min within the same well using a Nikon microscope driven by Metamorph (Molecular Devices, Chicago, IL, USA). A total of 80–100 cells per condition were tracked using the Fiji plugin Trackmate. Pre-processing was done using Mexican hat filter 3.0 radius to increase particle detection. Images were segmented using the fluorescence channel with the Laplacian of the Gaussian detector with a 30  $\mu\text{m}$  estimated particle diameter, a 10.0 threshold and median filter option selected. Segmented objects were linked from frame to frame with a Linear Assignment Problem (LAP) tracker with 45  $\mu\text{m}$  frame-to-frame linking distance and 2 frame gap closure. Criteria for track acceptance were track duration at least the 90% of the video. Tracks were visually inspected for completeness and accuracy of the tracking.

## Secretome mass spectrometry

Samples for secretome analysis were prepared as previously described<sup>130</sup>. In brief, 100  $\mu\text{g}$  of proteins were digested into peptides using trypsin and peptides were desalted using Oasis HLB extraction cartridges (Waters UK Ltd) and eluted with 50% acetonitrile (ACN) in 0.1% Trifluoroacetic acid (TFA).

Dried peptides were dissolved in 0.1% TFA and analysed by nano ACQUITY liquid chromatography (Waters Corp., Milford, MA, USA) coupled on-line to a tandem LTQ Orbitrap XL, mass spectrometer (Thermo Fisher Scientific)<sup>131</sup>. Gradient elution was from 5% to 25% buffer B in 180 min at a flow rate 300nL/min with buffer A being used to balance the mobile phase (buffer A was 0.1% formic acid in water and B was 0.1% formic acid in ACN). The mass spectrometer was controlled by Xcalibur software and operated in the positive mode. The spray voltage was 1.95 kV and the capillary temperature was set to 200 °C. The LTQ Orbitrap XL was operated in data dependent mode with one survey MS scan followed by 5 MS/MS scans. Label-free quantitative proteomics analysis was performed using three independent biological samples per group. Additionally, each sample was analysed in technical duplicates. To ensure robust quantitative analysis, we utilized LTQ Orbitrap XL tandem mass spectrometry (MS/MS) to generate six distinct mass spectral profiles from each group.

MS raw files were converted into Mascot Generic Format using Mascot Distiller (version 2.3.0) and searched against the SwissProt database (release December 2015) restricted to human entries using the Mascot search daemon (version 2.3.1). Allowed mass windows were 10 ppm and 600 mmu for parent and fragment mass to charge values, respectively. Variable modifications included in searches were oxidation of methionine, pyro-glu (N-term) and phosphorylation of serine, threonine and tyrosine.

Spectral counting quantification method relies on the number of times peptides are identified by tandem mass spectrometry (with expectancy value <0.05) from a given protein. Spectral counts were obtained from Mascot result (DAT) files using a python script written in house in the Mascot Parser Toolkit environment (version 2.4.x).

## Proteomic data analysis

The proteomic data obtained consisted of 6,844 peptides and 30 samples: Pik3ca<sup>WT/-</sup> and Pik3ca<sup>RBD/-</sup> BMDMs in steady state conditions (labelled as cell samples: WT/- and RBD/-), phagocytosing apoptotic cells (labelled as cell samples: WT/-Phag and RBD/-Phag), and the apoptotic LKR10 cells alone (labelled as LKR). For each of these samples, proteomic experiments were performed with 6 replicates (3 biological replicates x 2 technical replicates), yielding a data set of 30 samples.

The first analytical step was to remove all peptides for which there was no information contained in the proteomic raw data matrix and the peptides for which 85% or more of the signal values were missing. All these peptides were specific of mouse proteins and in many cases were unique. The corresponding proteins were annotated and labelled together with each measured peptide. Next, low-quality samples were also removed, testing the overall signal per sample to identify if there were clear outliers with very low signal or with a very different signal distribution. Comparison of the overall signal distributions of the 30 samples (comparing boxplots) and identified 3 samples that were very different were obtained and discarded (WT/-Phag\_s3r2 (sample 3, replicate 2), RBD/-\_s2r1 and LKRc\_s1r1). These 3 samples showed a median signal in their distributions that deviated >20% from the median signal of the distributions of all other samples.

Differential expression analysis for each peptide of each protein were next performed. The algorithm used to carry out this analysis was *limmaVoom* within EdgeR R package<sup>77, 78</sup>. Prior to this analysis, a Bartlett test was performed to see the homogeneity of variances, verifying that for this data we cannot consider equality of variances and this factor was included in the differential expression algorithm. With this algorithm, normalization factors to use *a-posteriori* were calculated and, transformation and calculation of the variance weights was performed. The model to fit before using *Voom* as specified since it uses the variances of the model's residuals (observed - fitted). Finally, an estimation of the contrast for each feature tested (i.e. each peptide) was carried out using the *Empirical Bayes* approach in *limma* as previously described<sup>79</sup>.

Peptides were ordered by the p.value of the *limma* test considering significant peptides changed only with a p.value below 0.05 and with a  $\log_2(\text{Fold-Change}) > |2|$ . All these analyses were performed using the statistical computing language R and packages or libraries obtained from R-cran ([cran.r-project.org](http://cran.r-project.org)) or Bioconductor ([www.bioconductor.org](http://www.bioconductor.org)).

Cytoscape software (v3.9)<sup>80</sup> including GeneMania app<sup>81</sup> was then used to generate and visualize protein-protein networks of the significantly altered proteins selected in secretome analysis of unstimulated and phagocytosing BMDMs. This tool provides information on protein-protein associations based in co-expression studies and also based in physical interaction studies.

## Western blot analysis

Immunoblot was performed per a general western-blot protocol (Abcam). Total protein was extracted using Cell Lysis Buffer (Cell Signaling Technology) supplemented with complete mini protease inhibitor cocktail (Roche), 50 mM sodium fluoride and 1 mM of PMSF. Protein was quantified using Bradford Method (Bio-Rad). 20  $\mu\text{g}$  of protein was separated by SDS-PAGE and transferred to 0.2  $\mu\text{m}$  pore-size PVDF membranes (Sigma-Aldrich). Blots were probed using the following antibodies, at a concentration 1:1000 unless otherwise stated: phospho-LIMK1 (Thr508)/LIMK2 (Thr505) (#3841, Cell signalling), cofilin (sc-376476, Santa Cruz Biotechnology), phospho-cofilin (Ser3) (#3311, Cell Signalling), cathepsin B (12216-1-AP, Proteintech), cathepsin D (21327-1-AP, Proteintech),  $\alpha$ -tubulin (ab15246, Abcam; concentration 1:5000). Horseradish peroxidase-conjugated secondary antibodies (Amersham) were used (1:5000) and detected using an enhanced chemiluminescent substrate (Amersham). Signal was detected using an iBright 1500 System (Invitrogen).

## Flow cytometry analysis

Single-cell suspensions from cultured cell, spleen or blood monocytes were generated from mice, washed twice in staining buffer and incubated with 1:100 Fc-block (BD Biosciences, #553142) diluted in FACS buffer. Cells were subjected to surface antibody staining with labelled antibodies diluted in staining buffer for 30 min at 4 °C: CD3-PE-Cy7 (#100328, Biolegend), CD4-BV605 (#100548, Biolegend), CD8-APC (#100712, Biolegend), CD19-PerCP-Cy5.5 (#115534, Biolegend), CD45-BV785 (#103149, Biolegend), Ly6C-PerCP-Cy5.5 (#128012, Biolegend), Ly6C-E450 (#48-5932-82, eBioscience), Ly6G-AF700 (#56-5931-82, eBioscience), CD11b-BV650 (#101239, Biolegend), F4/80-PE-Cy7 (#123114, Biolegend), F4/80-PE (#123110, Biolegend), CCR2 (CD192)-PE-Vio @ 770 (#130-108-724, Miltenyi Biotec). After incubation, cells were washed in staining buffer and analysed immediately. For all staining, isotype controls were used.

Samples were acquired on a BD LSR FORTESA FACS or FACS Aria III machine that uses FACS DIVA software (BD Biosciences). Compensation was performed using 1 drop of Ultracomp ebeads (eBioscience) in 300  $\mu\text{l}$  of FACS buffer. 1  $\mu\text{l}$  of each antibody used in the pool was mixed with 100  $\mu\text{l}$  of compensation beads solution and acquired. Analysis was performed with FlowJo software (FlowJo V10.4). Once the different pools were compensated samples were acquired.

## Phagocytosis assay

For phagocytosis assay, BMDMs in suspension in DMEM without FBS were labelled with 1:200 red cell tracker (Molecular Probes) for 30 minutes at 37°C. Cells were then centrifuged for 5 minutes at 300g at 4°C, supernatant was removed and labelled BMDMs were washed twice with 5ml of PBS and plated overnight in complete DMEM containing 20 ng/ml M-CSF. On the following day, 100 ng/ml LPS (Sigma Aldrich) was added to BMDMs overnight.

LKR10 cells (murine lung cancer cell line) were stained with red cell tracker (Molecular Probes) as previously described for macrophages. Stained cells were plated in complete DMEM medium and, after 12 hours, 50  $\mu\text{M}$  Cisplatin (MCE MedChemExpress) was added to the media and left

overnight. BMDMs were then incubated with apoptotic cancer cells at a 1:2 ratio and cultured at 37 °C for different time points in DMEM supplemented with 10% FBS.

Flow cytometry data were acquired using a BD LSR FORTRESSA FACS instrument with FACS DIVA software (BD Biosciences) and analysed using FlowJo V10.4 software. A minimum of  $2 \times 10^5$  events were acquired and analysed. Data analysis and interpretation was done using FlowJo software (FlowJo V10.4).

## Image analysis

Confocal images were post-processed and analysed using Fiji distribution of ImageJ version 1.53q. Cell shape descriptors such as “aspect ratio” (AR), “circularity” (C) and “cell area” were measured using Fiji. Specifically, aspect ratio is calculated as  $(\text{major axis} / \text{minor axis} - 1)$  therefore representing solely the degree of elongation, whereas circularity is calculated as  $[4\pi \times (\text{area} \times \text{perimeter} - 2)]$ , thus representing the degree of similarity to a circumference with a value ranging from 0 to 1 (perfect circle).

## Histology

Tissue was fixed using 4% formaldehyde for 48h, dehydrated and paraffin-embedded. Sections (3  $\mu\text{m}$ ) were cut and stained using hematoxylin-eosin. For immunodetection, citrate pH 6 buffer was used for antigen retrieval. Staining was used using the following primary antibodies: CD68 (ab125212, Abcam, 1:200), cathepsin B (12216-1-AP, Proteintech, 1:100), cathepsin D (21327-1-AP, Proteintech, 1:400). Dako EnVision+ System HRP labelled Polymer secondary antibodies (Dako) were used, and DAB+ Substrate Chromogen System (Dako) was used for color development.

## Note

This reviewed preprint has been updated to use the correct text; previously, a version prior to the one reviewed was presented here.



## References

- 1 Cuesta C., Arevalo-Alameda C., Castellano E. (2021) **The Importance of Being PI3K in the RAS Signaling Network** *Genes (Basel)* **12**
- 2 Cantley L.C. (2002) **The phosphoinositide 3-kinase pathway** *Science* **296**:1655–1657
- 3 Castellano E., Downward J. (2011) **RAS Interaction with PI3K: More Than Just Another Effector Pathway** *Genes Cancer* **2**:261–274
- 4 Madsen R.R., Vanhaesebroeck B. (2020) **Cracking the context-specific PI3K signaling code** *Sci Signal* **13**
- 5 Troutman T.D., Bazan J.F., Pasare C. (2012) **Toll-like receptors, signaling adapters and regulation of the pro-inflammatory response by PI3K** *Cell Cycle* **11**:3559–3567
- 6 Vergadi E., Ieronymaki E., Lyroni K., Vaporidi K., Tsatsanis C. (2017) **Akt Signaling Pathway in Macrophage Activation and M1/M2 Polarization** *J Immunol* **198**:1006–1014
- 7 Kok K., Geering B., Vanhaesebroeck B. (2009) **Regulation of phosphoinositide 3- kinase expression in health and disease** *Trends Biochem Sci* **34**:115–127
- 8 Hawkins P.T., Stephens L.R. (2015) **PI3K signalling in inflammation** *Biochim Biophys Acta* **1851**:882–897
- 9 Okkenhaug K. (2013) **Signaling by the phosphoinositide 3-kinase family in immune cells** *Annu Rev Immunol* **31**:675–704
- 10 et al., Tamura N. (2009) **Specific role of phosphoinositide 3-kinase p110alpha in the regulation of phagocytosis and pinocytosis in macrophages** *Biochem J* **423**:99–108
- 11 et al., Castellano E. (2013) **Requirement for interaction of PI3-kinase p110alpha with RAS in lung tumor maintenance** *Cancer Cell* **24**:617–630
- 12 et al., Murillo M.M. (2014) **RAS interaction with PI3K p110alpha is required for tumor-induced angiogenesis** *J Clin Invest* **124**:3601–3611
- 13 Bennett J.M., Reeves G., Billman G.E., Sturmborg J.P. (2018) **Inflammation-Nature’s Way to Efficiently Respond to All Types of Challenges: Implications for Understanding and Managing “the Epidemic” of Chronic Diseases** *Front Med (Lausanne)* **5**
- 14 Chaplin D.D. (2010) **Overview of the immune response** *J Allergy Clin Immunol* **125**:S3–23
- 15 et al., Chen L. (2018) **Inflammatory responses and inflammation-associated diseases in organs** *Oncotarget* **9**:7204–7218
- 16 Medzhitov R. (2008) **Origin and physiological roles of inflammation** *Nature* **454**:428–435
- 17 Medzhitov R. (2021) **The spectrum of inflammatory responses** *Science* **374**:1070–1075

- 18 Gerhardt T., Ley K. (2015) **Monocyte trafficking across the vessel wall** *Cardiovasc Res* **107**:321–330
- 19 Shi C., Pamer E.G. (2011) **Monocyte recruitment during infection and inflammation** *Nat Rev Immunol* **11**:762–774
- 20 Ballabio A., Bonifacino J.S. (2020) **Lysosomes as dynamic regulators of cell and organismal homeostasis** *Nat Rev Mol Cell Biol* **21**:101–118
- 21 Saftig P., Puertollano R. (2021) **How Lysosomes Sense, Integrate, and Cope with Stress** *Trends Biochem Sci* **46**:97–112
- 22 Gordon S. (2016) **Phagocytosis: An Immunobiologic Process** *Immunity* **44**:463–475
- 23 Kourtzelis I., Hajishengallis G., Chavakis T. (2020) **Phagocytosis of Apoptotic Cells in Resolution of Inflammation** *Front Immunol* **11**
- 24 et al., Gupta S. (2007) **Binding of ras to phosphoinositide 3-kinase p110alpha is required for ras-driven tumorigenesis in mice** *Cell* **129**:957–968
- 25 et al., Zhao J.J. (2006) **The p110alpha isoform of PI3K is essential for proper growth factor signaling and oncogenic transformation** *Proc Natl Acad Sci U S A* **103**:16296–16300
- 26 et al., Zhao H. (2021) **Inflammation and tumor progression: signaling pathways and targeted intervention** *Signal Transduct Target Ther* **6**
- 27 et al., Paulsen O. (2017) **The relationship between pro-inflammatory cytokines and pain, appetite and fatigue in patients with advanced cancer** *PLoS One* **12**
- 28 et al., Swirski F.K. (2009) **Identification of splenic reservoir monocytes and their deployment to inflammatory sites** *Science* **325**:612–616
- 29 Akashi K., Traver D., Miyamoto T., Weissman I.L. (2000) **A clonogenic common myeloid progenitor that gives rise to all myeloid lineages** *Nature* **404**:193–197
- 30 et al., Tsuyada A. (2012) **CCL2 mediates cross-talk between cancer cells and stromal fibroblasts that regulates breast cancer stem cells** *Cancer Res* **72**:2768–2779
- 31 Matsushima K., Larsen C.G., DuBois G.C., Oppenheim J.J. (1989) **Purification and characterization of a novel monocyte chemotactic and activating factor produced by a human myelomonocytic cell line** *J Exp Med* **169**:1485–1490
- 32 Orecchioni M., Ghosheh Y., Pramod A.B., Ley K. (2019) **Macrophage Polarization: Different Gene Signatures in M1(LPS+) vs. Classically and M2(LPS-) vs. Alternatively Activated Macrophages** *Front Immunol* **10**
- 33 Cui K., Ardell C.L., Podolnikova N.P., Yakubenko V.P. (2018) **Distinct Migratory Properties of M1, M2, and Resident Macrophages Are Regulated by alpha(D)beta(2) and alpha(M)beta(2) Integrin-Mediated Adhesion** *Front Immunol* **9**
- 34 Auffray C., Sieweke M.H., Geissmann F. (2009) **Blood monocytes: development, heterogeneity, and relationship with dendritic cells** *Annu Rev Immunol* **27**:669–692

- 35 et al., Geissmann F. (2010) **Development of monocytes, macrophages, and dendritic cells** *Science* **327**:656–661
- 36 et al., Soler A. (2013) **Inhibition of the p110alpha isoform of PI 3-kinase stimulates nonfunctional tumor angiogenesis** *J Exp Med* **210**:1937–1945
- 37 Faust N., Varas F., Kelly L.M., Heck S., Graf T. (2000) **Insertion of enhanced green fluorescent protein into the lysozyme gene creates mice with green fluorescent granulocytes and macrophages** *Blood* **96**:719–726
- 38 et al., Schwartz A.B. (2021) **Elucidating the Biomechanics of Leukocyte Transendothelial Migration by Quantitative Imaging** *Front Cell Dev Biol* **9**
- 39 Vicente-Manzanares M., Sanchez-Madrid F. (2004) **Role of the cytoskeleton during leukocyte responses** *Nat Rev Immunol* **4**:110–122
- 40 McWhorter F.Y., Wang T., Nguyen P., Chung T., Liu W.F. (2013) **Modulation of macrophage phenotype by cell shape** *Proc Natl Acad Sci U S A* **110**:17253–17258
- 41 Pollard T.D., Cooper J.A. (2009) **Actin, a central player in cell shape and movement** *Science* **326**:1208–1212
- 42 et al., Ronzier E. (2022) **The Actin Cytoskeleton Responds to Inflammatory Cues and Alters Macrophage Activation** *Cells* **11**
- 43 Biswas S.K., Mantovani A. (2010) **Macrophage plasticity and interaction with lymphocyte subsets: cancer as a paradigm** *Nat Immunol* **11**:889–896
- 44 et al., Mercurio A. (2013) **Rapamycin unbalances the polarization of human macrophages to M1** *Immunology* **140**:179–190
- 45 Hajishengallis G., Lambris J.D. (2016) **More than complementing Tolls: complement-Toll-like receptor synergy and crosstalk in innate immunity and inflammation** *Immunol Rev* **274**:233–244
- 46 Reis E.S., Mastellos D.C., Hajishengallis G., Lambris J.D. (2019) **New insights into the immune functions of complement** *Nat Rev Immunol* **19**:503–516
- 47 Underhill D.M., Goodridge H.S. (2012) **Information processing during phagocytosis** *Nat Rev Immunol* **12**:492–502
- 48 Yadati T., Houben T., Bitorina A., Shiri-Sverdlov R. (2020) **The Ins and Outs of Cathepsins: Physiological Function and Role in Disease Management** *Cells* **9**
- 49 Shinoda H., Shannon M., Nagai T. (2018) **Fluorescent Proteins for Investigating Biological Events in Acidic Environments** *Int J Mol Sci* **19**
- 50 Fredman G., Li Y., Dalli J., Chiang N., Serhan C.N. (2012) **Self-limited versus delayed resolution of acute inflammation: temporal regulation of pro-resolving mediators and microRNA** *Sci Rep* **2**
- 51 Meizlish M.L., Franklin R.A., Zhou X., Medzhitov R. (2021) **Tissue Homeostasis and Inflammation** *Annu Rev Immunol* **39**:557–581

- 52 Bamburg J.R., Bernstein B.W. (2010) **Roles of ADF/cofilin in actin polymerization and beyond** *F1000 Biol Rep* **2**
- 53 et al., Wilson K. (2013) **Mechanisms of leading edge protrusion in interstitial migration** *Nat Commun* **4**
- 54 Aderem A., Underhill D.M. (1999) **Mechanisms of phagocytosis in macrophages** *Annu Rev Immunol* **17**:593–623
- 55 Edwards D.C., Sanders L.C., Bokoch G.M., Gill G.N. (1999) **Activation of LIM-kinase by Pak1 couples Rac/Cdc42 GTPase signalling to actin cytoskeletal dynamics** *Nat Cell Biol* **1**:253–259
- 56 et al., Ohashi K. (2000) **Rho-associated kinase ROCK activates LIM-kinase 1 by phosphorylation at threonine 508 within the activation loop** *J Biol Chem* **275**:3577–3582
- 57 Soriano O., Alcon-Perez M., Vicente-Manzanares M., Castellano E. (2021) **The Crossroads between RAS and RHO Signaling Pathways in Cellular Transformation, Motility and Contraction** *Genes (Basel)* **12**
- 58 Hoon J.L., Tan M.H., Koh C.G. (2016) **The Regulation of Cellular Responses to Mechanical Cues by Rho GTPases** *Cells* **5**
- 59 Wolfenson H., Yang B., Sheetz M.P. (2019) **Steps in Mechanotransduction Pathways that Control Cell Morphology** *Annu Rev Physiol* **81**:585–605
- 60 Kamnev A., Lacouture C., Fusaro M., Dupre L. (2021) **Molecular Tuning of Actin Dynamics in Leukocyte Migration as Revealed by Immune-Related Actinopathies** *Front Immunol* **12**
- 61 et al., Scerra G. (2022) **Lysosomal positioning diseases: beyond substrate storage** *Open Biol* **12**
- 62 Ferreira C.R., Gahl W.A. (2017) **Lysosomal storage diseases** *Transl Sci Rare Dis* **2**:1–71
- 63 Liu T., Zhang L., Joo D., Sun S.C. (2017) **NF-kappaB signaling in inflammation** *Signal Transduct Target Ther* **2**
- 64 Tak P.P., Firestein G.S. (2001) **NF-kappaB: a key role in inflammatory diseases** *J Clin Invest* **107**:7–11
- 65 et al., de Mingo A. (2016) **Cysteine cathepsins control hepatic NF-kappaB-dependent inflammation via sirtuin-1 regulation** *Cell Death Dis* **7**
- 66 Yang N., Wang P., Wang W.J., Song Y.Z., Liang Z.Q. (2015) **Inhibition of cathepsin L sensitizes human glioma cells to ionizing radiation in vitro through NF-kappaB signaling pathway** *Acta Pharmacol Sin* **36**:400–410
- 67 Chang C.P., Su Y.C., Hu C.W., Lei H.Y. (2013) **TLR2-dependent selective autophagy regulates NF-kappaB lysosomal degradation in hepatoma-derived M2 macrophage differentiation** *Cell Death Differ* **20**:515–523
- 68 et al., Liu N. (2003) **NF-kappaB protects from the lysosomal pathway of cell death** *EMBO J* **22**:5313–5322

- 69 Kast D.J., Dominguez R. (2017) **The Cytoskeleton-Autophagy Connection** *Curr Biol* **27**:R318–R326
- 70 et al., Taunton J. (2000) **Actin-dependent propulsion of endosomes and lysosomes by recruitment of N-WASP** *J Cell Biol* **148**:519–530
- 71 Ettelt R.V., Vucak G., Didusch S., Riemelmoser B., Ehrenreiter K., Hartl M., Huber L.A., Baccarini M. (2023) **Interplay between PLEKHG3-regulated actin dynamics and lysosomal trafficking in cell motility** *BioRxiv*
- 72 et al., Stockinger W. (2006) **Differential requirements for actin polymerization, calmodulin, and Ca<sup>2+</sup> define distinct stages of lysosome/phagosome targeting** *Mol Biol Cell* **17**:1697–1710
- 73 Hryciw D.H., Lee E.M., Pollock C.A., Poronnik P. (2004) **Molecular changes in proximal tubule function in diabetes mellitus** *Clin Exp Pharmacol Physiol* **31**:372–379
- 74 et al., Gong G.Q. (2023) **A small-molecule PI3Kalpha activator for cardioprotection and neuroregeneration** *Nature* **618**:159–168
- 75 Jorba I., Uriarte J.J., Campillo N., Farre R., Navajas D. (2017) **Probing Micromechanical Properties of the Extracellular Matrix of Soft Tissues by Atomic Force Microscopy** *J Cell Physiol* **232**:19–26
- 76 et al., Rico F. (2005) **Probing mechanical properties of living cells by atomic force microscopy with blunted pyramidal cantilever tips** *Phys Rev E Stat Nonlin Soft Matter Phys* **72**
- 77 McCarthy D.J., Chen Y., Smyth G.K. (2012) **Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation** *Nucleic Acids Res* **40**:4288–4297
- 78 Robinson M.D., McCarthy D.J., Smyth G.K. (2010) **edgeR: a Bioconductor package for differential expression analysis of digital gene expression data** *Bioinformatics* **26**:139–140
- 79 et al., Campos-Laborie F.J. (2019) **DECO: decompose heterogeneous population cohorts for patient stratification and discovery of sample biomarkers using omic data profiling** *Bioinformatics* **35**:3651–3662
- 80 Killcoyne S., Carter G.W., Smith J., Boyle J. (2009) **Cytoscape: a community-based framework for network modeling** *Methods Mol Biol* **563**:219–239
- 81 et al., Franz M. (2018) **GeneMANIA update 2018** *Nucleic Acids Res* **46**:W60–W64

## Article and author information

### Alejandro Rosell

Tumour-Stroma Signalling Lab., Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain

### Agata A. Krygowska

Centre for Cancer and Inflammation, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom



**Marta Alcón Pérez**

Tumour-Stroma Signalling Lab., Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain

**Mathieu-Benoit Voisin**

Centre for Microvascular Research, William Harvey Research Institute, Queen Mary University of London, Charterhouse Square, London, EC1M 6BQ, United Kingdom

**Juan de Paz**

Tumour-Stroma Signalling Lab., Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain

**Héctor Sanz-Fraile**

Unit of Biophysics and Bioengineering, Department of Biomedicine, School of Medicine and Health Sciences, Universitat de Barcelona, Barcelona, Spain

**Vinothini Rajeeve**

Centre for Cancer Genomics and Computational Biology, Cell Signalling and Proteomics Laboratory, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom  
ORCID iD: [0000-0002-6361-4291](https://orcid.org/0000-0002-6361-4291)

**Alberto Berral-González**

Bioinformatics and Functional Genomics, Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain

**Cristina Cuesta**

Tumour-Stroma Signalling Lab., Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain

**Ottillie Swinyard**

Centre for Cancer and Inflammation, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom

**Enrique Gabandé-Rodriguez**

Centre for Cancer and Inflammation, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom  
ORCID iD: [0000-0002-9715-8714](https://orcid.org/0000-0002-9715-8714)

**Julian Downward**

Oncogene Biology Laboratory, Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK  
ORCID iD: [0000-0002-2331-4729](https://orcid.org/0000-0002-2331-4729)

**Jordi Alcaraz**

Unit of Biophysics and Bioengineering, Department of Biomedicine, School of Medicine and Health Sciences, Universitat de Barcelona, Barcelona, Spain, Institute for Bioengineering of Catalonia (IBEC), The Barcelona Institute for Science and Technology (BIST), Barcelona 08028, Spain

**Javier De Las Rivas**

Bioinformatics and Functional Genomics, Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain

**Pedro Cutillas**

Centre for Cancer Genomics and Computational Biology, Cell Signalling and Proteomics Laboratory, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom

ORCID iD: [0000-0002-3426-2274](https://orcid.org/0000-0002-3426-2274)

**Esther Castellano**

Tumour-Stroma Signalling Lab., Centro de Investigación del Cáncer, Instituto de Biología Molecular y Celular del Cáncer, Consejo Superior de Investigaciones Científicas (CSIC)-Universidad de Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain, Centre for Cancer and Inflammation, Barts Cancer Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, United Kingdom

**For correspondence:** [ecastellano@usal.es](mailto:ecastellano@usal.es)

ORCID iD: [0000-0002-8449-4081](https://orcid.org/0000-0002-8449-4081)

**Copyright**

© 2024, Rosell et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

**Editors**

Reviewing Editor

**Tadatsugu Taniguchi**

University of Tokyo, Tokyo, Japan

Senior Editor

**Tadatsugu Taniguchi**

University of Tokyo, Tokyo, Japan

**Reviewer #1 (Public Review):**

In this study, Alejandro Rosell et al. uncovers the immunoregulation functions of RAS-p110 $\alpha$  pathway in macrophages, including the extravasation of monocytes from the bloodstream and subsequent lysosomal digestion. Disrupting RAS-p110 $\alpha$  pathway by mouse genetic tools or by pharmacological intervention, hampers the inflammatory response, leading to delayed resolution and more severe acute inflammatory reactions. The authors proposed that activating p110 $\alpha$  using small molecules could be a promising approach for treating chronic

inflammation. This study provides insights into the roles and mechanisms of p110 $\alpha$  on macrophage function and the inflammatory response, while some conclusions are still questionable because of several issues described below.

(1) Fig. 1B showed that disruption of RAS-p110 $\alpha$  causes the decrease in the activation of NF- $\kappa$ B, which is a crucial transcription factor that regulates the expression of proinflammatory genes. However, the authors observed that disruption of RAS-p110 $\alpha$  interaction results in an exacerbated inflammatory state *in vivo*, in both localized paw inflammation and systemic inflammatory mediator levels. Also, the authors introduced that "this disruption leads to a change in macrophage polarization, favouring a more proinflammatory M1 state" in introduction according to reference 12. The conclusions drew from the signaling and the models seemed contradictory and puzzling. Besides, it is not clear why the protein level of p65 was decreased at 10' and 30'. Was it attributed to the degradation of p65 or experimental variation?

(2) In Fig 3, the authors used bone-marrow derived macrophages (BMDMs) instead of isolated monocytes to evaluate the ability of monocyte transendothelial migration, which is not sufficiently convincing. In Fig. 3B, the authors evaluated the migration in Pik3caWT/- BMDMs, and Pik3caWT/WT BMDMs treated with BYL-719'. Given that the dose effect of gene expression, the best control is Pik3caWT/- BMDMs treated with BYL-719.

(3) In Fig. 4E-4G, the authors observed that elevated levels of serine 3 phosphorylated Cofilin in Pik3caRBD/- BMDMs both in unstimulated and in proinflammatory conditions, and phosphorylation of Cofilin at Ser3 increase actin stabilization, it is not clear why disruption of RAS-p110 $\alpha$  binding caused a decrease in the F-actin pool in unstimulated BMDMs?

<https://doi.org/10.7554/eLife.94590.1.sa1>

## Reviewer #2 (Public Review):

### Summary:

Cell intrinsic signaling pathways controlling the function of macrophages in inflammatory processes, including in response to infection, injury or in the resolution of inflammation are incompletely understood. In this study, Rosell et al. investigate the contribution of RAS-p110 $\alpha$  signaling to macrophage activity. p110 $\alpha$  is a ubiquitously expressed catalytic subunit of PI3K with previously described roles in multiple biological processes including in epithelial cell growth and survival, and carcinogenesis. While previous studies have already suggested a role for RAS-p110 $\alpha$  signaling in macrophages function, the cell intrinsic impact of disrupting the interaction between RAS and p110 $\alpha$  in this central myeloid cell subset is not known.

### Strengths:

Exploiting a sound previously described genetically mouse model that allows tamoxifen-inducible disruption of the RAS-p110 $\alpha$  pathway and using different readouts of macrophage activity *in vitro* and *in vivo*, the authors provide data consistent with their conclusion that alteration in RAS-p110 $\alpha$  signaling impairs the function of macrophages in a cell intrinsic manner. The study is well designed, clearly written with overall high-quality figures.

### Weaknesses:

My main concern is that for many of the readouts, the difference between wild-type and mutant macrophages *in vitro* or between wild-type and Pik3caRBD mice *in vivo* is rather modest, even if statistically significant (e.g. Figure 1A, 1C, 2A, 2F, 3B, 4B, 4C). In other cases, such as for the analysis of the H&E images (Figure 1D-E, S1E), the images are not quantified, and it is hard to appreciate what the phenotype in samples from Pik3caRBD mice is or

whether this is consistently observed across different animals. Also, the authors claim there is a 'notable decrease' in Akt activation but 'no discernible change' in ERK activation based on the western blot data presented in Figure 1A. I do not think the data shown supports this conclusion.

To further substantiate the extent of macrophage function alteration upon disruption of RAS-p110 $\alpha$  signaling, the manuscript would benefit from testing macrophage activity in vitro and in vivo across other key macrophage activities such as bacteria phagocytosis, cytokine/chemokine production in response to titrating amounts of different PAMPs, inflammasome function, etc. This would be generally important overall but also useful to determine whether the defects in monocyte motility or macrophage lysosomal function are selectively controlled downstream of RAS-p110 $\alpha$  signaling.

Furthermore, given the key role of other myeloid cells besides macrophages in inflammation and immunity it remains unclear whether the phenotype observed in vivo can be attributed to impaired macrophage function. Is the function of neutrophils, dendritic cells or other key innate immune cells not affected?

Compelling proof of concept data that targeting RAS-p110 $\alpha$  signalling constitutes indeed a putative approach for modulation of chronic inflammation is lacking. Addressing this further would increase the conceptual advance of the manuscript and provide extra support to the authors' suggestion that p110 $\alpha$  inhibition or activation constitute promising approaches to manage inflammation.

Finally, the analysis by FACS should also include information about the total number of cells, not just the percentage, which is affected by the relative change in other populations. On this point, Figure S2B shows a substantial, albeit not significant (with less number of mice analysed), increase in the percentage of CD3<sup>+</sup> cells. Is there an increase in the absolute number of T cells or does this apparent relative increase reflect a reduction in myeloid cells?

<https://doi.org/10.7554/eLife.94590.1.sa0>