

Reviewed Preprint

v1 • October 13, 2025

Not revised

Reviewed Preprint

v2 • February 20, 2026

Revised by authors

Death receptor 6 does not regulate axon degeneration and Schwann cell injury responses during Wallerian degeneration

✉ For correspondence:

bogdan.beirowski@osumc.edu

elisabetta.babetto@osumc.edu

Competing interests:

No competing interests declared

Funding:

See [page 18](#)

Reviewing editor:

Lu Chen, Stanford University, United States

© 2025, Beirowski et al. This article is

distributed under the terms of the

[Creative Commons Attribution](#)

[License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Bogdan Beirowski ✉, Haoran Huang, Elisabetta Babetto ✉

Department of Neurology, The Neuroscience Research Institute, College of Medicine, The Ohio State University Wexner Medical Center, Columbus, United States

eLife Assessment

In this **valuable** study, through carefully executed and rigorously controlled experiments, the authors challenged a previously reported role of the Death Receptor 6 (DR6/Tnfrsf21) in Wallerian degeneration (WD). Using two DR6 knockout mouse lines and multiple WD assays, both in vitro and in vivo, the authors provided **convincing** evidence that loss of DR6 in mice does not protect peripheral axons from WD after injury, at least in the specific contexts of the mice and analyses performed in this study. Due to the lack of certain specific parameters from previous studies (sex, age, mouse strains etc.), the exact reasons underlying the observed inconsistencies between current and previous reports on the protective effects of DR6 remains to be determined. Overall, this is a carefully executed study providing invaluable information toward understanding DR6's role (or lack thereof) in axon degeneration.

<https://doi.org/10.7554/eLife.108389.2.sa3>

Abstract

Axon degeneration (AxD), accompanied by glial remodeling, is a pathological hallmark of many neurodegenerative diseases, leading to the disruption of neuronal connectivity [1–3]. Understanding the mechanisms in neurons and glia that regulate AxD is essential for developing therapeutic strategies to prevent or mitigate axon loss. Wallerian degeneration (WD) is a well-established model to study the mechanisms of nerve injury-induced AxD, glial responses, and axon-glia interactions. We recently showed that Schwann cells (SCs), the axon-associated glia of the peripheral nervous system, exert protective effects on axons through their rapid metabolic injury response [4]. Enhancing this SC response promotes axon protection during WD [4]. A prior study reported that eliminating the orphan tumor necrosis factor receptor DR6 (death receptor 6, Tnfrsf21) strongly delays AxD and alters SC injury responses during WD [5], suggesting a possible intersection with our findings. Here, we rigorously revisit the role of DR6 in WD using two independent DR6 knockout mouse lines including the same model used in the previous study. Surprisingly, in striking contrast to the earlier report, we observed no impact of DR6 deletion on AxD kinetics or SC injury responses across a range of WD assays. Moreover, injured axons in primary neuronal cultures lacking DR6 degenerated at a similar rate as wild-type axons. We conclude that DR6 is dispensable for the regulation of AxD and glial nerve injury responses during WD. Our data argue that any therapeutic benefit from DR6 suppression in neurodegeneration models occurs through mechanisms independent of WD.

Results

AxD following nerve injury proceeds normally in mice lacking DR6

DR6 has been implicated in AxD mechanisms in several neurodegeneration models [6–9]. Given the proposed mechanistic overlap with WD [10], we first sought to confirm that DR6 loss protects axons after injury [5]. The previous work reported that disconnected axons from DR6 null mice are preserved for up to 4 weeks after nerve injury, with a phenotypic penetrance in approximately 38.5% of the animals studied [5]. Quantitative real-time (qRT) PCR and western blot analysis validated successful disruption of DR6 expression in constitutive DR6 null mice (DR6^{-/-}), and in another DR6 knockout model with a distinct targeting strategy in which we deleted exon 2 of the *DR6* gene in all tissues including germ cells through Cre-mediated recombination (DR6^{-/-}, CMV-Cre) (Fig. S1). Unlike earlier studies using DR6 null mutants [5, 11], we found no significant abnormalities in early developmental myelin formation at postnatal day 1 (Fig. S2), or in g-ratio and axon- and fiber-caliber distributions as assessed by nerve histomorphometry of sciatic and tibial nerve axons from adult DR6^{-/-} and DR6^{-/-}, CMV-Cre animals (Fig. S3). This ensures a normal baseline for our comparative WD studies that follow below.

We transected sciatic nerves of overall 38 DR6^{-/-} and 18 DR6^{-/-}, CMV-Cre mice and assessed axon survival in the distal nerve stump 3 days post-axotomy by light and electron microscopy in comparison to littermate controls. Surprisingly, we found that all studied DR6^{-/-} and DR6^{-/-}, CMV-Cre mice consistently exhibited an almost complete breakdown of axonal structure, identical to control littermate mice (Fig. 1A). In contrast, nerve stumps from mice expressing one copy of the *Wld^S* gene, and mice with loss of the pro-degenerative molecules Phr1/Myebp2 or SARM1 displayed profound preservation of the vast majority of axons, in line with previous work (Fig. 1B).

To detect a potentially smaller delay of AxD in the DR6 null mouse models, we next performed lower stringency assays and studied axon survival 30h after nerve injury. This is a time point at which many disconnected axons just start to disintegrate, and ~50% of myelinated axons in tibial nerves remain structurally intact in wild-type mice as classified on high-resolution light and electron micrographs [4]. Fiber quantification in axotomized nerve segments from DR6^{-/-} and DR6^{-/-}, CMV-Cre mutants demonstrated that the axon survival was statistically indistinguishable from control samples (Fig. 2A). By contrast, *Wld^S* mice, and mutants lacking Phr1/Myebp2 or SARM1, displayed significantly enhanced axon survival (Fig. 2B).

We next asked if disconnected nerve stumps from DR6^{-/-} and DR6^{-/-}, CMV-Cre mutants show reduced site-specific phosphorylation (Thr183/Tyr185) and thus activation of c-Jun N-terminal kinases (JNK), as reported previously [5]. Reduced activation of JNKs, members of the mitogen-activated protein kinase (MAPK) family, has been also observed in axotomized optic nerves under conditions of axon protection conferred by SARM1 loss, or expression of a *Wld^S* variant [12]. In accord with the absence of any structural axon protection, we found that the JNK activation in axotomized nerves from DR6^{-/-} and DR6^{-/-}, CMV-Cre mutants was indistinguishable from control preparations (Fig. S4).

Together, these data indicate that the loss of DR6 *in vivo* has no obvious effect on the progression of AxD in axotomized nerves.

Normal Schwann cell injury responses and myelin remodeling dynamics during WD in the absence of DR6

SCs rapidly transdifferentiate and dismantle their myelin sheaths during WD, a process that is largely orchestrated by the transcription factor c-Jun [13, 14]. The deletion of DR6 has been claimed to alter the typical SC myelin reorganization observed after nerve lesion [5], suggesting abnormal SC transdifferentiation. To determine if DR6-deficient SCs show abnormal reactions to nerve injury, we analyzed c-Jun expression by immunofluorescence on sciatic nerve cross sections distal to the site of nerve injury (Fig. 3A). Additionally, we examined the ultrastructure of dedifferentiated SCs (Fig. S5A), and quantified the area occupied by myelin sheaths and myelin




Figure 1. Rapid Wallerian degeneration in mice lacking DR6 in comparison to multiple mutants with delayed degeneration, assessed at 3 days after axotomy

(A) Representative semithin (1st and 2nd columns) and electron micrographs (3rd column) of transverse sciatic nerve sections of distal nerve stumps from wild-type (WT) and the indicated mutant mice 3 days after sciatic nerve transection. Note complete structural disintegration of transected axons with absent or floccular cytoskeleton and collapsed myelin sheaths in the preparations from WT, DR6^{-/-}, and DR6^{-/-}, CMV-Cre mice. In contrast, the majority of disconnected axons from heterozygous Wld^S mice, and from Phr1 and SARM1 knockout mice, are structurally preserved with uniform cytoskeleton and intact myelin sheaths. Scale bars: 10 μ m (B) Quantification of preserved axons in transverse sciatic nerve sections of distal nerve stumps from mice with the indicated genotypes. Each symbol in the scatter dot plots represents the quantification from one animal (% of control axon numbers in micrographs from uninjured contralateral nerve preparations for each animal). All data were obtained from experimental animals between 3 and 12 months of age.




Figure 2. Normal rates of nerve injury-induced axon disintegration in DR6-deficient mice at earlier stages of Wallerian degeneration

(A) Representative semithin micrographs of transverse tibial nerve sections of distal nerve stumps from 3-months-old mice with the indicated genotypes (top) 30h after sciatic nerve transection with pseudocoloring of intact (turquoise) and structurally degenerated (magenta) myelinated fibers, and corresponding quantifications of relative axon survival (bottom). (B) Representative semithin micrographs of transverse tibial nerve sections of distal nerve stumps from 3-months-old mice with the indicated genotypes (top) 36h after sciatic nerve transection with pseudocoloring of intact (turquoise) and structurally degenerated (magenta) myelinated fibers, and corresponding quantifications of relative axon survival (bottom). Scale bars: 10 μ m

debris profiles on osmium tetroxide and toluidine blue stained nerve sections (Fig. S5B, C). We did not detect any significant differences in any of these assays at any time point investigated in DR6^{-/-} and DR6^{-/-},CMV-Cre samples as compared to control preparations (Fig. 3, S5). These data demonstrate that DR6-deficient SCs show normal injury responses following nerve lesion.

Injured DRG neurites in primary neuronal cultures from DR6 knockout mice degenerate rapidly

The results above contrast sharply with the previously reported dramatically delayed AxD during WD in DR6 null mice [5]. In an attempt to reconcile the contradictory findings, we finally considered the possibility that more subtle changes in the stability of injured axons could be potentially unveiled *in vitro* using embryonic neuronal explant cultures from the DR6 mutant models. Such assays were previously extensively used to reveal and characterize axonal DR6 phenotypes, although a key publication in this regard was recently retracted [15]. Importantly, in contrast to the *in vivo* data, no phenotypic penetrance effects in delaying AxD have been reported using *in vitro* assays in the previous DR6 knockout study focusing on WD [5].

We confirmed the loss of DR6 mRNA and protein in dorsal root ganglia explants prepared from DR6^{-/-} embryos, and additionally DR6^{LoxP/LoxP} embryos infected with a lentivirus expressing Cre recombinase (Fig. 4A, D, and not shown). However, time-lapse analysis of neurite disintegration after axotomy demonstrated no statistically significant differences between mutant and control preparations from the two models with the only exception of slightly accelerated AxD in the DR6^{LoxP/LoxP} preparations with the lentiviral Cre recombinase expression six hours after axotomy (Fig. 4B, C, E, F). This contrasts with the robust axonal protection observed in similar primary culture models devoid of Phr1/Mycbp2, or in cultures with expression of Wld^S protein variants we previously reported [16–18]. Together, these findings indicate that the loss of DR6 does not confer resistance to AxD in an established neuronal culture model.

Discussion

The main motivation for this study was to determine if the glial injury responses previously suggested as abnormal in DR6-deficient mice play a role in the extended axon survival observed during WD [5]. Unexpectedly, despite using the same DR6 mutant mouse line as the original study, we did not observe delayed AxD or the aberrant myelin remodeling described in that work. Our findings demonstrate that DR6 does not function as a pro-degenerative component within the axonal auto-destruction pathway, nor does it play a significant role in regulating the glial injury responses and myelin remodeling dynamics characteristic of WD in the peripheral nervous system (PNS). Our study underscores the importance of rigorous validation when assigning functional relevance to molecules with possible roles in the WD pathway.

The axon with its associated glial cells forms a specialized structural and functional unit that enables long-range neural communication. Much like how compromised cells initiate self-destruction, injured axons undergo a related but molecularly distinct process during WD, characterized by rapid metabolic collapse [12, 19]. We recently showed that SCs modulate this process through early injury responses that enhance metabolic coupling with axons [4].

Importantly, the WD pathway of axonal self-destruction is believed to contribute significantly to the progression of various neurodegenerative diseases, highlighting both its cell-autonomous and non-cell-autonomous suppression as promising therapeutic strategies. To date, apart from expression of the aberrant fusion protein Wld^S, only four endogenous molecules – Nmnat2, SARM1, Phr1/Mycbp2, and DR6 - have been implicated to play a significant role in the WD pathway in mammals [5, 16, 20, 21]. Their overexpression or inactivation has been reported to confer drastic resistance to experimentally-induced WD, with disconnected axons in the PNS surviving for days to weeks, unlike wild-type axons, which fully degenerate within 48–72 hours on structural level. While Wld^S, Nmnat2, SARM1, and Phr1/Mycbp2 are cell-autonomously involved in the common WD pathway leading to axonal metabolic failure, DR6 has emerged as a more elusive candidate. Initial reports suggested that DR6, along with its putative ligand amyloid precursor




Figure 3. Normal Schwann cell c-Jun injury responses during WD in mice lacking DR6

(A, C) Representative immunofluorescence confocal micrographs for c-Jun with DAPI nuclear counterstaining on transverse frozen sections from contralateral uninjured nerve (uncut) and distal sciatic nerve stumps 3 days and 30 h following axotomy in 3-months-old mice with the indicated genotypes. (B, D) Corresponding quantifications of percentage of c-Jun immunoreactive and DAPI⁺ cells on nerve sections. Scale bars: 50μm.




Figure 4. Normal degeneration kinetics of injured DRG neurites in primary neuronal cultures from DR6 knockout mice

(A, D) Western blot analysis (cropped blot images) of DR6 protein expression in dorsal root ganglia (DRGs) from DR6^{-/-} embryos and from DR6^{LoxP/LoxP} embryos infected with a lentivirus expressing Cre recombinase (DR6^{Cre}), together with the respective control preparations. (B, E) Time course of neurite fragmentation, quantified as degeneration index (DI) (see Experimental Procedures) in DRG preparations from embryos with the indicated genotypes or lentiviral infection conditions. The data points shown in (B) represent the averaged neurite DI values calculated from multiple DRG preparations from each embryo at the indicated time point after injury (one data point = one embryo). The data points shown in (E) represent the averaged neurite DI values calculated from multiple micrographs acquired from each DRG preparation in a cell culture well at the indicated time point after injury (one data point = one well). (C, F) Representative phase-contrast micrographs of disconnected DRG neurites from embryos with the indicated genotypes and lentiviral infection conditions at the indicated time points after axotomy. Scale bars: 50μm.

protein (APP), mediates developmental axon pruning [22–24]. However, this interpretation has been complicated by a recent retraction [15] and conflicting findings [25]. A later study proposed a striking new role for DR6 in WD, reporting that severed axons in DR6 knockout mice remained structurally preserved for up to four weeks post-injury even though separated from their parent neuronal cell bodies [5]. However, this phenotype was observed in fewer than half the animals, suggesting incomplete penetrance. The axonal phenotype, when present, was similar to that in *Wld^S* mice, which also showed incomplete phenotypic penetrance in this study [5]. Moreover, it was found that the subset of DR6-deficient mice with axon protection displayed abnormal patterns of myelin remodeling during delayed WD, suggesting aberrant SC injury responses that could have contributed to the axonal phenotype. Furthermore, another study reported that neuronal DR6 expression regulates SC proliferation and inhibits developmental myelin formation in SCs [11]. These findings together raised the possibility that inactivation of DR6 might influence both neuronal and glial functions and their crosstalk during WD. However, to date, no subsequent studies have expanded upon or validated these discoveries.

In surprising contrast to the earlier data, our results provide compelling evidence that DR6 does not regulate AxD or glial injury responses during WD in the PNS. We employed a robust experimental design and have extensively characterized the kinetics of WD in two independent DR6 mouse knockout models using a high number of experimental animals and multiple *in vivo* and *in vitro* assays. For the first time, we present a comprehensive side-by-side comparison of key mouse mutants that have previously been shown to exhibit markedly delayed WD following nerve injury. We found no evidence that loss of DR6 expression, as confirmed by qRT-PCR and western blotting, antagonizes AxD during WD. Even under low-stringency conditions designed to detect subtle axonal protection, axon death progressed normally. These findings align with a prior electrophysiological study in the central nervous system, which suggested normal degeneration of optic nerve axons in DR6-deficient mice [26]. We also did not find any changes in the SC injury responses and myelin reorganization known to be orchestrated by the transcription factor c-Jun during WD in the PNS [13, 14] in our DR6 mutant models at two post-injury time points that are frequently used to assess c-Jun expression and SC dedifferentiation characteristics. We therefore did not examine SC injury responses and myelin reorganization at 14 days after nerve injury, as performed in the previous study [5], because no evidence of axonal protection was seen at earlier post-lesion stages.

The prominent discrepancy between our results and the prior study [5] remains unresolved. One possibility is that a spontaneous germline mutation arose within the previously used mouse colony, conferring axonal protection in a subset of animals through a mechanism unrelated to DR6. Such mutation, perhaps conceptually similar to the one responsible for the *Wld^S* phenotype that occurred spontaneously in the Olac breeding colony in England [27], could then have propagated within the colony via autosomal dominant inheritance. Although we obtained our DR6^{-/-} breeders directly from the laboratory that initially reported the protective phenotype [5], it is conceivable that the specific mice we received did not carry the hypothetical second-site mutation. Alternatively, the phenotypic discrepancies may reflect the influences of unknown genetic modifiers, epigenetic factors, environmental conditions, or age-related differences that could affect the WD pathway. These variables could potentially explain the reported partial penetrance and the previously described developmental anomalies in DR6 null nerves [5, 11].

We also considered if changes in the nerve immune microenvironment might account for the discrepancies. For example, altered macrophage recruitment or activation, which is known to influence axon integrity, could contribute to differences in WD dynamics [28]. However, despite these possibilities, we note that we have never observed incomplete phenotypic penetrance effects in *Wld^S* mice we developed [17, 18, 29–31] or in mutants lacking SARM1 or Phr1/Mycbp2 [16], either in our hands or in reports from other laboratories over the last decades. This supports the view that the manipulation of these core components of an evolutionary conserved pathway produces fully penetrant protective phenotypes. Nonetheless, environmental modulation of axon protection in *Wld^S* mice has been described, although the reported effects are relatively modest [32].

DR6 inhibition has shown therapeutic benefit in several models of neurodegeneration, including amyotrophic lateral sclerosis [7, 8], multiple sclerosis [33, 34], prion disease [9], and *in vitro* amyloid-beta toxicity [6]. However, no protective effects were observed in two separate mouse models of Alzheimer's disease [24]. While these effects have often been attributed to reduced neuronal death [6, 8], changes in immune function [34], or enhanced myelination [33], the suggestion that DR6 suppression stabilizes injured axons raised the possibility that axon protection itself may have contributed to disease amelioration. Our data argue against this interpretation in the context of WD. Instead, as mentioned earlier, DR6 may play a regulatory role in alternative forms of AxD, such as axon pruning, which is known to shape and refine neural circuits during development and may also be reactivated under pathological conditions [35, 36]. Future studies using disease-specific models will be necessary to explore this hypothesis.

Author contributions

E.B. and B.B. conceived and supervised the study. E.B. and B.B. designed and planned the experiments. E.B., H.H., and B.B. performed the experiments and analyzed and interpreted the data shown in the figures. B.B. and E.B. wrote the manuscript.

Materials and Methods

Mice

All experiments were performed in compliance with the Association for Assessment of Laboratory Animal Care policies and approved by The Ohio State University College of Medicine Animal Care and Use Committee. Mice were housed under specific pathogen-free conditions at 70 °F, 50% room humidity, 12-h light/12-h dark cycle and received ad libitum access to water and food. Mice from different genotypes were group-housed in separate cages. The mice used in this study did not undergo any procedures before their stated use. The mice were of mixed sexes. In previous studies we found no significant differences in the rates of injury-induced axon degeneration between female and male mice. Mice within individual comparative *in vivo* experiments were littermates or, if use of littermates was not possible because of litter sizes not large enough, mice from different litters were age-matched for comparative experiments. Controls were regarded as littermates or age-matched mice from different litters carrying wild-type alleles or floxed alleles with no Cre recombinase expression for the respective genotype groups. Constitutive DR6 knockout mice (DR6^{-/-}) with deletion of exons 2 and 3 (on C57BL/6J.129S mixed background) and conditional DR6 mice with exon 2 flanked by loxP sites (DR6^{LoxP/LoxP}) were generated by Genentech [37, 38] (genetic background unknown). To generate DR6^{-/-}, CMV-Cre mice, we crossed DR6^{LoxP/LoxP} mice to CMV-Cre transgenic mice (strain no. 006054, The Jackson Laboratory, made congenic on C57Bl/6J background). For this study, we additionally used heterozygous Wld^S [29], Phr1 aKO [16], and SARM1^{-/-} mice (strain no. 018069, The Jackson Laboratory, made congenic on C57Bl/6J background). Genotyping was performed by PCR strategies using standard procedures and appropriate primers (sequences available upon request).

Multiplex TaqMan quantitative real-time PCR

Brain tissue from DR6^{-/-} and DR6^{-/-}, CMV-Cre mutants and the respective control mice were homogenized using a Bullet Blender (Next Advance, model BBX24B) and RNAase-free zirconium oxide beads (0.5mm diameter, Next Advance). Total brain RNA was extracted using PureLink RNA Mini Kit (Thermo Fisher Scientific) according to the manufacturer's guidelines, and RNA concentration and purity were assessed using a NanoDrop One spectrophotometer (Thermo Fisher Scientific). To eliminate potential genomic DNA contamination, 4 µg of total RNA per sample were treated with EzDNAase (Invitrogen) according to the manufacturer's instructions. The resulting RNA was split into two aliquots: 2 µg were reverse transcribed into cDNA using SuperScript VILO Master Mix (Thermo Fisher Scientific), and the remaining 2 µg underwent the same reaction without the reverse transcriptase enzyme to generate a no-RT control. Reaction times and volumes followed the manufacturer's guidelines. DR6 expression was tested with TaqMan gene expression

predesigned assay Mm00446361_m1 with PAM-MGB probe [6], and the housekeeping gene peptidyl-prolyl isomerase A (PPIA) was tested with TaqMan gene expression assay ID Mm02342430_g1 (Thermo Fisher Scientific) with VIC-MGP probe. Briefly, in each well of a 96-well plate (MicroAmp Optical 96-well reaction plate with barcode, Applied Biosystems), 1 μ l of cDNA was tested with 9 μ l of Mastermix containing 0.5 μ l of TaqMan probe, using a QuantStudio 3 Real Time PCR instrument (Applied Biosystems). The thermal-cycling conditions included an Uracil-N glycosylase incubation step (50°C for 2 minutes), followed by Polymerase activation (95°C for 20 seconds), and 40 cycles of PCR reaction (denaturation at 95°C for 3 seconds and annealing/extension at 60°C for 30 seconds). ROX reference reading (passive reference) and Cycle-Threshold (CT) values were collected and visualized with QuantStudio 3 software and analyzed with the DeltaDeltaCT method. DR6 expression levels were calculated and normalized to PPIA levels in the same well of the plate.

Western blot analysis

Mouse brains were quickly dissected and homogenized using a Polytron homogenizer (PT 1200C), followed by sonication with a Qsonica sonicator (Q500) equipped with a microtip. Sciatic nerves were quickly dissected, the epineurium removed in ice-cold phosphate buffered saline (PBS), and directly sonicated as above. Both brains and nerves were homogenized in RIPA buffer supplemented with protease and phosphatase inhibitors. Embryonic dorsal root ganglia preparations were washed with PBS to remove culture medium and then manually collected with forceps, placed in a tube with RIPA buffer supplemented with protease and phosphatase inhibitors, and then vortexed. After lysis, samples were spun at 10,000g for 10 minutes at 4°C to remove insolubilized material. Western blotting of tissue lysates was performed using Bolt Mini gels and Mini Blot wet transfer modules (Invitrogen) according to the manufacturer's instructions. For detection of the DR6 protein, tissue and cell homogenates were processed under denaturing conditions on Tris-Acetate gels according to the manufacturer's protocol optimized for large molecular weight proteins (Thermo Fisher Scientific). For detection of JNK and p-JNK (Thr183/Tyr185) in nerve lysates, proteins were separated on 4-12% Bis-Tris gradient gels (Thermo Fisher Scientific). The following antibodies were used in a solution of 5% bovine serum albumin (BSA) in Tris-buffered saline (TBS) with 0.1% of Tween20: anti-DR6 clone 6B6 (1:500, catalog no. MABC1594, Millipore Sigma), anti-phospho-SAPK/JNK (Thr183/Tyr185) clone 81E11 (1:1000, catalog no. 4668, Cell Signaling), anti-SAPK/JNK (1:1000, catalog no. 9252, Cell Signaling), anti-b-actin (1:10,000, catalog no. A5316, Millipore Sigma). Horseradish peroxidase-conjugated secondary antibodies from Cell Signaling Technology and Jackson ImmunoResearch were used for signal detection. Blot documentation was performed with a G:Box mini 6 digital imaging system (Syngene). The integrated band intensities of protein bands were determined using ImageJ software to calculate p-JNK(Thr183/Tyr185)/JNK ratios.

Histomorphometry of intact nerves

The embedding of nerve samples in Araldite 502 epoxy resin (Polysciences) and subsequent cutting and staining of 500-nm semithin cross sections was performed as described previously [4, 39, 40]. High-resolution tile scans of the entire sciatic or tibial nerve cross sections were acquired with a DMI8 digital imaging system equipped with a 100x high-numerical aperture objective and oil immersion condenser using the Application Suite X 3.7.2 software (Leica Microsystems). The quantification of thinly myelinated axons on entire sciatic nerves from P1 mouse pups was performed manually using the ImageJ Cell Counter plug-in. The *g* ratios of individual myelinated axons in sciatic or tibial nerves as a measure of myelin thickness were determined using a plug-in for the ImageJ software that allows semiautomated analysis of randomly selected axons on nerve transverse sections (<http://gratio.efil.de>). One hundred randomly chosen fibers were measured per mouse nerve. Cumulative *g* ratios were calculated for each mouse by averaging all individual *g* ratios. Axon and fiber caliber distributions were quantified from sciatic or tibial nerve micrographs using ImageJ. The Wand tracing tool was used to manually delineate individual regions of interest (ROIs). For axon caliber measurements, the tool was applied to select the axonal area (excluding the myelin sheath), with the tolerance adjusted to accurately define each axon's

boundary. For fiber caliber measurements, the entire fiber (axon plus myelin sheath) was selected using the same approach. The resulting ROI data were exported to Microsoft Excel, and calibers were calculated using the formula $A = \pi (d/2)^2$, where A is the area and d is the diameter. Binned distributions of axon and fiber calibers were calculated and visualized using GraphPad Prism software. All quantifications were conducted blinded to the genotypes of the animals.

Electron microscopy

Transmission electron microscopy was carried out as previously described [4, 39]. Electron micrographs of randomly selected areas on 85 μm thick ultrathin nerve cross sections stained with uranyl acetate and lead citrate were taken with a FEI Tecnai transmission electron microscope.

Unilateral sciatic nerve transection

Mice were deeply anesthetized using the SomnoSuite Low-Flow Anesthesia delivery system (Kent Scientific) operated with isoflurane. Right sciatic nerves were exposed and transected with surgical micro-scissors close to the sciatic notch with the contralateral nerve serving as the control. The wound was closed with surgical thread or clips and buprenorphine was administered as a postsurgery analgesic. Upon nerve removal from humanely killed mice, the lesion site was inspected to verify complete transection. Distal sciatic and tibial nerve stumps and contralateral uninjured nerve segments were processed for semithin/electron microscopy, or for immunofluorescence as described below.

Analysis of axonal survival in axotomized nerve stumps

Axon survival after unilateral sciatic nerve transection was determined using highly established methods for the evaluation of axonal integrity [4, 18, 29, 30]. Axon survival analysis 3 days post-axotomy was performed on semithin micrographs from randomly selected areas on sciatic nerve cross sections acquired with a DMI8 digital imaging system equipped with a 100x high-numerical aperture objective. One micrograph per mouse nerve was quantified. On each micrograph, the number of all structurally intact myelinated axons was manually counted using the Cell counter plug-in of the ImageJ software. Axon survival was expressed as percentage of intact axons relative to the total number of intact myelinated axons in corresponding uninjured control nerve micrographs. Axon survival at 30-36h after axotomy was assessed on montaged tile scan images of entire tibial nerves, acquired with a Leica DMI8 digital imaging system as described above, and similar to our previously published work [4]. Because Wallerian degeneration at such early stages following axotomy is asynchronous resulting in substantial heterogeneity of axons disintegration within a nerve [41], this method prevents the sampling bias that is associated with quantification of axons from individual micrographs taken from select areas of the nerve. Criteria for structurally intact axons after axotomy were uniform axoplasm with presence of non-swollen mitochondria and normal myelin sheaths. Criteria for degenerated axons were degraded axoplasm, absence of mitochondria and collapsed myelin sheaths. All structurally intact and all degenerated myelinated axons were counted on each tibial nerve montage, and their ratio was determined for each experimental animal (relative axon survival). Scoring was documented with the ImageJ software using the Cell counter plug-in. The experimenter was blind to the genotypes of the mice during data acquisition. Relative axonal survival is reported as the percentage of the respective control group.

Analysis of myelin remodeling in axotomized nerve stumps

Quantification of the area occupied by myelin (sheaths and myelin debris) in transverse semithin sections of axotomized sciatic and tibial nerves was performed using ImageJ. This analysis leverages the high contrast provided by osmium tetroxide and toluidine blue staining of lipid-rich myelin, which enables reliable thresholding and binarization of myelin structures on nerve cross sections. The binarized images were used to calculate the proportion of the nerve cross-sectional area occupied by myelin. We used axotomized nerves from mice lacking c-Jun in Schwann cells (c-Jun^{loxP/loxP}, p0^{Cre}) with suppressed Schwann cell transdifferentiation as a positive reference

which show abnormal myelin remodeling and clearance during Wallerian degeneration. This results in significantly larger myelin areas occupied during Wallerian degeneration as compared to control preparations.

c-Jun immunofluorescence and quantification

Segments of control and injured sciatic nerves were immersion fixed in 4% paraformaldehyde/0.1M PBS for 1h at 4°C, washed in PBS, cryoprotected in 30% sucrose, embedded in O.C.T. compound, and sectioned at 12 µm on a Leica cryostat (Leica Biosystems). The cross sections on adhesive glass slides were washed with TBS, permeabilized with 0.25% Triton X-100 and 0.5M NH₄Cl in 0.05M TBS for 10 min, blocked with 5% goat serum in 0.05M TBS for 1 h, and then incubated with anti-c-Jun antibody (1:200, catalog no. 9165, Cell Signaling Technology) diluted in 5% goat serum/0.05M TBS overnight at 4 °C. After washing in TBS, anti-rabbit secondary antibody coupled to Alexa Fluor 488 (1:500, catalog no. 115-545-003) was applied for 1 h, the sections counterstained with DAPI (1:10,000, catalog no. 62247, Thermo Fisher Scientific) and mounted in VECTASHIELD Antifade Mounting Medium. Micrographs of the stained cross sections were captured with an Andor Dragonfly 200 spinning disc confocal imaging system (Oxford Instruments) and a 63x high numerical aperture objective. Adjustments of brightness and contrast were applied with ImageJ and Microsoft Powerpoint equally across the entire image and were applied equally to control preparations for all the data presented. Two representative maximum intensity z-projection images from randomly selected equal volumes were taken per nerve cross section and used for the subsequent quantification. The quantification of c-Jun⁺/DAPI⁺ cells versus c-Jun⁻/DAPI⁺ cells used to calculate the percentage of cells showing c-Jun immunoreactivity was carried out blind to the mouse genotypes.

Preparation of mouse embryonic dorsal root ganglia cultures

Embryonic dorsal root ganglia (eDRGs) were isolated from mouse E13.5 embryos. DR6^{+/-} females were timed-mated with DR6^{+/-} males, and embryos from each litter were processed individually and genotyped after eDRG plating. eDRGs from DR6^{+/-} embryos were used as control preparations. DR6^{loxP/loxP} females were time-mated with DR6^{loxP/loxP} males, and all extracted embryos from these litters were pooled together for eDRG isolation. eDRGs were dissociated and plated as previously described [42]. A suspension of about 25,000 cells/µl was plated as a 2µl droplet into a well within a 24-well plate. Neurons were cultured on poly-D-lysine and laminin coated wells in Neurobasal medium supplemented with Glutamine (2mM), Penicillin/Streptomycin, serum-free B27, nerve growth factor (NGF, 50ng/ml), and the anti-mitotic agents uridine and 5-Fluoro-2'-deoxyuridine (1µM) to inhibit non-neuronal cell proliferation. At 2 days *in-vitro* (DIV 2), neurons from DR6^{loxP/loxP} embryos were infected with either a control lentivirus (FCIV), or a lentivirus expressing Cre recombinase (CRE), both based on the same vector backbone and co-expressing EGFP, as previously described [16]. Infection efficiency was monitored via fluorescence microscopy based on EGFP expression.

Analysis of *in vitro* axon degeneration

At DIV 7, when eDRG neurites had developed an extended radial arborization pattern, neurites were transected from their neuronal cell bodies using a microblade under microscopic guidance. The region distal to the transection site was imaged by phase contrast at the indicated time points with a Leica DMI8 digital imaging system and processed as previously described [42]. The micrographs were analyzed in ImageJ using a previously published degeneration index (DI) macro [43]. This macro binarizes the images and calculates the area occupied by fragmented neurites, normalized to the total neurite area. The resulting DI value models the extent of structural axon disintegration during Wallerian degeneration, with values above 0.8 reflecting complete axonal disintegration. For the comparison of DR6^{+/-} and DR6^{-/-} embryos, the time-course of neurite fragmentation for each embryo was obtained by averaging the DI values calculated from multiple phase contrast micrographs taken from each cell culture well at each time point. For the

comparison of lentiviral infected neurons from DR6^{loxP/loxP} embryos, the time-course of neurite fragmentation in each eDRG cell culture well preparation was obtained by averaging the DI values calculated from multiple phase contrast micrographs taken from each cell culture well.

Statistics and reproducibility

All shown micrographs (semithin and electron microscopy, immunofluorescence, phase contrast microscopy) are representative of at least three biological replicates (mice or cell culture preparations). All statistical analyses and data visualization were performed using Graphpad Prism software. An unpaired, two-tailed Student's t-test was used to compare two groups. A two-way analysis of variance (ANOVA) with Sidak's multiple comparisons test was used to compare more than two groups. Data are presented as arithmetic mean $\pm$ standard error of the mean (SEM). Statistical differences were considered to be significant when $P < 0.05$.

Supplementary figures




Figure S1. Disrupted DR6 expression in two DR6 knockout mouse models. (A, C) Quantitative real-time PCR analysis of relative brain DR6 mRNA levels from control and mutant mice with the indicated genotypes. Each dot represents the measurement from one mouse. (B, D) Western blot analysis (cropped blot images) of brain lysates from control and mutant mice with the indicated genotypes probed with the shown antibodies. Each western blot lane represents the brain lysate data from one individual mouse.




Figure S2. Normal developmental myelination in sciatic nerves from mice lacking DR6 at postnatal day 1.

(A, C) Representative semithin micrographs (A, C) of transverse sciatic nerve sections from control and mutant mouse pups at postnatal day 1 with the indicated genotypes. Red arrows depict examples of axons with nascent myelination. Scale bars: 10 μm . (B, D) Quantification of myelinated axons in transverse sciatic nerve sections from control and mutant pups at postnatal day 1 with the indicated genotypes. Each dot represents the quantification obtained from one mouse pup.




Figure S3. Normal nerve histomorphometry in DR6-deficient mice.

(A, C) Quantification of g ratios (left: scatter plots show g ratios of individual myelinated axons as function of axon diameter, right: corresponding cumulative g ratios per animal) in tibial (A) and sciatic nerves (C) from 3-months-old control and mutant mice with the indicated genotypes. (B, D) Axon caliber and fiber caliber (axon plus myelin sheath) distributions in tibial (B) and sciatic nerves (D) from 3-months-old control and mutant mice with the indicated genotypes. Each dot represents the measurement from one mouse.




Figure S4. Normal JNK activation in sciatic nerves from DR6-deficient mice.

(A, C) Representative western blots (cropped blot images) of lysates from uninjured (0 min) and axotomized distal sciatic nerve stumps (30 min after axotomy) from 1-2-months-old control and mutant mice with the indicated genotypes probed with the shown antibodies. Each lane represents the result from one mouse sciatic nerve. (B, D) Densitometric quantifications with statistical analysis of western blot data for all experimental animals used. Each dot represents a measurement from sciatic nerve lysate from one mouse.




Figure S5. Normal ultrastructure of transdifferentiated Schwann cells and normal myelin remodeling dynamics during Wallerian degeneration in DR6-deficient mice.

(A) Representative electron micrographs of transdifferentiated Schwann cells (red arrowheads) that lost axonal contact from distal sciatic nerve stumps 3 days following nerve transection in 3-months-old control and DR6 knockout mice with the shown genotypes. Note similar ultrastructure with marked cytoplasmic expansion, increased organelle content, and myelin debris (myelin ovoids) in Schwann cell bodies in all genotypes. Scale bar: 2µm. (B, C) Quantification of area occupied by myelin sheaths and myelin debris in transverse nerve sections from 3-months-old control and mutant mice with the indicated genotypes. Each dot represents the quantification obtained from one nerve.

Acknowledgements

We thank Genentech and Christopher Deppmann (University of Virginia) for providing the constitutive and conditional DR6 mutant mice, Konstantinos Tsemmis for technical experimental assistance, Ekaterina Stepanova and Alessio Colombo for genotyping information, Devin Donich for help with the g-ratio measurements, and Sara Bombardelli and the University Laboratory and Animal Resources (ULAR) at The Ohio State University College of Medicine for mouse husbandry assistance. This study was supported by the NIH-NINDS grant R01NS123450 (to E.B. and B.B.).

Additional information

Data availability

Figs. 1-4, S1-5 - source data contain the numerical data used to generate the figures.

Funding

Funder	Grant reference number	Author
HHS NIH National Institute of Neurological Disorders and Stroke (NINDS)	R01NS123450	Bogdan Beirowski Elisabetta Babetto Haoran Huang

Author ORCID iDs

Bogdan Beirowski: <https://orcid.org/0000-0002-1241-1777>

Additional files

Figure 1—source data 1. [↗](#)

Figure 2—source data 1. [↗](#)

Figure 3—source data 1. [↗](#)

Figure 4—source data 1. [↗](#)

Figure S1—source data 1. [↗](#)

Figure S2—source data 1. [↗](#)

Figure S3—source data 1. [↗](#)

Figure S4—source data 1. [↗](#)

Figure S5—source data 1. [↗](#)

References

1. Coleman M.P., Hoke A. (2020) Programmed axon degeneration: from mouse to mechanism to medicine. *Nat Rev Neurosci* **21**:183-196
2. Krauss R., et al. (2020) Axons Matter: The Promise of Treating Neurodegenerative Disorders by Targeting SARM1-Mediated Axonal Degeneration. *Trends Pharmacol Sci* **41**:281-293
3. Wilson D.M., et al. (2023) Hallmarks of neurodegenerative diseases. *Cell* **186**:693-714
4. Babetto E., Wong K.M., Beirowski B. (2020) A glycolytic shift in Schwann cells supports injured axons. *Nat Neurosci* **23**:1215-1228
5. Gamage K.K., et al. (2017) Death Receptor 6 Promotes Wallerian Degeneration in Peripheral Axons. *Curr Biol* **27**:890-896
6. Hu Y., et al. (2013) A DR6/p75(NTR) complex is responsible for beta-amyloid-induced cortical neuron death. *Cell Death Dis* **4**:e579

7. Huang G., et al. (2013) Death receptor 6 (DR6) antagonist antibody is neuroprotective in the mouse SOD1G93A model of amyotrophic lateral sclerosis. *Cell Death Dis* **4**:e841
8. Mishra V., et al. (2020) Systematic elucidation of neuron-astrocyte interaction in models of amyotrophic lateral sclerosis using multi-modal integrated bioinformatics workflow. *Nat Commun* **11**:5579
9. Wang Y., et al. (2015) Death Receptor 6 and Caspase-6 Regulate Prion Peptide-Induced Axonal Degeneration in Rat Spinal Neurons. *J Mol Neurosci* **56**:966-976
10. Conforti L., Gilley J., Coleman M.P. (2014) Wallerian degeneration: an emerging axon death pathway linking injury and disease. *Nat Rev Neurosci* **15**:394-409
11. Colombo A., et al. (2018) Non-cell-autonomous function of DR6 in Schwann cell proliferation. *EMBO J* **37**
12. Yang J., et al. (2015) Pathological axonal death through a MAPK cascade that triggers a local energy deficit. *Cell* **160**:161-76
13. Jessen K.R., Mirsky R. (2019) The Success and Failure of the Schwann Cell Response to Nerve Injury. *Front Cell Neurosci* **13**:33
14. Arthur-Farraj P.J., et al. (2012) , *c-Jun reprograms Schwann cells of injured nerves to generate a repair cell essential for regeneration*. *Neuron* **75**:633-47
15. Nikolaev A., et al. (2024) Retraction Note: APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature* **625**
16. Babetto E., et al. (2013) The Phr1 ubiquitin ligase promotes injury-induced axon self-destruction. *Cell Rep* **3**:1422-9
17. Babetto E., et al. (2010) Targeting NMNAT1 to Axons and Synapses Transforms Its Neuroprotective Potency In Vivo. *J Neurosci* **30**:13291-304
18. Beirowski B., et al. (2009) Non-nuclear Wld(S) determines its neuroprotective efficacy for axons and synapses in vivo. *J Neurosci* **29**:653-68
19. Gerdts J., et al. (2015) SARM1 activation triggers axon degeneration locally via NAD(+) destruction. *Science* **348**:453-7
20. Osterloh J.M., et al. (2012) dSarm/Sarm1 is required for activation of an injury-induced axon death pathway. *Science* **337**:481-4
21. Milde S., et al. (2013) Deletions within its subcellular targeting domain enhance the axon protective capacity of Nmnat2 in vivo. *Sci Rep* **3**:2567
22. Marik S.A., et al. (2013) Death receptor 6 regulates adult experience-dependent cortical plasticity. *J Neurosci* **33**:14998-5003
23. Nikolaev A., et al. (2009) APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature* **457**:981-9
24. Kallop D.Y., et al. (2014) A death receptor 6-amyloid precursor protein pathway regulates synapse density in the mature CNS but does not contribute to Alzheimer's disease-related pathophysiology in murine models. *J Neurosci* **34**:6425-37
25. Olsen O., et al. (2014) Genetic analysis reveals that amyloid precursor protein and death receptor 6 function in the same pathway to control axonal pruning independent of beta-secretase. *J Neurosci* **34**:6438-47
26. Fernandes K.A., et al. (2018) Role of SARM1 and DR6 in retinal ganglion cell axonal and somal degeneration following axonal injury. *Exp Eye Res* **171**:54-61
27. Lunn E.R., et al. (1989) Absence of Wallerian Degeneration does not Hinder Regeneration in Peripheral Nerve. *Eur J Neurosci* **1**:27-33
28. Vargas M.E., Barres B.A. (2007) Why is Wallerian degeneration in the CNS so slow?. *Annu Rev Neurosci* **30**:153-79

29. Mack T.G., et al. (2001) Wallerian degeneration of injured axons and synapses is delayed by a Ube4b/Nmnat chimeric gene. *Nat Neurosci* **4**:1199-1206
30. Beirowski B., et al. (2010) WldS can delay Wallerian degeneration in mice when interaction with valosin-containing protein is weakened. *Neuroscience* **166**:201-11
31. Conforti L., et al. (2009) WldS protein requires Nmnat activity and a short N-terminal sequence to protect axons in mice. *J Cell Biol* **184**:491-500
32. Perry V.H., et al. (1990) Evidence that the Rate of Wallerian Degeneration is Controlled by a Single Autosomal Dominant Gene. *Eur J Neurosci* **2**:408-413
33. Mi S., et al. (2011) Death receptor 6 negatively regulates oligodendrocyte survival, maturation and myelination. *Nat Med* **17**:816-21
34. Schmidt C.S., et al. (2005) Resistance to myelin oligodendrocyte glycoprotein-induced experimental autoimmune encephalomyelitis by death receptor 6-deficient mice. *J Immunol* **175**:2286-92
35. Riccomagno M.M., Kolodkin A.L. (2015) Sculpting neural circuits by axon and dendrite pruning. *Annu Rev Cell Dev Biol* **31**:779-805
36. Luo L., O'Leary D.D. (2005) Axon retraction and degeneration in development and disease. *Annu Rev Neurosci* **28**:127-56
37. Zhao H., et al. (2001) Impaired c-Jun amino terminal kinase activity and T cell differentiation in death receptor 6-deficient mice. *J Exp Med* **194**:1441-8
38. Tam S.J., et al. (2012) Death receptors DR6 and TROY regulate brain vascular development. *Dev Cell* **22**:403-17
39. Beirowski B., et al. (2017) mTORC1 promotes proliferation of immature Schwann cells and myelin growth of differentiated Schwann cells. *Proc Natl Acad Sci U S A* **114**:E4261-E4270
40. Beirowski B., et al. (2011) Sir-two-homolog 2 (Sirt2) modulates peripheral myelination through polarity protein Par-3/atypical protein kinase C (aPKC) signaling. *Proc Natl Acad Sci U S A* **108**:E952-61
41. Beirowski B., et al. (2004) Quantitative and qualitative analysis of Wallerian degeneration using restricted axonal labelling in YFP-H mice. *J Neurosci Methods* **134**:23-35
42. Shin J.E., Cho Y. (2020) Assessing Axonal Degeneration in Embryonic Dorsal Root Ganglion Neurons In Vitro. *Methods Mol Biol* **2143**:41-54
43. Sasaki Y., et al. (2009) Nicotinamide mononucleotide adenylyl transferase-mediated axonal protection requires enzymatic activity but not increased levels of neuronal nicotinamide adenine dinucleotide. *J Neurosci* **29**:5525-35

Peer reviews

Reviewer #1 (Public review):

Summary:

The authors show that genetic deletion of the orphan tumor necrosis factor receptor DR6 in mice does not protect peripheral axons against degeneration after axotomy. Similarly, Schwann cells in DR6 mutant mice react to axotomy similarly to wild type controls. These negative results are important because previous work has indicated that loss or inhibition of DR6 is protective in disease models and also against Wallerian degeneration of axons following injury. This carefully executed counterexample is important for the field to consider.

Strengths:

A strength of the paper is the use of two independent mouse strains that knockout DR6 in slightly different ways. The authors confirm that DR6 mRNA is absent in these models

(western blots for DR6 protein are less convincingly null, but given the absence of mRNA, this is likely an issue of antibody specificity). One of the DR6 knockout strains used is the same strain used in a previous paper examining the effects of DR6 on Wallerian degeneration.

The authors use a series of established assays to evaluate axon degeneration, including light and electron microscopy on nerve histological samples and cultured dorsal root ganglion neurons in which axons are mechanically severed and degeneration is scored in time lapse microscopy. These assays consistently show a lack of effect of loss of DR6 on Wallerian degeneration in both mouse strains examined.

Additional strengths are that the authors examine both the axonal response and the Schwann cell response to axotomy and use both in vivo and in vitro assays.

Therefore, these experiments, the author's data support their conclusion that loss of DR6 does not protect against Wallerian degeneration.

Weaknesses:

A weakness of this paper is that no effort is made to determine why the results presented here may differ from previous studies. A notable possibility is that the original mouse strain that showed 5 of 13 mice being protected from Wallerian degeneration was studies on a segregating C57BL/6.129S background.

Finally, it is important to note that previously reported effects of DR6 inhibition, such as protection of cultured cortical neurons from beta-amyloid toxicity, are not necessarily the same as Wallerian degeneration of axons distal to an injury studied here. The negative results presented here showing that loss of DR6 is not protective against Wallerian degeneration induced by injury are important given the interest in DR6 as a therapeutic target. However, care should be taken in attempting to extrapolate these results to other disease contexts such as ALS or Alzheimer's disease.

<https://doi.org/10.7554/eLife.108389.2.sa2>

Reviewer #3 (Public review):

Summary:

The authors revisit the role of DR6 in axon degeneration following physical injury (Wallerian degeneration), examining both its effects on axons and its role in regulating the Schwann cell response to injury. Surprisingly, and in contrast to previous studies, they find that DR6 deletion does not delay the rate of axon degeneration after injury, suggesting that DR6 is not a mediator of this process.

Overall, this is a valuable study. As the authors note, the current literature on DR6 is inconsistent, and these results provide useful new data and clarification. This work will help other researchers interpret their own data and re-evaluate studies related to DR6 and axon degeneration.

Strengths:

(1) The use of two independent DR6 knockout mouse models strengthens the conclusions, particularly when reporting the absence of a phenotype.

(2) The focus on early time points after injury addresses a key limitation of previous studies. This approach reduces the risk of missing subtle protective phenotypes and avoids confounding results with regenerating axons at later time points after axotomy.

Comments on revisions:

I thank the authors for their thorough responses to my previous comments. The revisions have addressed the points raised and have improved the clarity and overall quality of the manuscript. I appreciate the effort taken to strengthen the presentation of the work.

<https://doi.org/10.7554/eLife.108389.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public review):

Summary:

The authors show that genetic deletion of the orphan tumor necrosis factor receptor DR6 in mice does not protect peripheral axons against degeneration after axotomy. Similarly, Schwann cells in DR6 mutant mice react to axotomy similarly to wild-type controls. These negative results are important because previous work has indicated that loss or inhibition of DR6 is protective in disease models and also against Wallerian degeneration of axons following injury. This carefully executed counterexample is important for the field to consider.

Strengths:

A strength of the paper is the use of two independent mouse strains that knock out DR6 in slightly different ways. The authors confirm that DR6 mRNA is absent in these models (western blots for DR6 protein are less convincingly null, but given the absence of mRNA, this is likely an issue of antibody specificity). One of the DR6 knockout strains used is the same strain used in a previous paper examining the effects of DR6 on Wallerian degeneration.

The authors use a series of established assays to evaluate axon degeneration, including light and electron microscopy on nerve histological samples and cultured dorsal root ganglion neurons in which axons are mechanically severed and degeneration is scored in time-lapse microscopy. These assays consistently show a lack of effect of loss of DR6 on Wallerian degeneration in both mouse strains examined.

Therefore, in the specific context of these experiments, the author's data support their conclusion that loss of DR6 does not protect against Wallerian degeneration.

Weaknesses:

(1) The major weaknesses of this paper include the tone of correcting previously erroneous results and the lack of reporting on important details around animal experiments that would help determine whether the results here really are discordant with previous studies, and if so, why.

The authors do not report the genetic strain background of the mice used, the sex distributions of their experimental cohorts, or the age of the mice at the time the experiments were performed. All of these are important variables.

(Response 1) We thank the reviewer for emphasizing the importance of reporting the sex, age, and genetic background of the experimental animals used in our axon protection analyses. We have incorporated this information into the revised manuscript wherever available. The sole exception concerns the genetic background of the conditional DR6 mice

generated by Genentech, which remains unknown. The original publication describing these mice (Tam et al., 2012, Dev Cell, PMID 22340501) did not report this information, and we were unable to obtain it directly from Genentech. Details regarding the genetic background of the Wld^S and aPhr1 mutant mice are provided in their respective original publications, which are cited in our manuscript. Because the Gamage et al. study from the Deppmann laboratory did not report the sex or age of the animals used, we were unable to assess whether these variables might contribute to the differences observed between the two studies. Moreover, we are not aware of published evidence identifying sex or age as modifiers of structural axon preservation in axotomized peripheral nerve stumps in mouse models of delayed Wallerian degeneration. Furthermore, in the original publications describing the phenotypes of transgenic Nmnat2 and Wld^S mice, as well as Sarm1 or Phr1 knockout mice, sex and age of the animals used in the Wallerian degeneration assays were not reported (PMIDs 23995269, 12106171, 22678360, 23665224). Although, to our knowledge, no large-scale systematic studies have been conducted, over the last 15 years we have never observed any sex-based differences in Wallerian degeneration phenotypes in these mutants exhibiting prominent axon protection. This topic was discussed informally at conferences, and we are also not aware of other investigators having observed such effects.

In response to the reviewer's comment regarding "tone", we made sure that our data and interpretations are presented in a professional, balanced, and objective manner, including a detailed discussion of potential alternative explanations for the discrepant findings.

(2) The DR6 knockout strain reported in Gamage et al. (2017) was on a C57BL/6.129S segregating background. Gamage et al. reported that loss of DR6 protected axons from Wallerian degeneration for up to 4 weeks, but importantly, only in 38.5% (5 out of 13) mice they examined. In the present paper, the authors speculate on possible causes for differences between the lack of effect seen here and the effects reported in Gamage et al., including possible spontaneous background mutations, epigenetic changes, genetic modifiers, neuroinflammation, and environmental differences. A likely explanation of the incomplete penetrance reported by Gamage et al. is the segregating genetic background and the presence of modifier loci between C57BL/6 and 129S. The authors do not report the genetic background of the mice used in this study, other than to note that the knockout strain was provided by the group in Gamage et al. However, if, for example, that mutation has been made congenic on C57BL/6 in the intervening years, this would be important to know. One could also argue that the results presented here are consistent with 8 out of 13 mice presented in Gamage et al.

(Response 2) As noted above, we now provide information on the genetic background of the mice in the revised manuscript, where available. We have not backcrossed the constitutive DR6 knockout mice obtained from the Deppmann laboratory (Gamage et al.) to a C57BL/6 background; our colony was maintained primarily through intercrosses of heterozygous animals. Similarly, the conditional DR6 mutant mice used in this study were also not backcrossed to C57BL/6 mice.

We respectfully hold a different view regarding the reviewer's final point. We understand it is not appropriate to infer consistency between two datasets by disregarding the subset of results that do not align. By the same logic, it would be flawed to draw conclusions from the Gamage et al. study based solely on the single Wld^S mouse out of five that did not show axon preservation after nerve injury. Selectively omitting conflicting data does not provide a valid basis for establishing phenotype concordance across studies.

To further strengthen our study, we note that we performed additional analyses on three more nerve samples from constitutive DR6 null mice during the revision process and have incorporated the resulting data in Fig. 1.

(3) Age is also an important variable. The protective effects of the spontaneous WldS mutation decrease with age, for example. It is unclear whether the possible protective effects of DR6 also change with age; perhaps this could explain the variable response seen in Gamage et al. and the lack of response seen here.

(Response 3) As discussed above, we now provide the age information for the mice used for the Wallerian degeneration assessments in the respective figure legends. To our knowledge, there are no prior reports suggesting that age is a significant determinant of structural axon preservation in the indicated mutants. Electrophysiological function and neuromuscular junction preservation decrease with age in axotomized Wld^S mice (e.g., PMIDs 12231635, 19158292, 15654865), but these parameters are not subject of our study, and we have not studied them. Unfortunately, a direct comparison of ages between our DR6 mutant mice and those used in Gamage et al. (2017) is not possible, as the earlier study from the Deppmann laboratory did not report this information.

(4) It is unclear if sex is a factor, but this is part of why it should be reported.

(Response 4) We now report the requested sex information for our axon preservation analyses during nerve injury-induced Wallerian degeneration in the DR6 mouse models in Figs. 1 and 2.

(5) The authors also state that they do not see differences in the Schwann cell response to injury in the absence of DR6 that were reported in Gamage et al., but this is not an accurate comparison. In Gamage et al., they examined Schwann cells around axons that were protected from degeneration 2 and 4 weeks post-injury. Those axons had much thinner myelin, in contrast to axons protected by WldS or loss of Sarm1, where the myelin thickness remained relatively normal. Thus, Gamage et al. concluded that the protection of axons from degeneration and the preservation of Schwann cell myelin thickness are separate processes. Here, since no axon protection was seen, the same analysis cannot be done, and we can only say that when axons degenerate, the Schwann cells respond the same whether DR6 is expressed or not.

(Response 5) We appreciate the reviewer's detailed comments. Our intention was not to directly compare our findings with those of Gamage et al. regarding the myelin behavior at these time points (because we never observed axon protection), but rather to note that we did not observe any DR6-dependent alterations in Schwann cell responses under conditions where axons undergo normal Wallerian degeneration. As the reviewer correctly points out, Gamage et al. analyzed Schwann cell myelin surrounding axons that were protected from degeneration for extended periods, a context fundamentally different from the complete lack of axon protection observed in our DR6-deficient models. Therefore, the specific dissociation between axon preservation and myelin maintenance claimed by Gamage et al. cannot be evaluated in our study. A statement to make this point clearer has been incorporated in the revised manuscript.

We fully agree with the reviewer's concluding point: in our experiments, once axons degenerate, Schwann cell responses proceed similarly regardless of DR6 expression. This agreement reinforces one of the central conclusions of our work.

(6) The authors also take issue with Colombo et al. (2018), where it was reported that there is an increase in axon diameter and a change in the g-ratio (axon diameter to fiber diameter - the axon + myelin) in peripheral nerves in DR6 knockout mice. This change resulted in a small population of abnormally large axons that had thinner myelin than one would expect for their size. The change in g-ratio was specific to these axons and driven by the increased axon diameter, not decreased myelin thickness, although those two factors are normally loosely correlated. Here, the authors report no changes in axon size or g-ratio, but this could also be due to how the distribution of axon sizes was binned

for analysis, and looking at individual data points in supplemental figure 3A, there are axons in the DR6 knockout mice that are larger than any axons in wild type. Thus, this discrepancy may be down to specifics and how statistics were performed or how histograms were binned, but it is unclear if the results presented here are dramatically at odds with the results in Colombo et al. (2018).

(Response 6) Several points raised by the reviewer appear to reflect differences in interpretation of the findings reported in Colombo et al. (2018). That study did not report altered myelination in DR6 null mice at stages when myelination is largely complete (P21). Instead, modest changes were observed at P1, which were reduced by P7, and P21 mutants were reported to be indistinguishable from controls. No analyses of peripheral nerves in older animals were presented, and the authors concluded in the discussion that myelination in young adult DR6 null mice appears normal. In contrast, our analysis of constitutive DR6 null mice at P1 does not reproduce the increase in the number of myelinated fibers per unit area reported by Colombo et al. We obtained similar results in the independent conditional DR6 knockout mouse line. Differences in nerve tissue processing, embedding, staining, or in the microscopic imaging and quantification of thinly myelinated axons in P1 sciatic nerve cross-sections may have contributed to the observed discrepancy. However, because the relevant methodological details were not described in Colombo et al., the underlying reasons for these differences cannot be determined and remain speculative.

(7) Finally, it is important to note that previously reported effects of DR6 inhibition, such as protection of cultured cortical neurons from beta-amyloid toxicity, are not necessarily the same as Wallerian degeneration of axons distal to an injury studied here. The negative results presented here, showing that loss of DR6 is not protective against Wallerian degeneration induced by injury, are important given the interest in DR6 as a therapeutic target, but they are specific to these mice and this mechanism of induced axon degeneration. The extent to which these findings contradict previous work is difficult to assess due to the lack of detail in describing the mouse experiments, and care should be taken in attempting to extrapolate these results to other disease contexts, such as ALS or Alzheimer's disease.

(Response 7) We agree with the reviewer's point and emphasize that our manuscript carefully differentiates our data regarding the function of DR6 in Wallerian degeneration from the potential involvement of DR6 in other forms of axon degeneration. Our findings do not conflict with previous work on DR6 in the context of *in vitro* beta-amyloid and prion toxicity as well as *in vitro* models of ALS and multiple sclerosis. We believe these distinctions are explicitly and appropriately articulated throughout the entire manuscript and in more detail in the discussion section.

Reviewer #1 (Recommendations for the authors):

(1) The authors should include additional information about the mice used, including strain background for both the DR6 mice and the Cre transgenes crossed into the DR6 conditional knockout, the age of the mice when the nerve crush experiments were performed, and the sex distributions of the experimental cohorts. This information is critical for reproducibility in animal experiments, and that point is compounded here, where the major focus of this paper is taking issue with the reproducibility of previous work.

(Response 8) This information has been included in the revision. See above responses.

(2) In the abstract, reference 5 is cited as a study on the response to Schwann cells to injury in a DR6 background, but this probably should be reference 10.

(Response 9) This typo has been corrected.

(3) *"Site-by-site comparison" in line 201 should be side-by-side?*

(Response 10) This typo has been corrected.

(4) *The paper contains a lot of self-evaluative wording, "surprising contrast," "compelling evidence," "robust results." Whether those adjectives apply should be for the reader to decide, and a drier, more objective tone in the presentation would improve the paper.*

(Response 11) We agree that excessive self-evaluative wording can weaken objectivity. In the manuscript, such phrasing is used sparingly and intentionally to highlight differences from previously published studies, guide the reader, and convey scholarly judgment. We do not consider this limited use to be counterproductive. The adjectives “surprising,” “compelling,” and “robust” each appear only one to three times across the entire manuscript, and the specific phrase “robust results” does not appear at all.

(5) *In Figure 2A, DR6^{-/-}, there is no significant difference, but there is also a lot of variability, and one could argue the authors are seeing axon protection comparable to WldS in 40% of their samples (2/5), which is very similar to Gamage et al.*

(Response 12) We respectfully disagree with this reasoning as it relies on selectively emphasizing only a subset of the data. Please also see our response #2 for more detailed discussion.

(6) *Overall, the data presented here are convincing and support the conclusions drawn, but the paper needs to focus more on the negative results at hand and less on bashing previous studies, particularly when the results presented here do definitively show that the previous studies were incorrect and plausible explanations for differences in outcome exist.*

(Response 13) We have carefully revisited the wording of the manuscript and are confident that our emphasis remains on the central negative finding that DR6 does not regulate axon degeneration and Schwann cell injury responses during Wallerian degeneration. We do not believe the manuscript “bashes” previous studies; nonetheless, we thoroughly re-examined all relevant sections to ensure that our language is neutral, accurate, and non-inflammatory. We believe the current phrasing presents our interpretations in an appropriately balanced, objective, and professional manner.

Reviewer #2 (Public review):

Summary:

This manuscript by Beirowski, Huang, and Babetto revisits the proposed role of Death Receptor 6 (DR6/Tnfrsf21) in Wallerian degeneration (WD). A prior study (Gamage et al., 2017) suggested that DR6 deletion delays axon degeneration and alters Schwann cell responses following peripheral nerve injury. Here, the authors comprehensively test this claim using two DR6 knockout mouse models (the line used in the earlier report plus a CMV-Cre derived floxed ko line) and multiple WD assays in vivo and in vitro, aligned with three positive controls, Sarm1 WldS and Phr1/Mycbp2 mutants. Contrary to the prior findings, they find no evidence that DR6 deletion affects axon degeneration kinetics or Schwann cell dynamics (assessed by cJun expression or [intact+degenerating] myelin abundance after injury) during WD. Importantly, in DRG explant assays, neurites from DR6-deficient mice degenerated at rates indistinguishable from controls. The authors conclude that DR6 is dispensable for WD, and that previously reported protective effects may have been due to confounding factors such as genetic background or spontaneous mutations.

Strengths:

The authors employ two independently generated DR6 knockout models, one overlapping with the previously published study, and confirm loss of DR6 expression by qPCR and Western blotting. Multiple complementary readouts of WD are applied (structural, ultrastructural, molecular, and functional), providing a robust test of the hypothesis.

Comparisons are drawn with established positive controls (Wld^S, SARM1, Phr1/Mycbp2 mutants), reinforcing the validity of the assays.

By directly addressing an influential but inconsistent prior report, the manuscript clarifies the role of DR6 and prevents potential misdirection of therapeutic strategies aimed at modulating WD in the PNS. The discussion thoughtfully considers possible explanations for the earlier results, including colony-specific second-site mutations that could explain the incomplete penetrance of the earlier reported phenotype of only 36%.

Weaknesses:

(1) The study focuses on peripheral nerves. The manuscript frequently refers to CNS studies to argue for consistency with their findings. It would be more accurate to frame PNS/CNS similarities as reminiscences rather than as consistencies (e.g., line 205ff in the Discussion).

(Response 14) Axon protection in all key genetic models of delayed axon degeneration, including Wld^S, SARM1, Phr1/Mycbp2 mutants, has been demonstrated in both the peripheral and central nervous systems. This observation supports the view that core molecular mechanisms regulating axon degeneration are conserved across neuronal populations throughout the entire nervous system. We have scrutinized the wording in our manuscript and are not aware that we frequently refer to CNS studies in regards to axon degeneration. Nevertheless, we have replaced the term “consistent” to avoid potential ambiguity when we discuss the earlier study showing normal Wallerian degeneration in the optic nerves from DR6 knockout mice.

(2) The DRG explant assays are convincing, though the slight acceleration of degeneration in the DR6 floxed/Cre condition is intriguing (Figure 4E). Could the authors clarify whether this is statistically robust or biologically meaningful?

(Response 15) We thank the reviewer for noting this aspect of our *in vitro* data in Fig. 4. The difference observed in the DR6 floxed/Cre condition is statistically significant at the 6h time point following disconnection, as indicated by the p value shown in Fig. 4E. However, a similarly statistically significant acceleration of axon degeneration was not observed in DRG axotomy experiments using constitutive DR6 knockout preparations, although a trend toward more rapid axon breakdown is apparent at 6 h post-axotomy (Fig. 4B). These observations may suggest reduced stability of DR6-deficient axons in this specific neuron-only *in vitro* context. Further investigation would be required to determine the biological significance of this effect. In contrast, our *in vitro* quantitative analyses of the initiation and early phases of Wallerian degeneration (Fig. 2) revealed no evidence of accelerated axon disintegration in the DR6 mutant mouse models, highlighting potential differences between *in vitro* and *in vitro* systems.

(3) In the summary (line 43), the authors refer to Hu et al. (2013) (reference 5) as the study that previously reported AxD delay and SC response alteration after injury. However, this study did not investigate the PNS, and I believe the authors intended to reference Gamage et al. (2017) (reference 10) at this point.

(Response 16) Thanks for pointing this out. We have corrected this typo in the revised manuscript.

(4) In line 74ff of the results section, the authors claim that developmental myelination is not altered in DR6 mutants at postnatal day 1. However, the variability in Figure S2 appears substantial, and the group size seems underpowered to support this claim. Colombo et al. (2018) (reference 11) reported accelerated myelination at P1, but this study likewise appears underpowered. Possible reasons for these discrepancies and the large variability could be that only a defined cross-sectional area was quantified, rather than the entire nerve cross-section.

(Response 17) We confirm that the quantification of thinly myelinated axons was performed on entire sciatic nerves from P1 mouse pups, as described in the methods section in our original manuscript. The data shown in Fig. S2 were obtained from 5-9 pups per experimental group. Sample sizes were determined based on a priori power analyses using pilot data, which indicated that a minimum of five biological replicates was sufficient to detect statistically significant differences with acceptable confidence. Comparable sample sizes have been used in our previous studies and by other groups to assess early postnatal myelination (e.g., PMIDs 21949390, 28484008). Several published studies have reported analyses using 3-4 animals per group (e.g., PMIDs 28484008, 25310982, 29367382). For comparison, the study by Colombo et al. used 3-8 pups for the analysis presented in their Fig. 3. We note that the apparent variability in Fig. S2 may be accentuated by the scaling of the y-axis, which was chosen to ensure that individual data points are clearly resolved and visible.

(5) The authors stress the data of Gamage et al. (2017) on altered SC responses in DR6 mutants after injury. They employed cJun quantification to show that SC reprogramming after injury is not altered in DR6 mutants. This approach is valid and the conclusion trustworthy. Here, the addition of data showing the combined abundance of intact and degenerated myelin does not add much insight. However, Gamage et al. (2017) reported altered myelin thickness in a subset of axons at 14 days after injury, which is considerably later than the time points analyzed in the present study. While, in the Reviewer's view, the thin myelin observed by Gamage et al. in fact resembles remyelination, the authors may wish to highlight the difference in the time points analyzed.

(Response 18) We consider the additional quantification of the area occupied by intact myelin and myelin debris to provide complementary information that supports the c-Jun-based conclusion that Schwann cell injury responses are normal in DR6-deficient nerves following lesion. We agree with this reviewer that the thin myelin observed by Gamage et al. resembles remyelination, raising the possibility that axon regeneration occurred into the distal nerve stump at the studied 14d post-injury time point (see their Fig. 3). This may have been interpreted as axon protection in this study. In our study, it was impossible to examine such myelin effects since axon protection was never observed in any of the DR6 mutant models at any of the time point we investigated. We have incorporated appropriate additional text to highlight this difference. See also response #5 above.

Reviewer #3 (Public review):

Summary:

The authors revisit the role of DR6 in axon degeneration following physical injury (Wallerian degeneration), examining both its effects on axons and its role in regulating the Schwann cell response to injury. Surprisingly, and in contrast to previous studies, they find that DR6 deletion does not delay the rate of axon degeneration after injury, suggesting that DR6 is not a mediator of this process.

Overall, this is a valuable study. As the authors note, the current literature on DR6 is inconsistent, and these results provide useful new data and clarification. This work will

help other researchers interpret their own data and re-evaluate studies related to DR6 and axon degeneration.

Strengths:

(1) The use of two independent DR6 knockout mouse models strengthens the conclusions, particularly when reporting the absence of a phenotype.

(2) The focus on early time points after injury addresses a key limitation of previous studies. This approach reduces the risk of missing subtle protective phenotypes and avoids confounding results with regenerating axons at later time points after axotomy.

Weaknesses:

(1) The study would benefit from including an additional experimental paradigm in which DR6 deficiency is expected to have a protective effect, to increase confidence in the experimental models, and to better contextualize the findings within different pathways of axon degeneration. For example, DR6 deletion has been shown in more than one study to be partially axon protective in the NGF deprivation model in DRGs *in vitro*.

Incorporating such an experiment could be straightforward and would strengthen the paper, especially if some of the neuroprotective effects previously reported are confirmed.

(Response 19) We thank the reviewer for these suggestions. We would like to highlight that our study addresses the role of DR6 in Wallerian degeneration, whereas *in vitro* NGF deprivation has been used to model developmental axon pruning. Previous work indicates fundamental biological differences between these regressive pathways regulating the stereotyped removal of axon segments. We feel that studying this alternative form of axon degeneration is beyond the scope of the current work and could be addressed in a separate manuscript. Although additional tests will be needed, we note that our preliminary data using samples from both DR6 knockout mouse models suggest no axon protection after NGF-deprivation in DRG neuron preparations in our hands (deprivation of the growth factor and administration of anti-NGF antibody).

(2) The quality of some figures could be improved, particularly the EM images in Figure 2. As presented, they make it difficult to discern subtle differences.

(Response 20) We have pseudocolored intact (turquoise) and degenerated (magenta) myelinated fibers on the high-resolution semithin micrographs (not electron micrographs) in the new Fig. 2 to make the distinction between the two fiber categories clearer.

Reviewer #3 (Recommendations for the authors):

(1) Line 121: The authors mention toluidine blue staining, but it does not appear to be shown in Figure S5.

(Response 21) This appears to be a misunderstanding. Fig. S5A shows the ultrastructure of dedifferentiated Schwann cells in transmission electron micrographs, while Figs. S5B and C show quantification of the area occupied by myelin sheaths and myelin debris profiles on osmium tetroxide and toluidine blue stained nerve sections from the two DR6 mutant models, based on semithin light microscopy. These are two different aspects of the analysis. The text has been modified in the revised manuscript to make the distinction clearer.

(2) Line 175: The authors should add NMNAT2 to the list of enzymes implicated in the regulation of Wallerian degeneration in mammals.

(Response 22) Nmnat2 and a literature reference (Milde et al., 2013) has been incorporated in the discussion of the revised manuscript to address this point.

| (3) Line 201: Please correct the typo "site-by-site" to "side-by-side."

(Response 23) This typo has been corrected.

<https://doi.org/10.7554/eLife.108389.2.sa0>