

## The genome sequence of the colonial chordate, *Botryllus schlosseri*

Ayelet Voskoboynik<sup>1</sup>, Norma F Neff<sup>2</sup>, Debashis Sahoo<sup>3</sup>, Aaron M Newman<sup>4</sup>, Dmitry Pushkarev<sup>5</sup>, Winston Koh<sup>6</sup>, Benedetto Passarelli<sup>7</sup>, H Christina Fan<sup>8</sup>, Gary L Mantalas<sup>9</sup>, Karla J Palmeri<sup>10</sup>, Katherine J Ishizuka<sup>11</sup>, Carmela Gissi<sup>12</sup>, Francesca Griggio<sup>13</sup>, Rachel Ben-Shlomo<sup>14</sup>, Daniel M Corey<sup>15</sup>, Lolita Penland<sup>16</sup>, Richard A White III<sup>17</sup>, Irving L Weissman<sup>18</sup>, and Stephen R Quake<sup>19</sup>Ayelet Voskoboynik<sup>1</sup>, Norma F Neff<sup>2</sup>, Debashis Sahoo<sup>3</sup>, Aaron M Newman<sup>4</sup>, Dmitry Pushkarev<sup>5</sup>, Winston Koh<sup>6</sup>, Benedetto Passarelli<sup>7</sup>, H Christina Fan<sup>8</sup>, Gary L Mantalas<sup>9</sup>, Karla J Palmeri<sup>10</sup>, Katherine J Ishizuka<sup>11</sup>, Carmela Gissi<sup>12</sup>, Francesca Griggio<sup>13</sup>, Rachel Ben-Shlomo<sup>14</sup>, Daniel M Corey<sup>15</sup>, Lolita Penland<sup>16</sup>, Richard A White III<sup>17</sup>, Irving L Weissman<sup>18</sup>, and Stephen R Quake<sup>19</sup>


## **Abstract**


Botryllus schlosseri is a colonial urochordate that follows the chordate plan of development following sexual reproduction, but invokes a stem cellmediated budding program during subsequent rounds of asexual reproduction. As urochordates are considered to be the closest living invertebrate relatives of vertebrates, they are ideal subjects for whole genome sequence analyses. Using a novel method for high-throughput sequencing of eukaryotic genomes, we sequenced and assembled 580 Mbp of the B. schlosseri genome. The genome assembly is comprised of nearly 14,000 intron-containing predicted genes, and 13,500 intron-less predicted genes, 40% of which could be confidently parceled into 13 (of 16 haploid) chromosomes. A comparison of homologous genes between B. schlosseri and other diverse taxonomic groups revealed genomic events underlying the evolution of vertebrates and lymphoid-mediated immunity. The *B. schlosseri* genome is a community resource for studying alternative modes of reproduction, natural transplantation reactions, and stem cellmediated regeneration.

**DOI:** http://dx.doi.org/10.7554/eLife.00569.001

The tunicates are an evolutionary group that includes species such as sea squirts and sea tulips. Their name comes from the structure known as a















*eLife* 2() | DOI: http://dx.doi.org/10.7554/eLife.00569 | 02 July 20132013






























eLife 2() | DOI: http://dx.doi.org/10.7554/eLife.00569 | 02 July 20132013














eLife 2() | DOI: http://dx.doi.org/10.7554/eLife.00569 | 02 July 20132013
































 $\textit{eLife} \ 2() \ | \ \mathsf{DOI:http://dx.doi.org/10.7554/eLife.00569} \ | \ \mathsf{02} \ \mathsf{July} \ 20132013$ 











 $\textit{eLife}\ 2()\ |\ \mathsf{DOI}\ : \mathsf{http://dx.doi.org/10.7554/eLife.00569}\ |\ \mathsf{02}\ \mathsf{July}\ \mathsf{20132013}$ 



