

```

# -*- coding: utf-8 -*-
"""
Created on February 16, 2024
@author:
    Elif KARTAL, Istanbul University, Faculty of Economics, Department of
    Management Information Systems
    Beyaz Basak ESKISEHIRLI, Istanbul University, Faculty of Science,
    Department of Mathematics
    Fatma CALISKAN, Istanbul University, Faculty of Science, Department of
    Mathematics
    Zeki OZEN, Istanbul University, Faculty of Economics, Department of
    Management Information Systems

@title: Examples with p-adic calculations
"""

import p_adic

x = 25

# p_adic_val() function returns the p-adic absolute value of a given number.

x_p2 = p_adic.p_adic_val(x, p_num=2)

print("2-adic absolute value of ", x, " is ", x_p2)
2-adic absolute value of 25 is 1

x_p3 = p_adic.p_adic_val(x, p_num=3)

print("3-adic absolute value of ", x, " is ", x_p3)
3-adic absolute value of 25 is 1

x_p5 = p_adic.p_adic_val(x, p_num=5)

print("5-adic absolute value of ", x, " is ", x_p5)
5-adic absolute value of 25 is 0.04

y = 16

y_p2 = p_adic.p_adic_val(y, p_num=2)

print("2-adic absolute value of ", y, " is ", y_p2)
2-adic absolute value of 16 is 0.0625

# p_adic_pow_val() function returns both the p-adic order and the p-adic
absolute value of the given number.

y_p2_ord, y_p2_val = p_adic.p_adic_ord_val(y, p_num=2)

print(y_p2_ord)
4

print(y_p2_val)
0.0625

```

```

y_p3 = p_adic.p_adic_val(y, p_num=3)

print("3-adic absolute value of ", y, " is ", y_p3)
3-adic absolute value of 16 is 1

y_p5 = p_adic.p_adic_val(y, p_num=5)

print("5-adic absolute value of ", y, " is ", y_p5)
5-adic absolute value of 16 is 1

z = -16

z_p = p_adic.p_adic_val(z, p_num=5)

print("5-adic absolute value of ", z, " is ", z_p)
5-adic absolute value of -16 is 1

t = 0

print("2-adic absolute value of ", t, " is ", p_adic.p_adic_val(t, p_num=2))
2-adic absolute value of 0 is 0

print("3-adic absolute value of ", t, " is ", p_adic.p_adic_val(t, p_num=3))
3-adic absolute value of 0 is 0

print("5-adic absolute value of ", t, " is ", p_adic.p_adic_val(t, p_num=5))
5-adic absolute value of 0 is 0

print("17-adic absolute value of ", t, " is ", p_adic.p_adic_val(t,
p_num=17))
17-adic absolute value of 0 is 0

p = 1

print("2-adic absolute value of ", p, " is ", p_adic.p_adic_val(p, p_num=2))
2-adic absolute value of 1 is 1

print("3-adic absolute value of ", p, " is ", p_adic.p_adic_val(p, p_num=3))
3-adic absolute value of 1 is 1

print("5-adic absolute value of ", p, " is ", p_adic.p_adic_val(p, p_num=5))
5-adic absolute value of 1 is 1

print("17-adic absolute value of ", p, " is ", p_adic.p_adic_val(p,
p_num=17))
17-adic absolute value of 1 is 1

r1 = -54

print("3-adic absolute value of ", r1, " is ", p_adic.p_adic_val(r1,
p_num=3))
3-adic absolute value of -54 is 0.037037

```

```

s = -(24/16)

print("2-adic absolute value of ", s, " is ", p_adic.p_adic_val(s, p_num=2))
2-adic absolute value of  -1.5  is  2

print(p_adic.p_adic_ord_val(s, p_numb=2))
(-1, 2)

s1 = 8

print(p_adic.p_adic_ord_val(s1, p_numb=2))
(3, 0.125)

s2 = 100

print(p_adic.p_adic_ord_val(s2, p_numb=5))
(2, 0.04)

s3 = 0.270

print(p_adic.p_adic_ord_val(s3, p_numb=3))
(3, 0.037037)

s4 = 0.270

print(p_adic.p_adic_ord_val(s4, p_numb=5))
(-2, 25)

# p-parameter of p_adic_val() function must be prime!

m = 16

print(p_adic.p_adic_val(m, p_num=4))
p should be a prime number!
None

```