

Black-box, accurate, and efficient prediction of band structures with Koopmans functionals

Edward Linscott

Laboratory for Materials Simulations
Paul Scherrer Institute



PSI



DPG conference



21 March 2024



Accurate band structures

How can we calculate charged excitations (i.e. band structures, photoemission) accurately and efficiently?

Accurate band structures

How can we calculate charged excitations (i.e. band structures, photoemission) accurately and efficiently?

DFT intrinsic failure

Accurate band structures

How can we calculate charged excitations (i.e. band structures, photoemission) accurately and efficiently?

DFT intrinsic failure

GW accurate but more expensive;
diagrammatic


Accurate band structures

How can we calculate charged excitations (i.e. band structures, photoemission) accurately and efficiently?

DFT intrinsic failure

GW accurate but more expensive;
diagrammatic

Koopmans accurate band structures with a
functional theory



Accurate band structures


How can we calculate charged excitations (i.e. band structures, photoemission) accurately and efficiently?

DFT intrinsic failure

GW accurate but more expensive;
diagrammatic

Koopmans accurate band structures with a functional theory

	PBE	G_0W_0	KI	KIPZ	$QS\tilde{G}W$
E_{gap}	2.54	0.56	0.27	0.22	0.18
IP	1.09	0.39	0.19	0.21	0.49



$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

- a correction to DFT that ensures eigenvalues match total energy differences

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

- a correction to DFT that ensures eigenvalues match total energy differences
- is orbital-density-dependent (gives rise to, and is reliant on, a localised set of orbitals)

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

- a correction to DFT that ensures eigenvalues match total energy differences
- is orbital-density-dependent (gives rise to, and is reliant on, a localised set of orbitals)
- requires screening parameters α_i to be calculated

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

- a correction to DFT that ensures eigenvalues match total energy differences
- is orbital-density-dependent (gives rise to, and is reliant on, a localised set of orbitals)
- requires screening parameters α_i to be calculated

In order to evaluate this functional, one must...

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

- a correction to DFT that ensures eigenvalues match total energy differences
- is orbital-density-dependent (gives rise to, and is reliant on, a localised set of orbitals)
- requires screening parameters α_i to be calculated

In order to evaluate this functional, one must...

1. perform a Wannierisation

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

- a correction to DFT that ensures eigenvalues match total energy differences
- is orbital-density-dependent (gives rise to, and is reliant on, a localised set of orbitals)
- requires screening parameters α_i to be calculated

In order to evaluate this functional, one must...

1. perform a Wannierisation
2. calculate the screening parameters $\{\alpha_i\}$

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$

General features:

- a correction to DFT that ensures eigenvalues match total energy differences
- is orbital-density-dependent (gives rise to, and is reliant on, a localised set of orbitals)
- requires screening parameters α_i to be calculated

In order to evaluate this functional, one must...

1. perform a Wannierisation
2. calculate the screening parameters $\{\alpha_i\}$
3. minimize the functional

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{DFT}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) df + f_i \eta_i \right)$$


General features:

- a correction to DFT that ensures eigenvalues match total energy differences
- is orbital-density-dependent (gives rise to, and is reliant on, a localised set of orbitals)
- requires screening parameters α_i to be calculated

In order to evaluate this functional, one must...


1. perform a Wannierisation
2. calculate the screening parameters $\{\alpha_i\}$
3. minimize the functional
4. diagonalize the Hamiltonian

What needs to be in the box?



For a black box, we need...


What needs to be in the box?



For a black box, we need...

- automated start-to-finish calculations

What needs to be in the box?




For a black box, we need...

- automated start-to-finish calculations
- minimal input required from the user

koopmans

- v1.0 released last year¹
- implementations of Koopmans functionals within Quantum ESPRESSO
- automated workflows
 - Koopmans calculations
 - Wannierisation
 - dielectric tensor
 - ...
- built on top of ASE²
- does not require expert knowledge

koopmans-functionals.org



¹ E. B. Linscott et al. *J. Chem. Theory Comput.* 19.20 (2023), 7097

² A. H. Larsen et al. *J. Phys. Condens. Matter* 29.27 (2017), 273002

koopmans: An Open-Source Package for Accurately and Efficiently Predicting Spectral Properties with Koopmans Functionals

Edward B. Linscott,*[△] Nicola Colonna,[△] Riccardo De Gennaro, Ngoc Linh Nguyen, Giovanni Borghi, Andrea Ferretti, Ismaila Dabo, and Nicola Marzari*



Cite This: <https://doi.org/10.1021/acs.jctc.3c00652>



Read Online


ACCESS |

Metrics & More


Article Recommendations

Supporting Information

ABSTRACT: Over the past decade we have developed Koopmans functionals, a computationally efficient approach for predicting spectral properties with an orbital-density-dependent functional framework. These functionals impose a generalized piecewise linearity condition to the entire electronic manifold, ensuring that orbital energies match the corresponding electron removal/addition energy differences (in contrast to semilocal DFT, where a mismatch between the two lies at the heart of the band gap




What needs to be in the box?



For a black box, we need...

- automated start-to-finish calculations
- minimal input required from the user

What needs to be in the box?



For a black box, we need...


- automated start-to-finish calculations
- minimal input required from the user

One step still very manual: Wannierisation
Can we automate it?

One step still very manual: Wannierisation


Can we automate it?

- use PAOs found in pseudopotential files as initial guesses for Wannier functions




One step still very manual: Wannierisation
Can we automate it?

- use PAOs found in pseudopotential files as initial guesses for Wannier functions
- projectability-based disentanglement instead of energy-based disentanglement




One step still very manual: Wannierisation
Can we automate it?


- use PAOs found in pseudopotential files as initial guesses for Wannier functions
- projectability-based disentanglement instead of energy-based disentanglement
- use a parallel transport algorithm to separate the occupied and empty manifolds



Automating Wannierisation



Automating Wannierisation



```
{  
    "workflow": {  
        "task": "wannierize",  
        "init_orbitals": "mlwfs",  
        "pseudo_library": "pseudo_dojos_standard_v0.4.1"},  
    "atoms": {  
        "cell_parameters": {  
            "ibrav": 6,  
            "celldms": {"1": 8.675923, "3": 0.645248},  
            "periodic": true},  
        "atomic_positions": {  
            "positions": [  
                ["Ti", 0.5, 0.5, 0.5],  
                ["Ti", 0.0, 0.0, 0.0],  
                ["O", 0.1814, 0.8186, 0.5],  
                ["O", 0.8186, 0.1814, 0.5],  
                ["O", 0.3186, 0.3186, 0.0],  
                ["O", 0.6814, 0.6814, 0.0]  
            ],  
            "units": "crystal"}},  
    "kpoints": {"grid": [3, 3, 4]},  
    "calculator_parameters": {  
        "w90": {"auto_projections": true}}}}
```

A drawback: the number of Wannier functions is determined by the number of PAOs in the pseudopotentials

A drawback: the number of Wannier functions is determined by the number of PAOs in the pseudopotentials... which can be a problem e.g. LiF


A drawback: the number of Wannier functions is determined by the number of PAOs in the pseudopotentials... which can be a problem e.g. LiF

element	configuration	PAOs
Li	$1s^2 2s^1$	1s, 2s
F	$[1s^2] 2s^2 2p^5$	2s, 2p _x , 2p _y , 2p _z

Automating Wannierisation

A drawback: the number of Wannier functions is determined by the number of PAOs in the pseudopotentials... which can be a problem e.g. LiF

element	configuration	PAOs
Li	$1s^2 2s^1$	$1s, 2s$
F	$[1s^2] 2s^2 2p^5$	$2s, 2p_x, 2p_y, 2p_z$




Automating Wannierisation


A drawback: the number of Wannier functions is determined by the number of PAOs in the pseudopotentials... which can be a problem e.g. LiF

element	configuration	PAOs
Li	$1s^2 2s^1$	$1s, 2s$
F	$[1s^2] 2s^2 2p^5$	$2s, 2p_x, 2p_y, 2p_z$

If we want a better representation of the conduction bands, we're gonna need a bigger boat more PAOs...




Automating Wannierisation



For cases such as LiF:

- re-solve the radial Schrödinger equation for higher subshells


Automating Wannierisation



For cases such as LiF:

- re-solve the radial Schrödinger equation for higher subshells
- apply a confining potential while doing so


Automating Wannierisation



For cases such as LiF:

- re-solve the radial Schrödinger equation for higher subshells
- apply a confining potential while doing so
- optimize the confining potential to maximise projectability


Automating Wannierisation





For cases such as LiF:

- re-solve the radial Schrödinger equation for higher subshells
- apply a confining potential while doing so
- optimize the confining potential to maximise projectability
- N.B. only once per pseudopotential


Automating Wannierisation




Automating Wannierisation



Automating Wannierisation






Koopmans functionals can accurately and efficiently predict the band structure of materials

To make these calculations black box, we have...

- automated start-to-finish calculations


- minimal input required from the user



Koopmans functionals can accurately and efficiently predict the band structure of materials

To make these calculations black box, we have...

- automated start-to-finish calculations *with the koopmans package*
- minimal input required from the user



Koopmans functionals can accurately and efficiently predict the band structure of materials

To make these calculations black box, we have...

- automated start-to-finish calculations *with the koopmans package*
- minimal input required from the user *enabled via automated Wannierisation*

Acknowledgements



Nicola Marzari



Nicola Colonna



Junfeng Qiao



**Swiss National
Science Foundation**

MARVEL
The logo for MARVEL consists of four red hexagons arranged in a horizontal row, with the second and third hexagons partially overlapping.

NATIONAL CENTRE OF COMPETENCE IN RESEARCH

Want to find out more? Go to koopmans-functionals.org


Follow  [@ed_linscott](https://twitter.com/ed_linscott) for updates | Slides available at  github/elinscott

SPARE SLIDES


Consequences of ODD:

Consequences of ODD:

- variational (localized, minimizing) vs canonical (delocalized, diagonalizing) orbitals




(a) variational




(b) canonical

Consequences of ODD:

- variational (localized, minimizing) vs canonical (delocalized, diagonalizing) orbitals



(a) variational




(b) canonical


- Practically we can often use MLWFs

Consequences of ODD:

- variational (localized, minimizing) vs canonical (delocalized, diagonalizing) orbitals




(a) variational




(b) canonical

- Practically we can often use MLWFs
- localized variational orbitals naturally allow us to treat bulk systems

Localised states




Localised states



In the bulk limit for one cell $\Delta E_{\text{one cell}} = E(N - \delta N) - E(N)$

Localised states




In the bulk limit for one cell $\Delta E_{\text{one cell}} = E(N - \delta N) - E(N)$

Across all the cells $\Delta E_{\text{all cells}} = \frac{1}{\delta N} (E(N - \delta N) - E(N)) = -\frac{dE}{dN} = -\varepsilon_{\text{HO}}$


Workflows

(a) finite difference calculations using a supercell



Workflows

(a) finite difference calculations using a supercell




(b) DFPT using a primitive cell




Workflows

(a) finite difference calculations using a supercell



(b) DFPT using a primitive cell



All implemented in Koopmans

Accurate band structures


	PBE	$G_0W_0^1$	scGW ²	KI@[PBE,MLWFs]	KIPZ@PBE	exp ³
E_g	0.49	1.06	1.14	1.16	1.15	1.17
$\Gamma_{1v} \rightarrow \Gamma_{25'v}$	11.97	12.04		11.97	12.09	12.5 ± 0.6
$X_{1v} \rightarrow \Gamma_{25'v}$	7.82			7.82		7.75
$X_{4v} \rightarrow \Gamma_{25'v}$	2.85	2.99		2.85	2.86	2.90
$L_{2'v} \rightarrow \Gamma_{25'v}$	9.63	9.79		9.63	9.74	9.3 ± 0.4
$L_{1v} \rightarrow \Gamma_{25'v}$	6.98	7.18		6.98	7.04	6.8 ± 0.2
$L_{3'v} \rightarrow \Gamma_{25'v}$	1.19	1.27		1.19		1.2 ± 0.2
$\Gamma_{25'v} \rightarrow \Gamma_{15c}$	2.48	3.29		3.17	3.20	3.35 ± 0.01
$\Gamma_{25'v} \rightarrow \Gamma_{2'c}$	3.28	4.02		3.95	3.95	4.15 ± 0.05
$\Gamma_{25'v} \rightarrow X_{1c}$	0.62	1.38		1.28	1.31	1.13
$\Gamma_{25'v} \rightarrow L_{1c}$	1.45	2.21		2.12	2.13	2.04 ± 0.06
$\Gamma_{25'v} \rightarrow L_{3c}$	3.24	4.18		3.91	3.94	3.9 ± 0.1
MSE	0.35	0.02		0.01	0.03	
MAE	0.44	0.21		0.14	0.17	

¹ M. Shishkin et al. *Phys. Rev. Lett.* 99.24 (2007), 246403 for E_g and M. S. Hybertsen et al. *Phys. Rev. B* 34.8 (1986), 5390 for the transitions;

² M. Shishkin et al. *Phys. Rev. B* 75.23 (2007), 235102.

³ O. Madelung. *Semiconductors*. 3rd ed. Berlin: Springer-Verlag, 2004.


Koopmans functionals give accurate band structures




ZnO	LDA	HSE	GW_0	$scG\tilde{W}$	KI	exp
E_{gap} (eV)	0.79	2.79	3.0	3.2	3.62	3.60
$\langle \varepsilon_d \rangle$ (eV)	-5.1	-6.1	-6.4	-6.7	-6.9	-7.5/-8.0

Koopmans functionals: results for molecules

Ionisation potentials = $E(N - 1) - E(N) \stackrel{?}{=} -\varepsilon_{HO}$ of 100 molecules (the GW100 set) cf. CCSD(T)




Ultraviolet photoemission spectra




Koopmans functionals: results for toy systems

For Hooke's atom (two electrons in a harmonic confining potential with Coulombic repulsion)



Koopmans functionals: results for toy systems

For Hooke's atom (two electrons in a harmonic confining potential with Coulombic repulsion)



The projectability is

$$p_{m\mathbf{k}} = \sum_n |\langle \varphi_n | \psi_{m\mathbf{k}} \rangle|^2$$

We want a set of projectors such that...

- they should have a large overlap with some Kohn-Sham states and a small overlap with all the others
- they should lie within the span of the Kohn-Sham states

We can meet these criteria if we maximise the functional

$$F[\{\varphi_n\}] = \frac{1}{N_{\mathbf{k}} N_w} \sum_{\mathbf{k}} \sum_{m \in S_{\mathbf{k}}} p_{m\mathbf{k}}$$

where $S_{\mathbf{k}}$ corresponds to the N_w -largest values of $p_{m\mathbf{k}}$.

Bayesian optimisation

