

PSI


Center for Scientific Computing,
Theory and Data

Koopmans functionals in practice

minimisation, screening coefficients, automation, and more...

Edward Linscott

ICTP–MARVEL College, 27 May 2026



$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(\underbrace{- \int_0^{f_i} \varepsilon_i(f) \, df}_{\text{removes curvature}} + \underbrace{f_i \eta_i}_{\text{restores linearity}} \right)$$

I. Dabo *et al.* *Phys. Rev. B* **82**, 115121 (2010), G. Borghi *et al.* *Phys. Rev. B* **90**, 75135 (2014), N. Colonna *et al.* *J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(\underbrace{- \int_0^{f_i} \varepsilon_i(f) \, df}_{\text{removes curvature}} + \underbrace{f_i \eta_i}_{\text{restores linearity}} \right)$$

Differences to semi-local functionals:

I. Dabo *et al.* *Phys. Rev. B* **82**, 115121 (2010), G. Borghi *et al.* *Phys. Rev. B* **90**, 75135 (2014), N. Colonna *et al.* *J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(\underbrace{- \int_0^{f_i} \varepsilon_i(f) \, df}_{\text{removes curvature}} + \underbrace{f_i \eta_i}_{\text{restores linearity}} \right)$$

Differences to semi-local functionals:

- screening

I. Dabo *et al.* *Phys. Rev. B* **82**, 115121 (2010), G. Borghi *et al.* *Phys. Rev. B* **90**, 75135 (2014), N. Colonna *et al.* *J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(\underbrace{- \int_0^{f_i} \varepsilon_i(f) \, df}_{\text{removes curvature}} + \underbrace{f_i \eta_i}_{\text{restores linearity}} \right)$$

Differences to semi-local functionals:

- screening
- different flavours

I. Dabo *et al.* *Phys. Rev. B* **82**, 115121 (2010), G. Borghi *et al.* *Phys. Rev. B* **90**, 75135 (2014), N. Colonna *et al.* *J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(\underbrace{- \int_0^{f_i} \varepsilon_i(f) \, df}_{\text{removes curvature}} + \underbrace{f_i \eta_i}_{\text{restores linearity}} \right)$$

Differences to semi-local functionals:

- screening
- different flavours
- orbital-density dependence

I. Dabo *et al.* *Phys. Rev. B* **82**, 115121 (2010), G. Borghi *et al.* *Phys. Rev. B* **90**, 75135 (2014), N. Colonna *et al.* *J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

Screening

Screening

In Hartree-Fock

$$E_{\text{ee}}^{\text{HF}} = \frac{1}{2} \sum_{ij} f_i f_j \int d\mathbf{r} d\mathbf{r}' \frac{|\psi_i(\mathbf{r})|^2 |\psi_j(\mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|} - \frac{\psi_i^*(\mathbf{r}) \psi_j^*(\mathbf{r}') \psi_i(\mathbf{r}') \psi_j(\mathbf{r})}{|\mathbf{r} - \mathbf{r}'|}$$

C. Li *et al.* *J. Chem. Phys.* **146**, 74107 (2017)

T. Koopmans. *Physica* **1**, 104–113 (1934)

In Hartree-Fock


$$E_{\text{ee}}^{\text{HF}} = \frac{1}{2} \sum_{ij} f_i f_j \int d\mathbf{r} d\mathbf{r}' \frac{|\psi_i(\mathbf{r})|^2 |\psi_j(\mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|} - \frac{\psi_i^*(\mathbf{r}) \psi_j^*(\mathbf{r}') \psi_i(\mathbf{r}') \psi_j(\mathbf{r})}{|\mathbf{r} - \mathbf{r}'|} \rightarrow \frac{d^2 E_{\text{ee}}^{\text{HF}}}{df_i^2} \stackrel{?}{=} 0$$

C. Li *et al.* *J. Chem. Phys.* **146**, 74107 (2017)

T. Koopmans. *Physica* **1**, 104–113 (1934)

In Hartree-Fock

$$E_{\text{ee}}^{\text{HF}} = \frac{1}{2} \sum_{ij} f_i f_j \int d\mathbf{r} d\mathbf{r}' \frac{|\psi_i(\mathbf{r})|^2 |\psi_j(\mathbf{r}')|^2}{r - r'} - \frac{\psi_i^*(\mathbf{r}) \psi_j^*(\mathbf{r}') \psi_i(\mathbf{r}') \psi_j(\mathbf{r})}{r - r'} \rightarrow \frac{d^2 E_{\text{ee}}^{\text{HF}}}{df_i^2} \stackrel{?}{=} 0$$



In Hartree-Fock, the first ionisation energy of a system is equal to the negative of the highest occupied orbital's energy $-\epsilon_{\text{HOMO}}^{\text{HF}}$.

— Koopmans' theorem, 1934


$$\text{IP} = E(N - 1) - E(N) = -\epsilon_{\text{HOMO}}^{\text{HF}}$$

C. Li *et al.* *J. Chem. Phys.* **146**, 74107 (2017)

T. Koopmans. *Physica* **1**, 104–113 (1934)

In Hartree-Fock

$$E_{\text{ee}}^{\text{HF}} = \frac{1}{2} \sum_{ij} f_i f_j \int d\mathbf{r} d\mathbf{r}' \frac{|\psi_i(\mathbf{r})|^2 |\psi_j(\mathbf{r}')|^2}{r - r'} - \frac{\psi_i^*(\mathbf{r}) \psi_j^*(\mathbf{r}') \psi_i(\mathbf{r}') \psi_j(\mathbf{r})}{r - r'} \rightarrow \frac{d^2 E_{\text{ee}}^{\text{HF}}}{df_i^2} \stackrel{?}{=} 0$$



In Hartree-Fock, the first ionisation energy of a system is equal to the negative of the highest occupied orbital's energy $-\epsilon_{\text{HOMO}}^{\text{HF}}$.

— Koopmans' theorem, 1934

$$\text{IP} = E(N - 1) - E(N) = -\epsilon_{\text{HOMO}}^{\text{HF}}$$

... exact only if the remaining orbitals do not relax.

C. Li *et al.* *J. Chem. Phys.* **146**, 74107 (2017)

T. Koopmans. *Physica* **1**, 104–113 (1934)


Koopmans functionals


$$E_{\text{KI}}[\{\rho_i\}] = E_{\text{DFT}}[\rho] + \sum_i \left(- \int_0^{f_i} \varepsilon_i(f) \, df + f_i \int_0^1 \varepsilon_i(f) \, df \right)$$


Koopmans functionals

$$\begin{aligned} E_{\text{KI}}[\{\rho_i\}] &= E_{\text{DFT}}[\rho] + \sum_i \left(- \int_0^{f_i} \varepsilon_i(f) \, df + f_i \int_0^1 \varepsilon_i(f) \, df \right) \\ &= E_{\text{DFT}}[\rho] + \sum_i \left(- \left(E_{\text{DFT}}[\rho] - E_{\text{DFT}}[\rho^{f_i \rightarrow 0}] \right) + f_i \left(E_{\text{DFT}}[\rho^{f_i \rightarrow 1}] - E_{\text{DFT}}[\rho^{f_i \rightarrow 0}] \right) \right) \end{aligned}$$


cannot be evaluated directly total energy differences









Screening: how to calculate the ideal α ?




Screening: how to calculate the ideal α ?



Screening: how to calculate the ideal α ?




Screening: how to calculate the ideal α ?




$$\lambda_{ii}(\alpha) \equiv \langle \varphi_i | \hat{h}^{\text{DFT}} + \alpha \hat{v}^{\text{Koopmans}} | \varphi_i \rangle = \left. \frac{dE^{\text{Koopmans}}}{df_i} \right|_{f_i=s}$$

Screening: how to calculate the ideal α ?




$$\lambda_{ii}(\alpha) \equiv \left\langle \varphi_i \left| \hat{h}^{\text{DFT}} + \alpha \hat{v}^{\text{Koopmans}} \right| \varphi_i \right\rangle = \left. \frac{dE^{\text{Koopmans}}}{df_i} \right|_{f_i=s}$$

Screening: how to calculate the ideal α ?




$$\lambda_{ii}(\alpha) \equiv \left\langle \varphi_i \left| \hat{h}^{\text{DFT}} + \alpha \hat{v}^{\text{Koopmans}} \right| \varphi_i \right\rangle = \left. \frac{dE^{\text{Koopmans}}}{df_i} \right|_{f_i=s}$$

Screening: how to calculate the ideal α ?



Given this, how to work out the ideal α ?

Screening: how to calculate the ideal α ?



Given this, how to work out the ideal α ?

$$\alpha_{n+1} = \alpha_n \frac{E_i(N-1) - E(N) - \lambda_{ii}(0)}{\lambda_{ii}(\alpha_n) - \lambda_{ii}(0)}$$

$$\alpha^{n+1} = \alpha^n \frac{E_i(N-1) - E(N) - \lambda_{ii}(0)}{\lambda_{ii}(\alpha^n) - \lambda_{ii}(0)}$$

trial screening
parameter

total energy with electron
removed from orbital i

$$\alpha^{n+1} = \alpha^n \frac{E_i(N-1) - E(N) - \lambda_{ii}(0)}{\lambda_{ii}(\alpha^n) - \lambda_{ii}(0)}$$

trial screening
parameter

$$\alpha^{n+1} = \alpha^n \frac{E_i(N-1) - E(N) - \lambda_{ii}(0)}{\lambda_{ii}(\alpha^n) - \lambda_{ii}(0)}$$

Diagram illustrating the iterative screening coefficient α^{n+1} calculation:

- $E_i(N-1)$: total energy with electron removed from orbital i
- $E(N)$: total energy of neutral system
- $\lambda_{ii}(0)$: trial screening parameter

$$\alpha^{n+1} = \alpha^n \frac{E_i(N-1) - E(N) - \lambda_{ii}(0)}{\lambda_{ii}(\alpha^n) - \lambda_{ii}(0)}$$

Diagram illustrating the iterative screening coefficient α^{n+1} calculation:

- $E_i(N-1)$: total energy with electron removed from orbital i
- $E(N)$: total energy of neutral system
- $\lambda_{ii}(0)$: expectation value of $\hat{H}^{\text{Koopmans}}$ at $\alpha=0$
- $\lambda_{ii}(\alpha^n)$: expectation value of $\hat{H}^{\text{Koopmans}}$ at α^n
- α^n : trial screening parameter

$$\alpha^{n+1} = \alpha^n \frac{E_i(N-1) - E(N) - \lambda_{ii}(0)}{\lambda_{ii}(\alpha^n) - \lambda_{ii}(0)}$$

Diagram illustrating the screening coefficient formula with annotations:

- $E_i(N-1)$: total energy with electron removed from orbital i
- $E(N)$: total energy of neutral system
- $\lambda_{ii}(0)$: expectation value of $\hat{H}^{\text{Koopmans}}$
- $\lambda_{ii}(\alpha^n)$: expectation value of $\hat{H}^{\text{DFT}}$
- α^n : trial screening parameter

Screening via DFPT

How can we avoid explicit charged defect calculations in a supercell?

N. Colonna *et al. J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

N. Colonna *et al. J. Chem. Theory Comput.* **18**, 5435–5448 (2022)

Screening via DFPT

How can we avoid explicit charged defect calculations in a supercell?

Reformulate in terms of DFPT...

$$\alpha_i = 1 + \frac{\langle v_{\text{pert}}^i | \Delta^i n \rangle}{\langle n_i | v_{\text{pert}}^i \rangle}$$

N. Colonna *et al.* *J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

N. Colonna *et al.* *J. Chem. Theory Comput.* **18**, 5435–5448 (2022)

Screening via DFPT

How can we avoid explicit charged defect calculations in a supercell?

Reformulate in terms of DFPT...

$$\alpha_i = 1 + \frac{\langle v_{\text{pert}}^i \mid \Delta^i n \rangle}{\langle n_i \mid v_{\text{pert}}^i \rangle}$$


... in reciprocal space

$$\alpha_{0i} = 1 + \frac{\sum_{\mathbf{q}} \langle v_{\text{pert},\mathbf{q}}^{0i} \mid \Delta_{\mathbf{q}}^{0i} n \rangle}{\sum_{\mathbf{q}} \langle n_{\mathbf{q}}^{0i} \mid v_{\text{pert},\mathbf{q}}^{0i} \rangle}$$

N. Colonna *et al.* *J. Chem. Theory Comput.* **15**, 1905–1914 (2019)

N. Colonna *et al.* *J. Chem. Theory Comput.* **18**, 5435–5448 (2022)

Screening via DFPT



Orbital-density-dependence

Orbital-density-dependence



$$\begin{aligned} & - \int_0^{f_i} \varepsilon_i(f) \, df + f_i \int_0^1 \varepsilon_i(f) \, df \\ & = E_{\text{Hxc}}[\rho] + E_{\text{Hxc}}[\rho - \rho_i] + f_i(-E_{\text{Hxc}}[\rho - \rho_i] + E_{\text{Hxc}}[\rho - \rho_i + n_i]) \end{aligned}$$

Orbital-density-dependence



$$\begin{aligned} & - \int_0^{f_i} \varepsilon_i(f) \, df + f_i \int_0^1 \varepsilon_i(f) \, df \\ & = E_{\text{Hxc}}[\rho] + E_{\text{Hxc}}[\rho - \rho_i] + f_i(-E_{\text{Hxc}}[\rho - \rho_i] + E_{\text{Hxc}}[\rho - \rho_i + n_i]) \end{aligned}$$

For filled orbitals with KI:

$$v_i^{\text{KI}}/\alpha_i = -E_{\text{H}}[n_i] + E_{\text{xc}}[\rho] - E_{\text{xc}}[\rho - n_i] - \int d\mathbf{r}' v_{\text{xc}}(\mathbf{r}', [\rho]) n_i(\mathbf{r}')$$

Orbital-density-dependence

Minimise $E[\rho, \rho_i]$ with CG minimisation:

G. Borghi *et al.* *Phys. Rev. B* **91**, 155112 (2015)

Orbital-density-dependence

Minimise $E[\rho, \rho_i]$ with CG minimisation:

outer loop $\varphi_i^{(n+1)} = \varphi_i^{(n)} + \Delta_i$

Orbital-density-dependence

Minimise $E[\rho, \rho_i]$ with CG minimisation:

outer loop $\varphi_i^{(n+1)} = \varphi_i^{(n)} + \Delta_i$
inner loop $\varphi_i^{(n+1)} = \sum_j U_{ij} \varphi_j^{(n)}$

Orbital-density-dependence

Minimise $E[\rho, \rho_i]$ with CG minimisation:

outer loop $\varphi_i^{(n+1)} = \varphi_i^{(n)} + \Delta_i$

inner loop $\varphi_i^{(n+1)} = \sum_j U_{ij} \varphi_j^{(n)}$

For more details see Borghi *et al.* (2015)

Orbital-density-dependence

Minimise $E[\rho, \rho_i]$ with CG minimisation:

outer loop $\varphi_i^{(n+1)} = \varphi_i^{(n)} + \Delta_i$

inner loop $\varphi_i^{(n+1)} = \sum_j U_{ij} \varphi_j^{(n)}$

For more details see Borghi *et al.* (2015)

- gives rise to a set of **minimising orbitals** (localised/variational)

Orbital-density-dependence



Minimise $E[\rho, \rho_i]$ with CG minimisation:


outer loop $\varphi_i^{(n+1)} = \varphi_i^{(n)} + \Delta_i$

inner loop $\varphi_i^{(n+1)} = \sum_j U_{ij} \varphi_j^{(n)}$


For more details see Borghi *et al.* (2015)

- gives rise to a set of **minimising orbitals** (localised/variational)
- diagonalising at the minimum gives rise to **diagonalising orbitals** (delocalised/canonical)

Orbital-density-dependence




variational



canonical


N. L. Nguyen *et al.* *Phys. Rev. X* **8**, 21051 (2018)

Importance of localisation




N. L. Nguyen *et al.* *Phys. Rev. X* **8**, 21051 (2018)

Importance of localisation



In the bulk limit for one cell $\Delta E = E(N - \delta N) - E(N)$


Importance of localisation



In the bulk limit for one cell $\Delta E = E(N - \delta N) - E(N)$


Across all the cells $\Delta E = \frac{1}{\delta N} (E(N - \delta N) - E(N)) = -\frac{dE}{dN} = -\varepsilon_{\text{HO}}$

Importance of localisation



$$\lim_{n_i(r) \rightarrow 0} v_i^{\text{KI}} / \alpha_i = \lim_{n_i(r) \rightarrow 0} -E_{\text{H}}[n_i] + E_{\text{xc}}[\rho] - E_{\text{xc}}[\rho - n_i] - \int d\mathbf{r}' v_{\text{xc}}(\mathbf{r}', [\rho]) n_i(\mathbf{r}') = 0$$

Importance of localisation



$$\lim_{n_i(r) \rightarrow 0} v_i^{\text{KI}} / \alpha_i = \lim_{n_i(r) \rightarrow 0} -E_{\text{H}}[n_i] + E_{\text{xc}}[\rho] - E_{\text{xc}}[\rho - n_i] - \int d\mathbf{r}' v_{\text{xc}}(\mathbf{r}', [\rho]) n_i(\mathbf{r}') = 0$$

Solution: apply Koopmans correction to localised orbitals

Orbital-density-dependence

Other features of orbital-density-dependence

A. Ferretti *et al.* *Phys. Rev. B* **89**, 195134 (2014)

Orbital-density-dependence



Other features of orbital-density-dependence

- ODD functional means that we know $\hat{H}|\varphi_i\rangle$ for variational orbitals $\{|\varphi_i\rangle\}$ but we don't know $\hat{H}$ in general

Orbital-density-dependence



Other features of orbital-density-dependence

- ODD functional means that we know $\hat{H}|\varphi_i\rangle$ for variational orbitals $\{|\varphi_i\rangle\}$ but we don't know $\hat{H}$ in general
- Practically we can often use MLWFs

Orbital-density-dependence



Other features of orbital-density-dependence

- ODD functional means that we know $\hat{H}|\varphi_i\rangle$ for variational orbitals $\{|\varphi_i\rangle\}$ but we don't know $\hat{H}$ in general
- Practically we can often use MLWFs
- a natural generalisation in the direction of spectral functional theory

Flavours

Flavours of Koopmans functionals

$$E_{\text{Koopmans}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) \, df + f_i \eta_i \right)$$

One degree of freedom: what should be the gradient of this linear term?

$$E_{\text{KI}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) \, df + f_i \int_0^1 \varepsilon_i(f) \, df \right)$$

One degree of freedom: what should be the gradient of this linear term?

- the base functional $\rightarrow$ “KI” (Koopmans integral). Enforces IP theorem. Does not affect energy/density!

You might also see...

$$E_{\text{KIPZ}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) \, df + f_i \left\{ \int_0^1 \varepsilon_i(f) \, df - E_{\text{Hxc}}[n_i] \right\} \right)$$

One degree of freedom: what should be the gradient of this linear term?

- the base functional $\rightarrow$ “KI” (Koopmans integral). Enforces IP theorem. Does not affect energy/density!
- with a PZ correction $\rightarrow$ “KIPZ”

You might also see...

- “pKIPZ” = KIPZ Hamiltonian evaluated on the KI solution

Flavours of Koopmans functionals

$$E_{\text{KIPZ}}[\rho, \{f_i\}, \{\alpha_i\}] = E_{\text{DFT}}[\rho] + \sum_i \alpha_i \left(- \int_0^{f_i} \varepsilon_i(f) \, df + f_i \left\{ \int_0^1 \varepsilon_i(f) \, df - E_{\text{Hxc}}[n_i] \right\} \right)$$

One degree of freedom: what should be the gradient of this linear term?

- the base functional $\rightarrow$ “KI” (Koopmans integral). Enforces IP theorem. Does not affect energy/density!
- with a PZ correction $\rightarrow$ “KIPZ”

You might also see...

- “pKIPZ” = KIPZ Hamiltonian evaluated on the KI solution
- “K” = an earlier iteration based off half-filling rather than integer endpoints (no longer used)

To summarise...

To summarise...

When performing a Koopmans calculation, you must decide...

To summarise...

When performing a Koopmans calculation, you must decide...

- which flavour (KI, KIPZ)?

To summarise...

When performing a Koopmans calculation, you must decide...

- which flavour (KI, KIPZ)?
- how are we treating the orbital-density-dependence?
 - how are we initialising our variational orbitals? (N.B. depends on the flavour!)
 - are we going to explicitly minimise the ODD?


To summarise...

When performing a Koopmans calculation, you must decide...


- which flavour (KI, KIPZ)?
- how are we treating the orbital-density-dependence?
 - how are we initialising our variational orbitals? (N.B. depends on the flavour!)
 - are we going to explicitly minimise the ODD?
- how are we calculating the screening parameters? (finite differences, DFPT)

Results

Ionisation potentials of 100 molecules cf. CCSD(T)




Ultraviolet photoemission spectra



N. Colonna *et al.* *J. Chem. Theory Comput.* **14**, 2549–2557 (2018), N. L. Nguyen *et al.* *Phys. Rev. Lett.* **114**, 166405 (2015)

Koopmans functionals: results for solids




Mean absolute error (eV) across prototypical semiconductors and insulators

	PBE	G_0W_0	KI	KIPZ	QSGW $\tilde{W}$
E_{gap}	2.54	0.56	0.27	0.22	0.18
IP	1.09	0.39	0.19	0.21	0.49

Toy systems



For Hooke's atom (two electrons in a harmonic confining potential with Coulombic repulsion)



Y. Schubert *et al.* *J. Chem. Phys.* **158**, 144113 (2023)

Toy systems

For Hooke's atom (two electrons in a harmonic confining potential with Coulombic repulsion)



Koopmans functionals: caveats

N. Q. Su *et al.* *J. Phys. Chem. Lett.* **11**, 1528–1535 (2020)

Koopmans functionals: caveats

- restricted to systems with a non-zero band gap

N. Q. Su *et al.* *J. Phys. Chem. Lett.* **11**, 1528–1535 (2020)

Koopmans functionals: caveats

- restricted to systems with a non-zero band gap
- empty state localisation in the bulk limit

Koopmans functionals: caveats




- restricted to systems with a non-zero band gap
- empty state localisation in the bulk limit
- can potentially break the crystal point group symmetry

N. Q. Su *et al.* *J. Phys. Chem. Lett.* **11**, 1528–1535 (2020)

Computational cost and scaling

Computational cost and scaling



Computational cost and scaling

The vast majority of the computational cost: determining screening parameters

$$\alpha_i = \frac{\langle n_i | \varepsilon^{-1} f_{\text{Hxc}} | n_i \rangle}{\langle n_i | f_{\text{Hxc}} | n_i \rangle}$$

N. L. Nguyen *et al.* *Phys. Rev. X* **8**, 21051 (2018), R. De Gennaro *et al.* *Phys. Rev. B* **106**, 35106 (2022)

N. Colonna *et al.* *J. Chem. Theory Comput.* **18**, 5435–5448 (2022), N. Colonna *et al.* *J. Chem. Theory Comput.* **14**, 2549–2557 (2018)

Computational cost and scaling

The vast majority of the computational cost: determining screening parameters

$$\alpha_i = \frac{\langle n_i | \varepsilon^{-1} f_{\text{Hxc}} | n_i \rangle}{\langle n_i | f_{\text{Hxc}} | n_i \rangle}$$

- one screening parameter per orbital

N. L. Nguyen *et al.* *Phys. Rev. X* **8**, 21051 (2018), R. De Gennaro *et al.* *Phys. Rev. B* **106**, 35106 (2022)

N. Colonna *et al.* *J. Chem. Theory Comput.* **18**, 5435–5448 (2022), N. Colonna *et al.* *J. Chem. Theory Comput.* **14**, 2549–2557 (2018)

Computational cost and scaling

The vast majority of the computational cost: determining screening parameters

$$\alpha_i = \frac{\langle n_i | \varepsilon^{-1} f_{\text{Hxc}} | n_i \rangle}{\langle n_i | f_{\text{Hxc}} | n_i \rangle}$$

- one screening parameter per orbital
- must be computed *ab initio* via **embarrassingly parallel** steps...

N. L. Nguyen *et al.* *Phys. Rev. X* **8**, 21051 (2018), R. De Gennaro *et al.* *Phys. Rev. B* **106**, 35106 (2022)

N. Colonna *et al.* *J. Chem. Theory Comput.* **18**, 5435–5448 (2022), N. Colonna *et al.* *J. Chem. Theory Comput.* **14**, 2549–2557 (2018)

Computational cost and scaling

The vast majority of the computational cost: determining screening parameters

$$\alpha_i = \frac{\langle n_i | \varepsilon^{-1} f_{\text{Hxc}} | n_i \rangle}{\langle n_i | f_{\text{Hxc}} | n_i \rangle}$$

- one screening parameter per orbital
- must be computed *ab initio* via **embarrassingly parallel** steps...
 - ΔSCF : $\mathcal{O}(N_{\text{SC}}^3) \sim \mathcal{O}(N_{\mathbf{k}}^3 N^3)$

N. L. Nguyen *et al.* *Phys. Rev. X* **8**, 21051 (2018), R. De Gennaro *et al.* *Phys. Rev. B* **106**, 35106 (2022)

N. Colonna *et al.* *J. Chem. Theory Comput.* **18**, 5435–5448 (2022), N. Colonna *et al.* *J. Chem. Theory Comput.* **14**, 2549–2557 (2018)

Computational cost and scaling

The vast majority of the computational cost: determining screening parameters


$$\alpha_i = \frac{\langle n_i | \varepsilon^{-1} f_{\text{Hxc}} | n_i \rangle}{\langle n_i | f_{\text{Hxc}} | n_i \rangle}$$

- one screening parameter per orbital
- must be computed *ab initio* via **embarrassingly parallel** steps...
 - ΔSCF : $\mathcal{O}(N_{\text{SC}}^3) \sim \mathcal{O}(N_{\mathbf{k}}^3 N^3)$
 - DFPT: $\mathcal{O}(N_{\mathbf{k}}^2 N^3)$

N. L. Nguyen *et al.* *Phys. Rev. X* **8**, 21051 (2018), R. De Gennaro *et al.* *Phys. Rev. B* **106**, 35106 (2022)

N. Colonna *et al.* *J. Chem. Theory Comput.* **18**, 5435–5448 (2022), N. Colonna *et al.* *J. Chem. Theory Comput.* **14**, 2549–2557 (2018)

Machine-learned electronic screening



power spectrum decomposition

$$\begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ \vdots \end{bmatrix}$$


ridge regression

$$\alpha_i$$


$$c_{nlm,k}^i = \int d\mathbf{r} g_{nl}(r) Y_{lm}(\theta, \varphi) n^i(\mathbf{r} - \mathbf{R}^i)$$

$$p_{n_1 n_2 l, k_1 k_2}^i = \pi \sqrt{\frac{8}{2l+1}} \sum_m c_{n_1 l m, k_1}^{i*} c_{n_2 l m, k_2}^i$$

Machine-learned electronic screening










water



CsSnI_3

Machine-learned electronic screening

The use-case




...

train

predict

Machine-learned electronic screening

The use-case





train


predict

N.B. not a general model


Machine-learned electronic screening



Machine-learned electronic screening



accurate to within $\mathcal{O}(10 \text{ meV})$ cf. typical band gap accuracy of $\mathcal{O}(100 \text{ meV})$



speedup of $\mathcal{O}(10)$ to $\mathcal{O}(100)$

Running a Koopmans functional calculation


The workflows

The general workflow:

- define/initialise a set of variational orbitals
- calculate the screening parameters $\{\alpha_i\}$
- construct and diagonalise the Hamiltonian

The workflows


(a) finite difference calculations using a supercell



The workflows


The workflows

(a) finite difference calculations using a supercell



(b) DFPT in a primitive cell

The workflows




koopmans

The koopmans package

- v1.0 released in 2023 (now v1.2)
- implementations of Koopmans functionals
- automated workflows
 - start-to-finish Koopmans calculations
 - Wannierisation
 - dielectric tensor
 - ...
- built on top of ASE
- under the hood, calls Quantum ESPRESSO
- does not require expert knowledge

koopmans-functionals.org



E. B. Linscott *et al.* *J. Chem. Theory Comput.* **19**, 7097–7111 (2023)

A. H. Larsen *et al.* *J. Phys. Condens. Matter* **29**, 273002 (2017)

koopmans: the input file

```
{
  "workflow": {
    "task": "singlepoint",
    "functional": "ki",
    "method": "dscf",
    "init_orbitals": "mlwfs",
    "alpha_guess": 0.1
  },
  "atoms": {
    "atomic_positions": {
      "units": "crystal",
      "positions": [
        ["Si", 0.00, 0.00, 0.00],
        ["Si", 0.25, 0.25, 0.25]
      ]
    },
    "cell_parameters": {
      "periodic": true,
      "ibrav": 2,
      "celldm(1)": 10.262
    }
  },
}
```

```
  "k_points": {
    "grid": [8, 8, 8],
    "path": "LGXKG"
  },
  "calculator_parameters": {
    "ecutwfc": 60.0,
    "w90": {
      "projections": [
        [{"fsite": [0.125, 0.125, 0.125],
          "ang_mtm": "sp3"}],
        [{"fsite": [0.125, 0.125, 0.125],
          "ang_mtm": "sp3"}]
      ],
      "dis_froz_max": 11.5,
      "dis_win_max": 17.0
    }
  }
}
```


 **koopmans v2 is coming...** 

Koopmans AiiDA




- remote compute
- parallel step execution
- provenance-tracking
- error recovery
- and more!

Koopmans AiiDA

- remote compute
- parallel step execution
- provenance-tracking
- error recovery
- and more!



Take home messages



- Koopmans functionals are more complicated than a simple semi-local DFT calculation, because of...
 - orbital-density-dependence
 - screening parameters
- Koopmans functionals are implemented in Quantum ESPRESSO
- the complexity of the workflows are handled by the koopmans package
- keep your eyes peeled for v2!

koopmans: An Open-Source Package for Accurately and Efficiently Predicting Spectral Properties with Koopmans Functionals

Edward B. Linscott,^{*,△} Nicola Colonna,[△] Riccardo De Gennaro, Ngoc Linh Nguyen, Giovanni Borghi, Andrea Ferretti, Ismaila Dabo, and Nicola Marzari^{*}



Cite This: <https://doi.org/10.1021/acs.jctc.3c00652>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: Over the past decade we have developed Koopmans functionals, a computationally efficient approach for predicting spectral properties with an orbital-density-dependent functional framework. These functionals impose a generalized piecewise linearity condition to the entire electronic manifold, ensuring that orbital energies match the corresponding electron removal/addition energy differences (in contrast to semilocal DFT, where a mismatch between the two lies at the heart of the band gap problem and, more generally, the unreliability of Kohn–Sham orbital energies). This strategy has proven to be very powerful.



koopmans-functionals.org

Thank you!


Acknowledgements



Nicola
Colonna



Miki
Bonacci




Aleksandr
Poliukhin



Marija
Stojkovic



Giovanni
Cistaro



Junfeng
Qiao



Yannick
Schubert



Nicola
Marzari



Want to find out more? Go to koopmans-functionals.org

slides available at  [github/elinscott-talks](https://github.com/elinscott/talks)

- I. Dabo *et al.* Koopmans' Condition for Density-Functional Theory. *Phys. Rev. B* **82**, 115121 (2010).
- G. Borghi, A. Ferretti, N. L. Nguyen, I. Dabo & N. Marzari. Koopmans-Compliant Functionals and Their Performance against Reference Molecular Data. *Phys. Rev. B* **90**, 75135 (2014).
- N. Colonna, N. L. Nguyen, A. Ferretti & N. Marzari. Koopmans-Compliant Functionals and Potentials and Their Application to the GW100 Test Set. *J. Chem. Theory Comput.* **15**, 1905–1914 (2019).
- C. Li & W. Yang. On the Piecewise Convex or Concave Nature of Ground State Energy as a Function of Fractional Number of Electrons for Approximate Density Functionals. *J. Chem. Phys.* **146**, 74107 (2017).
- T. Koopmans. Über die Zuordnung von Wellenfunktionen und Eigenwerten zu den einzelnen Elektronen eines Atoms. *Physica* **1**, 104–113 (1934).
- N. Colonna, R. De Gennaro, E. Linscott & N. Marzari. Koopmans Spectral Functionals in Periodic Boundary Conditions. *J. Chem. Theory Comput.* **18**, 5435–5448 (2022).
- G. Borghi, C. H. Park, N. L. Nguyen, A. Ferretti & N. Marzari. Variational Minimization of Orbital-Density-Dependent Functionals. *Phys. Rev. B* **91**, 155112 (2015).
- N. L. Nguyen, N. Colonna, A. Ferretti & N. Marzari. Koopmans-Compliant Spectral Functionals for Extended Systems. *Phys. Rev. X* **8**, 21051 (2018).
- A. Ferretti, I. Dabo, M. Cococcioni & N. Marzari. Bridging Density-Functional and Many-Body Perturbation Theory: Orbital-Density Dependence in Electronic-Structure Functionals. *Phys. Rev. B* **89**, 195134 (2014).
- N. Colonna, N. L. Nguyen, A. Ferretti & N. Marzari. Screening in Orbital-Density-Dependent Functionals. *J. Chem. Theory Comput.* **14**, 2549–2557 (2018).
- N. L. Nguyen, G. Borghi, A. Ferretti, I. Dabo & N. Marzari. First-Principles Photoemission Spectroscopy and Orbital Tomography in Molecules from Koopmans-Compliant Functionals. *Phys. Rev. Lett.* **114**, 166405 (2015).
- Y. Schubert, N. Marzari & E. Linscott. Testing Koopmans Spectral Functionals on the Analytically Solvable Hooke's Atom. *J. Chem. Phys.* **158**, 144113 (2023).
- N. Q. Su, A. Mahler & W. Yang. Preserving Symmetry and Degeneracy in the Localized Orbital Scaling Correction Approach. *J. Phys. Chem. Lett.* **11**, 1528–1535 (2020).

- R. De Gennaro, N. Colonna, E. Linscott & N. Marzari. Bloch's theorem in orbital-density-dependent functionals: Band structures from Koopmans spectral functionals. *Phys. Rev. B* **106**, 35106 (2022).
- Y. Schubert, S. Luber, N. Marzari & E. Linscott. Predicting electronic screening for fast Koopmans spectral functional calculations. *npj Comput Mater* **10**, 1–12 (2024).
- E. B. Linscott *et al.* koopmans: an open-source package for accurately and efficiently predicting spectral properties with Koopmans Functionals. *J. Chem. Theory Comput.* **19**, 7097–7111 (2023).
- A. H. Larsen *et al.* The Atomic Simulation Environment—a Python Library for Working with Atoms. *J. Phys. Condens. Matter* **29**, 273002 (2017).
- S. P. Huber *et al.* AiiDA 1.0, a scalable computational infrastructure for automated reproducible workflows and data provenance. *Sci Data* **7**, 300 (2020).
- J. Qiao, G. Pizzi & N. Marzari. Projectability disentanglement for accurate and automated electronic-structure Hamiltonians. *npj Comput Mater* **9**, 208 (2023).
- J. Qiao, G. Pizzi & N. Marzari. Automated mixing of maximally localized Wannier functions into target manifolds. *npj Comput Mater* **9**, 206 (2023).

Spare slides

Recap from earlier

Key idea: construct a functional such that the *variational* orbital energies

$$\varepsilon_i^{\text{Koopmans}} = \langle \varphi_i | H | \varphi_i \rangle = \partial E_{\text{Koopmans}} / \partial f_i$$

are...

- independent of the corresponding occupancies f_i
- equal to the corresponding total energy difference $E_i(N-1) - E(N)$

zero band gap $\rightarrow$ occupancy matrix for variational orbitals is off-diagonal

Automated Wannierisation

Koopmans functionals rely heavily on Wannier functions...

- to initialise the minimising orbitals, *or*
- in place of the minimising orbitals entirely


J. Qiao *et al.* *npj Comput Mater* **9**, 208 (2023)

J. Qiao *et al.* *npj Comput Mater* **9**, 206 (2023)


Automated Wannierisation

Koopmans functionals rely heavily on Wannier functions...


- to initialise the minimising orbitals, *or*
- in place of the minimising orbitals entirely



projectability-based
disentanglement



use PAOs found in
pseudopotentials



parallel transport to separate manifolds

J. Qiao *et al.* *npj Comput Mater* **9**, 208 (2023)

J. Qiao *et al.* *npj Comput Mater* **9**, 206 (2023)