



# Frost Impressions of ACCESS-S2 Antarctic Sea Ice Forecast

Elio Campitelli<sup>1, 2</sup>,

- (1) School of Earth, Atmosphere and Environment, Monash University, Australia.
- (2) Securing Antarctica's Environmental Future, Monash University, Australia.
- (3) Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales. Buenos Aires, Argentina.
- (4) CONICET – Universidad de Buenos Aires. Centro de Investigaciones del Mar y la Atmósfera (CIMA). Buenos Aires, Argentina.
- (5) CNRS – IRD – CONICET – UBA. Instituto Franco-Argentino para el Estudio del Clima y sus Impactos (IFAECI). Buenos Aires, Argentina.




## ACCESS-S2

The Australian Bureau of Meteorology's current seasonal prediction system. Replaced ACCESS-S1 in October 2021. Same model configuration:

- Global Atmosphere 6.0 (GA6) N216 (~60km) 85 levels.
- Global Land 6.0 N216 4 levels.
- Global Ocean 5.0 1/4° 75 levels.
- Global Sea Ice 6.0 (CICE 4.1) 1/4° 5 ice categories.

Initial conditions (hindcast):  
S1: ERA-interim + Met Office FOAM (includes sea ice)  
S2: ERA-interim + BOM analysis (doesn't include sea ice)

## Results




**Figure 1:** Median sea ice extent for al hindcasts initialised the first of the month for ACCESS-S2 and ACCESS-S1 in colours representing the start month. In black, the median sea ice extent of NSIDC CDR.

## Data


ACCESS-S2 and S1 hindcasts (1981–2014 and 1990–2012, respectively). Initialised at the 1st of every month.  
Observations: NSIDC CDRv4 sea ice concentrations.

## Metrics

**Sea Ice Extent:** Area covered with at least 15% sea ice.  
**RMSE:** Root Mean Squared Error of sea ice concentration anomalies.  
**IIIE** (Integrated Ice Edge Error): Area in which model and observations disagree on the presence or absence of sea ice (concentration >15%).



**Figure 1:** ACCESS-S2 sea ice concentration bias with NSIDC sea ice concentration.



**Figure 1:** Median and 95% coverage of sea ice concentration anomalies RMSE as a function of forecast lag for all forecast initialised on the first of each month compared with a reference forecast of persistence of anomalies.