

Are planetary waves the same thing as quasi-stationary waves?

Elio Campitelli ¹Carolina Vera ^{1,2}Leandro Díaz ^{1,2}

¹Centro de Investigaciones del Mar y la Atmosfera, UMI-IFAECI (CONICET-UBA-CNRS)

²Departamento de Ciencias de la Atmósfera y los Océanos (FCEyN, UBA)

Key Points:

- Zonal waves and Quasi-stationary waves are distinct but related phenomena
- This distinction has theoretical and practical implications
- The relationship between the mean ZW amplitude and QS amplitude yields an estimate of stationarity

Corresponding author: Elio Campitelli, elio.campitelli@cima.fcen.uba.ar

Abstract

In the meteorological literature the analysis of the zonally asymmetric it is very common to analyse

1 Introduction

Many atmospheric variables have a strong dependence with latitude, so it is often natural to decompose them into a zonal mean component and a deviation from it. If ϕ is a generic variable, then

$$\phi(x, y, z, t) = [\phi]_{(y, z, t)} + \phi^*_{(x, y, z, t)} \quad (1)$$

where $[\phi]$ is the mean zonal field and ϕ^* , the deviations from it. This zonally asymmetric part is sometimes called “zonal wave” or “planetary wave”. The names “stationary wave” or “quasi-stationary wave”, on the other hand, are generally reserved to the zonal asymmetries of the time mean field ($\overline{\phi^*}$). However, these terms are sometimes used interchangeably in the literature (e.g. Irving & Simmonds, 2015; Kravchenko et al., 2012; Lastovicka, Krizan, & Kozubek, 2018; Rao, Fernandez, & Franchito, 2004; Raphael, 2004; Turner, Hosking, Bracegirdle, Phillips, & Marshall, 2017) which could lead to some confusion.

Given a set of atmospheric fields, we define *zonal waves* (ZW) as waves observed in each individual “instantaneous” field and *quasi-stationary waves* (QS) as the resulting waves in the mean field. While these definitions depend on which are the “instantaneous field” in question (monthly, daily, subdaily, etc. . .) and the averaging timescales used, they illustrate that ZWs are properties of the *elements* of the set, while the QSS are properties of the set as a whole. This is an important distinction with theoretical and methodological implications that is not always appreciated in the literature.

2 Story

To illustrate the distinction between ZWs and QS, Figure 1 shows the monthly seasonal cycle of amplitude of planetary waves at 60°S using monthly fields from the NCEP/NCAR reanalysis (Kalnay et al., 1996) between 1950 and 1998. The left column (\overline{ZW}) reproduces Figure 3 from Rao et al. (2004) and is computed by taking –for each month and level– the average amplitude of the 49 individual amplitudes. The right column (QS),





Figure 1. Seasonal cycle of amplitude of the geopotential planetary waves 1 to 3 at 60°S computed as the mean amplitude of the monthly waves (\overline{ZW}) and as the amplitude of the mean wave (QS). The period of analysis is 1950 to 1998. The left column reproduces Figure 3 from ©Rao2004.

on the other hand, is computed by taking the amplitude of the average geopotential field for each month and level.

The resulting fields convey different information. First, the amplitude of \overline{ZW} fields is always greater than the one for QS fields. This is a mathematical necessity (*xxDeberia demostrar eso? Vale la pena una demostracion en un material suplementario?xx*) that explains Rao et al. (2004)’s observation that their Wave 1 amplitude was greater than that reported by Hurrell, van Loon, and Shea (1998). Secondly, they have different annual cycles and vertical structures. QS2 has a strong minimum in the low stratosphere during the austral autumn that is not apparent in $\overline{ZW2}$. Similarly, the austral winter mid-tropospheric maximum is very well defined in $\overline{ZW3}$ but not so in QS3. Thirdly, the relative importance between each wave number vary. \overline{ZW} fields show a preponderance of wave 2 over 3 in almost every level and month. However, the QS3 has greater ampli-

56 tude than QS2 in the first half of the year. In contrast with wave-numbers 2 and 3, $\overline{ZW1}$
 57 and QS1 fields are very similar.



58 **Figure 2.** Seasonal cycle of amplitude of the geopotential planetary waves 2 at 300hPa com-
 59 puted as the mean amplitude of the monthly waves (\overline{ZW}) and as the amplitude of the mean
 60 wave (QS). From monthly NCEP/NCAR Reanalysis, 1979 to 2017.

61 These differences are related to the degree of stationarity of zonal waves and are
 62 location-dependent. Figure 2 show the same variable that Figure 1 but for 300hPa. The
 63 contrast between the northern and southern hemisphere is not only evident in the am-
 64 plitude of the planetary waves, but also in the comparison between \overline{ZW} and QS. Spe-
 65 cially for wave-numbers 2 and 3, \overline{ZW} and QS fields are very similar in the north but they
 66 have significant differences in the south.

67 2.1 Stationarity

68 Another important consequence of the distinction between \overline{ZW} and QS is that the
 69 quotient between the two can be used as a measure of stationarity. As an analogy with

the constancy of the wind (Singer, 1967), planetary wave stationarity can be estimated as

$$\hat{S} = \frac{QS}{\overline{ZW}} \quad (2)$$

It can be shown that $\hat{S} = 1$ for completely stationary waves and that $E(\hat{S}) = n^{-1/2}$ for completely non-stationary waves (where n is the sample size).




Figure 3. Seasonal cycle of stationarity of the 300hPa geopotential QS2 computed using Equation 2 (shaded) and $\overline{ZW2}$ (contours). From monthly NCEP/NCAR Reanalysis, 1979 to 2017.

As an example, Figure 3 shows \hat{S} for QS2 computed using Equation 2. The southern hemisphere clearly shows a lower degree of QS2 stationarity than the northern hemisphere or the tropics. In the northern mid latitudes there is a seasonal cycle of stationarity that follows the seasonal cycle of \overline{ZW} (Figure 2). In the southern hemisphere, instead, the June maximum of \overline{ZW} at 60°S coincides with a minimum of stationarity.

\hat{S} can equivalently be defined as the amplitude weighted mean correlation between each wave phase and the stationary phase (*xx de nuevo, esto podría demostrarse en un material suplementario xx*). This definition allows one to construct a time series of $\hat{S}(t)$ by computing a running (weigthed) mean.

While \hat{S} is used –sometimes as $2/\pi \arcsin(\hat{S})$ transformation (Singer, 1967)– in the meteorological literature in the context of wind steadiness, to our knowledge this is the first time it has been applied to the study of atmospheric waves. Furthermore, its statistical properties are not well studied. One problem with \hat{S} , is that its estimation from

a finite sample has a positive bias that is inversely proportional to the population stationarity, but its convergence properties are not explored.

2.2 QS activity

Defining quasi-stationary waves as a climatological property of a set of atmospheric fields, precludes, in principle, the possibility of quantifying a QS metric that applies to instantaneous fields. It would seem impossible to, for example, construct a time series of QS activity that could be used as a basis for correlations with other variables, compositions or for use in other methodologies. But there are ways of solving this issue.

One possibility is recognising that individual fields can be characterised by their degree of similarity with the climatological QS. The index produced by Raphael (2004) for the QS3 is an example. While not expressly a measure of similarity, it is sensitive to wave 3 patterns with phase close to the stationary phase. Yuan and Li (2008) use Principal Component Analysis on the meridional wind field; the spatial pattern of the leading mode is very similar to the QS3 and so a time series can be obtained by projecting each instantaneous field to it.

Another way of constructing a time series is to exploit the fact the timescale dependence of QS. By applying a running mean with a suitable window before computing wave amplitudes, one obtains the QS wave amplitude of that window. This is the methodology applied by Wolf, Brayshaw, Klingaman, and Czaja (2018) who performed a 15 day low pass filter before computing wave envelopes. Each data time represented, then, the mean field of the set of fields covered the 15 day window and thus waves computed from it are actually QS waves for each of those sets. (*xx no estoy seguro que se entienda bien xx*)

3 Conclusions

The fact that zonal waves (ZW) and quasi-stationary waves (QS) are two distinct but related phenomena has both practical and theoretical implications.

First, researchers should be aware of which phenomena they want to study and use the appropriate methods. The mean amplitude of the ZW could be appropriate to study the vertical propagation of Rossby waves, for example. But ZW amplitude could lead to misleading results if used as the basis of local impacts studies because they are prob-

ably more influenced by phase effects. For clarity and reproducibility, we encourage researchers in the field to describe if they are using the mean amplitude of the individual waves or the amplitude of the mean wave.

Secondly, comparison between results should also be made having this issues in mind. For instance, Irving and Simmonds (2015) compare their planetary wave activity index with Raphael (2004)’s wave 3 index and conclude that the later cannot account for events with waves far removed from their climatological position. However, by understanding it as an index of QS3 similitude, this limitation becomes a feature, not a bug.

Since planetary waves are generally more stationary in the northern hemisphere, these issues are more critical for studies of the southern hemisphere.

Thirdly, the explorations of both ZW and QS can lead to novel levels of analysis. Here, we showed it can be used to define a metric of stationarity of quasi-stationary waves, but other applications are also possible. Smith and Kushner (2012) used the phase relationship between ZW1 and QS1 to show that linear interference between the QS1 and ZW1 was related to vertical wave activity transport at the tropopause.

xx me falta un final acá xx

References

- Hurrell, J. W., van Loon, H., & Shea, D. J. (1998). The Mean State of the Troposphere. In D. J. Karoly & D. G. Vincent (Eds.), *Meteorology of the southern hemisphere* (pp. 1–46). Boston, MA: American Meteorological Society. Retrieved from https://doi.org/10.1007/978-1-935704-10-2{_}1 doi: 10.1007/978-1-935704-10-2_1
- Irving, D., & Simmonds, I. (2015, dec). A novel approach to diagnosing Southern Hemisphere planetary wave activity and its influence on regional climate variability. *Journal of Climate*, 28(23), 9041–9057. Retrieved from <http://journals.ametsoc.org/doi/10.1175/JCLI-D-15-0287.1> doi: 10.1175/JCLI-D-15-0287.1
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., ... Joseph, D. (1996). The NCEP/NCAR 40-year reanalysis project. *Bulletin of the American Meteorological Society*, 77(3), 437–471. doi: 10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2

- 152 Kravchenko, V. O., Evtushevsky, O. M., Grytsai, A. V., Klekociuk, A. R., Mi-
 153 linevsky, G. P., & Grytsai, Z. I. (2012). Quasi-stationary planetary waves
 154 in late winter Antarctic stratosphere temperature as a possible indicator of
 155 spring total ozone. *Atmospheric Chemistry and Physics*, 12(6), 2865–2879.
 156 doi: 10.5194/acp-12-2865-2012
- 157 Lastovicka, J., Krizan, P., & Kozubek, M. (2018). Longitudinal structure of station-
 158 ary planetary waves in the middle atmosphere - Extraordinary years. *Annales*
 159 *Geophysicae*, 36(1), 181–192. doi: 10.5194/angeo-36-181-2018
- 160 Rao, V. B., Fernandez, J. P. R., & Franchito, S. H. (2004). Quasi-stationary waves
 161 in the southern hemisphere during El Nina and La Nina events. *Annales Geo-*
 162 *physicae*, 22(3), 789–806.
- 163 Raphael, M. N. (2004, dec). A zonal wave 3 index for the Southern Hemisphere.
 164 *Geophysical Research Letters*, 31(23), 1–4. Retrieved from [http://doi.wiley](http://doi.wiley.com/10.1029/2004GL020365)
 165 [.com/10.1029/2004GL020365](http://doi.wiley.com/10.1029/2004GL020365) doi: 10.1029/2004GL020365
- 166 Singer, I. A. (1967). *Steadiness of the Wind* (Vol. 6) (No. 6). Retrieved from
 167 [http://journals.ametsoc.org/action/doSearch?AllField=steadiness+](http://journals.ametsoc.org/action/doSearch?AllField=steadiness+of+the+wind{%&}filter=AllField{%}%5Cnhttp://dx.doi.org/10.1175{%}%2F1520-0450{%}%281967{%}%29006{%}%3C1033{%}%3ASOTW{%}%3E2.0.CO{%}%3B2)
 168 [of+the+wind{%&}filter=AllField{%}%5Cnhttp://dx.doi.org/10.1175{%}%](http://journals.ametsoc.org/action/doSearch?AllField=steadiness+of+the+wind{%&}filter=AllField{%}%5Cnhttp://dx.doi.org/10.1175{%}%2F1520-0450{%}%281967{%}%29006{%}%3C1033{%}%3ASOTW{%}%3E2.0.CO{%}%3B2)
 169 [2F1520-0450{%}%281967{%}%29006{%}%3C1033{%}%3ASOTW{%}%3E2.0.CO{%}%](http://journals.ametsoc.org/action/doSearch?AllField=steadiness+of+the+wind{%&}filter=AllField{%}%5Cnhttp://dx.doi.org/10.1175{%}%2F1520-0450{%}%281967{%}%29006{%}%3C1033{%}%3ASOTW{%}%3E2.0.CO{%}%3B2)
 170 [3B2](http://journals.ametsoc.org/action/doSearch?AllField=steadiness+of+the+wind{%&}filter=AllField{%}%5Cnhttp://dx.doi.org/10.1175{%}%2F1520-0450{%}%281967{%}%29006{%}%3C1033{%}%3ASOTW{%}%3E2.0.CO{%}%3B2) doi: 10.1175/1520-0450(1967)006<1033:sotw>2.0.co;2
- 171 Smith, K. L., & Kushner, P. J. (2012). Linear interference and the initiation of ex-
 172 tratropical stratosphere-troposphere interactions. *Journal of Geophysical Re-*
 173 *search Atmospheres*, 117(13), 1–16. doi: 10.1029/2012JD017587,2012
- 174 Turner, J., Hosking, J. S., Bracegirdle, T. J., Phillips, T., & Marshall, G. J. (2017).
 175 Variability and trends in the Southern Hemisphere high latitude, quasi-
 176 stationary planetary waves. *International Journal of Climatology*, 37(5),
 177 2325–2336. doi: 10.1002/joc.4848
- 178 Wolf, G., Brayshaw, D. J., Klingaman, N. P., & Czaja, A. (2018). Quasi-stationary
 179 waves and their impact on European weather and extreme events. *Quar-*
 180 *terly Journal of the Royal Meteorological Society*, 1–18. Retrieved from
 181 <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3310>
 182 doi: 10.1002/qj.3310
- 183 Yuan, X., & Li, C. (2008). Climate modes in southern high latitudes and their im-
 184 pacts on Antarctic sea ice. *Journal of Geophysical Research: Oceans*, 113(6),

