




Figure 1: **Projection on TICA and free energy profile of Chignolin.** Projection on the first 2 TIC coordinates (left) and free energy computed as logarithm of the probability for a folding trajectory of the protein Chignolin. The initial string is the string obtained just by transport ($\gamma = 0$). The MEP and MEP have a lower free energy than the initial string.




Figure 2: **Projection on TICA and free energy profile of BBA.** Projection on the first 2 TIC coordinates (left) and free energy computed as logarithm of the probability for a folding trajectory of the protein BBA. The initial string is the string obtained just by transport ($\gamma = 0$). The MEP and MEP have a lower free energy than the initial string.




Figure 3: **Histograms of the likelihood for the different methods.** Histogram of the likelihoods of the images for strings obtained using the different methods compared to the histogram of the likelihood of the validation set of Imagenet. Here are showed the histograms for the Minimum energy path and for the principal curve at three different temperatures

In the next we computed different strings with the method at finite temperature. For the intensity of the finite temperature algorithm with the interpolant time we had

$$\gamma(t) = 15 \frac{6(t_{final}-t)(t-t_{init})}{(t_{final}-t_{init})^3} \mathbb{1}_{\{t \in [t_{init}, t_{final}]\}}$$




Figure 4: **Images on the principal curve for different schedulings.** The principal curves have been computed with $T = 0.9$ and $\gamma(t) = 15 \frac{6(t_{final}-t)(t-t_{init})}{(t_{final}-t_{init})^3} \mathbf{1}_{\{t \in [t_{init}, t_{final}]\}}$ for $t_{final} = 0.95$ and a different value of t_{init} at every row.




Figure 5: **Walker Images for different schedulings.** The walker are associated to the principal curve in Fig4. The principal curves have been computed with $T = 0.9$ and $\gamma(t) = 15 \frac{6(t_{final}-t)(t-t_{init})}{(t_{final}-t_{init})^3} \mathbf{1}_{\{t \in [t_{init}, t_{final}]\}}$ for $t_{final} = 0.95$ and a different value of t_{init} at every row.