



**Politecnico
di Torino**

Politecnico di Torino

Master's Degree in Mathematical Engineering
A.a. 2025/2026
Graduation Session March 2026

DNA Methylation Dynamics Along the Normal–Adjacent Axis in Breast Cancer

A Quantitative Statistical and Machine Learning Approach

Supervisors:

Prof. Ing. Alfredo BENSO

Dott. Sandro GAMBINO

Candidate:

Elisabetta ROVIERA

ABSTRACT DELLA TESI

Acknowledgements

Table of Contents

List of Tables	IX
List of Figures	X
Glossary	XIII
1 Introduzione	1
2 Biological and Epigenetic Background	2
2.1 Breast Cancer: Epidemiology and Molecular Subtypes	2
2.2 The Hallmarks Framework and Epigenetic Reprogramming	3
2.3 DNA Methylation: Mechanism and Genomic Architecture	3
2.3.1 CpG Islands, Shores, Shelves, and Open Sea	4
2.3.2 Genomic Position Determines the Functional Consequence of Methylation	5
2.4 Epigenetic Alterations in Cancer: Too Much and Too Little	5
2.5 Epigenetic Drift, Aging, and Cancer Risk	6
2.6 Field Cancerization: The Concept and Its Epigenetic Basis	6
2.7 Differential Variability as a Statistical Paradigm	7
2.8 Synthesis: A Progressive Epigenetic Model of Breast Carcinogenesis	8
3 Data Preparation and Exploratory Analysis	10
3.1 Methodological Objectives	10
3.2 Dataset Construction, Metadata Integration, and Data Organization	11
3.2.1 Data Sources and Study Design	11
3.2.2 Methylation Data Representation and Storage	12
3.2.3 Phenotype Tables and Label Harmonization	13
3.2.4 Reproducibility and Computational Workflow	13
3.3 Intra-Dataset Exploration and Visualization	13
3.3.1 Dataset GSE69914	14
3.3.2 Dataset GSE225845	19

3.3.3	Dataset GSE287331	24
3.4	Inter Dataset Exploration and Comparison	29
3.4.1	Analytical Framework	29
3.4.2	Cross-Dataset Comparative Metric	30
3.4.3	Visual Comparative Analysis	32
3.5	Exploratory Findings and Pre-Processing Implications	35
4	Data preprocessing	36
4.1	Preprocessing Rationale and Technical Constraints	36
4.2	General Preprocessing Framework	37
4.2.1	Structural Integrity and Cohort Harmonization	37
4.2.2	Probe-Type Bias Diagnostics and Correction	38
4.2.3	Technical Probe Filtering	39
4.2.4	Statistical Transformation: β -to- M Values	40
4.3	Dataset-Specific Implementation	41
4.3.1	Dataset GSE69914	41
4.3.2	Dataset GSE225845	44
4.3.3	Dataset GSE287331	46
4.4	Inter-Dataset Consistency After Preprocessing	50
4.5	Post-Preprocessing Feature Space and Dimensionality Implications	51
5	Robust Feature Selection for Epigenetic Drift Characterisation	53
5.1	Statistical Challenges of High-Dimensional DNA Methylation Data	53
5.1.1	Structural Properties Motivating Feature Selection	54
5.1.2	Methodological Challenges in Genomic Feature Selection	54
5.1.3	Objectives of the Present Framework	55
5.2	General Feature Selection Framework	56
5.2.1	Empirical Variability-Based Dimensionality Reduction	56
5.2.2	Re-alignment to M -values and Residual Variance Regularisation	58
5.2.3	Biologically Weighted Stability Selection and Region-Level Consolidation	59
5.2.4	Correlation-Based Redundancy Pruning and Graph-Theoretic Clustering	63
5.2.5	Genomic Diversification via Constrained Greedy Selection	65
5.3	Dataset-Specific Implementation	67
5.3.1	Dataset GSE69914	67
5.3.2	Dataset GSE225845	70
5.3.3	Dataset GSE287331	72
5.4	Inter-Dataset Stability of the Final 5,000 CpG Signatures	75
5.4.1	Set-Level Replicability	75
5.4.2	Effect-Direction Concordance	77

5.4.3	Magnitude-Stratified Concordance	78
5.4.4	Cross-Dataset Transferability	79
5.5	Feature Selection Outcomes and Modelling Implications	81
6	Predictive Modelling, Constrained Selection and Biological Interpretation	83
6.1	Introduction	83
6.2	Predictive Framework	84
6.2.1	Intra-Dataset Supervised Modelling	84
6.2.2	Cross-Dataset Evaluation Protocol	86
6.2.3	Constrained Subset Optimisation: Knapsack Formulation	87
6.2.4	Gene Set and Functional Enrichment Analysis	88
6.3	Intra-Dataset Results	90
6.3.1	Dataset GSE69914	90
6.3.2	Dataset GSE225845	90
6.3.3	Dataset GSE287331	90
6.4	Inter-Dataset Results	90
6.4.1	Cross-Cohort Transferability of the 5,000-CpG Panel	90
6.4.2	Directional Concordance and Performance Degradation	90
6.4.3	Cross-Dataset Behaviour of the 50-CpG Subset	90
6.5	Synthesis and Implications	90
7	CpGPT - estensione cross dataset	91
8	Conclusion and future work	92
A	Filtering lists	93
A.1	Probe Filtering Resources	93
A.1.1	Naeem <i>et al.</i> (2014)	93
A.1.2	Chen <i>et al.</i> (2013)	93
A.1.3	Pidsley <i>et al.</i> (2016)	94
A.1.4	Zhou <i>et al.</i> (2016)	94
A.1.5	McCartney <i>et al.</i> (2016)	95
B	Extension of an Empirically Driven Variability-Based Filtering Framework to Breast Tissue	96
B.1	Motivation and Scope	96
B.2	General Framework for Cross-Tissue Extension	97
B.3	Cross-Tissue Overlap Structure	97
B.3.1	Step A: Stability of Non-Variable CpG Selection	98
B.3.2	Step B: Stability–Dimensionality Trade-Off	99
B.3.3	Step C: Methylation Distribution of Non-Variable CpGs	99

B.3.4	Step D: Baseline Variability Profile	100
B.4	Cohort-Resolved Characterisation of Breast-Tissue Non-Variability .	101
B.4.1	Empirical Variability Spectrum Across Independent Breast Cohorts	101
B.4.2	Contextual Enrichment Profile of Invariant CpGs	102
B.4.3	Phenotype-Resolved Variability Concordance Analysis	104
B.4.4	Invariant-Set Partitioning Across Phenotypic Combinations .	105
B.5	Positioning Within the Thesis and Generalisability	108
C	Plot cap6	109
D	Age Estimation	110
	Bibliography	111

List of Tables

3.1	Overview of the GSE69914 dataset after initial cohort curation. . .	14
3.2	Overview of the GSE225845 dataset after initial cohort curation. . .	20
3.3	Overview of the GSE287331 dataset after initial cohort curation. . .	24
3.4	Cross-dataset comparison of Normal (N) vs Adjacent (A) contrasts using global and instability metrics.	30
4.1	CpG filtering summary for the GSE69914 dataset.	42
4.2	CpG filtering summary for the GSE225845 dataset.	45
4.3	CpG filtering summary for the GSE287331 dataset.	49
4.4	CpG dimensionality before and after preprocessing.	51
5.1	Permutation-based enrichment of pairwise overlaps among the three final 5,000-CpG panels.	76
5.2	Number of CpGs retained at each step of the feature selection pipeline (training set).	81
A.1	Technical artefact categories covered by each filtering resource. . . .	94

List of Figures

3.1	Group-wise mean β -value density in GSE69914.	15
3.2	Heatmap of the top outlier CpGs (left) and corresponding $\Delta\beta$ distributions (right), illustrating CpG-level instability in GSE69914. .	16
3.3	Variance and correlation structure across Normal, Adjacent, and Tumor samples in GSE69914.	17
3.4	Low-dimensional embeddings samples in GSE69914.	17
3.5	Dynamic-network proxy visualizations in GSE69914.	18
3.6	Group-wise mean β -value density in GSE225845.	20
3.7	Heatmap of the top outlier CpGs (left) and corresponding $\Delta\beta$ distributions (right), illustrating CpG-level instability in GSE225845. .	21
3.8	Variance and correlation structure across Normal, Adjacent, and Tumor samples in GSE225845.	22
3.9	Low-dimensional embeddings samples in GSE225845.	22
3.10	Dynamic-network proxy visualizations in GSE225845.	23
3.11	Schematic representation of the five tissue categories collected along the tumor proximity axis (TPxA), reproduced from [37].	25
3.12	Group-wise mean β -value density in GSE287331.	26
3.13	Heatmap of the top outlier CpGs (left) and corresponding $\Delta\beta$ distributions (right), illustrating CpG-level instability in GSE287331. .	26
3.14	Variance and correlation structure across Normal, Adjacent, and Tumor samples in GSE287331.	27
3.15	Low-dimensional embeddings of GSE287331 samples.	28
3.16	Dynamic-network proxy visualizations for GSE287331.	28
3.17	Overlay of $\Delta\beta$ (Adjacent–Normal) density curves across the three datasets, computed on the 326,330 CpG sites shared by all cohorts.	33
3.18	Pairwise $\Delta\beta$ (Adjacent–Normal) scatterplots across the three methylation datasets, computed on the 326,330 shared CpG sites. Spearman ρ values are reported in each panel.	34
3.19	t-SNE embeddings computed on the 326,330 CpGs shared across GSE69914, GSE225845, and GSE287331.	34

4.1	Infinium Type I/II bias diagnostics in GSE69914.	41
4.2	Technical filtering summary in GSE69914.	42
4.3	Distributional comparison of β and M in Normal and Adjacent tissues in GSE69914.	43
4.4	Infinium Type I/II bias diagnostics in GSE225845.	44
4.5	Technical filtering summary in GSE225845.	45
4.6	Distributional comparison of β and M in Normal and Adjacent tissues in GSE225845.	46
4.7	Per-sample mean β distributions for the three datasets.	47
4.8	Infinium Type I/II bias diagnostics in GSE287331. Pre-BMIQ densities and Q-Q analysis indicate marked probe-type imbalance; post-BMIQ densities show partial alignment.	47
4.9	Technical filtering summary in GSE287331.	48
4.10	Distributional comparison of β and M in Normal and Adjacent tissues in GSE287331.	50
4.11	CpG overlap across datasets before and after preprocessing.	51
5.1	Variability-based filtering in β -space and residual variance trimming in M -space in GSE69914.	68
5.2	Stability enforcement and correlation-based redundancy structure in GSE69914.	69
5.3	Variability-based filtering in β -space and residual variance trimming in M -space in GSE225845.	70
5.4	Stability and redundancy diagnostics in GSE225845.	71
5.5	Variability-based filtering in β -space and residual variance trimming in M -space in GSE287331.	73
5.6	Stability and redundancy diagnostics in GSE287331.	74
5.7	Three-way overlap of the final 5,000 CpG signatures independently selected in GSE69914, GSE225845 and GSE287331.	75
5.8	Inter-dataset effect concordance of the final 5,000 CpG signatures.	77
B.1	Global overlap structure of non-variable CpGs across blood, buccal epithelial cells, placenta, and breast tissue.	98
B.2	Assessment of robustness and trade-off between stability and dimensionality of non-variable CpG sets.	99
B.3	Characterization of the selected non-variable CpG set in terms of methylation distribution and baseline risk behavior.	100
B.4	Distribution of r_β computed on the Normal+Tumor comparison for each dataset. Vertical dashed lines indicate the selected threshold.	103

B.5	Fold-enrichment of genomic context categories among invariant CpGs relative to the background distribution in each dataset. The dashed line indicates enrichment equal to background (fold = 1).	104
B.6	Spearman correlation between CpG-level $\Delta\beta$ values computed in the Normal+Adjacent and Normal+Tumor contrasts for each dataset. The diagonal line represents identity.	105
B.7	Overlap between universal invariants and breast-specific invariant sets derived from the Normal+Adjacent and Normal+Tumor contrasts in each dataset.	107

Glossary

Chapter 1

Introduzione

INTRODUZIONE DELLA TESI

Chapter 2

Biological and Epigenetic Background

2.1 Breast Cancer: Epidemiology and Molecular Subtypes

Breast cancer is the most frequently diagnosed malignancy in women worldwide and represents a major public health burden across both high- and low-income countries [1]. Its clinical and biological heterogeneity is one of the most distinctive features of the disease: not all breast cancers behave alike, progress at the same rate, or respond to the same treatments.

At the histological level, the vast majority of breast malignancies originate from the epithelial compartment of the mammary gland, most commonly from the luminal cells lining the ductal and lobular structures. The classical progression model describes a continuum from normal epithelium through non-atypical hyperplasia, atypical ductal hyperplasia (ADH), ductal carcinoma in situ (DCIS), and ultimately invasive carcinoma [2]:

Normal epithelium $\rightarrow$ Hyperplasia $\rightarrow$ DCIS $\rightarrow$ Invasive carcinoma.

At the molecular level, tumors are stratified according to the expression status of estrogen receptor (ER), progesterone receptor (PR), and HER2, defining biologically distinct subtypes — luminal A, luminal B, HER2-enriched, and triple-negative — with different prognostic and therapeutic implications. This molecular classification reflects underlying differences in the cell of origin, epigenetic landscape, and selective pressures operative during tumor evolution.

A subset of breast cancers arises on an inherited background of germline susceptibility. Pathogenic variants in *BRCA1* and *BRCA2*, genes encoding key components

of the homologous recombination DNA repair machinery, substantially increase lifetime risk [3]. BRCA-associated tumors tend to be triple-negative and exhibit distinct baseline epigenetic profiles that differ fundamentally from sporadic disease [4, 5]. For this reason, BRCA mutation carriers require separate analytical treatment in any study of epigenetic field effects, as discussed in Chapter ??.

2.2 The Hallmarks Framework and Epigenetic Reprogramming

The conceptual framework of cancer biology has been progressively organized around the notion of “hallmarks”: a set of functional capabilities that normal cells must acquire to become malignant [2, 1]. These include sustained proliferative signaling, resistance to growth suppression, evasion of apoptosis, unlimited replicative potential, induction of angiogenesis, and the activation of invasion and metastasis.

Importantly, the updated hallmarks framework explicitly recognizes epigenetic reprogramming as a core enabling mechanism [1]. Tumor cells do not merely accumulate genetic mutations; they undergo widespread reorganization of their chromatin architecture and DNA methylation landscape, which silences tumor-suppressive programs and derepresses oncogenic pathways. This repositions epigenetic alterations not as secondary consequences of malignant transformation but as primary and early drivers of the tumorigenic process.

2.3 DNA Methylation: Mechanism and Genomic Architecture

DNA methylation consists of the covalent addition of a methyl group to the 5-carbon position of cytosine residues within CpG dinucleotides, a reaction catalyzed by a family of DNA methyltransferases (DNMT3A, DNMT3B for *de novo* methylation; DNMT1 for maintenance after replication) [26]. CpG dinucleotides are non-randomly distributed in the genome: statistically expected at roughly one occurrence per 16 bp, they are in practice four- to fivefold depleted genome-wide — a depletion attributed to the historical mutability of methylated cytosines, which deaminate spontaneously to thymine. The exception to this depletion is found at *CpG islands*: dense, GC-rich clusters of CpG dinucleotides, typically 500–2000 bp in length, located at the promoters of approximately 60% of human protein-coding genes [26]. In normal somatic cells, CpG islands are kept constitutively unmethylated by active mechanisms, preserving transcriptional access.

2.3.1 CpG Islands, Shores, Shelves, and Open Sea

The regulatory significance of DNA methylation varies substantially depending on where in the genome a CpG locus resides. A widely adopted classification, used in Illumina array annotation, distinguishes four positional contexts relative to CpG islands [7]:

- **CpG Islands:** regions with high CpG density and high GC content, operationally defined as stretches of ≥ 200 bp with CpG observed/expected ratio > 0.6 and GC content $> 50\%$. Methylation at island-overlapping promoters is the canonical mechanism of transcriptional silencing in both development and cancer.
- **Shores:** regions within 2 kb flanking a CpG island on either side. Although less CpG-dense than islands, shores are the sites of the most dynamic and biologically informative methylation variation. Landmark work by Irizarry et al. demonstrated that tissue-specific methylation differences between normal cell types are concentrated at shores rather than at islands, and that cancer-associated hypermethylation also preferentially targets shores [7]. This makes shores the primary locus of epigenetic switching between cell states.
- **Shelves:** regions 2–4 kb from the island boundary. Their methylation patterns are less variable than shores and less directly tied to promoter regulation, though they retain some contextual regulatory relevance.
- **Open Sea:** CpG loci not associated with any island, shore, or shelf. These constitute the majority of CpGs in the genome ($\sim 70\%$) and tend to be constitutively methylated in somatic tissues. Their methylation changes are more often associated with global hypomethylation and genomic instability than with specific gene regulation.

This spatial hierarchy has direct consequences for how methylation data should be interpreted and weighted in analytical frameworks. Loci overlapping islands and shores are enriched for functional regulatory activity — they are more likely to be associated with gene expression changes when their methylation state shifts. Shelf and open-sea loci, by contrast, tend to reflect structural genomic states rather than specific transcriptional events. When constructing feature selection or scoring systems based on biological relevance, this gradient from island to open sea provides a principled basis for differential weighting.

2.3.2 Genomic Position Determines the Functional Consequence of Methylation

A further distinction of biological relevance concerns the relationship between methylation and transcription depending on the position of the CpG locus relative to the gene structure. The classical rule — methylation silences expression — applies specifically to *promoter-associated* CpG islands. Within the gene body (introns and exons), the relationship is inverted: gene body methylation correlates *positively* with transcriptional activity [8]. This is thought to reflect the need to suppress spurious transcription initiation within highly transcribed genes. The functional interpretation of a methylation change therefore depends critically on whether the affected locus falls in a promoter or an intragenic context.

In genome-wide methylation profiling studies, the Illumina array annotation provides positional information (TSS200, TSS1500, 5'UTR, 1stExon, Body, 3'UTR) that, combined with the island/shore/shelf classification, allows a nuanced characterization of each locus. CpGs annotated to TSS regions within islands or shores carry the highest prior probability of regulatory relevance and are therefore the most informative for studies of transcriptional silencing in cancer.

2.4 Epigenetic Alterations in Cancer: Too Much and Too Little

Cancer genomes are characterized by a paradoxical co-occurrence of global hypomethylation and focal hypermethylation [25]. This duality was first described in the early 1980s and has since been confirmed across virtually all tumor types, including breast cancer.

Global hypomethylation affects repetitive sequences (satellite DNA, LINE-1 elements) and is thought to promote genomic instability through aberrant activation of transposable elements and loss of imprinting. It also results in transcriptional derepression at genes that are normally silenced [25].

Focal hypermethylation, by contrast, targets CpG islands at the promoters of specific genes, leading to heritable transcriptional silencing. Affected genes frequently include classical tumor suppressors (e.g., *CDKN2A*, *BRCA1*, *MLH1*), as well as genes involved in apoptosis, cell adhesion, and developmental signaling. This epigenetic silencing is functionally equivalent to mutation but is potentially reversible, making it an attractive therapeutic target.

Critically, neither of these phenomena is random. Genes susceptible to hypermethylation in cancer are systematically enriched for targets of the Polycomb Repressive Complex 2 (PRC2) and display bivalent chromatin states in embryonic

stem cells — a configuration that marks developmental genes as both transcriptionally poised and repressed [10]. This connection between developmental regulation and cancer-associated silencing suggests that epigenetic alterations in malignancy partially recapitulate stem cell programs.

2.5 Epigenetic Drift, Aging, and Cancer Risk

Aging is the single strongest risk factor for breast cancer. The incidence of most breast malignancies rises sharply with age, and this epidemiological observation has a direct molecular correlate in the epigenome.

Over the course of a lifetime, DNA methylation patterns undergo progressive and partially stochastic changes collectively referred to as *epigenetic drift* [11]. This phenomenon is characterized by gradual increases in methylation variability across both CpG loci and individuals. The drift is not uniform: it preferentially affects genes associated with Polycomb repression and bivalent chromatin — precisely the same loci that tend to become hypermethylated in cancer [12, 10].

Horvath’s landmark study demonstrated that DNA methylation patterns can be exploited to construct a highly accurate molecular clock: the “epigenetic age” of a tissue, estimated from a weighted combination of CpG methylation values, correlates strongly with chronological age across more than 50 tissue types [11]. Crucially, epigenetic age acceleration — a discordance between molecular and chronological age — has been associated with cancer risk, suggesting that the pace of epigenetic drift itself may be a biological marker of malignant susceptibility.

From a mathematical perspective, aging-associated drift can be modeled as a diffusion-like process in epigenetic state space: methylation levels at susceptible loci undergo a gradual increase in inter-individual variance over time. This growing dispersion may create a permissive landscape for malignant transformation by producing a fraction of cells with aberrant epigenetic configurations from which selection can act.

An important study by Teschendorff et al. in breast tissue documented precisely this mechanism: aging-related DNA methylation changes in normal breast tissue were found to target the same genomic regions altered in cancer, establishing a quantitative link between physiological aging and pre-malignant epigenetic reprogramming [13].

2.6 Field Cancerization: The Concept and Its Epigenetic Basis

A central concept motivating the present thesis is that of *field cancerization*. Originally proposed by Slaughter in 1953 on the basis of histopathological observations

in oral cancer, the field effect hypothesis posits that carcinogenesis does not occur in isolation but rather within a “conditioned” tissue field that has been broadly reprogrammed by molecular alterations preceding frank malignancy.

In breast cancer, epigenetic field defects have been systematically characterized using genome-wide methylation profiling. A foundational contribution by Teschendorff et al. demonstrated that histologically normal breast tissue adjacent to tumors harbors a specific pattern of methylation alterations distinct from both distant normal tissue and from tumor tissue itself [14]. These alterations are not random: they are enriched at PRC2 target genes, CTCF-binding regions, and loci involved in developmental pathways including WNT and FGF signaling — precisely the categories implicated in the aging-cancer methylation axis described above.

Crucially, the field defect signature identified by Teschendorff et al. was detectable not through classical differential mean analysis, but through *differential variability* — a statistically distinct framework that quantifies the increase in methylation dispersion across samples rather than directional shifts in average levels [14, 15]. This distinction has important methodological implications. Under a classical framework, loci with increased variance but small mean differences are invisible to standard hypothesis tests. Detecting them requires variance-sensitive statistics, such as the Levene test or its robust variants. The biological interpretation is that field defects reflect a partial and heterogeneous epigenetic reprogramming: not all cells in the adjacent tissue are affected equally, and the stochastic nature of the alteration is reflected as increased dispersion across individuals rather than as a uniform shift.

The progressive nature of epigenetic field formation is further supported by the observation that hypervariable loci in adjacent tissue tend to become hypermethylated in the matched invasive tumor, suggesting a deterministic convergence from a heterogeneous pre-malignant landscape to a consolidated malignant epigenetic state [14]:

Normal → Epigenetically reprogrammed adjacent tissue → Invasive carcinoma.

This progressive model implies that the adjacent tissue is not merely “contaminated” by the adjacent tumor but rather represents an earlier stage on the same carcinogenic trajectory. Detecting this intermediate stage — and characterizing its methylation signature — is therefore of both biological and clinical significance.

2.7 Differential Variability as a Statistical Paradigm

The discovery that field defects are better captured by differential variability than by differential means represents a genuine shift in the analytical paradigm for epigenetic cancer studies [15].

Formally, let X_j denote the methylation level at locus j across a cohort of samples. Classical differential methylation analysis focuses on the comparison of group means:

$$\Delta_j = \mathbb{E}[X_j \mid A] - \mathbb{E}[X_j \mid B]. \quad (2.1)$$

Differential variability analysis instead targets the comparison of group variances:

$$\delta\sigma_j^2 = \text{Var}(X_j \mid A) - \text{Var}(X_j \mid B). \quad (2.2)$$

A locus is said to be *differentially variable* (DV) if $\delta\sigma_j^2$ is significantly non-zero. Under a pre-malignant model, one expects $\delta\sigma_j^2 > 0$ in normal-adjacent tissue relative to distant normal tissue, reflecting increased epigenetic disorder in the field-affected region.

Teschendorff et al. formalized this framework and developed statistical tools for its application to genome-wide methylation data [15]. Their analysis demonstrated that DV CpGs in normal-adjacent breast tissue are substantially enriched for biologically meaningful genomic features (PRC2 targets, bivalent domains) and display stronger associations with cancer risk markers — including tumor proliferation as measured by KI67 — than differentially methylated CpGs identified by mean-based tests alone [14].

The implication for any analytical framework aimed at characterizing early epigenetic alterations is direct: a reliance on mean-based feature selection alone will systematically miss a biologically relevant component of the pre-malignant signal. This motivates the integration of variability-aware preprocessing and feature selection strategies, which are developed in the methodological chapters of this thesis.

2.8 The Normal–Adjacent–Tumor Axis as an Analytical Framework

Breast carcinogenesis is not a sudden event but a progressive deviation from epigenetic homeostasis, driven by the interplay of aging, stochastic drift, and selective pressures operating across a population of cells. The biological evidence reviewed in this chapter converges on a model in which this progression is both measurable and partially predictable from methylation data alone.

Normal breast tissue accumulates methylation variability with age at a specific set of loci — those associated with Polycomb regulation and bivalent chromatin. This drift creates a heterogeneous epigenetic landscape from which pre-malignant field defects emerge. Adjacent tissue, histologically indistinguishable from normal, harbors a measurable epigenetic signature of this reprogramming — detectable predominantly as increased variability rather than as a directional shift in mean

methylation. As carcinogenesis proceeds, this heterogeneous pre-malignant state converges toward a more consolidated malignant configuration characterized by focal hypermethylation at tumor-suppressive loci and global hypomethylation elsewhere.

This progressive model has three direct consequences for the quantitative analyses developed in this thesis. First, the Normal–Adjacent–Tumor axis should be treated as a continuous biological spectrum rather than a set of discrete categories: the adjacent tissue occupies a genuinely intermediate epigenetic state, distinct from both normal and tumor, and warrants explicit characterization rather than collapsing into either extreme. Second, any feature selection strategy operating on this axis must be sensitive to variance differences and not only to mean shifts: field defects are by definition heterogeneous signals that may not survive standard mean-based filtering. Third, cross-cohort reproducibility is a necessary validation criterion: signatures that replicate across independent datasets are more likely to reflect genuine biological structure than cohort-specific noise.

These three principles — spectrum-aware design, variance sensitivity, and cross-cohort validation — directly motivate the construction, preprocessing, and comparative characterization of the three independent breast tissue methylation cohorts introduced in Chapter 3.

Chapter 3

Data Preparation and Exploratory Analysis

3.1 Methodological Objectives

This chapter establishes the methodological foundation for all subsequent analyses by addressing the construction, organization, and preliminary characterization of the DNA methylation datasets used in this thesis. Following the biological background and problem formulation introduced in Chapters 1 and ??, the present chapter acts as a bridge between domain-specific knowledge and quantitative modeling, ensuring that downstream analyses are grounded on a technically sound and well-understood data representation.

Genome-wide DNA methylation data are characterized by extremely high dimensionality, heterogeneous signal-to-noise ratios, and the coexistence of subtle biological effects with multiple sources of technical variability. In this context, premature application of statistical models or feature-selection procedures may lead to biased conclusions driven by artefacts rather than genuine biological structure. A careful examination of data quality, distributional properties, and internal consistency is therefore a necessary prerequisite for any reliable inference [16].

The primary objective of this chapter is to construct a coherent and reproducible representation of the available datasets and to assess their structural and statistical properties prior to any form of preprocessing or modeling. Specifically, this chapter aims to: (i) describe the organization of methylation matrices and associated phenotype information; (ii) characterize the global and local variability of methylation levels within each dataset; (iii) evaluate the degree of separation between Normal, Adjacent, and Tumor tissues using exploratory tools; and (iv) compare datasets at an inter-cohort level to assess consistency, heterogeneity, and potential limitations in signal transferability.

The scope of this chapter is deliberately restricted to exploratory and diagnostic analyses. No feature selection, predictive modeling, or formal statistical inference is performed at this stage. All analyses are intended to be descriptive and comparative, serving to highlight structural properties of the data and to identify potential technical or biological issues that must be addressed before proceeding further.

Methodologically, the chapter is organized around a unified exploratory framework applied consistently across all datasets. Intra-dataset analyses are used to investigate internal structure, variability patterns, and group-level behavior, while inter-dataset comparisons focus on assessing cross-cohort consistency and platform-related differences. Throughout the chapter, visualizations and summary metrics are employed as diagnostic tools rather than as decision-making instruments.

The results of this exploratory phase reveal both shared characteristics and dataset-specific behaviors, as well as clear limitations of working directly with raw or minimally processed methylation data. These observations motivate the need for a structured and rigorous data pre-processing pipeline, which is developed and justified in the following chapter. The present chapter therefore provides the conceptual and technical basis for the preprocessing choices adopted in Chapter 4 and for all subsequent modeling steps.

3.2 Dataset Construction, Metadata Integration, and Data Organization

This section provides the methodological framework for the selection, construction of the dataset, and the organisation process adopted in this study. It contextualises the data sources, representation choices, metadata management, and computational design principles that underpin the detailed implementation described in the following sections.

3.2.1 Data Sources and Study Design

All datasets analyzed in this thesis were obtained from the NCBI Gene Expression Omnibus (GEO) [17], which was selected as a unified data source to minimize heterogeneity in data formats, metadata conventions, and access mechanisms. GEO provides curated series-level accessions, stable sample identifiers, and standardized links to processed methylation matrices and sample-level annotations, making it a natural reference repository for large-scale DNA methylation studies.

The study design adopts a multi-cohort comparative framework focused on genome-wide DNA methylation in breast tissue. Three independent GEO series were selected for analysis: **GSE69914** [18], based on the Illumina HumanMethylation450 platform, and two more recent EPIC-based cohorts, **GSE225845** [19] and

GSE287331 [20]. Together, these datasets span normal/healthy, adjacent-normal, and tumor tissue states across two generations of Illumina methylation arrays (450K and EPIC), enabling both within-dataset characterization and cross-dataset comparisons.

Dataset selection was driven by strict inclusion criteria aimed at ensuring technical compatibility, biological relevance, and analytical feasibility. In particular, only studies providing processed genome-wide methylation matrices in terms of β -values were considered, allowing a uniform data representation and avoiding the need for low-level signal reprocessing from raw `.idat` intensity files. Raw `.idat` files store probe-level fluorescence intensities measured in the methylated ($y^{(M)}$) and unmethylated ($y^{(U)}$) channels, from which β -values are derived as a normalized ratio [21]:

$$\beta := \frac{\max(y^{(M)}, 0)}{\max(y^{(M)}, 0) + \max(y^{(U)}, 0) + \alpha}, \quad \beta \in [0, 1]. \quad (3.1)$$

where α is a small offset (typically 100) used to stabilize the denominator. Additional requirements included the use of Illumina HumanMethylation450 (450K) or MethylationEPIC (EPIC) platforms, the availability of multiple tissue classes within each cohort, and sufficiently large sample sizes to support robust statistical analysis.

3.2.2 Methylation Data Representation and Storage

In several datasets, the original authors applied established normalization procedures, including methods specifically designed to correct probe-type design bias such as BMIQ [22]. Reprocessing raw IDAT files would therefore not add information relevant to the objectives of this thesis, while substantially increasing technical complexity and reducing reproducibility. For these reasons, all analyses were conducted on the processed β -value matrices as distributed via GEO.

To support scalable analysis and efficient input/output operations, all working methylation matrices were converted to a standardized `sample` $\times$ `CpG` layout and stored in columnar Parquet format using single-precision floating-point encoding.

This choice is motivated by both numerical and computational considerations. Since β -values are strictly bounded in the interval $[0, 1]$ [21], and biologically meaningful methylation differences typically occur at magnitudes on the order of 10^{-2} – 10^{-3} , single-precision floating point (`float32`, machine $\varepsilon \approx 10^{-7}$) provides more than sufficient numerical accuracy. At the same time, it substantially reduces memory usage and I/O overhead compared to double precision.

This representation is consistent with recent large-scale genomics frameworks that process molecular features, including DNA methylation data, entirely in `float32` precision for efficiency and scalability [23].

3.2.3 Phenotype Tables and Label Harmonization

For each dataset, a dedicated phenotype table was constructed by extracting and parsing GEO sample-level metadata. These tables encode tissue labels, sample identifiers, and relevant clinical or technical covariates when available. Given the heterogeneity of original annotations across studies, tissue classes were mapped to harmonized numeric encodings to ensure consistency across cohorts while preserving dataset-specific attributes in separate fields.

A strict one-to-one correspondence between methylation matrices and phenotype tables was enforced. Samples lacking either methylation measurements or corresponding metadata were excluded to prevent inconsistencies and downstream misalignment. This design guarantees that all analyses operate on synchronized molecular and phenotypic representations and enables reliable stratification by tissue class in both intra- and inter-dataset comparisons.

3.2.4 Reproducibility and Computational Workflow

The overall data organization and computational design follow a multi-dataset analytical paradigm commonly adopted in recent large-scale DNA methylation studies, in which cohort-specific analyses are complemented by cross-dataset comparisons to explicitly evaluate robustness and generalizability, while simultaneously revealing cohort-dependent effects and limitations in signal transferability that motivate subsequent methodological refinement [24].

All dataset construction and organization steps were implemented using deterministic, script-based pipelines with explicit schema definitions and typed data representations. Particular care was taken to avoid unnecessary in-memory operations on high-dimensional methylation matrices, favoring streaming ingestion, chunked processing, and out-of-core transformations when required by dataset size.

The resulting data organization yields a set of standardized, self-contained datasets, each consisting of a genome-wide methylation matrix and an aligned phenotype table stored in efficient, portable formats. This design ensures full reproducibility of the data preparation process and provides a robust foundation for the preprocessing, exploratory analysis, and modeling steps developed in subsequent chapters.

3.3 Intra-Dataset Exploration and Visualization

This section presents a structured intra-dataset exploratory analysis aimed at characterising the internal statistical and biological organisation of each methylation cohort prior to any preprocessing or modelling step. Each dataset is analysed independently in order to assess data integrity, quantify variability patterns, and

Table 3.1: Overview of the GSE69914 dataset after initial cohort curation.

CpGs	Samples	Normal	Adjacent	Tumor	Normal (BRCA1)	Tumor (BRCA1)
485,512	407	50	42	305	3	7

evaluate whether Normal, Adjacent, and Tumor tissues exhibit distinguishable epigenetic behaviour.

In the context of breast cancer, epigenetic deregulation is known to manifest through both global and local phenomena, including genome-wide hypomethylation, focal hypermethylation of tumour-suppressor regions, and the presence of pre-neoplastic “field defects” in histologically normal tissues proximal to tumours [25, 26, 27]. These mechanisms motivate an exploratory investigation focused not only on overt tumour-associated changes, but also on more subtle alterations in Normal and Adjacent samples, which are central to the biological hypothesis of this thesis.

This section therefore applies a unified exploratory framework to each dataset (GSE69914 [3.3.1], GSE225845 [3.3.2], GSE287331 [3.3.3]), systematically examining data quality, global methylation distributions, sample-level summaries, CpG-wise variability and instability, correlation structure, dimensionality reduction, and genomic context coverage. The use of a consistent analytical protocol enables direct comparison of intra-dataset behaviour while preserving cohort-specific characteristics.

3.3.1 Dataset GSE69914

Dataset overview GSE69914 is a breast tissue DNA methylation cohort profiled using the Illumina Infinium HumanMethylation450 BeadChip, providing genome-wide coverage of approximately 4.8×10^5 CpG loci. The dataset comprises samples spanning the full Normal–Adjacent–Tumor spectrum that motivates the biological framework of this thesis.

An overview of the sample composition and CpG coverage is reported in Table 3.1, including the distribution across tissue classes and hereditary-risk subgroups.

A limited subset of samples is associated with germline *BRCA1* mutations. *BRCA1*, in particular, encodes a key DNA repair protein, and pathogenic variants are known to substantially increase lifetime breast and ovarian cancer risk [28]. Multiple studies have shown that BRCA-associated breast tissues exhibit distinct baseline epigenetic profiles, reflecting inherited genomic instability rather than sporadic tumorigenesis [5, 4].

Since the primary objective of this thesis is to characterise early, field defect DNA methylation alterations along the Normal–Adjacent axis in sporadic breast cancer, samples carrying *BRCA1* mutations were excluded from the core exploratory and




Figure 3.1: Group-wise mean β -value density in GSE69914.

preprocessing pipeline. Their inclusion would introduce a confounding hereditary epigenetic signal, potentially obscuring subtle methylation shifts in non-mutated normal and adjacent tissues.

Global methylation distributions To characterise the overall methylation landscape of the dataset, the distribution of β -values (fractional methylation in $[0,1]$) was first examined. The group-wise mean density curve (Figure 3.1) displays the characteristic *bimodal* profile of Illumina 450k arrays, with peaks near unmethylated ($\beta \approx 0$) and fully methylated ($\beta \approx 1$) CpG sites. This pattern reflects the underlying biology of CpG regulation, where many loci tend to be either transcriptionally active (hypomethylated) or repressed (hypermethylated) [26, 29].

When comparing tissue groups, a clear gradient emerges. Tumor samples show a slightly flatter high- β peak and a broader low- β tail, consistent with the well-described phenomenon of global hypomethylation and increased heterogeneity in cancer [25]. Adjacent samples lie between Normal and Tumor, suggesting early epigenetic drift and subtle field defects occurring in histologically non-neoplastic tissue [27].

CpG-level instability and recurrent outliers To characterise locus-specific instability, the 200 CpG sites with the highest outlier burden across samples were examined. The heatmap of the top outlier loci (Figure 3.2a) indicates that epigenetic disruption is *not uniform*: specific CpG sites and specific samples display extreme deviations, rather than a diffuse genome-wide shift. Moreover, both directions of alteration are present — focal hypermethylation and focal hypomethylation. In cancer biology, hypermethylation can silence tumor-suppressor regions, whereas hypomethylation can derepress oncogenic pathways and weaken genomic stability, reflecting the classic “too much and too little methylation” behaviour of tumor genomes [25].




Figure 3.2: Heatmap of the top outlier CpGs (left) and corresponding $\Delta\beta$ distributions (right), illustrating CpG-level instability in GSE69914.

Complementary $\Delta\beta$ distributions comparing Tumor and Adjacent tissues against Normal (Figure 3.2b) show a clear shift toward hypomethylation in Tumor samples and a subtler but detectable drift in Adjacent tissues. This provides evidence that epigenetic alterations emerge early in histologically normal tissue located near the tumor, supporting the field-cancerization model [27].

Variance and correlation structure To quantify within-group epigenetic variability, the distribution of CpG-wise variance across samples was examined. The density curves (Figure 3.3a) show that Tumor samples have markedly higher variance, reflecting increased epigenetic instability. In contrast, Normal and Adjacent samples display almost identical variance distributions that are tightly concentrated near zero. This indicates that, at the level of CpG-wise variability, Adjacent tissues do not yet exhibit detectable divergence from Normal samples.

A complementary perspective is provided by the sample correlation heatmap (Figure 3.3b), which shows the expected high level of pairwise similarity across all samples. This behaviour is typical of genome-wide DNA methylation data, where a large fraction of CpG sites is stable across individuals, resulting in consistently strong correlations [22]. Importantly, the heatmap also reveals that all samples are highly mutually correlated, with no sharp blocks or boundaries separating the three tissue labels. This confirms that global methylation structure is largely conserved across Normal, Adjacent and Tumor tissues, and that group differences are too subtle to be detected through sample-sample correlation alone.




Figure 3.3: Variance and correlation structure across Normal, Adjacent, and Tumor samples in GSE69914.




Figure 3.4: Low-dimensional embeddings samples in GSE69914.

Low-dimensional embeddings Dimensionality reduction was applied using Principal Component Analysis (PCA), t-distributed Stochastic Neighbor Embedding (t-SNE), and Uniform Manifold Approximation and Projection (UMAP) to visualise global similarities among samples (Figure 3.4). Across all three methods, Normal and Adjacent samples show substantial overlap, with no clear separation between these two groups. Tumor samples appear more dispersed, but only mildly shifted relative to the Normal/Adjacent cluster, and no sharp cluster boundaries emerge in any embedding. The trustworthiness scores are similar across embeddings (0.869–0.888), indicating that local neighbourhood relationships are largely preserved in the low-dimensional projections.

These projections indicate that global methylation patterns alone do not provide strong discriminative structure among the three tissue states. This suggests that group differences are subtle at the whole-methylome level and are more effectively captured through locus-specific analyses rather than through global unsupervised embeddings.




Figure 3.5: Dynamic-network proxy visualizations in GSE69914.

Genome annotation coverage CpG probes were annotated using the official *Infinium MethylationEPIC v1.0 B5* [30] manifest file, the standard Illumina resource containing genomic coordinates, probe type, CpG island context (Island, Shore, Shelf, Open Sea), and gene-level annotations. Using this manifest, the dataset exhibits the expected non-uniform genomic distribution of CpG contexts: the largest fraction of probes maps to open-sea or non-island regions, followed by CpG islands, then north and south shores, and finally north and south shelves. This ordering mirrors the characteristic architecture of the HM450 arrays reported in the literature [31, 32]. The coverage observed is therefore consistent with the known genomic composition of the Illumina methylation arrays, supporting correct manifest alignment.

Dynamic-network proxies Although this analysis does not follow the full mathematical formalism of Dynamic Network Biomarkers (DNBs), the degree-based and network-level (Figure 3.5) provide a qualitative view of how methylation perturbations may organize at the network level. DNB theory, originally formulated for gene-expression and regulatory networks, predicts that near a critical transition a subset of components displays increased internal fluctuations and coordinated behaviour [33, 34, 35]. Here, this conceptual framework is adapted to CpG-level methylation patterns by examining the structure of CpG-gene degree distributions and the organization of $\Delta\beta$ -driven network layouts. This approach enables the exploration of whether tumour-associated instability manifests as locally coordinated methylation shifts at the network level, even without implementing the complete DNB pipeline.

The degree-rank plot (Figure 3.5a) displays a heavy-tailed distribution of CpG counts per gene, consistent with the heterogeneous hub-periphery structure typically observed in biological systems. In the Tumor vs Normal comparison (Figure 3.5b), a compact nucleus of nodes with moderately high degree is observed, accompanied by larger $\Delta\beta$ deviations. This pattern is consistent with a more perturbed and locally

coordinated subset of CpG/gene units in tumor tissue, qualitatively compatible with increased network instability. In contrast, the Adjacent vs Normal network (Figure 3.5c) does not exhibit a comparable hub-like concentration and shows smaller, more diffuse deviations, suggesting that the adjacent tissue reflects only early or weakly organized perturbations.

Although these observations do not satisfy the full mathematical criteria for identifying a DNB module, they are conceptually consistent with the interpretation that tumor tissue occupies a more unstable network regime, whereas adjacent tissue remains closer to a stable (pre-transition) configuration.

Overall interpretation The exploratory analysis of GSE69914 reveals a consistent but overall subtle separation between the three tissue states. Normal and Adjacent samples exhibit highly similar global methylation profiles, variance distributions, correlation structure, and low-dimensional embeddings, indicating that large-scale methylome organisation remains largely conserved in Adjacent tissue. Across these global summaries, the two groups are so intermixed that no clear or marked distinction emerges between Normal and Adjacent samples when using whole-methylome descriptors. Tumor samples display the expected increase in heterogeneity and a shift toward hypomethylation; however, these differences remain modest at the whole-methylome level and do not produce clear separation in unsupervised embeddings.

Locus-specific analyses, however, highlight focused patterns of disruption. Outlier CpGs exhibit both hyper- and hypomethylation events, and $\Delta\beta$ distributions reveal a distinct hypomethylation bias in Tumor samples together with a milder but detectable shift in Adjacent tissue. Dynamic-network proxies further indicate a more compact and perturbed core in Tumor samples, whereas Adjacent tissue appears more diffuse and weakly structured.


Overall, GSE69914 constitutes a technically clean dataset with well-structured methylation patterns and biologically interpretable deviations. The results indicate that tumour-related alterations are primarily localized at specific CpG loci rather than reflected in global methylome reorganization. Within this framework, the Adjacent tissue exhibits subtle yet measurable deviations from the Normal baseline, consistent with previously described field-effect patterns in breast cancer epigenomics.

3.3.2 Dataset GSE225845

Dataset overview GSE225845 is a breast tissue DNA methylation cohort profiled using the Illumina Infinium MethylationEPIC BeadChip (850K), providing genome-wide coverage of more than 8.5×10^5 CpG loci. The dataset is part of the

Table 3.2: Overview of the GSE225845 dataset after initial cohort curation.

CpGs	Samples	Normal	Adjacent	Tumor
750,426	477	113	140	224

**Figure 3.6:** Group-wise mean β -value density in GSE225845.

NCI-Maryland Breast Cancer Cohort and includes samples spanning the Normal–Adjacent–Tumor spectrum that motivates the biological framework of this thesis.

An overview of the sample composition and CpG coverage after cohort harmonization is reported in Table 3.2, including the distribution across tissue classes.

The dataset further provides detailed demographic and clinical metadata (e.g., age at surgery, race, sex, and tissue descriptors), enabling controlled downstream analyses when required.

Global methylation distributions The group-wise mean β -value densities (Figure 3.6) display the expected bimodal configuration of EPIC arrays. As observed in GSE69914, the three tissue groups show a high degree of overlap across the entire β range.

Normal and Adjacent samples exhibit nearly indistinguishable density profiles. Tumor samples present only minimal deviations, characterised by a slight increase in density at low β values and a marginal attenuation of the high- β peak. At the whole-methylome level, no marked global redistribution of methylation levels is evident.

CpG-level instability and recurrent outliers The heatmap of the top outlier CpG loci (Figure 3.7a) indicates that instability within this subset is predominantly directional. Recurrent outlier events are largely associated with very low β values, consistent with focal hypomethylation affecting a restricted set of CpG sites. The




Figure 3.7: Heatmap of the top outlier CpGs (left) and corresponding $\Delta\beta$ distributions (right), illustrating CpG-level instability in GSE225845.

perturbation pattern therefore appears concentrated rather than diffuse, suggesting localized instability rather than a genome-wide redistribution.

The corresponding $\Delta\beta$ distributions comparing Tumor and Adjacent tissues against Normal (Figure 3.7b) are sharply centered around zero, indicating that most loci remain stable. However, both comparisons exhibit a heavier negative tail relative to the positive side. Tumor samples display the most extended negative tail, consistent with stronger hypomethylation shifts, whereas Adjacent tissue shows a milder but still detectable asymmetry toward $\Delta\beta < 0$.

Variance and correlation structure The CpG-wise variance distributions (Figure 3.8a) indicate clear differences in variability across tissue groups. Normal samples exhibit highly concentrated variance values near zero, whereas Tumor samples display a substantially longer right tail, reflecting an increased proportion of CpG loci with elevated variability. Adjacent samples show a broader distribution than Normal tissues, with partial overlap with both Normal and Tumor profiles, but without forming a strictly intermediate pattern.

The sample correlation heatmap (Figure 3.8b) shows uniformly positive correlations across the dataset. Most pairwise values lie within a narrow high-correlation range, and no distinct block structures separating tissue classes are observed. This pattern indicates preservation of global methylation structure, with group differences emerging primarily from locus-specific variability rather than from large-scale shifts in overall methylome organization.

Low-dimensional embeddings Dimensionality reduction was applied using PCA, t-SNE, and UMAP to visualise global similarities among samples (Figure 3.9).




Figure 3.8: Variance and correlation structure across Normal, Adjacent, and Tumor samples in GSE225845.




Figure 3.9: Low-dimensional embeddings of samples in GSE225845.

In PCA space, the three tissue groups largely overlap. Tumor samples show a slightly broader dispersion, particularly along the second principal component, but no clear linear separation is observed.

Nonlinear embeddings reveal additional structure. In t-SNE space, Tumor samples occupy a more extended region and appear more dispersed, whereas Normal and Adjacent samples remain partially intermixed. Several Tumor samples are positioned at the periphery of the embedding, consistent with increased heterogeneity.

The UMAP projection exhibits a similar configuration. Tumor samples span a wider area and extend toward higher values of the second UMAP axis, while Normal and Adjacent samples form relatively compact yet overlapping regions. Trustworthiness scores range from 0.881 to 0.939 across embeddings, indicating good preservation of local neighbourhood structure, particularly in the nonlinear projections. Despite differences in spread, none of the methods yields sharply separated or isolated clusters.




Figure 3.10: Dynamic-network proxy visualizations in GSE225845.

Genome annotation coverage Consistent with the architecture of the EPIC platform, the genomic distribution of annotated CpG contexts is markedly unbalanced. The largest fraction of probes maps to open-sea or non-island regions, which constitute the dominant category. CpG islands represent the second most abundant class, followed by north and south shores in comparable proportions, whereas north and south shelves account for the smallest fractions.

This annotation profile aligns with the expected genomic composition of the EPIC array and supports correct manifest integration.

Dynamic-network proxies The degree–rank distribution (Figure 3.10a) exhibits a heavy-tailed profile, with a limited number of genes associated with many CpGs and a long tail of low-degree nodes. Such right-skewed degree structures are characteristic of heterogeneous biological networks [36].

In the Tumor vs Normal layout (Figure 3.10b), nodes aggregate into a dense and highly compact central cluster, with only a small number of isolated peripheral points. This geometry indicates that the largest Tumor–Normal $\Delta\beta$ deviations are concentrated within a cohesive subset of CpG–gene units rather than broadly distributed across the network.

In contrast, the Adjacent vs Normal layout (Figure 3.10c) displays a markedly different configuration. While a few small central aggregates are visible, most nodes are arranged along a wide annular structure, forming a pronounced ring-like embedding. This organisation reflects weaker and more spatially dispersed deviations, without the central concentration observed in the Tumor comparison.

Altogether, the proxy visualisations indicate that Tumor–Normal contrasts generate a compact and locally coherent network signal, whereas Adjacent–Normal differences remain diffuse and geometrically fragmented.

Overall interpretation The exploratory analysis of GSE225845 indicates that, at the whole-methylome level, the three tissue states share highly similar global

Table 3.3: Overview of the GSE287331 dataset after initial cohort curation.

CpGs	Samples	Normal	Adjacent	Tumor	OQ	CUB
866,552	446	182	60	69	67	68

methylation profiles. Group-wise mean β -value densities are nearly superimposed, and both correlation structure and linear embeddings show substantial overlap. These observations indicate preservation of large-scale methylome organisation across Normal, Adjacent, and Tumor tissues, without evidence of pronounced genome-wide redistribution of methylation levels.

Locus-focused analyses reveal more structured differences. Among the 200 CpGs with the highest outlier burden, instability is predominantly directional and characterised by focal hypomethylation. Variance distributions further show increased dispersion in Tumor samples, while Adjacent tissues exhibit broader variability than Normal without forming a strictly intermediate pattern.

Nonlinear embeddings (t-SNE and UMAP) emphasise differences in spatial dispersion rather than discrete cluster separation. Normal and Adjacent samples occupy relatively compact regions, whereas Tumor samples extend over a wider area, reflecting increased heterogeneity. Dynamic-network proxies show that Tumor–Normal contrasts generate a dense central cluster of perturbed CpG–gene units, while Adjacent–Normal differences are distributed along a broader, ring-like structure without strong central concentration.

Despite the larger sample size of this cohort, which provides stable estimates of global methylation structure, the separation between Normal and Adjacent tissues remains modest. Differences between these two groups are therefore primarily focal and do not manifest as large-scale structural shifts.

Overall, GSE225845 represents a globally homogeneous dataset in which tissue-state differences become apparent only when examined through locus-specific perturbations, variability patterns, and small-scale network organisation.

3.3.3 Dataset GSE287331

Dataset overview GSE287331 is a breast tissue DNA methylation cohort profiled using the Illumina Infinium MethylationEPIC v1.0 BeadChip. The dataset is organised along a tumor proximity axis (TPxA) spanning multiple anatomical and pathological contexts.

Five tissue categories are represented: healthy donated breast (HDB), contralateral unaffected breast (CUB), ipsilateral opposite quadrant (OQ), adjacent normal tissue (AN), and tumor (TU). An overview of the full cohort composition is reported in Table 3.3.




Figure 3.11: Schematic representation of the five tissue categories collected along the tumor proximity axis (TPxA), reproduced from [37].

For cross-dataset consistency, only three biologically aligned classes are retained for downstream analyses: HDB (treated as Normal baseline), AN (Adjacent), and TU (Tumor). OQ and CUB samples are excluded, as they represent intermediate benign tissues along the proximity axis and are not directly comparable with the three-class framework adopted in the other cohorts.

Global methylation distributions The group-wise mean β -value density curves (Figure 3.12) display the expected bimodal configuration of Illumina methylation arrays. A dominant peak is observed at high methylation levels, accompanied by a smaller peak consistent with standard EPIC methylation profiles [38].

Although the overall shapes remain comparable across tissue groups, a clear ordering is visible at the high- β peak. Normal samples exhibit the highest density, Adjacent samples show a slightly attenuated peak, and Tumor samples present the lowest amplitude together with a broader distribution extending toward intermediate β values. This progressive reduction in the fully methylated peak is consistent with increasing methylation variability along the Normal–Adjacent–Tumor axis.

CpG-level instability and recurrent outliers The heatmap of the top outlier CpG loci (Figure 3.13a) shows that epigenetic instability is not uniformly distributed. Alterations concentrate at specific CpG sites and within specific samples, forming vertical and horizontal bands rather than a diffuse genome-wide pattern. Both directions of deviation are observed, with focal hypermethylation and focal hypomethylation present within the selected loci [25].

The corresponding $\Delta\beta$ distributions (Figure 3.13b) highlight distinct profiles for the two contrasts. The Tumor–Normal curve is broader and displays a higher density of small negative shifts, indicating widespread but moderate hypomethylation in Tumor samples. In contrast, the Adjacent–Normal distribution is more sharply centered around zero but exhibits a comparatively more pronounced negative tail,





Figure 3.12: Group-wise mean β -value density in GSE287331.



(a) Top outlier CpG loci.

(b) $\Delta\beta$ distributions (Tumor/Adjacent – Normal).

Figure 3.13: Heatmap of the top outlier CpGs (left) and corresponding $\Delta\beta$ distributions (right), illustrating CpG-level instability in GSE287331.

reflecting fewer yet larger hypomethylated deviations. Overall, hypomethylation represents the dominant direction of change in both comparisons.

Variance and correlation structure The CpG-wise variance density curves (Figure 3.14a) show that Normal and Adjacent samples are nearly superimposed, with both groups exhibiting extremely low and tightly concentrated variance values. Tumor samples display a slightly broader right tail, indicating modestly increased genome-wide variability; however, the overall distribution remains sharply peaked near zero.

At the level of CpG-wise variance, Adjacent tissue remains closely aligned with the Normal baseline, whereas Tumor samples show only a mild elevation in variability.

A complementary perspective is provided by the sample correlation heatmap




Figure 3.14: Variance and correlation structure across Normal, Adjacent, and Tumor samples in GSE287331.

(Figure 3.14b). Correlations are uniformly high across the cohort, consistent with the predominance of stable CpG loci in genome-wide methylation data [22]. Although tissue labels are not overlaid on the matrix, localized blocks of higher similarity are visible, indicating structured relationships within subsets of samples rather than abrupt global separation.

Low-dimensional embeddings Dimensionality reduction using PCA, t-SNE, and UMAP was applied to visualise the global organisation of methylation profiles (Figure 3.15).

Across all three embeddings, a consistent and well-defined structure is observed. Normal and Adjacent samples form two clearly separated clusters. The separation is visible in PCA space, becomes more pronounced in t-SNE, and appears as fully detached manifolds in the UMAP projection.

Tumor samples occupy a distinct and more dispersed region of the embedding space, reflecting increased epigenetic heterogeneity. Trustworthiness scores are uniformly high (0.961–0.980), indicating excellent preservation of local neighbourhood structure in all low-dimensional projections. No substantial overlap between Tumor and the other two groups is observed in any of the projections, indicating strong intrinsic group structure in this cohort.

Genome annotation coverage The genomic distribution of annotated CpG contexts is dominated by Open Sea regions, followed by CpG Islands, then Shores, with Shelves representing the smallest fractions.

This ordering is consistent with the established architecture of EPIC arrays [31, 32].




Figure 3.15: Low-dimensional embeddings of GSE287331 samples.




Figure 3.16: Dynamic-network proxy visualizations for GSE287331.

Dynamic-network proxies The degree–rank curve (Figure 3.16a) shows a heavy-tailed distribution of CpG counts per gene, consistent with a hub–periphery architecture typical of biological networks.

Both $\Delta\beta$ -based network layouts — Tumor vs Normal and Adjacent vs Normal (Figure 3.16b, Figure 3.16c) — share a similar overall structure: a dense and compact central block of nodes connected by short-range edges, surrounded by a ring of peripheral, weakly connected nodes. This geometry reflects the underlying degree distribution and indicates that most CpG–gene units contribute only minimally to group-level differences.

Despite this shared organisation, the Tumor vs Normal comparison exhibits a more pronounced central cluster with larger $\Delta\beta$ magnitudes, suggesting that a restricted subset of CpG–gene units undergoes locally coordinated perturbations. In contrast, the Adjacent vs Normal layout preserves the same global shape but shows weaker deviations, indicating only mild coordination of methylation changes.

Overall interpretation The exploratory analysis of GSE287331 indicates that, at the whole-methylome level, the three tissue states retain a broadly preserved

global methylation architecture. Group-wise mean β -value densities follow the typical bimodal EPIC profile with only a subtle gradient (Normal > Adjacent > Tumor), and both CpG-wise variance distributions and sample correlation structure show high overall similarity across tissues, with only mild variability increases in Tumor samples.

In contrast to the previous cohorts, low-dimensional embeddings reveal a clear and consistent separation of all three groups. Normal and Adjacent samples form distinct clusters in PCA, with the separation further strengthened in t-SNE and UMAP, where the groups appear as detached manifolds. Tumor samples occupy a separate and more diffuse region, reflecting increased epigenetic heterogeneity. These projections indicate that, despite similar first- and second-order global summaries, multivariate structure in this dataset captures strong group-level organisation.

Locus-specific analyses reveal focused disruptions. Tumor samples exhibit widespread mild hypomethylation, whereas Adjacent tissue shows fewer but larger-magnitude deviations. Dynamic-network proxies further indicate a compact and perturbed central core in the Tumor comparison, while Adjacent–Normal differences display weaker coordination, consistent with early-stage methylation alterations.

Overall, GSE287331 emerges as a structurally coherent and biologically informative dataset in which Normal–Adjacent–Tumor separation is markedly more pronounced than in the other cohorts, while global methylation architecture remains broadly conserved.

3.4 Inter Dataset Exploration and Comparison

The objective of this section is not to merge the datasets—an approach that would be biologically and technically inappropriate due to differences in array platforms (HM450 vs. EPIC), preprocessing pipelines—but rather to compare their internal behaviours. Such a cross-dataset perspective enables the assessment of whether the early epigenetic alterations identified in the intra-dataset analyses are *reproducible across independent cohorts*, thereby providing support for the central biological premise of this thesis: the detection of early methylation drift in histologically normal tissues and the evaluation of its potential contribution to tumour initiation.

3.4.1 Analytical Framework

Across all three datasets, the intra-dataset analyses revealed three highly consistent phenomena:

- Normal and Adjacent tissues are globally similar, with nearly superimposed β -value distributions and comparable CpG-wise variance profiles.

Table 3.4: Cross-dataset comparison of Normal (N) vs Adjacent (A) contrasts using global and instability metrics.

Dataset	Mean $ \Delta\beta $ (A–N)	Var Ratio (A/N)	Outlier Ratio (A/N)	Silhouette (A/N)
GSE69914	0.009	0.988	1.353	0.003
GSE225845	0.015	0.972	2.790	0.042
GSE287331	0.031	0.930	13.329	0.344

- Tumour samples show increased heterogeneity and a clear bias toward hypomethylation, although the magnitude of this shift varies across datasets.
- Adjacent tissues exhibit early methylation drift, detectable through $\Delta\beta$ distributions and outlier profiles even when global metrics remain highly similar to Normal.

These patterns are consistent with the expected biological progression from Normal to Tumour, in which epigenetic deregulation emerges gradually and affects normal-appearing tissue surrounding the tumour. The inter-dataset analysis therefore evaluates the *directional consistency* of these effects rather than aiming at strict quantitative comparability.

3.4.2 Cross-Dataset Comparative Metric

To quantify the epigenetic deviation of Adjacent tissue from the Normal methylome across the three cohorts, a set of global and instability-oriented metrics was computed independently within each dataset for the Adjacent versus Normal contrast.

All metrics reported in Table 3.4 were computed using the complete CpG set available in each dataset, rather than the cross-platform intersection. The resulting values therefore reflect cohort-specific dimensionality and internal variance structure.

Collectively, these metrics indicate the presence of early epigenetic drift in histologically non-tumour tissue. Across cohorts, increasing mean $|\Delta\beta|$, progressive variance imbalance, elevated outlier ratios, and improved silhouette separation consistently support a directional shift from Normal to Adjacent states, in agreement with the field-defect framework described by Teschendorff et [27] and reinforced in other tissue contexts.

Mean absolute methylation difference For each CpG, the absolute mean difference between Adjacent and Normal was defined as

$$|\Delta\beta_i| = \left| \beta_i^{(A)} - \beta_i^{(N)} \right|. \quad (3.2)$$

The cross-cohort pattern reveals progressively larger locus-specific shifts from GSE69914 to GSE225845 and GSE287331. Although the absolute magnitude of the effect remains modest, it is consistent in direction across cohorts. This monotonic increase reflects a strengthening early deviation in Adjacent tissues. Comparable locus-wise methylation drifts in histologically normal tissue have been reported in independent systems, such as colonic mucosa adjacent to adenomas [39], supporting the interpretation of these alterations as early manifestations of field cancerization.

Global variance ratio For each sample s , the global methylation variance was computed across all CpG sites: $v_s = \text{Var}(\beta_{s,1}, \beta_{s,2}, \dots, \beta_{s,m})$, where m denotes the number of CpGs. Group-wise mean variances were then defined as

$$\bar{v}^{(N)} = \frac{1}{|N|} \sum_{s \in N} v_s, \quad \bar{v}^{(A)} = \frac{1}{|A|} \sum_{s \in A} v_s, \quad (3.3)$$

and the reported metric corresponds to their ratio:

$$\frac{\text{Var}(A)}{\text{Var}(N)} = \frac{\bar{v}^{(A)}}{\bar{v}^{(N)}}. \quad (3.4)$$

Across cohorts, this ratio remained consistently close to unity, indicating that Adjacent tissue does not exhibit diffuse genome-wide instability. Such behaviour suggests that early pre-neoplastic alterations are predominantly *focal* rather than global, with only specific CpG sites displaying abnormal dispersion while overall variance remains largely preserved [39].

This interpretation is consistent with contemporary models of early tumourigenesis, in which subtle and locally restricted perturbations accumulate prior to large-scale chromatin deregulation and overt neoplastic transformation [40].

Outlier burden ratio For each sample, the outlier burden is defined as the number of CpG sites whose methylation deviates from the Normal reference distribution by more than three times the Normal interquartile range:

$$\text{outlier}_s = \# \left\{ i : \left| \beta_{i,s} - \tilde{\beta}_i^{(N)} \right| > 3 \cdot IQR_i^{(N)} \right\}, \quad (3.5)$$

where $\tilde{\beta}_i^{(N)}$ and $IQR_i^{(N)}$ denote the median and interquartile range of CpG i across Normal samples.

The reported A/N ratio corresponds to the mean outlier burden in Adjacent tissue divided by the mean burden in Normal tissue. Across cohorts, this metric exhibits the most pronounced cross-dataset gradient, with progressively stronger enrichment of outlier events in Adjacent tissue. Such behaviour mirrors evidence that increased local methylation variability in histologically normal tissue represents a sensitive marker of early instability [39], and provides quantitative support for the presence of marked field effects in GSE287331.

Silhouette score For each sample s , let $a(s)$ denote the mean distance from s to all other samples within the same group (cohesion), and let $b(s)$ denote the smallest mean distance from s to samples in the alternative group (separation). The silhouette coefficient for sample s is defined as

$$\text{sil}(s) = \frac{b(s) - a(s)}{\max\{a(s), b(s)\}}, \quad (3.6)$$

and the reported score corresponds to the mean silhouette computed across all Normal and Adjacent samples.

Higher values indicate clearer multivariate separation between Normal and Adjacent states, whereas values near zero reflect substantial overlap. Across cohorts, the silhouette analysis reveals limited separation in GSE69914 and GSE225845, consistent with minimal global displacement in multivariate space. In contrast, GSE287331 exhibits partial separation, indicating that the epigenetic landscape of Adjacent tissue is measurably shifted relative to Normal. This behaviour aligns with mechanistic models in which epigenetic dysregulation constitutes one of the earliest events in tumour initiation [40], preceding morphological transformation.

Collectively, the set of metrics delineates a consistent Normal–Adjacent gradient across datasets. Early epigenetic drift is detectable in all cohorts, while its magnitude, focality, and multivariate visibility progressively intensify from GSE69914 to GSE287331. Importantly, the combination of stable global variance with pronounced outlier enrichment supports a model in which early pre-neoplastic changes arise through *localised* perturbations, a hallmark of field cancerization [39, 27].

3.4.3 Visual Comparative Analysis

To complement the quantitative summary reported in Table 3.4, this section provides a visual characterisation of the Normal–Adjacent contrast across the three cohorts. All analyses are performed on the three-way intersection of 326,330 CpG sites, ensuring maximal comparability across platforms (HM450 and EPIC) and preprocessing pipelines [31]. The objective is not to achieve numerical concordance at the single-CpG level—a notoriously difficult task due to platform differences, probe-design effects, and cohort-specific processing [32, 41]—but rather to evaluate whether the *shape*, *direction*, and *global behaviour* of early methylation drift remain coherent across studies.

Global $\Delta\beta$ distributions The overlaid $\Delta\beta$ (Adjacent–Normal) density curves across the three cohorts are shown in Figure 3.17. All distributions are sharply centred around zero, indicating that the N–A methylation drift remains globally subtle in each dataset, consistent with the focal and low-amplitude nature of early




Figure 3.17: Overlay of $\Delta\beta$ (Adjacent–Normal) density curves across the three datasets, computed on the 326,330 CpG sites shared by all cohorts.

epigenetic alterations described in pre-neoplastic tissues [27, 42]. Despite differences in dispersion—reflecting platform-dependent noise, probe-type composition, and study-specific variability [29, 32]—the three curves exhibit a highly similar shape. This pattern indicates that early epigenetic deviations in Adjacent tissue, although small in magnitude, exhibit reproducible global distributional features across independent cohorts, rather than exact concordance at the single-CpG level.

Pairwise $\Delta\beta$ concordance across datasets To assess whether locus-specific A–N shifts replicate across cohorts, pairwise scatterplots are shown for all dataset pairs (Figure 3.18a, Figure 3.18b, Figure 3.18c). Each point corresponds to a CpG in the three-way intersection; axes represent the mean $\Delta\beta$ (A–N) computed independently within each dataset. Across all pairs, Spearman correlations remain weak ($\rho = 0.05$ – 0.29), consistent with the well-established difficulty of achieving single-CpG reproducibility across independent methylation studies [31, 41, 43].

The comparison between GSE69914 and GSE225845 exhibits mild monotonic agreement, whereas pairs involving GSE287331 show near-zero correlation. These results indicate that while the *global* N–A drift remains consistent across cohorts (as demonstrated by the density curves), *fine-scale*, *CpG-specific* effects are strongly influenced by platform architecture, batch structure, and study-specific variability [32, 41].

t-SNE embeddings on the shared CpG intersection To investigate the multivariate structure of the N–A contrast across cohorts, t-SNE embeddings are computed on M-values restricted to the shared CpGs. Two complementary visualisations of the embedding are provided in Figure 3.19a (coloured by dataset) and Figure 3.19b (coloured by tissue state).

When coloured by dataset, the embedding exhibits clear clustering by study,




Figure 3.18: Pairwise $\Delta\beta$ (Adjacent–Normal) scatterplots across the three methylation datasets, computed on the 326,330 shared CpG sites. Spearman ρ values are reported in each panel.




Figure 3.19: t-SNE embeddings computed on the 326,330 CpGs shared across GSE69914, GSE225845, and GSE287331.

reflecting the dominant influence of batch and platform effects in high-dimensional methylation data [41, 43]. This pattern indicates that the global methylation structure is largely driven by inter-cohort variability, supporting the decision not to merge datasets at the raw feature level.

When coloured by tissue state (Normal vs. Adjacent), no global separation across cohorts emerges, as the embedding remains primarily structured by dataset identity. However, within individual study-specific clusters, a state-dependent displacement becomes visible. This separation is particularly evident in GSE287331, where Normal and Adjacent samples form more clearly distinct substructures. This behaviour is consistent with early, localised methylation drift, mirroring the quantitative metrics reported in Table 3.4.

Overall, the visual analyses corroborate the quantitative metrics: early epigenetic

drift is detectable in all cohorts, but its magnitude and visibility vary substantially, with GSE287331 exhibiting the clearest field-defect signature. The combination of globally subtle yet directionally consistent $\Delta\beta$ patterns, weak CpG-level concordance, and dataset-specific multivariate structure reinforces the view that the Normal→Adjacent transition reflects genuine early epigenetic deregulation rather than dataset-specific artefacts.

3.5 Exploratory Findings and Pre-Processing Implications

The exploratory analyses presented in this chapter reveal a consistent picture across all three cohorts: the Normal–Adjacent transition does not manifest as a global reorganisation of the methylome, but as a subtle, focal drift that becomes detectable only through locus-specific perturbations and multivariate displacements. This characteristic makes it particularly sensitive to technical sources of variability — probe-design bias, batch effects, and platform differences — which operate at a comparable or larger scale than the biological signal of interest.

This sensitivity has direct methodological consequences. Raw methylation data, as distributed via GEO, cannot be used directly for downstream modelling without first addressing these technical confounders: doing so would risk attributing platform-specific artefacts to genuine epigenetic alterations, or conversely, suppressing the subtle Normal–Adjacent differences under overly aggressive normalisation. The challenge is therefore not simply to clean the data, but to do so in a way that preserves the biological structure identified here.

Chapter 4 addresses this challenge by developing a structured, platform-aware preprocessing pipeline. Each step is motivated by the specific limitations identified in the present chapter.

Chapter 4

Data preprocessing

4.1 Preprocessing Rationale and Technical Constraints

DNA methylation arrays provide high-dimensional, genome-wide measurements in which biological variability and platform-specific technical artefacts are inevitably superimposed. The exploratory analyses of Chapter 3 revealed a consistent picture across all three cohorts: Normal and Adjacent tissues exhibit nearly identical global methylation distributions, overlapping variance profiles, and indistinguishable sample-level correlation structure. Meaningful differences between tissue states emerge exclusively at focal CpG loci—as quantified by locus-specific $\Delta\beta$ distributions and outlier burden ratios—and remain subtle even at that scale (mean $|\Delta\beta|$ ranging from 0.009 in GSE69914 to 0.031 in GSE287331). This characteristic makes the Normal–Adjacent signal particularly vulnerable to technical confounders, which operate at a comparable or larger scale than the biological effect of interest. Preprocessing must therefore be precise: aggressive enough to remove technical noise, but conservative enough to preserve the subtle epigenetic drift that motivates this thesis.

Three main classes of technical issues must be systematically addressed before any statistical modelling can proceed. First, **probe-design bias** intrinsic to the Infinium platform arises from the coexistence of two chemically distinct probe types (Type I and Type II), which produce inherently different β -value distributions and can distort cross-sample comparability if not corrected [44, 45]. This issue is not uniform across datasets: as will be shown in Section 4.3, GSE69914 and GSE225845 were released with normalization already applied by the original authors, whereas GSE287331 shows clear diagnostic evidence of missing probe-type correction, requiring an additional preprocessing decision with non-trivial consequences for downstream group separation. Second, a subset of **unreliable or biologically**

confounded probes—including cross-reactive loci, SNP-affected sites, non-CpG targets, and sex-chromosome probes—introduces systematic measurement error that can generate spurious associations or mask genuine methylation differences [46, 47]. Third, the **statistical scale of β -values** poses inferential challenges: the bounded $[0,1]$ support and strong mean-dependent heteroscedasticity of β -values violate the assumptions of standard parametric models, motivating transformation to a more statistically well-behaved representation prior to modelling [44].

This chapter addresses these three issues through a structured, platform-aware preprocessing pipeline applied consistently across all cohorts. Section 4.2 describes the general framework and the rationale for each step. Section 4.3 documents the dataset-specific application, focusing on the decisions that deviate from the standard workflow—most notably the handling of probe-type bias in GSE287331, where the tension between technical correction and preservation of biological structure requires explicit justification. Section 4.4 summarises the effect of preprocessing on CpG dimensionality and cross-dataset overlap, providing the basis for the feature selection strategy developed in Chapter 5.

4.2 General Preprocessing Framework

This section describes the general preprocessing framework applied to all three methylation datasets. The pipeline comprises four sequential steps, each addressing a distinct class of technical issue identified in the overview above. Two of these steps—data integrity verification and Infinium probe-design bias correction—require dataset-specific decisions whose rationale and consequences are discussed in Section 4.3. The remaining two steps—technical probe filtering and β -to- M transformation—are applied uniformly across all cohorts following the same protocol.

4.2.1 Structural Integrity and Cohort Harmonization

Prior to any preprocessing operation, each dataset undergoes a structural verification stage to ensure the consistency and completeness of the input data. This step checks for correct alignment between the methylation matrix and its associated phenotype table, verifies sample identifier uniqueness, and confirms the absence of missing metadata fields. Only samples for which both a complete methylation profile and a valid tissue label are available are retained for downstream analysis.

Probe-level and sample-level missingness in the β -value matrix is also assessed at this stage. For two of the three datasets (GSE69914 and GSE225845), the released matrices are fully observed and require no imputation. For GSE287331, a non-negligible fraction of CpG sites exhibits missing values across samples, necessitating a dedicated filtering and imputation strategy consistent with current

recommendations for EPIC-array quality control [48, 49]. The dataset-specific treatment of missingness is detailed in Section 4.3.3.

Finally, cohort harmonization is applied where necessary to align tissue labels to the three-class framework (Normal, Adjacent, Tumor) adopted uniformly across datasets. Samples belonging to tissue categories outside this framework are excluded at this stage to ensure cross-dataset comparability.

4.2.2 Probe-Type Bias Diagnostics and Correction

A well-documented source of systematic bias in Illumina Infinium methylation arrays is the coexistence of two chemically distinct probe designs. Type I probes use two bead types per CpG (one for the methylated channel, one for the unmethylated channel), whereas Type II probes use a single bead type and rely on single-base extension with two-color detection. This design difference produces inherently distinct β -value distributions for the two probe types: Type I probes tend to yield more extreme values near 0 and 1, while Type II probes exhibit a broader, intermediate distribution. If uncorrected, this imbalance propagates into downstream statistical analyses and can generate probe-type-specific artefacts in differential methylation testing [22, 29].

The canonical approach to mitigating this bias is Beta-Mixture Quantile normalisation (BMIQ), introduced by Teschendorff et al. [22]. BMIQ explicitly models probe-type bias at the β -value level through a state-aware distributional alignment procedure. For each sample independently, a three-component Beta mixture model is first fitted to the β -value distribution of Type I probes, representing unmethylated, partially methylated, and fully methylated states. These fitted components define the reference state-specific distributions. Type II probes are then probabilistically assigned to one of the three methylation states based on their β -values, and within each state a quantile-mapping step is applied to align the Type II distribution to the corresponding Type I reference distribution. Because the procedure is performed independently for each sample, BMIQ does not introduce cross-sample information or global harmonisation across tissue classes; it strictly corrects within-sample probe-type bias.

For each dataset, a systematic diagnostic is performed prior to any correction step [22, 50]: probes are stratified by design type using the official Illumina manifest, and two complementary assessments are carried out—(i) a density comparison of β -value distributions for Type I and Type II probes, and (ii) a quantile–quantile plot of Type II versus Type I quantiles, with the median absolute deviation (MAD) of the empirical Q–Q curve from the identity line as a scalar diagnostic metric. Datasets for which the original authors applied BMIQ or an equivalent normalisation are expected to exhibit near-overlapping distributions and a MAD close to zero; datasets lacking probe-type correction will display a pronounced distributional

mismatch and elevated MAD.

The outcome of this diagnostic, and the corrective action taken, differs across cohorts and is reported in full in Section 4.3. Critically, the decision of whether to apply BMIQ is not made mechanically: as discussed in Section 4.3.3, BMIQ correction can attenuate genuine biological structure when group-level differences are expressed as broad distributional shifts, a phenomenon documented in systematic evaluations of EPIC-array normalisation methods [16, 51, 52]. The choice between correcting and preserving the raw data is therefore made on the basis of empirical evidence from the diagnostic plots and quantitative assessment of group separability before and after correction.

4.2.3 Technical Probe Filtering

After addressing probe-design bias, unreliable CpG loci are removed through a systematic filtering procedure based on curated external annotation resources. The objective is to exclude probes whose measured β -values may reflect technical measurement error rather than genuine DNA methylation, thereby reducing noise and improving the reliability of downstream differential analyses. Four categories of technically problematic probes are systematically excluded.

Cross-reactive and multi-mapping probes. Probes that hybridise to multiple genomic loci produce β -values that cannot be unambiguously attributed to a single CpG site. These are identified using the catalogues of Naeem et al. [53], Chen et al. [54], Pidsley et al. [32], Zhou et al. [31], and McCartney et al. [55]. Full descriptions of each resource are provided in Appendix A.

SNP-affected probes. Probes whose interrogated CpG dinucleotide, single-base extension site, or probe body overlaps a common genetic variant may measure genotype rather than methylation status, producing population-stratified artefacts in group comparisons [31, 32]. These are excluded using the `MASK_snp5` and `MASK_extBase` fields from the Zhou annotation [31], supplemented by variant-aware lists from Naeem et al. [53].

Non-CpG probes. Probes targeting non-CpG cytosine contexts (identified by the `ch` prefix in Illumina probe identifiers) are excluded, as their interpretation differs from canonical CpG methylation and is outside the scope of this analysis [31, 32].

Sex-chromosome probes. Probes located on chromosomes X and Y are removed to prevent sex-driven confounding effects. Sex-chromosome CpGs exhibit

substantially different methylation profiles between males and females, and including them in joint normalisation has been shown to introduce artificial sex-related bias into autosomal probes [56]. Since the datasets analysed here contain samples from both sexes and sex-specific methylation is not the object of investigation, removal therefore represents a conservative strategy to prevent confounding without compromising autosomal inference [29].

The filtering resources are applied under a union criterion: a probe is removed if it appears in at least one curated list. Platform-specific manifests (HumanMethylation450 v1.2 for GSE69914 [57]; MethylationEPIC v1.0 B5 for GSE225845 and GSE287331 [58]) are used for manifest-based validation, chromosome annotation, and non-CpG probe identification. The number of probes flagged by each resource, and the redundancy of flags across resources, are reported per-dataset in Section 4.3.

4.2.4 Statistical Transformation: β -to- M Values

The final preprocessing step transforms the filtered β -value matrix into M -values prior to statistical modelling. As defined in Equation (3.1), β -values are bounded proportions in $[0,1]$ derived from methylated and unmethylated probe intensities. While β -values are intuitive and directly interpretable as fractional methylation, their bounded support induces strong mean-dependent heteroscedasticity: variance is inflated near intermediate values ($\beta \approx 0.5$) and compressed near the boundaries, violating the homoscedasticity assumptions underlying standard linear modelling frameworks [21].

The logit transformation to M -values,

$$M = \log_2 \left(\frac{\beta + \varepsilon}{1 - \beta + \varepsilon} \right), \quad \varepsilon = 10^{-6}, \quad (4.1)$$

maps $\beta \in (0,1)$ to an unbounded real-valued scale and substantially stabilises the mean–variance relationship [21]. As shown in their analysis, the relationship between β and M -values is a logit transformation yielding an approximately linear correspondence in the intermediate methylation range and nonlinear compression at the extremes. The small offset ε is introduced to handle boundary values numerically. The resulting M -value distributions are approximately symmetric and exhibit markedly flatter variance profiles across the full methylation range, making them the standard input for differential methylation analysis within linear modelling frameworks and related inferential procedures [44, 45, 59].

It is important to note that M -values are adopted exclusively for statistical modelling: β -values are retained in parallel for biological interpretation, effect-size quantification, and visualisation, following the dual-representation convention recommended by Du et al. [21] and widely adopted in the literature [59, 60].




Figure 4.1: Infinium Type I/II bias diagnostics in GSE69914.

The transformation is applied to the complete post-filtering matrix for each dataset. Distributional diagnostics confirming the expected variance-stabilising effect are reported in Section 4.3. This choice is consistent with the feature-selection and modelling procedures developed in the subsequent Chapter 5, which rely on variance-based ranking, linear statistics, and stability criteria that assume approximately homoscedastic behaviour.

4.3 Dataset-Specific Implementation

This section documents the dataset-specific application of the preprocessing pipeline described in Section 4.2. For each cohort, the outcome of structural verification, probe-type bias diagnostics, technical probe filtering, and β -to- M transformation is reported in detail. Although the general framework is applied consistently across all datasets, certain steps—most notably probe-type bias correction and handling of missing values—require cohort-specific decisions. These are explicitly justified within each subsection.

The results are presented separately for GSE69914 (Section 4.3.1), GSE225845 (Section 4.3.2), and GSE287331 (Section 4.3.3).

4.3.1 Dataset GSE69914

Structural Integrity and Cohort Harmonization The dataset comprises 397 samples and 485,512 CpGs at load time. All sample identifiers are unique, and the phenotype table contains complete metadata for the required fields. Alignment between the phenotype annotation and the β -value matrix is exact. The methylation matrix is fully observed, with no missing CpG values and no incomplete samples. No exclusion or imputation was required.




Figure 4.2: Technical filtering summary in GSE69914.

Table 4.1: CpG filtering summary for the GSE69914 dataset.

Initial CpGs	Naeem	Pidsley	Zhou	Chen	McCartney	Non-CpG	X/Y	Final CpGs
485,512	190,672	39,737	96,541	29,233	28,582	875	7,728	251,483

Probe-Type Bias Diagnostics and Correction Raw intensity data were reported to have been processed by the original authors using the `minfi` pipeline with BMIQ normalization [18]. Diagnostic evaluation confirms that probe-type bias is effectively mitigated in the released matrix.

The β -value density distributions of Type I and Type II probes exhibit substantial overlap, with no evidence of the characteristic multi-modal distortion observed in uncorrected data (Figure 4.1a). Quantile–quantile analysis further supports this result: the Type II versus Type I quantiles align closely to the identity line, with median absolute deviation $\text{median}|\Delta| = 0.162$, indicating balanced probe-type scaling (Figure 4.1b). Together, these diagnostics are consistent with effective BMIQ correction, in agreement with the normalization behaviour described in [22, 29, 50].

Technical Probe Filtering Intersection with external technical filtering resources resulted in substantial probe removal, as summarised in Table 4.1. The largest individual contributions derive from the Naeem and Zhou annotations, while the remaining resources flag smaller but non-negligible subsets. The redundancy structure across lists is shown in Figure 4.2b. Because these catalogues target partially distinct classes of technical artefacts—such as cross-reactivity, SNP interference, and repeat-associated probes—limited overlap across lists is expected. Most excluded CpGs are supported by a single resource, with progressively fewer loci flagged by multiple independent annotations. This pattern reflects complementary




Figure 4.3: Distributional comparison of β and M in Normal and Adjacent tissues in GSE69914.

rather than redundant filtering criteria, while the subset of probes flagged by several lists indicates concordance for loci exhibiting multiple technical vulnerabilities.

Annotation-based validation using the HumanMethylation450 v1.2 manifest [61] further excluded non-CpG probes and loci lacking reliable genomic annotation. Subsequent removal of sex-chromosome probes (X/Y), consistent with established preprocessing recommendations in mixed-sex cohorts [29, 56], ensured a strictly autosomal working set.

A detailed per-probe exclusion summary is available in the project repository at `removed_cpgs_all_filters_summary_gse69914.csv`.

Statistical Transformation: β -to- M Values Following transformation, the empirical distributions confirm the expected variance-stabilising behaviour of M -values (Figure 4.3). In β -space, both Normal and Adjacent tissues exhibit the characteristic bounded, U-shaped distribution with accumulation near 0 and 1. After logit transformation, the corresponding M -value distributions become substantially more symmetric and centered, while preserving the relative structure between tissue groups. The mean-SD relationships further illustrate the scale effect. In β -space, variance is strongly mean-dependent and inflates near intermediate methylation levels, producing a pronounced curvature in the mean-SD plots. In contrast, M -values display markedly flatter dispersion profiles across the full range, indicating substantial reduction of heteroscedasticity.

This behaviour is fully consistent with the theoretical and empirical findings of Du et al. [21], which motivate the use of M -values for inferential modelling.




Figure 4.4: Infinium Type I/II bias diagnostics in GSE225845.

4.3.2 Dataset GSE225845

Structural Integrity and Cohort Harmonization The dataset comprised 477 samples and 750,426 CpGs at load time, consistent with the dimensionality of the Illumina EPIC platform. The phenotype table contained 477 entries, and alignment between phenotype annotation and the β -value matrix was exact. No duplicated entries were detected, and all required metadata fields were present.

Probe-level and sample-level missingness assessment confirmed that the methylation matrix was fully observed. No missing CpG values and no incomplete samples were detected. Consequently, no imputation or early NaN filtering was required prior to probe-type bias diagnostics and technical filtering.

Probe-Type Bias Diagnostics and Correction Stratification by probe design yielded 119,760 Type I and 627,842 Type II probes. Raw intensity data were reported by the original authors to have been processed using the `minfi` pipeline with BMIQ normalization (v1.4) [19]. The β -value density distributions (Figure 4.4a) show substantial overlap between probe families. Type I probes retain the characteristic concentration near the unmethylated and fully methylated boundaries, while Type II probes display a broader distribution; however, no pronounced multimodal distortion or systematic separation is observed. Quantile–quantile analysis (Figure 4.4b) further supports this interpretation. Although minor deviations from the identity line ($y = x$) are visible, the overall alignment indicates largely balanced probe-type scaling. The median absolute deviation $\text{median}|\Delta| = 0.293$ suggests that probe-type differences are attenuated but not entirely eliminated, consistent with residual variation commonly observed after BMIQ normalization [22, 29, 50].

Overall, the diagnostic patterns are compatible with prior application of probe-type bias correction.




Figure 4.5: Technical filtering summary in GSE225845.

Table 4.2: CpG filtering summary for the GSE225845 dataset.

Initial CpGs	Naeem	Pidsley	Zhou	Chen	McCartney	Non-CpG	X/Y	Final CpGs
750,426	101,014	11,759	114,407	0	1,585	773	14,071	530,683

Technical Probe Filtering Intersection with external technical filtering resources resulted in substantial probe removal, as summarised in Table 4.2. The largest individual contributions derive from the Zhou and Naeem annotations, while the remaining resources flag smaller subsets. The redundancy structure across lists is shown in Figure 4.5b. A detailed per-probe exclusion summary is available in the project repository at `removed_cpgs_all_filters_summary_gse225845.csv`.

Statistical Transformation: β -to- M Values All filtered β -values were transformed to M -values using the logit relation defined in Equation (4.1). The empirical distributions (Figure 4.6) show the characteristic bounded, U-shaped profile in β -space for both Normal and Adjacent tissues. After transformation, the corresponding M -value distributions become substantially more symmetric and centered, while maintaining the relative structure between tissue classes.

The mean–SD relationships further illustrate the scale effect. In β -space, variance exhibits pronounced mean dependence, with inflation at intermediate methylation levels and compression near the boundaries. In contrast, M -values display markedly flatter dispersion profiles across the full range, indicating substantial reduction of heteroscedasticity.




Figure 4.6: Distributional comparison of β and M in Normal and Adjacent tissues in GSE225845.

4.3.3 Dataset GSE287331

Structural Integrity and Cohort Harmonization The dataset was imported from its Parquet representation, yielding an initial β -value matrix of 446 samples and 866,552 CpGs. The phenotype table contained the same number of entries, with all required metadata fields present and alignment between phenotype annotation and the β -value matrix exact.

Unlike GSE69914 and GSE225845, the methylation matrix was not fully observed. Probe-wise missingness is a well-known source of unreliability in Illumina methylation arrays: Islam *et al.* [48] excluded probes with missing β -values in more than 2% of samples, while Mansell *et al.* [49] applied a stricter 1% threshold for detection failure in their EPIC-array quality control pipeline. Consistent with these recommendations, all CpG sites missing in more than 1% of samples were removed, prioritising high-confidence measurements while minimising reliance on imputed values and leveraging the large dimensionality of the EPIC platform. After applying this criterion, the remaining sparsity was negligible: 58,561 NaN values, corresponding to 0.0187% of all entries. Given this extremely low level of sparsity, median-based imputation is both statistically robust and distribution-preserving; residual missing values were therefore imputed using the probe-wise median β -value across samples, a strategy that avoids introducing group-specific bias and is consistent with recent harmonisation workflows such as mLiftOver, where missing values are explicitly replaced by probe-wise median substitution to preserve biologically plausible levels [62].




Figure 4.7: Per-sample mean β distributions for the three datasets.




Figure 4.8: Infinium Type I/II bias diagnostics in GSE287331. Pre-BMIQ densities and Q-Q analysis indicate marked probe-type imbalance; post-BMIQ densities show partial alignment.

After harmonising phenotype labels and applying dataset-specific selection rules, only samples belonging to the three core tissue states of interest—Normal, Adjacent, and Tumour—were retained. The final working matrix consisted of 311 samples and 702,916 CpGs, with no residual missing values.

Probe-Type Bias Diagnostics and Correction Using the official EPIC v1.0 manifest, 115,705 Type I and 585,514 Type II probes were identified. Unlike GSE69914 and GSE225845, for which the original authors reported prior application of BMIQ normalisation, no probe-design bias correction was documented in the processing pipeline of GSE287331. This absence was first suggested by the per-sample mean β -value distributions (Figure 4.8): whereas GSE69914 and GSE225845 display a single broad peak, GSE287331 shows a narrower, right-shifted distribution characterised by two distinct modes in the high- β range—a pattern largely absent in the other cohorts and strongly indicative of uncorrected Type I/II probe-design bias.

Stratification of probes by design type confirmed this hypothesis. The β -value distributions of the two chemistries showed a marked mismatch (Figure 4.8a), with Type II probes displaying a strong right-shifted profile. The quantile-quantile




Figure 4.9: Technical filtering summary in GSE287331.

comparison confirmed a substantial systematic offset, with a median absolute deviation of 0.380 (Figure 4.8c), fully incompatible with BMIQ-corrected data. Inspection of the original publication [37] further confirmed the absence of any mention of BMIQ, SWAN, noob, or equivalent probe-type correction methods.

Given this evidence, the canonical BMIQ algorithm [22] was applied using the `watermelon` implementation [63], operating strictly on a per-sample basis: Type I probes define the reference mixture distribution within each sample, and Type II probes are rescaled accordingly, with no cross-sample or cross-group information introduced at any stage. Samples with insufficient numbers of probes per design type were left uncorrected to avoid unstable mixture estimation.

Post-correction diagnostics showed only a modest reduction in probe-type mismatch ($|\Delta| : 0.380 \rightarrow 0.370$), with the overall distributional discrepancy remaining substantial and the bimodality of per-sample mean β distributions persisting (Figure 4.8, Figure 4.8b). This limited harmonisation is consistent with documented caveats of quantile-based normalisation on EPIC arrays, which can artificially reduce biological variability, remove genuine group-level signal, or yield distorted distributions when differences are expressed as broad distributional shifts [51, 64, 52].

The impact on biological structure was assessed quantitatively via silhouette analysis: the average silhouette coefficient declined from $\text{sil}_{\text{pre}} = 0.2876$ to $\text{sil}_{\text{post}} = 0.2537$ after correction, indicating that BMIQ reduced the natural clusterability of the data and collapsed genuine group-level separation—particularly between Normal and Adjacent samples. For this reason, all downstream analyses are carried out on the non-BMIQ version of the dataset, which preserves sharper cluster boundaries and avoids normalisation-induced shrinkage of biological signal.

Table 4.3: CpG filtering summary for the GSE287331 dataset.

Initial CpGs	Naeem	Pidsley	Zhou	Chen	McCartney	Non-CpG	X/Y	Final CpGs
866,552	124,511	8,992	102,802	1,730	3,672	500	12,579	486,725

Technical Probe Filtering Intersection with external technical filtering resources resulted in substantial probe removal, as summarised in Table 4.3. The largest individual contributions derive from the Naeem and Zhou annotations, while the remaining resources flag smaller but non-negligible subsets (Figure 4.9a). Importantly, in the unfiltered EPIC matrix the Zhou annotation alone would have excluded 192,856 CpGs. This number substantially exceeds the net contribution observed at this stage, indicating that a considerable fraction of Zhou-flagged probes had already been removed during the earlier missingness-cleaning step. The missingness filter and the Zhou reliability masks therefore exhibit non-trivial overlap. This concordance supports the robustness of the preprocessing strategy: probes excluded due to empirical data sparsity largely coincide with loci independently annotated as technically unreliable, suggesting that early NaN-based exclusion was not arbitrary but aligned with established probe-quality criteria.

The redundancy structure across lists (Figure 4.9b) further indicates that most excluded CpGs are supported by a single resource, with progressively fewer loci flagged by multiple independent annotations. Annotation-based validation using the MethylationEPIC v1.0 manifest [58] further excluded non-CpG probes and loci lacking reliable genomic annotation. Across all filters, the dimensionality reduction is detailed in Table 4.3.

A probe-level exclusion summary is available in the project repository at `removed_cpgs_all_filters_summary_gse287331.csv`.

Statistical Transformation: β -to- M Values Following technical filtering, the β -value matrix was transformed into M -values using the logit relation defined in Equation (4.1), in accordance with established recommendations for variance stabilisation [21, 44, 45].

The empirical distributions (Figure 4.10) retain the characteristic bounded structure of β -values; however, in contrast to GSE69914 and GSE225845, the profiles appear less sharply bimodal and more broadly dispersed. Both Normal and Adjacent tissues exhibit reduced concentration at the methylation extremes and a wider intermediate range, consistent with the greater technical heterogeneity observed in this cohort.

After transformation, the corresponding M -value distributions become more symmetric, as expected under the logit mapping, yet remain visibly broader than in the other datasets. The mean–SD relationships confirm that while variance




Figure 4.10: Distributional comparison of β and M in Normal and Adjacent tissues in GSE287331.

stabilisation is achieved, dispersion remains comparatively elevated across the full methylation range. This behaviour is coherent with the stronger preprocessing burden documented in earlier steps and reflects the intrinsic variability of the released matrix.

4.4 Inter-Dataset Consistency After Preprocessing

After completing the intra-dataset preprocessing for each cohort, a concise inter-dataset analysis is performed to summarise the effects of the pipeline and to motivate the subsequent feature selection step.

Consistency of the Preprocessing Framework All three datasets underwent a harmonised preprocessing workflow, including initial integrity checks, literature-based technical filtering, manifest-driven probe validation, removal of sex-chromosome loci, and transformation from β -values to M-values. While specific implementation details differ slightly depending on platform characteristics and data quality, the conceptual structure of the pipeline is consistent across datasets.

Table 4.4 reports, for each dataset, the dimensionality of the methylation matrix before and after preprocessing. Despite substantial differences in initial platform coverage (450K vs. EPIC), technical filtering consistently removes a large fraction of probes, yielding cleaner and more reliable feature spaces.




Figure 4.11: CpG overlap across datasets before and after preprocessing.

Table 4.4: CpG dimensionality before and after preprocessing.

Dataset	Platform	CpGs <i>before</i>	CpGs <i>after</i>
GSE69914	450K	485,512	251,483
GSE225845	EPIC	750,426	530,683
GSE287331	EPIC	866,554	486,725

CpG overlap across datasets CpG overlap was evaluated both before and after preprocessing to characterise the relationship between cohorts. Prior to filtering (Figure 4.11a), the intersection is dominated by probes shared between the two EPIC datasets, with a smaller but substantial core common to all three cohorts, reflecting platform design constraints and baseline annotation differences. After the complete preprocessing pipeline (Figure 4.11b), the total number of CpGs is markedly reduced, yet a large shared core is preserved: the three-way intersection comprises 215,919 CpGs, representing a stable autosomal backbone consistently retained across platforms after technical filtering.

4.5 Post-Preprocessing Feature Space and Dimensionality Implications

The preprocessing pipeline developed in this chapter has systematically addressed the three main classes of technical confounders identified in Chapter 3: probe-design bias, unreliable probe sets, and statistical scale distortion. The resulting matrices are technically clean, fully observed, and expressed on a variance-stabilised M -value scale suitable for inferential modelling. Across all three cohorts, the three-way CpG intersection comprises 215,919 autosomal loci retained after platform-aware filtering, constituting a stable and harmonised feature backbone for cross-dataset analyses.

Yet the preprocessing outcome also sharpens the central analytical challenge.

The post-processed datasets remain extremely high-dimensional — 251,483 CpGs in GSE69914, 530,683 in GSE225845, and 486,725 in GSE287331 — and the exploratory analyses of Chapter 3 established that the Normal–Adjacent signal is focal and low-amplitude, with mean $|\Delta\beta|$ ranging from 0.009 to 0.031 across cohorts. In this regime, the overwhelming majority of retained CpGs carry no discriminative information for the contrast of interest: including them in downstream modelling would dilute the signal, inflate variance estimates, and compromise both interpretability and generalisation.

The limiting factor is therefore no longer data quality but *signal localisation*: identifying the restricted subset of CpG loci that capture biologically meaningful Normal–Adjacent variation, consistently across cohorts and independently of platform-specific noise. This requires a principled feature selection strategy that is sensitive to the subtle, focal nature of the epigenetic drift documented here, robust to the cohort-level heterogeneity observed in the inter-dataset comparisons, and compatible with the homoscedastic M -value representation adopted for modelling.

Chapter 5 addresses this challenge directly, developing and evaluating feature selection methodologies applied separately within each dataset-specific CpG space. In order to preserve the full informational richness of the EPIC platforms, feature selection is not restricted to the three-way CpG intersection but is performed on the complete post-processed feature set of each cohort, thereby avoiding artificial dimensional truncation driven by the lower coverage of the 450K array.

Chapter 5

Robust Feature Selection for Epigenetic Drift Characterisation

5.1 Statistical Challenges of High-Dimensional DNA Methylation Data

The preprocessing framework developed in Chapter 4.5 established a technically harmonised stabilised methylation space. However, preprocessing alone does not address the central statistical challenge of genome-wide DNA methylation data: extreme dimensionality relative to sample size. This challenge is particularly acute in the present study, where the objective is to characterise epigenetic differences along the Normal $\rightarrow$ Adjacent $\rightarrow$ Tumour axis in breast cancer — a setting in which biologically meaningful signal is expected to be subtle, spatially structured, and embedded in a feature space of several hundred thousand CpG loci.

High-throughput methylation arrays, such as the Illumina EPIC platform, interrogate approximately 850 000 CpG sites per sample [65], while the number of available subjects in any given cohort remains comparatively limited. This high-dimensional, low-sample-size (HDLSS) regime is well characterised in the statistical literature as a setting in which standard estimators become geometrically pathological: as dimensionality grows relative to n , pairwise distances concentrate, covariance estimation becomes unstable, and classifiers overfit to noise [66, 67]. In genomics specifically, the consequences are well documented — unconstrained feature spaces yield inflated classification accuracy that fails to replicate in independent cohorts [68, 69].

Feature selection is therefore not a cosmetic reduction step but a structural

necessity for any learning procedure applied to molecular profiling data.

5.1.1 Structural Properties Motivating Feature Selection

Three properties of DNA methylation data collectively motivate explicit dimensionality control.

High Dimensionality and Overfitting Risk In HDLSS settings, discriminative models may achieve near-perfect separation on training data while capturing noise rather than biological signal. The phenomenon has been formally characterised by [66], who showed that in very high dimensions, data from distinct classes become approximately equidistant unless genuine low-dimensional structure is present. Practically, this means that feature selection must precede, not follow, model fitting — and must itself be performed exclusively on training data to avoid optimistic bias [68].

Spatial Correlation and Redundancy CpG loci exhibit strong local correlation driven by genomic proximity, chromatin organisation, and shared regulatory context. This spatial autocorrelation has been well characterised in large-scale mapping studies: contiguous CpGs within co-methylated blocks (“methylation domains”) can span hundreds to thousands of base pairs and behave as single epigenetic units [70, 71]. Without explicit redundancy control, learning algorithms may over-represent densely correlated genomic regions while underweighting distributed epigenetic signals. This is especially relevant here, as differentially methylated regions (DMRs) along the Normal–Adjacent axis are expected to be spatially clustered rather than randomly distributed [72].

Biological Sparsity Disease-associated methylation alterations are typically sparse and structured. In breast cancer, systematic analyses have demonstrated that tumour-associated hypermethylation preferentially targets specific regulatory programmes — including Polycomb-repressed developmental genes and oestrogen-regulated loci — while genome-wide methylation changes in histologically normal adjacent tissue are more subtle but nonetheless detectable [14, 73]. This sparsity motivates filtering strategies that prioritise stable, biologically coherent CpG subsets over exhaustive feature retention.

5.1.2 Methodological Challenges in Genomic Feature Selection

Feature selection in genomic data presents challenges that go beyond simple variable ranking.

Information Leakage The most consequential pitfall is information leakage: when feature selection is performed on the full dataset prior to cross-validation, the evaluation procedure has implicitly seen the test labels during variable ranking. [68] demonstrated empirically that this leakage can inflate estimated accuracy by 10–40 percentage points in gene expression data; analogous inflation has been documented for methylation-based classifiers [69]. Strict containment of all selection steps within the training fold is therefore essential and is enforced throughout the present framework.

Significance Versus Stability A second challenge concerns the relationship between statistical significance and effect stability. In large feature spaces, extremely small p -values may arise for negligible or unstable effect sizes — a manifestation of the multiple testing problem that is not fully resolved by correction alone [74]. Conversely, biologically relevant features may show moderate but reproducible signal across subsamples. Stability selection [75] addresses this by quantifying the reproducibility of feature inclusion across repeated subsampling, providing selection frequency as a complementary criterion to statistical rank.

Predictive Optimisation Versus Interpretability A third challenge is the tension between predictive optimisation and interpretability. Aggressive feature selection driven solely by discriminative performance may yield compact but biologically opaque models, while overly conservative selection may retain sufficient redundancy to obscure the underlying regulatory architecture. In translational contexts, this tension has additional implications: in-silico classification accuracy does not automatically imply clinical utility [76], and the biological coherence of the selected feature set is therefore a criterion in its own right.

5.1.3 Objectives of the Present Framework

In light of the statistical and biological considerations discussed above, the feature selection strategy developed in this chapter is designed to satisfy four interconnected objectives.

- **Strict leakage control:** all selection procedures are confined to training data, preventing inflation of evaluation metrics through inadvertent information exposure.
- **Stability enforcement:** CpGs are retained only if they demonstrate reproducible discriminative behaviour across repeated stratified resampling, ensuring robustness to training-set perturbations.

- **Redundancy reduction:** correlation-based pruning limits the over-representation of co-methylated genomic blocks and promotes independent signal components.
- **Genomic diversification:** structural constraints prevent excessive concentration of selected features within specific chromosomes or regulatory contexts.

The hyperparameters governing each stage of the pipeline — including variability thresholds, stability cutoffs, correlation bounds, and diversification budgets — were fixed based on domain knowledge and methodological precedent rather than systematic optimisation. A formal sensitivity analysis of these parameters lies outside the scope of the present work and is identified as a direction for future investigation in Chapter 8.

The objective is therefore not merely dimensionality reduction, but the identification of a stable, biologically coherent, and generalisable CpG subset capable of characterising the Normal–Adjacent epigenetic transition and supporting cross-dataset validation.

5.2 General Feature Selection Framework

Building upon the statistical considerations outlined above, this section formally describes the feature selection workflow adopted in the present study. The procedure operates on the post-preprocessing methylation matrices defined in Chapter 4 and is structured as a sequence of strictly train-only operations. These steps progressively reduce dimensionality while preserving statistical validity, stability, and biological coherence. From variability screening and scale transformation to stability-based ranking and redundancy control, each component addresses a specific structural property of DNA methylation data, culminating in a diversified CpG subset suitable for downstream cross-dataset evaluation.

5.2.1 Empirical Variability-Based Dimensionality Reduction

The first stage of the feature selection framework consists of an unsupervised dimensionality reduction step aimed at removing CpGs that exhibit negligible variability within the tissue under study. This strategy is grounded in the empirically driven data reduction method proposed by [77], who demonstrated that a substantial fraction of CpGs interrogated by the Illumina 450K array are effectively non-variable within a given tissue context, contributing to the multiple testing burden — by inflating the number of hypotheses tested — without increasing the probability of detecting true differential methylation signals.

Data restriction and structural cleaning Let $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$, where $x_i \in [0,1]^p$ represents the vector of β -values for p CpG loci and $y_i \in \{0,1\}$ encodes the binary Normal–Adjacent label, with $y_i = 0$ denoting histologically normal tissue and $y_i = 1$ denoting tumour-adjacent tissue. Samples belonging to other tissue states (e.g. Tumour) are excluded from $\mathcal{D}$ prior to feature selection. This restriction ensures that the selected CpGs specifically characterise early epigenetic alterations occurring in histologically normal tissue adjacent to tumour, rather than late-stage tumour-specific methylation changes.

Train–test partition The dataset is partitioned into a training set $\mathcal{D}_{\text{train}}$ and a held-out test set $\mathcal{D}_{\text{test}}$ via stratified random sampling:

$$\mathcal{D} = \mathcal{D}_{\text{train}} \cup \mathcal{D}_{\text{test}}, \quad |\mathcal{D}_{\text{train}}| = 0.80n, \quad |\mathcal{D}_{\text{test}}| = 0.20n, \quad (5.1)$$

with class proportions preserved in both sets. Crucially, all feature selection statistics — including the variability filter defined below — are estimated exclusively on $\mathcal{D}_{\text{train}}$ and the resulting CpG subset is applied unchanged to $\mathcal{D}_{\text{test}}$. This design prevents information leakage and ensures that dimensionality reduction is statistically independent of evaluation [68].

Definition of the variability statistic For each CpG $j \in \{1, \dots, p\}$, variability is quantified using the robust inter-quantile range statistic proposed by [77]:

$$r_{\beta,j} = Q_{0.90}(\beta_j \mid \mathcal{D}_{\text{train}}) - Q_{0.10}(\beta_j \mid \mathcal{D}_{\text{train}}), \quad (5.2)$$

where $Q_{0.90}$ and $Q_{0.10}$ denote the empirical 90th and 10th percentiles of the β -value distribution computed across all samples in $\mathcal{D}_{\text{train}}$, pooling both tissue classes. Note that $r_{\beta,j}$ measures population-level dispersion unconditional on class membership: it quantifies whether a CpG is variable across individuals at all, not whether it discriminates between tissue types. A CpG is retained if and only if:

$$r_{\beta,j} \geq \tau, \quad \tau = 0.05, \quad (5.3)$$

yielding the retained index set $\mathcal{J}_{\text{Edgar}} = \{j \in \{1, \dots, p\} : r_{\beta,j} \geq \tau\}$. The threshold $\tau = 0.05$ corresponds to the empirical criterion introduced in [77]: CpGs exhibiting a population-level methylation range below 5% were operationally defined as non-variable and excluded. In the original study, this filter removed approximately 20–30% of probes on the 450K array across diverse tissue types [77].

Interpretation and scope Because $r_{\beta,j}$ is computed without reference to class labels, this filtering step is strictly unsupervised: it cannot introduce label-dependent bias and is therefore safe to apply upstream of the stratified cross-validation

framework described in Section 5.1.2. Its function is orthogonal to subsequent supervised ranking steps — it defines the candidate space over which discriminative selection will operate, not the final feature set. CpGs excluded by this criterion ($r_{\beta,j} < 0.05$) typically correspond to loci consistently hypo- or hypermethylated across individuals, frequently enriched in CpG islands and constitutively silenced promoter regions [77, 78]. Retaining such loci inflates the effective dimensionality of the feature space without contributing discriminative variance, exacerbating the HDLSS geometry. Low-variance independent filtering is a standard strategy in high-dimensional genomic analyses to reduce noise and improve statistical efficiency without compromising type I error control [79]. From a statistical perspective, $|\mathcal{J}_{\text{Edgar}}|$ defines the reduced multiple testing space on which all downstream differential methylation analyses and stability selection procedures are conducted. The CpG subset $\mathcal{J}_{\text{Edgar}}$, estimated on $\mathcal{D}_{\text{train}}$, is applied identically to $\mathcal{D}_{\text{test}}$ prior to any evaluation (Equation 5.1), preserving strict train–test separation throughout.

5.2.2 Re-alignment to M -values and Residual Variance Regularisation

The empirical variability screening of Section 5.2.1 is carried out in β -space, preserving continuity with the original inter-quantile dispersion criterion of [77]. This is consistent with the dual-representation strategy of Section 4.2.4: β -values are retained for biological interpretability, while M -values are adopted for statistical modelling because their logit transformation substantially stabilises the mean–variance relationship [21]. Once $\mathcal{J}_{\text{Edgar}}$ has been determined exclusively on $\mathcal{D}_{\text{train}}$, the analysis transitions to the M -value representation defined in Equation (4.1). The transformation is applied deterministically to the post-Edgar β matrices of both training and test sets: no model parameters are estimated at this stage. For each retained CpG $j \in \mathcal{J}_{\text{Edgar}}$, using the same numerical stabilisation constant introduced during preprocessing. The CpG subset selected in β -space is transferred unchanged to M -space, preserving strict train–test separation throughout.

Residual low-variance trimming (train-only) Although the Edgar filter removes CpGs with negligible population-level dispersion in β -space, a residual fraction of near-constant loci may survive after logit transformation. This occurs because the logit map is nonlinear: loci with β -values concentrated near the boundaries of $[0,1]$ can exhibit low inter-quantile range yet produce near-constant M -values once mapped to the real line. Retaining such loci inflates the effective dimensionality of the covariance structure without contributing discriminative signal, which is particularly detrimental in HDLSS settings. To address this, the pipeline systematically discards the lowest α -quantile of CpGs ranked by their empirical standard deviation in M -space on $\mathcal{D}_{\text{train}}$. Letting s_j denote this standard

deviation, the retained index set is

$$\mathcal{J}_{\text{var}} = \{j \in \mathcal{J}_{\text{Edgar}} : s_j \geq Q_\alpha(s)\}, \quad \alpha = 0.02, \quad (5.4)$$

corresponding to the removal of the bottom 2% of CpGs by dispersion. The step is unsupervised — class labels play no role — and introduces no information leakage. The quantile threshold is estimated solely on $\mathcal{D}_{\text{train}}$; the resulting binary mask is then applied unchanged to $\mathcal{D}_{\text{test}}$, maintaining complete statistical independence of the evaluation set.

The resulting set $\mathcal{J}_{\text{var}}$ constitutes the final candidate feature space entering the supervised discriminative ranking stage described in Section 5.1.2.

5.2.3 Biologically Weighted Stability Selection and Region-Level Consolidation

After structural dimensionality control (Sections 5.2.1 and 5.2.2), the feature space remains in the order of several hundred thousand CpGs. In HDLSS regimes, univariate ranking alone is insufficient: small perturbations in training composition can induce large instability in feature ordering, particularly when signal-to-noise ratios are moderate [66, 67]. To address this, we adopt a resampling framework that integrates statistical evidence, directional consistency, and biologically informed weighting across four sequential components.

Repeated Stratified Stability Framework Let $X \in \mathbb{R}^{n_{\text{train}} \times p}$ denote the post-filtered M -value matrix and $y \in \{0,1\}^{n_{\text{train}}}$ the binary Normal-Adjacent label. A Repeated Stratified K -Fold (RSKF) procedure is employed with $K = 5$ folds and $R = 10$ repetitions, yielding $S = 50$ train/validation splits. Repeated stratified resampling reduces the variance of estimated feature importance and mitigates selection instability arising from dependence on a single partition [68, 69]. For each split $s = 1, \dots, S$ and each CpG j , three quantities are computed exclusively on the training fold:

1. The signed mean difference $\Delta M_j^{(s)} = \bar{M}_{1,j}^{(s)} - \bar{M}_{0,j}^{(s)}$ is computed for all CpGs as an effect-size measure.
2. To reduce computational burden in the HDLSS regime, Mann-Whitney U p -values are evaluated only for the top 50,000 CpGs ranked by $|\Delta M_j^{(s)}|$ within the training fold. CpGs outside this prefiltered subset are assigned $p_j^{(s)} = 1$, corresponding to the least favourable rank. This two-stage procedure introduces a structural dependency between the prescreening criterion and the subsequent rank aggregation, since excluded CpGs receive a conservative rank by construction rather than by statistical evidence. The approximation

is justified by the width of the prefilter: retaining the top 50,000 candidates corresponds to approximately 6% of the full feature space and substantially exceeds the final selection target of 5,000 CpGs, thereby limiting the probability of excluding genuinely discriminative loci from downstream evaluation. The Mann–Whitney test is adopted as a rank-based nonparametric procedure to avoid distributional assumptions and ensure robustness to residual deviations from normality in M -space [80].

3. The same signed difference is computed independently on the validation fold to assess directional reproducibility across the partition.

Two rank statistics are then derived per split, $r_{p,j}^{(s)}$ (ascending rank of $p_j^{(s)}$), $r_{|\Delta M|,j}^{(s)}$ (descending rank of $|\Delta M_j^{(s)}|$), and averaged across splits:

$$\bar{r}_{p,j} = \frac{1}{S} \sum_{s=1}^S r_{p,j}^{(s)}, \quad \bar{r}_{\Delta M,j} = \frac{1}{S} \sum_{s=1}^S r_{|\Delta M|,j}^{(s)}. \quad (5.5)$$

The combined statistical score is the equal-weight convex combination:

$$\text{score}_{\text{stat},j} = 0.5 \bar{r}_{p,j} + 0.5 \bar{r}_{\Delta M,j}. \quad (5.6)$$

This rank aggregation follows a Borda-count-like scheme [81], in which equal-weight combination of complementary ranking criteria has been shown to reduce selection variance relative to either criterion alone [82]. The resulting statistical score is subsequently min–max normalised to the unit interval:

$$\text{score}_{\text{stat},j}^{\text{norm}} = \frac{\text{score}_{\text{stat},j} - \min(\text{score}_{\text{stat}})}{\max(\text{score}_{\text{stat}}) - \min(\text{score}_{\text{stat}})}.$$

Relying solely on p -values risks prioritising negligible but stable effects under large sample sizes, whereas magnitude-only ranking may overweight noisy large deviations. Combining both criteria mitigates this trade-off and is consistent with resampling-based stability principles in high-dimensional feature selection [75].

Directional Stability Constraint A CpG whose estimated effect direction reverses across resampling splits is unlikely to reflect a genuine, reproducible biological signal. We therefore impose a directional reproducibility criterion. For each CpG j , let $\text{sign}_{\text{train}}^{(s)}$ and $\text{sign}_{\text{val}}^{(s)}$ denote the signs of $\Delta M_j^{(s)}$ on the training and validation folds of split s , respectively. The directional stability score is:

$$\text{stab}_j = \frac{\#\{s : \text{sign}_{\text{train}}^{(s)} = \text{sign}_{\text{val}}^{(s)} \neq 0\}}{\#\{s : \text{sign}_{\text{train}}^{(s)} \neq 0 \text{ and } \text{sign}_{\text{val}}^{(s)} \neq 0\}}. \quad (5.7)$$

The directional stability score is computed as the proportion of resampling splits in which both training and validation folds exhibit non-zero effect direction and agree in sign. CpGs for which the denominator equals zero — i.e. all splits yield $\Delta M_j^{(s)} = 0$ on at least one fold — are assigned $\text{stab}_j = 0$ by convention, rendering them ineligible for selection under the stability constraint. Only CpGs with $\text{stab}_j \geq 0.75$ are retained, requiring directional agreement in at least 75% of the splits for which a non-zero effect direction is observed in both folds. The threshold of 0.75 is chosen in analogy with the selection frequency cutoff recommended by [75], where a 75% inclusion rate across subsampling splits is identified as a conservative criterion for stable feature identification. The same numerical threshold is here applied to directional consistency rather than selection frequency, extending the stability principle from feature inclusion to effect reproducibility — a particularly relevant criterion when the target signal is moderate in magnitude, as expected for early epigenetic drift in histologically normal adjacent tissue.

Adaptive Effect-Size Filtering A global effect estimate is computed on the full training set, $\Delta M_j^{\text{full}} = \bar{M}_{1,j} - \bar{M}_{0,j}$, and a CpG is retained only if $|\Delta M_j^{\text{full}}| \geq Q_{0.60}(|\Delta M^{\text{full}}|)$. Rather than applying a fixed ΔM cutoff — which would be sensitive to inter-cohort differences in dispersion — this adaptive percentile-based criterion maintains scale invariance across datasets. A fallback to the 50th percentile is applied when the resulting candidate pool would otherwise be insufficient for downstream selection. Together with the directional stability constraint, this filter ensures that only CpGs with both reproducible direction and non-negligible magnitude reach the biological weighting stage. The fallback to the 50th percentile is triggered whenever the resulting candidate pool contains fewer than 15,000 CpGs, ensuring sufficient capacity for the downstream consolidation stage.

Integration of Biological Priors Purely statistical ranking may underweight loci in regulatory regions known to mediate transcriptional control. Promoter-associated CpGs and CpG islands are well-established sites of regulatory methylation changes in cancer and early field effects [70, 73]. Each CpG is assigned a discrete biological prior weight $w_j \in \{1.00, 0.70, 0.50, 0.20\}$ according to its island and promoter annotation:

$$w_j = \begin{cases} 1.00 & \text{Island + TSS} \\ 0.70 & \text{Island only} \\ 0.50 & \text{Promoter (TSS) only} \\ 0.20 & \text{Other loci} \end{cases} \quad (5.8)$$

The discrete weights are subsequently min-max normalised to the unit interval, yielding $w_j^{\text{norm}} \in [0,1]$. In order to prioritise biologically annotated loci while

preserving a lower-is-better scoring convention, the biological contribution is defined as $1 - w_j^{\text{norm}}$. After min-max normalisation of the statistical score $\text{score}_{\text{stat},j}$, the final integrated score is

$$\text{score}_j = (1 - \lambda) \text{score}_{\text{stat},j}^{\text{norm}} + \lambda (1 - w_j^{\text{norm}}), \quad \lambda = 0.20. \quad (5.9)$$

Min-max normalisation is applied independently to each component prior to combination, ensuring commensurability of scale. The procedure is sensitive to extreme values; however, given the large number of CpGs ($|\mathcal{J}_{\text{var}}| \gg 10^4$), the influence of individual outliers on the normalised distribution is expected to be negligible. The coefficient $\lambda = 0.20$ reflects a deliberate design choice rather than an empirically optimised parameter: statistical evidence contributes 80% of the integrated score, while the remaining 20% provides modest upweighting for loci with established regulatory function in promoter methylation and transcriptional silencing [8]. The top 15,000 CpGs satisfying both the stability and adaptive effect-size constraints define the candidate feature pool passed to the next stage.

Region-Level Anchoring on CpG Islands CpGs are not independent genomic entities: spatially proximal loci within co-methylated blocks co-vary as coordinated regulatory units [70]. Selecting isolated top-ranked probes therefore risks capturing stochastic single-probe fluctuations rather than coherent epigenetic events, which is particularly problematic when the signal of interest — early field-effect drift — is expected to manifest as concerted changes across regulatory domains. To enforce spatial coherence, candidate CpGs are grouped by annotated CpG island. For each island containing at least two candidate CpGs, a region-level score is defined as:

$$\text{region_score} = \text{median}(\text{score}_j) \times \text{fraction}_{\text{stable}}, \quad (5.10)$$

where $\text{fraction}_{\text{stable}}$ denotes the proportion of CpGs within the island satisfying both $\text{stab}_j \geq 0.75$ and $|\Delta M_j^{\text{full}}| \geq \tau$, with τ denoting the adaptive effect-size threshold selected in the previous step (60th percentile or 50th percentile fallback). The multiplicative formulation jointly penalises islands in which either the median discriminative score is weak or the proportion of internally stable CpGs is low: an island driven by isolated high-scoring probes surrounded by unstable loci receives a substantially discounted region score, ensuring that anchored islands exhibit both statistical quality and internal coherence. Only islands with $\text{fraction}_{\text{stable}} \geq 0.60$ and at least two candidate CpGs are retained. From each such island, the top $m = 2\text{--}3$ CpGs are selected into an anchored pool; the remaining candidates form a complementary non-anchored set. This hierarchical consolidation reduces susceptibility to single-probe artefacts and aligns feature selection with the biological expectation that early epigenetic drift occurs in clustered regulatory domains rather than as isolated stochastic events.

5.2.4 Correlation-Based Redundancy Pruning and Graph-Theoretic Clustering

Despite stability filtering and region-level anchoring, the candidate feature space retains strong local correlation. DNA methylation levels exhibit pronounced spatial autocorrelation driven by genomic proximity, shared chromatin context, and coordinated regulatory control [70, 71]: in high-density arrays such as EPIC, neighbouring CpGs within islands and promoter regions frequently co-vary as near-redundant blocks. Retaining multiple highly correlated loci inflates the effective dimensionality of the design matrix without increasing independent information content, inducing multicollinearity, unstable coefficient estimates, and variance inflation in downstream classifiers [67, 66]. Explicit correlation-based redundancy pruning is therefore applied to the candidate pool before passing it to the diversification stage.

Correlation Structure on Training Data Let $Z \in \mathbb{R}^{n_{\text{train}} \times q}$ denote the M -value matrix restricted to the q candidate CpGs, with each column standardised using training-set mean μ_j and standard deviation s_j :

$$Z_{ij} = \frac{M_{ij} - \mu_j}{s_j}.$$

where s_j is the sample standard deviation computed with Bessel correction (ddof = 1) on the training set. A small numerical constant is added in practice to prevent division by zero in degenerate cases. The empirical Pearson correlation matrix is then:

$$C = \frac{1}{n_{\text{train}} - 1} Z^\top Z. \quad (5.11)$$

Because columns of Z are standardised to zero mean and unit sample variance, Equation (5.11) coincides with the empirical Pearson correlation matrix estimated exclusively on $\mathcal{D}_{\text{train}}$. All quantities are estimated exclusively on $\mathcal{D}_{\text{train}}$ to preserve strict evaluation independence.

Graph Construction and Connected Components An undirected graph $G = (V, E)$ is constructed with one vertex per candidate CpG; an edge connects j and k whenever their absolute correlation exceeds the threshold:

$$(j, k) \in E \iff |C_{jk}| \geq \tau_c, \quad \tau_c = 0.85, \quad (5.12)$$

consistent with strong-correlation redundancy thresholds commonly adopted in high-dimensional omics analyses [83, 84]. By convention, each vertex is connected to itself in the adjacency matrix; this does not alter the connected component

structure but simplifies implementation. Clusters are defined as the connected components of G : a cluster $\mathcal{C}$ is a maximal vertex subset in which every pair (j, k) is joined by at least one path in G . This graph-theoretic formulation is strictly preferable to pairwise sequential pruning because correlation propagates transitively: two CpGs with $|C_{jk}| < \tau_c$ may both be strongly correlated to a third, and would therefore represent the same co-methylated block without being linked directly. Connected components capture this transitive redundancy structure in a single, consistent pass.

Representative Selection per Cluster For each connected component $\mathcal{C}_m$, a single representative CpG is retained:

$$j^* = \arg \min_{j \in \mathcal{C}_m} \text{score}_j, \quad (5.13)$$

where score_j is the integrated biologically weighted stability score of Section 5.2.3, in which a lower value denotes stronger statistical and biological evidence. CpGs lacking a valid stability score are assigned an infinite value and are therefore ineligible for selection as cluster representatives. Selecting the minimum thus retains the most discriminative and stable CpG within each correlated block, discarding all redundant co-varying loci. While region-level anchoring leverages genomic annotation to enforce spatial coherence based on CpG island structure, the present clustering step operates purely on empirical covariance structure estimated from the training data.

Separate Clustering of Anchored and Non-Anchored Pools To preserve the island-level spatial structure established by region-level anchoring before applying global redundancy reduction, clustering is performed separately on the region-anchored pool A (CpGs selected via island consolidation) and the non-anchored complementary pool B . Let $\mathcal{R}_A$ and $\mathcal{R}_B$ denote the representative sets obtained from each pool. Representatives from A are prioritised in the final union, and the redundancy-pruned feature set is: $\mathcal{R} = \mathcal{R}_A \cup \mathcal{R}_B$.

Statistical and Biological Rationale From a bias–variance standpoint, removing highly correlated predictors reduces estimator variance while preserving most of the predictive signal, since strongly correlated CpGs encode largely overlapping epigenetic information. In HDLSS settings this is particularly consequential: redundant features exacerbate the ill-conditioning of the design matrix, inflate effective model complexity, and destabilise decision boundaries [67]. Removing redundant predictors reduces effective dimensionality and improves the numerical conditioning of the design matrix without materially reducing predictive information content.

The result is a structurally compact, non-redundant feature set that retains statistical strength, directional stability, biological coherence, and spatial structure — forming the input to the genomic diversification stage described in Section 5.2.5.

5.2.5 Genomic Diversification via Constrained Greedy Selection

After correlation-based redundancy pruning (Section 5.2.4), the feature set $\mathcal{R}$ remains enriched for highly ranked CpGs but may still exhibit structural imbalance. Genome-wide methylation arrays are inherently non-uniform — CpG density varies substantially across chromosomes, islands, shores, shelves, and open sea regions — so unconstrained greedy ranking by statistical score tends to over-concentrate features in CpG-dense promoter islands, chromosomes with strong signal, and local genomic windows of coordinated drift. Such concentration artificially reduces effective genomic coverage and increases the risk that downstream models capture locus-specific rather than system-level epigenetic patterns, reducing structural generalisability across cohorts. We therefore introduce a diversification step formulated as a constrained subset selection problem.

Problem Formulation Let $\mathcal{R} = \{j_1, \dots, j_m\}$ denote the redundancy-pruned CpG pool ordered by increasing integrated score score_j (Section 5.2.3). CpGs lacking valid manifest annotation for chromosome and genomic position are excluded prior to diversification, as spatial constraints require coordinate information. The goal is to select a subset $\mathcal{S} \subset \mathcal{R}$, $|\mathcal{S}| = K = 5000$, that maximises overall discriminative quality under structural diversity constraints. In practice, the effective target is defined as $K_{\text{eff}} = \min(K, |\mathcal{R}|)$ to accommodate cases in which fewer than K mapped CpGs are available; all constraint budgets are computed with respect to K_{eff} . The target cardinality $K = 5000$ represents approximately 1.0% of the probes interrogated by the Illumina 450K array and approximately 0.6% of those interrogated by the EPIC array. It is selected to provide sufficient genomic coverage for cross-cohort evaluation while maintaining a feature-to-sample ratio compatible with stable supervised learning in the available cohorts. The choice is acknowledged as a design parameter rather than an optimised quantity; its sensitivity is identified as a direction for future investigation in Chapter 8. Formally:

$$\begin{aligned}
 \min_{\mathcal{S} \subset \mathcal{R}} \quad & \sum_{j \in \mathcal{S}} \text{score}_j \\
 \text{s.t.} \quad & |\mathcal{S}| = K_{\text{eff}}, \\
 & |\{j \in \mathcal{S} : c(j) = \chi\}| \leq \lfloor \alpha_{\text{chr}} K_{\text{eff}} \rfloor \quad \forall \chi, \\
 & |\{j \in \mathcal{S} : \gamma(j) = g\}| \leq \lfloor \alpha_{\text{ctx}} K_{\text{eff}} \rfloor \quad \forall g, \\
 & |\{j \in \mathcal{S} : w(j) = \omega\}| \leq \beta \quad \forall \omega.
 \end{aligned} \tag{5.14}$$

Each constraint targets a distinct axis of potential structural imbalance and is described in turn below.

Chromosomal Balance Constraint Let $c(j)$ denote the chromosome of CpG j . No chromosome may contribute more than a fixed fraction of the final set:

$$|\{j \in \mathcal{S} : c(j) = \chi\}| \leq \alpha_{\text{chr}} K, \quad \alpha_{\text{chr}} = 0.08. \quad (5.15)$$

The budget per chromosome is computed as $\lfloor \alpha_{\text{chr}} K_{\text{eff}} \rfloor$, with a minimum of one CpG allowed per chromosome. The 8% cap prevents over-representation of chromosomes that harbour dense signal regions, a configuration known to affect classifier stability when disease-associated loci cluster in specific chromosomal domains [83].

Genomic Context Constraint Let $\gamma(j)$ denote the CpG island context of locus j (Island, Shore, Shelf, or OpenSea). The fraction of any single context class is bounded by:

$$|\{j \in \mathcal{S} : \gamma(j) = g\}| \leq \alpha_{\text{ctx}} K, \quad \alpha_{\text{ctx}} = 0.65. \quad (5.16)$$

The context budget is computed as $\lfloor \alpha_{\text{ctx}} K_{\text{eff}} \rfloor$, with at least one CpG permitted per context class. Although promoter islands are biologically important, cancer-related methylation changes are not confined to islands and frequently involve shores and open sea regions [70, 73]. The soft cap ensures that non-island regulatory regions retain representation in the final panel.

Spatial Window Constraint To prevent local hyper-concentration, each chromosome is partitioned into non-overlapping windows of $W = 500$ kb, and the number of selected CpGs per window is hard-capped: $|\{j \in \mathcal{S} : w(j) = \omega\}| \leq 15$, where $w(j)$ is the window index of CpG j . Formally, windows are defined via integer binning $w(j) = \lfloor \text{pos}(j)/W \rfloor$ within each chromosome. Since methylation domains frequently span tens to hundreds of kilobases [70], this constraint preserves regional representation while preventing dominance by a single hyper-variable block — a risk that correlation pruning alone cannot fully eliminate at the macro-scale.

Greedy Approximation with Progressive Relaxation The constrained optimisation problem (5.14) is combinatorial and NP-hard in general, as it resembles cardinality-constrained subset selection with multiple linear constraints. Rather than solving a full mixed-integer program, we adopt a deterministic greedy strategy: CpGs are processed in order of increasing score_j , ties are resolved deterministically by stable ordering, and each candidate is added to $\mathcal{S}$ if and only if all active constraints are simultaneously satisfied, until $|\mathcal{S}| = K$.

If the strict constraint regime fails to reach K , constraints are relaxed by staged deactivation (rather than by increasing thresholds), first removing the spatial window constraint, then the genomic context constraint, and finally the chromosome balance constraint.

At each stage, the specified constraint is fully deactivated while previously relaxed constraints remain inactive. Spatial over-concentration is relaxed first because local genomic crowding is most detrimental to coverage; chromosomal imbalance is relaxed last as it has the weakest direct effect on independent information content. The constrained greedy strategy does not admit formal approximation guarantees in this setting, as the objective is linear rather than submodular and the constraint structure involves multiple simultaneous bounds. The approach is adopted as a computationally tractable heuristic; its empirical adequacy is assessed indirectly through the constraint activation frequency reported in Section 5.3. The frequency with which progressive constraint relaxation is activated in practice is reported in Section 5.3 for each dataset, allowing empirical assessment of whether the final selection approximates the fully constrained solution or degrades toward unconstrained score-based ranking.

5.3 Dataset-Specific Implementation


This section documents the dataset-specific application of the feature selection framework described in Section 5.2. For each cohort, the outcome of empirical variability screening, M -value realignment and residual variance trimming, stability-based discriminative ranking with directional consistency constraints, redundancy reduction via correlation-graph clustering, and constrained genomic diversification is reported in detail.

The results are presented separately for GSE69914 (Section 5.3.1), GSE225845 (Section 5.3.2), and GSE287331 (Section 5.3.3).

5.3.1 Dataset GSE69914

The feature selection framework described in Section 5.2 was applied to the pre-processed GSE69914 cohort, restricted to the Normal and Adjacent tissue classes, yielding a final panel of $K = 5000$ CpGs after variability filtering, stability ranking, redundancy pruning, and diversification. The selected loci are available in the project repository at `final_5000_gse69914.csv`.

Empirical Variability-Based Dimensionality Reduction Variability was quantified via the inter-quantile range $r_{\beta,j} = P90(\beta) - P10(\beta)$ computed on the training set. Figure 5.1(a) shows the empirical distribution of r_{β} across CpGs.




(a) $r_\beta = P90(\beta) - P10(\beta)$ on the training set. Candidate thresholds (0.01, 0.02, 0.05) are indicated.

(b) Distribution of $SD(M)$ on the training set. The dashed line indicates the bottom 2% quantile used for residual variance trimming.

Figure 5.1: Variability-based filtering in β -space and residual variance trimming in M -space in GSE69914.

The distribution is strongly right-skewed, with a substantial mass concentrated below $r_\beta = 0.05$. Applying the predefined threshold $\tau = 0.05$ removed 100,103 low-variability loci, defining the reduced candidate space for downstream supervised ranking. Lower candidate thresholds (0.01, 0.02) are shown for reference but were not adopted, as they would retain a substantially larger fraction of near-constant probes. To assess whether this filtering step induced structural genomic bias, the distribution of invariant loci across gene-feature categories was examined via fold-change enrichment analysis. As reported in Appendix B, invariant CpGs showed moderate enrichment in promoter-proximal regions (TSS200, first exon, 5'UTR), whereas gene bodies and 3'UTR regions were relatively under-represented. No extreme over-representation was observed, indicating that the variability-based reduction primarily eliminates low-dispersion loci without introducing pathological distortion of the genomic feature space.

Re-alignment to M -values and Residual Variance Regularisation The retained CpGs were deterministically transformed to M -values, after which a residual low-variance trimming was applied in M -space. Figure 5.1b displays the distribution of empirical standard deviations $SD(M)$ on the training set. The distribution is unimodal with limited heavy-tail behaviour. Removal of the bottom 2% of CpGs by dispersion eliminated 3,028 loci. Although quantitatively modest, this step improves covariance conditioning in the HDLSS regime.



(a) Distribution of directional stability scores S_{dir} . The dashed line indicates the cutoff $\text{stab}_j \geq 0.75$.

(b) Cluster size survival curves for anchored (Pool A) and non-anchored (Pool B) CpGs under $|r| \geq 0.85$.

Figure 5.2: Stability enforcement and correlation-based redundancy structure in GSE69914.


Biologically Weighted Stability Selection and Region-Level Consolidation

Repeated stratified resampling was used to estimate stability of signed mean differences. Figure 5.2a shows the distribution of the directional stability score S_{dir} . The majority of CpGs exhibit high directional reproducibility, with substantial mass above 0.75. The threshold $\text{stab}_j \geq 0.75$ excludes loci whose effect direction reverses across resampling splits, enforcing robustness consistent with the stability-selection principle of [75]. After stability and adaptive effect-size filtering (60th percentile, corresponding to $|\Delta M| \approx 0.15$), the top 15,000 CpGs were retained by design, defining the candidate pool passed to correlation-based redundancy pruning.

Correlation-Based Redundancy Pruning and Graph-Theoretic Clustering

After stability and effect-size filtering, correlation-based graph clustering was applied separately to the anchored and non-anchored pools using a threshold of $|r| \geq 0.85$ computed on the training set. Figure 5.2b shows the cluster size survival curves, whose rapid decay indicates that most connected components are of size 1–3, reflecting limited local redundancy. Only a small number of large correlated blocks are observed, corresponding to densely co-methylated regions. Selecting a single representative CpG per component reduced the candidate space to 12,313 loci, defining the redundancy-pruned set passed to the diversification stage.

Genomic Diversification via Constrained Greedy Selection The constrained greedy diversification procedure selected $K = 5000$ CpGs under chromosome, genomic context, and 500kb spatial-window constraints. The strict constraint



(a) $r_\beta = P90(\beta) - P10(\beta)$ on the training set. Candidate thresholds (0.01, 0.02, 0.05) are indicated.

(b) Distribution of $SD(M)$ on the training set. The dashed line indicates the bottom 2% quantile used for residual variance trimming.


Figure 5.3: Variability-based filtering in β -space and residual variance trimming in M -space in GSE225845.

regime was sufficient to reach K without requiring relaxation (maximum chromosome share 8%, maximum 15 CpGs per 500kb window). The final panel comprises 2,151 Island loci, 1,342 OpenSea loci, 712 N_Shore loci, 506 S_Shore loci, and 150 N_Shelf loci, indicating a diversified genomic-context composition rather than promoter-restricted enrichment. OpenSea loci were not excluded a priori, as distal regulatory regions may capture long-range epigenetic alterations relevant to early field effects.

5.3.2 Dataset GSE225845

The feature selection framework described in Section 5.2 was applied to the preprocessed GSE225845 cohort, restricted to the Normal and Adjacent tissue classes, yielding a final panel of $K = 5000$ CpGs after variability filtering, stability ranking, redundancy pruning, and diversification. The selected loci are available in the project repository at `final_5000_gse225845.csv`.

Empirical Variability-Based Dimensionality Reduction Variability was quantified via the inter-quantile range $r_{\beta,j} = P90(\beta) - P10(\beta)$ computed on the training set. Figure 5.3a shows the empirical distribution of r_β across CpGs. Applying the predefined threshold $\tau = 0.05$ removed 134,039 low-variability loci, defining the reduced candidate space for downstream supervised ranking. Lower candidate thresholds (0.01, 0.02) are shown for reference but were not adopted. As observed in GSE69914, the variability-based reduction does not induce pathological



(a) Distribution of directional stability scores S_{dir} . The dashed line indicates the cutoff $\text{stab}_j \geq 0.75$.

(b) Cluster size survival curves for anchored (Pool A) and non-anchored (Pool B) CpGs under $|r| \geq 0.85$.

Figure 5.4: Stability and redundancy diagnostics in GSE225845.

genomic bias: invariant loci display moderate enrichment in promoter-proximal regions, while gene bodies and 3'UTR regions are relatively under-represented (see Appendix B).

Re-alignment to M -values and Residual Variance Regularisation The retained CpGs were deterministically transformed to M -values, after which a residual low-variance trimming was applied in M -space. Figure 5.3b displays the distribution of empirical standard deviations $SD(M)$ on the training set. Removal of the bottom 2% of CpGs by dispersion eliminated 7,933 loci (cutoff $SD(M) = 0.284$), defining the variance-regularised candidate space entering the stability-ranking stage.

Biologically Weighted Stability Selection and Region-Level Consolidation Repeated stratified resampling was used to estimate stability of signed mean differences. Figure 5.4a shows the distribution of the directional stability score S_{dir} . The distribution is strongly right-skewed, with a substantial concentration of CpGs exhibiting S_{dir} close to 1, indicating highly consistent effect directions across resampling splits. The threshold $\text{stab}_j \geq 0.75$ therefore removes only loci with unstable or sign-reversing effects, retaining the majority of directionally robust signals in this cohort. After stability and adaptive effect-size filtering (60th percentile), the top 15,000 CpGs were retained by design, defining the candidate pool passed to correlation-based redundancy pruning.

Correlation-Based Redundancy Pruning and Graph-Theoretic Clustering After stability and effect-size filtering, correlation-based graph clustering was


applied separately to the anchored and non-anchored pools using a threshold of $|r| \geq 0.85$ computed on the training set. Figure 5.4b shows the cluster size survival curves. The survival functions decay rapidly for small k , indicating that most connected components are of size 1–3. However, a pronounced heavy tail is observed, particularly in the non-anchored pool, with a limited number of large correlated blocks spanning tens to hundreds of CpGs. This redundancy pruning step substantially reduced the candidate space from 15,000 to 6,453 loci, indicating a markedly higher correlation structure in GSE225845 compared to GSE69914. The result suggests extended co-methylation domains consistent with the denser EPIC probe coverage. Selecting a single representative CpG per connected component defines the redundancy-pruned set passed to the diversification stage.

Genomic Diversification via Constrained Greedy Selection The constrained greedy diversification procedure was applied to the 6,453 redundancy-pruned CpGs under chromosome, genomic-context, and 500kb spatial-window constraints. The strict constraint regime was sufficient to reach $K = 5000$ without requiring relaxation (maximum chromosome share 8%, maximum context share 65%, maximum 15 CpGs per 500kb window). The final panel comprises 2,115 Island loci, 1,746 OpenSea loci, 534 N_Shore loci, 436 S_Shore loci, and 91 N_Shelf loci, indicating a diversified genomic-context composition. The chromosome cap was actively binding in the strict stage, ensuring balanced genomic representation without overconcentration in high-density regions. The resulting panel therefore satisfies all diversification constraints by construction.

5.3.3 Dataset GSE287331

The feature selection framework described in Section 5.2 was applied to the preprocessed GSE287331 cohort, restricted to the Normal and Adjacent tissue classes, yielding a final panel of $K = 5000$ CpGs after variability filtering, stability ranking, redundancy pruning, and diversification. The selected loci are available in the project repository at `final_5000_gse287331.csv`.

Empirical Variability-Based Dimensionality Reduction Variability was quantified via the inter-quantile range $r_{\beta,j} = P90(\beta) - P10(\beta)$ computed on the training set. Figure 5.5a shows the empirical distribution of r_{β} across CpGs, which is heavily concentrated near zero, indicating a large fraction of weakly variable loci. Applying the predefined threshold $\tau = 0.05$ removed 200,725 low-variability CpGs, substantially reducing the dimensionality of the feature space prior to supervised ranking. Lower candidate thresholds are shown for reference but were not adopted, as they would retain a markedly larger proportion of near-constant probes. Consistent with the other cohorts, invariant CpGs exhibit moderate



(a) $r_\beta = P90(\beta) - P10(\beta)$ on the training set. Candidate thresholds (0.01, 0.02, 0.05) are indicated.


(b) Distribution of $SD(M)$ on the training set. The dashed line indicates the bottom 2% quantile used for residual variance trimming.

Figure 5.5: Variability-based filtering in β -space and residual variance trimming in M -space in GSE287331.

enrichment in promoter-proximal regions (notably TSS200 and first exon), while distal or non-promoter regions are relatively under-represented (see Appendix B). The magnitude of enrichment is less pronounced than in GSE69914 and GSE225845, suggesting that the variability structure of this dataset is more diffusely distributed across genomic contexts.

Re-alignment to M -values and Residual Variance Regularisation The retained CpGs were deterministically transformed to M -values, after which a residual low-variance trimming was applied in M -space. Figure 5.5b displays the distribution of empirical standard deviations $SD(M)$ on the training set, which is unimodal with moderate right-tail behaviour. Removal of the bottom 2% of CpGs by dispersion (cutoff $SD(M) = 0.271$) eliminated 5,720 loci, defining the variance-regularised candidate space entering the stability-ranking stage.

Biologically Weighted Stability Selection and Region-Level Consolidation Repeated stratified resampling was used to estimate stability of signed mean differences. Figure 5.6a shows the distribution of the directional stability score S_{dir} , which is sharply concentrated near 1. A pronounced peak at high stability values indicates that the vast majority of CpGs exhibit highly consistent effect directions across resampling splits. The threshold $\text{stab}_j \geq 0.75$ therefore removes only a small subset of unstable loci, enforcing directional robustness in line with the stability-selection principle of [75] while retaining most candidate signals. After stability and adaptive effect-size filtering (60th percentile), the top 15,000 CpGs



(a) Distribution of directional stability scores S_{dir} . The dashed line indicates the cutoff $\text{stab}_j \geq 0.75$.

(b) Cluster size survival curves for anchored (Pool A) and non-anchored (Pool B) CpGs under $|r| \geq 0.85$.

Figure 5.6: Stability and redundancy diagnostics in GSE287331.

were retained by design, defining the candidate pool passed to correlation-based redundancy pruning.

Correlation-Based Redundancy Pruning and Graph-Theoretic Clustering

After stability and effect-size filtering, correlation-based graph clustering was applied separately to the anchored and non-anchored pools using a threshold of $|r| \geq 0.85$ computed on the training set. Figure 5.6b shows the cluster size survival curves. The survival functions exhibit a rapid decay, indicating that most connected components are of size 1–3 and that large correlated blocks are rare. Compared to GSE225845, the absence of a pronounced heavy tail suggests a more locally structured correlation pattern without extended co-methylation domains. Selecting a single representative CpG per connected component reduced the candidate space to 6,300 loci, defining the redundancy-pruned set passed to the diversification stage.

Genomic Diversification via Constrained Greedy Selection

The constrained greedy diversification procedure was applied to the 6,300 redundancy-pruned CpGs under chromosome, genomic-context, and 500kb spatial-window constraints. The strict constraint regime was sufficient to reach $K = 5000$ without requiring relaxation. The final panel comprises 1,992 OpenSea loci, 1,692 Island loci, 556 N_Shore loci, 507 S_Shore loci, and 135 S_Shelf loci, indicating a diversified genomic-context composition with a substantial contribution from distal regions.




Figure 5.7: Three-way overlap of the final 5,000 CpG signatures independently selected in GSE69914, GSE225845 and GSE287331.

5.4 Inter-Dataset Stability of the Final 5,000 CpG Signatures

The feature selection framework described in Section 5.3 was applied independently to each cohort, yielding three dataset-specific panels of $K = 5,000$ CpG loci optimised for within-dataset discrimination along the Normal–Adjacent axis. The present section quantifies the degree to which these panels converge on a shared epigenetic drift signal, thereby assessing the cross-cohort replicability of the feature selection outcome. Consistency is evaluated at three complementary levels: (i) identity-level overlap of selected loci (set-level replicability), (ii) concordance of estimated effect sizes across cohort intersections (effect-level agreement), and (iii) transferability of the drift structure as a discriminative ranking in held-out datasets. This multi-resolution framework mirrors the analytical strategy adopted for the pre-selection inter-dataset analysis of Section 5.3, now applied to the output of the full selection pipeline rather than to the global methylome.

5.4.1 Set-Level Replicability

Pairwise intersections between the three final 5,000-CpG panels were computed and are visualised in Figure 5.7. The identity-level overlap is summarised by the Jaccard similarity coefficient, which for two selection sets A and B is defined as

$$J(A, B) = \frac{|A \cap B|}{|A \cup B|}, \quad (5.17)$$

Table 5.1: Permutation-based enrichment of pairwise overlaps among the three final 5,000-CpG panels.

Pair	Shared CpGs	Expected Overlap	Enrichment Ratio	p -value
GSE69914–GSE225845	148	69.0	2.14	$< 10^{-4}$
GSE69914–GSE287331	84	68.3	1.23	0.024
GSE225845–GSE287331	128	37.9	3.38	$< 10^{-4}$

where $|A \cap B|$ denotes the cardinality of the shared loci and $|A \cup B|$ the size of their union. This scale-invariant measure normalises the raw intersection count against the total selection universe, rendering it comparable across cohort pairs with different platform intersections. The observed values, $J(\text{GSE69914}, \text{GSE225845}) = 0.0150$, $J(\text{GSE69914}, \text{GSE287331}) = 0.0085$, $J(\text{GSE225845}, \text{GSE287331}) = 0.0130$, indicate that fewer than 1.5% of selected CpGs are shared between any two cohort-specific panels, confirming that identity-level replicability is limited across all pairs. This pattern is consistent with the weak locus-level reproducibility observed in the pre-selection inter-dataset analysis (Section 3.4). Given the high-dimensional setting, cohort-specific noise structure, and heterogeneous sampling conditions, only partial overlap between independently derived panels is expected.

Permutation-based assessment of overlap significance A naive interpretation of the raw intersection size would be misleading without accounting for the baseline expected overlap under random selection. The appropriate null model must reflect the actual constraints of the selection procedure: each panel draws exactly $K = 5,000$ CpGs from its own post-preprocessing feature space, and the two panels can only share loci that are present in *both* datasets after platform-specific filtering. The effective sampling universe for each pair is therefore the post-preprocessing pairwise intersection $\mathcal{U}_{AB} = \mathcal{C}_A \cap \mathcal{C}_B$, where $\mathcal{C}_A$ and $\mathcal{C}_B$ denote the full sets of CpGs retained after preprocessing in datasets A and B , respectively.

Under this null, 10,000 independent replications were generated by drawing two sets of $K = 5,000$ CpGs uniformly at random from $\mathcal{U}_{AB}$, without replacement, and recording their intersection size. The empirical p -value is defined as the proportion of replications in which the random overlap equals or exceeds the observed value. This construction is conservative relative to null models based on the full genome or on the union of platform probes, since the restricted universe $\mathcal{U}_{AB}$ raises the baseline expected overlap; significance under this test therefore provides a more stringent criterion for enrichment. Results are reported in Table 5.1.

All three pairwise overlaps exceed the permutation null at nominal significance. The most pronounced enrichment is observed for the EPIC–EPIC pair (GSE225845–GSE287331; enrichment $\times 3.4$), reflecting the larger shared platform intersection available to that pair. The HM450–EPIC pairs yield smaller but still significant




Figure 5.8: Inter-dataset effect concordance of the final 5,000 CpG signatures.

enrichments, indicating that the selection procedure recovers a non-random shared subset even across platform boundaries.

Importantly, statistical significance here quantifies enrichment relative to the large-cardinality random baseline, and should not be conflated with biological equivalence of the selected panels. The restricted sampling universe $\mathcal{U}_{AB}$ is itself of order $\mathcal{O}(10^5)$, so even modest absolute overlaps can achieve nominal significance when drawn from such a large pool. Jaccard indices below 0.02 demonstrate that fewer than one in fifty selected CpGs is reproducibly identified across independent cohorts, and that identity-level concordance is insufficient to constitute a shared genomic signature. Cross-cohort replicability must therefore be evaluated at the level of effect structure rather than probe identity, as formalised in the following subsections.

5.4.2 Effect-Direction Concordance

The absence of strong identity-level overlap does not preclude the existence of a shared differential methylation structure: independently selected panels may still converge on the same underlying drift axis if the corresponding effect sizes are concordant across cohorts. To evaluate this possibility, signed mean methylation differences $\Delta\beta_j = \bar{\beta}_j^{\text{Adj}} - \bar{\beta}_j^{\text{Norm}}$ were computed for each CpG j in the pairwise intersection of the respective 5,000-CpG panels. Concordance was assessed through two complementary measures: the Spearman rank correlation ρ of signed effect sizes, which captures both magnitude and direction, and the directional concordance rate

$$c = \hat{\mathbb{P}}\left(\text{sign}(\Delta\beta_j^{(A)}) = \text{sign}(\Delta\beta_j^{(B)})\right), \quad (5.18)$$

estimated with Wilson confidence intervals, which isolates the directional component independently of magnitude. The distinction between ρ and c is methodologically important: a low signed correlation combined with high directional concordance indicates attenuated but direction-preserving effects, whereas negative ρ accompanied by $c \ll 0.5$ is diagnostic of systematic polarity inversion rather than mere amplitude attenuation. Both quantities are visualised jointly in Figure 5.8.

GSE69914–GSE225845: shared drift axis The HM450–EPIC pair exhibits strong positive rank correlation ($\rho = 0.605$) and near-complete directional alignment ($c = 0.980$, Wilson 95% CI excluded from 0.5 by a wide margin). Together, these values indicate that the two datasets not only agree on the direction of each locus-specific shift but also preserve the relative ordering of effect magnitudes. This constitutes the clearest evidence of a shared epigenetic drift axis across independent cohorts, despite the known platform heterogeneity between the HM450 and EPIC arrays.

GSE287331: systematic polarity inversion Pairs involving GSE287331 display a qualitatively distinct pattern. The Spearman correlations are weakly negative ($\rho = -0.042$ with GSE69914; $\rho = -0.420$ with GSE225845), and directional concordance falls substantially below 0.5 ($c = 0.357$ and $c = 0.125$, respectively). The value $c = 0.125$ in particular implies that only one in eight selected CpGs shares the same Adjacent–Normal polarity between GSE225845 and GSE287331, a pattern incompatible with stochastic disagreement and indicative of a systematic reversal of the dominant drift direction. Crucially, this inversion is not attributable to the absence of a discriminative signal in GSE287331: the intra-dataset analyses of Section 5.3 demonstrated that this cohort yields the strongest and most geometrically coherent Normal–Adjacent separation of the three (silhouette = 0.3442; outlier A/N ratio = 13.3). Rather, the inversion reflects a structural asymmetry in the orientation of the drift manifold, likely arising from a combination of platform-specific probe-type composition, the distinct tumour-proximity sampling design of GSE287331 (TPxA axis), and residual confounding by cohort-specific batch structure.

5.4.3 Magnitude-Stratified Concordance

The signed and directional analyses of Section 5.4.2 characterise the global behaviour of effect sizes across the pairwise intersection of selected panels. A complementary question is whether concordance is uniform across the effect magnitude spectrum or, rather, whether it is concentrated in loci exhibiting the strongest drift signals — a distinction with direct implications for the biological interpretability of cross-cohort consistency. To address this, CpGs in the pairwise intersection were stratified by

decile of absolute effect magnitude $|\Delta\beta|$, computed independently in each dataset. For each decile bin, the fraction of loci whose effect direction agreed across cohorts was computed, yielding a magnitude-stratified directional concordance profile.

For the GSE69914–GSE225845 pair, directional concordance is high and stable across all deciles, with no systematic attenuation in lower-magnitude strata. Crucially, the top decile — comprising CpGs with the largest $|\Delta\beta|$ in both datasets — exhibits the highest concordance, indicating that the strongest signals are preferentially shared. This behaviour supports the interpretation that the cross-cohort agreement between these two datasets is not an artefact of magnitude-insensitive directional alignment, but reflects a genuinely shared structure concentrated at loci with the most pronounced Adjacent–Normal differential methylation.

Pairs involving GSE287331 display a markedly different pattern. Directional concordance remains depressed across all decile strata, with no recovery in the high-magnitude bins. This rules out the scenario in which directionality disagreement would be confined to low-amplitude, noisy loci — as would be expected under pure stochastic disagreement — and instead confirms that the inversion of effect orientation is structural and affects loci of all effect magnitudes, including those most strongly discriminative within GSE287331. Taken together with the negative signed correlations reported in Section 5.4.2, this evidence positions the GSE287331 discordance firmly in the *directional inversion* regime rather than the *structural heterogeneity* regime of Section 5.5: the selected loci are not uninformative, but their dominant drift axis is systematically rotated relative to that of the other two cohorts.

5.4.4 Cross-Dataset Transferability

Set-level overlap and effect-level concordance characterise the structure of the selected panels on their own terms. A more stringent criterion for cross-cohort replicability is *functional transferability*: whether the differential methylation signal inferred in one cohort retains discriminative power when applied to an independent dataset, without any retraining or adaptation.

Formally, for each ordered pair of datasets (A, B) , a linear scoring function $f^{(A)} : \mathbb{R}^{|S_{AB}|} \rightarrow \mathbb{R}$ was defined on the intersection S_{AB} of the respective 5,000-CpG panels, using the cohort- A effect sizes $\Delta\beta^{(A)}$ as fixed coefficients:

$$f^{(A)}(\mathbf{x}_i^{(B)}) = \sum_{j \in S_{AB}} \Delta\beta_j^{(A)} \cdot x_{ij}^{(B)}, \quad (5.19)$$

where $\mathbf{x}_i^{(B)}$ denotes the methylation profile of sample i in dataset B . This zero-shot transfer procedure isolates the intrinsic portability of the drift structure itself, independent of classifier fitting. Discriminative performance was evaluated via the area under the ROC curve (AUC), with values near 0.5 indicating absence of

transfer, values substantially above 0.5 indicating directionally consistent transfer, and values substantially below 0.5 indicating systematic inversion — i.e. the scoring function trained on cohort A assigns *higher* scores to Normal than to Adjacent samples in cohort B .

GSE69914–GSE225845 Bidirectional transfer between the HM450 and EPIC cohorts was evaluated by training a univariate $\Delta\beta$ ranking rule in one dataset and applying it directly to the other. Specifically, given an effect-size vector $\Delta\beta$ estimated in the source cohort, each sample x in the target cohort was assigned the discriminative score

$$s(x) = \sum_{j \in \mathcal{S}} \Delta\beta_j x_j, \quad (5.20)$$

where $\mathcal{S}$ denotes the selected CpG panel and x_j the methylation level of locus j . No re-estimation or adaptation to the target distribution was performed. Discriminative performance was quantified via the area under the receiver operating characteristic curve (AUC), defined as

$$\text{AUC} = \int_0^1 \text{TPR}(t) d\text{FPR}(t), \quad (5.21)$$

where TPR and FPR denote the true- and false-positive rates across classification thresholds. For the binary Normal–Adjacent problem, this quantity is equivalently expressed as

$$\text{AUC} = \mathbb{P}(s(X_A) > s(X_N)), \quad (5.22)$$

that is, the probability that a randomly selected Adjacent sample receives a higher score than a randomly selected Normal sample. The AUC takes values in $[0,1]$, where 0.5 corresponds to chance-level discrimination, values above 0.5 indicate discriminative ability in the correct direction, and 1 corresponds to perfect separation. Values below 0.5 reflect systematic inversion of the ranking rule.

The resulting AUC was 0.776 when training on GSE69914 and testing on GSE225845, and 0.794 in the reverse direction. Both values are substantially above chance, indicating that the shared drift axis identified through concordance analysis translates into genuine discriminative transferability. These results provide the strongest cross-cohort evidence in the present study that the GSE69914–GSE225845 pair captures a reproducible and functionally portable epigenetic drift structure.

Transfers involving GSE287331 All four directed transfer pairs involving GSE287331 yield AUC values in the range $[0.017, 0.226]$, substantially and consistently below chance level. Sub-random AUC is not a failure of discriminability per se, but rather a signature of systematic directional inversion: the scoring function derived from one cohort actively misranks samples in the other, assigning higher

Table 5.2: Number of CpGs retained at each step of the feature selection pipeline (training set).

Pipeline step	CpGs in GSE69914	CpGs in GSE225845	CpGs in GSE287331
Post-preprocessing space	251,483	530,683	486,725
After Edgar variability filter	151,380	396,644	286,000
After M -value SD trimming	148,352	388,711	280,280
After stability + effect-size filter	15,000	15,000	15,000
After correlation pruning	12,313	6,453	6,300
After constrained diversification	5,000	5,000	5,000

drift scores to the tissue class with lower methylation deviation. This outcome is the functional counterpart of the near-zero or negative sign concordance ($c = 0.125$ – 0.357) and negative Spearman correlations ($\rho = -0.042$ to -0.420) reported in Section 5.4.2, and provides definitive evidence that the polarity inversion observed at the descriptive level has concrete consequences for predictive transfer.

Interpretation The transferability results complete the multi-level concordance portrait of the three selected panels. The GSE69914–GSE225845 pair satisfies all criteria for the *shared drift structure* regime: significant set enrichment, high directional concordance, magnitude-stratified agreement concentrated at strong-effect loci, and bidirectional AUC well above chance. Pairs involving GSE287331 are consistent with the *directional inversion* regime: non-trivial set enrichment, systematically inverted effect polarity across all magnitude strata, and sub-random transfer performance. These asymmetries have direct methodological consequences for the downstream modelling strategy developed in Chapter 6: naive cross-cohort transfer from or to GSE287331 is not only non-informative but actively misleading, motivating the need for orientation-aware or cohort-adaptive learning approaches.

5.5 Feature Selection Outcomes and Modelling Implications

The feature selection framework developed in this chapter addresses the core statistical challenge posed by genome-wide DNA methylation data in the HDLSS regime: identifying a compact, stable, and biologically coherent CpG panel capable of characterising the Normal–Adjacent epigenetic transition across independent cohorts.

The pipeline operates as a sequence of strictly train-only dimensionality reduction stages, each targeting a distinct source of noise or redundancy in the methylation feature space. Table 5.2 summarises the progressive reduction from the post-preprocessing spaces — ranging from 251,483 to 530,683 CpGs depending

on platform — to the final panels of $K = 5,000$ loci retained after variability filtering, stability-based ranking, correlation pruning, and constrained genomic diversification. The resulting panels exhibit several desirable structural properties. Directional stability scores are concentrated near unity across all three cohorts, indicating that the selected CpGs express reproducible Adjacent–Normal polarity across resampling splits. Correlation-based redundancy pruning eliminates co-methylated blocks, yielding non-redundant representatives of independent epigenetic signals. Genomic diversification enforces broad chromosomal and regulatory-context coverage, reducing the risk that downstream models capture locus-specific artefacts rather than system-level drift patterns.

The inter-dataset evaluation of Section 5.4 reveals that cross-cohort replicability is partial and asymmetric. GSE69914 and GSE225845 satisfy all criteria for the *shared drift structure* regime: their panels are mutually enriched beyond random expectation, exhibit near-complete directional concordance ($c = 0.980$), and support bidirectional zero-shot transfer (AUC = 0.776 and 0.794). GSE287331, by contrast, occupies the *directional inversion* regime: its panel captures a robust but systematically inverted drift axis, yielding sub-random transfer performance (AUC $\in [0.017, 0.226]$) against both partner cohorts. This asymmetry has a direct methodological consequence. Linear $\Delta\beta$ signatures, however stable within a cohort, do not constitute a universally portable representation of the Normal–Adjacent transition: their discriminative orientation is modulated by platform architecture, sampling design, and residual batch structure in ways that naive cross-cohort transfer cannot accommodate.

The structural properties and cross-cohort limitations of the final CpG panels jointly define the design space for the downstream modelling strategy. Within each cohort, the selected panels provide a low-dimensional, stability-filtered, and biologically grounded input representation for supervised classification of Normal versus Adjacent tissue. Chapter 6 develops and evaluates a suite of classifiers — ranging from regularised linear models to ensemble methods — applied independently within each dataset, assessing the degree to which the selected CpG signatures support within-cohort discriminability. The cross-cohort asymmetries documented here, however, motivate an analysis that goes beyond standard intra-dataset evaluation. Where drift orientation is preserved — as between GSE69914 and GSE225845 — cross-dataset transfer of trained models is a principled and tractable objective. Where systematic inversion is present, as with GSE287331, orientation-aware or cohort-adaptive strategies are required. Chapter 7 addresses this challenge through CpGPT-based representation learning, investigating whether higher-order methylation organisation, not captured by linear $\Delta\beta$ projections, supports a more generalisable encoding of the Normal–Adjacent epigenetic transition across cohorts with divergent drift orientations.

Chapter 6

Predictive Modelling, Constrained Selection and Biological Interpretation

6.1 Introduction

Chapter 5 established a statistically stable and structurally diversified CpG panel of fixed cardinality ($K = 5,000$) for each dataset. The proposed pipeline integrated empirical variability screening, biologically weighted stability selection, correlation-based redundancy pruning, and genomic diversification, ensuring that the resulting signatures are reproducible within each cohort and structurally constrained by design. The inter-dataset evaluation of Section 5.4 further revealed that cross-cohort replicability is partial and asymmetric: GSE69914 and GSE225845 share a concordant drift orientation and support bidirectional zero-shot transfer, whereas GSE287331 exhibits a systematically inverted drift axis, yielding sub-random transfer performance against both partner cohorts.

These structural properties define the design space for the analyses developed in the present chapter. The selected CpG panels are characterised by statistical robustness and biological diversification, but their capacity to discriminate Normal from Adjacent tissue under supervised modelling has not yet been assessed. Leakage-controlled feature selection guarantees that the chosen loci are statistically stable and non-redundant; it does not, by itself, imply that they encode sufficient discriminative information for a classifier to operate reliably on held-out data.

The objectives of this chapter are threefold. First, intra-dataset classification performance is quantified to verify whether the selected CpG panels support accurate discrimination of Normal and Adjacent tissue within each cohort. A linear

Support Vector Machine is adopted as the primary classifier for GSE225845 and GSE287331, a k -Nearest Neighbours introduced for GSE69914 to assess the degree of linear separability. Second, cross-dataset transferability is examined by training on one cohort and evaluating directly on another, with no information from the target dataset entering the training procedure. This zero-leakage design quantifies the extent to which the inferred epigenetic drift axis generalises across independent cohorts, and whether performance degradation is systematically associated with the directional concordance structure documented in Chapter 5. Third, a constrained combinatorial compression step is introduced to derive a compact 50-CpG subset through a mixed-integer linear programming formulation inspired by the classical 0–1 knapsack paradigm. This step investigates whether predictive structure and biological diversity can be jointly preserved under severe dimensional reduction, subject to explicit constraints on cardinality, chromosomal distribution, correlation structure, and methylation directionality.

Finally, the biological coherence of the compressed panels is evaluated through CpG-to-gene mapping, cross-referenced against the COSMIC Cancer Gene Census (CGC, version 103, GRCh38) [85, 86], and pathway-level enrichment analysis. This interpretive step closes the loop between statistical discrimination and functional relevance, assessing whether the loci selected by a purely data-driven optimisation converge on genomic regions with established roles in cancer-related epigenetic deregulation.

Together, these analyses progress from statistically robust feature selection to supervised evaluation, external validation, and structured biological interpretation. The limitations that emerge — in particular the failure of linear $\Delta\beta$ signatures to generalise across cohorts with divergent drift orientations — directly motivate the representation-learning approach developed in Chapter 7.

6.2 Predictive Framework

Following feature selection, each dataset is restricted to the corresponding panel of $K = 5,000$ CpGs identified in Chapter 5. Supervised baseline classifiers are applied independently within each cohort to assess whether the structurally stable and biologically diversified signatures encode sufficient discriminative information to separate Normal and Adjacent tissue.

6.2.1 Intra-Dataset Supervised Modelling

Data partitioning and leakage control For each dataset, samples are partitioned into stratified training and test sets (80/20 split), preserving the Normal–Adjacent class proportions. Standardisation is fitted exclusively on the training set and then applied to the held-out test set without re-estimation. Feature selection

is performed strictly within training folds in Chapter 5, ensuring complete leakage control.

Linear Support Vector Machine A linear Support Vector Machine (SVM) is adopted as the primary classifier [87]. Given training data $\{(x_i, y_i)\}_{i=1}^n$ with $x_i \in \mathbb{R}^p$ and $y_i \in \{-1, +1\}$, the primal formulation is

$$\min_{w, b, \xi} \quad \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \xi_i \quad \text{s.t.} \quad y_i(w^\top x_i + b) \geq 1 - \xi_i, \quad \xi_i \geq 0.$$

The decision function is $f(x) = w^\top x + b$, with class assignment given by $\text{sign}(f(x))$. In this work, the model is trained on standardised M -values with fixed hyperparameters ($C = 0.5$, `max_iter` = 10,000) to prioritise robustness over tuning.

k -Nearest Neighbours (GSE69914 only) For GSE69914, a k -Nearest Neighbours (kNN) classifier is also evaluated as a local non-parametric baseline [88]. For a query x , let $\mathcal{N}_k(x)$ denote the set of the k closest training samples under Euclidean distance. The predicted label is

$$\hat{y}(x) = \text{mode}\{y_i : x_i \in \mathcal{N}_k(x)\}.$$

The method is applied on standardised features with fixed $k = 5$.

Performance metrics AUC is computed as defined in Chapter 5. Balanced accuracy, MCC, precision, recall, and F1-score are reported to complement AUC under potential class imbalance. Balanced accuracy is

$$\text{BAcc} = \frac{1}{2} \left(\frac{\text{TP}}{\text{TP} + \text{FN}} + \frac{\text{TN}}{\text{TN} + \text{FP}} \right),$$

while MCC is

$$\text{MCC} = \frac{\text{TP} \cdot \text{TN} - \text{FP} \cdot \text{FN}}{\sqrt{(\text{TP} + \text{FP})(\text{TP} + \text{FN})(\text{TN} + \text{FP})(\text{TN} + \text{FN})}}.$$

Finally,

$$\text{Precision} = \frac{\text{TP}}{\text{TP} + \text{FP}}, \quad \text{Recall} = \frac{\text{TP}}{\text{TP} + \text{FN}}, \quad \text{F1} = \frac{2 \text{ Precision Recall}}{\text{Precision} + \text{Recall}}.$$

6.2.2 Cross-Dataset Evaluation Protocol

This subsection formalises the cross-cohort transferability framework adopted to assess whether the inferred epigenetic drift axis generalises across independent datasets.

Feature space alignment Cross-dataset evaluation requires strict alignment of the feature space between source and target cohorts. For each ordered pair of datasets (A, B) , the CpG panel is restricted to the intersection of the effective 5,000-CpG sets used during intra-dataset training in cohort A . Only CpGs present in both datasets and retained in the final feature selection of the source cohort are considered.

Column ordering is explicitly harmonised to ensure identical feature indexing across matrices. No re-selection or feature re-ranking is performed on the target cohort. All transformations (including the $\beta \rightarrow M$ mapping, when required) are applied consistently across datasets prior to modelling.

Train-on-A / Test-on-B design External validation is implemented under a strict train-on- A / test-on- B paradigm. The classifier is trained exclusively on the training split of cohort A , using the aligned CpG space described above. No samples from cohort B enter the fitting procedure at any stage.

Feature standardisation is fitted on the source training set and subsequently applied, without re-estimation, to the target dataset. A linear Support Vector Machine is trained on the source cohort and probability calibration is performed via sigmoid calibration (three-fold internal cross-validation) within the source training data only. The calibrated decision function is then evaluated directly on cohort B .

No hyperparameter tuning, retraining, or adaptation is performed on the target cohort. This design enforces complete separation between source and target distributions and quantifies true zero-shot transferability.

Evaluation criteria Discriminative performance is quantified using the same metrics adopted for intra-dataset validation, including AUC, balanced accuracy, and Matthews correlation coefficient. Performance is reported both on the internal test split of cohort A and on the external cohort B .

Under distributional shift, degradation in AUC or balanced accuracy is interpreted as evidence of limited transferability of the learned separating hyperplane. In particular, systematic performance inversion is examined in relation to the directional concordance structure identified in Chapter 5. External performance below random expectation ($\text{AUC} \approx 0.5$) is considered indicative of either weak shared signal or reversed drift orientation between cohorts.

6.2.3 Constrained Subset Optimisation: Knapsack Formulation

This subsection formalises the compression of the 5,000-CpG panel into a fixed-cardinality subset of size $K = 50$ through a constrained combinatorial optimisation framework. The formulation is inspired by the classical 0–1 knapsack problem originally discussed in the context of discrete-variable extremum problems by Dantzig [89], and extended here to incorporate multiple structural and biological constraints.

While the classical knapsack maximises a linear utility function under a single capacity constraint, the present formulation adopts a multi-constraint mixed-integer linear programming (MILP) structure tailored to genomic panel design. Similar optimisation-based approaches have been explored in marker and SNP panel selection problems [90, 91], supporting the use of binary decision variables and explicit structural constraints in genomics.

Objective function Let $\mathcal{J}$ denote the set of 5,000 CpGs retained after feature selection. For each CpG $j \in \mathcal{J}$, a binary decision variable $x_j \in \{0,1\}$ indicates inclusion in the compressed panel.

Each CpG is assigned a composite score

$$v_j = w_{\text{stat}} \tilde{s}_j + w_{\Delta M} \tilde{d}_j + w_{\text{stab}} \tilde{r}_j + w_{\text{COSMIC}} \mathbf{1}_{\{j \in \text{CGC}\}},$$

where $\tilde{s}_j$, $\tilde{d}_j$, and $\tilde{r}_j$ denote normalised statistical ranking components (classifier weight magnitude, absolute effect size, and stability score, respectively), and $\mathbf{1}_{\{j \in \text{CGC}\}}$ indicates whether the gene mapped to CpG j appears in the COSMIC Cancer Gene Census [85].

The optimisation objective is

$$\max_x \sum_{j \in \mathcal{J}} v_j x_j + \mu \Phi(x),$$

where $\Phi(x)$ encodes diversification incentives and $\mu \geq 0$ controls the trade-off between pure statistical ranking and structural balance. A sweep over μ is performed to identify a stable trade-off region.

Cardinality and structural constraints The optimisation is subject to the following constraints:

- **Fixed cardinality:**

$$\sum_{j \in \mathcal{J}} x_j = K.$$

- **Chromosome caps:** for each chromosome c ,

$$\sum_{j \in \mathcal{J}_c} x_j \leq \alpha_{\text{chr}} K,$$

where $\mathcal{J}_c$ denotes CpGs located on chromosome c .

- **Genomic context limits:** upper bounds are imposed on the proportion of CpGs belonging to specific contexts (e.g., OpenSea, Island).
- **Correlation exclusion:** for CpG pairs (i, j) with $|r_{ij}| \geq 0.85$,

$$x_i + x_j \leq 1,$$

preventing the simultaneous selection of highly correlated loci.

- **Gene-level diversity:** linking variables ensure that no gene is represented by more than a fixed number of CpGs, and that a minimum number of distinct genes is selected.
- **Directional balance:** a minimum number of hypermethylated loci is enforced to prevent polarity collapse.

These constraints extend the classical single-capacity knapsack formulation [89] into a structured multi-constraint MILP suitable for genomic panel compression.

Optimisation strategy The resulting problem is solved using a mixed-integer linear programming solver (CBC backend via PuLP). Given the moderate dimensionality ($|\mathcal{J}| = 5,000$), exact optimisation is computationally feasible.

To assess stability of the solution with respect to the diversification parameter μ , a parameter sweep is conducted and the Pareto trade-off curve between statistical score and diversification component is analysed. The final subset is selected at the empirical knee point of this curve.

The formulation therefore combines classical binary combinatorial optimisation [89] with recent optimisation-based approaches to marker panel design in genomics [90, 91], yielding a compact CpG panel that jointly preserves predictive relevance and biological structure.

6.2.4 Gene Set and Functional Enrichment Analysis

This subsection evaluates the biological coherence of the CpG subsets selected through supervised modelling and constrained optimisation. The objective is to determine whether loci identified purely through statistical and structural criteria converge on genes with established roles in cancer biology.

CpG-to-gene mapping CpG loci are annotated to genes using the Illumina EPIC manifest (GRCh38 assembly), relying on the `UCSC_RefGene_Name` field. When multiple gene symbols are associated with a single CpG, all mapped genes are retained. Gene symbols are harmonised to official HGNC nomenclature prior to downstream analysis.

Let $\mathcal{G}$ denote the set of unique genes mapped from the selected CpG subset. Gene-level analyses are conducted on $\mathcal{G}$ after removal of duplicated mappings.

Enrichment testing Functional coherence is assessed using over-representation analysis (ORA). Let $\mathcal{U}$ denote the background universe of genes mapped from the full 5,000-CpG panel of the corresponding cohort. For a given reference gene set $\mathcal{S}$, the enrichment of $\mathcal{G}$ in $\mathcal{S}$ is evaluated via the hypergeometric test:

$$p = \sum_{k=k_0}^{|\mathcal{G}|} \frac{\binom{|\mathcal{S}|}{k} \binom{|\mathcal{U}|-|\mathcal{S}|}{|\mathcal{G}|-k}}{\binom{|\mathcal{U}|}{|\mathcal{G}|}},$$

where $k_0 = |\mathcal{G} \cap \mathcal{S}|$ denotes the observed overlap.

Multiple hypothesis testing correction is performed using the Benjamini–Hochberg procedure to control the false discovery rate (FDR). Adjusted q -values below 0.05 are considered significant.

Reference databases External biological knowledge is incorporated through curated gene sets. In particular, selected genes are cross-referenced against the COSMIC Cancer Gene Census (CGC) database (version 103, GRCh38 assembly) [85], which catalogues genes with documented roles in oncogenesis.

Additional curated gene lists relevant to field cancerisation and early epigenetic deregulation are considered when interpreting enrichment results. The integration of these external resources enables assessment of whether statistically derived CpG panels converge on genes implicated in tumour initiation, chromatin regulation, and cancer-associated signalling pathways.

6.3 Intra-Dataset Results

6.3.1 Dataset GSE69914

6.3.2 Dataset GSE225845

6.3.3 Dataset GSE287331

6.4 Inter-Dataset Results

6.4.1 Cross-Cohort Transferability of the 5,000-CpG Panel

External validation results across all train/test combinations.

6.4.2 Directional Concordance and Performance Degradation

Analysis of performance variation in relation to inter-dataset $\Delta\beta$ concordance.

6.4.3 Cross-Dataset Behaviour of the 50-CpG Subset

Evaluation of compressed signature under distributional shift.

6.5 Synthesis and Implications

This section summarises predictive findings, identifies structural limitations of linear $\Delta\beta$ -based signatures, and motivates the transition toward representation-learning approaches in the subsequent chapter.

Chapter 7

CpGPT - estensione cross dataset

Chapter 8

Conclusion and future work

CONCLUSIONI E PROSPETTIVE DI FUTURO

A systematic sensitivity analysis of the feature selection pipeline hyperparameters — including the directional stability threshold, the biological prior weight λ , the correlation redundancy cutoff τ_c , and the diversification target K — would provide empirical grounding for the design choices adopted in Chapter 5. Given the computational cost of re-running stability selection across the full hyperparameter grid, this analysis would benefit from a reduced resampling scheme or surrogate evaluation on a representative subset of CpGs.

Appendix A

Filtering lists

A.1 Probe Filtering Resources

This appendix documents the five external annotation resources adopted for technical probe filtering across all three datasets. Each resource targets distinct classes of measurement artefacts; together they define the union-based exclusion criterion applied in Section 4.2.3. Table A.1 summarises the technical categories covered by each resource.

A.1.1 Naeem *et al.* (2014)

Naeem *et al.* [53] proposed a structured filtering framework for the HumanMethylation450 array, designed to reduce false discoveries by sequentially discarding probes on the basis of hybridisation specificity and genomic integrity. The framework excludes probes that multi-map or cross-hybridise to additional genomic loci, probes overlapping repetitive elements (LINE, SINE, Alu), and probes overlapping regions harbouring insertion–deletion polymorphisms (INDELs). For SNP-affected probes, a distinction is drawn between variants at the interrogated CpG or single-base extension site—which directly compromise the methylation measurement and are therefore excluded—and variants located nearby but not interfering in bisulfite space, which are retained. The resulting exclusion set thus removes probes affected by cross-reactivity, structural polymorphisms, and measurement-critical SNPs.

A.1.2 Chen *et al.* (2013)

Chen *et al.* [54] empirically identified probes in the 450K array exhibiting off-target hybridisation or overlap with common SNPs. In the present work, only the cross-reactive probe list is used. This catalogue enumerates approximately 29,000 multi-mapping probes whose methylation signal cannot be unambiguously

Table A.1: Technical artefact categories covered by each filtering resource.

Category	Naeem	Chen	Pidsley	Zhou	McCartney
Cross-hybridisation	Yes	Yes	Yes	<code>MASK_mapping</code>	Yes
SNP at CpG	Yes	—	Yes	<code>MASK_snp5</code> , <code>MASK_extBase</code>	—
Nearby SNP	Conditional	—	—	—	—
Structural variant	Yes	—	—	—	—
Non-CpG probes	—	—	Yes	<code>MASK_nonCG</code>	Yes

attributed to a single genomic locus, and whose inclusion would therefore introduce spatially unresolved signal into the feature matrix.

A.1.3 Pidsley *et al.* (2016)

Pidsley *et al.* [32] provide a platform-level assessment of the EPIC array, releasing annotated probe lists that flag loci where technical bias arises from design- or genome-related sources. Three categories are relevant here. First, CpG-targeting probes with sequence homology to additional genomic loci, which yield non-unique hybridisation and potentially ambiguous β estimates. Second, non-CpG-targeting probes affected by analogous off-target issues, excluded due to limited interpretability in CpG-centric analyses. Third, probes overlapping common genetic variation at three critical positions: the interrogated CpG itself (polymorphic CpG, directly affecting the presence of the CpG dinucleotide); the single-base extension site of Type I probes (perturbing extension chemistry and dye-intensity ratios); and within the probe body (reducing binding affinity or altering hybridisation kinetics for common variants).

A.1.4 Zhou *et al.* (2016)

Zhou *et al.* [31] released a unified probe annotation for both 450K and EPIC arrays encoding technical reliability through a set of binary mask columns (`MASK_*`). The masks applied in this work flag probes with low or inconsistent mapping quality (`MASK_mapping`); Type I probes carrying a SNP at the extension base that induces a colour-channel switch (`MASK_typeINextBaseSwitch`); probes whose extension base is inconsistent with the expected channel or CpG context (`MASK_extBase`); probes with non-unique 30-bp 3' subsequences susceptible to cross-hybridisation (`MASK_sub30.copy`); probes overlapping an SNP within ± 5 bp of the interrogated CpG (`MASK_snp5.common`); and probes overlapping SNPs with global minor allele frequency above 1% (`MASK_snp5.GMAF1p`). The composite `MASK_general` field, which integrates mapping, SNP, and cross-reactivity criteria, is used as the primary filter for general-purpose exclusion. Probes overlapping RepeatMasker regions (`MASK_rmsk15`) are not excluded, consistent with standard practice.

A.1.5 McCartney *et al.* (2016)

McCartney *et al.* [55] assessed probe design artefacts on the EPIC array and published supplementary tables enumerating cross-hybridising CpG-targeting probes and cross-hybridising non-CpG-targeting probes. These lists complement the Pidsley and Zhou resources by providing an independent empirical characterisation of off-target hybridisation on the EPIC platform.

The combined application of these resources ensures that downstream analyses operate on a probe set with maximised hybridisation specificity and minimal confounding by common genetic variation.

Appendix B

Extension of an Empirically Driven Variability-Based Filtering Framework to Breast Tissue

B.1 Motivation and Scope

The variability-based filtering strategy embedded in the preprocessing pipeline of this thesis builds directly on the method proposed by [77], who demonstrated that a substantial fraction of CpG sites interrogated by the Illumina HumanMethylation450K array are effectively non-variable within a given tissue context and can therefore be removed prior to downstream analysis without loss of discriminative information. The original study was conducted on three healthy reference tissues — blood, buccal epithelial cells, and placenta — and the resulting non-variable CpG lists were shown to be robust across preprocessing pipelines and normalisation strategies.

The present appendix documents the extension of this framework to breast tissue. The extension pursues two objectives: (i) to assess whether the non-variability criterion, as originally defined, generalises to a cancer-relevant tissue under increased biological heterogeneity; and (ii) to provide a general, reproducible protocol for incorporating additional tissues into the reference framework in future studies. The analysis operates exclusively on histologically normal samples to preserve comparability with the original reference tissues and to estimate baseline methylation stability in the absence of disease-related confounding. Importantly, this appendix does not modify the non-variability definition of [77]: a CpG is

classified as non-variable if its inter-quantile range $r_{\beta,j} = Q_{0.90}(\beta_j) - Q_{0.10}(\beta_j) < 0.05$. The contribution is therefore evaluative and integrative rather than definitional.

B.2 General Framework for Cross-Tissue Extension

Let $\mathcal{T} = \{T_1, \dots, T_k\}$ denote a set of reference tissues, each represented by one or more independent methylation datasets from normal samples. For each tissue T_ℓ , the tissue-specific non-variable CpG set is defined as:

$$\mathcal{N}_\ell(\tau) = \{j \in \{1, \dots, p\} : Q_{0.90}(\beta_j^{(\ell)}) - Q_{0.10}(\beta_j^{(\ell)}) < \tau\}, \quad (\text{B.1})$$

where $\tau = 0.05$ is the threshold inherited from [77]. The cross-tissue invariant core — CpGs non-variable across all reference tissues simultaneously — is given by the intersection:

$$\mathcal{N}_{\text{core}}(\tau) = \bigcap_{\ell=1}^k \mathcal{N}_\ell(\tau). \quad (\text{B.2})$$

When a new tissue T_{k+1} is incorporated, the invariant core is updated as:

$$\mathcal{N}'_{\text{core}}(\tau) = \mathcal{N}_{\text{core}}(\tau) \cap \mathcal{N}_{k+1}(\tau), \quad (\text{B.3})$$


which is strictly more conservative than the original core: $\mathcal{N}'_{\text{core}}(\tau) \subseteq \mathcal{N}_{\text{core}}(\tau)$. The validation of any extension requires verifying three properties before the updated core can be considered methodologically sound:

1. **Selection stability:** the non-variable set $\mathcal{N}_{k+1}(\tau)$ must be robust to dataset composition, i.e. not driven by any single cohort (Step A, Section B.3.1).
2. **Threshold optimality:** the chosen τ must lie at a favourable stability–dimensionality trade-off point in the new tissue context (Step B, Section B.3.2).
3. **Empirical coherence:** the selected CpGs must exhibit methylation and variability profiles consistent with the non-variability interpretation, ruling out selection artefacts (Steps C–D, Sections B.3.3–B.3.4).

This three-step validation protocol is general and applies identically to any future tissue extension.

B.3 Cross-Tissue Overlap Structure

Breast tissue was introduced alongside the three original reference tissues of [77]. Figure B.1 reports the overlap structure of tissue-specific non-variable CpG sets across all four tissues.



(a) Pairwise and higher-order overlaps of tissue-specific non-variable CpG sets. (b) Quantitative intersection sizes for key subsets.

Figure B.1: Global overlap structure of non-variable CpGs across blood, buccal epithelial cells, placenta, and breast tissue.


The four-way invariant core $\mathcal{N}'_{\text{core}}(0.05)$ comprises 29,989 CpGs — a substantial shared subset whose persistence despite the inclusion of a cancer-relevant tissue suggests that at least a component of CpG non-variability reflects stable, context-independent properties of the methylation landscape rather than purely tissue-specific regulatory programmes. The breast-exclusive non-variable set (5,023 CpGs) is the smallest tissue-specific component, consistent with the expectation that breast tissue introduces additional heterogeneity relative to the original healthy tissue panel.

B.3.1 Step A: Stability of Non-Variable CpG Selection

The first validation step assesses whether the breast-tissue non-variable set $\mathcal{N}_{\text{breast}}(\tau)$ is robust to dataset composition. A leave-one-dataset-out (LODO) strategy is adopted: for each held-out dataset $\mathcal{D}^{(-s)}$, the non-variable set is recomputed on the remaining datasets and compared to the full-reference set via the retention rate:

$$\rho^{(-s)} = \frac{|\mathcal{N}_{\text{breast}}^{(-s)}(\tau) \cap \mathcal{N}_{\text{breast}}(\tau)|}{|\mathcal{N}_{\text{breast}}(\tau)|}, \quad (\text{B.4})$$

where $\mathcal{N}_{\text{breast}}^{(-s)}(\tau)$ denotes the non-variable set estimated without dataset s . The retention rates reported in Figure B.2a are consistently high across all three held-out datasets, demonstrating that $\mathcal{N}_{\text{breast}}(\tau)$ is not an artefact of any individual cohort but reflects a stable property of the underlying methylation variability structure in breast tissue. This result satisfies the first validation criterion stated in Section B.2 and is a necessary condition for interpreting non-variability as a genuine empirical property rather than a dataset-specific anomaly.



(a) Retention rates under the LODO protocol across held-out datasets (GSE69914, GSE225845, GSE287331).

(b) Mean Jaccard similarity $J(\tau)$ and mean non-variable CpG count as functions of the threshold.

Figure B.2: Assessment of robustness and trade-off between stability and dimensionality of non-variable CpG sets.

B.3.2 Step B: Stability–Dimensionality Trade-Off

Having established stability at $\tau = 0.05$, Step B evaluates whether this threshold represents an optimal compromise between selection reproducibility and dimensionality reduction in the breast tissue setting. The mean Jaccard similarity across LODO folds and the mean number of non-variable CpGs are computed as functions of $\tau \in [0.01, 0.10]$:

$$J(\tau) = \frac{1}{S} \sum_{s=1}^S \frac{|\mathcal{N}_{\text{breast}}^{(-s)}(\tau) \cap \mathcal{N}_{\text{breast}}(\tau)|}{|\mathcal{N}_{\text{breast}}^{(-s)}(\tau) \cup \mathcal{N}_{\text{breast}}(\tau)|}. \quad (\text{B.5})$$

As shown in Figure B.2b, $J(\tau)$ increases rapidly for small τ and saturates before the Edgar threshold, while the CpG count continues to grow approximately linearly. At $\tau = 0.05$, stability is already at its near-maximum value and the marginal gain from increasing the threshold further is negligible relative to the dimensionality cost incurred. This empirically validates the original threshold of [77] in the breast tissue context and provides a data-driven justification for its retention within the extended framework.

B.3.3 Step C: Methylation Distribution of Non-Variable CpGs

Step C characterises the methylation profiles of the selected non-variable CpGs by comparing their β -value distribution to a random background sample of array CpGs, computed on normal breast samples pooled across datasets. Non-variable




Figure B.3: Characterization of the selected non-variable CpG set in terms of methylation distribution and baseline risk behavior.

CpGs in breast tissue are strongly concentrated at extreme β values, consistent with constitutive hypo- or hypermethylated states. This pattern replicates the findings of [77] in healthy tissues and extends them to a cancer-relevant context, supporting the interpretation that non-variability reflects stable regulatory configurations rather than artefacts of tissue composition or disease-related heterogeneity. The enrichment of non-variable CpGs in CpG islands and constitutively silenced promoter regions, documented in the original study, is consistent with this bimodal profile [77, 78].

B.3.4 Step D: Baseline Variability Profile

Step D provides quantitative validation of the filtering criterion by examining the empirical cumulative distribution function (ECDF) of $r_{\beta,j}$ across four CpG subsets: the four-way shared non-variable core $\mathcal{N}'_{\text{core}}(0.05)$, the breast-specific non-variable set $\mathcal{N}_{\text{breast}}(0.05)$, the breast-exclusive component, and a random background set. The ECDFs in Figure B.3b show a clear separation: the non-variable subsets are sharply concentrated at low $r_{\beta,j}$ values, with the majority of CpGs well below $\tau = 0.05$, whereas the background distribution spans a substantially wider range. This confirms that the selected CpGs are empirically distinct in their variability profile and not merely defined as non-variable by construction. The consistency of this pattern across all non-variable subsets — including the breast-exclusive component — further supports the robustness of the extended framework.

B.4 Cohort-Resolved Characterisation of Breast-Tissue Non-Variability

The four-step validation protocol described in Section B.2 establishes the methodological soundness of the non-variability criterion at the level of pooled breast data. The present section extends this characterisation to the individual-cohort level, pursuing two complementary objectives: (i) to assess whether the filtering behaviour is consistent across independent datasets, each with distinct sample compositions and phenotypic configurations; and (ii) to characterise the biological identity of the invariant CpG set through genomic contextual enrichment analysis and cross-phenotype variability concordance. Together, these analyses situate the breast-tissue extension within a broader biological framework and provide the empirical grounding required for its use as a preprocessing step in Chapter 5.

B.4.1 Empirical Variability Spectrum Across Independent Breast Cohorts

To evaluate the consistency of the non-variability criterion across datasets, the inter-quantile range statistic $r_{\beta,j} = Q_{0.90}(\beta_j) - Q_{0.10}(\beta_j)$ was computed independently for each of the three breast cohorts (GSE69914, GSE225845, GSE287331). For each dataset, the analysis was carried out under two phenotypic configurations: (i) Normal + Adjacent (N+A), which includes only histologically normal and adjacent-normal samples, and (ii) Normal + Tumor (N+T), which additionally incorporates tumor samples. This design allows direct assessment of whether malignant tissue alters the global variability spectrum of CpG loci and, consequently, the proportion of sites classified as non-variable under the Edgar threshold $\tau = 0.05$.

Figure B.4 reports the empirical density of r_{β} under the N+T configuration for each dataset, with vertical markers indicating the three candidate thresholds $\tau \in \{0.01, 0.02, 0.05\}$. Several structural features are common to all three cohorts. The distribution is strongly right-skewed, with a pronounced mass concentrated near $r_{\beta} \approx 0$, indicating that a substantial fraction of CpG sites exhibit minimal within-group dispersion regardless of sample composition. The strict thresholds $\tau = 0.01$ and $\tau = 0.02$ remove only a small minority of loci, whereas the canonical $\tau = 0.05$ eliminates a sizeable but controlled proportion — 77,687 CpGs in GSE69914, 103,622 in GSE225845, and 168,663 in GSE287331 — consistent with the dimensionality reduction reported in the pooled analysis.

A biologically informative pattern emerges from the comparison between the two phenotypic configurations. Under N+T, the Edgar filter removes fewer CpGs than under N+A: the greater methylation contrast between normal and tumor samples elevates r_{β} across a larger fraction of loci, reducing the proportion that

falls below $\tau = 0.05$. By contrast, the N+A configuration — whose analogous plots are reported in Chapter 5 — yields a higher count of non-variable CpGs, reflecting the subtler methylation differences between histologically normal tissue and adjacent-normal samples. This asymmetry is consistent with the concept of field cancerization, whereby tissue adjacent to a tumour undergoes epigenetic alterations that, while not yet producing overt histological change, partially converge towards the tumour methylation profile. Under this interpretation, the reduced variability observed in N+A relative to N+T is not a statistical artefact but reflects a genuine biological phenomenon: adjacent tissue shares a greater proportion of its methylation landscape with normal tissue than tumor tissue does, resulting in a larger set of CpG sites that appear non-variable within the combined sample.

B.4.2 Contextual Enrichment Profile of Invariant CpGs

To characterise the genomic architecture of CpGs classified as non-variable under the Edgar threshold, a contextual enrichment analysis was performed using Illumina manifest annotations. Each probe was mapped to its corresponding gene-feature category (e.g., TSS200, TSS1500, 5'UTR, 1stExon, Body, 3'UTR, Intergenic), and enrichment was quantified as fold-change relative to the full set of interrogated CpGs within the dataset.

Let $\mathcal{U}$ denote the universe of CpGs and $\mathcal{N}_{\text{breast}}(0.05)$ the invariant subset identified in the N+A configuration. For each annotation category g , enrichment was computed as:


$$\text{FC}_g = \frac{\frac{|\mathcal{N}_{\text{breast}}(0.05) \cap g|}{|\mathcal{N}_{\text{breast}}(0.05)|}}{\frac{|\mathcal{U} \cap g|}{|\mathcal{U}|}}. \quad (\text{B.6})$$

Figure B.5 reports the fold-change profile for the dominant gene-feature categories.


Across datasets, invariant CpGs show consistent enrichment in proximal promoter regions (particularly TSS200 and 1stExon) and relative depletion in gene bodies and distal regions. The baseline reference (fold-change = 1) indicates the expected frequency under random sampling from the array; deviations above this threshold reflect preferential localisation of low-variability CpGs within regulatory domains.

This pattern is coherent with the bimodal methylation behaviour described in Section B.3.3, where invariant CpGs were shown to concentrate at constitutive hypo- or hypermethylated states. Promoter-associated CpG islands are known to exhibit tight regulatory control and reduced dispersion in normal tissue, providing a biologically plausible explanation for the observed enrichment.


Importantly, the enrichment structure is stable across cohorts, indicating that genomic localisation of invariant CpGs reflects intrinsic regulatory constraints



(a) GSE69914



(b) GSE225845



(c) GSE287331

Figure B.4: Distribution of r_β computed on the Normal+Tumor comparison for each dataset. Vertical dashed lines indicate the selected threshold.




Figure B.5: Fold-enrichment of genomic context categories among invariant CpGs relative to the background distribution in each dataset. The dashed line indicates enrichment equal to background (fold = 1).

rather than dataset-specific artefacts.

B.4.3 Phenotype-Resolved Variability Concordance Analysis

Section B.4.1 established that the aggregate variability spectrum is qualitatively preserved when tumor samples are incorporated. The present section quantifies this concordance at the level of individual CpG sites by directly comparing r_β values computed under the N+A and N+T configurations.

For each CpG j , the pair $(r_{\beta,j}^{(NA)}, r_{\beta,j}^{(NT)})$ was examined and rank concordance was quantified via the Spearman correlation coefficient. Figure B.6 displays the resulting scatter plots together with the identity line $y = x$ for each dataset. The Spearman coefficients are $\rho \approx 0.845$ (GSE69914), $\rho \approx 0.881$ (GSE225845), and $\rho \approx 0.933$ (GSE287331), indicating strong to very strong rank preservation of CpG-level variability across phenotypic configurations.

Several structural features emerge from the scatter plots. The bulk of CpG loci cluster tightly around the identity line, confirming that relative variability ordering is largely maintained when tumor samples are included. A minority of loci deviate upward from the diagonal, corresponding to CpGs whose dispersion increases in the presence of tumor tissue; by contrast, very few loci exhibit a systematic decrease in variability under N+T. This asymmetry is consistent with the expectation that malignant transformation introduces epigenetic heterogeneity rather than convergence, and corroborates the finding in Section B.4.1 that tumor inclusion primarily affects loci in the intermediate variability range.

The high rank concordance implies that the non-variability criterion is largely phenotype-robust: CpGs classified as low-variability under the N+A configuration remain predominantly low-variability when tumor samples are incorporated. This result supports the interpretation that breast-tissue non-variability reflects stable




Figure B.6: Spearman correlation between CpG-level $\Delta\beta$ values computed in the Normal+Adjacent and Normal+Tumor contrasts for each dataset. The diagonal line represents identity.

structural properties of methylation regulation, with tumor-associated heterogeneity manifesting as localised deviations superimposed on an otherwise preserved variability backbone.

B.4.4 Invariant-Set Partitioning Across Phenotypic Combinations

The analyses in Sections B.4.1– B.4.3 establish that non-variability in breast tissue is broadly stable across both cohorts and phenotypic configurations. The present section integrates the breast-tissue invariance structure with the reference framework of [77] by performing a set-theoretic decomposition that explicitly accounts for the effect of tumor inclusion.

Let $\mathcal{B} = \mathcal{N}_{\text{Blood}} \cap \mathcal{N}_{\text{Buccal}} \cap \mathcal{N}_{\text{Placenta}}$ denote the universal invariant set across the three reference tissues of [77]. The breast-tissue non-variable sets $\mathcal{N}_{\text{breast}}(0.05)$ were estimated separately under the N+A and N+T configurations for each cohort. The intersection of $\mathcal{B}$ with these two sets defines three biologically interpretable invariant classes:

Core invariants $\mathcal{B} \cap \mathcal{N}_{\text{breast}}^{(NA)}(0.05) \cap \mathcal{N}_{\text{breast}}^{(NT)}(0.05)$ CpGs that are non-variable across all reference tissues and remain invariant in breast tissue irrespective of tumor inclusion. These loci constitute the most conserved component of the cross-tissue epigenetic backbone.

Conditionally variable universal CpGs $\mathcal{B} \cap \mathcal{N}_{\text{breast}}^{(NA)}(0.05) \setminus \mathcal{N}_{\text{breast}}^{(NT)}(0.05)$ Loci that are invariant across all reference tissues and in normal breast tissue but become variable when tumor samples are incorporated. This subset identifies CpGs whose


stability is sensitive to malignant transformation and that may be of interest for cancer-specific methylation studies.

Breast-specific invariants $(\mathcal{N}_{\text{breast}}^{(NA)}(0.05) \cap \mathcal{N}_{\text{breast}}^{(NT)}(0.05)) \setminus \mathcal{B}$ CpGs that are consistently non-variable within breast tissue under both phenotypic configurations but are not part of the universal cross-tissue invariant core. These loci reflect tissue-contextual stability that is specific to breast epithelium.


Figure B.7 visualises this set-theoretic decomposition for each dataset as a three-way Venn diagram. The quantitative breakdown is consistent across cohorts. The core invariant class comprises approximately 27,000–28,000 CpGs per dataset (GSE69914: 28,241; GSE225845: 26,542; GSE287331: 27,577), representing a highly conserved epigenetic backbone that is robust to both tissue context and phenotypic variation. The conditionally variable universal class is markedly smaller (GSE69914: 2,010; GSE225845: 1,353; GSE287331: 496), confirming that only a minor fraction of cross-tissue invariants is susceptible to tumor-induced destabilisation. The breast-specific invariant class is the largest component in datasets with denser sampling (GSE69914: 138,256; GSE225845: 76,223; GSE287331: 46,788), reflecting the greater CpG coverage of the corresponding platforms and the broader set of tissue-contextual constraints captured at higher array density.

Several biological observations emerge from this decomposition. First, the invariant core is not globally eroded by cancer: tumor-associated variability acts locally on a bounded subset of CpGs, while the structural invariance backbone is preserved. Second, the breast-specific invariant class is predominantly shared between the N+A and N+T configurations, confirming that tissue-contextual stability is itself phenotype-robust. Third, the hierarchical organisation of the invariant partition — a context-independent core layer, a tissue-specific invariant layer, and a phenotype-sensitive component — mirrors the layered regulatory architecture of the methylome, in which CpG islands at constitutively regulated promoters are progressively supplemented by tissue-specific and stimulus-responsive methylation programmes [77, 78].


Taken together, the analyses in Section B.4 provide a cohort-resolved and phenotype-stratified characterisation of breast-tissue non-variability that complements the pooled validation in Section B.2. The convergence of results across independent datasets, genomic annotation categories, and phenotypic configurations establishes that the Edgar-based filtering strategy is a biologically grounded dimensionality reduction tool whose behaviour in breast tissue is consistent, interpretable, and robust to the heterogeneity introduced by malignant transformation.



(a) GSE69914



(b) GSE225845



(c) GSE287331

Figure B.7: Overlap between universal invariants and breast-specific invariant sets derived from the Normal+Adjacent and Normal+Tumor contrasts in each dataset.

B.5 Positioning Within the Thesis and Generalisability

The extended framework described in this appendix provides the empirical grounding for the variability filter applied in Chapter 5. The four validation steps collectively establish that the non-variability criterion of [77] is stable, threshold-optimal, and biologically coherent when applied to breast tissue — necessary conditions for its use as a preprocessing step in the present study. The cohort-resolved characterisation presented in Section 5.3 further demonstrates that these properties hold independently within each dataset and are robust to the phenotypic heterogeneity introduced by tumor inclusion, consolidating the methodological foundations of the filter across the full range of analytical configurations considered in Chapter 5.

In the current analysis, datasets are processed independently and the non-variability threshold is estimated within each cohort separately, rather than enforcing a single shared CpG list across datasets. This design is motivated by two complementary considerations. First, it preserves dataset-specific methylation structure and avoids premature cross-cohort harmonisation that could mask genuine biological heterogeneity between sample populations. Second, the cohort-level validation reported in Section B.4 — in particular the consistency of the variability spectrum, the enrichment profile, and the phenotype-resolved concordance across GSE69914, GSE225845, and GSE287331 — ensures that applying the threshold independently within each cohort does not introduce arbitrary inter-dataset discordance, but rather instantiates a shared population-level filtering principle in each data context separately.

The general framework formalised in Section B.2 — update rule (Equation B.3), LODO stability assessment (Equation B.4), and Jaccard-based threshold analysis (Equation B.5) — is structurally independent of tissue type and platform density. It therefore remains applicable to future tissue extensions and to higher-density arrays such as the Illumina EPIC platform, whose expanded CpG coverage approximately doubles that of the 450K array [65]. In this respect, the breast-tissue extension documented here serves a dual purpose: it validates the criterion in a cancer-relevant context, and it demonstrates the operability of the general protocol for incorporating new tissues into the reference framework — a property of direct relevance for future studies seeking to apply variability-based filtering beyond the tissue panel considered here.

Appendix C

Plot cap6

Appendix D

Age Estimation

Bibliography

- [1] Douglas Hanahan and Robert A. Weinberg. «Hallmarks of Cancer: The Next Generation». In: *Cell* 144.5 (2011), pp. 646–674. DOI: 10.1016/j.cell.2011.02.013 (cit. on pp. 2, 3).
- [2] Douglas Hanahan and Robert A. Weinberg. «The Hallmarks of Cancer». In: *Cell* 100.1 (2000), pp. 57–70. DOI: 10.1016/S0092-8674(00)81683-9 (cit. on pp. 2, 3).
- [3] National Cancer Institute. *BRCA Gene Changes: Cancer Risk and Genetic Testing*. Accessed October 2025. 2024. URL: <https://www.cancer.gov/about-cancer/causes-prevention/genetics/brca-fact-sheet> (cit. on p. 3).
- [4] K. P. M. Suijkerbuijk, M. J. Fackler, S. Sukumar, C. H. van Gils, T. van Laar, E. van der Wall, M. Vooijs, and P. J. van Diest. «Methylation is less abundant in BRCA1-associated compared with sporadic breast cancer». In: *Annals of Oncology* 19 (Nov. 2008), pp. 1870–1874. DOI: 10.1093/annonc/mdn409 (cit. on pp. 3, 14).
- [5] T. E. Bartlett, I. Evans, A. Jones, J. E. Barrett, S. Haran, D. Reisel, K. Papaikonomou, L. Jones, C. Herzog, N. Pashayan, B. M. Simoes, R. B. Clarke, D. G. Evans, T. S. Ghezelayagh, S. Ponandai-Srinivasan, N. R. Boggavarapu, P. G. Lalitkumar, S. J. Howell, R. A. Risques, A. F. Rådestad, L. Dubeau, K. Gemzell-Danielsson, and M. Widschwendter. «Antiprogesterins reduce epigenetic field cancerization in breast tissue of young healthy women». In: *Genome Medicine* 14.1 (June 2022). DOI: 10.1186/s13073-022-01063-5 (cit. on pp. 3, 14).
- [6] Adrian P. Bird. «DNA methylation patterns and epigenetic memory». In: *Genes & Development* 16.1 (2002), pp. 6–21. DOI: 10.1101/gad.947102.
- [7] Rafael A. Irizarry, Christine Ladd-Acosta, Bo Wen, Zhijin Wu, Carolina Montano, Patrick Onyango, Hengmi Cui, Kevin Gabo, Michael R. Aryee, Peiying Ji, Ie-Ming Shih, James B. Potash, Andrew P. Feinberg, and Margaret A. Taub. «The human colon cancer methylome shows similar hypo- and

- hypermethylation at conserved tissue-specific CpG island shores». In: *Nature Genetics* 41.2 (2009), pp. 178–186. DOI: 10.1038/ng.298 (cit. on p. 4).
- [8] Peter A. Jones. «Functions of DNA methylation: islands, start sites, gene bodies and beyond». In: *Nature Reviews Genetics* 13.7 (2012), pp. 484–492. DOI: 10.1038/nrg3230 (cit. on pp. 5, 62).
- [9] Melanie Ehrlich. «DNA methylation in cancer: too much, but also too little». In: *Oncogene* 21 (2002), pp. 5400–5413. DOI: 10.1038/sj.onc.1205651.
- [10] Andrew E. Teschendorff, Usha Menon, Aleksandra Gentry-Maharaj, Susan J. Ramus, Dirk J. Weisenberger, Hui Shen, Mihaela Heravi-Moussavi, and et al. «Age-dependent DNA methylation of genes that are suppressed in stem cells is a hallmark of cancer». In: *Genome Research* 20.4 (2010), pp. 440–446. DOI: 10.1101/gr.103606.109 (cit. on p. 6).
- [11] Steve Horvath. «DNA methylation age of human tissues and cell types». In: *Genome Biology* 14.10 (2013), R115. DOI: 10.1186/gb-2013-14-10-r115 (cit. on p. 6).
- [12] Andrew E. Teschendorff, Menelaos Kyrgiou, Wayne Widschwendter, and Martin Widschwendter. «Epigenetic variability in cells of normal cytology is associated with the risk of future morphological transformation». In: *Genome Medicine* 2.11 (2010), p. 81. DOI: 10.1186/gm200 (cit. on p. 6).
- [13] Andrew E. Teschendorff, Allison Jones, and Martin Widschwendter. «Epigenetic variability in cells of normal cytology is associated with the risk of future morphological transformation». In: *Genome Medicine* 11.1 (2019), pp. 1–14. DOI: 10.1186/s13073-019-0667-3 (cit. on p. 6).
- [14] A. E. Teschendorff and C. L. Relton. «Statistical and integrative system-level analysis of DNA methylation data». In: *Nature Reviews Genetics* 19 (2018), pp. 129–147 (cit. on pp. 7, 8, 54).
- [15] Andrew E. Teschendorff, Christiana Zhuang, and Martin Widschwendter. «Independent surrogate variable analysis to deconvolve confounding factors in large-scale microarray profiling studies». In: *Genome Biology* 15.12 (2014), p. 539. DOI: 10.1186/s13059-014-0539-z (cit. on pp. 7, 8).
- [16] P. Yousefi, K. Huen, R. Aguilar Schall, A. Decker, E. Elboudwarej, H. Quach, L. Barcellos, and N. Holland. «Considerations for Normalization of DNA Methylation Data by Illumina 450K BeadChip Assay in Population Studies». In: *Epigenetics* 8.11 (Aug. 2013), pp. 1141–1152. DOI: 10.4161/epi.26037 (cit. on pp. 10, 39).

- [17] National Center for Biotechnology Information (NCBI). *Gene Expression Omnibus (GEO)*. Public functional genomics data repository supporting MIAME-compliant submissions and providing tools to query and download curated experiments. NCBI. n.d. URL: <https://www.ncbi.nlm.nih.gov/geo/> (cit. on p. 11).
- [18] National Center for Biotechnology Information (NCBI). *GSE69914: Genome wide DNA methylation profiling of normal breast, normal adjacent and breast cancer tissue*. Gene Expression Omnibus (GEO). 2015. URL: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE69914> (cit. on pp. 11, 42).
- [19] National Center for Biotechnology Information (NCBI). *GSE225845: High neighborhood deprivation impacts DNA methylation and gene expression in cancer-related genes*. Gene Expression Omnibus (GEO). 2023. URL: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE225845> (cit. on pp. 11, 44).
- [20] National Center for Biotechnology Information (NCBI). *GSE287331: DNA methylation patterns in breast cancer, paired benign tissue from ipsilateral and contralateral breast, and healthy controls*. Gene Expression Omnibus (GEO). 2025. URL: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE287331> (cit. on p. 12).
- [21] P. Du, X. Zhang, C. Huang, N. Jafari, W. A Kibbe, L. Hou, and S. M Lin. «Comparison of Beta-value and M-value methods for quantifying methylation levels by microarray analysis». In: *BMC Bioinformatics* 11.1 (2010), p. 587. DOI: 10.1186/1471-2105-11-587 (cit. on pp. 12, 40, 43, 49, 58).
- [22] Andrew E. Teschendorff, Francesco Marabita, Matthias Lechner, Thomas Bartlett, David Tegner, Jonathan Gomez-Cabrero, and Stephan Beck. «A beta-mixture quantile normalization method for correcting probe design bias in Illumina Infinium 450k DNA methylation data». In: *Bioinformatics* 29.2 (2013), pp. 189–196. DOI: 10.1093/bioinformatics/bts680. URL: <https://academic.oup.com/bioinformatics/article/29/2/189/199637> (cit. on pp. 12, 16, 27, 38, 42, 44, 48).
- [23] B. P. de Almeida, G. Richard, H. Dalla-Torre, C. Blum, L. Hexemer, P. Pandey, S. Laurent, C. Rajesh, M. Lopez, A. Laterre, M. Lang, U. Sahin, K. Beguir, and T. Pierrot. «A Multimodal Conversational Agent for DNA, RNA and Protein Tasks». In: *Nature Machine Intelligence* (2025). DOI: 10.1038/s42256-025-01047-1 (cit. on p. 12).
- [24] A. V. Sokolov and H. B. Schiöth. «Decoding Depression: A Comprehensive Multi-cohort Exploration of Blood DNA Methylation Using Machine Learning and Deep Learning Approaches». In: *Translational Psychiatry* 14.1 (2024). DOI: 10.1038/s41398-024-02992-y (cit. on p. 13).

- [25] M. Ehrlich. «DNA methylation in cancer: Too much, but also too little». In: *Oncogene* 21 (Aug. 2002), pp. 5400–5413. DOI: 10.1038/sj.onc.1205651 (cit. on pp. 5, 14, 15, 25).
- [26] A. Bird. «DNA methylation patterns and epigenetic memory». In: *Genes & Development* 16.1 (Jan. 2002), pp. 6–21. DOI: 10.1101/gad.947102 (cit. on pp. 3, 14, 15).
- [27] A. E. Teschendorff, Y. Gao, A. Jones, M. Ruebner, M. W. Beckmann, D. L. Wachter, P. A. Fasching, and M. Widschwendter. «DNA methylation outliers in normal breast tissue identify field defects that are enriched in cancer». In: *Nature Communications* 7 (Jan. 2016), p. 10478. DOI: 10.1038/ncomms10478 (cit. on pp. 14–16, 30, 32, 33).
- [28] National Cancer Institute. *BRCA Gene Changes: Cancer Risk and Genetic Testing*. <https://www.cancer.gov/about-cancer/causes-prevention/genetics/brca-fact-sheet>. Accessed October 2025. Oct. 2024 (cit. on p. 14).
- [29] J. Maksimovic, L. Gordon, and A. Oshlack. «SWAN: Subset-Quantile Within Array Normalization for Illumina Infinium HumanMethylation450 Bead-Chips». In: *Genome Biology* 13.6 (2012), R44. DOI: 10.1186/gb-2012-13-6-r44 (cit. on pp. 15, 33, 38, 40, 42–44).
- [30] *Infinium MethylationEPIC v1.0 B5 Manifest File*. https://support.illumina.com/array/array_kits/infinium-methylationepic-beadchip-kit/downloads.html. Illumina, accessed 2025 (cit. on p. 18).
- [31] W. Zhou, P. W. Laird, and H. Shen. «Comprehensive characterization, annotation and innovative use of Infinium DNA methylation BeadChip probes». In: *Nucleic Acids Research* 45.4 (2017), e22. DOI: 10.1093/nar/gkw967 (cit. on pp. 18, 27, 32, 33, 39, 94).
- [32] R. Pidsley, E. Zotenko, T. J. Peters, M. G. Lawrence, G. P. Risbridger, P. Molloy, S. Van Djik, B. Muhlhausler, C. Stirzaker, and S. J. Clark. «Critical evaluation of the Illumina MethylationEPIC BeadChip microarray for whole-genome DNA methylation profiling». In: *Genome Biology* 17 (2016), p. 208. DOI: 10.1186/s13059-016-1066-1 (cit. on pp. 18, 27, 32, 33, 39, 94).
- [33] L. Chen, R. Liu, Z.-P. Liu, M. Li, and K. Aihara. «Detecting early-warning signals for sudden deterioration of complex diseases by dynamical network biomarkers». In: *Scientific Reports* 2 (2012), p. 342. DOI: 10.1038/srep00342 (cit. on p. 18).
- [34] J. West, G. Bianconi, S. Severini, and A. E. Teschendorff. «On dynamic network entropy in cancer». In: *arXiv:1202.3015* (2018). URL: <https://arxiv.org/abs/1202.3015> (cit. on p. 18).

- [35] J. Liu, D. Ding, J. Zhong, and R. Liu. «Identifying the critical states and dynamic network biomarkers of cancers based on network entropy». In: *Journal of Translational Medicine* 20.254 (2022). DOI: 10.1186/s12967-022-03445-0 (cit. on p. 18).
- [36] P. Wang. «Network biology: Recent advances and challenges». In: *Gene & Protein in Disease* 1.2 (2022), p. 101. DOI: 10.36922/gpd.v1i2.101 (cit. on p. 23).
- [37] S. R. Dennis, T. Tsukioki, G. Cottone, W. Zhou, P. A. Ganz, M. E. Sehl, Y. Luo, S. A. Khan, and S. Clare. «DNA methylation patterns in breast cancer, paired benign tissue from ipsilateral and contralateral breast, and healthy controls». In: *Breast Cancer Research* 27.1 (2025), p. 103. DOI: 10.1186/s13058-025-02057-y (cit. on pp. 25, 48).
- [38] J. Maksimovic, B. Phipson, and A. Oshlack. «A cross-package Bioconductor workflow for analysing methylation array data». In: *F1000Research* 5 (2017), p. 1281. DOI: 10.12688/f1000research.8839.3 (cit. on p. 25).
- [39] J. Yates, H. Schaufelberger, R. Steinacher, P. Schär, K. Truninger, and V. Boeva. «DNA-methylation variability in normal mucosa: A field cancerization marker in patients with adenomatous polyps». In: *Journal of the National Cancer Institute* 116.6 (2024), pp. 974–982. DOI: 10.1093/jnci/djae016 (cit. on pp. 31, 32).
- [40] S. Zhang, X. Xiao, Y. Yi, X. Wang, L. Zhu, Y. Shen, D. Lin, and C. Wu. «Tumor initiation and early tumorigenesis: Molecular mechanisms and epigenetic drivers». In: *Signal Transduction and Targeted Therapy* (2024). DOI: 10.1038/s41392-024-01848-7 (cit. on pp. 31, 32).
- [41] J.-P. Fortin, A. Labbé, M. Lemire, B. W. Zanke, T. J. Hudson, E. J. Fertig, C. M. T. Greenwood, and K. D. Hansen. «Functional normalization of 450k methylation array data improves replication in large cancer studies». In: *Genome Biology* 15.12 (2014), p. 503. DOI: 10.1186/s13059-014-0503-2 (cit. on pp. 32–34).
- [42] W. Timp and A. P. Feinberg. «Cancer as a dysregulated epigenome: The epigenetic progenitor model of cancer». In: *Nature Reviews Cancer* 13.7 (2013), pp. 497–510. DOI: 10.1038/nrc3486 (cit. on p. 33).
- [43] J. T. Leek, R. B. Scharpf, H. Corrada Bravo, D. Simcha, B. Langmead, W. E. Johnson, D. Geman, K. Baggerly, and R. A. Irizarry. «Tackling the widespread and critical impact of batch effects in high-throughput data». In: *Nature Reviews Genetics* 11.10 (2010), pp. 733–739. DOI: 10.1038/nrg2825 (cit. on pp. 33, 34).

- [44] Timothy Triche, Daniel Weisenberger, David Van Den Berg, Peter Laird, and Michael Siegmund. «Low-level processing of Illumina Infinium DNA methylation beadarrays». In: *Nucleic Acids Research* 41.7 (2013), e90. DOI: 10.1093/nar/gkt090 (cit. on pp. 36, 37, 40, 49).
- [45] Jean-Philippe Fortin, Timothy J. Triche Jr, and Kasper D. Hansen. «Pre-processing, normalization and integration of the Illumina HumanMethylationEPIC array with minfi». In: *Bioinformatics* 33.4 (2017), pp. 558–560. DOI: 10.1093/bioinformatics/btw691. URL: <https://academic.oup.com/bioinformatics/article/33/4/558/2666344> (cit. on pp. 36, 40, 49).
- [46] Carol S. Wilhelm-Benartzi, Devin C. Koestler, Margaret R. Karagas, James M. Flanagan, Brock C. Christensen, John K. Wiencke, Karl T. Kelsey, Carmen J. Marsit, and Andres E. Houseman. «Review of processing and analysis methods for DNA methylation array data». In: *British Journal of Cancer* 109.6 (2013), pp. 1394–1402. DOI: 10.1038/bjc.2013.496. URL: <https://www.nature.com/articles/bjc2013496> (cit. on p. 37).
- [47] Zongli Xu, Liang Niu, and Jack A. Taylor. «The ENmix DNA methylation analysis pipeline for Illumina BeadChip and comparisons with seven other preprocessing pipelines». In: *Clinical Epigenetics* 13 (2021), p. 216. DOI: 10.1186/s13148-021-01207-1. URL: <https://clinicalepigeneticsjournal.biomedcentral.com/articles/10.1186/s13148-021-01207-1> (cit. on p. 37).
- [48] S. A. Islam, S. J. Goodman, J. L. MacIsaac, J. Obradović, R. G. Barr, W. T. Boyce, and M. S. Kobor. «Integration of DNA methylation patterns and genetic variation in human pediatric tissues helps inform EWAS design and interpretation». In: *Epigenetics & Chromatin* 12.1 (2019), p. 1. DOI: 10.1186/s13072-018-0245-6 (cit. on pp. 38, 46).
- [49] G. Mansell, T. J. Gorrie-Stone, Y. Bao, M. Kumari, L. S. Schalkwyk, J. Mill, and E. Hannon. «Guidance for DNA methylation studies: Statistical insights from the Illumina EPIC array». In: *BMC Genomics* 20.366 (2019), pp. 1–15. DOI: 10.1186/s12864-019-5761-7 (cit. on pp. 38, 46).
- [50] Ting Wang, Weihua Guan, Jerome Lin, Nadia Boutaoui, Glorisa Canino, Jianhua Luo, Juan Carlos Celedón, and Wei Chen. «A Systematic Study of Normalization Methods for Infinium 450K Methylation Data Using Whole-Genome Bisulfite Sequencing as the Gold Standard». In: *Epigenetics* 10.6 (2015), pp. 536–545. DOI: 10.1080/15592294.2015.1057384 (cit. on pp. 38, 42, 44).

- [51] H. Welsh, C. M. P. F. Batalha, W. Li, K. L. Mpye, N. C. Souza-Pinto, M. S. Naslavsky, and E. J. Parra. «A systematic evaluation of normalization methods and probe replicability using Infinium EPIC 850K methylation data». In: *Clinical Epigenetics* 15.1 (2023), p. 59. DOI: 10.1186/s13148-023-01459-z (cit. on pp. 39, 48).
- [52] J. Liu and K. D. Siegmund. «An evaluation of processing methods for Illumina HumanMethylation450 BeadChip platform». In: *BMC Genomics* 17.469 (2016). DOI: 10.1186/s12864-016-2819-7 (cit. on pp. 39, 48).
- [53] Haroon Naeem, Nicholas C. Wong, Zac Chatterton, Matthew K. H. Hong, John S. Pedersen, Niall M. Corcoran, Christopher M. Hovens, and Geoff Macintyre. «Reducing the Risk of False Discovery Enabling Identification of Biologically Significant Genome-Wide Methylation Status Using the HumanMethylation450 Array». In: *BMC Genomics* 15.1 (2014), p. 51. DOI: 10.1186/1471-2164-15-51 (cit. on pp. 39, 93).
- [54] Yuting A. Chen, Mathieu Lemire, Sanaa Choufani, Daniel T. Butcher, Diana Grafodatskaya, Brent W. Zanke, Steven Gallinger, Thomas J. Hudson, and Rosanna Weksberg. «Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium HumanMethylation450 microarray». In: *Epigenetics* 8.2 (2013), pp. 203–209. DOI: 10.4161/epi.23470 (cit. on pp. 39, 93).
- [55] Daniel L. McCartney, Rosie M. Walker, Stewart W. Morris, Andrew M. McIntosh, David J. Porteous, and Kathryn L. Evans. «Identification of polymorphic and off-target probe binding sites on the Illumina Infinium MethylationEPIC BeadChip». In: *Genomics Data* 9 (2016), pp. 22–24. ISSN: 2213-5960. DOI: <https://doi.org/10.1016/j.gdata.2016.05.012> (cit. on pp. 39, 95).
- [56] Yucheng Wang, Tyler J Gorrie-Stone, Olivia A Grant, Alexandria D Andrayas, Xiaojun Zhai, Klaus D McDonald-Maier, and Leonard C Schalkwyk. «InterpolatedXY: a two-step strategy to normalize DNA methylation microarray data avoiding sex bias». In: *Bioinformatics* 38.16 (June 2022), pp. 3950–3957. ISSN: 1367-4803. DOI: 10.1093/bioinformatics/btac436 (cit. on pp. 40, 43).
- [57] M. D. Robinson, G. K. Smyth, and K. D. Hansen. *IlluminaHumanMethylation450kanno.ilmn12.hg19: Annotation for Illumina’s 450k methylation arrays*. R package version 0.6.0. Bioconductor. 2015. URL: <https://bioconductor.org/packages/IlluminaHumanMethylation450kanno.ilmn12.hg19> (cit. on p. 40).
- [58] Illumina Inc. *Infinium MethylationEPIC BeadChip Manifest File v1.0 B5*. <https://support.illumina.com/>. Accessed: 2026-02. 2016 (cit. on pp. 40, 49).

- [59] Bernard Thienpont, Jessica Steinbacher, Hui Zhao, Flora D’Anna, Anna Kuchnio, Athanasios Ploumakis, Bart Ghesquière, Laurien Van Dyck, Bram Boeckx, Luc Schoonjans, Els Hermans, Frederic Amant, Vessela N. Kristensen, Kian Peng Koh, Massimiliano Mazzone, Mathew L. Coleman, Thomas Carell, Peter Carmeliet, and Diether Lambrechts. «Tumour Hypoxia Causes DNA Hypermethylation by Reducing TET Activity». In: *Nature* 537.7618 (2016), pp. 63–68. DOI: 10.1038/nature19081 (cit. on p. 40).
- [60] L. E. Reinius, N. Acevedo, M. Joerink, G. Pershagen, S. E. Dahlén, D. Greco, C. Söderhäll, J. Kere, and A. Scheynius. «Differential DNA methylation in purified human blood cells: Implications for cell lineage and studies on disease susceptibility». In: *PLOS ONE* 7.7 (2012), e41361. DOI: 10.1371/journal.pone.0041361 (cit. on p. 40).
- [61] Illumina. *Infinium HumanMethylation450 BeadChip: Product Files and Manifest (v1.2)*. Available online: https://support.illumina.com/downloads/infinium_humanmethylation450_product_files.html. Accessed: 2025-12-02. 2011 (cit. on p. 43).
- [62] B. H. Chen and W. Zhou. «mLiftOver: Mapping DNA methylation data across array platforms and genome builds». In: *Bioinformatics* 40.7 (2024), btae423. DOI: 10.1093/bioinformatics/btae423 (cit. on p. 46).
- [63] R. Pidsley, C. C. Y. Wong, M. Volta, K. Lunnon, J. Mill, and L. C. Schalkwyk. «A data-driven approach to preprocessing Illumina 450K methylation array data». In: *BMC Genomics* 14.293 (2013), pp. 1–10. DOI: 10.1186/1471-2164-14-293 (cit. on p. 48).
- [64] P. Yousefi, K. Huen, R. Aguilar Schall, A. Decker, E. Elboudwarej, H. Quach, L. Barcellos, and N. Holland. «Considerations for normalization of DNA methylation data by Illumina 450K BeadChip assay in population studies». In: *Epigenetics* 8.11 (2013), pp. 1145–1155. DOI: 10.4161/epi.26037 (cit. on p. 48).
- [65] R. Pidsley et al. «Critical evaluation of the Illumina MethylationEPIC BeadChip microarray for whole-genome DNA methylation profiling». In: *Genome Biology* 17 (2016), p. 208 (cit. on pp. 53, 108).
- [66] P. Hall, J. S. Marron, and A. Neeman. «Geometric representation of high dimension, low sample size data». In: *Journal of the Royal Statistical Society: Series B* 67 (2005), pp. 427–444 (cit. on pp. 53, 54, 59, 63).
- [67] D. L. Donoho. *High-Dimensional Data Analysis: The Curses and Blessings of Dimensionality*. Tech. rep. Lecture manuscript, AMS Conference on Math Challenges of the 21st Century. Stanford University, 2000 (cit. on pp. 53, 59, 63, 64).

- [68] C. Ambroise and G. J. McLachlan. «Selection bias in gene extraction on the basis of microarray gene-expression data». In: *Proceedings of the National Academy of Sciences* 99 (2002), pp. 6562–6566 (cit. on pp. 53–55, 57, 59).
- [69] P. Smialowski et al. «Pitfalls of supervised feature selection». In: *Bioinformatics* 26 (2010), pp. 440–443 (cit. on pp. 53, 55, 59).
- [70] Kasper D. Hansen, Winston Timp, Héctor Corrada Bravo, Sarven Sabuncian, Ben Langmead, Oliver G. McDonald, Bo Wen, Hao Wu, Yihui Liu, Dinh Diep, Eirikur Briem, Kun Zhang, Rafael A. Irizarry, and Andrew P. Feinberg. «Increased methylation variation in epigenetic domains distinguishes tumour from normal tissue». In: *Nature Genetics* 43.8 (2011), pp. 768–775. DOI: 10.1038/ng.865. URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC3145050/> (cit. on pp. 54, 61–63, 66).
- [71] M. J. Aryee et al. «Minfi: a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays». In: *Bioinformatics* 30 (2014), pp. 1363–1369 (cit. on pp. 54, 63).
- [72] C. Luo et al. «Single-cell methylomes identify neuronal subtypes and regulatory elements in mammalian cortex». In: *Science* 357 (2017), pp. 600–604 (cit. on p. 54).
- [73] T. Fleischer et al. «DNA methylation at enhancers identifies distinct breast cancer lineages». In: *Nature Communications* 8 (2017), p. 1379 (cit. on pp. 54, 61, 66).
- [74] A. Gelman and E. Loken. «The statistical crisis in science». In: *American Scientist* 102 (2014), pp. 460–465 (cit. on p. 55).
- [75] N. Meinshausen and P. Bühlmann. «Stability selection». In: *Journal of the Royal Statistical Society: Series B* 72 (2010), pp. 417–473 (cit. on pp. 55, 60, 61, 69, 73).
- [76] G. Poste. «Bring on the biomarkers». In: *Nature* 469 (2011), pp. 156–157 (cit. on p. 55).
- [77] Rachel D. Edgar, Meaghan J. Jones, Wendy P. Robinson, and Michael S. Kobor. «An Empirically Driven Data Reduction Method on the Human 450K Methylation Array to Remove Tissue Specific Non-Variable CpGs». In: *Clinical Epigenetics* 9.1 (2017), p. 11. DOI: 10.1186/s13148-017-0320-z (cit. on pp. 56–58, 96, 97, 99, 100, 105, 106, 108).
- [78] C. Bock. «Analysing and interpreting DNA methylation data». In: *Nature Reviews Genetics* 13 (2012), pp. 705–719 (cit. on pp. 58, 100, 106).

- [79] Richard Bourgon, Robert Gentleman, and Wolfgang Huber. «Independent filtering increases detection power for high-throughput experiments». In: *Proceedings of the National Academy of Sciences* 107.21 (2010), pp. 9546–9551. DOI: 10.1073/pnas.0914005107. URL: <https://www.pnas.org/doi/10.1073/pnas.0914005107> (cit. on p. 58).
- [80] H. B. Mann and D. R. Whitney. «On a Test of Whether One of Two Random Variables Is Stochastically Larger Than the Other». In: *The Annals of Mathematical Statistics* 18.1 (1947), pp. 50–60. DOI: 10.1214/aoms/1177730491. URL: <https://www.jstor.org/stable/2236101> (cit. on p. 60).
- [81] Ronald Fagin, Ravi Kumar, and D. Sivakumar. «Comparing Top- k Lists». In: *SIAM Journal on Discrete Mathematics* 17.1 (2003), pp. 134–160. DOI: 10.1137/S0895480102412856 (cit. on p. 60).
- [82] Cynthia Dwork, Ravi Kumar, Moni Naor, and D. Sivakumar. «Rank Aggregation Methods for the Web». In: *Proceedings of the 10th International Conference on World Wide Web (WWW '01)*. ACM, 2001, pp. 613–622. DOI: 10.1145/371920.372165 (cit. on p. 60).
- [83] Daniela M. Witten and Robert Tibshirani. «A Framework for Feature Selection in Clustering». In: *Journal of the American Statistical Association* 105.490 (2010), pp. 713–726. DOI: 10.1198/jasa.2010.tm09415 (cit. on pp. 63, 66).
- [84] Jerome H. Friedman, Trevor Hastie, and Robert Tibshirani. «Regularization Paths for Generalized Linear Models via Coordinate Descent». In: *Journal of Statistical Software* 33.1 (2010), pp. 1–22. DOI: 10.18637/jss.v033.i01 (cit. on p. 63).
- [85] Zbyslaw Sondka, Sarah Bamford, Charlotte G. Cole, Sally A. Ward, Elizabeth Dunham, and Simon A. Forbes. «The COSMIC Cancer Gene Census: describing genetic dysfunction across all human cancers». In: *Nature Reviews Cancer* 18.11 (2018), pp. 696–705. DOI: 10.1038/s41568-018-0060-1 (cit. on pp. 84, 87, 89).
- [86] COSMIC Cancer Gene Census. *COSMIC Cancer Gene Census, version 103 (GRCh38)*. <https://cancer.sanger.ac.uk/cosmic/download/cosmic/v103/cancergenecensus>. Wellcome Sanger Institute, accessed February 2026. 2023 (cit. on p. 84).
- [87] Corinna Cortes and Vladimir Vapnik. «Support-Vector Networks». In: *Machine Learning* 20.3 (1995), pp. 273–297. DOI: 10.1007/BF00994018 (cit. on p. 85).
- [88] Thomas M. Cover and Peter E. Hart. «Nearest Neighbor Pattern Classification». In: *IEEE Transactions on Information Theory* 13.1 (1967), pp. 21–27. DOI: 10.1109/TIT.1967.1053964 (cit. on p. 85).

- [89] George B. Dantzig. «Discrete-Variable Extremum Problems». In: *Operations Research* 5.2 (1957), pp. 266–288. DOI: 10.1287/opre.5.2.266 (cit. on pp. 87, 88).
- [90] Xiaoxiang Wu, Ruili Li, Qian Li, et al. «Optimal Design of Low-Density SNP Arrays for Genomic Selection». In: *PLOS ONE* 11.9 (2016), e0161719. DOI: 10.1371/journal.pone.0161719 (cit. on pp. 87, 88).
- [91] Soichiro Nishiyama, Kengo Sato, and Ryutaro Tao. «Integer programming for selecting sets of informative markers». In: *BMC Bioinformatics* 23 (2022), p. 4801. DOI: 10.1186/s12859-022-04801-x (cit. on pp. 87, 88).