

RESEARCH

Open Access



DNA methylation patterns in breast cancer, paired benign tissue from ipsilateral and contralateral breast, and healthy controls

Saya R. Dennis^{1†}, Takahiro Tsukioki^{2†}, Gannon Cottone³, Wanding Zhou^{4,5}, Patricia A. Ganz^{6,7}, Mary E. Sehl^{6,8}, Yuan Luo¹, Seema A. Khan^{3,9*†} and Susan Clare^{3,9*†}

Abstract

Background Epigenetic changes, particularly DNA methylation, are crucial to breast cancer development. Tumor-adjacent normal (AN) tissue frequently serves as a reference for characterizing genomic alterations but is reported to share some characteristics with tumors. However, it is unclear whether AN's epigenetic profiles reflect a predisposition to cancer or a response to the presence of the tumor. We address this gap by systematically comparing methylation profiles of tumor, AN, and matched-benign tissues from both breasts, as well as to healthy donated breast tissue.

Methods We studied four different sample categories from 69 cancer cases: tumor (TU), AN, ipsilateral opposite quadrant (OQ), and contralateral unaffected breast (CUB); and healthy donated breast (HDB) tissue from 182 cancer-unaffected donors. These constitute a “tumor proximity axis” (TPxA): HDB→CUB→OQ→AN→TU. Methylation profiles were assayed using Illumina's Infinium Methylation EPICv1.0 BeadChip. Differential methylation (DM) analysis was conducted, and the significantly DM CpGs were analyzed for enrichment of transcription factor binding sites (TFBS) and other features.

Results Following data processing and quality control, there were 69 TU, 60 AN, 67 OQ, 68 CUB, and 182 HDB samples for analysis. DM analysis showed distinct methylation profiles of TU relative to benign tissues, whereas case-benign tissues were similar to each other but distinct from HDB. Hypomethylated sites in case-benign versus HDB were enriched for TF binding sites of TP63, GATA3, ESR1, PR, AR, NR3C1, and GREB1. TU hypermethylation events were enriched for Polycomb-repressive complex 2 (PRC2) binding, including EZH2, SUZ12, and JARID2, with hypermethylation enrichment for PRC2-related binding motifs in both ER+ and ER- tumors. TU methylation profiles were otherwise highly distinct by ER status: TFBS enrichment of hypomethylation events for hormone receptor-

[†]Saya R. Dennis and Takahiro Tsukioki contributed equally to this work.

[†]Seema A. Khan and Susan Clare contributed equally to this work.

*Correspondence:

Seema A. Khan
s-khan2@northwestern.edu
Susan Clare
susan.clare@northwestern.edu

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

related pathways in ER+ tumors and for hematopoiesis/immune-related pathways in ER- tumors. We found no differential methylation between benign tissues from patients with ER+ vs. ER- tumors.

Conclusions DNA methylation profiles differ profoundly at two points: tumor to case-benign and case-benign to HDB, with clear distinction between ER+ and ER- tumors. Case-benign tissues are not epigenetically “normal”, are similar across both breasts, and do not differ by ER status of paired tumors.

Keywords DNA methylation, Breast cancer, Tumor-adjacent normal tissue, Epigenetic profiling, Illumina methylation EPIC BeadChip, Differential methylation analysis, Hormone-receptor signaling

Background

Epigenetic changes play a crucial role in cancer development [1]. Among these changes, DNA methylation is one of the most significant due to its impact on gene expression and genome stability [2–4]. DNA methylation occurs at sites in the genome where the sequence is cytosine followed by guanine (CpG), and in which cytosine acquires a methyl group at the 5th position of its ring structure. This modification affects transcription, with hyper-methylation generally suppressing transcription and hypo-methylation promoting it [2].

In breast cancer research, studying the epigenetic alterations of the tumor requires a non-cancerous reference. The tumor-adjacent normal tissue, typically defined as the morphologically normal tissue located at least 3 cm from the tumor margin [5], is often used as a reference. However, field cancerization—also known as the field effect or field defect—has been a topic of study for decades. Advances in molecular technologies have identified field cancerization in breast cancer, including changes in telomeres, allelic imbalances, DNA methylation, and gene expression, as extensively reviewed by Danforth 2016 [6] and Gadaleta et al. 2022 [7]. These findings support the concept that tumors arise from normal tissue gradually accumulating genomic aberrations, leading to clonal expansion of genomically unstable cells [2, 8]. Earlier DNA methylation studies were limited to a moderate number of CpG loci due to technological constraints [9, 10]; however, with the development of the HumanMethylation 450 K (HM450) array, studies examining methylation at over 450,000 CpG loci across the genome became possible [5, 11–13]. More recently, the EPIC BeadChip array—surveying nearly twice the CpGs of the HM450—offers a more comprehensive approach.

Breast cancer is heterogeneous, particularly in its subtypes defined by hormone receptor and HER2-neu expression. In previous studies, DNA methylation differences among tumor subtypes have been characterized using the HM450 array [14, 15] but whether the DNA methylation profiles of benign tissues from cancer patients reflect characteristics of their corresponding tumor subtypes remain unexplored. We have employed the more comprehensive EPIC array to fully profile epigenetic differences by estrogen receptor subtype, along

with matched benign tissues at various proximities to the tumor and reference control tissues from unaffected donors. We have then systematically compared DNA methylation profiles across tumors, paired benign ipsilateral and contralateral breast samples, and benign breast samples from women with no breast cancer history to investigate epigenetic changes associated with tumor proximity and examine whether case-benign tissues at different locations vary in their similarity or differences from paired tumors. By incorporating both tumor and non-tumor contexts and considering estrogen receptor status, we demonstrate distinct differences in the methylation profile of benign breast tissue remote from the tumor in cases when compared to tissue from healthy controls; we establish that case-benign tissues from three different locations spanning the affected and unaffected control breast have similar methylation profiles; and we identify specific differences in the transition from adjacent normal to tumor tissue based on the ER status of the tumor. Overall, these data provide valuable insights into the complex landscape of breast cancer epigenetics.

Methods

Sample processing, library construction and methylation profiling

This study was conducted as a collaboration between investigators at Northwestern University (NU) and the University of California, Los Angeles (UCLA). The case samples (tumor and paired benign) were collected at NU, and the unaffected control samples were obtained by UCLA investigators from the Susan G Komen Tissue Bank at the IU Simon Cancer Center under protocols approved by NU and UCLA's Institutional Review Boards (IRB # STU00051134 and 16–000853, respectively). Tissues were obtained at NU between September 2011 and September 2022. A total of 334 patients were enrolled, providing mammary tissue samples without impact on clinical care. From cancer-affected participants (cases), tissue was collected from the tumor (TU), the tumor-adjacent normal (AN; at least 2 cm tumor margin), the opposite quadrant (OQ), and from the contralateral unaffected breast (CUB). A total of 79 matched sets were available, with sufficient tissue from the tumor and all

three benign tissue locations (hereon referred to as “case-benign” tissue categories).

The mammary tissue samples were stored in Optimal Cutting Temperature compound, which was removed prior to homogenization and DNA extraction using QIAGEN’s AllPrep DNA/RNA Kit. DNA quality was assessed using a NanoDrop 1000 Spectrophotometer; only samples meeting a minimum concentration of 50 ng/μL and a volume of 20 μL were retained for analysis. After excluding samples that did not meet these criteria, DNA from 69 patients was available for methylation profiling using Illumina’s Infinium MethylationEPIC v1.0 BeadChip, covering over 850,000 CpG sites. The assay was performed at the University of Minnesota’s Genomics Center. The first batch assayed included TU, OQ, and CUB samples, and the second batch consisted of AN samples.

To minimize chip-to-chip bias during methylation analysis, we randomized sample placement across the 96-well plates. DNA from different tissues of the same individual was randomly assigned to different columns. To validate that this approach effectively eliminated chip-to-chip bias, we placed randomly selected DNA samples in empty wells and confirmed that methylation profiles were consistent across different chips. This validation confirmed the removal of chip-to-chip bias, enabling us to proceed with the analysis.

Normal mammary tissue was initially collected from 192 cancer-unaffected individuals (controls) who had donated tissue to the Susan G Komen Tissue Bank at the IU Simon Cancer Center. DNA methylation profiling was conducted at UCLA as described in Sehl et al. [16]. To match the age distribution between case and control individuals, the 10 youngest patients were excluded, leaving 182 healthy donated breast (HDB) samples for analysis.

Our five tissue categories form an order along the “tumor proximity axis” (TPxA): HDB → CUB → OQ → AN → TU. (Fig. 1a)

Data processing and quality control

The raw data was processed using the SeSAmE software (version 1.20.0) [17] with R (version 4.3.0). To minimize technical variation in methylation signal, we applied SeSAmE’s built-in non-linear dye bias correction and used out-of-band signal for background subtraction and detection *p*-value masking. Samples that had a detection success rate less than 90% were excluded from all following analyses.

Following processing, methylation levels are represented as beta values, which are the ratio of methylated signal intensity to the total signal intensity (methylated plus unmethylated) at each CpG site for each sample. Beta values range from 0 (completely unmethylated) to 1 (completely methylated).

Differential methylation analysis

Differential methylation between tissue categories was determined by fitting a linear model to the methylation levels, with tissue category, age, and BMI included as covariates. All *p*-values were adjusted for false discovery rate (FDR) of 0.05 to correct for multiple testing. A stringent significance threshold was applied, defined as the FDR-adjusted *p*-value < 0.01 and slope < -0.1 or slope > 0.1, using an F-test for each contrast variable, unless specified otherwise. We conducted two sets of four pairwise comparisons: (1) comparing tumor samples to the four benign tissue categories, and (2) comparing tissue categories that are adjacent along the tumor proximity axis.

Probe set enrichment analysis

To understand the trends in differentially methylated (DM) CpG sets, we performed probe set enrichment analysis using built-in functions of the SeSAmE software [17]. We queried our DM CpG sets to 5 categories of databases: transcription factor binding sites (TFBS), regions defined in relation to CpG Islands (CGI), histone modification consensus sites (HMconsensus), chromatin states identified by chromHMM [18], and genes (Ensembl). Enrichment was assessed by determining whether the CpGs in our query set were significantly enriched for specific database category probe sets via Fisher’s exact test. Statistical significance was defined as an FDR-adjusted *p*-value of less than 0.05, and fold-enrichment greater than 1.05.

Results

Data quality assessment and patient/sample summary

Prior to quality control of the methylation signals, we started with 69 TU, 60 AN, 69 OQ, 69 CUB, and 182 HDB samples. (Fig. 1a) Overall, our samples had uniformly high-quality scores, and only two OQ samples and one CUB sample were removed due to signal detection rates below 0.90. (Fig. 1b; Supplementary Fig. 1) The cases and cancer-unaffected donors had a mean age of 52.5 (± 10.0 SD) and 52.3 (± 10.0 SD) respectively, and mean BMI of 28.4 (± 7.1 SD) and 28.8 (± 6.9 SD), respectively. (Fig. 1c) Although age and BMI distributions were comparable between the cancer patients and cancer-free individuals, we correct for these two variables in all differential methylation analyses since these are well-documented confounders [19, 20]. The tumor characteristics of our cases are summarized in Fig. 1d, and the racial distribution of all patients is summarized in Fig. 1e. The estimated fraction of leukocytes was highest in TU and lowest in HDB. (Supplementary Fig. 2)




Fig. 1 Study design and sample characteristics. **(a)** Illustration of the five tissue categories which were collected. **(b)** Sample availability for each tissue category. **(c)** Age and BMI distributions for cancer cases and cancer-unaffected controls. **(d)** Overview of tumor characteristics for cancer cases. **(e)** Racial distribution of the cohort. **(f)** Illustration of the two sets of four pairwise comparisons: (i) tumor versus the four benign tissue categories, and (ii) comparisons between tissue categories neighboring along the tumor proximity axis (TPxA)

The effects of reference selection

Sample clustering via PCA and t-SNE showed that tumor samples were distinctly separated from all case-benign samples (CUB, OQ, and AN), which were in turn highly distinct from HDB samples. (Fig. 2a) When examining the number of differentially methylated CpGs (Fig. 1f – (i)), we observed more hypomethylation events than hypermethylation events in tumor samples compared to any benign tissue category. (Fig. 2b) Additionally, using HDB samples as a reference resulted in nearly double the number of hypomethylation events and moderately fewer hypermethylation events compared to using case-benign samples as a reference. (Fig. 2b)

Differential methylation along the tumor proximity axis

Next, we analyzed the differential methylated (DM) CpGs along the tumor proximity axis (TPxA) (Fig. 1f – (ii)). The numbers of hyper- and hypomethylated CpGs are summarized in Fig. 2c. TU was distinguished from AN by tens of thousands of both hyper- and hypomethylation events. CUB was also found to be highly distinct from HDB, although this difference was dominated by hypomethylation events. The three case-benign tissue categories (CUB, OQ, and AN) were largely similar to each other, with <300 DM CpGs between AN and OQ and no DM between OQ and CUB (Fig. 2c).

The results of the probe set enrichment analysis for DM CpGs along the TPxA are presented in Table 1, with their detailed metrics in Supplementary Tables 1–3. Notably, transcription factor binding site (TFBS) enrichment analysis revealed that hypomethylation events distinguishing CUB from HDB included transcription factor (TF) hits such as TP63 and TP73, along with breast cancer-associated TFs such as ESR1, PR, AR, GREB1, and GATA3. Many of these TFBS enrichment hits were also observed in the hypomethylation events in TU compared to AN, although the enrichment hits were not necessarily driven by the same genes or with the same CpG loci within a gene (Supplementary Fig. 3). The pathway enrichment analysis results for genes driving recurrent TFBS enrichment hits in the hypomethylated sites of CUB vs. HDB and TU vs. AN comparisons are shown in Supplementary Fig. 4. Both hyper- and hypo-methylated sets from CUB vs. HDB were enriched for enhancer regions (Table 1). The proportions of hyper- and hypo-methylated CpGs with their corresponding chromatin states defined by chromHMM for the CUB vs. HDB and TU vs. AN comparisons are shown in Fig. 2d.

TFBS enrichment hits in the hypermethylation events in TU compared to AN included EZH2, SUZ12, and JARID2—key components of the Polycomb-repressive complex 2 (PRC2) and particularly the JARID2-associated variant, known for H3K27me3 activity. The

hypomethylated CpGs in this comparison were enriched for Quiescent sites (Table 1, Supplementary Table 2).

Although methylation profiles in the case-benign tissue categories were similar to each other, the AN vs. OQ comparison yielded 167 hyper- and 78 hypo-methylated CpGs (Fig. 2c). The hypermethylation set showed enrichment for binding targets of CDCA2. The hypomethylated CpGs were highly enriched for CpG Islands as well as active transcription start sites (TssA), with 59/78 (75.6%) CpGs mapping to a CpG Island and 64/78 (82.1%) CpGs mapping to active TssA (Table 1, Supplementary Table 2).

Overall, histone modification consensus site (HMconsensus) enrichments in the hypermethylation set corresponded to repressive marks (H1.0, H3K9me3, H4K20me3, H3K27me3, H2AK119, H3K23me2), whereas those in the hypomethylation set corresponded to activation marks (H3R26Cit, H3K18cr, H3K4me3, H3K23ac).

Although most DM CpGs showed a monotonic increase or decrease in methylation levels along the TPxA, a notable subset (6,663 CpGs) was hypomethylated in case-benign tissues compared to both HDB and TU (Supplementary Fig. 5: Non-monotonic down & up pattern A). This subset was significantly enriched for genes in the HOXA gene cluster (HOXA2-7; Supplementary Fig. 6) and known tumor suppressor genes such as TSC2, EGR1, TP53AIP1, FOXO3 [21–23], RAD51B, and DAXX (Supplementary Table 4). This set was also enriched for consensus sites of H2BK120ub, associated with gene activation, and H4K20me1, associated with both activation and repression (Supplementary Table 4). Additionally, it was significantly enriched for the binding sites of transcription factors TP63, TCF21, TP73, TEAD1, and NEUROG2 (Supplementary Table 4).

Biomarker-related differences in methylation profiles

PCA and t-SNE plots of case samples (CUB, OQ, AN, TU) revealed that TU samples clustered distinctly by ER status, whereas case-benign samples (CUB, OQ, AN) did not show such separation (Fig. 3a). No distinct clustering was observed for HER2 status across any tissue categories (Fig. 3b).

Consistent with this lack of clustering, the comparison between HER2+ and HER2- TU samples identified only 5 differentially methylated (DM) CpGs, all hypomethylated in HER2+ samples. These CpGs mapped to genes ERBB2, MIEN1, OSBPL3, EBF1, and PRKN (Supplementary Table 5).

To thoroughly describe methylation trends related to ER status, we defined two sets of contrasts for our differential methylation (DM) analysis: (a) ER-status-based contrasts (vertical contrasts in Fig. 3c), and (b) contrasts along the TPxA stratified by ER status (horizontal contrasts in Fig. 3c). For ease of notation, we denote the




Fig. 2 Global methylation trends and number of differentially methylated CpGs, **(a)** PCA and t-SNE plots of methylation profiles by tissue category. **(b)** Number of differentially methylated (DM) CpGs in tumor vs. benign tissue comparisons. The second category represents the reference (e.g., in “TU vs HDB,” HDB is the reference category and TU the comparison category). **(c)** Sankey plot depicting the number of DM CpGs in comparisons along the tumor proximity axis (TPxA). As in panel (b), “A vs B” indicates B as the reference category and A as the comparison category. The height of nodes represents the number of DM CpGs, which is also labeled numerically. The “n.d.” represents CpGs that have no significant difference between categories. **(d)** Proportions of chromatin states (as defined by chromHMM) for the DM CpGs in the CUB vs. HDB and TU vs. AN comparisons

Table 1 Enrichment analysis results of DM CpG sets along the TPxA

		TU vs. AN	AN vs. OQ	CUB vs. HDB
TFBS	Hyper	LYL1, LMO2, MEIS1, ZBTB16, FUS, PBX1, PPARG, ETV6, RXRG, JMJD1C, PRDM6, STAT5, NCOR2, RUNX1T1, GATA1, MEF2B, LDB1, TAL1, TCF4, GATA2... (29)	CDCA2, CNOT3	CDCA2, JARID2, TCF21, EZH2, SUZ12, ATF7IP, NEUROG2, SP140, ZFP57, PCGF2, TEAD1
	Hypo	TP63, ZNF217, AHR, TP73, TLE3, GREB1, GRHL2, FOSL2, ESR1, PR, AR, GATA3, TEAD1, NR3C1, HES2	NCAPH2, CNOT3, ARID2, RAG2, CBX8, FAM208A, INTS11, EGR3, ZNF202, ZBTB14, UHRF2, ASXL1, KDM4C, TAF3, LMO1, CCNT2, MAPK1, HCFC1R1, TRIM25, HMGN3... (142)	TLE3, AHR, ZNF217, AR, FOXA1, GREB1, HOXB13, ESR1, GATA3, EBNA3C, GRHL2, PIAS1, PR, HES2, FOSL2, FOXA2, NR3C1
chromHMM	Hyper	7_Enh	9_Het	13_ReprPC, 7_Enh, 10_TssBiv, 11_Biv-Flnk, 12_EnhBiv, 2_TssAFlnk, 6_EnhG
	Hypo	7_Enh, 3_TxFlnk, 6_EnhG	1_TssA, 10_TssBiv	15_Quies
HMconsensus	Hyper	N/A	H1.0, H3K9me3, H4K20me3, H3K36me2	H3K27me3, H2AK119, H3K9me3, H3K23me2
	Hypo	H3R26Cit	H3K18cr, H3K4me3, H3K23ac	H3R26Cit

samples that come from patients with ER+ or ER- tumors with superscript such as TU^{ER+} and AN^{ER-}. Since unaffected controls have no associated biomarker status, they are simply treated as a single group (HDB).

As for the ER-status-based contrasts, TU^{ER+} vs. TU^{ER-} resulted in 40,717 hypermethylated and 32,703 hypomethylated CpGs. No significant DM was detected with case-benign tissues (CUB, OQ, AN) from patients with ER+ versus ER- tumors (vertical contrasts in Fig. 3c). The probe-set enrichment analysis results are summarized in Table 2 and Supplementary Tables 6–9. The CpGs hypermethylated in TU^{ER+} compared to TU^{ER-} were enriched for transcription factor binding sites (TFBS) related to chromatin remodeling and epigenetic regulation (HMG20A, ZNF597, NCOR2, JMJD1C), stem cell regulation (LYL1, ETV6, LMO2, RUNX1T1, MEIS1, PBX1, TLX1), and immune modulation (SPIB, NFKB2, RELB, TBX21, BACH2, TERF1). The hypomethylated CpGs were enriched for TFBS associated with hormone receptor (HR) signaling (ESR1, PR, AR, FOXA1, GATA3, NR3C1, HOXB13) and their transcriptional co-regulators (GREB1, PIAS1, TLE3, ZXDC). Both hyper- and hypomethylation events were enriched for enhancer regions identified by chromHMM. The CpGs that were hypomethylated in TU^{ER+} compared to TU^{ER-} were significantly enriched for consensus sites of the histone modification H3R26Cit.

Contrasts along the TPxA stratified by ER status (horizontal contrasts in Fig. 3c) showed that ER+ contrasts generally yielded a higher number of DM CpGs compared to their corresponding ER- contrasts. Notably, when we assessed the overlaps between ER-stratified and pooled DM CpG sets (Fig. 3d), many CpGs were only identified as DM when stratified by ER status (TU^{ER+} vs. AN^{ER+} or TU^{ER-} vs. AN^{ER-}) but not in the combined comparison (Pooled TU^{all} vs. AN^{all}). (Fig. 3d) Specifically, 18,626 hyper- and 23,313 hypomethylated CpGs were unique to the ER+ comparison, while 9,773 hyper- and

25,380 hypomethylated CpGs were unique to the ER- comparison (Fig. 3d). Similar trends were seen in the AN vs. OQ comparison, with 46 hyper- and 26 hypomethylated CpGs unique to the ER+ comparison, and 25 hyper- and 7 hypomethylated CpGs unique to the ER- comparison.

TFBS enrichment analysis of TU^{ER+} vs. AN^{ER+} and TU^{ER-} vs. AN^{ER-} included common hits such as PRC2.2 components (SUZ12, EZH2, JARID2) and other factors (TCF21, NEUROG2). However, there was no overlap between hypomethylation TFBS hit of the TU^{ER+} vs. AN^{ER+} and TU^{ER-} vs. AN^{ER-} comparisons. Hypomethylated sites in TU^{ER+} vs. AN^{ER+} were enriched for HR-related transcription factors (e.g., ESR1, PR, AR, TLE3, ZNF217, GREB1, FOXA1, HOXB13, GATA3). Conversely, hypomethylated sites in TU^{ER-} vs. AN^{ER-} were enriched for TFBS of transcriptional regulators (e.g., HMG20A, NCOR2, RUNX1T1) and stem cell regulators (e.g., LYL1, ETV6, LMO2, MEIS1). Histone modification consensus site enrichment for H3R26Cit was only observed in TU^{ER+} vs. AN^{ER+} and not in TU^{ER-} vs. AN^{ER-}.

Interestingly, although there was no significant DM between benign tissues from patients with ER+ and ER- tumors (AN^{ER+} vs. AN^{ER-}; OQ^{ER+} vs. OQ^{ER-}; CUB^{ER+} vs. CUB^{ER-}), the number of DM CpGs differed between the CUB^{ER+} vs. HDB comparison and the CUB^{ER-} vs. HDB comparison (Fig. 3c). When we further investigated any trends in the non-overlapping hyper- or hypo-methylated CpGs between these comparisons (Supplementary Fig. 7), we found that the CpGs that were only hypomethylated in the ER+ comparison were enriched for TFBS of FOSL2, TLE3, TEAD1, NR3C1, and AR, whereas those only hypomethylated in the ER- comparison were enriched for TFBS of TP63, GREB1, AHR, TP73, PR, and GATA3 (Supplementary Table 10).




Fig. 3 Biomarker-based differences in methylation patterns. **(a, b)** PCA and t-SNE plots of case samples, separated by ER status (a) and HER2 status (b). Light gray points represent samples from the opposing ER or HER2 category, respectively. **(c)** Number of differentially methylated (DM) CpGs in ER-related comparisons. Horizontal comparisons show ER-stratified contrasts along the tumor proximity axis (TPxA), with samples more distal from TU used as the reference. Vertical comparisons represent differences between samples paired to ER+ and ER- tumors. Those paired to ER-negative tumors were used as the reference category. **(d)** Upset plots displaying overlaps between hyper- and hypo-methylated DM CpG sets from pooled or ER-stratified TU vs. AN comparison. The vertical bars represent the size of intersections, indicated by the solid circles at the bottom. The horizontal bars in the bottom left indicate the size of each individual DM CpG set by themselves

Table 2 TFBS enrichment analysis results of ER-based comparisons

		Hyper	Hypo
Within-tissue comparison	TU ^{ER+} vs. TU ^{ER-}	LYL1, EBNA3C, ETV6, LMO2, RUNX1T1, HMG20A, PBX1, ZNF597, MEIS1, SND1, TLX1, SPIB, ZBTB16, NCOR2, NFKB2, TERF1, TBX21, BACH2, RELB, JMJD1C... (24)	TLE3, AHR, ZNF217, AR, GATA3, NKX3-1, ESR1, FOXA1, HOXB13, GREB1, PIAS1, PR, H2AFX, FOXA2, NR3C1, APO-BEC3B, ZXDC, GATA6, HES2, ARRB1... (40)
Between-tissue comparison	TU ^{ER+} vs. AN ^{ER+}	JARID2, EZH2, SUZ12, TCF21, ATF7IP, NEUROG2, SP140, PCGF2, ZFP57	TLE3, ZNF217, AHR, AR, GREB1, FOXA1, ESR1, HOXB13, GATA3, PR, PIAS1, GRHL2, FOXA2, HES2, FOSL2, NR3C1
	TU ^{ER-} vs. AN ^{ER-}	TCF21, JARID2, NEUROG2, TP63, EZH2, TEAD1, TP73, ZFP57, SUZ12, PRDM6, NR3C1	EBNA3C, ETV6, HMG20A, LYL1, LMO2, RUNX1T1, MEIS1, SPIB, BACH2, NCOR2
	AN ^{ER+} vs. OQ ^{ER+}	CDCA2, CNOT3, GRHL3	NCAPH2, CNOT3, ARID2, ZBTB14, CBX8, RAG2, INTS11, FAM208A, ASXL1, ZNF202, UHRF2, EGR3, LMO1, KDM4C, TAF3, SATB1, CCNT2, HMG3, HEY1, MAPK1... (188)
	AN ^{ER-} vs. OQ ^{ER-}	CDCA2, CNOT3	NCAPH2, ARID2, CNOT3, ZNF202, ZBTB14, RAG2, FAM208A, CBX8, UHRF2, ASXL1, LMO1, INTS11, ERCC6, EGR3, HMG3, ZBTB26, SATB1, CCNT2, PCGF1, HEY1... (63)
	CUB ^{ER+} vs. HDB	LYL1, LMO2, ZBTB16, MEIS1, ETV6, FUS, PBX1, PPARG, JMJD1C, RUNX1T1, PRDM6, RXRG, TLX1, STAT5, EBNA3C, TCF4, NCOR2, MEF2B, LDB1, ZNF384... (28)	TP63, ZNF217, AHR, TLE3, TP73, GREB1, GRHL2, FOSL2, ESR1, AR, GATA3, PR, TEAD1, NR3C1, HES2
	CUB ^{ER-} vs. HDB	LYL1, LMO2, PBX1, PRDM6, PPARG, ETV6, MEIS1, ZBTB16, RXRG, FUS, JMJD1C, CEBPG, MEF2B, GATA4, LDB1, STAT5, GATA2, TAL1, GATA1, ZNF318... (32)	TP63, ZNF217, AHR, TP73, TLE3, GREB1, GRHL2, PR, ESR1, GATA3, AR, FOSL2, TEAD1, NR3C1, HES2

“Within-tissue comparison” corresponds to the vertical comparisons in Fig. 3c, and “Between-tissue comparison” corresponds to the horizontal comparisons in Fig. 3c

Discussion

This study provides the most comprehensive characterization to date of methylation profiles in tumors, three categories of case-benign tissues from 69 women, and normal breast tissue from 182 controls unaffected by cancer. Using the EPICv1 BeadChip technology, we assayed over 850,000 CpG sites across the genome.

Our findings reveal significantly different DNA methylation profiles at two critical points along the tumor proximity axis (TPx): between healthy donated breast (HDB) and contralateral unaffected breast (CUB) tissues, and between adjacent-normal (AN) and tumor (TU) tissues. This difference was evident in PCA/t-SNE visualizations and the count of differentially methylated (DM) CpGs. (Fig. 2a, c) Notably, TFBS enrichment of CUB-HDB differences underscored important breast cancer-related factors, including ESR1, PR, AR, GREB1, and GATA3. (Table 1) These results indicate the unaffected tissues in cancer cases are not epigenetically “normal”. In all our DM comparisons, we generally observed a larger number of hypomethylation events than hypermethylation events in TU compared to other tissue categories. (Fig. 2b) This finding aligns with our current understanding of DNA methylation in tumors, which is characterized by global hypomethylation and localized hypermethylation [24, 25].

Theoretically, two possible explanations could account for the observed differences between HDB and case-benign tissues. The first, more common explanation is that case-benign tissues reflect the conditions under which tumors develop. Tumor initiation is rarely a single event; instead, it results from the accumulation of

multiple factors, both exogenous (e.g., behavioral, environmental) and endogenous (e.g., aging, genetic susceptibility, inflammation) [26]. These factors may predispose larger regions of tissue (or the whole body) to malignancy, leading to epigenetic patterns associated with cancer risk [27]. The second explanation suggests reversed causality, where the epigenetic profiles in case-benign tissues develop in response to the tumor’s presence. Such tumor-responsive trends have primarily been studied in the context of tumor microenvironment [28]. The fact that the methylation profiles of AN, OQ and CUB are remarkably similar favors the first explanation. A broad response to the tumor cannot be ruled out, but is considered unlikely especially in the contralateral breast.

Although further research is required to clarify the mechanisms behind our findings, we provide novel data supporting a critical concept: while the TU vs. AN comparison remains informative for certain questions, they do not fully capture epigenetic distinctions between cancerous and normal tissues. Given the important role of the epigenetic environment on gene expression and protein content, these data have similar implications for molecular profiling in these additional dimensions. We emphasize the value of these findings for future research design, particularly in the selection of reference tissues for epigenomic studies. Our results suggest that benign tissues from breast cancer patients differ substantially from healthy donor breast tissue, underscoring the need to account for these differences when interpreting molecular profiles or constructing control groups in clinical and translational research.

The hypermethylation events distinguishing TU from AN were enriched for PRC2 transcription factor binding sites (TFBS), as well as chromHMM's Polycomb-repressed sites, reinforcing PRC2's known role in breast and other cancers. (Table 1) [29–31] Notably, we observed significant TFBS enrichment not only for core PRC2 components, EZH2 and SUZ12, but also of its optional subunit, JARID2, which recent studies have shown to promote breast tumorigenesis [32–35]. A study in prostate cancer suggests that PRC2 and DNA methylation may serve as a switch between transient and permanent gene silencing [30]. This may explain why PRC2 targets show unique enrichment in TU-AN differences but not in CUB-HDB differences. (Table 1) Certain genes transiently silenced by PRC2 in HDB, CUB, OQ, and AN may become locked in a repressed state via DNA methylation in TU. Although no specific PRC2 variant has been linked directly to DNA methylation, our findings imply the JARID2-associated variant (PRC2.2).

Interestingly, using HDB as a reference to TU resulted in 15–30% fewer hyper-methylated CpGs than when case-benign tissues were used as a reference (Fig. 2b). This reduction is due to a subset of CpGs being hypomethylated in case-benign tissues compared to both HDB and TU (Supplementary Fig. 5: “Non-monotonic down & up pattern A”). One notable group showing this pattern was the HOXA gene cluster (Supplementary Fig. 6). HOXA genes have been demonstrated to be hypermethylated in breast tumors [36], with some showing tumor-suppressor effects [37, 38], which may explain their hypomethylation in case-benign tissues to counteract the proliferative signals. Some CpGs in this set also mapped to other tumor-suppressor genes, including BAP1 [39], FBXW7 [40], FOXO3 [21–23], and TSC2 [41]. (Additional File 2: Table S4)

The TU^{ER+} vs. TU^{ER-} comparison exhibited striking methylation profile differences, with tens of thousands of DM CpGs, while TU^{HER2+} vs. TU^{HER2-} showed only five DM CpGs (Fig. 3a–c). This aligns with previous studies that identified fewer DM CpGs in HER2+ tumors compared to luminal subtypes [42, 43], although the exact CpG counts vary due to assay differences and threshold choices. Interestingly, no differential methylation was observed between benign tissues from women with ER+ vs. ER- tumors, suggesting that differences related to ER status of the tumor emerge later in the course of tumor development. (Fig. 3c) While prior studies reported that gene expression in cancer-adjacent tissue can reflect tumor subtype [44], we observed minimal epigenetic differences. Hypomethylated CpGs in TU^{ER+} vs. TU^{ER-} were, as expected, enriched for hormone receptor (HR)-related TFBS (Table 2).

Importantly, stratifying the TU-AN comparison by the tumor's ER status revealed tens of thousands of DM

CpGs that were not detected when ER groups were combined (Fig. 3d). By comparing the stratified enrichment results (Table 2) with the pooled results (Table 1), we identified which findings were consistent across ER groups and which were distinct. For example, PRC2.2 TFBS, quiescent sites, and H3K27me3/H2AK119 sites were consistently enriched in both TU^{ER+} vs. AN^{ER+} and TU^{ER-} vs. AN^{ER-} comparisons. In contrast, enrichment of H3R26Cit sites and many hypomethylation TFBS hits were predominantly seen in ER+ comparisons. (Tables 1 and 2) These findings highlight the importance of stratifying by ER status to more accurately capture the biological diversity among breast tumors.

The hypomethylated CpGs in the CUB-HDB and TU-AN comparisons shared many enrichment hits, such as TFBS of HR signaling proteins and cofactors (e.g., ESR1, PR, AR, GREB1, GATA3, ZNF217, TLE3, GRHL2) as well as the consensus histone modification sites of H3R26Cit (Table 1). These findings gain added value when interpreted in conjunction with those of the tumor ER status-related comparisons (Table 2). Specifically, many enrichment hits in the TU-AN comparison appear specific to ER+ samples: HR-related TFBS and H3R26Cit sites are enriched in TU^{ER+} vs. TU^{ER-} and TU^{ER+} vs. AN^{ER+} , but not in TU^{ER-} vs. AN^{ER-} (Table 2; Supplementary Tables 6–9). However, in the CUB vs. HDB comparison, HR-related TFBS and H3R26Cit enrichment are present in both CUB^{ER+} vs. HDB^{all} and CUB^{ER-} vs. HDB^{all} comparisons (Tables 1 and 2). This suggests that case-benign tissues are experiencing elevated HR-related signaling, regardless of the ER status of the tumor.

Our exploration of the non-overlapping DM CpGs between the CUB^{ER+} vs. HDB and CUB^{ER-} vs. HDB comparisons (Supplementary Fig. 7) showed that the non-overlapping DM CpGs between these two comparisons were enriched for many HR-related factors, such as AR, TLE3, GATA3, GREB1, PR, and TP63 (Supplementary Table 10). These findings imply that, although there were no significant differences in the direct DM analysis between CUB^{ER+} vs. CUB^{ER-} (Fig. 3c), there may be subtle signs of the paired tumor's ER status in the case-benign epigenetic profiles as well.

The enrichment of H3R26 citrullination (H3R26Cit) sites (Table 1) highlights the need for further research into potential interactions between DNA methylation and H3R26Cit in breast cancer. H3R26Cit is induced by estradiol [45] and facilitates ER binding to DNA [46]. While several studies have examined links between DNA methylation and histone methylation/deacetylation [47–49], no direct interactions between DNA methylation and histone citrullination have been reported. Our observed enrichment of H3R26Cit sites in both CUB-HDB and ER-based TU-AN comparisons, together with these known

links between estrogen pathways and H3R26Cit, suggest a promising direction of further research.

One limitation of our study is that the majority of our patient population is of European descent (Fig. 1e), which restricts the generalizability of our findings across other racial and ethnic groups. Another concern is potential batch effect, as HDB tissue processing and methylation assays were conducted at a different institution. We mitigated this by applying probe quality masking, dye bias correction, detection *p*-value masking, and background subtraction [17, 50]. Furthermore, breast cancer-related trends in our TFBS enrichment results suggest the major findings reflect meaningful biology (Tables 1 and 2), though technical artifacts in a small subset of DM CpGs, particularly in HDB comparisons, cannot be fully excluded. Finally, our study lacked external validation. Future investigations would benefit from either experimental validation (e.g. RNA expression analysis) or the validation of our findings in an independent cohort.

Conclusions

Our study revealed substantial epigenetic differences at two steps along the tumor proximity axis, underscoring the importance of tissue context and ER status in breast cancer biology. By comparing both case-benign and unaffected control tissues, we demonstrate that case-benign tissues display hormone receptor-related epigenetic changes independent of the tumor's ER status, suggesting a predisposition to malignancy and breast tissue remote from the tumor. Additionally, our findings of PRC2 and H3R26 citrullination enrichment, particularly in ER+ tumors, hint at a complex interplay between DNA methylation and histone modifications, providing a basis for future studies to explore these mechanisms in breast cancer progression. Together, these findings offer valuable insights to facilitate an understanding the multifaceted roles of epigenetic regulation in breast tumorigenesis and highlight the importance of thoughtful study design in both biological and clinical research, particularly in the selection of reference tissues and consideration of ER status.

Abbreviations

AN	Tumor-adjacent normal breast tissue
BMI	Body Mass Index
CpG	Sites in the genome where the sequence is cytosine followed by guanine
CUB	Contralateral unaffected breast tissue
DM	Differential methylation / differentially methylated
ER	Estrogen receptor
FDR	false discovery rate
HDB	Healthy donated breast tissue
HER2	Human epidermal growth factor receptor 2
HR	Hormone receptor
OQ	Opposite quadrant of the ipsilateral breast tissue
PCA	Principal component analysis
PRC2	Polycomb Repressive Complex 2
TF	Transcription factor

TFBS	Transcription factor binding site(s)
TPxA	Tumor Proximity Axis
tSNE	t-distributed stochastic neighbor embedding
TU	Tumor

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13058-025-02057-y>.

Supplementary Material 1: Supplementary Figures. This file contains all supplementary figures and their captions.

Supplementary Material 2: Supplementary Tables. This file contains all supplementary tables. Description by sheet: **S1** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in the CUB vs. HDB comparison. **S2** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in the AN vs. OQ comparison. **S3** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in the TU vs. AN comparison. **S4** – Details of the probe set enrichment analysis results for the CpGs that are hypomethylated in case-benign tissue categories. **S5** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in the TU^{HER2+} vs. TU^{HER2-} comparison. **S6** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in the TU^{ER+} vs. TU^{ER-} comparison. **S7** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in TU vs. AN stratified by ER status. **S8** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in AN vs. OQ stratified by ER status. **S9** – Details of the probe set enrichment analysis results for the hyper- and hypo-methylated CpGs in CUB vs. HDB stratified by ER status. **S10** – Details of the probe set enrichment analysis results for the CpGs only hypomethylated in CUB^{ER+} vs. HDB or in CUB^{ER-} vs. HDB

Acknowledgements

This research was supported in part through the computational resources and staff contributions provided for the Quest high performance computing facility at Northwestern University which is jointly supported by the Office of the Provost, the Office for Research, and Northwestern University Information Technology. We received support from the Pathology Core Facility of the Lurie Cancer Center which is supported by the Cancer Center Support Grant (CCSG) P30 CA060553.

Author contributions

S.C. and S.A.K. conceptualized the study. T.T. conducted sample processing and experimental assays to generate data. G.C. performed the initial data analysis. S.C., S.A.K. and Y.L. supervised the study. M.S. and P.G. contributed data from a previous study. S.D. performed data processing, analyses, visualizations, and interpretations with guidance from S.C., S.A.K., and W.Z. S.D. drafted the manuscript, which was primarily edited by S.C. and S.A.K. All authors reviewed and approved the final manuscript.

Funding

This study was supported in part by the Bluhm Family Foundation, the Cancer Center Support Grant (CCSG) P30 CA060553, and the NIH grant R01LM013337.

Data availability

The datasets generated and/or analysed during the current study will be available in NCBI's Gene Expression Omnibus (GEO) under accession GSE287331 upon publication. (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE287331>).

Declarations

Ethics approval and consent to participate

This study was conducted with the approval of the Institutional Review Boards of Northwestern University (IRB #STU00051134) and UCLA (IRB #16-000853).

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Preventive Medicine, Feinberg School of Medicine, Northwestern University, Chicago, IL 60611, USA

²Department of Breast and Thyroid Surgery, Okayama University Hospital, Okayama, Japan

³Department of Surgery, Feinberg School of Medicine, Northwestern University, Chicago, IL 60611, USA

⁴Center for Computational and Genomic Medicine, The Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA

⁵Department of Pathology and Laboratory Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

⁶Division of Hematology-Oncology, Department of Medicine, UCLA David Geffen School of Medicine, Los Angeles, CA, USA

⁷Department of Health Policy and Management, Fielding School of Public Health, University of California, Los Angeles, Los Angeles, CA, USA

⁸Department of Computational Medicine, UCLA David Geffen School of Medicine, Los Angeles, CA, USA

⁹Robert H. Lurie Cancer Center of Northwestern University, Chicago, IL, USA

Received: 28 January 2025 / Accepted: 27 May 2025

Published online: 11 June 2025

References

1. Esteller M. Epigenetics in cancer. *N Engl J Med*. 2008;358(11):1148–59.
2. Baylin SB, Jones PA. Epigenetic determinants of cancer. *Cold Spring Harb Perspect Biol*. 2016;8(9).
3. Robertson KD. DNA methylation and human disease. *Nat Rev Genet*. 2005;6(8):597–610.
4. Klutstein M, et al. DNA methylation in cancer and aging. *Cancer Res*. 2016;76(12):3446–50.
5. Teschendorff AE et al. DNA methylation outliers in normal breast tissue identify field defects that are enriched in cancer. *Nat Commun*. 2016;7.
6. Danforth DN Jr. Genomic changes in normal breast tissue in women at normal risk or at high risk for breast cancer. *Breast Cancer (Auckl)*. 2016;10:109–46.
7. Gadaleta E, et al. Field cancerization in breast cancer. *J Pathol*. 2022;257(4):561–74.
8. Greaves M, Maley CC. Clonal evolution in cancer. *Nature*. 2012;481(7381):306–13.
9. Cheng AS, et al. Epithelial progeny of estrogen-exposed breast progenitor cells display a cancer-like methylome. *Cancer Res*. 2008;68(6):1786–96.
10. Yan PS, et al. Mapping geographic zones of cancer risk with epigenetic biomarkers in normal breast tissue. *Clin Cancer Res*. 2006;12(22):6626–36.
11. Gao Y, Widschwendter M, Teschendorff AE. DNA methylation patterns in normal tissue correlate more strongly with breast cancer status than copy-number variants. *EBioMedicine*. 2018;31:243–52.
12. Muse ME, et al. Enrichment of CpG island shore region hypermethylation in epigenetic breast field cancerization. *Epigenetics*. 2020;15(10):1093–106.
13. Panjarian S, et al. Accelerated aging in normal breast tissue of women with breast cancer. *Breast Cancer Res*. 2021;23(1):58.
14. Stefansson OA, et al. A DNA methylation-based definition of biologically distinct breast cancer subtypes. *Mol Oncol*. 2015;9(3):555–68.
15. Chen L et al. Identifying methylation pattern and genes associated with breast cancer subtypes. *Int J Mol Sci*. 2019;20(17).
16. Sehl ME, et al. The effects of lifetime estrogen exposure on breast epigenetic age. *Cancer Epidemiol Biomarkers Prev*. 2021;30(6):1241–9.
17. Zhou W, et al. SeSAME: reducing artifactual detection of DNA methylation by Infinium beadchips in genomic deletions. *Nucleic Acids Res*. 2018;46(20):e123.
18. Ernst J, Kellis M. Chromatin-state discovery and genome annotation with chromHMM. *Nat Protoc*. 2017;12(12):2478–92.
19. Wahl S, et al. Epigenome-wide association study of body mass index, and the adverse outcomes of adiposity. *Nature*. 2017;541(7635):81–6.
20. Jones MJ, Goodman SJ, Kobor MS. DNA methylation and healthy human aging. *Aging Cell*. 2015;14(6):924–32.
21. Tsuji T, et al. FOXO3 is a latent tumor suppressor for FOXO3-positive and cytoplasmic-type gastric cancer cells. *Oncogene*. 2021;40(17):3072–86.
22. Mei W, et al. Role and regulation of FOXO3a: new insights into breast cancer therapy. *Front Pharmacol*. 2024;15:1346745.
23. Khan MA, et al. FOXO3 gene hypermethylation and its marked down-regulation in breast cancer cases: a study on female patients. *Front Oncol*. 2022;12:1078051.
24. Lehmann U. Aberrant DNA methylation of MicroRNA genes in human breast cancer - a critical appraisal. *Cell Tissue Res*. 2014;356(3):657–64.
25. Suijkerbuijk KPM, van Diest PJ, van der Wall E. Improving early breast cancer detection: focus on methylation. *Ann Oncol*. 2011;22(1):24–9.
26. Wu S, et al. Evaluating intrinsic and non-intrinsic cancer risk factors. *Nat Commun*. 2018;9(1):3490.
27. Zhang S, et al. Tumor initiation and early tumorigenesis: molecular mechanisms and interventional targets. *Signal Transduct Target Ther*. 2024;9(1):149.
28. Troester MA, et al. Activation of host wound responses in breast cancer microenvironment. *Clin Cancer Res*. 2009;15(22):7020–8.
29. Schlesinger Y, et al. Polycomb-mediated methylation on Lys27 of histone H3 pre-marks genes for de novo methylation in cancer. *Nat Genet*. 2007;39(2):232–6.
30. Gal-Yam EN, et al. Frequent switching of polycomb repressive marks and DNA hypermethylation in the PC3 prostate cancer cell line. *Proc Natl Acad Sci U S A*. 2008;105(35):12979–84.
31. Cai Y, et al. Epigenetic alterations to polycomb targets precede malignant transition in a mouse model of breast cancer. *Sci Rep*. 2018;8(1):5535.
32. Liu W, et al. JARID2 coordinates with the NuRD complex to facilitate breast tumorigenesis through response to adipocyte-derived leptin. *Cancer Commun (Lond)*. 2023;43(10):1117–42.
33. Zhang X, et al. Tumor mutation burden and JARID2 gene alteration are associated with short disease-free survival in locally advanced triple-negative breast cancer. *Ann Transl Med*. 2020;8(17):1052.
34. Li J, et al. JARID2 activation by NFYA promotes stemness of triple-negative breast cancer cells through the PI3K/AKT pathway. *Expert Rev Anticancer Ther*. 2024;24(10):1029–40.
35. Zhang H, et al. PBX3-activated DLG1-AS1 can promote the proliferation, invasion, and migration of TNBC cells by sponging miR-16-5p. *Mol Ther Oncolytics*. 2022;25:201–10.
36. Novak P, et al. Epigenetic inactivation of the HOXA gene cluster in breast cancer. *Cancer Res*. 2006;66(22):10664–70.
37. Pai P, et al. HOXA5-mediated stabilization of IκappaBα inhibits the NF-κappaB pathway and suppresses malignant transformation of breast epithelial cells. *Cancer Res*. 2022;82(20):3802–14.
38. Jin K, Sukumar S. BRCA1: linking HOX to breast cancer suppression. *Breast Cancer Res*. 2010;12(4):306.
39. Han A, Purwin TJ, Aplin AE. Roles of the BAP1 tumor suppressor in cell metabolism. *Cancer Res*. 2021;81(11):2807–14.
40. Jimenez-Izquierdo R, et al. FBXW7 tumor suppressor regulation by dual specificity tyrosine-regulated kinase 2. *Cell Death Dis*. 2023;14(3):202.
41. Zhu QY, et al. The role of TSC2 in breast cancer: a literature review. *Front Oncol*. 2023;13:1188371.
42. Fiegl H, et al. Breast cancer DNA methylation profiles in cancer cells and tumor stroma: association with HER-2/neu status in primary breast cancer. *Cancer Res*. 2006;66(1):29–33.
43. Chen X, Zhang J, Dai X. DNA methylation profiles capturing breast cancer heterogeneity. *BMC Genomics*. 2019;20(1):823.
44. Casbas-Hernandez P, et al. Tumor intrinsic subtype is reflected in cancer-adjacent tissue. *Cancer Epidemiol Biomarkers Prev*. 2015;24(2):406–14.
45. Zhang X, et al. Peptidylarginine deiminase 2-catalyzed histone H3 arginine 26 citrullination facilitates estrogen receptor alpha target gene activation. *Proc Natl Acad Sci U S A*. 2012;109(33):13331–6.
46. Guertin MJ, et al. Targeted H3R26 deimination specifically facilitates estrogen receptor binding by modifying nucleosome structure. *PLoS Genet*. 2014;10(9):e1004613.
47. Fuks F, et al. DNA methyltransferase Dnmt1 associates with histone deacetylase activity. *Nat Genet*. 2000;24(1):88–91.
48. Geiman TM, et al. DNMT3B interacts with hSNF2H chromatin remodeling enzyme, HDACs 1 and 2, and components of the histone methylation system. *Biochem Biophys Res Commun*. 2004;318(2):544–55.
49. Shen WC, et al. Methylation and PTEN activation in dental pulp mesenchymal stem cells promotes osteogenesis and reduces oncogenesis. *Nat Commun*. 2019;10(1):2226.

50. Leek JT, et al. Tackling the widespread and critical impact of batch effects in high-throughput data. *Nat Rev Genet.* 2010;11(10):733–9.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.