

Homework 3

Elise LaFreniere

2026-05-28

Part 1. Set up tasks

```
# reading in packages
library(tidyverse)
library(janitor)
library(here)

# reading in kelp data
kelp <- read.csv(
  here("data", "temp-kelp.csv")
)

# reading in personal data finalized
```

Part 2. Problems

Problem 1. Giant kelp fronds

Climate change threatens all ecosystems, including giant kelp (*Macrocystis pyrifera*) forests.

You are interested in the potential relationship between ocean temperature (measured in °C) and giant kelp frond elongation rate (also known as growth rate, measured in cm day⁻¹).


You conduct a field study in which you track giant kelp individuals growing in different temperatures (on average) and measure their elongation rates. Admittedly, this is not a perfect study, but you do the best with what you have!

a. An appropriate test

The appropriate tests to determine the strength of the relationship between ocean temperature and giant kelp frond elongation rate are Pearson's correlation (r) and Spearman's rank correlation (ρ) test. Pearson's correlation (r) evaluates a linear relationship between the continuous variables and assumes normality of the data, whereas Spearman's rank correlation (ρ) assesses the monotonic relationship by ranking the data points. Since biological growth rates like kelp elongation often do not follow a typical linear pattern, Spearman's rank correlation (ρ) may provide a more accurate measure of the relationship strength.

b. Create a visualization

```
# base layer: ggplot
ggplot(kelp, # from data source
      mapping = aes(
        x = temp_c, # x axis temp
        y = kelp_elong # y axis elongation
      )) +
# first layer : data points showing individual observation
geom_point(size = 2,
           alpha = 0.7,
           color = "palegreen4") + # adding custom color
# adding a linear trend line
# adding ribbon to represent confidence interval
geom_smooth(method = "lm",
            color = "sienna",
            fill = "gray",
            alpha = 0.3) +
# renaming axes, adding a title
labs(
  x = "Temperature (°C)",
  y = "Kelp Elongation (cm/day)",
  title = "Giant kelp frond elongation rate in
response to ocean temperature"
) +
# adding theme
theme_minimal()
```



Ocean temperature increase leads to an approximate decrease in kelp elongation.

c. Check your assumptions and run your test

Checking assumptions for Pearson's correlation (r)

```
# writing out model in code
kelp_model <- lm(
  kelp_elong ~ temp_c,
  data = kelp)
# looking at model results
summary(kelp_model)
```

Call:

```
lm(formula = kelp_elong ~ temp_c, data = kelp)
```

Residuals:

Min	1Q	Median	3Q	Max
-1.8643	-0.5017	0.1790	0.5659	1.2177

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	14.08645	1.49219	9.440	1.72e-10	***
temp_c	-0.43179	0.08321	-5.189	1.37e-05	***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.8665 on 30 degrees of freedom

Multiple R-squared: 0.473, Adjusted R-squared: 0.4554

F-statistic: 26.93 on 1 and 30 DF, p-value: 1.366e-05

```
# looking at diagnostics
par(mfrow = c(2,2)) # creating 2 x 2 grid
plot(kelp_model) # plotting diagnostics
```



Diagnostic Assessment

- **Residuals vs Fitted & Scale-Location:** The residuals vs fitted and scale-location appear to be homoscedastic through the approximate match of the red line to the trend line and scatter of points in relation to the line, indicating constant variance.
- **QQ-Residuals:** The QQ residuals points approximately follow the reference line with slight deviation at the left and right tails, allowing the assumption of normality to be met.
- **Residuals vs Leverage:** No outliers are past Cook's distance, indicating that outliers are not influencing the model coefficient estimates.

Since the assumptions of normality and homoscedasticity are met, and no influential outliers are present, a linear model framework is appropriate for this data, validating the use of a

Pearson's correlation test (r).

Running a Pearson's correlation test (r)

```
# creating a new object to display correlation test results
kelp_correlation <- cor.test(kelp$temp_c,
                             kelp$kelp_elong,
                             # pearson's correlation
                             method = "pearson")
# viewing correlation test results
kelp_correlation
```

Pearson's product-moment correlation

```
data: kelp$temp_c and kelp$kelp_elong
t = -5.189, df = 30, p-value = 1.366e-05
alternative hypothesis: true correlation is not equal to 0
95 percent confidence interval:
 -0.8359665 -0.4460157
sample estimates:
      cor
-0.6877489
```

d. Results communication

To evaluate the strength of the relationship between between temperature and giant kelp frond elongation rate I used a Pearson's correlation test (r) due to the model matching assumptions of linearity through approximate normality and homoscedasticity and no outliers having an influence on the model coefficients. We found a moderate-strong relationship between ocean temperature and kelp elongation (Pearson's $r = -0.69$, 95% CI $[-0.84, -0.45]$, $t(30) = -5.19$, $p < 0.001$, $\alpha = 0.05$). This indicates that there is a negative relationship between temperature and kelp elongation where for every 1 °C increase, kelp elongation decreases by 0.43 cm day⁻¹.