

# Introduction to Data Science - 1MS041

Benny Avelin

Department of Mathematics

HT 2023

## Recall from last time

---

- The Joint Distribution Function for  $X = (X_1, \dots, X_n)$  is the function

$$F(x) = \mathbb{P}(X_1 \leq x_1; \dots; X_n \leq x_n) \quad x = (x_1, \dots, x_n)$$

## Recall from last time

---

- The Joint Distribution Function for  $X = (X_1, \dots, X_n)$  is the function

$$F(x) = \mathbb{P}(X_1 \leq x_1; \dots; X_n \leq x_n) \quad x = (x_1, \dots, x_n)$$

- Random variables  $Z = (X, Y)$  are said to be independent if

$$F(z) = F(X \leq x)F(Y \leq y) \quad z = (x, y).$$

# Recall from last time

---

- The Joint Distribution Function for  $X = (X_1, \dots, X_n)$  is the function

$$F(x) = \mathbb{P}(X_1 \leq x_1; \dots; X_n \leq x_n) \quad x = (x_1, \dots, x_n)$$

- Random variables  $Z = (X, Y)$  are said to be independent if

$$F(z) = F(X \leq x)F(Y \leq y) \quad z = (x, y).$$

- A sequence of random variables  $X_1, \dots, X_n$  is simply a random vector  $X = (X_1, \dots, X_n)$

# Recall from last time

---

- The Joint Distribution Function for  $X = (X_1, \dots, X_n)$  is the function

$$F(x) = \mathbb{P}(X_1 \leq x_1; \dots; X_n \leq x_n) \quad x = (x_1, \dots, x_n)$$

- Random variables  $Z = (X, Y)$  are said to be independent if

$$F(z) = F(X \leq x)F(Y \leq y) \quad z = (x, y).$$

- A sequence of random variables  $X_1, \dots, X_n$  is simply a random vector  $X = (X_1, \dots, X_n)$
- The sequence is independent if

$$F_{X_1, \dots, X_n}(x_1, \dots, x_n) = F_{X_1}(x_1) \cdots F_{X_n}(x_n).$$

## Recall from last time

---

- The Joint Distribution Function for  $X = (X_1, \dots, X_n)$  is the function

$$F(x) = \mathbb{P}(X_1 \leq x_1; \dots; X_n \leq x_n) \quad x = (x_1, \dots, x_n)$$

- Random variables  $Z = (X, Y)$  are said to be independent if

$$F(z) = F(X \leq x)F(Y \leq y) \quad z = (x, y).$$

- A sequence of random variables  $X_1, \dots, X_n$  is simply a random vector  $X = (X_1, \dots, X_n)$
- The sequence is independent if
$$F_{X_1, \dots, X_n}(x_1, \dots, x_n) = F_{X_1}(x_1) \cdots F_{X_n}(x_n).$$
- The sequence is identically distributed if  $F_{X_i} = F_{X_j}$ .

## Recall from last time

---

- The Joint Distribution Function for  $X = (X_1, \dots, X_n)$  is the function

$$F(x) = \mathbb{P}(X_1 \leq x_1; \dots; X_n \leq x_n) \quad x = (x_1, \dots, x_n)$$

- Random variables  $Z = (X, Y)$  are said to be independent if

$$F(z) = F(X \leq x)F(Y \leq y) \quad z = (x, y).$$

- A sequence of random variables  $X_1, \dots, X_n$  is simply a random vector  $X = (X_1, \dots, X_n)$
- The sequence is independent if
$$F_{X_1, \dots, X_n}(x_1, \dots, x_n) = F_{X_1}(x_1) \cdots F_{X_n}(x_n).$$
- The sequence is identically distributed if  $F_{X_i} = F_{X_j}$ .
- If both then IID (Independent and Identically Distributed)

# Learning from data

---

- What is the average weight of the population in Sweden?



# Learning from data

---

- What is the average weight of the population in Sweden?
- What is a reasonable experiment?

# Learning from data

---

- What is the average weight of the population in Sweden?
- What is a reasonable experiment?
- What is a the random variable? Is it discrete or continuous? Is it bounded?

# Learning from data

---

- What is the average weight of the population in Sweden?
- What is a reasonable experiment?
- What is a the random variable? Is it discrete or continuous? Is it bounded?
- Is our setup of the type IID?

# Learning from data

---

## Experiment

Randomly picking a Swedish person and weighing them.

## Random variable

$X$  represents the weight of the randomly picked individual. Lets assume that the weight is between 0 and 300. We can state this as  $\mathbb{P}(0 \leq X \leq 300) = 1$ .

## What do we want to learn?

# Learning from data

---

## Experiment

Randomly picking a Swedish person and weighing them.

## Random variable

$X$  represents the weight of the randomly picked individual. Lets assume that the weight is between 0 and 300. We can state this as  $\mathbb{P}(0 \leq X \leq 300) = 1$ .

## What do we want to learn?

We want to learn  $\mathbb{E}[X]$ .

## Design

How many people should we check the weight of? What do we use to estimate  $\mathbb{E}[X]$ ?

# Learning from data

---

## Repeat experiment

Lets now say that we choose to check  $n$  people. This is an  $n$ -product experiment and we can write the result as  $X = (X_1, \dots, X_n)$  where each  $X_i$  is the weight of person  $i$ .

# Learning from data

---

## Repeat experiment

Lets now say that we choose to check  $n$  people. This is an  $n$ -product experiment and we can write the result as  $X = (X_1, \dots, X_n)$  where each  $X_i$  is the weight of person  $i$ .

## Estimator

The empirical mean is a good candidate

$$\frac{1}{n} \sum_{i=1}^n X_i \approx \mathbb{E}[X]?$$

We say that the empirical mean is an **estimator** of  $\mathbb{E}[X]$ .

# Concentration

---

## Concentration of measure

How much is the empirical mean **concentrated** around  $\mathbb{E}[X]$ ?



# Concentration

---

## Concentration of measure

How much is the empirical mean **concentrated** around  $\mathbb{E}[X]$ ?

## Theorem (**Chebychev's inequality**, $L^2$ )

For **any** RV  $X$  and any  $\epsilon > 0$ ,

$$\mathbb{P}(|X - \mathbb{E}(X)| \geq \epsilon) \leq \frac{\mathbb{V}(X)}{\epsilon^2}$$

See simulation:

# Concentration

## Concentration of measure

How much is the empirical mean **concentrated** around  $\mathbb{E}[X]$ ?

## Theorem (Chebychev's inequality, $L^2$ )

For **any** RV  $X$  and any  $\epsilon > 0$ ,

$$\mathbb{P}(|X - \mathbb{E}(X)| \geq \epsilon) \leq \frac{\mathbb{V}(X)}{\epsilon^2}$$

See simulation:

Let  $\bar{X}_n$  be our empirical mean, and say we choose  $\epsilon = 10$ , and since  $X_i \leq 300$  then  $\mathbb{E}[|Z_n|] \leq 300$ , so we have

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq 10) \leq \frac{\mathbb{V}[\bar{X}_n]}{100}$$

# What does this tell us?

---

Let  $\bar{X}_n$  be our empirical mean, and say we choose  $n = 10$  and  $\epsilon = 1$ , and since  $X_i \leq 300$  then  $\mathbb{E}[|Z_n|] \leq 300$ , so we have

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq 10) \leq \frac{\mathbb{V}[\bar{X}_n]}{100}$$

We have to use our assumptions

1. What can we say about  $\mathbb{E}[\bar{X}_n]$ ?

# What does this tell us?

---

Let  $\bar{X}_n$  be our empirical mean, and say we choose  $n = 10$  and  $\epsilon = 1$ , and since  $X_i \leq 300$  then  $\mathbb{E}[|Z_n|] \leq 300$ , so we have

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq 10) \leq \frac{\mathbb{V}[\bar{X}_n]}{100}$$

We have to use our assumptions

1. What can we say about  $\mathbb{E}[\bar{X}_n]$ ?
2. What can we say about  $\mathbb{V}[\bar{X}_n]$ ?

# What does this tell us?

Let  $\bar{X}_n$  be our empirical mean, and say we choose  $n = 10$  and  $\epsilon = 1$ , and since  $X_i \leq 300$  then  $\mathbb{E}[|Z_n|] \leq 300$ , so we have

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq 10) \leq \frac{\mathbb{V}[\bar{X}_n]}{100}$$

We have to use our assumptions

1. What can we say about  $\mathbb{E}[\bar{X}_n]$ ?
2. What can we say about  $\mathbb{V}[\bar{X}_n]$ ?

That is

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| \geq 10) \leq \frac{\mathbb{V}[X]}{100n}$$

What does this mean?

# Building confidence intervals

---

That is

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| \geq 10) \leq \frac{\mathbb{V}[X]}{100n}$$

What does this mean?

- We can use the statement to build confidence intervals.

# Building confidence intervals

That is

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| \geq 10) \leq \frac{\mathbb{V}[X]}{100n}$$

What does this mean?

- We can use the statement to build confidence intervals.
- If we want the probability of our measurement landing within 10 from the true expectation to be larger than 95% we need to find  $n$  such that

$$\frac{\mathbb{V}[X]}{100n} \leq 1 - 0.95$$

# Building confidence intervals

That is

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| \geq 10) \leq \frac{\mathbb{V}[X]}{100n}$$

What does this mean?

- We can use the statement to build confidence intervals.
- If we want the probability of our measurement landing within 10 from the true expectation to be larger than 95% we need to find  $n$  such that

$$\frac{\mathbb{V}[X]}{100n} \leq 1 - 0.95$$

- $0 \leq X \leq 300$  implies  $\mathbb{V}[X] \leq 300^2/4 = 22500$ , thus we need  $22500/5 = 4500$  samples.



# Confidence interval

---

If we look at the statement from before

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| \geq \epsilon) \leq \delta$$

# Confidence interval

---

If we look at the statement from before

$$\mathbb{P}(|\overline{X}_n - \mathbb{E}[X]| \geq \epsilon) \leq \delta$$

If we use the complementary event we get

$$\mathbb{P}(|\overline{X}_n - \mathbb{E}[X]| < \epsilon) \geq 1 - \delta$$

# Confidence interval

---

If we look at the statement from before

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| \geq \epsilon) \leq \delta$$

If we use the complementary event we get

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| < \epsilon) \geq 1 - \delta$$

Now by rearranging we get

$$\mathbb{P}(\bar{X}_n - \epsilon < \mathbb{E}[X] < \bar{X}_n + \epsilon) \geq 1 - \delta$$

# Confidence interval

---

If we look at the statement from before

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| \geq \epsilon) \leq \delta$$

If we use the complementary event we get

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[X]| < \epsilon) \geq 1 - \delta$$

Now by rearranging we get

$$\mathbb{P}(\bar{X}_n - \epsilon < \mathbb{E}[X] < \bar{X}_n + \epsilon) \geq 1 - \delta$$

## Confidence interval

For this example the confidence interval is the interval

$$(\bar{X}_n - \epsilon, \bar{X}_n + \epsilon).$$

# Confidence interval

---

## Confidence interval

If we let

$$I = (\bar{X}_n - \epsilon, \bar{X}_n + \epsilon) \quad \text{then} \quad \mathbb{P}(\mathbb{E}[X] \in I) \geq 1 - \delta.$$

Which of the following is true?

1. Lets say we used data and got an interval of  $(0.1, 0.3)$ , then the probability that the confidence interval contains the expectation is greater than  $1 - \delta$ .

# Confidence interval

---

## Confidence interval

If we let

$$I = (\bar{X}_n - \epsilon, \bar{X}_n + \epsilon) \quad \text{then} \quad \mathbb{P}(\mathbb{E}[X] \in I) \geq 1 - \delta.$$

Which of the following is true?

1. Before we have computed the interval with data, the probability that the random interval contains  $\mathbb{E}[X]$  is greater than or equal to  $1 - \delta$ .

# Confidence interval

---

## Confidence interval

If we let

$$I = (\bar{X}_n - \epsilon, \bar{X}_n + \epsilon) \quad \text{then} \quad \mathbb{P}(\mathbb{E}[X] \in I) \geq 1 - \delta.$$

Which of the following is true?

1. If I repeat the experiment of collecting data and each time computing the confidence interval then I should see roughly  $1 - \delta$  or more of them containing  $\mathbb{E}[X]$ .

# Confidence interval

---

## Confidence interval

If we let

$$I = (\bar{X}_n - \epsilon, \bar{X}_n + \epsilon) \quad \text{then} \quad \mathbb{P}(\mathbb{E}[X] \in I) \geq 1 - \delta.$$

Which of the following is true?

1. Before I repeat the experiment of collecting data and each time computing the confidence interval, I expect to see roughly  $1 - \delta$  or more of them containing  $\mathbb{E}[X]$ .



# Confidence interval

---

## Confidence interval

If we let

$$I = (\bar{X}_n - \epsilon, \bar{X}_n + \epsilon) \quad \text{then} \quad \mathbb{P}(\mathbb{E}[X] \in I) \geq 1 - \delta.$$

Which of the following is true?

1. If I in the future will compute confidence intervals with  $1 - \delta$  for the rest of my professional life, then I will produce intervals covering the true expectation roughly  $1 - \delta$  or more of the time.

# Can we do better?

---

## Theorem (Hoeffdings inequality)

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability triple and let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} F$  be  $\mathbb{R}$ -valued RVs such that  $\mathbb{P}(X_i \in [a, b]) = 1$ , then for any  $\epsilon > 0$  we get for  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ ,

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq \epsilon) \leq 2e^{-\frac{2n\epsilon^2}{(b-a)^2}}.$$

# Can we do better?

---

## Theorem (Hoeffdings inequality)

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability triple and let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} F$  be  $\mathbb{R}$ -valued RVs such that  $\mathbb{P}(X_i \in [a, b]) = 1$ , then for any  $\epsilon > 0$  we get for  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ ,

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq \epsilon) \leq 2e^{-\frac{2n\epsilon^2}{(b-a)^2}}.$$

See simulation:

# Can we do better?

## Theorem (Hoeffdings inequality)

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability triple and let  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} F$  be  $\mathbb{R}$ -valued RVs such that  $\mathbb{P}(X_i \in [a, b]) = 1$ , then for any  $\epsilon > 0$  we get for  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$ ,

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq \epsilon) \leq 2e^{-\frac{2n\epsilon^2}{(b-a)^2}}.$$

See simulation:

Again choose  $\epsilon = 10$  and find  $n$  such that

$$2e^{-\frac{2n}{900}} = 0.05$$

The solution is given by

$$1700 \approx 450 * \ln(1/0.025) = n$$

# Can we do better still?

---

## Assumptions

If we make no further assumptions, we cannot do better!!

# Can we do better still?

---

## Assumptions

If we make no further assumptions, we cannot do better!!

## Small variance

It is not unreasonable to think that the variance is not as big as  $300^2/4$  as that would correspond to half the population having weight 0 and the other half having weight 300.

# Can we do better still?

---

## Assumptions

If we make no further assumptions, we cannot do better!!

## Small variance

It is not unreasonable to think that the variance is not as big as  $300^2/4$  as that would correspond to half the population having weight 0 and the other half having weight 300.

## Data

I could not find any weight data, but I could find some data on BMI instead. Here the variance is roughly 34.

# Assumptions

---

- For simple output like Bernoulli we get good bounds.
- For random variables with a large span, it is often better to use some guided assumptions about either "spread" or how heavy the tails are.



# Tail assumptions

---

For random variables with large range but has a small spread, we can use the following instead

## Theorem (Bennett's inequality)

*Let  $X_1, \dots, X_n$  be i.i.d. random variables with finite variance such that  $\mathbb{P}(X_i \leq b) = 1$  with mean zero. Let and  $\sigma^2 = \mathbb{V}[X_i]$ . Then for any  $\epsilon > 0$ ,*

$$\mathbb{P}(|\bar{X}_n - \mathbb{E}[\bar{X}_n]| \geq \epsilon) \leq 2 \exp \left( -\frac{n\sigma^2}{b^2} h \left( \frac{b\epsilon}{\sigma^2} \right) \right)$$

*where  $h(u) = (1 + u) \log(1 + u) - u$  for  $u > 0$ .*

Going back to our example of measuring weight, if we assume that  $\sigma = 20$  we get that  $n$  should be roughly 50 for  $\epsilon = 10$ .