

# Close Binary Stars

Mónica Zorotovic, [Claus Tappert](#), Maja Vučković




Instituto de Física y Astronomía  
Universidad de Valparaíso  
Valparaíso, Chile  
[claus.tappert@uv.cl](mailto:claus.tappert@uv.cl)



## Cataclysmic Variables Part 1 – Discs

# Long-term light curves




original: AAVSO

(Normal) Dwarf Novae, U Gem stars;

semi-regular brightenings (**outbursts**) with  $\Delta m \sim 3 - 4$  mag,  $\Delta t \sim 1 - 3$  d,  $\Delta t_{\text{rec}} \sim 20 - 100$  d

# Long-term light curves

SS Cyg




original: AAVSO

(Normal) Dwarf Novae, U Gem stars;

semi-regular brightenings (**outbursts**) with  $\Delta m \sim 3 - 4$  mag,  $\Delta t \sim 1 - 3$  d,  $\Delta t_{\text{rec}} \sim 20 - 100$  d

Z Cam




original: AAVSO

Z Cam stars; irregular **standstills** after outbursts at  $\sim 1$  mag below maximum

$\Delta m \sim 2 - 3$  mag,  $\Delta t \sim 1 - 3$  d,  $\Delta t_{\text{rec}} \sim 15 - 30$  d


# Long-term light curves



original: Greiner (1998)


VY Scl stars; irregular drops in brightness (low states);  $\Delta m \sim 3 - 4$  mag

# Long-term light curves



original: Greiner (1998)


VY Scl stars; irregular drops in brightness (**low states**);  $\Delta m \sim 3 - 4$  mag



data: AAVSO


(Normal) Nova-likes; UX UMa stars;  $\approx$  constant brightness

# Long-term light curves



original: Greiner (1998)

VY Scl stars; irregular drops in brightness (**low states**);  $\Delta m \sim 3 - 4$  mag




data: AAVSO

(Normal) Nova-likes; UX UMa stars;  $\approx$  constant brightness

fun fact: the first four discovered CVs, U Gem, SS Cyg, T Leo and Z Cam, belong to three different CV subtypes

# Colours




(data from the Calán-Tololo Survey; Tappert et al. 2004; Augusteijn et al. 2010)

⇒ CVs are blue, but also red




# Spectroscopy: Dwarf Nova in quiescence



(Stover 1981)

⇒ emission lines of Balmer, He series; double-peaked

# Spectroscopy: Dwarf Nova in quiescence



(Stover 1981)

⇒ emission lines of Balmer, He series; double-peaked ⇒ **disc!**

## Line profiles from a disc

assume optically thin, axisymmetric, flat accretion disc with a Keplerian velocity distribution

## Line profiles from a disc

assume optically thin, axisymmetric, flat accretion disc with a Keplerian velocity distribution

$$\text{gravitational force} = \text{centripetal force} \Rightarrow \frac{GMm}{R^2} = \frac{mv^2}{R}$$

$$\Rightarrow v_\varphi(\varphi, R) = v_\varphi(R) \propto R^{-1/2},$$

thickness  $H \ll R_D$  ( $R_D$  radius of the disc)

(thin disc approximation)

## Line profiles from a disc

assume optically thin, axisymmetric, flat accretion disc with a Keplerian velocity distribution

$$\text{gravitational force} = \text{centripetal force} \Rightarrow \frac{GMm}{R^2} = \frac{mv^2}{R}$$


$$\Rightarrow v_\varphi(\varphi, R) = v_\varphi(R) \propto R^{-1/2},$$

thickness  $H \ll R_D$  ( $R_D$  radius of the disc)

(**thin disc approximation**)

$v_{\text{rad}} = v_\varphi(R) \cos \varphi$ : component of  $v_\varphi$  along the line of sight

(also:  $v_{\text{rad}} = v_{\text{rad}}(i)$ ,  $i$ : inclination of the disc)



# Line profiles from a disc

assume optically thin, axisymmetric, flat accretion disc with a Keplerian velocity distribution

$$\text{gravitational force} = \text{centripetal force} \Rightarrow \frac{GMm}{R^2} = \frac{mv^2}{R}$$

$$\Rightarrow v_\varphi(\varphi, R) = v_\varphi(R) \propto R^{-1/2},$$

thickness  $H \ll R_D$  ( $R_D$  radius of the disc)

(thin disc approximation)

$v_{\text{rad}} = v_\varphi(R) \cos \varphi$ : component of  $v_\varphi$  along the line of sight

(also:  $v_{\text{rad}} = v_{\text{rad}}(i)$ ,  $i$ : inclination of the disc)


$\Rightarrow$  intensity distribution along the radial velocity  $v_{\text{rad}}$ :

$$I(v_{\text{rad}}) \propto \int_{R_1}^{R_2} j(R) R \frac{d\varphi}{dv_{\text{rad}}} dR$$

$j(R) \propto R^{-\alpha}$  is the local line emissivity ( $\alpha = [1..2]$ ),

$R_1$  and  $R_2$  are the limiting radii of the disc segment


minimum: inner disc radius  $R_i$ , maximum: outer disc radius  $R_D$



## Line profiles from a disc

using  $r = \frac{R}{R_D}$ , setting the origin of a cartesian coordinate system to the centre of the accretion disc, with the observer being at  $(x, y) = (0, < 0)$ , the Keplerian velocity law transforms to

$$v_\phi(r) = \left(\frac{GM}{R}\right)^{1/2} = \frac{\text{const}}{R^{1/2}} = \frac{\text{const}}{\sqrt{r}}$$



# Line profiles from a disc

$$v_{\varphi}(r) = \left(\frac{GM}{R}\right)^{1/2} = \frac{\text{const}}{r^{1/2}}; \text{ set const} = 1$$

$$v_{\varphi}(r) = \frac{1}{\sqrt{r}} = \frac{1}{\sqrt[4]{x^2+y^2}}$$

component along the line-of-sight

$$v_y = v_{\varphi} \cos \varphi = v_{\varphi} \frac{x}{\sqrt{x^2+y^2}} = \frac{x}{(x^2+y^2)^{3/4}}$$

$$\Rightarrow y = \pm \sqrt{\left(\frac{x}{v_y}\right)^{4/3} - x^2}$$

$\Rightarrow$  equation of a dipole



# Line profiles from a disc

$$v_{\varphi}(r) = \left(\frac{GM}{R}\right)^{1/2} = \frac{\text{const}}{r^{1/2}}; \text{ set const} = 1$$



$$v_{\varphi}(r) = \frac{1}{\sqrt{r}} = \frac{1}{\sqrt[4]{x^2+y^2}}$$

component along the line-of-sight

$$v_y = v_{\varphi} \cos \varphi = v_{\varphi} \frac{x}{\sqrt{x^2+y^2}} = \frac{x}{(x^2+y^2)^{3/4}}$$


$$\Rightarrow y = \pm \sqrt{\left(\frac{x}{v_y}\right)^{4/3} - x^2}$$

$\Rightarrow$  equation of a dipole










(Horne & Marsh 1986)

# The effect of different parameters on the line profile




Tappert (1999)

# The effect of eclipses




Frank et al. (2002)

# Eclipses




Wood et al. (1986)




Wood et al. (1986)

## (Disc) CV components




Warner (1995)

# Eclipse structure




([www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))



# Eclipse structure





([www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))


# Eclipse structure



# Eclipse structure




([www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




(www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html)




(Animation: www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html)

# Eclipse structure




([www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))



# Eclipse structure




([www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikdhillon.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikhiln.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhiln.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikhiln.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhiln.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))

# Eclipse structure




([www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))




(Animation: [www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html](http://www.vikhilov.staff.shef.ac.uk/seminars/exeter/eclipses.html))


# Disc formation




a) initial gas stream




b) formation of ring



c) ring spreads



d) disk is formed





d') side view

Verbunt (1982)

Accretion discs  
transport matter  
inwards and angular  
momentum outwards

# Magnetic Turbulence




(Heller 2001)


previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)

now: matter transport through disc = flow of ionised material (blobs)  
connected via magnetic field lines: flow along lines, drag lines along

# Magnetic Turbulence



(Heller 2001)




previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)

now: matter transport through disc = flow of ionised material (blobs)


connected via magnetic field lines: flow along lines, drag lines along

consider two connected blobs  $B_1, B_2$ , at (slightly) different radii  $R_1 < R_2$

# Magnetic Turbulence



(Heller 2001)



previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)

now: matter transport through disc = flow of ionised material (blobs)


connected via magnetic field lines: flow along lines, drag lines along

consider two connected blobs  $B_1$ ,  $B_2$ , at (slightly) different radii  $R_1 < R_2$


differential rotation  $\Rightarrow B_1$  orbits faster



# Magnetic Turbulence



(Hollner 2001)



previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)


now: matter transport through disc = flow of ionised material (blobs)

connected via magnetic field lines: flow along lines, drag lines along


consider two connected blobs  $B_1$ ,  $B_2$ , at (slightly) different radii  $R_1 < R_2$

differential rotation  $\Rightarrow B_1$  orbits faster  $\Rightarrow$  stretching field line

# Magnetic Turbulence



(Heller 2001)



previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)


now: matter transport through disc = flow of ionised material (blobs)

connected via magnetic field lines: flow along lines, drag lines along


consider two connected blobs  $B_1, B_2$ , at (slightly) different radii  $R_1 < R_2$

differential rotation  $\Rightarrow B_1$  orbits faster  $\Rightarrow$  stretching field line  $\Rightarrow$  field tries to bring blobs back together

# Magnetic Turbulence



(Heller 2001)



previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)


now: matter transport through disc = flow of ionised material (blobs)

connected via magnetic field lines: flow along lines, drag lines along

consider two connected blobs  $B_1, B_2$ , at (slightly) different radii  $R_1 < R_2$

differential rotation  $\Rightarrow B_1$  orbits faster  $\Rightarrow$  stretching field line  $\Rightarrow$  field tries to bring blobs back together  $\Rightarrow B_2$  is sped up ( $\dot{J} > 0$ ),  $B_1$  is slowed down ( $\dot{J} < 0$ )

# Magnetic Turbulence



(Hellier 2001)

previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)


now: matter transport through disc = flow of ionised material (blobs)

connected via magnetic field lines: flow along lines, drag lines along


consider two connected blobs  $B_1, B_2$ , at (slightly) different radii  $R_1 < R_2$

differential rotation  $\Rightarrow B_1$  orbits faster  $\Rightarrow$  stretching field line  $\Rightarrow$  field tries to bring blobs back together  $\Rightarrow B_2$  is sped up ( $\dot{J} > 0$ ),  $B_1$  is slowed down ( $\dot{J} < 0$ )  $\Rightarrow B_2$  moves outwards,  $B_1$  inwards

# Magnetic Turbulence



(Heller 2001)



previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)


now: matter transport through disc = flow of ionised material (blobs)

connected via magnetic field lines: flow along lines, drag lines along

consider two connected blobs  $B_1, B_2$ , at (slightly) different radii  $R_1 < R_2$

differential rotation  $\Rightarrow B_1$  orbits faster  $\Rightarrow$  stretching field line  $\Rightarrow$  field tries to bring blobs back together  $\Rightarrow B_2$  is sped up ( $\dot{J} > 0$ ),  $B_1$  is slowed down ( $\dot{J} < 0$ )  $\Rightarrow B_2$  moves outwards,  $B_1$  inwards  $\Rightarrow$  field line is stretched further

# Magnetic Turbulence



(Heller 2001)

previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)



now: matter transport through disc = flow of ionised material (blobs)

connected via magnetic field lines: flow along lines, drag lines along

consider two connected blobs  $B_1, B_2$ , at (slightly) different radii  $R_1 < R_2$

differential rotation  $\Rightarrow B_1$  orbits faster  $\Rightarrow$  stretching field line  $\Rightarrow$  field tries to bring blobs back together  $\Rightarrow B_2$  is sped up ( $\dot{J} > 0$ ),  $B_1$  is slowed down ( $\dot{J} < 0$ )  $\Rightarrow B_2$  moves outwards,  $B_1$  inwards  $\Rightarrow$  field line is stretched further  $\Rightarrow$  eventually dissipation of magnetic energy by reconnection (Balbus-Hawley instability)

# Magnetic Turbulence



(Heller 2001)

previously: viscosity  $\nu_K = \alpha c_s H$  (Shakura & Sunyaev 1973)

now: matter transport through disc = flow of ionised material (blobs)


connected via magnetic field lines: flow along lines, drag lines along

consider two connected blobs  $B_1, B_2$ , at (slightly) different radii  $R_1 < R_2$

differential rotation  $\Rightarrow B_1$  orbits faster  $\Rightarrow$  stretching field line  $\Rightarrow$  field tries to bring blobs back together  $\Rightarrow B_2$  is sped up ( $\dot{J} > 0$ ),  $B_1$  is slowed down ( $\dot{J} < 0$ )  $\Rightarrow B_2$  moves outwards,  $B_1$  inwards  $\Rightarrow$  field line is stretched further  $\Rightarrow$  eventually dissipation of magnetic energy by reconnection  
(Balbus-Hawley instability)

material follow field lines  $\Rightarrow$  "intermediate" magnetic loops effectively transport matter and angular momentum  $\Rightarrow$  feasible viscosity mechanism

## Disc Structure: Eclipse Mapping




During eclipse different parts of the disc are obscured

Simultaneous obs. in different filters give colour ( $T$ ) information

translate 1-D info to 2-D structure  $\Rightarrow$  maximum-entropy algorithm






# Disc Structure: Eclipse Mapping




During eclipse different parts of the disk are obscured

Simultaneous obs. in different filters give colour ( $T$ ) information


translate 1-D info to 2-D structure  $\Rightarrow$  maximum-entropy algorithm




Wood (1994)



Wood (1994)




original: Wood et al. (1986)




original: Horne & Cook (1985)

# Outburst observations



Rutten et al. (1992)



Rutten et al. (1992)

during outburst, hump disappears, eclipse becomes symmetric (disc eclipse)

⇒ dwarf-nova outbursts = increase of disc luminosity

# Disc-Radius Variation





Fig. 1. Variations of the disk-radius. Small symbols are of lower weight. Open circles are from eclipses observed at the onset of an outburst.

(Smak 1984)

during outburst, the radius of the accretion disc increases rapidly, and afterwards slowly decreases to its quiescence value


# Spectroscopy in quiescence and outburst



Shafter (1983)

accretion disc becomes bluer (= hotter) and optically thick during outburst

# Temperature evolution in outburst



Baptista & Catalán (2001)

## Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs  
in a cold disc, recombination produces neutral atoms

## Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs

in a cold disc, recombination produces neutral atoms

⇒ mechanism is needed that can flip between a cold, low-viscosity disc (quiescence) and a hot, high-viscosity disc (outburst)

## Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs

in a cold disc, recombination produces neutral atoms

⇒ mechanism is needed that can flip between a cold, low-viscosity disc (quiescence) and a hot, high-viscosity disc (outburst)

solution: **opacity**  $\kappa$

photons interact strongly with charged particles, and only weakly with neutral ones

⇒ neutral material has low opacity to radiation



# Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs

in a cold disc, recombination produces neutral atoms

⇒ mechanism is needed that can flip between a cold, low-viscosity disc (quiescence) and a hot, high-viscosity disc (outburst)

solution: **opacity**  $\kappa$

photons interact strongly with charged particles, and only weakly with neutral ones

⇒ neutral material has low opacity to radiation

temperature fluctuations can partially ionise H ( $T \sim 7000$  K)  $\Rightarrow \dot{\kappa} > 0$

# Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs

in a cold disc, recombination produces neutral atoms

⇒ mechanism is needed that can flip between a cold, low-viscosity disc (quiescence) and a hot, high-viscosity disc (outburst)

solution: **opacity**  $\kappa$

photons interact strongly with charged particles, and only weakly with neutral ones

⇒ neutral material has low opacity to radiation

temperature fluctuations can partially ionise H ( $T \sim 7000$  K)  $\Rightarrow \dot{\kappa} > 0$

further heating favours ionisation such that  $\kappa \propto T^{10} \Rightarrow$  energy is "trapped"

⇒ jump in temperature

# Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs

in a cold disc, recombination produces neutral atoms

⇒ mechanism is needed that can flip between a cold, low-viscosity disc (quiescence) and a hot, high-viscosity disc (outburst)

solution: **opacity**  $\kappa$

photons interact strongly with charged particles, and only weakly with neutral ones

⇒ neutral material has low opacity to radiation

temperature fluctuations can partially ionise H ( $T \sim 7000$  K)  $\Rightarrow \dot{\kappa} > 0$

further heating favours ionisation such that  $\kappa \propto T^{10} \Rightarrow$  energy is "trapped"

⇒ jump in temperature

once H is fully ionised  $\kappa \propto T^{-2.5} \Rightarrow$  stable configuration

# Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs

in a cold disc, recombination produces neutral atoms

⇒ mechanism is needed that can flip between a cold, low-viscosity disc (quiescence) and a hot, high-viscosity disc (outburst)

solution: **opacity**  $\kappa$

photons interact strongly with charged particles, and only weakly with neutral ones

⇒ neutral material has low opacity to radiation

temperature fluctuations can partially ionise H ( $T \sim 7000$  K)  $\Rightarrow \dot{\kappa} > 0$

further heating favours ionisation such that  $\kappa \propto T^{10} \Rightarrow$  energy is "trapped"

⇒ jump in temperature

once H is fully ionised  $\kappa \propto T^{-2.5} \Rightarrow$  stable configuration

but: high viscosity  $\Rightarrow$  high mass-throughput

# Thermal Instability

Balbus-Hawley instability only works effectively in hot, ionised discs

in a cold disc, recombination produces neutral atoms

⇒ mechanism is needed that can flip between a cold, low-viscosity disc (quiescence) and a hot, high-viscosity disc (outburst)

solution: **opacity**  $\kappa$

photons interact strongly with charged particles, and only weakly with neutral ones

⇒ neutral material has low opacity to radiation

temperature fluctuations can partially ionise H ( $T \sim 7000$  K)  $\Rightarrow \dot{\kappa} > 0$

further heating favours ionisation such that  $\kappa \propto T^{10} \Rightarrow$  energy is "trapped"

⇒ jump in temperature


once H is fully ionised  $\kappa \propto T^{-2.5} \Rightarrow$  stable configuration

but: high viscosity  $\Rightarrow$  high mass-throughput

now, mass transfer from secondary  $\dot{M}_2$  is smaller than mass transport through the disc  $\dot{M}_d$

⇒ disc eventually returns to its quiescent state


# S-curve



(Hellier 2001)

consider one annulus in the disc with given  $\Sigma$ ,  $T$   
(surface density, temperature)

# S-curve




(Hellier 2001)

consider one annulus in the disc with given  $\Sigma$ ,  $T$   
(surface density, temperature)

A: H neutral,  $\dot{M}_d < \dot{M}_2$  (traffic jam)  $\Rightarrow \dot{\Sigma} > 0$ ,  $\dot{T} > 0$

# S-curve



(Hellier 2001)


consider one annulus in the disc with given  $\Sigma$ ,  $T$   
(surface density, temperature)

A: H neutral,  $\dot{M}_d < \dot{M}_2$  (traffic jam)  $\Rightarrow \dot{\Sigma} > 0$ ,  $\dot{T} > 0$

B: H becomes partially ionised  $\Rightarrow \dot{\kappa} > 0$ ,  $T$  rises faster than the viscous timescale at which matter is transported inwards



# S-curve



(Hellier 2001)


consider one annulus in the disc with given  $\Sigma$ ,  $T$  (surface density, temperature)

A: H neutral,  $\dot{M}_d < \dot{M}_2$  (traffic jam)  $\Rightarrow \dot{\Sigma} > 0$ ,  $\dot{T} > 0$

B: H becomes partially ionised  $\Rightarrow \dot{\kappa} > 0$ ,  $T$  rises faster than the viscous timescale at which matter is transported inwards

C: H fully ionised, high viscosity  $\Rightarrow \dot{M}_d > \dot{M}_2$  (all lights are green)  $\Rightarrow \dot{\Sigma} < 0$ ,  $\dot{T} < 0$

# S-curve



(Hellier 2001)

consider one annulus in the disc with given  $\Sigma$ ,  $T$   
(surface density, temperature)

A: H neutral,  $\dot{M}_d < \dot{M}_2$  (traffic jam)  $\Rightarrow \dot{\Sigma} > 0$ ,  $\dot{T} > 0$

B: H becomes partially ionised  $\Rightarrow \dot{\kappa} > 0$ ,  $T$  rises faster than the viscous timescale at which matter is transported inwards


C: H fully ionised, high viscosity  $\Rightarrow \dot{M}_d > \dot{M}_2$  (all lights are green)  $\Rightarrow \dot{\Sigma} < 0$ ,  $\dot{T} < 0$

D: H recombines, disc cools down to A

**Disc-Instability Model (DIM)** (Osaki 1974)

alternative: Mass-Transfer Instability (Bath 1975)


# Outburst Cycle



(Ichikawa &amp; Osaki 1992)


# Interpretation of long-term light curves

SS Cyg




original: AAVSO


Z Cam



original: AAVSO



original: Greiner (1998)



data: AAVSO

U Gem dwarf novae continuously follow the S-curve

Z Cam dwarf novae follow the S-curve, but sometimes get "stuck" in a stable configuration


VY Scl NLs have stable discs, but sometimes slip into an unstable configuration

UX UMa NLs have stable discs

→  $\dot{M}_2$  vs  $\dot{M}_{\text{disc}}$

# Additional emission sources


Theory:



(Horne & Marsh 1986)


# Additional emission sources

Theory:



(Horne & Marsh 1986)


Observation:



(Stover 1981)


# Additional emission sources

Theory:




(Horne & Marsh 1986)

Observation:




(Stover 1981)




# Additional emission sources

Theory:




(Horne & Marsh 1986)

Observation:




(Slower 1981)



possible emission sources:  
bright spot, secondary, ...



# Consequences for radial velocity measurements




$$v_r(\varphi) = \gamma - K_1 \sin(2\pi\varphi), \quad \varphi = (t - T_0) \pmod{P}$$

$$\text{Kepler III: } \frac{P^2}{a^3} = \frac{4\pi^2}{G(M_1 + M_2)}; \quad K_1 = 2\pi \frac{a_1}{P} \sin i; \quad a = a_1 + a_2$$

$$\Rightarrow q = \frac{M_2}{M_1} = \frac{K_1}{K_2}$$

# Consequences for radial velocity measurements




$$v_r(\varphi) = \gamma - K_1 \sin(2\pi\varphi), \quad \varphi = (t - T_0) \pmod{P}$$

$$\text{Kepler III: } \frac{P^2}{a^3} = \frac{4\pi^2}{G(M_1 + M_2)}; \quad K_1 = 2\pi \frac{a_1}{P} \sin i; \quad a = a_1 + a_2$$

$$\Rightarrow q = \frac{M_2}{M_1} = \frac{K_1}{K_2}$$

assuming a constant line profile that is Doppler-shifted according to the orbital motion of the WD

# Consequences for radial velocity measurements




$$v_r(\varphi) = \gamma - K_1 \sin(2\pi\varphi), \quad \varphi = (t - T_0) \pmod{P}$$

$$\text{Kepler III: } \frac{P^2}{a^3} = \frac{4\pi^2}{G(M_1 + M_2)}; \quad K_1 = 2\pi \frac{a_1}{P} \sin i; \quad a = a_1 + a_2$$

$$\Rightarrow q = \frac{M_2}{M_1} = \frac{K_1}{K_2}$$

assuming a constant line profile that is Doppler-shifted according to the orbital motion of the WD





using a single broad Gaussian to measure the central wavelength of the emission line

$\Rightarrow$  the measured semi-amplitude does not reflect the motion of the WD and thus yields erroneous parameters


additional emission is confined to the central part  $\Rightarrow$  solution: measure line wings (but: lower S/N)

# Doppler Tomography

images of the emission distribution in velocity space



Marsh & Horne (1988)





Marsh & Horne (1988)

line profiles = 1-D projections of the 2-D velocity image

⇒ maximum-entropy algorithm

# Doppler Tomography: VW Hyi




⇒ emission from secondary + bright spot

Note: transformation into positional coordinates not possible without knowledge of the velocity law!

# IP Peg in outburst

He II 4686

Model



Harlaftis et al. (1999)

spiral waves as an additional driver of angular momentum

depends on the temperature (works principally in high  $\dot{M}$  discs)