

Vector potential and electric field of a rotating charged sphere

Solution: a. The charge density is,

$$\rho(\mathbf{r}) = \frac{Q}{4\pi R^2} \delta(r - R) ,$$

giving the current density,

$$\mathbf{j}(\mathbf{r}) = \rho(\mathbf{r})\mathbf{v}(\mathbf{r}) = \rho(\mathbf{r})[\vec{\omega} \times \mathbf{r}]$$

Now, we have at the surface,

$$\mathbf{r} = R\hat{\mathbf{e}}_r = R(\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta) .$$

With $\vec{\omega} = \omega\hat{\mathbf{e}}_z = \omega(0, 0, 1)$ we obtain,

$$\hat{\mathbf{e}}_z \times \hat{\mathbf{e}}_r = (-\sin \theta \sin \phi, \sin \theta \cos \phi, 0) = \sin \theta (-\sin \phi, \cos \phi, 0) = \sin \theta \hat{\mathbf{e}}_\phi$$

and therefore the current density,

$$\mathbf{j}(\mathbf{r}) = \frac{Q\omega}{4\pi R} \sin \theta \delta(r - R) \hat{\mathbf{e}}_\phi$$

b. For the potential vector we have,

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{j}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d^3r' = \frac{Q}{(4\pi)^2 R^2} \vec{\omega} \times \int d^3r' \delta(r' - R) \frac{\mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} .$$

For the integration we let \mathbf{r} go towards z (in this case, the direction of $\vec{\omega}$ is no longer the z -axis). Now we have, $|\mathbf{r} - \mathbf{r}'| = \sqrt{r^2 + r'^2 - 2rr' \cos \theta'}$ $e \mathbf{r}' = r'(\sin \theta' \cos \phi', \sin \theta' \sin \phi', \cos \theta')$. A integration over ϕ' then gives for the x and y components the value 0. Denoting $\cos' \theta'$ by u' and replacing $\hat{\mathbf{e}}_z$, we get again, after integration for $\hat{\mathbf{e}}_r$,

$$\begin{aligned} \mathbf{A}(\mathbf{r}) &= \frac{2\pi Q}{(4\pi)^2 R^2} (\vec{\omega} \times \hat{\mathbf{e}}_r) \int r'^3 \delta(r' - R) \int_{-1}^{+1} \frac{u' du'}{\sqrt{r^2 + r'^2 - 2rr'u'}} dr' \\ &= \frac{\mu_0 QR}{8\pi} (\vec{\omega} \times \hat{\mathbf{e}}_r) \int_{-1}^{+1} \frac{u' du'}{\sqrt{r^2 + R^2 - 2rRu'}} . \end{aligned}$$

The integral can be resolved by partial integration,

$$\begin{aligned} I &= \int_{-1}^{+1} \frac{u' du'}{\sqrt{r^2 + R^2 - 2rRu'}} = -\frac{1}{rR} \left[u' \sqrt{r^2 + R^2 - 2rRu'} \right]_{-1}^{+1} + \frac{1}{rR} \int_{-1}^{+1} \sqrt{r^2 + R^2 - 2rRu'} \\ &= -\frac{1}{rR} (|r - R| + |r + R|) + \frac{1}{rR} \left[-\frac{2}{3} \frac{1}{2rR} (r^2 + R^2 - 2rRu')^{3/2} \right]_{-1}^{+1} \\ &= -\frac{1}{rR} (|r - R| + |r + R|) - \frac{1}{3r^2 R^2} (|r - R|^3 - |r + R|^3) . \end{aligned}$$

For $r > R$ we get from this,

$$I = \frac{2R}{3r^2}$$

while for $r < R$

$$I = \frac{2r}{3R^2}$$

With this we get for the potential vector,

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0 QR}{12\pi} (\vec{\omega} \times \hat{\mathbf{e}}_r) \begin{cases} \frac{r}{R^2} & \text{if } r < R \\ \frac{R}{r^2} & \text{se } r > R \end{cases} = \frac{\mu_0 QR}{12\pi} \omega \sin \theta \hat{\mathbf{e}}_\phi \begin{cases} \frac{r}{R^2} & \text{if } r < R \\ \frac{R}{r^2} & \text{se } r > R \end{cases}$$

where we use again that $\vec{\omega} = \omega \hat{\mathbf{e}}_z$ e $\hat{\mathbf{e}}_z \times \hat{\mathbf{e}}_r = \sin \theta \hat{\mathbf{e}}_\phi$. Obviously, \mathbf{A} has only the component ϕ .

b. Now the magnetic induction is $\mathbf{B} = \nabla \times \mathbf{A}$. Using the rotation in spherical coordinates (1.91), we get immediately,

$$\mathbf{B} = \hat{\mathbf{e}}_r \left[\frac{1}{r \sin \theta} \frac{\partial A_\phi \sin \theta}{\partial \theta} \right] - \hat{\mathbf{e}}_\theta \left[\frac{1}{r} A_\phi + \frac{\partial A_\phi}{\partial r} \right] = \frac{\mu_0 Q \omega}{6\pi R} \begin{cases} (\hat{\mathbf{e}}_r \cos \theta - \hat{\mathbf{e}}_\theta \sin \theta) & \text{se } r < R \\ \frac{R^3}{r^3} (\hat{\mathbf{e}}_r \cos \theta + \frac{1}{2} \hat{\mathbf{e}}_\theta \sin \theta) & \text{se } r > R \end{cases}$$

With $\hat{\mathbf{e}}_r \cos \theta - \hat{\mathbf{e}}_\theta \sin \theta = \hat{\mathbf{e}}_z$ we find that, curiously, the magnetic field is homogeneous within the layer.