

## Article

## Responses of Microbial Communities to Single-Walled Carbon Nanotubes in Phenol Wastewater Treatment Systems

Yuanyuan Qu, Qiao Ma, Jie Deng, Wenli Shen, Xuwang Zhang,  
Zhili He, Joy D Van Nostrand, Jiti Zhou, and Jizhong Zhou

*Environ. Sci. Technol.*, **Just Accepted Manuscript** • DOI: 10.1021/es5053045 • Publication Date (Web): 09 Mar 2015

Downloaded from <http://pubs.acs.org> on March 13, 2015

### Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications  
High quality. High impact.

Environmental Science & Technology is published by the American Chemical Society.  
1155 Sixteenth Street N.W., Washington, DC 20036  
Published by American Chemical Society. Copyright © American Chemical Society.  
However, no copyright claim is made to original U.S. Government works, or works  
produced by employees of any Commonwealth realm Crown government in the course  
of their duties.

**Responses of Microbial Communities to Single-Walled Carbon Nanotubes in Phenol Wastewater Treatment Systems**

Yuanyuan Qu,<sup>†,‡,\*,||</sup> Qiao Ma,<sup>†,‡,||</sup> Jie Deng,<sup>‡,||</sup> Wenli Shen,<sup>†</sup> Xuwang Zhang,<sup>†</sup> Zhili He,<sup>‡</sup> Joy D. Van Nostrand,<sup>‡</sup> Jiti Zhou<sup>†</sup>, Jizhong Zhou<sup>‡,§,#</sup>


<sup>†</sup> Key Laboratory of Industrial Ecology and Environmental Engineering (Ministry of Education), School of Environmental Science and Technology, Dalian University of Technology, Dalian 116024, China

<sup>‡</sup> Institute for Environmental Genomics (IEG), Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma 73019, USA

<sup>§</sup> Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>#</sup> State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing 100084, China

13 TOC Art



14

15

**ABSTRACT:** The expanding use of single-walled carbon nanotubes (SWCNTs) raises environmental concerns. Wastewater treatment systems are potential recipients of SWCNTs containing influent, yet the impacts of SWCNTs on these systems are poorly documented. In this study, the microbial responses to SWCNTs in simulated phenol wastewater treatment systems were investigated. The phenol removal rates were improved in all SWCNTs-treated sequencing batch reactors during the first 20 days, but when facing higher phenol concentration (1000 mg/L) after 60 days, reactors with the highest concentration (3.5 g/L) of SWCNTs exhibited a notably decreased phenol removal capacity. Cell viability tests, scanning electron microscopy analysis and DNA leakage data suggested that SWCNTs protected microbes from inactivation possibly by producing more bound extracellular polymeric substances (EPS), which could create a protective barrier for the microbes. Illumina sequencing of 16S rRNA gene amplicons revealed that the bacterial diversity did not change significantly except for a minor reduction after the immediate addition of SWCNTs. Bacterial community structure significantly shifted after SWCNTs addition and did not recover afterwards. *Zoogloea* increased significantly upon SWCNTs shocking. At final stage, *Rudaea* and *Mobilicoccus* increased, while *Burkholderia*, *Singulisphaera*, *Labrys* and *Mucilaginibacter* decreased notably. The shifts of these dominant genera may be associated with altered sludge settling, aromatic degradation and EPS production. This study suggested that SWCNTs exerted protective rather than cytotoxic effects on sludge microbes of phenol wastewater treatment systems and they affected the bacterial community structure and diversity at test concentrations. These findings provide new insights into our understanding of the potential effects of SWCNTs on wastewater treatment processes.

## ■ INTRODUCTION

Carbon nanotubes (CNTs) are amongst the most promising engineering nanomaterials due to their unique physiochemical properties compared to bulk materials. They have been incorporated into a diverse array of commercial products such as pharmaceuticals, optical devices, cosmetics, electronics and antimicrobial coatings.<sup>1,2</sup> With the exponential increase in manufacturing and application of CNTs in nanotechnology, they will inevitably enter various environmental matrices.<sup>3-6</sup> However, the potential environmental risks are poorly understood and a comprehensive investigation is needed.

Cytotoxicity of CNTs to microbes has been demonstrated using pure microbial strains.<sup>7-13</sup> The underlying molecular mechanisms were proposed to be the synergistic impacts of cell membrane perturbation and oxidative stress. However, evidence has shown that CNT toxicity to pure microbial strains is a poor predictor of toxicity to microbial communities.<sup>14,15</sup> A number of studies have therefore investigated the effects of CNTs on microbial communities in aquatic and soil environments.<sup>16-18</sup> For instance, Chung et al. reported that high concentrations of CNTs significantly lowered biomass and some enzyme activities of microbial communities from an urban soil,<sup>19,20</sup> while Shrestha et al. suggested that soil respiration and extracellular enzyme activities were not significantly affected in a sandy loam soil even in the presence of extremely high concentrations of multi-walled carbon nanotubes (MWCNTs).<sup>21</sup> CNTs were also reported to affect microbial degradation of aromatic pollutants including 2,4-dichlorophenol and phenanthrene.<sup>22,23</sup> Therefore, it is necessary to investigate the responses of microbial communities to CNTs from various environments.

Wastewater treatment plants (WWTPs), as important receptors and the sink of waste streams, are amongst the most probable CNT recipients of industrial and domestic effluents.<sup>24</sup>

Moreover, CNTs are being tailored to work as catalyst supports, composite filters, adsorbents and antimicrobial agents in wastewater treatment processes, increasing the likelihood of contact with microbial communities in activated sludges.<sup>2,25</sup> Consequently, they may pose a risk to the microbial populations and their associated functions. In this respect, the effects of other nanoparticles (Cu-, Zn-, Ag-, Al<sub>2</sub>O<sub>3</sub>-, ZnO-, TiO<sub>2</sub>-, SiO<sub>2</sub>- nanoparticles), especially Ag-nanoparticles, have been extensively studied and most negatively affected waste removal efficacy.<sup>26-28</sup> Only a few studies have examined the effects of CNTs on wastewater microbial communities.<sup>29-32</sup> Luongo and Zhang found that MWCNTs inhibited the activated sludge respiration in a dose-dependent manner within three hours,<sup>30</sup> and that single-walled carbon nanotubes (SWCNTs) altered the structure of the bacterial communities in the sludge systems within five hours based on automated ribosomal intergenic spacer analysis (ARISA).<sup>29</sup> Very recently, two studies also revealed that CNTs affected bacterial community structure of activated sludge and its corresponding functions including methane production, nitrogen and phosphorus removal.<sup>33,34</sup> Notwithstanding, the ecological effects of CNTs on activated sludge system and microbial survival are far from clear.

Recently, reports have demonstrated that CNTs altered the composition and structure of microbial communities based on results from several culture-independent technologies including ARISA, PCR-denaturing gradient gel electrophoresis (DGGE), multiplex-terminal restriction fragment length polymorphism (M-TRFLP) and clone library analyses.<sup>18,29,35</sup> However, these methods only allowed identification of microbial populations at rather coarse taxonomic levels. Current advances in high-throughput sequencing techniques have not only increased sequencing depth at a lower cost, but also provided higher taxonomic resolution.<sup>36-38</sup> Especially, the Illumina MiSeq platform, which is able to achieve comparable or greater sequencing depth than related

pyrosequencing, has been widely used to examine the phylogenetic/taxonomic diversity, composition and structure of microbial communities from a variety of environments.<sup>21,39</sup>

In this study, we aimed to: (1) explore the impacts of SWCNTs on pollutant removal efficiency in activated sludge systems, (2) investigate the microbial survival and cytotoxic mechanism, (3) monitor diversity and structure shifts of microbial communities, and (4) identify the dominant microorganisms susceptible to SWCNTs. To achieve these goals, sequencing batch reactors (SBRs) were constructed for treating phenol containing wastewater which were dosed with 0.5, 1.5, 2.5 and 3.5 g/L SWCNTs. Results demonstrated that SWCNTs played protective roles for sludge microbes, and in the meantime changed the structure of sludge bacterial communities.

## ■ MATERIALS AND METHODS

**SWCNTs and Activated Sludge.** Commercially available SWCNTs (>95%) were purchased from Shenzhen Nanotech Port Co., Ltd. (Shenzhen, China) and were suspended in distilled water using ultrasonic treatment for 30 min to obtain a better dispersion. More details regarding the SWCNTs have been described previously.<sup>40</sup> Activated sludge was gathered from the secondary sedimentation tank of Chunliu River WWTP (Dalian, China).

**Experimental Design.** The reactors were 65 cm tall with an internal diameter of 8 cm and a working volume of 2.5 L. Fine air bubbles for aeration were supplied through an air pump at the reactor bottom with an airflow rate of 0.4 L/min. The synthetic wastewater consisted of 20 mg  $\text{KH}_2\text{PO}_4/\text{L}$ , 90 mg  $\text{NH}_4\text{Cl}/\text{L}$ , 10 mg  $\text{NaCl}/\text{L}$ , 12.5 mg  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}/\text{L}$ , 12 mg  $\text{CaCl}_2/\text{L}$ , 10 mg  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}/\text{L}$ , 785 mg glucose/L and 200 mg phenol/L. The SBRs ( $n=15$ ) were seeded with activated sludge (2.57 g/L, dry weight at 105 °C), and domesticated with the synthetic wastewater. The SBRs were operated under identical conditions. Each cycle of SBR was operated for 24 h, including 2 h fill, 18 h aeration, 2 h settling, and 2 h decant. Effluent was discharged with a volumetric exchange ratio of 50%. After 15 days of domestication, the SBRs were divided into 5 groups, each receiving different concentrations of SWCNT (g/L): 0 (G1), 0.5 (G2), 1.5 (G3), 2.5 (G4) and 3.5 (G5). Each group contained three replicates. During the two month operation period, phenol concentrations of the influent and effluent were monitored daily, and the concentration of mixed liquor suspended solid (MLSS), the sludge volume after 0.5 h of settling ( $\text{SV}_{30}$ ), pH and dissolved oxygen (DO) were measured every other day. Since the system was operated in good condition, and there was barely any sludge discharge during the whole process, the MLSS was relatively stable during the process (Figure S1), thus the sludge retention time was not considered herein. The influent phenol concentration was increased to 1000 mg/L



on Day 61 to compare the robustness of the control reactors with the SWCNTs-treated ones. Phenol removal was monitored on Day 1, 20, 40, 60, 61 and 67 (the last day of operation).

**Analytical Methods.** Since activated sludge used in this study was gathered from a municipal WWTPs and aerated immediately which was transparent and colorless after centrifugation, the concentration of phenol was measured directly using a UV-vis spectrophotometer (V-560, JASCO, Japan). MLSS and SV<sub>30</sub> were determined according to standard methods. The pH and DO were measured using a pH meter (S20, Mettler-Toledo, Switzerland) and a DO meter (FLX310, Flow Electronic, China), respectively. Scanning electron microscopy (SEM) images of the SWCNTs and activated sludges were recorded using a Field Emission Scanning Electron Microscope (KYKY-1000B, KYKY Technology, China). For the detection of live/dead cells, the 2-(4-amidinophenyl)-6-indolecarbamide dihydrochloride (DAPI, Beyotime, China) and propidium iodide (PI, Beyotime, China) staining assays were performed according to the manufacturer's instructions. The efflux DNA was determined by fluorescence spectroscopy (Hitachi, Japan) using DAPI as the fluorescent dye (excitation 364, emission 454 nm) after filtration through a 0.22 µm membrane. The bound extracellular polymeric substance (EPS) of activated sludge was extracted using ethylenediaminetetraacetic acid. Concentrations of protein and carbohydrate were measured using the Lowry and anthrone methods, respectively.<sup>41</sup>

**DNA Extraction, PCR Amplification and Sequencing.** Activated sludge samples were collected at Day 1, 20, 40, 60 and 67 before sludge settling, and the genomic DNA was extracted using a protocol based on Purkhold et al.<sup>42,43</sup> DNA concentration was measured with Pico Green assays using a FLUOstar OPTIMA fluorescence plate reader (BMG Labtech, Germany). For high-throughput sequencing, the primers 515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-GGACTACHVGGGTWTCTAAT-3') were used to amplify the V4 region of the 16S

141 rRNA gene.<sup>44</sup> PCR was conducted in a 25  $\mu$ L mixture containing 0.1  $\mu$ L AccuPrime High  
142 Fidelity Taq Polymerase (Invitrogen, USA), 2.5  $\mu$ L of 10 $\times$ AccuPrime PCR buffer II, 1  $\mu$ L of  
143 each primer (10  $\mu$ M), 1  $\mu$ L template DNA and 19.4  $\mu$ L nuclease-free water under the following  
144 conditions: 94  $^{\circ}$ C for 1 min; 35 cycles of 94  $^{\circ}$ C for 20 s, 53  $^{\circ}$ C for 25 s, and 68  $^{\circ}$ C for 45 s; and a  
145 final extension at 68  $^{\circ}$ C for 10 min. Each sample was amplified in triplicate. PCR products were  
146 pooled, purified through QIAquick Gel Extraction Kit (Qiagen), and quantified by Pico Green  
147 analysis. The 16S rRNA high-throughput sequencing was conducted on Illumina MiSeq platform  
148 at the Institute for Environmental Genomics, University of Oklahoma.


149 **Sequencing data analysis of 16S rRNA gene amplicons.** After sequencing, PhiX sequences  
150 were removed and primers were trimmed (mismatch 1.5), and the paired-end reads were joined  
151 using the Flash program (phredOffset 33, standard deviation of fragment lengths 20).<sup>45</sup>  
152 Sequences containing ambiguous reads (N) and reads shorter than 240 bp were removed.<sup>46</sup> The  
153 resulting sequences were screened for Chimeras using UCHIME.<sup>47</sup> Operational taxonomic units  
154 (OTUs) were categorized using CD-HIT at a 97% sequence similarity threshold,<sup>48</sup> and the  
155 taxonomic assignment of OTUs was performed by RDP classifier with 50% confidence.<sup>49</sup> The  
156 above processes were performed through a pipeline (<http://zhoulab5.rccc.ou.edu/>) (not  
157 published). Detrended correspondence analysis (DCA) and correlation tests were calculated  
158 using R v2.15.1 (<http://www.r-project.org/>). Three nonparametric tests, including multiple  
159 response permutation procedure (MRPP), permutational multivariate analysis of variance  
160 (Adonis), and analysis of similarity (ANOSIM) were performed to test dissimilarity among  
161 treatment groups based on bray-cutis distance index (<http://ieg.ou.edu/>).

## ■ RESULTS AND DISCUSSION

**Characterization of SWCNTs.** Cytotoxicity of CNTs was closely related to the physicochemical properties. The pristine SWCNTs used here were 5-16  $\mu\text{m}$  in length, and less than 2 nm in diameter, with a surface area of 500-700  $\text{m}^2/\text{g}$ . SEM and TEM micrographs were obtained to confirm the manufacturer's description.<sup>40</sup> The impact of low metal impurities (carbon content > 95%) was not considered since it showed insignificant antibacterial activity on soil and activated sludge communities.<sup>15,29,35</sup> With the increasing manufacturing and application of CNTs, more and more CNTs will inevitably enter WWTPs via CNT production facility's release, manufacturing and disposal of CNT containing products.<sup>4,5,50,51</sup> CNTs released from composites such as sports equipment, tires, electronics, etc. have been reported elsewhere.<sup>51</sup> CNTs may agglomerate and accumulate in activated sludge due to their high hydrophobic and biodegradation-resistant characteristics.<sup>5,34</sup> In previous reports, low concentrations of CNTs have been proven to exert no or little impact on microbial and enzyme activities of soils and activated sludge.<sup>19,33,35</sup> Our preliminary experiments also showed that the microbial communities were not changed with 0.1 g/L SWCNT, while a noticeable shift was observed in 2.0 g/L SWCNT group using PCR-DGGE analysis (data not shown). Therefore, the concentrations of SWCNTs used in the present study were set at a relative high level from 0.5 to 3.5 g/L.

**Effects of SWCNTs on Reactor Performances.** Acute and chronic influences of SWCNTs on the phenol wastewater treatments were mimicked over a course of two months' operation. Phenol was almost completely removed after one cycle of operation in all reactors (Figure 1). On the first 20 days, phenol removal rates were SWCNT-dose-dependent with the order of  $\text{G5} > \text{G4} > \text{G3} > \text{G2} > \text{G1}$  (Figure S2). On Day 1, the removal rate reached  $96.1 \pm 2.4\%$  within two

185 hours in G5, while it took eight hours for G1 to reach a similar removal rate, indicating a positive  
186 influence of SWCNTs on phenol removal in this system. After a 40-day operation, all groups  
187 reached similar phenol removal rates (Figure S2). Therefore, despite the potential toxicity,  
188 SWCNTs appeared to exhibit positive effects on sludge microbial communities, especially  
189 during the early stages of operation. SWCNTs have been widely used as adsorbents for heavy  
190 metals and aromatics removal by virtue of their unique properties.<sup>52-54</sup> Results from this study  
191 also indicated a dose-dependent relationship between phenol removal and SWCNTs, and 3.5 g/L  
192 SWCNTs could adsorb 58.9% phenol (initial concentration 180 mg/L) within 6 hours (Table S1).  
193 Thus, we initially speculated the elevated phenol removal rates on Day 1 and 20 were attributed  
194 to phenol adsorption by SWCNTs. However, when using autoclaved sludge systems, phenol  
195 adsorption did not improve in SWCNTs-treated groups (Table S1). It was previously proven that  
196 carbon-based nanomaterials could adsorb free EPS and improve EPS production, which could  
197 form a protective barrier for the microbes.<sup>12,31,34</sup> Therefore, we determined the bound protein and  
198 polysaccharide concentrations of the five groups (Figure S3), which revealed higher bound EPS  
199 concentrations in SWCNTs-treated SBRs compared to the control group. Altogether, our results  
200 suggested that the potential protective mechanism of SWCNTs possibly resulted from the higher  
201 bound EPS production.




**Figure 1.** Phenol removal performances of each group during the 67-day operation process. Each group was performed in triplicate with different concentrations of SWCNTs (g/L): 0 (G1), 0.5 (G2), 1.5 (G3), 2.5 (G4) and 3.5 (G5).

The phenol concentration in the influent was increased to 1000 mg/L on Day 61 to investigate the robustness of the constructed systems. Over 94% of the phenol was removed within 14 hours in all groups with slightly lower removal rate in G5 (Figure S2). On Day 67, all five groups showed different removal performances. G1 had a similar removal rate compared to Day 61; G2, G3 and G4 showed higher removal rates, in which 96% phenol was removed within 10 hours; G5 had a considerably lower removal rate, requiring over 18 hours to achieve 88% phenol removal (Figure S2). Therefore, different concentrations of SWCNTs had different effects on wastewater treatment systems. At low concentrations, they improved treatment performance. Once past a threshold, i.e. 3.5 g/L in the present study, they could result in a negative impact after a longtime interaction.

To determine cell survival rates upon addition of SWCNTs, cell viability tests (DAPI/PI staining) was performed at each sampling time (Figure2, Figure S4). The percentage of

218 inactivated cells in the control group was similar to G2 and was much higher than those in G3,  
219 G4 and G5. This phenomenon was consistent throughout the whole operation. Therefore, the  
220 addition of SWCNTs at relatively high concentrations apparently lowered phenol toxicity to cells.  
221 DNA leakage data also revealed a higher cell-free DNA intensity in G1, confirming the lower  
222 cell membrane damage rates in the SWCNT groups (Figure S5). The results further confirmed  
223 the positive influences of SWCNTs on the vitality of the microbial communities, which were  
224 also consistent with the EPS data. Three major cytotoxic mechanisms of CNTs have been  
225 recognized in previous studies, i.e. generation of oxidative stress, release of certain impurities  
226 and physical perturbation.<sup>33,34,55</sup> The reduced cell death and high purity of SWCNTs in the  
227 present study implied oxidative stress and metal purities should not be significant. SEM images  
228 showed a prevalence of aggregates of SWCNTs, largely reducing their direct contact with  
229 microbial cells (Figure S6). Only a few morphologically changed cells intertwined with  
230 SWCNTs, indicating limited physical toxicity of SWCNTs.<sup>15,29</sup> Meanwhile, the bound EPS  
231 concentrations in SWCNTs-treated groups were higher than those of the control group, further  
232 reducing the possibility of SWCNT physical toxicity. Therefore, the cytotoxicity of SWCNTs on  
233 sludge microbial communities was very limited and negligible. However, our results also  
234 suggested that G5 displayed a decreased phenol removal capability after phenol shock on Day 61,  
235 contrary to the results derived from microscopic observations. The distinct phenol degrading  
236 capability could be primarily, if not completely, due to community shifts resulting from  
237 SWCNTs addition.



**Figure 2.** DAPI/PI staining results of each group on Day 1. Blue parts represent the active cells stained with DAPI and red parts represent the inactivated cells stained with PI.

**Shifts of Bacterial Community Diversity and Structure.** The sludge samples from Day 1, 20, 40, 60 and 67 were sequenced using the Illumina Miseq platform. After removing low quality sequences and chimeras, the sequence number of each sample was rarefied to 15131, resulting in 226-579 OTUs at the clustering threshold of 0.97. Our results showed that 99.9% of the sequences belonged to bacteria. SWCNTs addition significantly reduced Shannon diversity and evenness (ANOVA  $P=0.027$  and  $0.025$ , respectively) on the first day (Table 1), while the richness (OTU and Chao1) did not change ( $P>0.05$ ). Thereafter, the Shannon indices, OTU richness and evenness were similar among all groups ( $P>0.05$ ), suggesting a shock loading effect upon SWCNTs addition, followed by a gradual recovery of bacterial diversity. Our results were in accordance with the report by Khodakovskaya et al. that CNTs did not affect soil bacterial diversity after a nine-week influence,<sup>39</sup> while Hai et al. reported that continuous

addition of 20 mg/L MWCNTs significantly decreased bacterial diversity,<sup>33</sup> indicating the influence of CNTs on microbial diversity was inconclusive.

**Table 1.** Alpha-diversity of all the groups at each sampling time


Day 1						Day 20				
Group	G1	G2	G3	G4	G5	G1	G2	G3	G4	G5
Shannon index	3.85	3.61	3.56	3.48	3.48	2.66	2.60	2.54	2.50	2.81
Evenness	0.61	0.57	0.56	0.56	0.56	0.49	0.46	0.44	0.43	0.48
Chao1	861	868	926	857	824	401	484	526	595	630
OTU	533	543	550	533	522	237	300	325	334	364
Day 40						Day 60				
Group	G1	G2	G3	G4	G5	G1	G2	G3	G4	G5
Shannon index	3.37	3.69	4.00	3.87	3.98	3.79	3.46	3.41	3.49	3.92
Evenness	0.57	0.61	0.65	0.63	0.66	0.65	0.65	0.59	0.57	0.59
Chao1	553	655	742	695	665	502	540	654	596	663
OTU	379	405	474	482	418	340	348	387	369	435
Day 67										
Group	G1	G2	G3	G4	G5					
Shannon index	3.10	3.00	2.84	2.87	3.16					
Evenness	0.64	0.61	0.59	0.59	0.65					
Chao1	578	559	565	567	579					
OTU	340	345	324	349	368					

Pearson correlations between taxa abundances and CNT concentrations were assessed to estimate the impact of SWCNTs on taxonomic groups (Table S2). On Day 1, more taxa were positively affected than negative ones at phylum, class, genus and OTU levels, whereas the opposite trend was observed at all other sampling time. For example, the majority of significantly impacted OTUs (78/106) were of lower abundances on Day 1, yet much smaller fractions of those were reduced on other sampling days (32/155, 27/93, 31/98, 55/126 for day 20, 40, 60 and 67, respectively). Therefore, the short-term exposure to SWCNTs repressed the growth of the majority of bacterial populations, while continuous interaction exhibited positive effects.

DCA analysis showed that the G1 samples at all sampling time separated from the other groups on a two-axes plot (Figure 3). Hierarchical clustering analysis also showed that the



triplicate samples of G1 clustered together and separated from the samples of all other groups (Figure S7). Dissimilarity analyses by Adonis, ANOSIM and MRPP confirmed that G1 was significantly distinct from the other samples (data not shown,  $P < 0.01$ ) at all time points. The changes in bacterial communities, especially the dominant populations, may lead to disturbed system functions.




276 **Figure 3.** DCA plots of all samples showing the relationships of microbial community structures  
277 among different groups at each sampling time (Day 1, 20, 40, 60 and 67). Symbols represent the  
278 samples from different groups. A distinct cluster can be defined from the samples of G1 based on  
279 the DCA ordination, which suggests the differences in microbial community structures between  
280 G1 and other four groups.

281 **Susceptible Genera in Response to SWCNTs.** The influences of SWCNTs on dominant genera  
282 (relative abundances over 1%) were further assessed at the initial (Day 1) and final (Day 60)  
283 stages of SWCNTs addition. On Day 1, there were 11 dominant genera in all groups, among  
284 which only *Zoogloea* exhibited different abundances among groups (ANOVA,  $P < 0.05$ ) and it  
285 did not survive at final stage (Figure S8). The  $SV_{30}$  values in our experiment were in the order of  
286 G1 (72.8% in average) > G2 (62.4%) > G3  $\approx$  G4  $\approx$  G5 (50.3%) in the first 10 days, which indicated  
287 that the addition of SWCNTs improved sludge settling ability (Figure S9). Yin et al. also  
288 reported SWCNTs addition improved sludge settling ability by 5.7-10.8% within five hours.<sup>31</sup>  
289 Species of *Zoogloea* (*Zoogloea ramigera*) were widely spread in activated sludge, and have been  
290 regarded as the key populations responsible for the flocculation of activated sludges.<sup>37,56,57</sup>  
291 Therefore, the increased *Zoogloea* upon SWCNTs addition in this study might have positively  
292 influenced the sludge settling ability.

293 There were 19 dominant genera on Day 60 (Figure S10), and 7 of them showed significant  
294 shifts (Figure 4). *Rudaea*, the most predominant population at final stage, remarkably increased  
295 from 3.17% in G1 to 10.90-26.55% in G2-G5 (Figure 4). The relative abundance of this genus on  
296 Day 67 (3.69% in G1, 11.47-26.48% in others) was similar to that of Day 60 (Figure S11).  
297 *Rudaea* has been identified in long-term contaminated soils with biphenyl, benzoate and  
298 naphthalene, as well as in a petroleum refinery wastewater treatment plant, thus it is of

considerable potential for aromatics biodegradation.<sup>58,59</sup> Meanwhile, it is also known for cellulose degradation and was detected as the predominate and sensitive genus susceptible to metal nanoparticles, especially nano-TiO<sub>2</sub>.<sup>60,61</sup> In our study, the increase of *Rudaea* in SWCNTs-treated groups might lead to improved aromatic (such as phenol) degradation capacity.

The abundances of *Burkholderia*, *Singulisphaera*, *Mucilaginibacter* and *Labrys* in SWCNTs-treated groups were notably lower than that in G1. Members of *Burkholderia* were known to play important roles in bioremediation of recalcitrant xenobiotics, as well as polyphosphate uptake and accumulation in activated sludge systems.<sup>62-64</sup> Type strains of *Labrys* were reported to be capable of degrading fluorobenzene, chlorobenzene and various pharmaceuticals.<sup>65,66</sup> Thus, the changes of these two bacteria might cause the fluctuation of xenobiotic removal performance. *Singulisphaera* was a newly established genus belonging to the order *Planctomycetales* with biopolymers-degrading ability.<sup>67</sup> Certain species from *Mucilaginibacter* including *Mucilaginibacter gracilis* and *Mucilaginibacter paludis* were also proficient in degrading various biopolymers (pectin, xylan, laminarin, etc.).<sup>68</sup> Therefore, the higher abundances of *Singulisphaera* and *Mucilaginibacter* in G1 might result in relatively more EPS degradation, which may explain the higher EPS concentration in SWCNTs-treated groups. *Mobilicoccus* was another increased genus in response to SWCNTs. It has so far only been isolated from fish intestinal tracts and its ecological role in WWTPs kept unknown.<sup>69</sup> *Geothrix* was an anaerobic Fe(III)-reducing bacterium which usually existed in hydrocarbon-contaminated matrix.<sup>70</sup> The roles of this genus also needed further investigation.



**Figure 4.** Relative abundances of significantly shifted genera in SWCNTs-treated groups compared with control group on Day 60. ANOVA analysis was adopted and the 7 genera had  $P<0.05$ .

*Rhodanobacter* was another predominant genus with relative abundance of over 10%. Although it showed no significant differences among G1 to G4 (ANOVA,  $P>0.05$ ), it was significantly lower in G5 (7.54%) compared with G1 (16.73%) at final stage (Figure S11). *Rhodanobacter* was reported to be capable of aromatics degradation and denitrification.<sup>71,72</sup> The decrease in abundance of this genus might negatively affect aromatic degradation and nitrogen removal performance in G5.

Most of the previous studies only searched the shifted microbes upon SWCNTs addition at one time point, yet the long-time and termly detection of community changes could bring us more useful insights. Our results revealed similar responses of *Rudaea*, *Burkholderia*, *Geothrix*, *Mobilicoccus* and *Labrys* at any sampling time, indicating that they were more susceptible to

SWCNTs in phenol wastewater treatment systems (Figure S11). However, SWCNTs exhibited varied or converse effects on some other taxa including *Rhodanobacter*, *Singulisphaer* and *Mucilaginibacter* at different time points. It suggested the impacts of SWCNTs were of temporal relations. Under *in situ* conditions, more complicated surroundings will induce disparate impacts on the community members. Therefore, the ecological effects of SWCNTs should be investigated case by case.

It was previously reported that high concentrations of SWCNTs could significantly reduce urban soil enzyme activity and affect 2,4-dichlorophenol mineralization, possibly by inhibiting the activity of soil endogenous microorganisms, whereas low concentrations of SWCNTs showed no or little influence on the microbes.<sup>19,22</sup> Based on our study, 3.5 g/L seemed to be the threshold for SWCNT toxicity, above which the performance of the SBRs could be unstable upon high phenol influent shock. For controllable engineered system-bioreactors, the performance and stability correlated with functional redundancy.<sup>73-75</sup> Hence, the specific contributions and interactions of the community members, and particularly the impacts of SWCNTs on related functional genes need further investigation to obtain an in-depth understanding of the underlying mechanisms of the SWCNT ecological effects. Since SWCNTs can significantly alter the microbial community structure, both improving and inhibiting wastewater treatment system performance, application of CNTs to wastewater treatment systems should be carefully considered for balancing both positive and negative effects.

**Implications.** Although SWCNTs were shown to be toxic to bacteria elsewhere, this study showed that addition of SWCNTs to phenol wastewater treatment systems could reduce cytotoxicity and increase phenol removal rates. However, the performance of bioreactors receiving a 3.5 g/L dose of SWCNT was significantly lowered upon loading with high

concentrations of phenol, suggesting SWCNTs also posed a potential threat to the treatment systems. SWCNTs addition significantly altered the composition and structure of indigenous activated sludge microbial communities. *Zoogloea*, *Rudaea*, *Mobilicoccus*, *Burkholderia*, *Singulisphaera*, *Labrys* and *Mucilaginibacter* were significantly shifted which might result in microbial community function fluctuations. The high-throughput sequencing technology has proven to be a feasible method for detecting subtle microbial changes resulting from CNT contamination in realistic environmental matrixes, which will contribute to promoting understanding of SWCNT nanoecotoxicology.

## ■ ASSOCIATED CONTENT

### Supporting Information

This file contains Figures S1-S11 and Tables S1-S2. The information is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*Phone: +86 411 84706251; Fax: +86 411 84706252; E-mail: [qyy@dlut.edu.cn](mailto:qyy@dlut.edu.cn)

<sup>||</sup>These authors contributed equally to this article.

### Notes

The authors declare no competing financial interest

## ■ ACKNOWLEDEMENT

This work is supported by the National Natural Science Foundation of China (No. 21176040), the Program for New Century Excellent Talents in University (No. NCET-13-

0077), the Fundamental Research Funds for the Central Universities (No. DUT14YQ107), the Office of the Vice President for Research at the University of Oklahoma and the Collaborative Innovation Center for Regional Environmental Quality. We also acknowledge Dr. Jingwei Wang, Hao Zhou and Zhaojing Zhang (Dalian University of Technology) and Yujia Qin (Oklahoma University) for technical help and useful discussions.

## ■ REFERENCES

- 1 Baughman, R. H.; Zakhidov, A. A.; de Heer, W. A. Carbon nanotubes--the route toward applications. *Science* **2002**, 297 (5582), 787-792.
- 2 Mauter, M. S.; Elimelech, M. Environmental applications of carbon-based nanomaterials. *Environ. Sci. Technol.* **2008**, 42 (16), 5843-5859.
- 3 Petersen, E. J.; Zhang, L. W.; Mattison, N. T.; O'Carroll, D. M.; Whelton, A. J.; Uddin, N.; Nguyen, T.; Huang, Q. G.; Henry, T. B.; Holbrook, R. D.; Chen, K. L. Potential release pathways, environmental fate, and ecological risks of carbon nanotubes. *Environ. Sci. Technol.* **2011**, 45 (23), 9837-9856.
- 4 Neale, P. A.; Jämting, Å. K.; Escher, B. I.; Herrmann, J. A review of the detection, fate and effects of engineered nanomaterials in wastewater treatment plants. *Water Sci. Technol.* **2013**, 68 (7), 1440-1453.
- 5 Sarma, S. J.; Bhattacharya, I.; Brar, S. K.; Tyagi, R. D.; Surampalli, R. Y. Carbon nanotube-bioaccumulation and recent advances in environmental monitoring. *Crit. Rev. Environ. Sci. Tech.* **2015**, 45 (9), 905-938.

- 398 6 Durán, N.; Guterres, S. S.; Alves, O. L.; Zucolotto, V., Eds. *Nanotoxicology: Materials,*  
399 *Methodologies, and Assessments*. Springer Science & Business Media, 2013.
- 400 7 Zhu, B.; Xia, X.; Xia, N.; Zhang, S.; Guo, X. Modification of fatty acids in membranes of  
401 bacteria: implication for an adaptive mechanism to the toxicity of carbon nanotubes.  
402 *Environ. Sci. Technol.* **2014**, *48*, 4086-4095.
- 403 8 Kang, S.; Mauter, M. S.; Elimelech, M. Physicochemical determinants of multiwalled  
404 carbon nanotube bacterial cytotoxicity. *Environ. Sci. Technol.* **2008**, *42* (19), 7528-7534.
- 405 9 Kang, S.; Pinault, M.; Pfefferle, L. D.; Elimelech, M. Single-walled carbon nanotubes  
406 exhibit strong antimicrobial activity. *Langmuir* **2007**, *23* (17), 8670-8673.
- 407 10 Pasquini, L. M.; Hashmi, S. M.; Sommer, T. J.; Elimelech, M.; Zimmerman, J. B. Impact of  
408 surface functionalization on bacterial cytotoxicity of single-walled carbon nanotubes.  
409 *Environ. Sci. Technol.* **2012**, *46* (11), 6297-6305.
- 410 11 Vecitis, C. D.; Zodrow, K. R.; Kang, S.; Elimelech, M. Electronic-structure-dependent  
411 bacterial cytotoxicity of single-walled carbon nanotubes. *ACS nano* **2010**, *4* (9), 5471-5479.
- 412 12 Rodrigues, D. F.; Elimelech, M. Toxic effects of single-walled carbon nanotubes in the  
413 development of *E. coli* biofilm. *Environ. Sci. Technol.* **2010**, *44* (12), 4583-4589.
- 414 13 Tu, Y. S.; Lv, M.; Xiu, P.; Huynh, T.; Zhang, M.; Castelli, M.; Liu, Z. R.; Huang, Q.; Fan, C.  
415 H.; Fang, H. P.; Zhou, R. H. Destructive extraction of phospholipids from *Escherichia coli*  
416 membranes by graphene nanosheets. *Nature Nanotechnol.* **2013**, *8* (8), 594-601.



- 417 14 Kang, S.; Mauter, M. S.; Elimelech, M. Microbial cytotoxicity of carbon-based  
418 nanomaterials: implications for river water and wastewater effluent. *Environ. Sci. Technol.*  
419 **2009**, *43* (7), 2648-2653.
- 420 15 Tong, Z.; Bischoff, M.; Nies, L. F.; Myer, P.; Applegate, B.; Turco, R. F. Response of soil  
421 microorganisms to as-produced and functionalized single-wall carbon nanotubes (SWNTs).  
422 *Environ. Sci. Technol.* **2012**, *46* (24), 13471-13479.
- 423 16 Farré, M.; Gajda-Schranz, K.; Kantiani, L.; Barceló, D. Ecotoxicity and analysis of  
424 nanomaterials in the aquatic environment. *Anal. Bioanal. Chem.* **2009**, *393* (1), 81-95.
- 425 17 Eckelman, M. J.; Mauter, M. S.; Isaacs, J. A.; Elimelech, M. New perspectives on  
426 nanomaterial aquatic ecotoxicity: production impacts exceed direct exposure impacts for  
427 carbon nanotubes. *Environ. Sci. Technol.* **2012**, *46* (5), 2902-2910.
- 428 18 Rodrigues, D. F.; Jaisi, D. P.; Elimelech, M. Toxicity of functionalized single-walled carbon  
429 nanotubes on soil microbial communities: implications for nutrient cycling in soil. *Environ.*  
430 *Sci. Technol.* **2012**, *47* (1), 625-633.
- 431 19 Jin, L.; Son, Y.; Yoon, T. K.; Kang, Y. J.; Kim, W.; Chung, H. High concentrations of  
432 single-walled carbon nanotubes lower soil enzyme activity and microbial biomass.  
433 *Ecotoxicol. Environ. Saf.* **2013**, *88* (1), 9-15.
- 434 20 Chung, H.; Son, Y.; Yoon, T. K.; Kim, S.; Kim, W. The effect of multi-walled carbon  
435 nanotubes on soil microbial activity. *Ecotoxicol. Environ. Saf.* **2011**, *74* (4), 569-575.

- 436 21 Shrestha, B.; Acosta-Martinez, V.; Cox, S. B.; Green, M. J.; Li, S.; Cañas-Carrell, J. E. An  
437 evaluation of the impact of multiwalled carbon nanotubes on soil microbial community  
438 structure and functioning. *J. Hazard. Mater.* **2013**, *261*, 188-197.
- 439 22 Zhou, W. Q.; Shan, J.; Jiang, B. Q.; Wang, L. H.; Feng, J. F.; Guo, H. Y.; Ji, R. Inhibitory  
440 effects of carbon nanotubes on the degradation of  $^{14}\text{C}$ -2,4-dichlorophenol in soil.  
441 *Chemosphere* **2013**, *90* (2), 527-534.
- 442 23 Cui, X. Y.; Jia, F.; Chen, Y. X.; Gan, J. Influence of single-walled carbon nanotubes on  
443 microbial availability of phenanthrene in sediment. *Ecotoxicology* **2011**, *20* (6), 1277-1285
- 444 24 Musee, N.; Thwala, M.; Nota, N. The antibacterial effects of engineered nanomaterials:  
445 implications for wastewater treatment plants. *J. Environ. Monit.* **2011**, *13* (5), 1164-1183.
- 446 25 Upadhyayula, V. K. K.; Deng, S.; Mitchell, M. C.; Smith, G. B. Application of carbon  
447 nanotube technology for removal of contaminants in drinking water: a review. *Sci. Total*  
448 *Environ.* **2009**, *408* (1), 1-13.
- 449 26 Brar, S. K.; Verma, M.; Tyagi, R. D.; Surampalli, R. Y. Engineered nanoparticles in  
450 wastewater and wastewater sludge--evidence and impacts. *Waste Manag.* **2010**, *30* (3), 504-  
451 520.
- 452 27 Chen, Y. G.; Wang, D. B.; Zhu, X. Y.; Zheng, X.; Feng, L. Y. Long-term effects of copper  
453 nanoparticles on wastewater biological nutrient removal and  $\text{N}_2\text{O}$  generation in the activated  
454 sludge process. *Environ. Sci. Technol.* **2012**, *46* (22), 12452-12458.

- 455 28 Yang, Y.; Quensen, J.; Mathieu, J.; Wang, Q.; Wang, J.; Li, M. Y.; Tiedje, J. M.; Alvarez, P.  
456 J. J. Pyrosequencing reveals higher impact of silver nanoparticles than  $\text{Ag}^+$  on the microbial  
457 community structure of activated sludge. *Water Res.* **2014**, *48*, 317-325.
- 458 29 Goyal, D.; Zhang, X. J.; Rooney - Varga, J. N. Impacts of single-walled carbon nanotubes  
459 on microbial community structure in activated sludge. *Lett. Appl. Microbiol.* **2010**, *51* (4),  
460 428-435.
- 461 30 Luongo, L. A.; Zhang, X. J. Toxicity of carbon nanotubes to the activated sludge process. *J.*  
462 *Hazard. Mater.* **2010**, *178* (1-3), 356-362.
- 463 31 Yin, Y.; Zhang, X. J.; Graham, J.; Luongo, L. Examination of purified single-walled carbon  
464 nanotubes on activated sludge process using batch reactors. *J. Environ. Sci. Health A. Tox.*  
465 *Hazard. Subst. Environ. Eng.* **2009**, *44* (7), 661-665.
- 466 32 Yin, Y.; Zhang, X. J. Evaluation of the impact of single-walled carbon nanotubes in an  
467 activated sludge wastewater reactor. *Water Sci. Technol.* **2008**, *58* (3), 623-628.
- 468 33 Hai, R.; Wang, Y.; Wang, X.; Du, Z.; Li, Y. Impacts of multiwalled carbon nanotubes on  
469 nutrient removal from wastewater and bacterial community structure in activated sludge.  
470 *PLoS ONE* **2014**, *9* (9), e107345.
- 471 34 Li, L. L.; Tong, Z. H.; Fang, C. Y.; Chu, J.; Yu, H. Q. Response of anaerobic granular  
472 sludge to single-wall carbon nanotube exposure. *Water Res.* **2015**, *70*, 1-8.
- 473 35 Jin, L.; Son, Y.; DeForest, J. L.; Kang, Y. J.; Kim, W.; Chung, H. Single-walled carbon  
474 nanotubes alter soil microbial community composition. *Sci. Total Environ.* **2014**, *466*, 533-  
475 538.

- 476 36 Luo, C.; Tsementzi, D.; Kyrpides, N.; Read, T.; Konstantinidis, K. T. Direct comparisons of  
477 Illumina vs. Roche 454 sequencing technologies on the same microbial community DNA  
478 sample. *PLoS ONE* **2012**, 7 (2), e30087.
- 479 37 Zhang, T.; Shao, M. F.; Ye, L. 454 Pyrosequencing reveals bacterial diversity of activated  
480 sludge from 14 sewage treatment plants. *ISME J.* **2012**, 6 (6), 1137-1147.
- 481 38 Ge, Y.; Schimel, J. P.; Holden, P. A. Identification of soil bacteria susceptible to TiO<sub>2</sub> and  
482 ZnO nanoparticles. *Appl. Environ. Microbiol.* **2012**, 78 (18), 6749-6758.
- 483 39 Khodakovskaya, M. V.; Kim, B. S.; Kim, J. N.; Alimohammadi, M.; Dervishi, E.; Mustafa,  
484 T.; Cernigla, C. E. Carbon nanotubes as plant growth regulators: effects on tomato growth,  
485 reproductive system, and soil microbial community. *Small* **2013**, 9 (1), 115-123.
- 486 40 Shen, E.; Qu, Y. Y.; Zhou, H.; Kong, C. L.; Ma, Q.; Zhang, X. W.; Zhou, J. T. Catalytic  
487 performance and stability of C-C bond hydrolase BphD immobilized onto single-wall  
488 carbon nanotubes. *Chinese J. Catal.* **2013**, 34 (4), 723-733.
- 489 41 Raunkjær, K.; Hvitved-Jacobsen, T.; Nielsen, P. H. Measurement of pools of protein,  
490 carbohydrate and lipid in domestic wastewater. *Water Res.* **1994**, 28 (2), 251-262.
- 491 42 Purkhold, U.; Pommerening-Röser, A.; Juretschko, S.; Schmid, M. C.; Koops, H. P.;  
492 Wagner, M. Phylogeny of all recognized species of ammonia oxidizers based on  
493 comparative 16S rRNA and *amoA* sequence analysis: implications for molecular diversity  
494 surveys. *Appl. Environ. Microbiol.* **2000**, 66, 5368-5382.

- 495 43 Ma, Q.; Qu, Y.; Shen, W.; Zhang, Z.; Wang, J.; Liu, Z.; Li, D.; Li, H.; Zhou, J. Bacterial  
496 community compositions of coking wastewater treatment plants in steel industry revealed  
497 by Illumina high-throughput sequencing. *Bioresour. Technol.* **2015**, 179, 436-443.
- 498 44 Bates, S. T.; Berg-Lyons, D.; Caporaso, J. G.; Walters, W. A.; Knight, R.; Fierer, N.  
499 Examining the global distribution of dominant archaeal populations in soil. *ISME J.* **2010**, 5  
500 (5), 908-917.
- 501 45 Magoč, T.; Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome  
502 assemblies. *Bioinformatics* **2011**, 27, 2957-2963.
- 503 46 Deng, Y.; He, Z.; Xu, M.; Qin, Y.; Van Nostrand, J. D.; Wu, L.; Roe, B. A.; Wiley, G.;  
504 Hobbie, S. E.; Reich, P. B.; Zhou, J. Elevated carbon dioxide alters the structure of soil  
505 microbial communities. *Appl. Environ. Microbiol.* **2012**, 78 (8), 2991-2995.
- 506 47 Edgar, R. C.; Haas, B. J.; Clemente, J. C.; Quince, C.; Knight, R. UCHIME improves  
507 sensitivity and speed of chimera detection. *Bioinformatics* **2011**, 27 (16), 2194-2200.
- 508 48 Li, W. Z.; Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of  
509 protein or nucleotide sequences. *Bioinformatics* **2006**, 22 (13), 1658-1659.
- 510 49 Wang, Q.; Garrity, G. M.; Tiedje, J. M.; Cole, J. R. Naïve Bayesian classifier for rapid  
511 assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.*  
512 **2007**, 73 (16), 5261-5267.
- 513 50 Bour, A.; Mouchet, F.; Silvestre, J.; Gauthier, L.; Pinelli, E. Environmentally relevant  
514 approaches to assess nanoparticles ecotoxicity: a review. *J. Hazard. Mater.* **2015**, 283, 764-  
515 777.

- 516 51 Nowack, B.; David, R. M.; Fissan, H.; Morris, H.; Shatkin, J. A.; Stintz, M.; Zepp, R.;  
517 Brouwer, D. Potential release scenarios for carbon nanotubes used in composites. *Environ.*  
518 *Int.* **2013**, 59, 1-11.
- 519 52 Ren, X. M.; Chen, C. L.; Nagatsu, M.; Wang, X. K. Carbon nanotubes as adsorbents in  
520 environmental pollution management: a review. *Chem. Eng. J.* **2011**, 170 (2-3), 395-410.
- 521 53 Parks, A. N.; Chandler, G. T.; Portis, L. M.; Sullivan, J. C.; Perron, M. M.; Cantwell, M. G.;  
522 Burgess, R. M.; Ho, K. T.; Ferguson, P. L. Effects of single-walled carbon nanotubes on the  
523 bioavailability of PCBs in field-contaminated sediments. *Nanotoxicology* **2013**, DOI:  
524 10.3109/17435390.2013.858794.
- 525 54 Lin, D.; Xingt, B. Adsorption of phenolic compounds by carbon nanotubes: role of  
526 aromaticity and substitution of hydroxyl groups. *Environ. Sci. Technol.* **2008**, 42 (19), 7254-  
527 7259.
- 528 55 Qu, X.; Alvarez, P. J.; Li, Q. Applications of nanotechnology in water and wastewater  
529 treatment. *Water Res.* **2013**, 47 (12), 3931-3946.
- 530 56 Rosselló-Mora, R. A.; Wagner, M.; Amann, R.; Schleifer, K. H. The abundance of *Zoogloea*  
531 *ramigera* in sewage treatment plants. *Appl. Environ. Microbiol.* **1995**, 61 (2), 702-707.
- 532 57 Salehizadeh, H.; Shojaosadati, S. A. Extracellular biopolymeric flocculants: recent trends  
533 and biotechnological importance. *Biotechnol. Adv.* **2001**, 19 (5), 371-385.
- 534 58 Uhlik, O.; Wald, J.; Strejcek, M.; Musilova, L.; Ridl, J.; Hroudova, M.; Vlcek, C.; Cardenas,  
535 E.; Mackova, M.; Macek, T. Identification of bacteria utilizing biphenyl, benzoate, and  
536 naphthalene in long-term contaminated soil. *PloS ONE*. **2012**, 7 (7), e40653.

- 537 59 Ibarbalz, F. M.; Figuerola, E. L.; Erijman, L. Industrial activated sludge exhibit unique  
538 bacterial community composition at high taxonomic ranks. *Water Res.* **2013**, *47* (11), 3854-  
539 3864.
- 540 60 Weon, H. Y.; Yoo, S. H.; Kim, Y. J.; Lee, C. M.; Kim, B. Y.; Jeon, Y. A.; Hong, S. B.;  
541 Anandham, R.; Kwon, S. W. *Rudaea cellulositytica* gen. nov., sp. nov., isolated from soil.  
542 *Int. J. Syst. Evol. Microbiol.* **2009**, *59*, 2308-2312.
- 543 61 Shah, V.; Jones, J.; Dickman, J.; Greenman, S. Response of soil bacterial community to  
544 metal nanoparticles in biosolids. *J. Hazard. Mater.* **2014**, *274*, 399-403.
- 545 62 Coenye, T.; Vandamme, P. Diversity and significance of *Burkholderia* species occupying  
546 diverse ecological niches. *Environ. Microbiol.* **2003**, *5* (9), 719-729.
- 547 63 Wang, J.; Quan, X.; Wu, L.; Qian, Y.; Werner, H. Bioaugmentation as a tool to enhance the  
548 removal of refractory compound in coke plant wastewater. *Process Biochem.* **2002**, *38* (5),  
549 777-781.
- 550 64 Mullan, A.; Quinn, J. P.; McGrath, J. W. Enhanced phosphate uptake and polyphosphate  
551 accumulation in *Burkholderia cepacia* grown under low-pH conditions. *Microb. Ecol.* **2002**,  
552 *44* (1), 69-77.
- 553 65 Carvalho, M. F.; De Marco, P.; Duque, A. F.; Pacheco, C. C.; Janssen, D. B.; Castro, P. M.  
554 *Labrys portucalensis* sp. nov., a fluorobenzene-degrading bacterium isolated from an  
555 industrially contaminated sediment in northern Portugal. *Int. J. Syst. Evol. Microbiol.* **2008**,  
556 *58* (3), 692-698.

- 557 66 Amorim, C. L.; Moreira, I. S.; Maia, A. S.; Tiritan, M. E.; Castro, P. M. Biodegradation of  
558 ofloxacin, norfloxacin, and ciprofloxacin as single and mixed substrates by *Labrys*  
559 *portucalensis* F11. *Appl. Microbiol. Biotechnol.* **2014**, 98 (7), 3181-3190.
- 560 67 Kulichevskaya, I. S.; Ivanova, A. O.; Baulina, O. I.; Bodelier, P. L.; Damsté, J. S. S.;  
561 Dedysh, S. N. *Singulisphaera acidiphila* gen. nov., sp. nov., a non-filamentous, Isosphaera-  
562 like planctomycete from acidic northern wetlands. *Int. J. Syst. Evol. Microbiol.* **2008**, 58 (5),  
563 1186-1193.
- 564 68 Pankratov, T. A.; Tindall, B. J.; Liesack, W.; Dedysh, S. N. *Mucilaginibacter paludis* gen.  
565 nov., sp. nov. and *Mucilaginibacter gracilis* sp. nov., pectin-, xylan- and laminarin-  
566 degrading members of the family *Sphingobacteriaceae* from acidic Sphagnum peat bog. *Int.*  
567 *J. Syst. Evol. Microbiol.* **2007**, 57 (10), 2349-2354.
- 568 69 Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E., Thompson, F., Eds. *The*  
569 *Prokaryotes: Actinobacteria*, 4th, ed.; Springer Berlin Heidelberg: Berlin, 2014.
- 570 70 Coates, J. D.; Ellis, D. J.; Gaw, C. V.; Lovley, D. R. *Geothrix fermentans* gen. nov., sp. nov.,  
571 a novel Fe (III)-reducing bacterium from a hydrocarbon-contaminated aquifer. *Int. J. Syst.*  
572 *Evol. Microbiol.* **1999**, 49 (4), 1615-1622.
- 573 71 Felföldi, T.; Székely, A. J.; Gorál, R.; Barkács, K.; Scheirich, G.; András, J.; Rácz, A.;  
574 Márialigeti, K. Polyphasic bacterial community analysis of an aerobic activated sludge  
575 removing phenols and thiocyanate from coke plant effluent. *Bioresour. Technol.* **2010**, 101  
576 (10), 3406-3414.



- 577 72 Bacosa, H. P.; Suto, K.; Inoue, C. Bacterial community dynamics during the preferential  
578 degradation of aromatic hydrocarbons by a microbial consortium. *Int. Biodeter. Biodegr.*  
579 **2012**, *74*, 109-115.
- 580 73 Briones, A.; Raskin, L. Diversity and dynamics of microbial communities in engineered  
581 environments and their implications for process stability. *Curr. Opin. Biotechnol.* **2003**, *14*  
582 (3), 270-276.
- 583 74 Pholchan, M. K.; Baptista, Jde. C.; Davenport, R. J.; Curtis, T. P. Systematic study of the  
584 effect of operating variables on reactor performance and microbial diversity in laboratory-  
585 scale activated sludge reactors. *Water Res.* **2010**, *44* (5), 1341-1352.
- 586 75 Cabrol, L.; Malhautier, L.; Poly, F.; Lepeuple, A. S.; Fanlo, J. L. Bacterial dynamics in  
587 steady-state biofilters: beyond functional stability. *FEMS Microbiol. Ecol.* **2012**, *79* (1),  
588 260-271.