

Cech (co)homology

Yury Elkin

Contents

1	Introduction	2
2	Background	3
2.1	Direct sum and direct product	3
2.2	Category theory	4
2.2.1	Dual category	7
2.2.2	Direct and inverse systems	8
2.2.3	Morphisms between systems	10
2.2.4	Finality properties of subsystems	13
2.3	Nerve of covering	15
2.3.1	Abstract simplicial complex	15
2.3.2	Coverings	17
2.3.3	The construction	20
3	Cech homology	23
3.1	Well-definedness of the Cech homology groups	24
3.2	Induced homomorphisms	24
3.3	Dimension axiom	25
3.4	Homotopy axiom	25
4	Applications	29
4.1	Lefschetz fixed-point theorem	29
4.2	Kakutani fixed point theorem	29
4.3	Continuous game theory	29
4.3.1	General version of resource optimization problem	30
	Appendices	31
A	Homology axioms	32

Chapter 1

Introduction

Chapter 2

Background

In this chapter we will recall a few details about algebra, category theory and abstract simplicial complexes. We assume that the reader is familiar with basic concepts of homology and cohomology theory and constructions related to them.

2.1 Direct sum and direct product

In this section we will recall a few details about direct sums and direct products.

Definition 2.1. Let $\{A_i\}_{i \in J}$ be a family of groups. The direct product of these groups is the Cartesian product $\prod_{i \in J} A_i$ where addition is defined component-wise $(a+b)_i = a_i + b_i$

Definition 2.2. Let $\{A_i\}_{i \in J}$ be a family of groups. The direct sum of these groups is the subgroup of direct product given by $\bigoplus_{i \in J} A_i = \{x \in \prod_{i \in J} A_i \mid x_i \neq 0 \text{ for only finitely many } i\}$.

For direct sums and direct products we have following universal properties:

Lemma 2.3. *Let $A = \prod_{i \in J} A_i$ be the direct product and D an arbitrary group. Then for every set of homomorphisms $\{f_i : D \rightarrow A_i\}_{i \in J}$ there exists a unique homomorphism $f : D \rightarrow A$ for which condition $pr_i \circ f = f_i$ holds for every $i \in J$.*

Proof. Proved in [6] theorem 3.7. □

Lemma 2.4. *Let $A = \bigoplus_{i \in J} A_i$ be the direct sum and D an arbitrary group. Then for every set of homomorphisms $\{f_i : A_i \rightarrow D\}_{i \in J}$ there exists a unique homomorphism $f : A \rightarrow D$ for which the condition $f \circ j_i = f_i$ holds for every $i \in J$.*

Proof. Proved in [6] theorem 3.6. □

2.2 Category theory

In this thesis we will present theory in a categorical way, which will make the construction more general. First lets recall the definition of a category.

Definition 2.5. A category C consist of the following ingredients: A class of objects $\text{ob}(C)$, a class of morphisms $\text{hom}(C)$ for which and for every objects A, B in $\text{ob}(C)$ there exists a subclass $\text{Hom}(A, B)$ and a rule of composition $\text{Comp} : \text{Hom}(A, B) \times \text{Hom}(B, D) \rightarrow \text{Hom}(A, D)$ for which following conditions hold:

- (1) Composition is associative. Let $f : A \rightarrow B$, $g : B \rightarrow D$ and $h : D \rightarrow E$ be morphisms between objects then $(f \circ g) \circ h = f \circ (g \circ h)$.
- (2) For every object $A \in \text{ob}(C)$ there exists identity morphism $1_A \in \text{hom}(A, A)$ for which the following condition holds: Let $f : A \rightarrow B$ and $g : B \rightarrow A$ be arbitrary morphisms then $f \circ 1_A = f$ and $1_A \circ g = g$.

Definition 2.6. We say that a category D is a subcategory of category C , if the following conditions are satisfied:

- (1) The collection of objects $\text{ob}(D)$ is a subcollection of $\text{ob}(C)$.
- (2) The collection of morphisms $\text{hom}(D)$ is a subcollection of $\text{hom}(C)$.

Definition 2.7. Let C be a category and A, B its objects and let $f : A \rightarrow B$ be some morphism between them. We say that f is an isomorphism, if there exists a morphism $g : B \rightarrow A$ for which the equations $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$ hold.

It is easy to see that topological spaces with continuous functions and abelian groups with homomorphisms form a category. We denote those categories by TOP and AB. Next we will define the concept of a functor between categories.

Definition 2.8. Let C and C' be categories and let F be a map between them. We say that F is a functor between categories if the following conditions hold:

- (1) For every object $A \in \text{ob}(C)$ there exists a unique object $F(A) \in C'$.
- (2) Let A and B be objects in $\text{ob}(C)$ and $f : A \rightarrow B$ a morphism between them. Then there exists a unique morphism $F(f) : F(A) \rightarrow F(B)$.
- (3) If f and g are morphisms in $\text{hom}(C)$, then the following condition holds: $F(f) \circ F(g) = F(f \circ g)$.

- (4) The identity element is mapped to the identity element. Let A be an object in $\text{ob}(C)$ and 1_A the identity morphism corresponding to it. Then $F(1_A)$ is the identity morphism of $F(A)$.

Homology and cohomology groups form a functor from TOP to AB. For details see Rotman [1]. Now we will introduce a new concept called natural transformation.

Definition 2.9. Let C and D be categories and let $F : C \rightarrow D$ and $G : C \rightarrow D$ be functors. Then the following family of functors is a natural transformation:

$$\{\phi_X : F(X) \rightarrow G(X)\}_{X \in C}$$

if the following conditions hold:

- (1) For every object $X \in C$ there exists a unique morphism $\phi_X : F(X) \rightarrow G(X)$
- (2) For every morphism $f : X \rightarrow Y$ the following diagram commutes:

$$\begin{array}{ccc} F(X) & \xrightarrow{\quad \phi_X \quad} & G(X) \\ \downarrow F(f) & & \downarrow G(f) \\ F(Y) & \xrightarrow{\quad \phi_Y \quad} & G(Y) \end{array}$$

or in other words $\phi_Y \circ F(f) = G(f) \circ \phi_X$ holds.

Next we will define the concepts of limit and colimit for arbitrary categories, which will be later applied to direct and inverse systems.

Definition 2.10. Let C and J be categories. We say that the category C is indexed by the category J if there exists a functor $F : J \rightarrow C$.

We call the functor F described above an indexing functor. It is easy to see that the functor induces a subcategory $F(J)$ of C . In the case where F is indexing functor it is relatively easy to see that it is injective and thus we can equate $F(J)$ with J .

Definition 2.11. Let C be a category indexed by J and let $F : J \rightarrow C$ be the indexing functor. Let N be a fixed object of the category C . We define the cone from N to F to be an indexed family of morphisms

$$\Omega_N = \{\omega_X : N \rightarrow F(X)\}_{X \in J}$$

which satisfies the following property: If $f : X \rightarrow Y$ is a morphism in C then the following diagram commutes

$$\begin{array}{ccc} & N & \\ \omega_X \swarrow & & \searrow \omega_Y \\ F(X) & \xrightarrow{F(f)} & F(Y) \end{array}$$

From now on we will denote the cone structure described above by $\Delta(F, N, \Omega_N)$.

Definition 2.12. Let C be a category which is indexed by a category J and let F be the indexing functor. We say that the object $D \in \text{ob}(C)$ is a limit of the functor F if the following conditions are satisfied:

- (1) There exists a family of morphisms Ω_D which induce the cone structure $\Delta(F, D, \Omega_D)$.
- (2) If $\Delta(F, N, \phi)$ is any other cone structure, then there exists a unique morphism $u : N \rightarrow D$ which makes the following diagram commute.

$$\begin{array}{ccc} & N & \\ \omega_X \swarrow & \downarrow u & \searrow \omega_Y \\ & D & \\ \phi_X \swarrow & & \searrow \phi_Y \\ F(X) & \xrightarrow{F(f)} & F(Y) \end{array}$$

Next we will prove that if we have two limits N and D of a functor F then there exists an isomorphism between those two objects. In other words N and D can be identified and the limit of the functor F can be denoted simply as $\lim_{\rightarrow} C$.

Theorem 2.13. Let $F : J \rightarrow C$ be an indexing functor. Let (N, ω) and (D, ϕ) be limits of a functor F . Then there exists an isomorphism between them.

Proof. Because N and D are both limits there exist morphisms $u : N \rightarrow D$ and $v : D \rightarrow N$ as above. Because of the symmetry it is enough to show that $v \circ u = id_N$. This follows directly from the equation $\omega_X \circ v \circ u = \phi_X \circ u = \omega_X$. Because the diagram commutes for id_N then by the uniqueness condition we see that $v \circ u = id_N$. \square

2.2.1 Dual category

To simplify definitions we will define the concept of a dual category.

Definition 2.14. The dual of any concept in category theory can be obtained in the following way:

- (1) Interchange every occurrence of source with target
- (2) Reorder every composition. That is, replace $a \circ b$ with $b \circ a$.

Next we will give a few important examples of duality.

Definition 2.15. Let $F : J \rightarrow C$ be the indexing functor. We define cocone to be the dual structure of the cone.

We will denote the cocone over an object N as $\nabla(F, N, \Omega_N)$

Definition 2.16. Let C be a category which is indexed by a category J and let F be the indexing functor. We say that the object $D \in \text{ob}(C)$ is a colimit of the functor F if the following conditions are satisfied:

- (1) There exists a family of morphisms Ω_D which induces a cocone structure $\nabla(F, D, \Omega_D)$.
- (2) If $\nabla(F, N, \phi)$ is any other cocone structure, then there exists a unique morphism $u : D \rightarrow N$ which makes the following diagram commute.

$$\begin{array}{ccccc}
 & & N & & \\
 & \nearrow \phi_X & \uparrow \text{---} & \nwarrow \phi_Y & \\
 & & D & & \\
 & \nwarrow \omega_X & \uparrow \text{---} & \nearrow \omega_Y & \\
 F(X) & \xleftarrow{F(f)} & F(Y) & &
 \end{array}$$

Like in the case of limits, colimits are unique.

Theorem 2.17. Let F be an indexing functor and let (N, ϕ) and (D, ω) be colimits of F . Then there exists an isomorphism between the objects N and D .

Proof. Because N and D are both colimits there exist morphisms $u : N \rightarrow D$ and $v : D \rightarrow N$ as above. Because of the symmetry it is enough to show that $u \circ v = id_D$ holds. This follows from the equation $u \circ v \circ \omega_X = u \circ \phi_X = \omega_X$. The diagram commutes for id_D and thus by the uniqueness condition we see that $u \circ v = id_D$. \square

2.2.2 Direct and inverse systems

In this section we will give a categorical definition of direct and inverse systems. It appears that those concepts are dual of each other. We will begin by defining a quasi-ordering relation:

Definition 2.18. Let λ be a category and let $a \leq b$ be a relation in the object class $\text{ob}(\lambda)$. We say that this relation is a quasi-ordering if the following conditions hold:

- (1) $a \leq a$ for all $a \in \text{ob}(\lambda)$.
- (2) If $a \leq b$ and $b \leq c$ then $a \leq c$ for all $a, b, c \in \text{ob}(\lambda)$.

Definition 2.19. Let λ be a category with a quasi-order relation. We say that the collection of morphisms correspond to the relation, if the classes $\text{hom}(a, b)$ are of the following form:

$$\text{hom}(a, b) = \begin{cases} \{\lambda_b^a(i)\}_{i \in I(a, b)} & \text{if } b \leq a \\ \emptyset & \text{else.} \end{cases}$$

We assumed that λ is a category then every $\lambda_b^a(i) \circ \lambda_c^b(j)$ is well-defined. We define the composition to be $\lambda_c^a(i * j)$.

Definition 2.20. Let λ be a quasi-ordered category with the corresponding class of morphisms. Let C be a category and let $I : \lambda \rightarrow C$ be a covariant functor for which the following conditions are satisfied:

- (1) For every non-empty class of morphisms $\text{hom}(a, b)$ there corresponds a morphism I_b^a in category C for which $I(\text{hom}(a, b)) = \{I_b^a\}$ holds.
- (2) Every element in the class of morphisms of the form $\text{hom}(a, a)$ is mapped to the morphism $id_{I(a)}$

Then we say that I is inverse system indexed by the category λ .

The dual of the concept inverse system is direct system.

Definition 2.21. Let λ be a quasi-ordered category with the corresponding class of morphisms. Let C be a category and let $I : \lambda \rightarrow C$ be a contra-variant functor for which the following conditions are satisfied:

- (1) For every non-empty class of morphisms $\text{hom}(a, b)$ there corresponds a morphism D_a^b in category C for which $D(\text{hom}(a, b)) = \{D_a^b\}$ holds.
- (2) Every element in the class of morphisms of the form $\text{hom}(a, a)$ is mapped to the morphism $id_{I(a)}$

Then we say that D is directed system indexed by the category λ .

Lemma 2.22. *The direct and inverse system functors are well-defined*

Proof. We write this proof only for inverse systems, the case of direct systems is similar. Let (I, λ) be arbitrary inverse system, then following conditions hold:

- (1) Combining conditions (1) and (2) of the definition 2.20 and using the fact that every identity morphism can be found in collection $\text{hom}_\lambda(a, a)$ we see that the condition $I(id_a) = id_{I(a)}$ is satisfied.
- (2) The condition $I(\lambda_b^a(i)) \circ I(\lambda_c^b(j)) = I_b^a \circ I_c^b = I_c^a = I(\lambda_c^a(i * j)) = I(\lambda_b^a(i) \circ \lambda_c^b(j))$

□

Example 2.23. Let SET be the category of sets with order relation defined by $U \leq V \Leftrightarrow U \subset V$. Now for every U, V we define $D_V^U : U \rightarrow V$ to be the inclusion from U to V . Clearly this forms a direct system.

Earlier we defined the concept of limit for general indexing functors. We can interpret direct and inverse system as indexing functors, taking λ to be the indexing set. In general systems the limit may not exist. However, for our purpose it is enough to show that such limit can be found in the category of abelian groups.

Theorem 2.24. *Let $D : \lambda \rightarrow AB$ be a direct system of groups. For every $a \in \text{ob}(\lambda)$ let $i_a : D(a) \rightarrow \bigoplus_{a \in \text{ob}(\lambda)} D(a)$ be the inclusion and let G be the subgroup of $D = \bigoplus_{a \in \lambda} D(a)$ which is generated by $\{i_a x_a - i_b D_a^b x_a\}$. Then the colimit of this system is the object $L = \bigoplus_{a \in \lambda} D(a) / G$.*

Proof. We denote p to be the projection map: $p : \bigoplus_{a \in \text{ob}(\lambda)} D(a) \rightarrow L$. Let v be the family of morphisms $\{v_a = p \circ i_a : D(a) \rightarrow L\}_{a \in \text{ob}(\lambda)}$ which induces the cocone $\nabla(D, L, v)$ and let K be any other object with family of morphisms $\phi = \{\phi_a : D(a) \rightarrow K\}_{a \in \text{ob}(\lambda)}$ which induces the cocone $\nabla(I, K, \phi)$. We have to prove that there exists a unique homomorphism $u : L \rightarrow K$ for which the condition $u \circ v_a = \phi_a$ holds for all $a \in \text{ob}(\lambda)$. We will first show that such homomorphism exists

Using lemma 2.4 we can find a unique homomorphism $u' : \bigoplus_{a \in \lambda} D(a) \rightarrow K$ for which $u' \circ i_a = \phi_a$ holds. Now we see that for every generator of the subgroup G the following equations hold:

$$u'(i_a x_a - i_b D_a^b x_a) = \phi_a(x_a) - \phi_b D_a^b x_a = \phi_a(x_a) - \phi_a(x_a) = 0.$$

Now we can define u in such way that it maps every element $x + D$ to element $\phi'(x)$. Now we see that

$$u \circ v_a = u \circ p \circ i_a = u' \circ i_a = \phi_a.$$

To prove that the homomorphism u is unique we notice that if $x \in L$, then $x = \sum_{a \in A \subset D} v_a(x_a) + G$. Then the following equations hold

$$u(x) = u\left(\sum_{a \in A \subset D} v_a(x_a)\right) = \sum_{a \in A \subset D} u \circ v_a(x_a) = \sum_{a \in A \subset D} \phi_a(x_a)$$

Thus $u(x)$ is determined by the sum of the homomorphisms ϕ_a . It follows that the map u is unique. \square

Theorem 2.25. *Let $I : \lambda \rightarrow AB$ be an inverse system of groups. Then the limit of this functor is the group $L = \{x \in \prod_{i \in \text{ob}(\lambda)} I_i \mid x_a = I_a^b x_b \text{ for all } a \leq b\}$*

Proof. Let v be a family of homomorphisms $\{v_a = pr_a : L \rightarrow I(a)\}_{a \in \lambda}$ which together with the group L induce the cone $\triangle(I, L, v)$. Let K be any other group and let ϕ be a family of homomorphisms $\{\phi_a : K \rightarrow I(a)\}_{a \in \lambda}$ which induces the cone $\triangle(I, K, \phi)$.

We construct a function $u : K \rightarrow L$ for which $v_a \circ u = \phi_a$ holds. Using lemma 2.3 we find a unique homomorphism $u' : K \rightarrow \prod_{i \in J} I_i$ for which $pr_a \circ u' = \phi_a$ is satisfied. Let u'' be the homomorphism, where the image is restricted to subgroup L . We will show that the homomorphism is well-defined, specifically u' maps every element inside the subgroup L . Assume that b and a are elements which satisfy the condition $a \leq b$. Then the following equation holds

$$I_a^b \circ pr_b \circ u' = I_a^b \circ \phi_b = \phi_a = pr_a \circ u'.$$

We can define u to be u'' . The uniqueness property of the homomorphism follows directly from the fact that u' is unique. \square

2.2.3 Morphisms between systems

2.2.3.1 Morphisms between inverse systems

To be able to define morphisms between Cech homology groups we need the concept of a limit homomorphism. In this section we will define the limit morphism between systems and investigate properties of it. Concepts used in this chapter can be presented in a more general way. However, for our use it is enough to restrict ourselves to the direct and inverse system.

Definition 2.26. Let C and D be categories with order relation. A functor $\phi : C \rightarrow D$ is order preserving if for every pair $a \leq b$ in the category C , the condition $\phi(a) \leq \phi(b)$ is satisfied.

Let I be an inverse system and $\phi : \lambda' \rightarrow \lambda$ an order preserving functor. Then the inverse system $(I\phi, \lambda')$ consist of objects $\{I(\phi(a)) \mid a \in \text{ob}(\lambda')\}$ and morphisms $\{I_{\phi(a)}^{\phi(b)} \mid a \leq b\}$.

Definition 2.27. Let (I, λ) and (I', λ') be inverse systems and $\phi : \lambda' \rightarrow \lambda$ an order and limit preserving functor. Let $\{f_a : I(\phi(a)) \rightarrow I'(a) \mid a \in \lambda\}$ be a natural transformation between $I\phi$ and I' . Then we say that $\{f_a\}_{a \in \lambda}$ is an inverse system of morphisms corresponding to the functor ϕ from the system I into the system I' .

In the next theorem we will prove that it is possible to define the limit of morphisms between systems in a unique way. We recall that exists of limit implies that for every objects of category I there exists unique functor $u_{I,a} : \lim I \rightarrow I(a)$.

Lemma 2.28. Let (I, λ) and (I', λ') be inverse systems for which limit exists. Let $\{f_a\}$ be inverse system of morphisms corresponding to that pair. Then there exists a unique morphism $\lim f : \lim I \rightarrow \lim I'$ for which the condition $\lim f \circ u_{I,\phi(a)} = u_{I',a} \circ f_a$ holds for every $a \in \lambda'$.

Proof. Consider the following diagram





Diagram 2.2.3.1

To prove that the limit morphism exists we form the following triangle out of the diagram described above:



Then by definition of limit there exists a unique morphism $\lim f : \lim I \rightarrow \lim I'$ for which the diagram commutes. We will prove now that the universal property $\lim f \circ u_{I,\phi(a)} = u_{I',a} \circ f_a$ holds for this morphism. By diagram 2.2.3.1 and the fact that the family of

functions $\{f_a\}$ is a natural transformation, in other words the condition $\omega'_a \circ f_n = f_a \circ \omega_{\phi(a)}$ holds, the following diagram commutes:

$$\begin{array}{ccc}
 & \lim I(\phi(n)) & \\
 f_a \omega_{\phi(a)} \swarrow & \downarrow & \searrow f_b \omega_{\phi(b)} \\
 & \lim I' & \\
 \rho'_a \swarrow & & \searrow \rho'_b \\
 I'(a) & \xrightarrow{I'_a} & I'(b)
 \end{array}$$

Both maps $\lim f \circ u_{I, \phi(a)}$ and $u_{I', a} \circ f_a$ complete the diagram. Thus by uniqueness of the map corresponding to limit, we can conclude that the functions are same. \square

For composition of two limits we have following result:

Lemma 2.29. *Let (I, λ) , (I', λ') and (I'', λ'') be inverse systems with limits and let $\phi : \lambda' \rightarrow \lambda$ and $\phi' : \lambda'' \rightarrow \lambda'$ functors between systems. Let $\{f_a : I(\phi(a)) \rightarrow I'(a) \mid a \in \phi'(\lambda'')\}$ and $\{g_a : I'(\phi(a)) \rightarrow I''(a) \mid a \in \lambda''\}$ be families of morphisms corresponding to ϕ and ϕ' . Let $\lim(f \circ g) : I \rightarrow I''$ be the limit of the family $\{g_a f_{\phi(a)} : I(\phi\phi'(a)) \rightarrow I''(a) \mid a \in \lambda''\}$. Then $\lim(f \circ g) = \lim f \circ \lim g$ holds.*

Proof. To prove this we will use previous lemma and following commutative diagram:

$$\begin{array}{ccccc}
 I(\phi(\phi'(a))) & \xrightarrow{f_{\phi'(a)}} & I'(\phi'(a)) & \xrightarrow{g_a} & I''(a) \\
 \downarrow u_{I, \phi\phi'a} & & \downarrow u_{I', \phi'a} & & \downarrow u_{I'', a} \\
 \lim I & \xrightarrow{\lim f} & \lim I' & \xrightarrow{\lim g} & \lim I''
 \end{array}$$

By definition of limit morphism the condition $\lim(f \circ g) \circ u_{I, \phi(\phi'(a))} = u_{I'', a} \circ (g_a \circ f_{\phi(a)})$ holds for the limit of the composition. By uniqueness of the limit homomorphism it is enough to show that the composition of two limit functors satisfies the condition. This follows from the following equations:

$$\lim g \circ \lim f \circ u_{I, \phi(\phi'(a))} = \lim g \circ u_{I', \phi'(a)} \circ f_{\phi'(a)} = u_{I'', a} \circ g_a \circ f_{\phi'(a)}$$

\square

Now we would like to define limit for the system of homomorphisms between inverse systems of groups.

Lemma 2.30. *Let (I, λ) and (I', λ') be inverse systems of groups. Then*

$$f : \varinjlim I \rightarrow \varinjlim I' : f(x) = \prod_{a \in \lambda} f_a(\text{proj}_{\phi(a)}(x))$$

is a well defined homomorphism between the limit groups.

Proof. We have to show that the image is in the subgroup

$$L = \{x \in \prod_{i \in \lambda'} I'(i) \mid x_a = I_a^b x_b \text{ for all } a \leq b\}.$$

Let b be some index, then: $I_a^b x'_b = I_a^b f_b(x_{\phi(b)}) = f_b I_{\phi(a)}^{\phi(b)}(x_{\phi(b)}) = f_a x_{\phi(a)} = x'_a$. In this equation we used the property which says that the morphisms define a natural transformation. \square

2.2.4 Finality properties of subsystems

Final and cofinal functors can be defined in a more general way [5]. However, because we are mainly interested in properties of systems, in this section we will investigate the properties of those specific functors only in that special case. In the case of general categories, cofinality includes a concept of connectedness defined in [5] page 217. In our definition we will use a stronger version of that concept.

Definition 2.31. Let λ be a category with a quasi-order relation on its objects. We say that λ is directed category, if for every two objects $a \in \lambda$ and $b \in \lambda$ we find object $c \in \lambda$ for which relations $a \leq b$ and $a \leq c$ hold.

Definition 2.32. Let λ be a category with a quasi-order relation on its objects and λ' its directed subcategory for which following condition holds:

$$\text{For every } a \in \lambda \text{ there exists } b \in \lambda' \text{ for which } a \leq b.$$

Then we say that λ' is dense in λ .

Definition 2.33. Let λ be a directed category with a quasi-order relation on its objects and let λ' be its subcategory. Then we say that λ' is a cofinal subcategory of λ , if λ' is dense in λ .

Lemma 2.34. *Let λ be a directed category and λ' its dense subcategory. Then λ' is directed category.*

Proof. Let α and β be objects of λ . Then there exists $t \in \lambda$ for which $\alpha \leq \gamma$ and $\beta \leq \gamma$ hold. By density property there exists $\gamma' \in \lambda'$ for which condition $\gamma \leq \gamma'$ holds. Now claim follows directly from the equations $\alpha \leq \gamma \leq \gamma'$ and $\beta \leq \gamma \leq \gamma'$. \square

Now for the inverse system we have the following theorem:

Theorem 2.35. *Let (I, λ) be an inverse system and let (I', λ') be its subsystem generated by a cofinal subcategory λ . Assume that there exists a limit object for I' . Then the limit of the system (I, λ) is $\lim I'$.*

Proof. Let $\lim I'$ be the limit of the inverse system I' . It is enough to show that $\lim I'$ is also the limit of I . We will first show that $\lim I'$ forms a cone in I and then prove that for any other cone induced by an object N in I there exists a unique morphism $u : N \rightarrow \lim I'$.

Let a be an arbitrary object of the indexing category λ . Then by the definition of cofinal subcategory there exists element $a' \in \lambda'$ for which $a \leq a'$ holds. Thus there exists a morphism $I_a^{a'}$. Because $\lim I'$ is the limit of the inverse system I' it forms a cone $\triangle(I', \lim I', \rho)$, where ρ is some collection of morphisms $\{\rho_a : \lim I' \rightarrow I'(a)\}_{a \in \text{ob}(\lambda')}$. Using ρ we form a new collection of morphisms ρ' defined by $\{I_a^{a'} \circ \rho_{a'}\}_{a \in \text{ob}(\lambda')}$. Because $I_a^{a'} \circ \rho_{a'}$ can vary on the choice of a , the composition may be not uniquely determined. However, if a'' is another object of λ' for which the relation $a \leq a''$ holds, by lemma 2.34 there exists an object $t \in \lambda'$ for which the conditions $a \leq t$ and $a'' \leq t$ are satisfied. Because the following equations hold

$$I_a^{a'} \circ \rho_{a'} = I_a^{a'} \circ I_{a'}^t \circ \rho_t = I_a^t \circ \rho_t = I_a^{a''} \circ I_{a''}^t \circ \rho_t = I_a^{a''} \circ \rho_{a''}$$

we can conclude that $I_a^{a'} \circ \rho_{a'} = I_a^{a''} \circ \rho_{a''}$ and thus the family of homomorphisms is well-defined and $\lim(I')$ forms a cone $\triangle(I, \lim I', \rho')$

Now if there exists another cone $\triangle(I, N, \omega)$, we have to construct a morphism $u : N \rightarrow \lim I'$ and prove that it is unique. Let a and b be objects of λ , then there exists objects a' and b' in λ' for which relations $a \leq a'$ and $b \leq b'$ hold. Restricting the collection ω to the morphisms indexed by λ' we get the cone $\triangle(I', N, \omega')$ of I' . Because $\lim I'$ is the limit of the functor I' , there exists a unique homomorphism $u : N \rightarrow \lim I'$. It is left to prove

that the following diagram commutes:

$$\begin{array}{ccccc}
 & & N & & \\
 \omega_a \swarrow & & \downarrow & \searrow \omega_{b'} & \omega_b \\
 & \lim I' & & & \\
 \omega_{a'} \swarrow & & \rho'_a & \searrow \rho'_b & \\
 I'(a') & \xrightarrow{I_{b'}^{a'}} & I'(b') & & \\
 I_a^{a'} \swarrow & & & \searrow I_b^{b'} & \\
 I(a) & \xrightarrow{I_b^a} & I(b) & &
 \end{array}$$

Epecially we have to prove that $I_a^{a'} \circ \omega_{a'} = \omega_a$. This claim follows directly from the fact that $\triangle(I, N, \omega)$ is a cone. Thus the u defined above is the map we were searching for. \square

2.3 Nerve of covering

In this chapter we will give a definition of an abstract simplicial complex.

2.3.1 Abstract simplicial complex

We assume that the reader knows already basic facts about simplicial complex. If not we suggest to take a look at [1].

Definition 2.36. Let S be some simplex and let X be a collection of its faces. We define operator $\triangleright : 2^X \rightarrow X$ to denote a simplex which is obtained by the intersection of all simplexes for which the union of all the faces in the domain are contained.

Definition 2.37. Let X be a set and let $\mathcal{A} = \{S_i\}_{i \in I}$ be some collection of finite subsets of the set X . We say that \mathcal{A} forms an abstract simplicial complex over X , if a subset of any set S_i in the collection \mathcal{A} is in the collection.

Definition 2.38. Assume that the set X together with the collection $\mathcal{A} = \{S_i\}_{i \in I}$ is an abstract simplicial complex. Let A be some subset of X and $\mathcal{A}' = \{S'_j\}_{j \in I'}$ a collection of subsets of the space A for which the condition

$$S'_j \in \mathcal{A}' \Leftrightarrow S'_j \subset S_i \text{ for some } i \in I$$

holds. Then we say that $\Gamma(A)_{\mathcal{A}'}$ is a subcomplex of $\Gamma(X)_{\mathcal{A}}$.

From now on we will denote the abstract simplicial pair $(\Gamma(X)_{\mathcal{A}}, \Gamma(A)_{\mathcal{A}'})$ simply as $\Gamma(X, A)_{\alpha}$, where α is the pair of collections $(\mathcal{A}, \mathcal{A}')$. In the following example we will see that a simplicial complex defined in the geometric way is an abstract simplicial complex.

Example 2.39. Let s be a simplicial structure, with edge set $\text{edge}(s)$ and simplices(s) a collection of simplices. Then the abstract simplicial complex is obtained by forgetting the geometric structure.

We will now define corresponding geometric structure for an abstract simplicial complex. This will give us the interface of original simplicial complex and will allow us to use the theory behind it.

Definition 2.40. Let J be a set and $\Gamma(J)_{\mathcal{A}}$ abstract simplicial complex induced by vertex set J and some collection of abstract simplices \mathcal{A} . Then the geometric realization of the abstract simplicial complex $\mathcal{S}(J)_{\mathcal{A}}$ is a subset of the set $[0, 1]^J$ for which the following conditions are satisfied: Let $x \in \mathcal{S}(J)_{\mathcal{A}}$ and let $\text{pr}_j : [0, 1]^J \rightarrow [0, 1]$ be the projection $p_j(x) = x_j$ then:

- (1) Relation $\text{pr}_j(x) > 0$ holds only for finitely many $j \in J$.
- (2) $\sum_{j \in J} \text{pr}_j(x) = 1$.
- (3) The set $\{j \in J \mid \text{pr}_j(x) > 0\}$ belongs to collection \mathcal{A} .

For the abstract simplicial pair $\Gamma(X, A)_{\alpha}$, we use the definition of geometric realization $\mathcal{S}(X, A)_{\alpha}$ separately for $\Gamma(X)_{\mathcal{A}}$ and $\Gamma(A)_{\mathcal{A}'}$ and treat set $[0, 1]^A$ as subset of $[0, 1]^X$, where for every element $a \in [0, 1]^A$ the projection $\text{pr}_i(a) = 0$ for every index $i \in X \setminus A$.

Lemma 2.41. *The geometric realization $\mathcal{S}(X, A)_{\alpha}$ of an abstract simplicial complex is a simplicial complex.*

Proof. The following conditions are satisfied:

- (1) The collection of simplexes of the simplicial complex is the geometric realizations of sets S_i of the collection \mathcal{A} . Respectively the subsimplex consists of the geometric realizations S'_i of collection \mathcal{A}' .
- (2) Any face is in the collection by definition of abstract simplicial complex
- (3) Intersection of any two simplexes in the collection is clearly a face or the empty set.

Thus $\mathcal{S}(X, A)_{\alpha}$ is simplicial complex. □

Next we will recall the definition of a simplicial map.

Definition 2.42. Let (X, A) and (Y, B) be pairs of simplicial complexes and let f be a continuous function between pairs. Then we say that the map f is simplicial if for any simplex in X the images of vertexes of the simplex span some simplex in Y and the images of vertexes of a simplex in A span some simplex in B .

Lemma 2.43. Let (X, A) and (Y, B) be pairs of abstract simplicial complex and let $f : (X, A) \rightarrow (Y, B)$ be a simplicial map between the complexes. Let (X', A') and (Y', B') be geometric realization of the abstract simplicial complexes (X, A) and (Y, B) . Then the map $f_* : (X', A') \rightarrow (Y', B')$ induced by images of the edges mapped by function f and defined to be piecewise linear map on all the other elements is well-defined and simplicial map.

Proof. Let x be an arbitrary element of (X, A) , by definition x has a unique representation $x = \sum_{i \in J} r_i a_i$, where r_i are coefficients and a_i belong to the set of edges. By piecewise linearity

$$f_*(x) = f_*\left(\sum_{i \in J} r_i a_i\right) = \sum_{i \in J} r_i f_*(a_i)$$

Because we assumed that the map f simplicial, we see that the sum is in the corresponding subset of $[0, 1]^J$. Thus $f_*(x)$ is uniquely determined. It is easy to see that the map is simplicial. Let $\{a_{t(1)}, \dots, a_{t(n)}\}$ be some simplex in the collection, then it is mapped to simplex $\{f(a_{t(1)}), \dots, f(a_{t(n)})\}$ by definition of the map f . \square

Definition 2.44. Let $f : (X, A) \rightarrow (Y, B)$ and $g : (X, A) \rightarrow (Y, B)$ be simplicial maps. Then we say that the maps are contiguous if for every simplex $s = \{a_1, \dots, a_n\}$ in (X, A) there exists a simplex k in (Y, B) for which relations $\text{span}(f(a_1), \dots, f(a_n)) \subset k$ and $\text{span}(g(a_1), \dots, g(a_n)) \subset k$ hold.

Lemma 2.45. If $f : (X, A) \rightarrow (Y, B)$ and $g : (X, A) \rightarrow (Y, B)$ are contiguous maps then they are homotopic.

Proof. We define $H : X \times [0, 1] \rightarrow Y$ in the following way:

$$H(x, t) = tf(x) + (1 - t)g(x).$$

It is easy to see that $H(x, 1) = f(x)$ and $H(x, 0) = g(x)$. Because $f(x)$ and $g(x)$ both belong to the same simplex, it follows that $H(x, t)$ is well defined at every point. \square

2.3.2 Coverings

We begin by recalling the definition of a covering

Definition 2.46. Let X be a topological space and let $\{A_i\}_{i \in J}$ be the collection of its open subsets. Then we say that $\{A_i\}_{i \in J}$ is a covering of X if $\bigcup_{i \in J} A_i = X$.

We will denote the set of all open coverings of the space X by $\text{COV}(X)$. Because in homology we are interested in a pair of spaces we give the definition of covering for the pair.

Definition 2.47. Let (X, A) be a topological pair. Then we say that $(\{U_i\}_{i \in J}, \{U_i\}_{i \in J'})$ is an open covering of (X, A) if the following conditions hold:

- (1) $\{U_i\}_{i \in J}$ is an open covering of X
- (2) $J' \subset J$.
- (3) $\{U_i\}_{i \in J'}$ covers A

A covering defined above will be from now on denoted as $\mathcal{C}(U_i, J, J')$

We define the collection of all coverings of the pair (X, A) to be $\text{COV}(X, A)$. Next we will see that we can define quasi order for coverings of topological pair (X, A) in a natural way. However, first we have to recall the definition of a refinement.

Definition 2.48. Let $\alpha = \mathcal{C}(U_i, I, I')$ and $\beta = \mathcal{C}(V_j, J, J')$ be coverings of some pair (X, A) . Then the covering β is called a refinement of the covering α , if for every $V_j \in \beta$ there exists some $U_i \in \alpha$ for which $V_j \subset U_i$ holds.

Now we are ready to define the quasi order in the collection of coverings of an arbitrary topological space.

Lemma 2.49. Let $\text{COV}(X, A)$ be the collection of all coverings of the pair (X, A) . Then the order \leq defined by $\alpha \leq \beta$ if and only if β is a refinement of α , is quasi-order.

Proof. Any covering is a refinement of itself and if we have coverings for which $a \leq b$ and $b \leq c$ holds, then for every set U_c we can find U_b for which $U_c \subset U_b$ and $U_b \subset U_a$ for some U_a . Then we see that $U_c \subset U_a$ so actually c is refinement of a . \square

2.3.2.1 Category of coverings

Definition 2.50. Let (X, A) be a topological pair and let α and β be its coverings for which the relation $\alpha \leq \beta$ holds. Then we define $\mathcal{P}(\alpha, \beta)$ to be the collection of all the following projection maps $p_\alpha^\beta : \beta \rightarrow \alpha$: Every $U_i \in \beta$ is mapped to some set $U'_i \in \alpha$ for which $U_i \subset U'_i$ holds.

We will now show that the family of projection maps is closed under composition.

Lemma 2.51. *Let α , β and γ be coverings of some topological pair, for which relation $\alpha \leq \beta \leq \gamma$ holds. Let p_α^β and p_β^γ be some projection maps. Then $p_\alpha^\beta \circ p_\beta^\gamma$ is also a projection map.*

Proof. Let $U \in \gamma$, then if $U' = p_\beta^\gamma(U) \in \beta$ and $U'' = p_\alpha^\beta \circ p_\beta^\gamma(U)$, by the definition condition $U \subset U' \subset U''$ holds. By properties of inclusion we can conclude that $U \subset U''$ and thus $p_\alpha^\beta \circ p_\beta^\gamma$ is a projection map. \square

Lemma 2.52. *The collection of coverings $COV(X, A)$ is a category defined in the following way:*

- (1) *The objects of $COV(X, A)$ are all open coverings of the pair (X, A) .*
- (2) *We define $Hom(\beta, \alpha)$ to be $\mathcal{P}(\alpha, \beta)$, if $\alpha \leq \beta$. Else we define $Hom(\beta, \alpha)$ to be the empty set.*
- (3) *The composition of maps is defined to be the usual composition of functions.*

Proof. By Lemma 2.51 the composition of two projection maps is a projection map and thus it is well-defined. For every α there exists a well-defined projection map p_α^α which acts like the identity map on $COV(X, A)$. The associativity condition holds, because every morphism is a function. \square

2.3.2.2 Directness property of coverings

For the order relation in the category of coverings we have following theorem

Lemma 2.53. *Let (X, A) be a topological pair. The category $COV(X, A)$ with the quasi-order defined in the definition 2.48 is directed.*

Proof. Let $\alpha = \mathcal{C}(U_i, I, I')$ and $\beta = \mathcal{C}(V_j, J, J')$ be objects of $COV(X, A)$. We will define a covering $\gamma = \mathcal{C}(W_a, I \times J, I' \times J')$ to be covering which consists of the following elements:

$$W_{(i,j)} = U_i \cap V_j$$

We will prove now that the covering constructed above satisfies the condition of this lemma. Let $W \in \gamma$, then $W = U \cap V$ for some $U \in \alpha$ and $V \in \beta$. Thus $W \subset U$ and $W \subset V$. Because we chose W to be an arbitrary element of the collection, relations $\alpha \leq \gamma$ and $\beta \leq \gamma$ hold. \square

2.3.3 The construction

Now we are ready to give the definition for the nerve of the covering.

Definition 2.54. Let X be a topological space with an open covering $\mathcal{U}(X) = \{U_i\}_{i \in J}$. Then the nerve of this covering is the following abstract simplicial complex:

- (1) The vertex set is defined to be the collection $\mathcal{U}(X)$.
- (2) Let $\text{CNI}(\mathcal{U}(X))$ be the collection of all finite non-empty intersections of sets in the collection $\mathcal{U}(X)$. The simplex set of the abstract simplicial complex is defined to be $\text{CNI}(\mathcal{U}(X))$.

The structure described above can be denoted as $\Gamma(\mathcal{U}(X))_{\text{CNI}(\mathcal{U}(X))}$. However, to simplify notation we will from now on denote the nerve of the covering as $\Gamma(X)_{\mathcal{U}(X)}$.

Lemma 2.55. *The abstract simplicial complex $\Gamma(X)_{\mathcal{U}(X)}$ described above is well-defined.*

Proof. We have to show that any subset of an element in the collection $\text{CNI}(\mathcal{U}(X))$ is in the collection. This follows from the fact that if $\bigcap_{i \in J} U_i \neq \emptyset$, then also $\bigcap_{i \in J'} U_i \neq \emptyset$ for any subcollection $J' \subset J$. \square

We will now give the definition of the nerve of the covering for topological pairs.

Definition 2.56. Let $\mathcal{L} = \mathcal{C}(U_i, J, J')$ be a covering of some topological pair (X, A) . We denote \mathcal{L}_X to be the covering taking elements from \mathcal{L} indexed by J and \mathcal{L}_A to be the covering, where we use elements from \mathcal{L} indexed by J' . Then the nerve of this covering $\mathcal{S}(X, A)_{\mathcal{L}}$ is the simplicial pair $(\mathcal{S}(\mathcal{L}_X)_{\text{CNI}(\mathcal{L}_X)}, \mathcal{S}(\mathcal{L}_A)_{\text{CNI}(\mathcal{L}_A)})$.

Lemma 2.57. *The simplicial pair described above is well-defined.*

Proof. By Lemma 2.55, the simplicial complexes are well-defined. We only have to show that $\mathcal{S}(\mathcal{L}_A)_{\text{CNI}(\mathcal{L}_A)}$ is a subsimplex of $\mathcal{S}(\mathcal{L}_X)_{\text{CNI}(\mathcal{L}_X)}$. This follows directly from the definition of the pair of coverings, as we defined the covering \mathcal{L}_A to be a sub covering of \mathcal{L}_X . \square

Lemma 2.58. *Let $\beta = \mathcal{C}(V_i, I, I')$ and $\alpha = \mathcal{C}(U_i, I, I')$ be coverings of (X, A) in such way that $\alpha \leq \beta$ and let p be a projection map belonging to $\mathcal{P}(\alpha, \beta)$. Let the map $p'_\alpha : \Gamma(X, A)_\beta \rightarrow \Gamma(X, A)_\alpha$ be a map between abstract simplicial complexes induced by the edges $V_i \in \beta$ and map p . The map $p'_\alpha : \Gamma(X, A)_\beta$ is well-defined simplicial map.*

Proof. Well-definiteness follows directly from the fact that the inducing map p is well-defined. Let s be a simplex in $\Gamma(X, A)_\beta$ spanned by some sets $\{U_i\}_{i \in J}$ such that $\bigcap_{i \in J} U_i \neq \emptyset$. Every U_i is mapped to some vertex V_i in $\Gamma(X, A)_\alpha$ for which $U_i \subset V_i$. Intersection $\bigcap_{i \in J} V_i$ is clearly non-empty, so edges of every simplex span some simplex in the image. \square

All the lemmas and proves have been presented only for the abstract simplicial complexes. The geometric realization of the nerve of the covering will be denoted simply as $\mathcal{S}(X, A)_\alpha$. The concept of projection map is being used in many different contexts. However, because it is clear from the context which definition is used in all the contexts we will denote the projection map to be simply p'_α . It is also worth noticing that from lemma 2.43 it follows that the induced map p_α^β between the geometric realizations $\mathcal{S}(X, A)_\beta$ and $\mathcal{S}(X, A)_\alpha$ is also well-defined and simplicial.

Projection maps are not uniquely determined by coverings α and β , but any of those maps are contiguous thus they define the same homomorphisms between corresponding homology groups of spaces $\mathcal{S}(X, A)_\alpha$ and $\mathcal{S}(X, A)_\beta$.

Lemma 2.59. *Let α and β be any coverings of (X, A) for which condition $\alpha \leq \beta$ is satisfied. Then any two projection maps p_α^β and p'_α^β are contiguous.*

Proof. To prove that the maps are contiguous we form a simplex in $\mathcal{S}(Y, B)$ and prove that the images of both maps belong to it. Every vertex x_i in $\mathcal{S}(X, A)$ is mapped to $x'_{s(i)}$ and $x''_{d(i)}$ by p_α^β and p'_α^β respectively. Let U_i be the open sets corresponding to vertexes x_i . Now for every simplex $\{x_{r(0)}, \dots, x_{r(n)}\}$ the corresponding intersection $\bigcap_{i \in J} U_i$ is non-empty. Let U'_i correspond to $x'_{s(i)}$ and U''_i to $x''_{d(i)}$. We see that $\bigcap_{i \in J} U'_i \cap \bigcap_{i \in J} U''_i \neq \emptyset$ and thus the corresponding simplex $\{x'_{s(r(1))}, \dots, x'_{s(r(n))}, x''_{d(r(1))}, \dots, x''_{d(r(n))}\}$ where the duplicates are removed is in the collection. In the case where the simplex belongs to $\mathcal{S}(A)$ it is easy to see that any projection map maps them to some simplex which lies inside $\mathcal{S}(B)$. \square

Corollary 2.60. *Let (X, A) be a topological pair. The projection maps in collection $\mathcal{P}_{(X, A)}(a, b)$ define same homomorphisms between singular homology groups.*

Proof. Follows directly using lemma 2.59 and the fact that contiguous simplicial maps induce the same homomorphisms between homology groups \square

Thus we can define the unique homomorphisms $I_\alpha^\beta : H_q(\mathcal{S}(X, A)_\beta) \rightarrow H_q(\mathcal{S}(X, A)_\alpha)$ for every $q \in \mathbb{N}$.

Definition 2.61. Let $F : (X, A) \rightarrow (Y, B)$ and let $\beta = \mathcal{C}(U_i, J_\beta, J'_\beta)$ be a covering of (Y, A) . The covering $f^{-1}\beta$ is a covering of (X, A) which is formed by the collection $\mathcal{C}(V_i, J_\beta, J'_\beta)$, where $V_i = f^{-1}U_i$ for every $i \in J_\beta$.

Note that the collection α above is defined in a such way, that for every element $U_i \in \beta$ we make copy, take preimage of it and add it to the collection α .

Lemma 2.62. *Let $f : (X, A) \rightarrow (Y, B)$ be a function between pairs and let $\alpha = \mathcal{C}(U_i, I, I')$ be a covering of (Y, B) . Let β be the covering $f^{-1}\alpha$ of (X, A) . Then the induced map $f_* : \mathcal{S}(X, A)_\alpha \rightarrow \mathcal{S}(Y, B)_\beta$, defined by $f^{-1}U_i \rightarrow U_i$ is well-defined.*

Proof. Let $\bigcap_{i \in I} f^{-1}U_i \neq \emptyset$ for some index set $I \subset J$. We see that $\bigcap_{i \in I} U_i \neq \emptyset$. Thus the lemma holds. \square

For projection maps and induced homomorphisms we have following useful lemma.

Lemma 2.63. *Let α be a covering of the pair (X, A) and let β be a covering of the pair (Y, B) for which $\alpha \leq \beta$ holds and let $f : (X, A) \rightarrow (Y, B)$ be a function between pairs. Then the pre-image $f^{-1}\beta$ of the covering β is a refinement of $f^{-1}\alpha$.*

Proof. This follows directly from the fact that if $U \subset V$, then also $f^{-1}U \subset f^{-1}V$. \square

Lemma 2.64. *Let $f : (X, A) \rightarrow (Y, B)$ and let α and β be coverings of (Y, B) such that $\alpha \leq \beta$. Let $\alpha' = f^{-1}\alpha$ and let $\beta' = f^{-1}\beta$ and let p_α^β be a projection map for the coverings. Then there exists a projection map $p_{\alpha'}^{\beta'}$ for which the following diagram commutes.*

$$\begin{array}{ccc} \mathcal{S}(X, A)_{\beta'} & \xrightarrow{\quad p_{\alpha'}^{\beta'} \quad} & \mathcal{S}(X, A)_{\alpha'} \\ \downarrow f_\beta & & \downarrow f_\alpha \\ \mathcal{S}(Y, B)_\beta & \xrightarrow{\quad p_\alpha^\beta \quad} & \mathcal{S}(Y, B)_\alpha \end{array}$$

Proof. Lemma 2.63 says that there exists a projection map $p_{\alpha'}^{\beta'}$, \square

Chapter 3

Cech homology

In this chapter we will construct Cech homology and prove the main results related to it. We will first define an inverse system of homology groups of the nerves.

Definition 3.1. Let α be an element of $\text{COV}(X, A)$ and let G be an abelian group. Then we can define the group $H_{q,\alpha}(X, A; G)$ using the singular homology groups in the following way:

$$H_{q,\alpha}(X, A; G) = H_q(\mathcal{S}(X, A)_\alpha; G).$$

For all coverings of α, β in (X, A) with $\alpha \leq \beta$ we define the maps to be

$$I_\alpha^\beta : H_{q,\beta} \rightarrow H_{q,\alpha}$$

maps I_β^α are induced by some maps p_α^β which are unique up to homotopy.

Lemma 3.2. Let $q \in \mathbb{N}$ be a fixed number. Let (X, A) be a topological pair and let G be an abelian group. Then the groups $H_{q,\alpha}(X, A; G)$ together with maps I_β^α form an inverse system of groups.

Proof. Lemma 2.52 says that $\text{COV}(X, A)$ with the morphisms induced by the quasi-order relation forms a category. Let I be a functor with maps every covering to a corresponding geometric realization of the nerve and every morphism between the coverings to the induced homomorphism. Using lemmas 2.60 and 2.22 we see that the functor defined in this way is a well-defined functor and forms an inverse system. \square

Now we are ready to define the Cech homology groups \check{H}_n .

Definition 3.3. Let (X, A) be a topological pair and G an abelian group. Then Cech homology groups with coefficients G are defined to be inverse limits of $H_{q,\alpha}(X, A; G)$ where α belongs to $\text{COV}(X, A)$. Simply denoted $\check{H}_q(X, A; G) = \varprojlim \{H_{q,\alpha}(X, A; G)\}$

Because coverings can be infinite the limits are not necessarily defined. However there is a way to evade this problem.

3.1 Well-definedness of the Čech homology groups

In this section we will provide condition for the spaces which will guarantee that the Čech groups are well-defined. ... From now on in this chapter we assume that the topological spaces are such for which the Čech groups are well-defined.

3.2 Induced homomorphisms

In this section we will investigate properties of induced homomorphism. Earlier we defined Čech homology groups to be limit of the groups induced by coverings of the pair. In section 2.2.3 we defined concept of limit homomorphism. In next theorem we will show that such homomorphism exists.

Theorem 3.4. *Let $f : (X, A) \rightarrow (Y, B)$ be continuous function between pairs of spaces. Let $\mathcal{A} = \{f_\alpha \mid \alpha \text{ is covering of pair } (Y, B)\}$ be family of functions for which every element of the open covering α is mapped to its preimage. This family induces maps between corresponding Čech homology groups.*

$$f'_{\alpha,q} : H_q(\mathcal{S}(X, A)_{f^{-1}\alpha}; G) \rightarrow H_q(\mathcal{S}(Y, B)_\alpha; G)$$

We denote a family of functions indexed by coverings of (Y, B) described above with \mathcal{A}_q . Then every such collection is natural transformation of the Čech homology groups and the limit function is well defined.

Proof. We can define a functor $\phi : \text{COV}(Y, B) \rightarrow \text{COV}(X, A)$ to map every object α to $f^{-1}\alpha$. By lemma 2.63 this functor preserves order. Let β and α be some coverings of pair (Y, B) . We denote α' and β' to be the preimage categories of α and β under map f . The diagram of lemma 2.64 shows that the corresponding maps between coverings commute. Thus we have a following commutable diagram:

$$\begin{array}{ccc} H_q(\mathcal{S}(X, A)_{\beta'}) & \xrightarrow{I_{\alpha'}^{\beta'}} & H_q(\mathcal{S}(X, A)_{\alpha'}) \\ \downarrow f'_{\beta,q} & & \downarrow f'_{\alpha,q} \\ H_q(\mathcal{S}(Y, B)_\beta) & \xrightarrow{I_\alpha^\beta} & H_q(\mathcal{S}(Y, B)_\alpha) \end{array}$$

The fact that collection \mathcal{A}_q is natural transformation of the groups follows directly from the above diagram. Thus by lemma 2.30 there exists limit homomorphism

$$f_q : \check{H}_q(X, A; G) \rightarrow \check{H}_q(Y, B; G)$$

□

We will show now that the Čech homology group is a functor.

Theorem 3.5. *Cech homology groups are functors from the category of the topological pairs to the category of abelian groups*

Proof. (1) Every pair of spaces is mapped uniquely to some abelian group.

(2) The limit of identity map is an identity homomorphism.

(3) Let $f : (X, A) \rightarrow (Y, B)$ and $g : (Y, B) \rightarrow (Z, C)$ be continuous maps between pairs. We know that $(g \circ f)_* = g_* \circ f_*$ holds for the singular homology groups. Then by using theorem 2.29 we see that the limit homomorphism $\lim((g \circ f)_*)$ equals $\lim g_* \circ \lim f_*$.

□

3.3 Dimension axiom

In this section we will prove that the Čech homology theory satisfies the dimension axiom.

Theorem 3.6. *Let $X = \{x\}$ be a space which consist of one point. Then the Čech homology groups $\check{H}_n(X; G)$ are trivial groups for $n \geq 1$ and isomorphic to G when $n = 0$.*

Proof. For the space X there exists only one open covering $\alpha = \{x\}$ and thus the inverse system of the corresponding singular homology groups is trivial. It is easy to see that the limit of this system is the group $H_q(\mathcal{S}(X)_\alpha; G)$ together with the identity homomorphisms. The simplex $\mathcal{S}(X)_\alpha$ consists only of one edge and thus is homeomorphic to a one point space. From the dimension axiom of singular homology it is easy to see that

$$\check{H}_n(X; G) = \begin{cases} G & \text{if } n = 0 \\ 0 & \text{if } n \geq 1 \end{cases}$$

□

3.4 Homotopy axiom

The main result of this section is the homotopy axiom of Čech homology which states that if two spaces are homotopic then their all Čech homology groups are isomorphic. In the beginning we will prove an important result regarding the singular homology groups of the nerve of the unit interval $[0, 1]$.

Definition 3.7. Let X be a topological space and α its finite open covering. Then $CLEAN(\alpha)$ is a covering of space X , which is formed by removing every such element $A \in \alpha$ for which $A \subset B$ holds for some $B \in \alpha$ from the collection.

Lemma 3.8. Let X be a topological space and α its finite open covering. Let $\beta = CLEAN(\alpha)$ be the covering defined in previous definition. Then singular homology $H_q(\mathcal{S}(X)_\alpha)$ and $H_q(\mathcal{S}(X)_\beta)$ are isomorphic for every $q \in \mathbb{N}$.

Proof. Because β is subcollection of α , the relation $\alpha \leq \beta$ holds. Relation $\beta \leq \alpha$ holds, because if we take any element from $a \in \alpha$ which is not in β by definition we will find an element $b \in \beta$ for which the relation $a \subset b$ holds.

Thus there exists morphisms I_α^β and I_β^α in the inverse system of abelian groups induced by coverings. By definition equations $I_\alpha^\beta \circ I_\beta^\alpha = I_\beta^\beta = id_\beta$ and $I_\beta^\alpha \circ I_\alpha^\beta = I_\alpha^\alpha = id_\alpha$ hold. Thus I_α^β is an isomorphism between $H_q(\mathcal{S}(X)_\alpha)$ and $H_q(\mathcal{S}(X)_\beta)$. \square

Definition 3.9. Let α be a cleaned finite open covering of $[0, 1]$, for which every element in the collection is connected. Trivial indexation of α is the following indexation: We order open intervals in the collection $(a, b) \in \alpha$ by the coordinate a and number coverings.

Lemma 3.10. Let α be a open covering of $[0, 1]$ which is cleaned, finite and connected. Then the trivial indexation $[m]$ preserves order of the coordinate b of open intervals.

Proof. If $b_n > b_{n+1}$ for some $n \in [m - 1]$, then $(a_{n+1}, b_{n+1}) \subset (a_n, b_n)$ which is a contradiction by the definition of the covering α . \square

Lemma 3.11. Let α be an open covering which is finite and every element in the collection is connected. Then every singular homology group of the nerve $\mathcal{S}([0, 1])_\alpha$ is trivial.

Proof. Using 3.8 we can reduce our problem to coverings which have property that no inclusion holds for any two element of the collection α . We define indexation to be trivial indexation for covering α . Let $\{f_n(x) : \mathcal{S}([0, 1])_\alpha \rightarrow \mathcal{S}([0, 1])_\alpha\}_{n \in [m]}$ be a collection of simplicial maps which are induced by the edges in the following way:

$$f_n(v_i) = \begin{cases} v_i & \text{if } i \leq n \\ v_n & \text{if } i \geq n \end{cases}$$

We will now show that the maps defined above induce same homomorphisms between the corresponding groups. Because the indexation set is finite, it is enough to show that f_n and f_{n+1} are contiguous for every n in the collection. Let s be some simplex in the nerve $\mathcal{S}([0, 1])_\alpha$. If simplex s is spanned only by elements whose index is less than n , maps f_n and f_{n+1} are both identity on it. Now assume that s contains atleast one edge the index of which is larger than $n - 1$. Then the image of the element s under the map f_n can be

presented as $\triangleright(s', v_n)$, where s' is simplex for which index of every edge is smaller than n . There are two possible representations for the image $f_{n+1}(s)$ namely $\triangleright(s', v_n, v_{n+1})$ and $\triangleright(s v_{n+1})$. If the representation of $f_{n+1}(s)$ is $\triangleright(s', v_n, v_{n+1})$, we can choose the simplex to be $\triangleright(s', v_n, v_{n+1})$, as it clear from definition that $\triangleright(s', v_{n+1})$ is its face. In the second case we notice that because intersection $\bigcap_{a \in s'} a \cap v_{n+1}$ is non-empty and connected. Thus it is an open interval (a_{n+1}, b_1) . Because condition $(a_n, b_n) \cap (a_{n+1}, b_1) \neq \emptyset$ holds, we can choose $\triangleright(s', v_n, v_{n+1})$ to be the simplex where image of s of the both functions are contained. \square

Next we define a class of nice coverings for the unit interval $[0, 1]$ and prove that the subcategory which consists of the coverings defined is a subcategory of the Čech inverse system.

Definition 3.12. Let $\alpha = \{U_i\}_{i \in [n]}$ be a connected open covering of unit interval $[0, 1]$ for which the following conditions are satisfied:

- (1) $0 \in U_0$ and $1 \in U_n$
- (2) $U_i \cap U_{i+1} \neq \emptyset$ for every $i \in [n - 1]$
- (3) $U_i \cap U_j = \emptyset$ for all $|i - j| \geq 1$

Then we say that covering α is regular.

Lemma 3.13. *The category of the regular coverings of $[0, 1]$ is cofinal in the $COV([0, 1])$.*

Proof. Let α be arbitrary covering of $[0, 1]$. We modify the covering in the following ways:

- (1) We form covering β by selecting some finite subcovering of α .
- (2) We split every non-connected element in the collection β into connected elements.
- (3) We search for the removable elements i.e the the collection after removing the element covers the space $[0, 1]$. We remove every such element and stop this process after no removable elements are found.
- (4) We define the indexation to be the trivial indexation.

The condition (1) and (2) are satisfied, because γ is trivially indexed covering. Condition (3) is satisfied, because if two open intervals with are not neighbors have non-empty intersection then all the coverings between can be removed and that collection will still cover the set. Because covering γ is obtained by removing and cutting elements, the relation $\beta \leq \gamma$ holds. \square

Definition 3.14. Let (X, A) be topological pair and $\alpha = \{(V_i, U_i)\}_{i \in J}$ its covering. Suppose that for every V_x there is a regular covering $\beta(x)$ of an unit interval $[0, 1]$ indexed by the set $[n]$ for some $n \in \mathbb{N}$. Covering γ of the pair $(X \times [0, 1], A \times [0, 1])$ defined in the following way:

$$\gamma = \{(V_x \times \beta(x), V_x \times \beta(x))\}_{x \in J}$$

is called a stacked covering over α .

Lemma 3.15. *The category of stacked coverings is cofinal in $COV((X \times [0, 1], A \times [0, 1]))$*

Proof. Let $\gamma = (V_i, U_i)_{i \in J}$ □

Chapter 4

Applications

4.1 Lefschetz fixed-point theorem

[Theory of this will be presented later]

4.2 Kakutani fixed point theorem

The fixed point theorem of Kakutani can be derived for Lefschetz fixed-point theorem.

Definition 4.1. Let X be vector space with topological structure and K topological field. If vector addition $+: X \times X \rightarrow X$ and scalar multiplication $\cdot: K \times X \rightarrow X$ are continuous we say that X is topological vector space over field K or shortly $L(X, K)$

Topological vector space has following property:

Theorem 4.2. *Kakutani fixed point theorem. Let S be a non empty, compact and convex subset of a locally convex Hausdorff space. Let $f: S \rightarrow 2^S$ be a multivalued function on S which has closed graph and the property that $f(x)$ is convex and non-empty for all $x \in S$. Then the set of fixed points of f is non-empty and compact.*

4.3 Continuous game theory

In this section we will apply theory developed in previous chapters to game theory. We define n player continuous game structure $Cgs(J, \mathcal{A})$ in a following way:

Definition 4.3. Let J be finite indexing set and \mathcal{A} collection of continuous utility functions $\{u_i: X_i \rightarrow \mathbb{R}\}_{i \in J}$ with a compact domain. Then we say that $Cgs(J, \mathcal{A})$ is a continuous game.

Now we give definition of Nash equilibrium for Cgs.

Definition 4.4. Let $Cgs(J, \mathcal{A})$ be continuous game we say that point $x^* \in \prod_{i \in J} X_i$ is Nash equilibrium if for every $k \in J$ and every $x'_k \in X_k$ following condition holds:

$$u_k(x^*) \geq u_k(x_1, \dots, x'_k, \dots, x_n)$$

Next we will prove important lemma which says that game has equilibrium only and only if specific function has fixed point.

Lemma 4.5. $Cgs(n, \mathcal{A})$ has Nash equilibrium only and only if following set value function

$$g : \prod_{i \in J} X_i \rightarrow \prod_{i \in J} 2^{X_i} : g_k(x) = \operatorname{argmax}_{t \in X_k} u_k(x_1, \dots, x_k, t, x_{k+1}, \dots, x_n)$$

has a fixed point $x \in g(x)$

Proof. Trivial (Just check definitions) □

4.3.1 General version of resource optimization problem

Consider a game where a set of players compete in a multiple competitions at the same. Every player has limited energy and they can allocate it freely. Every competition is won by a player who allocated most energy in it and the goal of the players is to maximize the amount of the competitions they win.

Definition 4.6. We define the continuous game structure of the game described above as follows:

1. The choice space for every player consists of continuous functions $f : [0, 1] \rightarrow \mathbb{R}$ for which $\int_0^1 f(x) dx = 1$
2. The evaluation function $u_n(x) : \prod_{i \in J} X_i \rightarrow [0, 1]$ is defined to be:

$$u_n(f) = m^*(\{x \in [0, 1] \mid f_n(x) \geq f_i(x)\})$$

We will show that in some cases there exists an equilibrium.

Appendices

Appendix A

Homology axioms

For homology theories we have a following interface:

Definition A.1. A homology theory consists of a family of the functors from category of topological pairs to category of abelian groups $\{H_n\}_{n \in \mathbb{N}}$ and natural transformations $\{\delta_n : H_n(X, A) \rightarrow H_{n-1}(A, \emptyset)\}_{n \in \mathbb{N}}$. For the homological theory the following axioms are satisfied:

1. text here

Bibliography

- [1] J. Rotman An Introduction to Algebraic Topology (Graduate Texts in Mathematics), Springer
- [2] L. Górniewicz, Topological Fixed Point Theory of Multivalued Mappings, Kluwer Academic Publishers, 1999.
- [3] I. L. Glicksberg, A further generalization of the Kakutani fixed point theorem with application to Nash equilibrium points, Proc. Amer. Math. Soc. 3 (1952), 170–174.
- [4] M. BAYE, G. TIAN, ANU J. ZHOU, “The Existence of Pure Nash Equilibrium in Games with Nonquasiconcave Payoffs,” Texas A M University, mimeo, 1990.
- [5] Saunders Mac Lane: Categories for the Working Mathematician (Graduate Texts in Mathematics)
- [6] Joke Häsa: Algebra II, 2016.