Final Exam Problems

MATH 598 / 784 · High-Dimensional Probability

November 23–27, 2026

Prepare all eight problems. The oral exam consists of two randomly selected problems presented at the blackboard in **Burnside 925**. There are **20 minutes for the entire examination**, within your booked 30-minute appointment. Arrive at the start of your appointment. The instructor may ask you to explain steps, justify assumptions, or discuss the cases specified in the problems. You may cite the facts explicitly supplied below without proving them.

Each of the two presentations is worth up to 25 points. A replacement problem costs 5 points; the total score is at least zero. For the complete procedure, rubric, and booking information, see the final exam page.

Throughout, $C, c > 0$ denote absolute constants, whose values may change. All logarithms are natural. Write $B_p^n = \{x \in \mathbb{R}^n : \|x\|_p \le 1\}$, and, for a nonempty bounded set T in a Euclidean space, write $w(T) = \mathbb{E}\sup_{t\in T}\langle g, t\rangle$ for standard Gaussian g. A standard Gaussian matrix has independent $N(0,1)$ entries; $\|A\|_F^2 = \sum_{i,j} A_{ij}^2$ denotes its squared Frobenius norm.

1. Estimating a trace without reading the matrix

Let A be a nonzero positive semidefinite $n \times n$ matrix. Suppose that we can compute Av for a chosen vector v, but cannot read the entries of A directly. Let $g \sim N(0, I_n)$ and let $\varepsilon \in \{-1, 1\}^n$ have independent uniform sign coordinates.

1. Show that $g^{\mathsf{T}} A g$ and $\varepsilon^{\mathsf{T}} A \varepsilon$ are unbiased estimators of $\operatorname{tr} A$, and prove
$$\operatorname{Var}(g^{\mathsf{T}} A g) = 2\|A\|_F^2, \qquad \operatorname{Var}(\varepsilon^{\mathsf{T}} A \varepsilon) = 2\sum_{i \ne j} A_{ij}^2.$$
2. Choose either estimator. Form b independent batches, each containing k independent copies, and let $\widehat{\tau}$ be a median of the b batch averages. For $0 < \delta, \rho < 1$, prove that
$$k \ge C\delta^{-2}, \qquad b \ge C\log(2/\rho) \quad \Longrightarrow \quad \mathbb{P}\{|\widehat{\tau} - \operatorname{tr} A| > \delta \operatorname{tr} A\} \le \rho.$$
Thus only $O(\delta^{-2}\log(2/\rho))$ matrix-vector products are needed, independently of n.

Hint. Use positivity of the eigenvalues to bound the variance relative to $(\operatorname{tr} A)^2$, then control how many batch averages are inaccurate.

2. Encoding a finite set with random hyperplanes

Let $T \subset S^{n-1}$ have $N \ge 2$ points. Take independent $g_1, \dots, g_m \sim N(0, I_n)$ and encode $x \in T$ by
$$\Phi(x) = \big(\operatorname{sign}\langle g_i, x\rangle\big)_{i=1}^m \in \{-1, 1\}^m,$$
where $\operatorname{sign}(0) = 1$. Write $d_H(u, v) = m^{-1}|\{i : u_i \ne v_i\}|$ and $d_{\mathrm{ang}}(x, y) = \pi^{-1}\arccos\langle x, y\rangle$.

1. Prove that $\mathbb{E}\, d_H(\Phi(x), \Phi(y)) = d_{\mathrm{ang}}(x, y)$ for every $x, y \in T$.
2. For $0 < \delta, \rho < 1$, prove that $m \ge C\delta^{-2}\log(N/\rho)$ implies, with probability at least $1 - \rho$,
$$\sup_{x,y\in T} |d_H(\Phi(x), \Phi(y)) - d_{\mathrm{ang}}(x, y)| \le \delta.$$
Deduce that on this event any two points assigned the same code have Euclidean distance at most $\pi\delta$.

Hint. Use rotational invariance in the plane spanned by x and y.

3. The ground-state energy of a Gaussian spin system

For $n \geq 2$, let $(Z_{ij})_{1 \leq j < i \leq n}$ be independent standard normal random variables. For $\sigma \in \{-1, 1\}^n$, define

$$H_\sigma = \sum_{i>j} Z_{ij}\sigma_i\sigma_j, \qquad E_n = \min_{\sigma \in \{-1,1\}^n} H_\sigma.$$

1. Prove that $\mathbb{E}E_n = -\mathbb{E}\max_\sigma H_\sigma$ and that $-\mathbb{E}E_n \leq Cn^{3/2}$.
2. Construct signs successively by taking $\sigma_1 = 1$ and choosing σ_i to maximize $\sigma_i \sum_{j<i} Z_{ij}\sigma_j$. Condition on the previous rows to show that the expected value attained is $\sqrt{2/\pi}\sum_{i=2}^n \sqrt{i-1}$. Deduce $cn^{3/2} \leq -\mathbb{E}E_n$.
3. Prove that
$$\mathbb{P}\left\{\left|\frac{E_n}{\mathbb{E}E_n} - 1\right| > \delta\right\} \leq 2e^{-c\delta^2 n} \qquad (\delta > 0).$$
In particular, $E_n/\mathbb{E}E_n \to 1$ in probability.

Facts you may use. For centered Gaussian variables with variances at most v^2, $\mathbb{E}\max_{1 \leq j \leq N} X_j \leq v\sqrt{2\log N}$, without any independence assumption.

4. Recovering a signal from noisy measurements

An unknown signal x belongs to a known nonempty compact set $T \subset \mathbb{R}^n$. We observe

$$y = Ax + \eta,$$

where A is an $m \times n$ standard Gaussian matrix and $\eta \in \mathbb{R}^m$ is a fixed noise vector. Reconstruct x by choosing any measurable constrained least-squares solution

$$\widehat{x} \in \arg\min_{z \in T} \|Az - y\|_2.$$

The expectation below is over A; x and η are fixed.

1. Prove that
$$\mathbb{E}\|\widehat{x} - x\|_2 \leq \frac{Cw(T) + 2\|\eta\|_2}{\sqrt{m}}.$$
2. If the only prior information is $\|x\|_1 \leq R$ for $R > 0$, take $T = RB_1^n$ and deduce
$$\mathbb{E}\|\widehat{x} - x\|_2 \leq \frac{CR\sqrt{\log(2n)} + 2\|\eta\|_2}{\sqrt{m}}.$$

Hint. Apply the following **Gaussian deviation theorem** to $T - T$: for a compact symmetric set S containing 0,

$$\mathbb{E}\sup_{h \in S} \left| \|Ah\|_2 - \sqrt{m}\|h\|_2 \right| \leq Cw(S).$$

You may use this theorem without proof.

5. Random sections of the octahedron

Let $1 \le d < n$, let A be a $d \times n$ standard Gaussian matrix, and let $E = \ker A$. All radii and diameters below are Euclidean.

1. Put $T_r = B_1^n \cap r B_2^n$. If $n^{-1/2} \le r \le 1$, prove
$$w(T_r) \le C\sqrt{\log(enr^2)}.$$
You may use $\mathbb{E}(\sum_{i=1}^{s}(g_i^*)^2)^{1/2} \le C\sqrt{s\log(en/s)}$, where $g_1^* \ge \cdots \ge g_n^*$ are the decreasing rearrangement of $|g_1|, \dots, |g_n|$.
2. Prove the Garnaev–Gluskin bound
$$\mathbb{E}\,\mathrm{diam}(B_1^n \cap E) \le C\sqrt{\frac{\log(en/d)}{d}}.$$

Hint. Apply the **Gaussian deviation theorem** to T_r for a suitable r. You may use the following form without proof: for compact symmetric S containing 0 and $u \ge 0$, with probability at least $1 - 2e^{-u^2}$,
$$\sup_{h \in S}\Big|\|Ah\|_2 - \sqrt{d}\|h\|_2\Big| \le C_0\big[w(S) + u\,\mathrm{rad}(S)\big], \qquad \mathrm{rad}(S) = \sup_{h\in S}\|h\|_2.$$

6. A large Euclidean section of the ℓ^1-ball

Let G be an $n \times k$ standard Gaussian matrix, where $1 \le k \le n$, and put $\mu = \mathbb{E}|g| = \sqrt{2/\pi}$ for $g \sim N(0,1)$.

1. Prove that there are absolute constants $c_0, c_1 > 0$ such that, if $k \le c_0 n$, then with probability at least $1 - 2e^{-c_1 n}$,
$$\frac{\mu}{2} n\|v\|_2 \le \|Gv\|_1 \le 2\mu n\|v\|_2 \qquad \text{for every } v \in \mathbb{R}^k.$$
2. Deduce that, for $k \le c_0 n$ (reducing the absolute constants if needed), with probability at least $1 - 2e^{-cn}$ the k-dimensional subspace $F = \mathrm{range}(G)$ satisfies
$$c\sqrt{n}\|z\|_2 \le \|z\|_1 \le \sqrt{n}\|z\|_2 \qquad \text{for every } z \in F.$$

Hint. For a fixed unit vector v, use Gaussian concentration for $\|Gv\|_1$, then pass from a sufficiently fine net to the whole sphere. You may use the Gaussian singular-value estimates from the course.

7. Generic chaining on a binary tree

For a finite metric space (T, d), define

$$\gamma_2(T,d) = \inf_{(T_j)} \sup_{t\in T} \sum_{j\ge 0} 2^{j/2} d(t, T_j), \qquad |T_0| = 1, \quad |T_j| \le 2^{2^j} \ (j \ge 1).$$

You may use **Talagrand's theorem**: for a centered Gaussian process with canonical metric $d(s,t) = (\mathbb{E}|X_s - X_t|^2)^{1/2}$,

$$c\gamma_2(T,d) \le \mathbb{E}\max_{t\in T} X_t \le C\gamma_2(T,d).$$

Let $T = \{0,1\}^m$, $m \ge 1$, be the leaves of a binary tree of depth m. Attach independent standard Gaussian variables g_e to its edges and put $X_t = \sum_{e\in[\varnothing,t]} g_e$.

1. If $\ell(s,t)$ is the length of the common prefix of two leaves, prove $d(s,t)^2 = 2(m - \ell(s,t))$.
2. Construct an admissible sequence by representing every prefix of length $\min\{m, 2^j\}$ at level $j \ge 1$ by one leaf, and taking a single leaf at level 0. Prove directly that $\gamma_2(T,d) \le Cm$.
3. Starting at the root, repeatedly follow the edge with the larger Gaussian label. Use independence of unexplored edges to show that the expected value at the resulting leaf is $m/\sqrt{\pi}$. Deduce $\gamma_2(T,d) \ge cm$ and $cm \le \mathbb{E}\max_t X_t \le Cm$.

Hint. If a, b are independent standard normals, $\max(a,b) = (a + b + |a-b|)/2$. In part 2, bound the nonzero sum by $C\sqrt{m}\sum_{2^j<m} 2^{j/2}$, keeping the level-zero term.

8. Generic chaining for a product of balls

Let $a_1, \dots, a_q > 0$ and integers $d_1, \dots, d_q \ge 1$ be given. In the orthogonal product $\mathbb{R}^{d_1} \times \cdots \times \mathbb{R}^{d_q}$, let

$$T = \prod_{j=1}^{q} a_j B_2^{d_j}$$

with its Euclidean metric. The definition and theorem in Problem 7 extend to compact Euclidean sets and their continuous Gaussian processes.

Prove that

$$w(T) = \sum_{j=1}^{q} a_j \mathbb{E}\|g^{(j)}\|_2, \qquad c\sum_{j=1}^{q} a_j\sqrt{d_j} \le \gamma_2(T) \le C\sum_{j=1}^{q} a_j\sqrt{d_j},$$

where $g^{(j)}$ is standard Gaussian in $\mathbb{R}^{d_j}$.

Fact you may use. For a standard Gaussian vector in $\mathbb{R}^d$, $c\sqrt{d} \le \mathbb{E}\|g\|_2 \le \sqrt{d}$.

Problems 3–5 develop past exam prompts supplied by the instructor. Problems 4 and 5 are related to the noisy-recovery and random-section exercises in Roman Vershynin, *High-Dimensional Probability*; the necessary matrix deviation statements are supplied in full here.