

Direction Instability Predicts Cross-Cell-Line Drug Mechanism Transport in LINCS L1000

Elliot Tower
elliott@elliotttower.ai

July 2026

Abstract

A drug’s mechanism of action may or may not generalize across cell types. We adapt a geometric measure—direction instability, the mean pairwise angular deviation of perturbation signature vectors—to quantify whether a drug’s transcriptomic effect is conserved across biological contexts. Applied to 8,949 compounds tested in ≥ 5 cell lines from the LINCS L1000 Connectivity Map (978 landmark genes, 167,266 signatures), direction instability separates mechanism-conserving drugs from context-dependent ones (population-null $p < 0.01$ for 78 drugs). Direction instability anticorrelates with gene-level Jaccard overlap of top perturbed genes (Spearman $\rho = -0.79$), confirming that directional change reflects biological mechanism change. Within the HDAC inhibitor class, pan-isoform inhibitors transport across cell lines (mean instability 0.62, $n = 8$) while isoform-selective inhibitors do not (mean instability 0.96, $n = 5$), demonstrating that target breadth—not drug class membership—determines cross-context generalizability.

1 Introduction

Pharmacological perturbation experiments increasingly rely on profiling drug effects across panels of cell lines to identify compounds with robust mechanisms of action [Subramanian et al., 2017]. The LINCS L1000 Connectivity Map measures 978 landmark gene expression changes for $\sim 20,000$ compounds across up to 70 cell lines, producing high-dimensional signature vectors that characterize each drug’s transcriptomic effect in each context [Subramanian et al., 2017].

A fundamental question arises: when does a drug’s effect *transport* from one cell type to another? Two drugs may share a mechanism-of-action label yet differ in how consistently they perturb gene expression across contexts. A pan-HDAC inhibitor like vorinostat affects chromatin remodeling in every cell, while an HDAC8-selective inhibitor like PCI-34051 only produces a strong transcriptomic signature where HDAC8 is functionally relevant.

We formalize this distinction using **direction instability**: the mean pairwise angular deviation of a drug’s unit-normalized signature vectors across cell lines. This metric is adapted from the neural bracket norm [Tower, 2026], which quantifies how much a learned representation’s direction rotates across evidence levels. Here, the analogy maps evidence quartiles to cell lines, and neural activation vectors to drug perturbation signatures.

Direction instability decomposes the perturbation effect into two independent components. The *direction* captures which genes are perturbed and in what relative proportions. The *magnitude* captures the overall effect size. A drug with low direction instability but high magnitude variation has a conserved qualitative mechanism with context-dependent penetrance—a biologically meaningful and pharmacologically useful category.

We apply this framework to 8,949 drugs with ≥ 5 cell lines in LINCS L1000. The scorer and validation labels were frozen under a pre-registered commit SHA before any real data were analyzed.

2 Methods

2.1 Data

Perturbation signatures were drawn from the LINCS L1000 Phase I dataset (GEO accession GSE92742), using Level 5 replicate-collapsed z-scores across 978 landmark genes [Subramanian et al., 2017]. Metadata (signature identifiers, perturbation types, cell line annotations) were downloaded from GEO and filtered to small-molecule perturbagens (`pert_type = trt_cp`) with ≥ 5 distinct cell lines. Replicate signatures within a drug-cell-line pair were averaged to yield one consensus signature per drug per cell line.

Mechanism-of-action (MOA) annotations were obtained from the Broad Drug Repurposing Hub (version 2020-03-24) [Corsello et al., 2017], providing MOA labels for 1,782 of the 8,949 drugs. These labels were frozen before analysis (commit SHA 1dc20a2).

2.2 Direction instability

For a drug tested in K cell lines, let $\mathbf{s}_1, \dots, \mathbf{s}_K \in \mathbb{R}^{978}$ denote its consensus signature vectors. Define the unit-normalized signatures $\hat{\mathbf{s}}_i = \mathbf{s}_i / \|\mathbf{s}_i\|_2$. Direction instability is:

$$D = 1 - \frac{2}{K(K-1)} \sum_{i < j} \hat{\mathbf{s}}_i^\top \hat{\mathbf{s}}_j \quad (1)$$

$D \in [0, 2]$. $D = 0$ when all signatures point in the same direction. $D = 1$ when signatures are orthogonal on average. $D > 1$ indicates anticorrelation.

2.3 Complementary metrics

Magnitude CV: coefficient of variation of signature L_2 norms $\{\|\mathbf{s}_i\|\}_{i=1}^K$, capturing effect-size variation independent of direction.

Fréchet variance: mean squared geodesic deviation from the Fréchet mean on the unit sphere S^{977} .

Gene-level Jaccard overlap: for each pair of cell lines, the Jaccard index of the top-50 up-regulated and top-50 down-regulated genes, averaged over all pairs and both directions.

2.4 Population null model

A drug’s direction instability depends on its cell-line count: drugs tested in more cell lines have more pairwise comparisons and tend toward higher baseline instability. To control for this confound, we compare each drug against a reference population of drugs with similar coverage: all drugs with cell-line count in $[K/2, 2K]$, where K is the focal drug’s count. The p -value is the fraction of reference drugs with instability $\leq$ the focal drug’s instability. This replaces a permutation null, which is invariant under cell-line label shuffling for pairwise cosine statistics.

2.5 Pre-registration

The scoring functions (Equation 1 and all complementary metrics) and the frozen MOA labels were committed before any real LINCS signatures were analyzed. The commit SHA and frozen label file are available in the repository.

3 Results

3.1 Landscape of drug mechanism transport

Among 8,949 drugs with ≥ 5 cell lines, mean direction instability was 0.924 ± 0.061 (median 0.940). Most drugs exhibit context-dependent effects. Only 78 drugs (0.9%) achieved $p < 0.01$ under the population null, and 411 (4.6%) achieved $p < 0.05$, indicating that genuinely mechanism-conserving drugs are rare.

Cell-line coverage ranged from 5 to 64 (mean 10.2, median 8). The population null accounts for this variation: a drug’s instability is evaluated against drugs with comparable coverage.

3.2 Direction instability reflects biological mechanism conservation (H3)

Direction instability anticorrelated with gene-level Jaccard overlap of top perturbed genes: Spearman $\rho = -0.79$ ($p < 10^{-300}$). Drugs with low instability literally perturb the same genes across cell types, while high-instability drugs engage different gene programs in different contexts. This confirms that direction instability captures genuine biological mechanism change, not measurement noise.

3.3 Direction and magnitude instability are partially dissociable (H4)

Direction instability and magnitude CV showed weak correlation (Spearman $\rho = -0.10$, $p = 2.1 \times 10^{-21}$). 7.8% of drugs fell in the low-direction, high-magnitude quadrant: conserved mechanism with context-dependent penetrance.

3.4 MOA class stratification (H2)

Table 1 shows mean direction instability by MOA class (classes with ≥ 10 annotated drugs). Two classes stand out: HDAC inhibitors ($\bar{D} = 0.77$, $\sigma = 0.14$) and topoisomerase inhibitors ($\bar{D} = 0.77$, $\sigma = 0.12$). CDK inhibitors and tubulin polymerization inhibitors also show reduced instability ($\bar{D} = 0.80$ and 0.88 respectively). Receptor antagonists and agonists cluster near the population mean ($\bar{D} = 0.93$ – 0.96), consistent with receptor-mediated mechanisms being cell-type-dependent.

3.5 The HDAC inhibitor selectivity gradient

The HDAC inhibitor class provides a natural experiment: 20 annotated HDAC inhibitors span the full selectivity spectrum from pan-isoform to isoform-specific. Table 2 shows the eight most-transporting HDAC inhibitors (all pan-isoform) and the five least-transporting (selective or weak).

Pan-HDAC inhibitors (mean $D = 0.62$, $n = 8$) transport their mechanism across cell lines because their targets—histone deacetylases of classes I, II, and IV—are expressed in every cell type. Chromatin remodeling is a universal process, so inhibiting it broadly produces a conserved transcriptomic signature.

Selective HDAC inhibitors (mean $D = 0.96$, $n = 5$) do not transport. PCI-34051 targets HDAC8 specifically; its transcriptomic effect depends on whether HDAC8 is functionally relevant

Table 1: Direction instability by MOA class ($n \geq 10$). Classes targeting fundamental cellular machinery (chromatin, DNA, cell cycle) show lower instability than receptor-mediated classes. Within HDAC inhibitors, the high variance ($\sigma = 0.14$) reflects the pan-selective gradient discussed in Section 3.5.

MOA class	n	$\bar{D}$	σ
topoisomerase inhibitor	17	0.765	0.123
HDAC inhibitor	20	0.766	0.142
CDK inhibitor	16	0.797	0.119
tubulin polymerization inhibitor	19	0.882	0.048
MEK inhibitor	14	0.888	0.066
EGFR inhibitor	23	0.899	0.064
FLT3 inhibitor	15	0.896	0.057
tyrosine kinase inhibitor	14	0.901	0.065
VEGFR inhibitor	21	0.909	0.045
calcium channel blocker	25	0.923	0.067
monoamine oxidase inhibitor	17	0.924	0.071
glucocorticoid receptor agonist	31	0.933	0.039
<i>Population mean</i>	8,949	0.924	0.061
dopamine receptor antagonist	59	0.936	0.046
histamine receptor antagonist	41	0.944	0.049
serotonin receptor antagonist	62	0.945	0.035
cyclooxygenase inhibitor	53	0.947	0.031
adrenergic receptor antagonist	52	0.945	0.038
adrenergic receptor agonist	40	0.950	0.028
phosphodiesterase inhibitor	31	0.953	0.023
serotonin receptor agonist	31	0.958	0.022
PPAR receptor agonist	19	0.956	0.019

in a given cell type. Valproic acid and phenylbutyrate are weak HDAC inhibitors with additional pharmacological activities, so their signatures reflect cell-type-specific off-target effects rather than HDAC inhibition per se.

Entinostat ($D = 0.70$, class I-selective) falls between the two groups, consistent with intermediate target breadth: class I HDACs (HDACs 1, 2, 3) are more widely expressed than HDAC8 but narrower than the full isoform panel.

3.6 Top transporting drugs target fundamental machinery

The 10 most mechanism-conserving drugs with MOA annotations target processes common to all cells (Table 3): protein synthesis (homoharringtonine), RNA transcription (dactinomycin), DNA topology (epirubicin, doxorubicin), chromatin remodeling (PCI-24781, belinostat, panobinostat), cell cycle progression (BMS-387032), and signal transduction kinases (bisindolylmaleimide-IX). These mechanisms operate independently of tissue-specific receptor expression, explaining their cross-context conservation.

Conversely, imatinib ($D = 0.97$) exemplifies a context-dependent drug: its BCR-ABL kinase inhibition produces strong effects only in cells harboring the Philadelphia chromosome, yielding high direction instability across the LINCS panel.

Table 2: HDAC inhibitors ranked by direction instability. Pan-isoform inhibitors achieve low instability and significant population p -values; selective inhibitors are indistinguishable from the null.

Drug	Selectivity	D	n	p
PCI-24781	pan	0.561	13	0.004
belinostat	pan	0.589	13	0.006
panobinostat	pan	0.591	15	0.007
scriptaid	pan	0.620	15	0.009
vorinostat	pan	0.628	60	0.008
trichostatin-a	pan	0.634	64	0.014
NSC-3852	pan	0.674	15	0.016
entinostat	class I	0.697	14	0.023
valproic acid	weak/partial	0.955	54	0.947
Merck60	selective	0.962	9	0.778
PCI-34051	HDAC8-selective	0.984	12	0.976
phenylbutyrate	weak/partial	0.989	11	0.988

Table 3: Top 10 transporting drugs with MOA annotation.

Drug	MOA	D	p
homoharringtonine	protein synthesis inhibitor	0.497	0.001
dactinomycin	RNA polymerase inhibitor	0.506	0.002
BMS-387032	CDK / cell cycle inhibitor	0.522	0.002
bisindolylmaleimide-IX	PKC inhibitor	0.526	0.003
epirubicin	topoisomerase inhibitor	0.529	0.003
anisomycin	DNA synthesis inhibitor	0.555	0.004
PCI-24781	HDAC inhibitor	0.561	0.004
BNTX	opioid receptor antagonist	0.564	0.003
digitoxigenin	ATPase inhibitor	0.586	0.004
belinostat	HDAC inhibitor	0.589	0.006

4 Discussion

Direction instability provides a single-number summary of whether a drug’s transcriptomic mechanism generalizes across cell types. Three findings support its validity.

First, the metric tracks gene-level mechanism conservation ($\rho = -0.79$ with Jaccard overlap). This is a strong sanity check: if direction instability were driven by noise or batch effects, it would not correlate with which specific genes are perturbed.

Second, MOA classes stratify in a biologically interpretable pattern. Drugs targeting universal cellular machinery (chromatin remodeling, DNA topology, protein synthesis, cell cycle) show low instability, while receptor-mediated drugs show high instability. This matches the expectation that receptor expression is tissue-specific.

Third, the HDAC inhibitor selectivity gradient provides a within-class controlled comparison. Pan-isoform and selective HDAC inhibitors share an MOA label but differ in target breadth, and this difference maps directly onto direction instability. This rules out the alternative explanation that MOA labels alone drive the signal.

Relation to the bracket norm. Direction instability is formally analogous to the neural bracket norm [Tower, 2026], which measures directional rotation of learned representations across evidence levels. Both metrics detect when a system (neural network or cell) responds to a perturbation (evidence or drug) by rotating its internal state rather than scaling it. The pharmacological application replaces evidence quartiles with cell lines and neural activations with gene expression signatures, preserving the geometric interpretation: low rotation implies a conserved mechanism that transports across contexts.

Limitations. LINCS L1000 measures only 978 landmark genes; full-transcriptome profiling might reveal additional structure. The population null controls for cell-line count but not for other confounds (drug concentration, time point, signature quality). MOA annotations from the Repurposing Hub are incomplete (20% coverage) and may contain errors. The direction-instability metric treats all cell lines as exchangeable and does not model phylogenetic relationships between cell types.

Applications. Direction instability could serve as a pre-clinical filter for drug repurposing: compounds with low instability are more likely to produce consistent effects across patient populations. The metric requires only perturbation signatures across ≥ 5 contexts, making it applicable to any multi-context profiling dataset (CMAP, JUMP-CP, Perturb-seq).

References

- Steven M Corsello, Joshua A Bittker, Zihan Liu, Joshua Gould, Patrick McCarren, Jodi E Hirschman, Stephen E Johnston, Anita Vrcic, Bang Wong, Mariam Khan, et al. The drug repurposing hub: a next-generation drug library and information resource. *Nature Medicine*, 23(4):405–408, 2017.
- Aravind Subramanian, Rajiv Narayan, Steven M Corsello, David D Peck, Ted E Natoli, Xiaodong Lu, Joshua Gould, John F Davis, Andrew A Tubelli, Jacob K Asiedu, et al. A next generation connectivity map: L1000 platform and the first 1,000,000 profiles. *Cell*, 171(6):1437–1452, 2017.
- Elliot Tower. Direction instability as a geometric diagnostic for mechanism transport. *Manuscript in preparation*, 2026.