

## 1 Binary operations (09/20)

This course is about the theory of groups. Groups are sets equipped with extra structure, a binary operation, which satisfies certain conditions, namely associativity, the existence of an identity, and the existence of inverses.

**Definition 1.1** (Products). Let  $S$  be a set. The **product** of  $S$  with itself, written  $S \times S$ , is the set of ordered pairs  $(a, b)$  where  $a$  and  $b$  are in  $S$ . Elements of  $S \times S$  are often called **ordered tuples**.

**Definition 1.2** (Binary operations). A binary operation on a set  $S$  is a function  $m: S \times S \rightarrow S$ . For  $a, b \in S$  we will often write  $a \cdot b$  or even  $ab$  for  $m(a, b)$ . This is multiplicative notation. We will also have occasion to use additive notation and write  $a + b$  for  $m(a, b)$ .

**Definition 1.3** (Properties of binary operations). Let  $m: S \times S \rightarrow S$  be a binary operation on a set  $S$ , written  $m(a, b) = a \cdot b$ .

- (a) We say  $m$  is **commutative** if  $a \cdot b = b \cdot a$  for all  $a, b \in S$ .
- (b) We say  $m$  is **associative** if  $a \cdot (b \cdot c) = (a \cdot b) \cdot c$  for all  $a, b, c \in S$ .
- (c) We say  $m$  is **unitary** if there exists a (two-sided) **identity element**, which is an element  $e \in S$  such that  $e \cdot a = a = a \cdot e$  for all  $a \in S$ . If  $m$  is unitary, then the identity element  $e$  is unique; see Lemma 1.5.
- (d) We say  $m$  has the **Latin square property** if for each  $a, b \in S$  there exist unique  $x, y \in S$  such that  $a \cdot x = b$  and  $y \cdot a = b$ .
- (e) We say that a unitary binary operation  $m$  has **inverses** if for each  $a \in S$  there exists  $b \in S$  such that  $a \cdot b = b \cdot a = e$  for an identity element  $e$  (which is unique by Lemma 1.5). Such an element  $b$  is called a (two-sided) **inverse** of  $a$  and is written as  $a^{-1}$ . Inverses are unique if  $m$  is additionally associative by Exercise 1.3.

**Example 1.4.** Binary operations can be very simple, too simple to be of interest. For example, let  $\mathbf{Z}$  be the **set of integers**. Define  $m: \mathbf{Z} \times \mathbf{Z} \rightarrow \mathbf{Z}$  by setting  $m(a, b) = 17$  for all integers  $a, b$ . In the notation above, we let  $a \cdot b = 17$  for all  $a, b \in \mathbf{Z}$ . This is a binary operation, which is commutative and associative, but not terribly useful.

**Lemma 1.5** (Identities are unique). *Suppose that  $m$  is a unitary binary operation on a set  $S$ . If  $e$  and  $e'$  are identity elements, then  $e = e'$ .*

*Proof.* We have  $e = e \cdot e' = e'$ , where the first equality uses the identity property of  $e'$  and the second equality uses the identity property of  $e$ .  $\square$

**Notation 1.6.** We will sometimes write 1 for the identities with respect to binary operations when writing multiplicatively; and we will sometimes write 0 for the identities with respect to binary operation written additively. Similarly, we might write  $-a$  for the inverse of  $a$  when writing additively.

**Remark 1.7** (Commutative diagrams). Associativity can be expressed as follows. Let  $m \times \text{id}_S: S \times S \times S \rightarrow S \times S$  be defined by  $(m \times \text{id}_S)(a, b, c) = (m(a, b), c)$  and let  $\text{id}_S \times m: S \times S \times S \rightarrow S \times S$  be defined by  $(\text{id}_S \times m)(a, b, c) = (a, m(b, c))$ . The functions  $m \circ (m \times \text{id}_S)$  and  $m \circ (\text{id}_S \times m)$  define two functions on the set  $S \times S \times S$  of ordered triples of elements of  $S$ . (These might be called ternary operations.) The binary

operation  $m$  is associative if these two functions are equal. In contemporary mathematics, it is common to express this via a **commutative diagram**. In this case, the diagram would be as follows:

$$\begin{array}{ccc}
 S \times S \times S & \xrightarrow{m \times \text{id}_S} & S \times S \\
 \text{id}_S \times m \downarrow & & \downarrow m \\
 S \times S & \xrightarrow{m} & S.
 \end{array}$$

Saying that the diagram is commutative amounts to asserting that the two ways of traversing the diagram from the upper left to the bottom right by composing functions result in the same function  $S \times S \times S \rightarrow S$ . Commutative diagrams need not be square. For example, let  $t: S \times S \rightarrow S \times S$  be defined by  $t(a, b) = (b, a)$ . Commutativity is the statement that the following triangular diagram commutes:

$$\begin{array}{ccc}
 S \times S & \xrightarrow{t} & S \times S \\
 & \searrow m & \swarrow m \\
 & S, &
 \end{array}$$

which means that  $m \circ t = m$ .

**Remark 1.8.** If  $m$  is a binary operation on  $S$  satisfying the Latin square property, then the multiplication table of  $m$  is a Latin square: each element of  $S$  appears exactly once in each row and column. In the context of binary operations, these are called Cayley tables. For example, the Latin square of Figure 1 can be viewed

+	0	1	2
0	0	1	2
1	1	2	0
2	2	0	1

Table 1: A Cayley table, which in this case represents a Latin square (the bottom right  $3 \times 3$  part of the table).

as the “addition table” of a binary operation  $m$  on the set  $S = \{0, 1, 2\}$ .

**Example 1.9.** Let  $\mathbf{N} = \{0, 1, 2, 3, 4, 5, \dots\}$  be the set of **natural numbers**, which we take to be the non-negative integers. On  $\mathbf{N}$  we have the binary operation of addition, given by  $m(a, b) = a + b$ . This binary operation is commutative, associative, and unital; it has neither the Latin square property nor inverses.

**Example 1.10.** Let  $\mathbf{Z}$  be the set of integers. On  $\mathbf{Z}$  the binary operation of addition has all of the properties (a)-(e) of Definition 1.3. We can also multiply integers: the binary operation of multiplication satisfies properties (a)-(c) but not (d) or (e).

**Example 1.11.** We can construct Cayley tables for the outcomes of simple games. For example, consider the two-player game of rock, paper, scissors. The plays are denoted by  $r$ ,  $p$ , and  $s$ . The outcomes of possible plays are listed in Figure 2. For example, if  $p \cdot s = s = s \cdot p$  represents the fact that scissor beats paper, no matter who plays it. Now, consider

$$(r \cdot p) \cdot s = p \cdot s = s \quad \text{and} \quad r \cdot (p \cdot s) = r \cdot s = r,$$

which shows that this commutative binary operation is not associative.

$\cdot$	$r$	$p$	$s$
$r$	$r$	$p$	$r$
$p$	$p$	$p$	$s$
$s$	$r$	$s$	$s$

Table 2: A Cayley table for rock, paper, scissors. The associated binary operation is commutative, but not associative.

## 1.1 Exercises

**Exercise 1.1.** If  $S$  and  $I$  are sets, let  $S^I$  be the set of functions  $f: I \rightarrow S$ . Let  $I = \{0, 1\}$ . Prove that for any set  $S$  there is a bijection  $p: S^I \rightarrow S \times S$ .

**Exercise 1.2.** Let  $S = \{1, \dots, n\}$  for some positive integer  $n$ . Compute the number of binary operations on  $S$ .

**Exercise 1.3.** Show that if  $m$  is a unital, associative binary operation on a set  $S$ , then inverses are unique when they exist: if  $a \in S$  and  $x, y \in S$  are inverses of  $a$ , then  $x = y$ .

**Exercise 1.4** (The Eckmann–Hilton argument). Let  $S$  be a set with two binary operations  $\bullet$  and  $\circ$  satisfying the following two axioms:

- (i)  $\bullet$  and  $\circ$  each has a two-sided identity element,  $\mathbf{1}_\bullet$  and  $\mathbf{1}_\circ$ , respectively;
- (ii) for each  $a, b, c, d \in S$ , there is the identity  $(a \circ b) \bullet (c \circ d) = (a \bullet c) \circ (b \bullet d)$ .

Prove that (a)  $\mathbf{1}_\bullet = \mathbf{1}_\circ$ , (b)  $\bullet = \circ$ , (c)  $\bullet$  is associative, and (d)  $\bullet$  is commutative.

**Exercise 1.5.** Find a binary operation which is not commutative and not associative.

## 2 Groups (09/22)

Algebraic structures are sets equipped with additional structures, often binary operations, which satisfy certain properties and are viewed as being part of the data of the algebraic structure.

**Definition 2.1** (Magma). A **magma**  $M$  is a pair  $(S, \cdot)$  where  $S$  is a set and  $\cdot$  is a binary operation on  $S$ . The binary operation could also be written as  $+$  or  $\bullet$  or  $\star$ , etc.

**Notation 2.2.** It is very convenient to write  $M$  for the magma *and* the underlying set. So, a magma  $M$  will be a set  $M$  equipped with a binary operation on  $M$ . This is an abuse of notation, but is harmless and will make everything a bit prettier.

**Remark 2.3.** While a set has varying binary operations, a magma has a single binary operation which is singled out and viewed as fixed.

**Definition 2.4** (Types of magmas). In general, one can say that a magma is commutative, associative, unital, and so forth if its binary operation has that property. In many cases, magmas possessing these properties have special names.

- (a) A **semigroup** is an associative magma.
- (b) A **monoid** is a unital semigroup (a unital associative magma).
- (c) A **group** is a monoid which has inverses (a unital associative magma with inverses).
- (d) An **abelian group** is a group whose underlying magma is commutative.<sup>1</sup>
- (e) A **quasigroup** is a magma with the Latin square property.
- (f) A **loop** is a unital quasigroup.

This course will focus on the theory of groups, although monoids are also sometimes useful.

**Definition 2.5.** A **finite group** is a group whose underlying set is finite.

**Example 2.6.** The set  $\mathbf{N} = \{0, 1, 2, \dots\}$  of natural numbers is a commutative monoid under addition. It is not a group.

**Example 2.7.** The set  $\mathbf{Z} = \{0, \pm 1, \pm 2, \dots\}$  of integers under addition is an abelian group. Unless otherwise specified, when we speak of  $\mathbf{Z}$  we will always mean this particular group.

**Warning 2.8.** There is another natural binary operation on  $\mathbf{Z}$ : multiplication. Under this operation,  $(\mathbf{Z}, \cdot)$  is a commutative monoid, but it is not a group. Taken together, the triple  $(\mathbf{Z}, +, \cdot)$  forms a **ring**: a set with an abelian group structure under  $+$ , a monoid structure under  $\cdot$ , and where  $+$  and  $\cdot$  interact in a prescribed way via the **distributivity laws**:  $(a + b) \cdot c = a \cdot c + b \cdot c$  and  $a \cdot (b + c) = a \cdot b + a \cdot c$ . This particular ring is commutative because the multiplicative monoid is. These algebraic structures are the subject of the second quarter of this sequence.

**Example 2.9.** The sets  $\mathbf{Q}$ ,  $\mathbf{R}$ ,  $\mathbf{C}$ , and  $\mathbf{R}^n$  under (vector) addition are abelian groups.

**Example 2.10.** If  $k$  is a field and  $V$  is a  $k$ -vector space, then addition makes  $V$  into an abelian group.

**Example 2.11.** If  $G = \{e\}$  is a set with a single element,  $e$ , then the unique binary operation on  $G$  (specified by  $e \cdot e = e$ ) makes  $G$  into a group (with identity element  $e$ ).

<sup>1</sup>One could call these commutative groups, but for historical reasons, abelian groups are used instead.

**Example 2.12.** The empty set  $\emptyset$  also admits a unique binary operation  $\emptyset \times \emptyset \rightarrow \emptyset$ . It is commutative, associative, and has the Latin square property, but is not unital as unitality asserts the existence of an element. So, it is a semigroup and a quasigroup, but it is not a group.

Now, we introduce two of the most important examples of groups: addition modulo  $N$  and symmetric groups.

**Lemma 2.13.** Fix a positive integer  $N \geq 1$ . Let  $\mathbf{Z}/N$  be the set  $\{0, 1, \dots, N-1\}$ . The binary operation on  $\mathbf{Z}/N$  defined by letting  $a +_N b = r$  where  $r$  is the unique integer in  $\{0, \dots, N-1\}$  such that  $a + b \equiv r \pmod{N}$  makes  $\mathbf{Z}/N$  into an abelian group.

*Proof.* The existence and uniqueness of  $c$  follows from the fact that for  $c \in \mathbf{Z}$  there are unique integers  $q$  and  $r \in \{0, \dots, N-1\}$  such that  $c = qN + r$  (this is often called **Euclidean division**). Applying this to  $c = a + b$  (where the sum is computed in  $\mathbf{Z}$ ) produces  $q$  and  $r$  such that  $a + b = qN + r$ . We define  $a +_N b = r$ . This operation is commutative since  $a + b = b + a = qN + r$ , so  $a +_N b = b +_N a$  and unital since  $a + 0 = 0 + a = 0 \cdot N + a = a$  for  $a \in \{0, \dots, N-1\}$ , so  $a +_N 0 = 0 +_N a = a$ . The inverse of  $a$  is computed by finding  $r \in \{0, \dots, N-1\}$  such that  $-a = qN + r$ . Then,  $0 = a + r = a + qN + r$  is divisible by  $N$  so that  $a + r = N$  and hence  $a + r = (q+1)N + 0$ , so  $a +_N r = 0$ . Thus,  $+_N$  has inverses. For associativity, suppose that  $a + b = q_0N + r_0$  and  $b + c = q_1N + r_1$ , where  $r_0, r_1 \in \{0, \dots, N-1\}$ . Then, assume that  $r_0 + c = q_2N + r_2$  and  $a + r_1 = q_3N + r_3$  for  $r_2, r_3 \in \{0, \dots, N-1\}$ . Then, by associativity of addition on  $\mathbf{Z}$ ,

$$(q_1 + q_3)N + r_3 = a + q_1N + r_1 = a + b + c = q_0N + r_0 + c = (q_0 + q_1)N + r_2.$$

By uniqueness of the remainder, we must have  $r_3 = r_2$ , so that  $a +_N (b +_N c) = (a +_N b) +_N c$ , which proves associativity and finally that  $\mathbf{Z}/N$  is an abelian group.  $\square$

**Notation 2.14.** We will typically write  $a + b \equiv c \pmod{N}$  instead of  $a +_N b = c$  when working in  $\mathbf{Z}/N$ .

**Example 2.15.** The Cayley table of  $\mathbf{Z}/3$  was already introduced in Remark 1.8. We reproduce it here for convenience.

+	0	1	2
0	0	1	2
1	1	2	0
2	2	0	1

Table 1: The Cayley table of  $\mathbf{Z}/3$ .

## 2.1 Exercises

**Exercise 2.1.** An associative loop is a group. Show that there exist non-associative loops.

**Exercise 2.2.** Let  $G$  be a group and fix  $a \in G$ . Prove that  $(a^{-1})^{-1} = a$ .

**Exercise 2.3.** Let  $G$  be a group and fix  $a, b \in G$ . Prove that  $(a \cdot b)^{-1} = b^{-1} \cdot a^{-1}$ .

**Exercise 2.4.** Let  $G$  be a group with identity element  $e$  and fix  $a \in G$  and  $n \in \mathbf{Z}$ . Set  $a^0 = e$ . For  $n > 0$ , define  $a^n$  inductively by  $a^n = a \cdot a^{n-1}$ . For  $n < 0$ , define  $a^n = (a^{-n})^{-1}$ . One has  $a^m \cdot a^n = a^{m+n}$  and  $(a^m)^n = a^{mn}$  for  $m, n \in \mathbf{Z}$ . Prove that if  $G$  is abelian, then  $(a \cdot b)^n = a^n \cdot b^n$  for all  $a, b \in G$ .

**Exercise 2.5.** Let  $G$  be a finite group with identity element  $e$ . Show that there exists an integer  $n > 0$  such that  $a^n = e$  for all  $a \in G$ .

### 3 Symmetric groups (09/25)

**Lemma 3.1.** *Let  $X$  be a set. Let  $S_X$  be the set of bijections  $f: X \rightarrow X$ . On  $S_X$  we define a binary operation via  $f \circ g$ , the composition of  $f$  and  $g$ . This makes  $S_X$  into a group.*

*Proof.* Let  $\text{id}_X: X \rightarrow X$  be the function  $\text{id}_X(x) = x$  for all  $x \in X$ . This is an identity element for  $S_X$ . Indeed, if  $f: X \rightarrow X$  is another function, then  $(f \circ \text{id}_X)(x) = f(\text{id}_X(x)) = f(x) = \text{id}_X(f(x)) = (\text{id}_X \circ f)(x)$  for all  $x \in X$ , so  $f \circ \text{id}_X = \text{id}_X \circ f = f$ .<sup>1</sup> Associativity follows from the fact that  $(f \circ (g \circ h))(x) = f((g \circ h)(x)) = f(g(h(x))) = (f \circ g)(h(x)) = ((f \circ g) \circ h)(x)$ . Finally, the existence of inverses follows because each  $f \in S_X$  is a bijection; the inverse of  $f$  is the inverse function  $f^{-1}$ .  $\square$

**Definition 3.2.** The group  $S_X$  is called the **group of permutations of  $X$** . When  $X = \{1, \dots, n\}$ , we write  $S_n$  for  $S_X$ . This is called the **permutation group on  $n$  symbols** or the **symmetric group of degree  $n$** . We write  $e$  for the identity element of  $S_n$ .

**Lemma 3.3.** *The symmetric group  $S_n$  on degree  $n$  has  $n! = n(n-1)(n-2) \cdots 1$  elements for  $n \geq 1$ .<sup>2</sup>*

*Proof.* We prove the result by induction. Let  $s_n$  be the number of bijections from a set with  $n$  elements to another set with  $n$  elements. We want to show  $s_n = n!$ . When  $n = 1$ , this is true because there is exactly 1 function from a set with 1 element to another set with 1 element. Now, suppose the result is true for  $1, \dots, n-1$ . In particular,  $s_{n-1} = (n-1)!$ . To specify a bijection  $f: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ , we must choose  $f(1)$ . Let  $Y = \{1, \dots, n\} - \{f(1)\}$ . Then, the rest of the values of  $f$  are determined by a bijective function  $f': \{2, \dots, n\} \rightarrow Y$ . There are  $n$  choices of  $f(1)$  and for each such choice  $s_{n-1} = (n-1)!$  for  $f'$ . Thus, there are  $n \cdot (n-1)! = n!$  bijections  $f$ , so  $s_n = n!$ , as desired.  $\square$

**Definition 3.4.** Fix  $n \geq 1$  and consider the symmetric group  $S_n$  of degree  $n$ . A **cycle** of order  $k$  is an ordered string  $(a_1 a_2 \cdots a_k)$  where  $a_1, \dots, a_k \in \{1, \dots, n\}$  are distinct. We view a cycle as a bijection  $\sigma = (a_1 \cdots a_k): \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ , and hence as an element of  $S_n$ , by letting

$$\sigma(x) = \begin{cases} a_{k+1} & \text{if } x = a_1, \dots, a_{k-1}, \\ a_1 & \text{if } x = a_k, \text{ and} \\ x & \text{otherwise.} \end{cases}$$

In words,  $\sigma = (a_1 \cdots a_k)$  is the function which takes  $a_1$  to  $a_2$ ,  $a_2$  to  $a_3$  and so on, all the way to  $a_k$  to  $a_1$ . It does not change other elements.

**Example 3.5.** If  $i \in \{1, \dots, n\}$ , then the cycle  $(i)$  of length 1 is equal to the identity element of  $S_n$ .

**Example 3.6.** Recall that if  $G$  is a group and  $a \in G$ , then the **order of  $a$** , if it exists, is the least integer  $k \geq 1$  such that  $a^k = e$ . Write  $|a| = k$  for the order of  $a$ . (Written additively, this would be the least  $n \geq 1$  such that  $na = 0$ .) If  $f = (a_1 \cdots a_k)$  is a cycle, then its order is  $k$ .

**Definition 3.7.** A **transposition** is a cycle  $(ab)$  of length 2. If  $f = (ab)$ , then  $f^2 = e$ , so  $f^{-1} = f$ .

**Proposition 3.8.** *If  $X$  is a set with at least 3 elements, then  $S_X$  is not abelian. In particular, if  $n \geq 3$  be an integer, then  $S_n$  is not abelian.*

<sup>1</sup>We use throughout that two functions  $f$  and  $g$  from  $X$  to  $Y$  are equal if and only if  $f(x) = g(x)$  for all  $x \in X$ .

<sup>2</sup>It also makes sense to write  $S_0$  for  $S_\emptyset$ ; this group has 1 element.

*Proof.* We can assume that  $X$  contains the set  $\{1, 2, 3\}$ . We compute the compositions

$$(12) \circ (23) = (123) \quad \text{and} \quad (23) \circ (12) = (132).$$

These cycles represent different functions on  $\{1, \dots, n\}$ , so  $(12) \circ (23) \neq (23) \circ (12)$ . (Here, as in Definition 3.4, the cycles given act as the identity away from  $\{1, 2, 3\}$ .)  $\square$

**Remark 3.9.** Note that as an element of  $S_n$  there is no difference between  $(a_1 a_2 \cdots a_n)$  and  $(a_2 a_3 \cdots a_n a_1)$ . But, as in the previous proof, if two cycles  $(a_1 \cdots a_k)$  and  $(b_1 \cdots b_m)$  start with the same element  $a_1 = b_1$ , then they are the same if and only if  $m = k$  and  $b_i = a_i$  for  $1 \leq i \leq k$ .

**Lemma 3.10** (Disjoint cycles commute). *Suppose that  $f = (a_1 \cdots a_k)$  and  $g = (b_1 \cdots b_m)$  are disjoint cycles, meaning that  $a_i \neq b_j$  for all  $1 \leq i \leq k$  and  $1 \leq j \leq m$ . Then,  $f \circ g = g \circ f$ .*

*Proof.* Fix  $x \in \{1, \dots, n\}$ . If  $x$  is not in  $\{a_1, \dots, a_k\}$ , then  $f(x) = x$  and  $g(x)$  is also not in  $\{a_1, \dots, a_k\}$  so that  $(f \circ g)(x) = f(g(x)) = g(x) = g(f(x)) = (g \circ f)(x)$ . The same holds if  $x$  is not in  $\{b_1, \dots, b_m\}$ . But, the union of the complements of  $\{a_1, \dots, a_k\}$  and  $\{b_1, \dots, b_m\}$  is all of  $\{1, \dots, n\}$ . So,  $f \circ g$  and  $g \circ f$  are equal on all of  $\{1, \dots, n\}$  and hence are equal.  $\square$

**Notation 3.11.** Since disjoint cycles commute, if  $(a_1 \cdots a_k)$  and  $(b_1 \cdots b_m)$  are disjoint cycles, we write  $(a_1 \cdots a_k)(b_1 \cdots b_m)$  for their composition, in any order. Thus, for example,  $(12)(34) = (12) \circ (34) = (34) \circ (12)$ . We also make this convention for compositions of multiple pairwise disjoint cycles.

### 3.1 Exercises

**Exercise 3.1.** Let  $f = (a_1 \cdots a_k)$  be a cycle of length  $k$  in  $S_n$ . Write the inverse of  $f$  as a cycle.

**Exercise 3.2.** Let  $f = (a_1 \cdots a_k)$  be a cycle of length  $k$  in  $S_n$ . Prove that  $f$  has order  $k$ .

**Exercise 3.3.** Let  $f = (a_1 \cdots a_k)$  be a cycle of length  $k$  in  $S_n$ . Fix  $s \geq 1$ . Find (and prove) necessary and sufficient conditions for  $f^s$  to be a cycle. Hint: first consider the case of  $s = 2$ .

**Exercise 3.4.** Let  $\mathbf{Z}/N = \{0, \dots, N-1\}$ . Equip  $\mathbf{Z}/N$  with the binary operation given by multiplication modulo  $N$ , so that if  $a, b \in \mathbf{Z}/N$ , then  $a \cdot_N b = r$  where  $ab = qN + r$  where  $r \in \{0, \dots, N-1\}$ . We write  $ab \equiv r \pmod{N}$ .

(a) Show that this binary operation makes  $\mathbf{Z}/N$  into a commutative monoid with identity element 1.

Let  $(\mathbf{Z}/N)^\times \subseteq \mathbf{Z}/N$  be the subset of elements  $a \in \mathbf{Z}/N$  such that there exists  $b \in \mathbf{Z}/N$  with  $ab \equiv ba \equiv 1 \pmod{N}$ .

(b) Show that  $(\mathbf{Z}/N)^\times$  is an abelian group.

(c) Show that  $(\mathbf{Z}/N)^\times$  consists of the elements of  $\mathbf{Z}/N$  which are relatively prime to  $N$ .

## 4 Cycle decomposition in cyclic groups (09/27)

**Theorem 4.1** (Cycle decomposition). *Let  $f \in S_n$  be an element of  $S_n$ . Then, for some  $1 \leq r \leq n$  there are  $r$  pairwise disjoint cycles  $(a_{11} \cdots a_{1,k_1}), (a_{21} \cdots a_{2,k_2}), \dots, (a_{r1} \cdots a_{r,k_r})$  such that*

$$f = (a_{11} \cdots a_{1,k_1}) \cdots (a_{r1} \cdots a_{r,k_r}).$$

*Proof.* As  $\{1, \dots, n\}$  is finite, there is some smallest  $k \geq 1$  for which  $f^{(k)}(1) = 1$ . Then,  $(1 f(1) f(f(1)) \cdots f^{(k-1)}(1))$  is a cycle of length  $k$ . Let this be  $(a_{11} \cdots a_{1,k_1})$ . Let  $a_{21}$  be the first element in  $\{1, \dots, n\}$  not in the cycle  $(a_{11} \cdots a_{1,k_1})$  and consider the cycle generated by  $a_{21}$ , say  $(a_{21} \cdots a_{2,k_2})$ . This is a disjoint cycle. Continue on in this way until every element of  $\{1, \dots, n\}$  appears in a cycle.  $\square$

**Remark 4.2.** As cycles of length 1 all correspond to the identity element of  $S_n$  it is standard to omit them from the final cycle decomposition of  $f$ . The cycle decomposition of  $f$  is unique up to cyclically rotating the terms in the cycles (Remark 3.9) and reordering the cycles themselves (Lemma 3.10).

**Example 4.3.** If  $f = (a_{11} \cdots a_{1,k_1}) \cdots (a_{r1} \cdots a_{r,k_r})$  is a decomposition of  $f$  into disjoint cycles, then the order of  $f$  is the least common multiple of  $k_1, \dots, k_r$ . For example, if  $f = (1\ 12\ 8\ 10\ 4)(2\ 13)(5\ 11\ 7)(6\ 9)$ , then  $|f| = 30$ .

Recall the following definition from last time.

**Definition 4.4.** A **transposition** is a cycle  $(ab)$  of length 2. If  $f = (ab)$ , then  $f^2 = e$ , so  $f^{-1} = f$ .

**Lemma 4.5.** Every element  $f \in S_n$  can be written as a product of transpositions.

*Proof.* Using cycle decomposition, it is enough to prove the result for cycles. Thus, assume that  $f = (a_1 \cdots a_k)$ . Then,  $f = (a_1 a_2) \circ (a_2 a_3) \circ \cdots \circ (a_{k-1} a_k)$ . Indeed, for  $a_i$  with  $1 \leq i \leq k-1$ , it is unchanged except by  $(a_i a_{i+1})$ , which sends it to  $a_{i+1}$ . For  $a_k$ ,  $(a_{k-1} a_k)$  sends it to  $a_{k-1}$ , then  $(a_{k-2} a_{k-1})$  sends it to  $a_{k-2}$ . This continues until finally  $(a_1 a_2)$  sends the result to  $a_1$ .  $\square$

**Example 4.6.** Write down the cycle decomposition of each element of  $S_3$  and compute the order of each element. See Table 1 for the solution.

$e$	1
$(1\ 2)$	2
$(1\ 3)$	2
$(2\ 3)$	2
$(1\ 2\ 3)$	3
$(1\ 3\ 2)$	3

Table 1: The cycle decompositions and orders of the  $6 = 3!$  elements of  $S_3$ .

**Example 4.7.** If  $f = (a_{11} \cdots a_{1,k_1}) \cdots (a_{r1} \cdots a_{r,k_r})$  is a decomposition of  $f$  into disjoint cycles, then the order of  $f$  is the least common multiple of  $k_1, \dots, k_r$ . For example, if  $f = (1\ 12\ 8\ 10\ 4)(2\ 13)(5\ 11\ 7)(6\ 9)$ , then  $|f| = 30$ .

**Example 4.8** (Dummit–Foote, Exercise 1.3.1). One way to write down permutations is using a kind of matrix notation: the permutation  $f \in S_5$  given by

$$1 \mapsto 3 \quad 2 \mapsto 4 \quad 3 \mapsto 5 \quad 4 \mapsto 2 \quad 5 \mapsto 1$$



can be written efficiently as

$$f = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 4 & 5 & 2 & 1 \end{pmatrix},$$

which is just a lookup table. The cycle decomposition of  $f$  is  $f = (1\ 3\ 5)(2\ 4)$ . If we consider

$$g = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 3 & 2 & 4 & 1 \end{pmatrix},$$

which has cycle decomposition  $g = (1\ 5)(2\ 3)$ , then we can compute the cycle decompositions

$$\begin{aligned} f^2 &= (1\ 5\ 3) \\ fg &= (2\ 5\ 3\ 4) \\ gf &= (1\ 2\ 4\ 3) \\ g^2f &= f = (1\ 3\ 5)(2\ 4). \end{aligned}$$

## 4.1 Exercises

**Exercise 4.1.** Justify Example 4.7. Fix pairwise commuting elements  $f_1, \dots, f_r$  of a group  $G$ , i.e., elements such that  $f_i f_j = f_j f_i$  for all  $1 \leq i, j \leq r$ . Prove that if each  $f_i$  has finite order  $n_i$ , then  $f = f_1 \cdots f_r$  has order the least common multiple of  $f_1, \dots, f_r$ .

**Exercise 4.2.** By Lemma 4.5, every element  $f \in S_n$  can be written as a product of transpositions. Suppose that  $f = g_1 \circ \cdots \circ g_k$  where  $g_1, \dots, g_k$  are transpositions. We say that  $f$  is **even** if  $k$  is even and we say that  $f$  is **odd** if  $k$  is odd. Show that this is well-defined by proving that if  $f = h_1 \circ \cdots \circ h_m$  is another way of writing  $f$  as a product of transpositions, then  $k \equiv m \pmod{2}$ .

**Exercise 4.3.** Let  $f = (a_1 \cdots a_k)$  be a cycle. Show that  $f$  is even if  $k$  is odd and that  $f$  is odd if  $k$  is even.

**Exercise 4.4.** Write down the cycle decomposition of each element of  $S_4$  and compute the order of each element.

**Exercise 4.5** (Dummit–Foote, Exercise 1.3.2). Let

$$f = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 \\ 13 & 2 & 15 & 14 & 10 & 6 & 12 & 3 & 4 & 1 & 7 & 9 & 5 & 11 & 8 \end{pmatrix}$$

and

$$g = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 \\ 14 & 9 & 10 & 2 & 12 & 6 & 5 & 11 & 15 & 3 & 8 & 7 & 4 & 1 & 13 \end{pmatrix}$$

be two elements of  $S_{15}$ . Find cycle decompositions for  $f$ ,  $g$ ,  $f^2$ ,  $f \circ g$ ,  $g \circ f$ , and  $g^2 \circ f$ .

## 5 Group homomorphisms (09/29)

**Definition 5.1** (Magma homomorphisms). Let  $M$  and  $N$  be two magmas. A function  $f: M \rightarrow N$  is a **magma homomorphism** if  $f(ab) = f(a)f(b)$  for all  $a, b \in M$ .

**Remark 5.2.** The magma homomorphisms are the functions between the underlying sets that *respect the algebraic structures* given by the binary operations on  $M$  and  $N$ .

**Definition 5.3.** If  $G$  and  $H$  are groups, a function  $f: G \rightarrow H$  is a **group homomorphism** if it is a homomorphism of the underlying magmas, i.e., if  $f(ab) = f(a)f(b)$  for all  $a, b \in G$ .

**Remark 5.4.** In the same way, one can define semigroup, monoid, quasigroup, and loop homomorphisms.

**Lemma 5.5.** If  $f: G \rightarrow H$  is a group homomorphism, then  $f(e_G) = e_H$  where  $e_G$  is the identity element of  $G$  and  $e_H$  is the identity element of  $H$ .

*Proof.* Since  $H$  is a group,  $f(e_G)$  possesses an inverse, say  $a$  so that  $af(e_G) = e_H$ . We have  $f(e_G) = f(e_G e_G) = f(e_G)f(e_G)$ ; multiplying both sides on the left by  $a$  we obtain  $e_H = af(e_G) = af(e_G)f(e_G) = e_H f(e_G) = f(e_G)$ , as desired.  $\square$

**Lemma 5.6.** If  $f: G \rightarrow H$  is a group homomorphism, then  $f(a)^{-1} = f(a^{-1})$  for all  $a \in G$ .

*Proof.* By uniqueness of inverses in groups, it is enough to show that  $f(a^{-1})$  is an inverse for  $f(a)$ . But,  $f(a^{-1})f(a) = f(a^{-1}a) = f(e_G) = e_H$ , by Lemma 5.5, and similarly  $f(a)f(a^{-1}) = e_H$ .  $\square$

**Example 5.7.** Consider the exponential function  $\exp: \mathbf{R} \rightarrow \mathbf{R}$  given by  $\exp(x) = e^x$ . As  $\exp(x+y) = \exp(x)\exp(y)$ , the map  $\exp$  is a commutative monoid homomorphism  $(\mathbf{R}, +) \rightarrow (\mathbf{R}, \times)$ . If we delete 0, the function  $\exp$  can be viewed as a group homomorphism  $\mathbf{R} \rightarrow \mathbf{R}^\times$ , where  $\mathbf{R}^\times = \mathbf{R} - \{0\}$  is the group of non-zero elements of  $\mathbf{R}$  under multiplication.

**Example 5.8.** We can also consider the function  $f: (\mathbf{R}, +) \rightarrow (\mathbf{R}, \times)$  given by  $f(x) = 0$  for all  $x$ . This is also a commutative monoid homomorphism. However, we do not have  $f(0) = 1$ , so it does not preserve the identity element of  $(\mathbf{R}, +)$ . This shows that the hypothesis that  $G$  and  $H$  be groups in Lemma 5.5 is necessary.

**Definition 5.9.** We say that a group homomorphism  $f: G \rightarrow H$  is injective (one-to-one), surjective (onto), or bijective if the underlying function of sets is injective, surjective, or bijective.

**Lemma 5.10.** A group homomorphism  $f: G \rightarrow H$  is injective if and only if  $f(x) = e$  implies  $x = e$ .

*Proof.* Suppose that  $f(x) = f(y)$  for some  $x, y \in G$ . Then,  $e = f(e) = f(x^{-1})f(x) = f(x^{-1})f(y) = f(x^{-1}y)$ , so  $x^{-1}y = e$ , or  $y = x$ .  $\square$

**Lemma 5.11.** Suppose that  $f: G \rightarrow H$  is a bijective group homomorphism. Let  $f^{-1}: H \rightarrow G$  be the inverse function. Then,  $f^{-1}$  is a group homomorphism (which is again bijective).

*Proof.* Let  $x, y \in H$ . We have to prove that  $f^{-1}(xy) = f^{-1}(x)f^{-1}(y)$ . Write  $x = f(a)$  and  $y = f(b)$ , for unique  $a, b \in G$ , using that  $f$  is a bijection. Then,  $f(ab) = f(a)f(b) = xy$ , so that  $f^{-1}(xy) = ab = f^{-1}(x)f^{-1}(y)$ .  $\square$

**Definition 5.12.** A bijective group homomorphism is called a **isomorphism**. Two groups  $G$  and  $H$  are called **isomorphic** if there exists a group isomorphism  $f: G \rightarrow H$ .

**Example 5.13.** Let  $\mathbf{R}_+^\times$  be the group of positive real numbers under multiplication. The exponential map  $\exp: \mathbf{R} \rightarrow \mathbf{R}_+^\times$  is an isomorphism, so  $\mathbf{R} \cong \mathbf{R}_+^\times$ .

**Remark 5.14.** If  $G$  is a group, then the identity function  $\text{id}_G$  is a group isomorphism. If  $f: G \rightarrow H$  and  $h: H \rightarrow K$  are group isomorphisms, then so is  $h \circ f: G \rightarrow K$ . Using these facts and Lemma 5.11, it follows that the relation  $G \cong H$  if  $G$  and  $H$  are isomorphic is an equivalence relation on the class of groups.

**Example 5.15.** Let  $G$  and  $H$  be groups with 1 element. Then,  $G \cong H$ . In particular,  $S_0 = S_\emptyset$  and  $S_1$  are isomorphic.

**Example 5.16.** There is an isomorphism  $\mathbf{Z}/2 \rightarrow S_2$ , so  $\mathbf{Z}/2 \cong S_2$ .

**Example 5.17.** If  $G$  is a group of order 2 (i.e., the underlying set has exactly 2 elements), then  $G \cong \mathbf{Z}/2$ .

**Example 5.18.** If  $G$  is a group of order 3, then  $G \cong \mathbf{Z}/3$ .

**Definition 5.19** (Cyclic groups). A group  $G$  is **cyclic** if  $G \cong \mathbf{Z}$  or  $G \cong \mathbf{Z}/N$  for some  $N \geq 1$ .

**Example 5.20.** Let  $K = \mathbf{Z}/2 \times \mathbf{Z}/2$  be the product of two copies of  $\mathbf{Z}/2$ , with addition defined componentwise, so that  $(a, b) + (c, d) = (a + c, b + d)$  where  $a + c$  and  $b + d$  are computed in  $\mathbf{Z}/2$ . This is a group with 4 elements, but  $K$  is not isomorphic to  $\mathbf{Z}/4$ . Indeed,  $\mathbf{Z}/4$  has an two elements of order 4, but  $K$  has no element of order 4.

## 5.1 Exercises

**Exercise 5.1.** Prove that if  $n \geq 3$ , then  $S_n$  is not cyclic.

**Exercise 5.2.** Recall the group  $(\mathbf{Z}/N)^\times$  from Exercise 3.4. Let  $\phi(N)$  be the number of elements of  $(\mathbf{Z}/N)^\times$ . The function  $\phi$  is called the **Euler totient function**.<sup>1</sup>

- (a) Show that if  $M, N \geq 1$  are relatively prime, then  $\phi(MN) = \phi(M)\phi(N)$ .
- (b) Show that if  $n \geq 1$ , then for every prime number  $p$  we have  $\phi(p^n) = p^{n-1}\phi(p)$ .
- (c) Show that  $\phi(p) = p - 1$  if  $p$  is prime.
- (d) What is  $\phi(3072)$ ?

**Exercise 5.3.** Let  $f: X \rightarrow Y$  be a bijection. Consider the permutation groups  $S_X$  and  $S_Y$  and the function  $g: S_X \rightarrow S_Y$  defined by  $g(h) = f \circ h \circ f^{-1}$  for  $h \in S_X$ . Prove that  $g$  is a group isomorphism.

---

<sup>1</sup>This is just a name. As far as I know, “totient” does not mean anything else.

## 6 Subgroups (10/02)

**Definition 6.1** (Subgroups). Let  $G$  be a group and let  $X$  be a subset of  $G$  we say that  $X$  is a **subgroup** if the following conditions hold:

- (i)  $X$  is nonempty,
- (ii) if  $a \in X$ , then  $a^{-1} \in X$ , and
- (iii) if  $a, b \in X$ , then  $ab \in X$ .

These conditions imply

- (iv)  $e \in X$ ,

**Example 6.2.** The group  $\mathbf{Z}$  is a subgroup of  $\mathbf{R}$ , while  $\mathbf{N}$  is not a subgroup of  $\mathbf{Z}$  because (ii) fails.

**Example 6.3.** If  $V$  is a vector space and  $W \subseteq V$  is a subspace, then  $W$  is a subgroup of  $V$ .

**Example 6.4.** The set of positive real numbers  $\mathbf{R}_+^\times$  is a subgroup of the group  $\mathbf{R}^\times$  of non-zero real numbers under multiplication.

**Remark 6.5.** If  $G$  is a group and  $X \subseteq G$  is a subgroup, then  $X$  is a group. Here, we use condition (iii) to view the restriction of the binary operation from  $G$  to  $X$  as a binary operation on  $X$ . Specifically, write  $a \cdot_G b$  for the binary operation in  $G$  and if  $a, b \in X$ , define  $X \times X \rightarrow X$  by  $a \cdot_X b = a \cdot_G b$ , viewed as an element of  $X$ . Then,  $X$  together with this binary operation is a group.

**Lemma 6.6.** If  $f: G \rightarrow H$  is a group homomorphism, then the image of  $f$ , written  $\text{im}(f)$  or  $f(G)$ , is a subgroup of  $H$  and  $f$  induces a group homomorphism  $G \rightarrow f(G)$ .

*Proof.* Since  $G$  has an identity element  $e$ , there is an element  $f(e) \in f(G)$ , so  $f(G)$  is nonempty. Similarly, if  $x, y \in f(G)$ , we can write  $x = f(g)$  and  $y = f(h)$  for some  $g, h \in G$  and hence  $xy = f(g)f(h) = f(gh)$ , so  $xy \in f(G)$  as well. Finally,  $x^{-1} = f(g^{-1})$ . That the induced function  $G \rightarrow f(G)$  is a group homomorphism follows from the fact that  $f: G \rightarrow H$  is.  $\square$



**Lemma 6.7.** If  $f: G \rightarrow H$  is an injective group homomorphism, then the induced function  $G \rightarrow f(G)$  is a group isomorphism.

*Proof.* It is surjective by definition and injective by hypothesis.  $\square$

**Example 6.8** (Subgroup lattice of  $S_3$ ). Figure 1 shows the subgroups of  $S_3$  arranged into what is called a subgroup lattice. The lines represent containment. The group on the middle left is isomorphic to  $\mathbf{Z}/3$  while the three groups on the middle right are isomorphic to  $\mathbf{Z}/2$ . These are all of the subgroups because one checks that if a subgroup of  $S_3$  has an element of order 2 and an element of order 3, then it is all of  $S_3$ . Note that two distinct elements of order 2 multiply to an element of order 3.

**Example 6.9** (Subgroup lattice of  $\mathbf{Z}/6$ ). Figure 2 shows the subgroup lattice of  $\mathbf{Z}/6$ . The middle left subgroup is isomorphic to  $\mathbf{Z}/3$  and the middle right to  $\mathbf{Z}/2$ .

**Theorem 6.10** (Cayley's theorem). If  $G$  is a group, then there is an injective group homomorphism  $\ell: G \rightarrow S_G$ , where  $S_G$  denotes the group of bijections from the set of elements of  $G$  to itself.

Figure 1: Subgroups of  $S_3$ .Figure 2: Subgroups of  $\mathbf{Z}/6$ .

*Proof.* Given  $g \in G$ , let  $\ell_g: G \rightarrow G$  be defined by  $\ell_g(h) = gh$ . This is a bijection by the Latin square property, which holds for all groups. Alternatively,  $\ell_g(g^{-1}h) = g(g^{-1}h) = h$ , and this is a unique solution to  $\ell_g(x) = h$ . Thus, the assignment  $g \mapsto \ell_g$  gives a function  $\ell: G \rightarrow S_G$  where  $\ell(g) = f_g$ . The claim is that this is an injective group homomorphism. If  $\ell_g = \ell_{g'}$  for  $g, g' \in G$ , then  $g = \ell_g(e) = \ell_{g'}(e) = g'$ , which proves injectivity. Now,  $(\ell_g \circ \ell_{g'})(h) = \ell_g(\ell_{g'}(h)) = \ell_g(g'h) = g(g'h) = (gg')h = \ell_{gg'}(h)$ , so  $\ell_g \circ \ell_{g'} = \ell_{gg'}$  and the function  $\ell$  is a group homomorphism.  $\square$

**Remark 6.11.** Cayley's theorem implies every group is a subgroup of a permutation group. However, this can be rather inefficient. For example, the injective group homomorphism  $\ell: \mathbf{Z}/N \rightarrow S_{\mathbf{Z}/N} \cong S_N$  embeds the group  $\mathbf{Z}/N$  of order  $N$  into a group of order  $N!$ . What does this embedding look like? It sends  $1 \in \mathbf{Z}/N$  to a cycle  $c = (0\ 1\ \dots\ N-1)$  (where we use  $\{0, \dots, N-1\}$  instead of  $\{1, \dots, N\}$  since these are the elements of  $\mathbf{Z}/N$ ) and  $a \in \mathbf{Z}/N$  to  $c^a$ .

**Example 6.12.** What about  $S_3$ ? This is a group with 6 elements, so the homomorphism from Cayley's theorem is a group homomorphism  $\ell: S_3 \rightarrow S_6$ . Let's label the elements of  $S_3$  as:

$$\begin{pmatrix} e & (12) & (13) & (23) & (123) & (132) \\ 1 & 2 & 3 & 4 & 5 & 6 \end{pmatrix}.$$

Then,  $e$  of  $S_3$  gets mapped to the identity element  $e$  of  $S_6$ . A cycle decomposition for  $\ell(12)$  is  $(12)(36)(45)$ .

**Remark 6.13** (Orders and group homomorphisms). If  $f: G \rightarrow H$  is a group homomorphism and  $a \in G$  has order  $n$ , then  $f(a)$  has order dividing  $n$ . Indeed,  $f(a)^n = f(a^n) = f(e) = e$ . Thus, in the example above  $\ell(12)$  has order dividing 2. But, it's clearly not of order 1, so its order must be exactly 2, which means the only cycles appearing in its cycle decomposition are of length 1 or 2.

## 6.1 Exercises

**Exercise 6.1.** Show that if  $G$  is a group and  $a \in G$  is an element satisfying  $a^n = e$  for some integer  $n \geq 1$ , then the order of  $a$  divides  $n$ .

**Exercise 6.2.** Draw the lattice of subgroups for the group  $K = \mathbf{Z}/2 \times \mathbf{Z}/2$ . (Sample LaTeX code is in Discord.)

**Exercise 6.3.** Draw the lattice of subgroups for the group  $\mathbf{Z}/12$ .

**Exercise 6.4.** Using Example 6.12, find a cycle decomposition for  $\ell(1\,2\,3)$ .

## 7 Group actions (10/04)

**Definition 7.1.** Let  $G$  be a group and  $X$  a set. An **action** of  $G$  on  $X$  is a function  $k: G \times X \rightarrow X$ , written  $a \cdot x = k(a, x)$  for  $a \in G$  and  $x \in X$ , satisfying the following axioms:

- (a)  $e \cdot x = x$  for all  $x \in X$  where  $e$  is the identity element of  $G$ ;
- (b)  $a \cdot (b \cdot x) = (ab) \cdot x$  for all  $a, b \in G$  and  $x \in X$ .

**Example 7.2.** The group  $\mathbf{Z}$  acts on  $\mathbf{R}$  by  $n \cdot x = n + x$  for  $n \in \mathbf{Z}$  and  $x \in \mathbf{R}$ .

**Example 7.3.** The group  $S_X$  acts on  $X$  by  $f \cdot x = f(x)$  for  $f \in S_X$  and  $x \in X$ . In particular,  $S_n$  acts on the set  $\{1, \dots, n\}$ .

**Example 7.4.** If  $V$  is a real vector space, then the group  $\mathbf{R}^\times$  of non-zero real numbers acts on  $V$  by scalar multiplication: if  $v \in V$  and  $\alpha \in \mathbf{R}^\times$ , then  $\alpha \cdot v = \alpha v$ .

**Example 7.5.** If  $G$  is a group, it acts on itself by left multiplication: for  $g, h \in G$ , we let  $g \cdot h = gh$ . Here, we view the  $G$  which acts as the *left*  $G$  in  $m: G \times G \rightarrow G$ . This is called the *left regular action* of  $G$  on itself. The formula  $g \cdot h = hg$  would not generally be a group action of  $G$  on itself. Why not?

**Example 7.6** (Return to Exercise 4.2). We can learn about a group  $G$  via its actions. For example, consider a symmetric group  $S_n$ . The symmetric group acts on the set  $F$  of functions  $\mathbf{R}^n \rightarrow \mathbf{R}$  as follows. Given  $a \in S_n$  and  $f: \mathbf{R}^n \rightarrow \mathbf{R}$ , we let  $(a \cdot f)(x_1, \dots, x_n) = f(x_{a(1)}, x_{a(2)}, \dots, x_{a(n)})$ , i.e., by reordering the inputs. Let  $g(x) = \prod_{1 \leq i < j \leq n} (x_i - x_j)$ . This polynomial is called the Vandermonde polynomial. Note that for any  $a \in S_n$ , either  $a \cdot g = g$  or  $a \cdot g = -g$ . For example, if  $n = 4$ , this polynomial is

$$g(x_1, x_2, x_3, x_4) = (x_1 - x_2)(x_1 - x_3)(x_1 - x_4)(x_2 - x_3)(x_2 - x_4)(x_3 - x_4).$$

The element  $a = (1\ 2\ 3\ 4)$  of  $S_4$  then acts as

$$(a \cdot g)(x_1, x_2, x_3, x_4) = (x_2 - x_3)(x_2 - x_4)(x_2 - x_1)(x_3 - x_4)(x_3 - x_1)(x_4 - x_1) = -g(x_1, x_2, x_3, x_4).$$

Let  $S_n$  act on  $\{1, -1\}$  by letting  $a \cdot \epsilon = \gamma$  if  $a \cdot (\epsilon g) = \gamma g$ . In the example above, the 4-cycle  $a$  has  $a \cdot 1 = -1$  and  $a \cdot (-1) = 1$ . If  $a \in S_n$  is a transposition, then  $a \cdot 1 = -1$ . To see this, suppose that  $a = (cd)$  where  $1 \leq c < d \leq n$ . If  $i < c$ , then  $a \cdot (x_i - c) = (x_i - d)$  and if  $d < j$ , then  $a \cdot (c - x_j) = (d - x_j)$ . We also have  $a \cdot (x_i - x_j) = (x_j - x_i) = -(x_i - x_j)$ . Finally, if  $c < i < d$ ,

$$a \cdot (x_c - x_i)(x_i - x_d) = (x_d - x_i)(x_i - x_c) = -(x_i - x_d)(-(x_c - x_i)) = (x_c - x_i)(x_i - x_d).$$

Collating these calculations, it follows that  $a \cdot v = -v$  for  $a = (cd)$ . Thus, by axiom (b) of a group action, if  $a$  is a product of  $k$  transpositions, then  $a \cdot 1 = (-1)^k$ . This proves the claim from Exercise 4.2 as if  $(-1)^k = (-1)^m$ , then  $k \equiv m \pmod{2}$ .

The next theorem says that group actions of  $G$  on  $X$  are “the same” as group homomorphisms  $G \rightarrow S_X$ .

**Theorem 7.7.** Let  $G$  be a group and  $X$  as set. There is a bijection

$$\{\text{actions } k \text{ of } G \text{ on } X\} \xrightarrow{k \mapsto f_k} \text{Hom}(G, S_X).$$

*Proof.* Next time. □

**Example 7.8.** The action of  $S_n$  on the Vandermonde polynomial induces, via the theorem, a surjective group homomorphism  $S_n \rightarrow S_{\{1, -1\}}$ , which we view as a group homomorphism  $\epsilon: S_n \rightarrow S_2 \cong \mathbf{Z}/2 \cong \{1, -1\}$ , where  $\{1, -1\}$  is a group under multiplication. The **sign** of an element  $a \in S_n$  is  $\epsilon(a) \in \{1, -1\}$ .

## 7.1 Exercises

**Exercise 7.1.** Suppose that  $G$  is a finite group of even order. Show that there exists  $x \neq e$  in  $G$  with  $x^2 = e$ .

**Exercise 7.2.** Show that every finite group  $G$  of order 4 is isomorphic to either  $\mathbf{Z}/4$  or to  $K = \mathbf{Z}/2 \times \mathbf{Z}/2$ .

**Exercise 7.3.** Show that a finite group  $G$  of order 5 is isomorphic to  $\mathbf{Z}/5$ .



## 8 The adjoint homomorphism (10/06)

Our next theorem says that group actions of  $G$  on  $X$  are “the same” as group homomorphisms  $G \rightarrow S_X$ .

**Theorem 8.1.** *Let  $G$  be a group and  $X$  as set. There is a bijection*

$$\{\text{actions } k \text{ of } G \text{ on } X\} \xrightarrow{k \mapsto f_k} \text{Hom}(G, S_X),$$

where  $\text{Hom}(G, S_X)$  denotes the set of group homomorphisms from  $G$  to  $S_X$ .

*Proof.* Let  $k: G \times X \rightarrow X$  be a group action; we will write  $g \cdot_k x$  for  $k(g, x)$  in this proof. For  $g \in G$ , let  $f_k(g)$  be the function  $X \rightarrow X$  defined by  $f_k(g)(x) = k(g, x) = g \cdot_k x$ . This is a bijection as one sees by observing that  $f_k(g^{-1})$  is an inverse using (a) and (b) from the definition of a group action. Therefore,  $f_k$  is a function  $G \rightarrow S_X$ . In fact, this is a group homomorphism. Indeed,  $f_k(gh)(x) = gh \cdot_k x = g \cdot_k (h \cdot_k x) = f_k(g)(f_k(h)(x))$  for all  $g, h \in G$  and  $x \in X$ . Therefore,  $f_k(gh) = f_k(g) \circ f_k(h)$ , as desired.

To show that the assignment  $k \mapsto f_k$  is bijective, assume first that  $k$  and  $n$  are distinct group actions. Then, there exists a pair  $(g, x) \in G \times X$  such that  $g \cdot_k x \neq g \cdot_n x$ . It follows that  $f_k(g) \neq f_n(g)$ . This shows injectivity.

Given a group homomorphism  $f: G \rightarrow S_X$ , we define a new group action  $k_f$  of  $G$  on  $X$  by letting  $g \cdot_{k_f} x = f(g)(x)$ . By definition,  $f_{k_f}(g)(x) = g \cdot_{k_f} x = f(g)(x)$ , so  $f_{k_f}(g) = f(g)$  for all  $g \in G$  and hence  $f_{k_f} = f$ , which proves surjectivity.  $\square$

**Definition 8.2.** If  $k$  is an action of  $G$  on  $X$ , then  $f_k: G \rightarrow S_X$  is called the **adjoint homomorphism**. If  $f: G \rightarrow S_X$  is a homomorphism, then  $k_f$  is called the **action associated to  $f$** .

**Example 8.3.** Let  $G$  be a group and consider its left regular action on itself  $m: G \times G \rightarrow G$ . The adjoint homomorphism  $\ell = f_m: G \rightarrow S_G$  is the homomorphism used in the proof of Cayley’s Theorem 6.10.

**Example 8.4.** Recall the group  $K = \mathbf{Z}/2 \times \mathbf{Z}/2$ , sometimes known as the **Klein four-group**. It has four elements, which we label as follows:

$$\begin{pmatrix} (0,0) & (1,0) & (0,1) & (1,1) \\ 1 & 2 & 3 & 4 \end{pmatrix}.$$

The adjoint homomorphism  $\ell: K \rightarrow S_K$  we view, using the labeling above, as a homomorphism  $\ell: K \rightarrow S_4$ . A cycle decomposition of  $\ell(1,0)$  is  $((0,0)(1,0))((0,1)(1,1)) = (1\,2)(3\,4)$ .

### 8.1 Exercises

**Exercise 8.1.** Say that an action of a group  $G$  on a set  $X$  is **trivial** if  $g \cdot x = x$  for all  $g \in G$  and  $x$  on  $X$ . Suppose that  $p$  is a prime and that  $X$  is a set with fewer than  $p$  elements. Show that all actions of  $\mathbf{Z}/p$  on  $X$  are trivial.

**Exercise 8.2.** Compute the set  $\text{Hom}(\mathbf{Z}/2, S_4)$  of group homomorphisms into  $S_4$ . Use your computation to describe all group actions of  $\mathbf{Z}/2$  on  $\{1, 2, 3, 4\}$ .

## 9 Dihedral groups and some properties of group actions (10/09)

**Example 9.1** (Dihedral groups). Fix  $n \geq 3$ . Let  $X$  be a regular  $n$ -gon with vertices labeled as  $\{1, \dots, n\}$  sitting in  $\mathbf{R}^2$  centered at the origin. Let  $D_{2n} \subseteq S_n$  be the set of permutations of the vertex set  $\{1, \dots, n\}$  consisting of those which can be achieved by a rigid motion of  $X$  in  $\mathbf{R}^3$  returning  $X$  bijectively to itself. Among these, we single out two. Let  $r$  denote the permutation obtained by counterclockwise rotation about the origin by  $\frac{2\pi i}{n}$ . Let  $s$  denote the reflection across the line between 1 and the origin. Geometrically, we see that  $rs = sr^{-1}$ . This implies that the elements  $\{e, r, r^2, \dots, r^{n-1}, s, sr, \dots, sr^{n-1}\}$  form a subgroup of  $S_n$ . Indeed,

$$(s^a r^b)(s^c r^d) = s^a s^c r^{(-1)^c b + d} = s^{a+c} r^{(-1)^c b + d} = s^e r^f,$$

where  $e \equiv a + c \pmod{2}$  is in  $\{0, 1\}$  and  $f \equiv (-1)^c b + d \pmod{2}$  is in  $\{0, \dots, n\}$ . We claim that this is all of  $D_{2n}$  and hence that  $D_{2n}$  is a subgroup of  $S_n$ , known as the **dihedral group of order  $2n$** . To see this, suppose that  $x \in D_{2n}$ . We want to show that  $x$  is in the list of  $2n$  elements above. We can compose with a rotation and assume that  $x$  sends the vertex 1 to itself. Then, since it arises from a rigid motion of  $\mathbf{R}^3$ , we must have that  $x$  sends either 2 to itself and  $n-1$  to itself, or it sends 2 to  $n-1$  and  $n-1$  to 2. In the first case, it must be the identity. In the second case, it must be the reflection  $s$ .

**Remark 9.2.** Let  $n = 4$  and consider the dihedral group  $D_8$  of order 8. Let  $s$  denote the reflection across the diagonal through 1 and 3 and let  $s'$  denote the reflection across the diagonal through 2 and 4. Then,  $ss'$  has cycle decomposition  $(13)(24)$ . But, so does  $r^2$ . So,  $ss' = r^2$ .

### 9.1 Exercises

**Exercise 9.1.** Make a list of all elements of  $D_8$ , their orders, and a cycle decomposition for each (with respect to the action above of  $D_8$  on  $\{1, 2, 3, 4\}$ ).

**Exercise 9.2.** Find the lattice of subgroups of  $D_8$ .

**Exercise 9.3.** Find the lattice of subgroups of  $D_{10}$ .

## 10 Some properties of group actions (10/11)

Recall the following definition from Section 8.

**Definition 10.1** (Trivial actions). Say that an action of  $G$  on  $X$  is **trivial** if  $g \cdot x = x$  for all  $x \in X$  and all  $g \in G$ . This is the case if and only if the adjoint homomorphism  $f: G \rightarrow S_X$  satisfies  $f(g) = e$  for all  $g \in G$ .

At the opposite extreme, we have the faithful actions.

**Definition 10.2.** The action of a group  $G$  on a set  $X$  is **faithful** if the adjoint homomorphism  $G \rightarrow S_X$  is injective.

**Remark 10.3.** In other words, an action of  $G$  on  $X$  is faithful if different elements of  $G$  produce different permutations on  $X$ . Unwinding, this means that for each pair of distinct elements  $f, g \in G$  there exists  $x \in X$  such that  $f \cdot x \neq g \cdot x$ .

**Remark 10.4.** If  $X$  is a set and  $S_X$  is the permutation group of  $X$ , then any subgroup  $G \subseteq S_X$  comes with an action on  $X$  which is faithful.

**Example 10.5.** As  $D_{2n}$  is a subgroup of  $S_n$ , its action on  $\{1, \dots, n\}$  is faithful.

**Definition 10.6** (Orbits and stabilizers). Let  $G$  be a group acting on a set  $X$ .

- (i) If  $x \in X$ , the **orbit** of  $G$  containing  $x$  is the set  $G \cdot x = \{g \cdot x | g \in G\}$ . Alternatively, if  $k: G \times X \rightarrow X$  denotes the action map, it is the image of  $G \times \{x\}$  under  $k$ .
- (ii) If  $x \in X$ , the **stabilizer** of  $x$  in  $G$  is the set  $G_x = \{g \in G | g \cdot x = x\}$ .

**Lemma 10.7.** If  $G$  acts on a set  $X$  and if  $x \in X$ , then the stabilizer  $G_x \subseteq G$  is a subgroup.

*Proof.* Of course,  $e \in G_x$ . We also have that if  $g \in G_x$ , then  $g^{-1} \in G_x$  as  $g^{-1} \cdot x = g^{-1} \cdot (g \cdot x) = (g^{-1}g) \cdot x = e \cdot x = x$ . Similarly, if  $g, h \in G_x$ , then  $(gh) \cdot x = g \cdot (h \cdot x) = g \cdot x = x$ , so  $gh \in G_x$ .  $\square$

**Example 10.8.** Consider  $D_{2n}$  acting on the  $n$ -gon  $X_n$  with vertex set  $\{1, \dots, n\}$  as in Definition 9.1. The orbit of any vertex is  $\{1, \dots, n\}$ . What about the orbit of a point on  $X$  that is not a vertex? The stabilizer of 1 in  $G$  is  $G_1 = \{e, s\}$ . Indeed, any rotation must “move” 1. Any element  $f$  which fixes 1 must either send 2 to itself, in which case  $f = 1$  or it sends 2 to  $n$  and  $n$  to 2, in which case  $sf = e$ , or  $f = s$ . The stabilizer of a point which is not a vertex is trivial if  $n$  is even and usually trivial if  $n$  is odd, the exception being the points opposite to vertices which are fixed by appropriate reflections.

**Philosophy 10.9.** The approach to defining the dihedral group is very helpful in finding new groups. For example, let  $T$  in  $\mathbf{R}^3$  be a regular tetrahedron with vertex set  $\{1, 2, 3, 4\}$ . Among all rigid motions of  $\mathbf{R}^3$ , there are those which act bijectively on  $T$ , and must send vertices to vertices, edges to edges, and faces to faces. How many are there? I can send 1 to any vertex  $i \in \{1, 2, 3, 4\}$ , which amounts to four choices of where 1 goes. Once that is fixed, 2 must go to one an element of  $\{1, 2, 3, 4\} - \{i\}$ , so there are three more choices. But, then it is fixed. For example, if 1 maps to 3 and 2 maps to 1, then one sees by rigidity that 3 maps to 2 and 4 maps to 3.

**Example 10.10** (The conjugation action). Let  $G$  be a group. We define a new action of  $G$  on itself, given by conjugation. Namely, let  $c: G \times G \rightarrow G$  be defined by  $c(g, h) = ghg^{-1}$ . This is the result of *conjugating*  $h$  by  $g$ . We have  $c(e, h) = ehe^{-1} = h$  for all  $h \in G$  and we have  $c(f, c(g, h)) = f(ghg^{-1})f^{-1} = (fg)h(fg)^{-1} = c(fg, h)$ . So, conjugation defines a group action of  $G$  on itself. The conjugation action is always different from the left regular action if  $G$  is not the trivial group  $\{e\}$ .

**Question 10.11.** When is the conjugation action trivial?

**Definition 10.12** (Orbit set). Let a group  $G$  act on a set  $X$ . For  $x, y \in X$ , write  $x \sim y$  if there exists  $g \in G$  such that  $g \cdot x = y$ . This defines an equivalence relation on  $X$ . Indeed,  $e \cdot x = x$  so  $x \sim x$  (reflexivity), if  $g \cdot x = y$ , then  $g^{-1} \cdot y = x$  (reflexivity), and if  $g \cdot x = y$  and  $h \cdot y = z$ , then  $(hg) \cdot x = z$  (transitivity). The equivalence classes are precisely the orbits. We write  $X/G$  for the set of orbits. The quotient function  $f: X \rightarrow X/G$  sends  $x \in X$  to  $G \cdot x \in X/G$ .

**Question 10.13.** What does the orbit set of  $D_{2n}$  acting on  $X_n$  look like? It is bijective to the half-open line segment  $L$  from vertex 1 (inclusive) to vertex 2 (not inclusive). Indeed, for each  $x \in X_n$  there is a unique  $y$  on  $L$  such that  $g \cdot x = y$  for some  $g \in D_{2n}$ . Note that this is the same orbit set as that corresponding to the action of  $\mathbf{Z}/n$  on  $X_n$  by rotations by multiples of  $\frac{2\pi}{n}$ .

**Definition 10.14** (Transitive actions). The action of a group  $G$  on a set  $X$  is **transitive** if  $X/G$  is a point or, equivalently, if there is only one orbit or, equivalently, if for all pairs  $x, y \in X$  there exists  $g \in G$  such that  $g \cdot x = y$ .

## 10.1 Exercises

**Exercise 10.1.** Let  $G = S_n$  act on  $X = \{1, \dots, n\}$  via permutations.

- (a) What is the orbit  $G \cdot 1$ ?
- (b) What is the stabilizer  $G_1$  of 1 in  $G$ ? (It is isomorphic to a group we have a name for.)
- (c) What is the set of orbits  $X/G$ ?
- (d) Is the action faithful?
- (e) Is the action transitive?

**Exercise 10.2.** Repeat Exercise 8.2(a)-(e) for the left regular action of a group  $G$  on itself (where 1 is replaced by  $e$  in parts (a) and (b)).

**Exercise 10.3.** Repeat Exercise 8.2(a)-(e) for the conjugation action of  $G = D_8$  on itself (where 1 is replaced by  $e$  in parts (a) and (b)).

**Exercise 10.4.** Arguing as in Philosophy 10.9, compute the order of the group of rigid motions of an icosahedron in  $\mathbf{R}^3$ .