

Linear Algebra

Math 314 Fall 2025

Today's poll code:

4LHPXC

Lecture 4

Office hours

Mondays 5–6 pm

Wednesdays 2–3 pm

in Avery 339 (Dr. Grifo)

Tuesdays 11–noon

Thursdays 1–2 pm

in Avery 337 (Kara)

To do list:

- WeBWorK 1.3 and 2.1 due tomorrow
- WebWork 2.2 due Friday
- Webwork 2.3 due Tuesday September 16

Quiz 3 on Friday
on lectures 4–5

Office hours

Mondays 5–6 pm and Wednesdays 2–3 pm
in Avery 339 (Dr. Grifo)

Tuesdays 11–noon and Thursdays 1–2 pm
in Avery 337 (Kara)

Quick Recap

Let v_1, \dots, v_p be vectors in \mathbb{R}^n .

span of v_1, \dots, v_p = set of all linear combinations of v_1, \dots, v_p

$$\text{span}\{v_1, \dots, v_p\} = \{c_1v_1 + \dots + c_pv_p \mid c_i \in \mathbb{R}\}$$


if $v_1 \neq 0$

$$\begin{aligned}\text{span}(\{v_1\}) &= \text{line determined by } v_1 \\ &= \text{all multiples of } v_1\end{aligned}$$

Let v_1, \dots, v_p be vectors in \mathbb{R}^n .

span of v_1, \dots, v_p = set of all linear combinations of v_1, \dots, v_p

$$\text{span}\{v_1, \dots, v_p\} = \{c_1 v_1 + \dots + c_p v_p \mid c_i \in \mathbb{R}\}$$



$\text{span}(\{v_1, v_2\})$ = plane determined by v_1 and v_2
if $v_1 \neq 0$ and $v_2 \neq 0$ and v_1 and v_2 not multiples of each other

Let v_1, \dots, v_p be vectors in \mathbb{R}^n .

span of v_1, \dots, v_p = set of all linear combinations of v_1, \dots, v_p

$$\text{span}\{v_1, \dots, v_p\} = \{c_1v_1 + \dots + c_pv_p \mid c_i \in \mathbb{R}\}$$

$$\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \end{bmatrix} \right\} =$$

Let v_1, \dots, v_p be vectors in \mathbb{R}^n .

span of v_1, \dots, v_p = set of all linear combinations of v_1, \dots, v_p

$$\text{span}\{v_1, \dots, v_p\} = \{c_1v_1 + \dots + c_pv_p \mid c_i \in \mathbb{R}\}$$

$$\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \end{bmatrix} \right\} = \mathbb{R}^2$$

Question:

Is $\begin{bmatrix} 7 \\ 4 \\ -3 \end{bmatrix}$ a linear combination of $\begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix}$ and $\begin{bmatrix} 2 \\ 5 \\ 6 \end{bmatrix}$?

Equivalently:

Does the system $x_1 \begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 7 \\ 4 \\ -3 \end{bmatrix}$ have a solution?

$$\begin{bmatrix} 1 & 2 & 7 \\ -2 & 5 & 4 \\ -5 & 6 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

No pivots in the last column

system is consistent

Answer: yes.

Question:

Is $\begin{bmatrix} 7 \\ 4 \\ -3 \end{bmatrix}$ in $\text{span}\left\{\begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ 6 \end{bmatrix}\right\}$?

Equivalently:

Does the system $x_1 \begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 7 \\ 4 \\ -3 \end{bmatrix}$ have a solution?

$$\begin{bmatrix} 1 & 2 & 7 \\ -2 & 5 & 4 \\ -5 & 6 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

No pivots in the last column

system is consistent

Answer: yes.

Question:

Is $\begin{bmatrix} 7 \\ 4 \\ -3 \end{bmatrix}$ in $\text{span}\left\{\begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ 6 \end{bmatrix}\right\}$?

$$\begin{bmatrix} 1 & 2 & 7 \\ -2 & 5 & 4 \\ -5 & 6 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

No pivots in the last column
system is consistent
Answer: yes.

Solution set:

$$\{(3, 2)\}$$

$$3 \begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 7 \\ 4 \\ -3 \end{bmatrix}.$$

Today's poll code:

4LHPXC

Today's poll code:

4LHPXC

Fact: $\begin{bmatrix} 3 & -1 & 1 & 3 \\ 0 & 1 & 2 & 6 \\ 1 & 2 & 2 & 9 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 2 \end{bmatrix}$

Is $\begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$ in $\text{span}\left\{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}\right\}$?

- A. Yes B. No

Today's poll code:

4LHPXC

What is $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$?

A. $\left\{ \begin{bmatrix} a \\ a \end{bmatrix} : a \text{ is any value} \right\}$

B. $\left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$

C. \mathbb{R}^2

Matrix Equations

$$Ax = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

Above the matrix A , there is a purple bracket labeled A . Above the vector x , there is a purple bracket labeled x .

$$\begin{array}{c}
A \\
\curvearrowleft \quad \curvearrowright \\
x
\end{array}$$

$$Ax = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \cdots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}$$

$$= \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix}.$$

$$\begin{aligned}
& A \\
& \overbrace{\quad}^A \quad \overbrace{\quad}^x \\
Ax &= \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix} + x_2 \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix} + \cdots + x_n \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix} \\
&= \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix}. \\
&= x_1a_1 + \cdots + x_na_n
\end{aligned}$$

Theorem. Let c be a scalar, let $u, v \in \mathbb{R}^n$,

and let A be an $m \times n$ matrix. Then

$$A(u + v) = Au + Av \quad \text{and} \quad A(cu) = c(Au).$$

$$\left\{ \begin{array}{l} a_{11}x_1 + \cdots + a_{1n}x_n = b_1 \\ \vdots \\ a_{m1}x_1 + \cdots + a_{mn}x_n = b_m \end{array} \right.$$

coefficient matrix

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

constant vector

$$\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

matrix equation

$$Ax = b$$

$$\begin{cases} x_1 + 3x_2 = 4 \\ -x_1 + x_2 = 1 \end{cases}$$



matrix equation

$$\begin{bmatrix} 1 & 3 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$$

The system $Ax = b$ has a solution

if and only if

b is a linear combination of the columns of A .

Theorem. Fix an $m \times n$ matrix A . The following are equivalent:

1. The system $Ax = b$ has a solution for *every* vector $b \in \mathbb{R}^m$.
2. Every vector $b \in \mathbb{R}^m$ is a linear combination of the columns of A .
3. The columns of A span \mathbb{R}^m .
4. The coefficient matrix A has a pivot in every row.

The last statement is about A , the coefficient matrix of the system,
not the augmented matrix.

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{pivot in every row}$$



$Ax = b$ has solutions for every $b \in \mathbb{R}^2$

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{pivot in every row}$$



$Ax = b$ has solutions for every $b \in \mathbb{R}^2$

$$\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} = \mathbb{R}^2$$

Homogeneous systems

A linear system is **homogeneous** if we can write it as

$$Ax = 0.$$

A linear system is **homogeneous** if we can write it as

$$Ax = 0.$$

A homogeneous system always has a solution:

the **trivial solution** $x = 0$.

A linear system is **homogeneous** if we can write it as

$$Ax = 0.$$

A homogeneous system always has a solution:

the **trivial solution** $x = 0$.

A solution $x \neq 0$ is called **nontrivial**.

$$\begin{cases} 3x_1 + 5x_2 - 4x_3 = 0 \\ -3x_1 - 2x_2 + 4x_3 = 0 \\ 6x_1 + x_2 - 8x_3 = 0 \end{cases} \quad \text{homogeneous system}$$

$$\begin{cases} 3x_1 + 5x_2 - 4x_3 = 0 \\ -3x_1 - 2x_2 + 4x_3 = 0 \\ 6x_1 + x_2 - 8x_3 = 0 \end{cases} \quad \text{homogeneous system}$$

$$\underbrace{\begin{bmatrix} 3 & 5 & -4 \\ -3 & -2 & 4 \\ 6 & 1 & -8 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}}_b \quad \text{matrix equation}$$

$$\begin{cases} 3x_1 + 5x_2 - 4x_3 = 0 \\ -3x_1 - 2x_2 + 4x_3 = 0 \\ 6x_1 + x_2 - 8x_3 = 0 \end{cases} \quad \text{homogeneous system}$$

$$\begin{array}{c}
\left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ -3 & -2 & 4 & 0 \\ 6 & 1 & -8 & 0 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + R_1} \left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ 0 & 3 & 0 & 0 \\ 6 & 1 & -8 & 0 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - 2R_1} \left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & -9 & 0 & 0 \end{array} \right] \\
\xrightarrow{R_2 \rightarrow \frac{1}{3}R_2} \left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -9 & 0 & 0 \end{array} \right] \xrightarrow{R_1 \rightarrow R_1 - 5R_2} \left[\begin{array}{ccc|c} 3 & 0 & -4 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -9 & 0 & 0 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 + 9R_2} \left[\begin{array}{ccc|c} 3 & 0 & -4 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \\
\xrightarrow{R_1 \rightarrow \frac{1}{3}R_1} \left[\begin{array}{ccc|c} 1 & 0 & -\frac{4}{3} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].
\end{array}$$

Note that the last column doesn't really add any information!
When the system is homogeneous, can row reduce A instead.

$$\begin{cases} 3x_1 + 5x_2 - 4x_3 = 0 \\ -3x_1 - 2x_2 + 4x_3 = 0 \\ 6x_1 + x_2 - 8x_3 = 0 \end{cases} \quad \text{homogeneous system}$$

$$\left[\begin{array}{ccc} 3 & 5 & -4 \\ -3 & -2 & 4 \\ 6 & 1 & -8 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + R_1} \left[\begin{array}{ccc} 3 & 5 & -4 \\ 0 & 3 & 0 \\ 6 & 1 & -8 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - 2R_1} \left[\begin{array}{ccc} 3 & 5 & -4 \\ 0 & 3 & 0 \\ 0 & -9 & 0 \end{array} \right]$$

$$\xrightarrow{R_2 \rightarrow \frac{1}{3}R_2} \left[\begin{array}{ccc} 3 & 5 & -4 \\ 0 & 1 & 0 \\ 0 & -9 & 0 \end{array} \right] \xrightarrow{R_1 \rightarrow R_1 - 5R_2} \left[\begin{array}{ccc} 3 & 0 & -4 \\ 0 & 1 & 0 \\ 0 & -9 & 0 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 + 9R_2} \left[\begin{array}{ccc} 3 & 0 & -4 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

$$\xrightarrow{R_1 \rightarrow \frac{1}{3}R_1} \left[\begin{array}{ccc} 1 & 0 & -\frac{4}{3} \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right].$$

$$\begin{cases} 3x_1 + 5x_2 - 4x_3 = 0 \\ -3x_1 - 2x_2 + 4x_3 = 0 \\ 6x_1 + x_2 - 8x_3 = 0 \end{cases} \quad \text{homogeneous system}$$

$$\begin{array}{c}
\left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ -3 & -2 & 4 & 0 \\ 6 & 1 & -8 & 0 \end{array} \right] \xrightarrow{R_2 \rightarrow R_2 + R_1} \left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ 0 & 3 & 0 & 0 \\ 6 & 1 & -8 & 0 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - 2R_1} \left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & -9 & 0 & 0 \end{array} \right] \\
\xrightarrow{R_2 \rightarrow \frac{1}{3}R_2} \left[\begin{array}{ccc|c} 3 & 5 & -4 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -9 & 0 & 0 \end{array} \right] \xrightarrow{R_1 \rightarrow R_1 - 5R_2} \left[\begin{array}{ccc|c} 3 & 0 & -4 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -9 & 0 & 0 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 + 9R_2} \left[\begin{array}{ccc|c} 3 & 0 & -4 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \\
\xrightarrow{R_1 \rightarrow \frac{1}{3}R_1} \left[\begin{array}{ccc|c} 1 & 0 & -\frac{4}{3} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].
\end{array}$$

x_3 is free

$$x_1 = \frac{4}{3}x_3$$

$$x_2 = 0$$

$$\underbrace{\begin{bmatrix} 3 & 5 & -4 \\ -3 & -2 & 4 \\ 6 & 1 & -8 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}}_b \quad \text{homogeneous system}$$

x_3 is free

$$x_1 = \frac{4}{3}x_3 \qquad x_2 = 0$$

General solution:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = x_3 \begin{bmatrix} \frac{4}{3} \\ 0 \\ 1 \end{bmatrix} \quad \text{or} \quad t \begin{bmatrix} \frac{4}{3} \\ 0 \\ 1 \end{bmatrix}$$

$t = 0$ gives the trivial solution

Each choice of $t \neq 0$ gives a nontrivial particular solution

the general solution to a homogeneous system is
a linear combination of vectors with the free variables as coefficients

$$\underbrace{\begin{bmatrix} 3 & 5 & -4 \\ -3 & -2 & 4 \\ 6 & 1 & -8 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}}_x = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}}_b$$

homogeneous system

General solution in parametric vector form

$$x = x_3 \begin{bmatrix} \frac{4}{3} \\ 0 \\ 1 \end{bmatrix} \quad \text{or} \quad x = t \begin{bmatrix} \frac{4}{3} \\ 0 \\ 1 \end{bmatrix}$$

A **nonhomogeneous linear system** is a linear system of the form

$$Ax = b \quad \text{with } b \neq 0$$

Theorem. The general solution to the nonhomogeneous system $Ax = b$ is

$$x = \text{one particular solution} + \text{general solution to the homogeneous system } Ax = 0$$

A homogeneous system always has a solution:
the **trivial solution** $x = 0$.

A nonhomogeneous system might not have a solution.

$$\begin{bmatrix} 3 & 5 & -4 \\ -3 & -2 & 4 \\ 6 & 1 & 8 \end{bmatrix} x = \begin{bmatrix} 7 \\ 1 \\ -4 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 5 & -4 & 7 \\ -3 & -2 & 4 & 1 \\ 6 & 1 & 8 & -4 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -\frac{4}{3} & -1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$A \xrightarrow{\quad} \begin{bmatrix} 3 & 5 & -4 \\ -3 & -2 & 4 \\ 6 & 1 & 8 \end{bmatrix} x = \begin{bmatrix} 7 \\ 1 \\ -4 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 5 & -4 & 7 \\ -3 & -2 & 4 & 1 \\ 6 & 1 & 8 & -4 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -\frac{4}{3} & -1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

General solution in
parametric vector form

$$x = \begin{bmatrix} -1 \\ 2 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} \frac{4}{3} \\ 0 \\ 1 \end{bmatrix}$$

particular
solution

solution
to $Ax = 0$

Important

To write the solution set of a consistent system:

- 1) Row-reduce the augmented matrix into reduced echelon form.
- 2) Write each non-free variable in terms of the free ones.
- 3) Write the general solution x as a vector whose entries depend on the free variables (if there are free variables).
- 4) Decompose this as a linear combination of vectors where each coefficient is a free variable (plus possibly one term with coefficient 1 for a particular solution).

Important

Caution! Given a linear system $Ax = b$, there is a big difference between the coefficient matrix A and the augmented matrix $[A \ b]$.

- Is the system $Ax = b$ consistent? \implies look at the augmented matrix.
- The system $Ax = 0$ is always consistent.

We can solve the system by focusing only on A and then finding a particular solution, but if we do so we must remember A is *not* the augmented matrix of the system.

Today's poll code:

4LHPXC

Today's poll code:

4LHPXC

A homogeneous linear system is always consistent.

A. True

B. False

Today's poll code:

4LHPXC

A non-homogeneous linear system is always consistent.

A. True

B. False

To do list:

- WeBWorK 1.3 and 2.1 due tomorrow
- WebWork 2.2 due Friday
- Webwork 2.3 due next Tuesday

On Friday:

Quiz 3

**at the beginning
of the recitation
on Lectures 4-5**

Office hours

Mondays 5–6 pm and Wednesdays 2–3 pm
in Avery 339 (Dr Grifo)

Tuesdays 11–noon and Thursdays 1–2 pm
in Avery 337 (Kara)