

Constrained evolution of a core winter proteome across independently cold-adapted PACMAD grasses

Authors

Elad Oren^{1*}, Jingjing Zhai¹, Travis E Rooney², Ruthie Angelovici³, Charles O. Hale⁴, Lara J. Brindisi^{1,6}, Sheng-Kai Hsu¹, Christine M. Gault¹, Jian Hua⁵, Thuy La¹, Nicholas Lepak⁶, Qin Fu⁷, Edward S. Buckler^{1,6}, M. Cinta Romay^{1*}

Affiliations

1. Institute for Genomic Diversity, Cornell University, Ithaca, NY 14853, USA
2. Sesaco Corporation, Austin, TX 78701, USA
3. Division of Biological Sciences and Interdisciplinary Plant Group, University of Missouri, Columbia, Missouri, 65211, USA
4. School of Integrative Plant Sciences, Plant Breeding and Genetics Section, Cornell University, Ithaca, New York, 14853, USA
5. School of Integrative Plant Science, Plant Biology Section, Cornell University, Ithaca, New York, 14853, USA
6. USDA-ARS, Ithaca, NY, USA 14853
7. Proteomics and Metabolomics Facility, Cornell University, Ithaca, New York, 14850, USA

***Corresponding authors:**

Elad Oren (eladoren@volcani.agri.gov.il)

M. Cinta Romay (mcr72@cornell.edu)

The author responsible for distribution of materials integral to the findings presented in this article is Elad Oren.

Summary

- Grasses in the PACMAD clade independently colonized cold environments from warm-climate ancestors, but whether their molecular responses to freezing reflect shared evolutionary solutions or lineage-specific innovations remains unknown. We used comparative proteomics to test whether protein-level cold responses show stronger cross-species conservation than previously observed at the transcript level.
- We quantified seasonal rhizome proteomes (winter vs summer) from five PACMAD grass species grown in a common garden exposed to sustained sub-zero temperatures, identified differentially abundant proteins, and compared fold-change magnitudes across species using orthogroup-based correlation analyses. We further examined LEA3 protein structure through hydropathy profiling and motif analysis.
- Shared cold-responsive proteins showed substantially higher cross-species fold-change correlation ($\rho = 0.80$) than background proteins ($\rho = 0.45$), despite greater divergence in baseline abundance. LEA3 was the only ortholog elevated across all five species. The cold-sensitive maize ortholog contained a 15-amino-acid insertion that disrupted the conserved 11-mer motif architecture maintained in all cold-tolerant species examined.
- Independent evolution of freezing tolerance in PACMAD grasses is governed by evolutionary constraints on protein-level response magnitude. The structural divergence of LEA3 in maize suggests that transcriptional induction alone does not

ensure freezing tolerance; functional protection likely requires intact motif architecture.

Keywords: Rhizome, *Tripsacum*, Cold acclimation, Freezing tolerance, Proteomics, Parallel evolution, LEA proteins, PACMAD grasses

Introduction

Maize (*Zea mays* ssp. *mays*), a vital crop for global food security, is susceptible to cold stress due to its tropical origins, limiting its yield in temperate regions. Early planting of maize can improve nitrogen capture, net primary productivity, and yield (Ojeda-Rivera *et al.*, 2025). However, this strategy requires cold tolerance, which is lacking in current elite maize varieties (Lyons, 1973; Aroca *et al.*, 2012). Cold and freezing tolerance has arisen independently multiple times within the *Panicoideae*, *Aristidoideae*, *Chloridoideae*, *Micrairoideae*, *Aristavideae*, and *Danthonioideae* (PACMAD) clade, a major lineage of predominantly C4 grasses (*Poaceae*) that diverged ~64 Mya, with ancestral roots in the Afrotropics (Sage *et al.*, 2011; Gallaher *et al.*, 2022). These parallel origins of freezing tolerance provide a natural comparative framework. Among these, *Tripsacum*, the closest perennial and cold-tolerant relative of maize, having diverged about 650 Kya (Chen *et al.*, 2022), is valuable for enhancing maize cold tolerance.

PACMAD grasses originated in warm-season climates unlike temperate diversified *Pooideae* and *Danthonioideae* (Kellogg, 2015). However, most transitions of these grasses to colder climates occurred in tropical mountains, where cold-adapted lineages converged with temperate migrants (Schubert *et al.*, 2020). Each PACMAD genus exhibits distinct evolutionary trajectories, polyploid histories, and biogeographic origins, having independently evolved robust cold and freezing tolerance over the past 20-30 million years (Gallaher *et al.*, 2022; Stitzer *et al.*, 2025).

Plant exposure to cold stress, including chilling ($>0^{\circ}\text{C}$) and freezing ($<0^{\circ}\text{C}$) temperatures, triggers physiological and molecular responses. Chilling stress disrupts membrane fluidity, impairs enzyme activity, and inhibits photosynthesis (Aroca *et al.*, 2012), whereas freezing stress induces ice crystal formation in extracellular spaces, leading to cellular dehydration, membrane damage, and ultimately, cell death (Pearce, 2001). Many temperate plants undergo cold acclimation. Adaptations resulting from cold acclimation include membrane lipid modifications (Degenkolbe *et al.*, 2012; Barnes *et al.*, 2016), accumulation of compatible solutes (Koster & Lynch, 1992), enhanced antioxidant enzyme activity (Suzuki & Mittler, 2006) and protective protein accumulation (Puhakainen *et al.*, 2004; Xu *et al.*, 2022). Plant cold perception involves plasma membrane modifications, including fluidity changes, Ca^{2+} channel activation, and cytosolic acidification, driving lipid and transcriptomic responses (Mori *et al.*, 2018; Barnes *et al.*, 2023). A widely conserved mechanism underpinning freezing tolerance is the activation of the C-repeat binding factor/dehydration-responsive element-binding (CBF/DREB) signaling pathway, which induces expression of cold-responsive (COR) genes (Thomashow, 1999; Nakashima & Yamaguchi-Shinozaki, 2006).

Studies in PACMAD grasses reveal both broadly conserved and species-specific molecular adaptations associated with this pathway. *Panicum virgatum* utilizes a zinc finger transcription factor, PvC3H72, to enhance ICE1-CBF-COR signaling, significantly increasing cold resilience (Xie *et al.*, 2019). Additionally, calcium signaling pathways involving calmodulin and MEKK1 provide early response to cold stress (Ranaweera *et al.*, 2022; Wu *et al.*, 2024). *Pennisetum giganteum* maintains membrane fluidity at low temperature through the accumulation of unsaturated fatty acids such as linoleic and α -linolenic acids (Zhang *et al.*, 2017; Li *et al.*, 2020). Protective proteins including late embryogenesis abundant (LEA) proteins, dehydrins, and antioxidant enzymes are accumulated in response to freezing

conditions, reducing oxidative injury and cellular dehydration (Liu *et al.*, 2016; Xie *et al.*, 2019). Species-specific adaptations include enhanced proline biosynthesis via the ornithine pathway in *Sorghum bicolor* (Vera-Hernández *et al.*, 2018) and elevated activity of phenylpropanoid pathway enzymes (PAL and CAD) in *Miscanthus* to protect cells during freezing stress (Domon *et al.*, 2013).

Previous studies in PACMAD grasses have focused mainly on transcriptomic and metabolomic adaptations. However, proteomic responses remain largely unexplored. As the functional products of genes, proteins integrate transcriptional, translational, and post-translational regulation. Given the persistent nature of cold stress, proteomic analysis uniquely captures the accumulation of proteins reflecting acclimation and sustained adaptive responses. Furthermore, (Cope *et al.*, 2025) demonstrated that proteins face stronger evolutionary constraints than transcripts, with protein abundances being highly conserved across species while mRNA levels can vary without substantially affecting protein output. This suggests that shared cold adaptive responses may be more readily detected at the protein level than at the transcript level, making proteomics essential for understanding adaptive processes such as those underlying cold or freezing tolerance. Our multi-species study examines rhizome protein accumulation across seasonal variation to identify conserved freezing-tolerance mechanisms in PACMAD grasses, testing whether independent evolution of freezing tolerance proceeded through shared or lineage-specific molecular mechanisms.

Materials and Methods

To investigate proteomic responses to seasonal changes across prairie grasses, we conducted a comparative analysis of rhizomes from five species under winter (January, with lows reaching -29°C) and summer (August, with highs of 26°C) conditions. The species analyzed

were *Tripsacum*, *Andropogon gerardii*, *Miscanthus giganteus*, *Panicum virgatum*, and *Sorghastrum nutans* (Figure 1A).

Plant material and environmental conditions

Plants were grown outdoors at two field sites in Ithaca, New York (USDA Hardiness Zone 6a), where they experienced natural seasonal fluctuations, with an average winter temperature of -5°C (lows reaching -29°C) and summer temperatures averaging 20°C (highs of 26°C) over the five years preceding sampling (Northeast Regional Climate Center, 2025). *Tripsacum* hybrids were cultivated at Cornell University Caldwell Field, while *Andropogon gerardii*, *Miscanthus giganteus*, *Panicum virgatum*, and *Sorghastrum nutans* were grown at the Cornell Botanic Gardens. Environmental data were obtained from the Northeast Regional Climate Center (Station USC00304174) and the New York State Mesonet (Station GROT).

Rhizome tissue was collected during two sampling campaigns in 2019 and 2022, timed to capture distinct physiological states: winter dormancy and summer active growth. In 2019, summer samples were collected on August 11 following 121 mm of precipitation in the preceding 30 days, and mean soil temperature at 5 cm depth was 19.9°C. Winter samples were collected on January 8, 2020, after 22 days with soil temperature below 2°C. In 2022, summer sampling occurred on August 15 following 93 mm of precipitation; 30-day mean soil temperature was 20.6°C. Winter sampling on December 9 followed 20 days with soil temperature below 5°C. At summer sampling, plants displayed active vegetative growth with no visible drought stress; winter samples were dormant with hardened exteriors (Supplemental Figure S7).

Sampling and experimental design

Our study was conducted in two phases. Initial sampling of *Tripsacum* hybrids in 2019 provided evidence of conserved winter protein accumulation, prompting expansion in 2022 to

both replicated *Tripsacum* sampling and four additional PACMAD species with independent evolutionary histories of cold adaptation.

Rhizome tissue (minimum 5 g fresh weight) was collected from the internal mid-sections of rhizomes using sterile pruning scissors. Sampling was standardized to exclude apical tips, fine roots, and necrotic zones to ensure reproducibility and tissue comparability across species (Figure 1C and Supplemental Figure S7). Samples were immediately placed in pre-labeled 15 ml tubes, transported on dry ice, and stored at -80°C within 4 hours of collection. Biological replicates were defined as tissue collected from independent clonal plants. Sample sizes and genotypes are detailed in Table 1. For *A. gerardii* ('Sentinel'), *M. giganteus*, and *P. virgatum* ('Dust Devil'), three biological replicates were sampled per season from established accessions at Cornell Botanic Gardens. *S. nutans* ('Golden Sunset') was represented by two biological replicates per season due to winter mortality of one clonal plant.

Tripsacum hybrid population

Tripsacum rhizomes were sampled from seven-year-old F_1 hybrid progenies that had survived multiple freezing winters at Cornell University's Caldwell Field. The hybrids were derived from controlled crosses between northern founders collected from Iowa, Kansas, Illinois, and Missouri, where winters experience heavy frost, and southern founders collected from Florida and Texas, with subtropical climates. Most crosses were interspecific, between *T. dactyloides* subsp. *dactyloides* (northern) and *T. floridanum* (southern); additional crosses were intraspecific, between northern and southern populations of *T. dactyloides* (Table 1). In 2019, we sampled eight individuals per season representing six distinct hybrid genotypes from five cross combinations, prioritizing coverage of genetic diversity across the hybrid population. In

2022, we sampled 24 individuals per season comprising eight hybrid genotypes with three clonal replicates each, balancing genetic diversity with robust statistical replication.

F₂ Seedlings for RNA sequencing

F₂ seeds used for seedling freezing tolerance screening and RNA sequencing were generated from the same *Tripsacum* hybrids described above. Due to the limited seed set achieved through self-pollination (~500 seeds over the entire season), F₁ plants were allowed to open-pollinate. Seeds were harvested from individual maternal plants, germinated in trays, and grouped into F₂ families based on maternal lineage. Six-week-old F₂ seedlings sharing the same maternal F₁ parent were selected as experimental material for freezing assays and RNA extraction.

Proteomics data

Proteomic profiling was conducted using tandem mass tag (TMTpro) labeling and Orbitrap Eclipse Real Time Search (RTS)-SPS-MS3 technology. The experimental design included samples collected in both winter (January) and summer (August), with each species represented by at least three biological replicates with a larger number of genotypes analyzed for *Tripsacum* across two years (see Table 1 and Figure 1D).

Protein Extraction and Digestion

Fresh rhizome tissue was ground in liquid nitrogen. The ground sample was suspended in a lysis buffer, composed of 1× PBS (pH 8), 10 mM EDTA, 900 mM sucrose, and 0.4% β-mercaptoethanol. The suspension was vortexed and incubated on ice, then mixed with an equal volume of phenol buffered to pH 8 with Tris-EDTA (TE). After centrifugation and phase separation, the phenol phase containing proteins was collected. Proteins were precipitated from the phenol phase using ice-cold ammonium acetate /methanol solution (100

mM w/v ammonium acetate in 100% methanol) with protease and phosphatase inhibitors. After precipitation the pellets were washed five times with three different solutions and frozen. Then protein pellets were dissolved in 8-M urea buffer. Protein concentration was determined using a BCA protein assay. A set amount of each protein sample underwent S-Trap digestion, including treatment with dithiothreitol and iodoacetamide. The sample underwent multiple rounds of centrifugation with the addition of 8-M urea, followed by 50-mM ammonium bicarbonate (ABC) or triethylammonium bicarbonate (TEAB). The sample was digested using a trypsin buffer, centrifuged, and then dried by vacuum freeze-drying for storage prior to subsequent TMT labeling.

TMT Labeling and LC–MS/MS Analysis

Digested peptide mixtures were labeled using TMTpro reagents. Due to the multi-year nature of the study, different TMT plex levels were used: TMTpro 16-plex for 2019 *Tripsacum* samples and TMTpro 18-plex for 2022 samples. The 2022 samples were processed across seven separate TMTpro 18-plex batches: four batches for *Tripsacum* and three batches for the remaining four species.

Labeled peptides were pre fractionated using an Ultimate 3000 instrument. Samples were injected into nanoLC tandem Orbitrap Eclipse by a RTS-SPS-MS³ method (Fu *et al.*, 2021). Raw MS data were processed in Proteome Discoverer 2.5 using the Sequest HT search engine. Spectra were searched against the UniProtKB database supplemented with common contaminants. Searches allowed up to two missed cleavages and included fixed modifications of TMTpro tags on peptide N-termini and lysine residues, and carbamidomethylation of cysteine. Dynamic modifications included methionine oxidation and asparagine/glutamine deamidation. Precursor and fragment mass tolerances were set to 20 ppm and 0.5 Da, respectively.

Peptide spectral matches were filtered using Percolator with a 1% false discovery rate (FDR) at the peptide level. Proteins were initially identified with a minimum of 1 unique peptide and subsequently filtered to retain only those supported by at least 2 distinct peptides for downstream analysis. Protein quantification was based on the summed reporter ion intensities of unique and razor peptides, filtered with a minimum signal-to-noise ratio of 10, SPS mass matches $\geq 65\%$, and a co-isolation threshold of 50%. No minimum channel occupancy threshold was applied, allowing quantification of proteins detected in subsets of samples. Total peptide amount normalization was applied across TMT channels. Given the inherent biological variability of a multi-species field study, we applied a stringent threshold for differential abundance. Differentially abundant proteins (DAPs) were determined based on $|\log_2 \text{ fold change}| \geq 1$ and adjusted P-value < 0.05 , to prioritize high-confidence candidates with robust seasonal responses over minor fluctuations that may reflect transient environmental noise rather than conserved adaptive strategies. This approach helps address the statistical power imbalance between the deeply sampled *Tripsacum* dataset ($n \approx 32$) and the other grass lineages ($n=3$). While the higher power in *Tripsacum* inherently yields a larger list of statistically significant changes, our comparative analysis defines conservation based on the intersection of species. Consequently, the identification of conserved proteins is constrained by the more limited statistical power of the smaller datasets, suggesting that the core response network we report represents a high-confidence set of shared adaptations rather than an exhaustive catalog.

Proteomics Fractionation

All samples underwent high-pH reversed-phase fractionation prior to LC-MS/MS analysis. The fractionation protocol collected 48 fractions which were concatenated into 10 pools using a meandering pooling scheme for analysis.

Peptide Mapping and Reference Proteome Assignment

Due to multiplexing constraints, species with <15% predicted proteome divergence, such as *Andropogon gerardii* and *Miscanthus giganteus*, were pooled in the same TMT batch and searched against the *A. gerardii* proteome. To improve protein recovery for *M. giganteus*, peptides were re-mapped to the *M. sinensis* proteome using DIAMOND BLASTp in “ultrasensitive” mode. Proteins were reassigned to their best-matching *M. sinensis* counterparts based on aggregated peptide evidence (see Supplemental Code S1–S3; [GitHub repository](#)).

All final analyses were conducted using species-specific reference proteomes: *Andropogon gerardii* (Ag-CAM1351-DRAFT-PanAnd-1.0), *Miscanthus giganteus* (Msinensis_497_v7.0), *Panicum virgatum* (Pvirgatumvar_WBCHAP1_778_v1.0), *Sorghastrum nutans* (Sn-CAM1369-DRAFT-PanAnd-1.0), and *Tripsacum dactyloides* (Td-FL_9056069_6-DRAFT-PanAnd-1.0); genome assemblies and annotations are listed in the Data Availability section.

Protein abundance normalization and statistical analysis

Protein intensities were normalized using the "total peptide amount" method in Proteome Discoverer 2.5, ensuring comparability across TMT channels. Differential protein abundance between winter and summer was assessed using one-way ANOVA, followed by Tukey's HSD post hoc test. For the multi-batch *Tripsacum* 2022 dataset, statistical testing was performed within each batch and P-values were combined using Fisher's method. To ensure high confidence in quantification, proteins in the *Tripsacum* 2022 dataset were only retained for downstream analysis if they were successfully quantified in at least 3 of the 4 TMT batches. Multiple testing correction was performed using the Benjamini-Hochberg procedure. For differential abundance analysis, only proteins quantified in both winter and summer samples

within each species were retained. Proteins were considered differentially abundant if $|\log_2 \text{fold change}| \geq 1$ and adjusted P-value < 0.05 .

To assess technical reproducibility and identify sources of variation, principal component analysis (PCA) was performed on $\log_2$ -transformed protein abundances (Supplemental Table S1) using the FactoMineR::PCA function in R with centering and scaling. For cross-species comparison at the orthogroup level, protein families among the five species were identified as shared using OrthoFinder (Emms & Kelly, 2019). This analysis identified a total of 51,442 orthogroups across the five proteomes. For the comparative proteomics analysis, 18,454 of the 20,737 quantified proteins were successfully assigned to 4,888 orthogroups. The remaining 2,283 proteins lacked clear orthologous assignment and were excluded from cross-species comparisons. In cases where multiple proteins belonged to the same orthogroup within a species, $\log_2$ fold-change values were averaged to generate a single orthogroup-level fold change.

PCA was performed on orthogroups detected in $>50\%$ of samples within each species. For *Tripsacum*, batch effects were removed by extracting residuals from a linear model of abundance on batch prior to PCA. For the multi-year *Tripsacum* dataset, batch and year effects were evaluated alongside seasonal variation; because season emerged as the dominant factor (Supplemental Figure S1B), all *Tripsacum* samples were analyzed as a pooled species group. All statistical analyses and visualization were performed in R version 4.4.3, using RStudio 2024.12.1+563 ("Kousa Dogwood" release).

Cross-species correlation analysis

To assess evolutionary constraint on cold-responsive proteins, we compared cross-species conservation of seasonal response magnitude between shared cold-responsive orthogroups and background proteins, following the framework of Cope et al. (2025).

For each orthogroup quantified in at least four of five species, we calculated pairwise Spearman correlations of $\log_2$ fold-change (winter/summer) across all species pairs. Orthogroups were classified as "shared DAPs" if differentially abundant ($|\log_2\text{FC}| \geq 1$, adjusted $P < 0.05$) in two or more species, or as "background" if differentially abundant in at most one species. Distributions of pairwise correlations were compared between shared DAPs and background using the Wilcoxon rank-sum test. The same approach was applied to basal protein abundance (summer and winter separately) to distinguish constraint on response dynamics from constraint on absolute abundance levels.

Functional annotation and GO enrichment

Orthogroups were assigned using OrthoFinder (v2.5.5, (Emms & Kelly, 2019), clustering protein sequences from all five species based on sequence similarity and inferred gene trees. DIAMOND (v2.1.9, (Buchfink *et al.*, 2021) was used for all-vs-all sequence comparisons. Orthogroups served as the basis for cross-species comparisons and functional propagation of annotations, particularly for proteins with limited annotation coverage outside *Tripsacum dactyloides*.

Functional annotation was performed separately for DAPs and non-significant proteins. DAP sequences were first annotated using DIAMOND BLASTp against the SwissProt database. Protein descriptions and Gene Ontology (GO) terms were extracted from UniProt using a custom script. For DAPs lacking SwissProt hits, functional predictions were generated using PANNZER2 (Törönen & Holm, 2022), run independently for each species. In contrast, for non-significant proteins, functional annotations were propagated from *Tripsacum dactyloides* to orthologous proteins within the same orthogroup when available. Remaining unannotated entries were manually curated based on sequence similarity and conserved domain features.

GO term enrichment analysis was performed using the topGO R package (v2.50.0), applying the “elim” algorithm with Fisher’s exact test to reduce redundancy and account for GO hierarchy. The background set included all proteins quantified across the experiment. Enrichment was conducted separately for proteins with elevated abundance and reduced abundance in winter. Terms with a minimum node size of 10 and a P-value < 0.01 were considered significant. The top 15 enriched Biological Process (BP) terms in each group were visualized using dot plots in ggplot2, incorporating log-transformed P-values, gene ratios, and counts of significant proteins.

Candidate protein exploration methods

(Barnes *et al.*, 2023) showed that cold-responsive transcriptional and proteomic changes often overlap with broader stress mechanisms involving damage, osmotic imbalance, and acidification. To improve biological resolution, we curated DAPs into nine functional categories based on known roles in freezing tolerance (Örvar *et al.*, 2000; Iwaya-Inoue *et al.*, 2018; Schubert *et al.*, 2020) : (1) protein aggregation and membrane stability, (2) protein folding and chaperone activity, (3) metabolism and osmoregulation, (4) detoxification and ROS scavenging, (5) lipid metabolism, (6) antifreeze and cell wall modification, (7) cold signal transduction, (8) cytoskeletal organization, and (9) other/uncategorized. Assignments were informed by protein descriptions, GO terms, PANNZER2 predictions, and manual review of domain features and relevant literature. This categorical framework enabled a clearer visualization of conserved and divergent cold responses across species (Supplemental Figure S5), particularly by consolidating functionally relevant proteins that may be dispersed across non-specific GO categories.

Selecting proteins for traits on trees

Proteins for phylogenetic trait mapping were identified by text-mining functional annotations produced by PANNZER2 for both the *Tripsacum dactyloides* proteome and the DAP sets from all five species, searching specifically for GO terms associated with late embryogenesis abundant (LEA) proteins, heat shock proteins (HSPs), and elongation factor 1 (EF1). Extracted orthogroup identifiers from our $\log_2$ fold-change matrix were then used to retrieve the corresponding protein IDs, which served as input for sequence extraction. Full-length amino-acid sequences were obtained from each species' reference proteome using seqkit grep (v2.0) with protein headers matching these IDs. Multiple sequence alignments were generated in MAFFT (v7.475) under default settings, and maximum-likelihood and maximum-likelihood trees were inferred using raxmlHPC (RAxML-8.2.13) PROTGAMEJTT model with 100 rapid bootstraps. Each resulting tree was rooted on a biologically appropriate outgroup (e.g., a chloroplastic class I small HSP) and visualized in R (v4.1.0) using the ggtree package, with $\log_2$ FC values mapped as a color gradient onto tip points.

RNA-seq processing and differential expression analysis

Transcriptome data for maize chilling responses across tissues (leaf, root, top-crown, and bottom-crown) were derived from (Xue *et al.*, 2021) using 3' RNA-seq (5C vs. 22C). For *Tripsacum*, we analyzed leaf transcriptome data from a segregating freezing-tolerant bulk of F₂ seedlings derived from the same *Tripsacum* hybrids used in our rhizome analysis. Plants from seven F₂ families were bulked as freezing-tolerant or intolerant based on survival after exposure to -8°C for three hours. Within the freezing-tolerant bulk only, RNA was collected from 6-week-old seedlings at control temperature (22°C) and after 7 days of cold acclimation (5°C), with expression compared between these conditions (tol_5C vs. tol_22C). The second

transcriptome from *Tripsacum* 'Pete' (leaf, root, top-crown, and bottom-crown) was generated using the same methodology (5C vs. 22C) as in (Xue *et al.*, 2021) and is currently unpublished.

F₂ seedling leaf RNA was extracted using the methods described in (Gault *et al.*, 2018). Paired-end RNA-seq reads were quality-filtered and trimmed using Trimmomatic (version 0.32, (Bolger *et al.*, 2014) to remove Illumina adapter bases discarding bases from beginning and end with a phred score <35 or <20, respectively. Reads with a minimum length of 36 bp were retained. Reads were aligned to the *Tripsacum dactyloides* genome (Td-FL_9056069_6-DRAFT-PanAnd-1.0.fasta) and annotation (Td-FL_9056069_6-DRAFT-PanAnd-1.0_final.gff3) using STAR v2.7.10b (`--twopassMode Basic`), allowing each read to map to up to 20 locations (`--outFilterMultimapNmax 20`). Gene-level counts were extracted from uniquely mapped reads using featureCounts v2.0.3 with default settings. Raw counts were normalized using the estimateSizeFactors method in DESeq2, which adjusts for sequencing depth and RNA composition bias. Differential gene expression analysis was conducted using DESeq2 (Love *et al.*, 2014). Contrasts were made between 7 °C and 22 °C within each genotype (eight F₂ families) with 5 °C as the reference, modeling both temperature and genotype as main effects ($\sim$ genotype + temperature). Genes were considered differentially expressed if they passed an adjusted P-value cutoff of 0.05 (Benjamini-Hochberg correction) and showed a minimum absolute log₂ fold change > 1.

Structural and Biophysical Analysis of LEA3 Proteins

Amino acid sequences of LEA3 orthologs from *Tripsacum dactyloides* (Td00001aa022594_T002) and *Zea mays* (Zm00001eb294480) were aligned using MAFFT v7.475 with default settings. Predicted three-dimensional structures were generated using

AlphaFold3 (Abramson *et al.*, 2024). LEA3 proteins are classified as intrinsically disordered proteins that are thought to adopt amphipathic alpha-helical conformations upon binding to membranes or other proteins during cellular dehydration (Kovacs *et al.*, 2008). Structures were predicted using default parameters with full-length protein sequences as input. Note that predicted structures may differ from AlphaFold2 models deposited in UniProt due to differences in the AlphaFold3 architecture. Hydrophobicity was visualized in PyMOL (v2.5) using a custom script based on the Eisenberg scale, which colored residues according to side-chain properties. Hydropathy plots were generated using both ProtScale (ExPASy) and AmphipaSeek (Sapay *et al.*, 2006) , with a sliding window size of 11 amino acids corresponding to the LEA3 motif length. Hydrophobic peaks associated with conserved LEA motifs were annotated across both sequence alignments and structural models.

Results

To examine seasonal proteomic responses in PACMAD grasses, we analyzed rhizome tissues of five species using a proteomic shotgun approach during the winter (January) and summer (August) of 2019 and 2022 (Figure 1) in Ithaca, New York (USDA Hardiness Zone 6a). The exposure to deep-freezing winters ensured that the proteomic responses captured were reflective of natural seasonal variation (Figure 1B).




Figure 1. Overview of sampling and proteomics workflow. (A) Rhizomes from the five species sampled in the study: *Tripsacum* spp, *Panicum virgatum*, *Andropogon gerardii*, *Miscanthus giganteus*, and *Sorghastrum nutans*. (B) Field conditions during the two sampling seasons, illustrating the active growth in August and dormancy in January. (C) Example of the rhizome tissue sampled for proteomics, shown here for *Tripsacum*. The red arrow highlights the specific region collected. (D) Schematic overview of the proteomics workflow. Rhizome samples were collected each season, followed by protein extraction, digestion, and TMTpro labeling. Peptides were fractionated and analyzed via RTS-SPS-MS³ using an Orbitrap Eclipse mass spectrometer. Each species' reference proteome was used for peptide and protein assignment, enabling quantification and identification. (E)

Phylogeny of species presented along with *Z. mays* for reference; snowflakes indicate freezing tolerance.

Protein abundance analysis was performed in batches (see Methods). Across all five species a total of 20,737 proteins were quantified, but for differential accumulation across seasons we included only those quantified in both winter and summer samples (Table 1). PCA of 745 shared orthogroups confirmed clustering by species and season; batch-corrected *Tripsacum* samples (1,889 proteins) clustered by season regardless of hybrid background (Supplemental Figure S1). The number of identified proteins ranged from 2,372 to 4,451 per species. Differentially abundant proteins (DAPs) varied across species, with 58–174 proteins significantly increased and 56–218 decreased in abundance under freezing conditions (Table 1). Although fewer proteins were identified in *Sorghastrum nutans*, reflecting its draft genome status, robust seasonal signatures were detectable across all five lineages (Supplemental Figure S2).

Cross-Species comparative proteomic analysis reveals conserved seasonal protein accumulation patterns

We restricted our comparative analysis to the 18,454 proteins assigned to 4,888 orthogroups identified across species (Supplemental Table S2). Seasonal fold-changes were positively correlated across all species pairs ($\rho = 0.37\text{--}0.69$; Figure 2A), with specific orthogroups—including LEA3, GST, HSP1, and PEBP—consistently identified as differentially abundant. To test whether this conservation reflects evolutionary constraint on cold response, we compared fold-change correlations between shared DAPs (differential abundance in ≥ 2 species; $n = 135$ orthogroups) and background proteins (≤ 1 species; $n = 1,548$). Shared DAPs showed substantially higher correlation in seasonal response magnitude ($\rho = 0.80$) than background ($\rho = 0.45$; $P = 1.1 \times 10^{-5}$; Figure 2B). Notably, this constraint did not extend to basal abundance, where shared DAPs showed slightly *lower* correlation than

background ($\rho = 0.62$ vs 0.68 ; $P = 0.004$). This pattern—tight constraint on response dynamics but flexibility in baseline levels—suggests that independent cold adaptations relied on parallel regulation of a core set of protective proteins.




Figure 2. Conservation of seasonal protein accumulation across independently cold-adapted PACMAD grasses. (A) Pairwise comparisons of $\log_2$ fold-change ($\log_2$ FC winter/summer) in differentially accumulated orthogroups across species. Scatterplots display the correlation of orthogroup-level $\log_2$ FC values between species, with shared DAPs elevated in winter shown in blue, reduced in red, and non-significant proteins in gray. Each panel represents a comparison between two species, with the number of shared orthogroups (n) and Spearman's correlation coefficient (ρ) displayed. (B) Pairwise Spearman correlations for orthogroups quantified in at least four of five species, comparing shared DAPs (red; differentially abundant in ≥ 2 species) versus background orthogroups (gray; differentially abundant in ≤ 1 species). Correlations are shown for summer abundance, winter abundance, and seasonal response magnitude ($\log_2$ FC winter/summer). Wilcoxon rank-sum test: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Ortholog and pathway sharing across species

We next examined orthogroup sharing across species (Supplemental Figure S3). Winter-elevated orthogroups showed greater conservation than winter-reduced: 27% of 330 winter-elevated orthogroups were shared across at least two species versus 21% of 470 winter-reduced orthogroups. Variation in identified proteins partly reflects reference proteome completeness—*S. nutans*, with a draft genome containing 26% ambiguous bases, yielded fewer identifications (Supplemental Table S12).

GO enrichment analysis revealed distinct functional signatures (Supplemental Figure S4). Winter-elevated proteins were enriched for protein folding, stress response, cold acclimation, and ROS scavenging (Supplemental Table S4). Winter-reduced proteins were enriched for microtubule-based processes, mitotic cell cycle, cytoskeleton organization, and cell wall biogenesis—processes associated with active growth (Supplemental Table S5). The greater conservation of winter-elevated orthogroups suggests that protective responses to freezing are more constrained than growth-related processes active during summer. To identify the most conserved functional themes, we examined the 50 most winter-elevated proteins per species (Figure 3; Supplemental Table S7). LEAs and heat shock proteins (HSPs) were present in at least four of five species. A chi-square test on standard GO categories revealed significant species heterogeneity ($\chi^2[70] = 97.5$, $P = 0.017$), but GO terms may conflate cold-specific processes with broader stress responses. (Barnes *et al.*, 2023). Using curated cold-related categories (see Methods), 63% of proteins mapped to one of nine functional groups (Supplemental Figure S7). Heterogeneity remained significant ($\chi^2[40] = 58.1$, $P = 0.032$; Figure 3A), indicating that while species share the same core functional categories, they differ in relative investment. Detoxification and ROS scavenging proteins were represented in all five species. Metabolism/osmoregulation proteins were less prominent in *A. gerardii*,

while protein aggregation and membrane stability proteins were enriched in *A. gerardii* and *P. virgatum*. These patterns indicate a shared functional core with lineage-specific variation in relative allocation (Figure 3A), particularly in metabolism/osmoregulation and protein folding/stability.

LEA3 (OG0022470) was the only ortholog ranked among the top 10 winter-elevated proteins in all five species, and was the single most elevated protein in *Tripsacum* ($\log_2\text{FC} = 5.0$; Figure 3B; Supplemental Table S7). Other functional categories showed species-specific representation: HSPs appeared in the top 10 for *Tripsacum* (HSP22, Class IV) and *S. nutans* (HSP23.2) but involved different family members; ROS scavenging was universal but employed glutathione transferases in four species and peroxidases in *A. gerardii*; putative antifreeze proteins (chitinases) ranked highly in *S. nutans* and *A. gerardii*. This pattern—a single conserved ortholog (LEA3) alongside functionally equivalent but distinct proteins in other categories—exemplifies parallel recruitment of LEA3 alongside functional convergence for chaperone and antioxidant functions.



Figure 3. Overview of cold-related functional categories represented among the top 50 most winter-enriched rhizome proteins across five PACMAD grasses. (A) Mosaic plot of cold-related

functional categories across species. This plot visualizes the distribution of proteins among cold-related functional categories for each species. The relative abundance of each category within a species is shown, highlighting distinct functional profiles and species-specific variations in cold tolerance mechanisms. A Chi-square test for independence ($\chi^2[40] = 58.1$, $P = 0.032$) was performed to evaluate variation across species. **(B)** Visual representation of cold-related functional categories in rhizome proteins of five *Andropogoneae* species. This figure illustrates the top 50 most abundant proteins within the rhizomes of five *Andropogoneae* species. Proteins are classified into functional categories relevant to cold-related mechanisms, such as metabolism/osmoregulation, cold signal transduction, membrane stability, lipid metabolism, and others. Color-coding facilitates a comparative visualization of cold tolerance mechanisms across species. Antifreeze proteins are marked by an asterisk, as their classification is based solely on protein homology.

Evolutionary patterns in accumulated protein families

Phylogenetic analysis revealed distinct evolutionary patterns between LEAs and HSP families (Figure 4). The control elongation factor 1 protein (EF1) family, previously shown to have stable expression at transcript and protein levels across conditions (Takamori *et al.*, 2017), Figure 4A), showed minimal seasonal accumulation and no phylogenetic clustering of abundance level patterns, consistent with their housekeeping function (Supplemental Table S8). HSPs showed greater diversity (212 proteins across 37 orthogroups, Supplemental Table S9) compared to LEAs (66 proteins in 20 orthogroups, Supplemental Table S10), with control EF1 comprising 33 proteins in 4 orthogroups. Both LEA and HSP phylogeny (Figures 4B and C) resolved into clear functional sub-families, but among HSPs the most strongly winter-elevated members ($\log_2FC > 2.5$) belonged to different clades across different species, with only 32% of HSP proteins significantly elevated in winter. In contrast, LEA proteins (Figure 4B) demonstrated more consistent winter abundance (47% of the proteins), particularly within the LEA3 subfamily, which displayed elevated abundance across all species despite having diverged earlier than other LEA clades.




Figure 4. Phylogenetic analysis and seasonal abundance dynamics of selected protein families across five PACMAD grasses (*Tripsacum* spp., *P. virgatum*, *A. gerardii*, *M. giganteus*, and *S. nutans*). Protein identifiers represent species-specific sequences, and brackets denote functional group annotations, merging adjacent orthogroups with similar functions for clarity. Heat maps illustrate log₂

fold-change ($\log_2FC$) in protein abundance between winter and summer conditions, where blue indicates higher abundance in winter, and red signifies higher summer abundance. Panels show distinct protein families: **(A)** translation elongation factor 1 proteins (EF1; control group), **(B)** late embryogenesis abundant (LEA) proteins, and **(C)** heat shock proteins (HSPs). Asterisks (*) mark proteins exhibiting statistically significant differential accumulation ($P < 0.05$).

*Beyond rhizomes: Comparison with maize & *Tripsacum* mRNA*

We next tested whether rhizome protein signatures corresponded to mRNA responses in other tissues. Cold-induced mRNA in F₂ seedling leaves of *Tripsacum*, derived from the same hybrids used in our rhizome study, showed weak but significant correlation with rhizome protein fold-changes ($\rho = 0.29$, $P < 0.001$; Supplemental Figure S5A). Other *Tripsacum* tissues showed weaker correlations ($\rho = 0.08$ – 0.16 ; Supplemental Figure S5B–E), as did maize orthologs ($\rho = 0.01$ – 0.09 ; Supplemental Figure S5F–I). Despite these weak overall correlations, LEA3 displayed strong cold induction in leaves of both species ($\log_2FC = 8$ in *Tripsacum*, and 7 in maize; Supplemental Figure S5A, F). That maize induces LEA3 transcription to levels comparable to cold-tolerant *Tripsacum*, yet remains freezing-sensitive, suggests that transcriptional induction alone does not confer protection.

Structural and Functional Insights into LEA3: A Key Player in Freezing Tolerance

LEA3 (OG0022470) was the only ortholog consistently elevated across all five species, ranking as the top DAP in *Tripsacum* rhizomes ($\log_2FC = 5.0$) and the most cold-induced transcript in leaves ($\log_2FC = 8$). Sequence alignment of the *Tripsacum* and *Zea mays* orthologs revealed a specific divergence in the maize protein affecting the periodicity of the conserved motifs and hydropathy profiles (Figure 5). LEA3 proteins are defined by a repeating 11-mer motif described by Dure as essential for desiccation and freezing tolerance (Dure, 1993). These motifs feature apolar residues at positions 1, 2, 5, and 9, forming the hydrophobic face of an amphipathic α -helix critical for stabilizing cellular membranes and preventing protein aggregation. Positions 6, 7, and 8 exhibit either a "+ – +" or "+ Q +"

charge arrangement. *Tripsacum* LEA3 (Td00001aa022594) contains six consecutive 11-mer motifs adhering closely to Dure's rules (Figure 5A). These motifs align with the α -helical domains in the AlphaFold3-predicted structure (Figure 5B). Kyte-Doolittle hydropathy analysis shows regular oscillations with ~ 11 -residue periodicity, consistent with intact amphipathic helices (Figure 5C). The maize ortholog (Zm00001eb294480) contains only four motifs interrupted by a 15-amino-acid insertion that disrupts regular motif spacing, producing an irregular hydropathy profile (Figure 5C).

Quantitative comparison of the motif region (positions 40–101) revealed that mean hydropathy at Dure's hydrophobic positions (1, 2, 5, 9) averaged approximately -1.1 in *Tripsacum*, whereas the maize ortholog showed values 15–20% more hydrophilic at corresponding positions. The disruption is most pronounced in positions 70–80, where individual position differences exceed $|\Delta| = 1.0$, and in positions 87–101, where the Td–Zm hydropathy pattern correlation becomes strongly negative ($r = -0.81$, $n = 15$ positions), indicating that the hydrophobic and hydrophilic faces are misaligned relative to the conserved architecture (Supplemental Table S11). Hydropathy profiles from other cold-tolerant species—including other PACMAD grasses, wheat (*TdLEA3*; (Koubaa *et al.*, 2019), and *Arabidopsis* (*AtLEA7*; (Popova *et al.*, 2015)—show oscillating patterns similar to *Tripsacum*, not maize (Supplemental Figure S6). These structural differences suggest that transcriptional induction of LEA3 alone may not ensure freezing tolerance; functional protection likely depends on both sufficient abundance and intact motif architecture.




Figure 5. LEA3 protein sequence alignment, structure and hydrophobicity. (A) The alignment shows the LEA3 proteins from *T. dactyloides* (Td00001aa022594_T002) and, *Z. mays* (Zm00001eb294480), with the repeating 11-mer LEA3 motifs highlighted and labeled numerically. The dashed motif deviates from Dure's LEA3 motifs in the first position. (B) Predicted secondary structure of LEA3 proteins using AlphaFold3, showing the conserved alpha-helical domains across *T. dactyloides* (top) and *Z. mays* (bottom). LEA motif regions corresponding to labeled sequence motifs (e.g., 1, 2–6) are boxed. Red and blue indicate regions of high hydrophobicity and hydrophilicity, respectively. (C) Kyte-Doolittle hydropathy plots for LEA3 proteins. The X-axis represents residue position, while the Y-axis indicates hydropathicity scores (positive values for hydrophobic regions, negative for hydrophilic regions). 11-mer motif regions are highlighted with red boxes, corresponding to key domains in (B).

Discussion

Our analysis of rhizome proteomes from five independently cold-adapted PACMAD grasses reveals signatures of both parallel and convergent evolution in protein-level responses to freezing. Parallel evolution is evident in the repeated recruitment of the same orthologs—most strikingly LEA3, accumulated across all five lineages—while convergent evolution is reflected in the use of phylogenetically distinct HSP family members to achieve similar chaperone function. Parallel recruitment of LEA3 implies that the ancestral protein was already suited to cryoprotection and has been maintained, whereas convergent HSP

recruitment suggests that chaperone-mediated protection permits greater flexibility in which family members are deployed. To our knowledge, this study provides the first quantitative comparison of cross-species fold-change conservation at the protein level in a cold-stress context, complementing transcript-level comparisons in *Pooideae* (Schubert *et al.*, 2019) and desiccation-tolerance studies in resurrection grasses (Marks *et al.*, 2024).

The most striking finding is the evolutionary constraint on response magnitude. Among shared cold-responsive orthogroups, cross-species fold-change correlations were substantially higher ($\rho = 0.80$, Figure 2B) than among background proteins ($\rho = 0.45$; $P = 1.1 \times 10^{-5}$), indicating that when freezing tolerance evolved independently across PACMAD lineages, it proceeded through quantitatively similar responses in a core set of protective proteins. These patterns reflect cold acclimation rather than drought responses: summer-collected tissues showed vigorous growth with actively emerging rhizome buds (Supplemental Figure S9) after >90 mm cumulative precipitation, while winter samples were collected after sustained sub-zero exposure reaching -16°C , with soil temperatures remaining below $2\text{--}5^{\circ}\text{C}$ for 20–22 days; (Brotzge *et al.*, 2020). Our common-garden design ensured that all genotypes experienced identical winter conditions, and the successful overwintering of all sampled plants confirms that their proteomic configurations were sufficient to survive USDA Hardiness Zone 6a winters.

With these seasonal responses established as cold-driven, their conservation contrasts sharply with transcript-level patterns: comparative transcriptomics in *Pooideae* found generally low similarity in cold-responsive gene expression across species ($r = 0.09\text{--}0.40$; (Schubert *et al.*, 2019), with nearly half of all cold-responsive genes being species-specific. That protein-level conservation substantially exceeds transcript-level conservation is consistent with the framework of (Cope *et al.*, 2025), who demonstrated that protein abundances are under

stronger stabilizing selection than mRNA abundances over macroevolutionary timescales, with mutations affecting mRNA levels largely buffered before they alter protein output. If this buffering operates on cold-responsive genes as it does genome-wide, then transcript-level comparisons would systematically underestimate the degree of evolutionary conservation in stress responses. Notably, this constraint applies specifically to the dynamic response: basal abundance correlations were slightly lower among shared cold-responsive proteins than background ($\rho = 0.62$ vs 0.68 ; $P = 0.004$; Figure 2B), indicating that while independent lineages converged on similar fold-change magnitudes, they achieved these responses from different baseline levels. Shared cold-responsive proteins are more divergent at baseline than background proteins, implying that there may be no single optimal resting level, but only a conserved requirement for how much to accumulate when freezing arrives.

LEA3 provides the clearest illustration of this constraint and connects the evolutionary pattern to a mechanistic hypothesis. LEA proteins function as molecular shields during freezing through membrane stabilization, ion sequestration, and chaperone activity (Kovacs *et al.*, 2008; Cuevas-Velazquez *et al.*, 2017). These functions depend on the repeating 11-mer motif architecture described by (Dure, 1993), in which apolar residues form the hydrophobic face of amphipathic α -helices that bind membranes and partially unfolded proteins during dehydration and freezing (Koubaa *et al.*, 2019; Singh & Graether, 2021). Among cold-tolerant PACMAD grasses and outgroup species including wheat (Koubaa *et al.*, 2019) and *Arabidopsis* (Popova *et al.*, 2015), LEA3 proteins maintain this architecture, producing regular hydropathy oscillations with ~ 11 -residue periodicity (Supplemental Figure S6). The maize ortholog, by contrast, contains a 15-amino-acid insertion that disrupts motif spacing, producing a hydropathy profile in which the hydrophobic and hydrophilic faces are misaligned relative to the conserved architecture (Figure 5C; Supplemental Table S11). Because amphipathic helix function depends on a continuous hydrophobic face for membrane

association (Singh & Graether, 2021), this disruption may directly impair the capacity of *ZmLEA3* to stabilize membranes, protect enzymes or form oligomers during freezing.

While *ZmLEA3* is cold-inducible (Liu *et al.*, 2013) with transcription to levels comparable to cold-tolerant *Tripsacum* in leaves (Supplemental Figure S5F) yet remains freezing-sensitive underscores a key point: transcriptional induction alone does not ensure protection. Functional efficacy likely depends on both sufficient abundance and intact motif structure. This interpretation is consistent with observations that cold-regulated cereal LEA3 proteins require conserved C-terminal domains for oligomerization (NDong *et al.*, 2002) and that *Arabidopsis* LEA3 proteins form higher-order assemblies under stress (Ginsawaeng *et al.*, 2021; Romero-Pérez *et al.*, 2024). AlphaFold3 multimer predictions for *TdLEA3* and *ZmLEA3* are consistent with this, though with notable differences in packing geometry (Supplemental Figure S8).

More broadly, several of the most consistently accumulated proteins across PACMAD rhizomes have well-characterized orthologs implicated in cold acclimation in *Arabidopsis*, including LEA proteins, small HSPs, and glutathione S-transferases (Marrs, 1996; Sun *et al.*, 2002; Hundertmark & Hinch, 2008), indicating deep conservation of core protective pathways across monocots and dicots. However, some highly accumulated proteins in PACMAD rhizomes—notably PEBP family members and certain chitinase-like proteins—have not been prominently featured in *Arabidopsis* cold studies, potentially reflecting grass-specific or rhizome-specific adaptations (Roeder *et al.*, 2025). The presence of ABA-inducible proteins and dehydrins among the most accumulated winter proteins across species is consistent with the hypothesis that freezing tolerance in PACMAD grasses evolved partly by co-opting existing drought-tolerance pathways. This aligns with phylogenetic evidence that drought tolerance served as an evolutionary precursor to frost tolerance in

grasses (Schat *et al.*, 2025). The early divergence of LEA3 from other LEA subfamilies (Figure 4B) suggests that the molecular properties enabling its cryoprotective function may have been present before the radiation of PACMAD grasses, representing a pre-existing capacity subsequently maintained under constraint in cold-adapted lineages and degraded where freezing selection was absent, as in maize. Similar patterns of pre-adaptation followed by lineage-specific elaboration have been documented in resurrection grasses (Marks *et al.*, 2024).

In conclusion, independent evolution of rhizome freezing tolerance in PACMAD grasses is governed by a combination of parallel recruitment of conserved orthologs and functional convergence among diverse protein families, with quantitative constraint on response magnitude that is substantially stronger at the protein level than previously observed at the transcript level. The structural differences between *Tripsacum* and maize LEA3—canonical motif architecture maintained in a cold-adapted lineage versus disrupted architecture in a tropical relative—offer a candidate mechanism for enhancing freezing resilience in maize. Future work should prioritize functional validation of LEA3 structural variants to translate these evolutionary insights into breeding applications that could extend growing seasons in temperate regions.

Acknowledgments

We thank Cornell Botanic Gardens for their support of this study and for maintaining the field plots that provided essential plant material, particularly Kristine Boys, Emily Detrick (Elizabeth Weaver Director of Horticulture), and Todd Bittner (Director of Natural Areas). We acknowledge the Proteomics and Metabolomics Facility at the Cornell Institute of Biotechnology (RRID:SCR_021743) for providing mass spectrometry services and data analysis. We are grateful to Prof. Hening Lin for funding support from the HHMI

Transformative Technology 2019 program for the Orbitrap Eclipse system, and to Dr. Sheng Zhang for NIH SIG grant support (1S10 OD017992-01 for the Orbitrap Fusion mass spectrometer and 1S10 RR025449-01 for the Orbitrap Velos/Elite mass spectrometer). We extend our appreciation to Elizabeth T. Anderson and Robert W. Sherwood at the Proteomics and Metabolomics Facility for their technical expertise and assistance with sample processing and analysis. We thank the Genomics Facility (RRID:SCR_021727) of the Biotechnology Resource Center at the Cornell Institute of Biotechnology for their assistance with sequencing experiments. Additional computational resources and data management were provided by the Bioinformatics Facility (RRID:SCR_021757) at the Cornell Institute of Biotechnology. This research is made possible by the New York State (NYS) Mesonet. Original funding for the NYS Mesonet (NYSM) buildup was provided by the Federal Emergency Management Agency grant FEMA-4085-DR-NY. The continued operation and maintenance of the NYSM is supported by the National Mesonet Program, University at Albany, Federal and private grants, and others. We acknowledge all who have contributed to the preservation, cultivation and study of wild and domesticated grasses

Author Contributions

M.C.R., T.E.R., and E.O. conceived and designed the study. N.L. and T.E.R. developed the long-term sampling strategy that enabled seasonal comparisons. N.L., T.E.R., T.L., S.-K.H., C.M.G., and E.O. performed sample collection and experimental work. T.L., R.A., and Q.F. conducted proteomics sample preparation, LC-MS/MS, and data processing. C.M.G., J.H., T.L., and J.Z. performed RNA-seq analysis. J.Z., S.-K.H., and E.O. carried out bioinformatics and statistical analysis. J.Z., S.-K.H., C.O.H., and E.O. prepared the figures. E.O., J.Z., L.J.B., and E.S.B. wrote the manuscript. C.O.H., L.J.B., S.-K.H., J.H., Q.F., E.S.B., and

M.C.R. reviewed and edited the manuscript. E.O. and M.C.R. supervised the project and E.S.B. and M.C.R. acquired funding.

Funding

This research was supported by the National Science Foundation under Grant Number 1822330, and the U.S. Department of Agriculture–Agricultural Research Service (USDA-ARS) under Project Number 5030-21000-072-00-D (Corn Insects and Crop Genetics Research Unit, Ames, Iowa), Project Number 6066-21310-005-00-D (Genomics and Bioinformatics Research Unit, Stoneville, MS), Project Number 8062-21000-043-00-D (Plant, Soil and Nutrition Research Unit, Ithaca, NY), and the United States - Israel Binational Agricultural Research and Development Fund, Vaadia-BARD Postdoctoral Fellowship Award No. FI-628-2022. C.M.G. was supported by the National Science Foundation Postdoctoral Research Fellowship in Biology under Grant No. 1523861. Research in the lab of J.H. is supported by USDA NIFA (2022-67013-37040). Q.F. was supported by HHMI Transformative Technology 2019 program for the OrbitrapEclipse system.

Data Availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol *et al.*, 2025) partner repository with the dataset identifier PXD063668 (Sample key file in Supplemental Table S12). RNA-seq data are available under BioProject accession numbers PRJNA705456 (*Zea mays*) and PRJNA1260937 (*Tripsacum dactyloides*) in the NCBI BioProject database (<https://www.ncbi.nlm.nih.gov/bioproject/>). Genome assemblies and annotations used in this study are publicly available from MaizeGDB (Woodhouse *et al.*, 2021) at https://maizegdb.org/PanAnd_project for *Tripsacum dactyloides* (Td-FL_9056069_6-DRAFT-PanAnd-1.0), *Andropogon gerardii*

(Ag-CAM1351-DRAFT-PanAnd-1.0), and *Sorghastrum nutans* (Sn-CAM1369-DRAFT-PanAnd-1.0), and from the JGI Data Portal (Goodstein *et al.*, 2012) at <https://data.jgi.doe.gov/> for *Miscanthus giganteus* (Msinensis_497_v7.0) and *Panicum virgatum* (Pvirgatumvar_WBCHAP1_778_v1.0), along with their corresponding GFF annotation files. Functional annotations, data tables, and analysis scripts are available from GitHub at <https://github.com/elor77/PACMAD-Rhizome-Proteomics>.

References

- Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, *et al.* 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**: 493–500.
- Aroca R, Porcel R, Ruiz-Lozano JM. 2012. Regulation of root water uptake under abiotic stress conditions. *Journal of experimental botany* **63**: 43–57.
- Barnes AC, Benning C, Roston RL. 2016. Chloroplast membrane remodeling during freezing stress is accompanied by cytoplasmic acidification activating SENSITIVE TO FREEZING2. *Plant physiology* **171**: 2140–2149.
- Barnes AC, Myers JL, Surber SM, Liang Z, Mower JP, Schnable JC, Roston RL. 2023. Oligogalactolipid production during cold challenge is conserved in early diverging lineages. *Journal of experimental botany* **74**: 5405–5417.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)* **30**: 2114–2120.
- Brotzge JA, Wang J, Thorncroft CD, Joseph E, Bain N, Bassill N, Farruggio N, Freedman JM, Hemker K, Johnston D, *et al.* 2020. A technical overview of the New York state mesonet standard network. *Journal of Atmospheric and Oceanic Technology* **37**: 1827–1845.
- Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nature methods* **18**: 366–368.
- Chen L, Luo J, Jin M, Yang N, Liu X, Peng Y, Li W, Phillips A, Cameron B, Bernal JS, *et al.* 2022. Genome sequencing reveals evidence of adaptive variation in the genus *Zea*. *Nature genetics* **54**: 1736–1745.
- Cope AL, Schraiber JG, Pennell M. 2025. Macroevolutionary divergence of gene expression driven by selection on protein abundance. *Science (New York, N.Y.)* **387**: 1063–1068.
- Cuevas-Velazquez CL, Reyes JL, Covarrubias AA. 2017. Group 4 late embryogenesis abundant proteins as a model to study intrinsically disordered proteins in plants. *Plant signaling & behavior* **12**: e1343777.
- Degenkolbe T, Giavalisco P, Zuther E, Seiwert B, Hinch DK, Willmitzer L. 2012. Differential remodeling of the lipidome during cold acclimation in natural accessions of *Arabidopsis thaliana*: Lipidomics of *Arabidopsis* cold acclimation. *The Plant journal: for cell and molecular biology* **72**:

972–982.

Domon J-M, Baldwin L, Acket S, Caudeville E, Arnoult S, Zub H, Gillet F, Lejeune-Hénaut I, Brancourt-Hulmel M, Pelloux J, et al. 2013. Cell wall compositional modifications of *Miscanthus* ecotypes in response to cold acclimation. *Phytochemistry* **85**: 51–61.

Dure L III. 1993. A repeating 11-mer amino acid motif and plant desiccation. *The Plant journal: for cell and molecular biology* **3**: 363–369.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome biology* **20**: 238.

Fu Q, Liu Z, Bhawal R, Anderson ET, Sherwood RW, Yang Y, Thannhauser T, Schroyen M, Tang X, Zhang H, et al. 2021. Comparison of MS2, synchronous precursor selection MS3, and real-time search MS3 methodologies for lung proteomes of hydrogen sulfide treated swine. *Analytical and bioanalytical chemistry* **413**: 419–429.

Gallaher TJ, Peterson PM, Soreng RJ, Zuloaga FO, Li D-Z, Clark LG, Tyrrell CD, Welker CAD, Kellogg EA, Teisher JK. 2022. Grasses through space and time: An overview of the biogeographical and macroevolutionary history of Poaceae. *Journal of systematics and evolution* **60**: 522–569.

Gault CM, Kremling KA, Buckler ES. 2018. Tripsacum De novo Transcriptome Assemblies Reveal Parallel Gene Evolution with Maize after Ancient Polyploidy. *The plant genome* **11**: 180012.

Ginsawaeng O, Heise C, Sangwan R, Karcher D, Hernández-Sánchez IE, Sampathkumar A, Zuther E. 2021. Subcellular Localization of Seed-Expressed LEA_4 Proteins Reveals Liquid-Liquid Phase Separation for LEA9 and for LEA48 Homo- and LEA42-LEA48 Heterodimers. *Biomolecules* **11**.

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, et al. 2012. Phytozome: a comparative platform for green plant genomics. *Nucleic acids research* **40**: D1178–86.

Hundertmark M, Hinch DK. 2008. LEA (Late Embryogenesis Abundant) proteins and their encoding genes in *Arabidopsis thaliana*. *BMC genomics* **9**: 118.

Iwaya-Inoue M, Sakurai M, Uemura M (Eds.). 2018. *Survival Strategies in Extreme Cold and Desiccation: Adaptation Mechanisms and Their Applications*. Berlin, Germany: Springer.

Kellogg EA. 2015. PACMAD Clade. In: Kellogg EA, ed. *Flowering Plants. Monocots: Poaceae*. Cham: Springer International Publishing, 267–269.

Koster KL, Lynch DV. 1992. Solute accumulation and compartmentation during the cold acclimation of Puma rye. *Plant physiology* **98**: 108–113.

Koubaa S, Bremer A, Hinch DK, Brini F. 2019. Structural properties and enzyme stabilization function of the intrinsically disordered LEA_4 protein TdLEA3 from wheat. *Scientific reports* **9**: 3720.

Kovacs D, Agoston B, Tompa P. 2008. Disordered plant LEA proteins as molecular chaperones. *Plant signaling & behavior* **3**: 710–713.

Liu Y, Liang J, Sun L, Yang X, Li D. 2016. Group 3 LEA Protein, ZmLEA3, Is Involved in Protection from Low Temperature Stress. *Frontiers in plant science* **7**: 1011.

Liu Y, Wang L, Xing X, Sun L, Pan J, Kong X, Zhang M, Li D. 2013. ZmLEA3, a Multifunctional

Group 3 LEA Protein from Maize (*Zea mays* L.), is Involved in Biotic and Abiotic Stresses. *Plant & cell physiology* **54**: 944–959.

Li Q, Xiang C, Xu L, Cui J, Fu S, Chen B, Yang S, Wang P, Xie Y, Wei M, et al. 2020. SMRT sequencing of a full-length transcriptome reveals transcript variants involved in C18 unsaturated fatty acid biosynthesis and metabolism pathways at chilling temperature in *Pennisetum giganteum*. *BMC genomics* **21**: 52.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* **15**: 550.

Lyons JM. 1973. Chilling Injury in Plants. *Annual review of plant physiology* **24**: 445–466.

Marks RA, Van Der Pas L, Schuster J, Gilman IS, VanBuren R. 2024. Convergent evolution of desiccation tolerance in grasses. *Nature plants* **10**: 1112–1125.

Marrs KA. 1996. THE FUNCTIONS AND REGULATION OF GLUTATHIONE S-TRANSFERASES IN PLANTS. *Annual review of plant physiology and plant molecular biology* **47**: 127–158.

Mori K, Renhu N, Naito M, Nakamura A, Shiba H, Yamamoto T, Suzaki T, Iida H, Miura K. 2018. Ca²⁺-permeable mechanosensitive channels MCA1 and MCA2 mediate cold-induced cytosolic Ca²⁺ increase and cold tolerance in *Arabidopsis*. *Scientific reports* **8**: 550.

Nakashima K, Yamaguchi-Shinozaki K. 2006. Regulons involved in osmotic stress-responsive and cold stress-responsive gene expression in plants. *Physiologia plantarum* **126**: 62–71.

NDong C, Danyluk J, Wilson KE, Pocock T, Huner NPA, Sarhan F. 2002. Cold-regulated cereal chloroplast late embryogenesis abundant-like proteins. Molecular characterization and functional analyses. *Plant physiology* **129**: 1368–1381.

Ojeda-Rivera JO, Barnes AC, Ainsworth EA, Angelovici R, Basso B, Brindisi L, Brooks MD, Busch W, Buttelmann GL, Castellano MJ, et al. 2025. Designing a nitrogen-efficient cold-tolerant maize for modern agricultural systems. *The plant cell* **37**.

Örvar BL, Sangwan V, Omann F, Dhindsa RS. 2000. Early steps in cold sensing by plant cells: the role of actin cytoskeleton and membrane fluidity. *The Plant journal: for cell and molecular biology* **23**: 785–794.

Pearce RS. 2001. Plant Freezing and Damage. *Annals of botany* **87**: 417–424.

Perez-Riverol Y, Bandla C, Kundu DJ, Kamatchinathan S, Bai J, Hewapathirana S, John NS, Prakash A, Walzer M, Wang S, et al. 2025. The PRIDE database at 20 years: 2025 update. *Nucleic acids research* **53**: D543–D553.

Popova AV, Rausch S, Hundertmark M, Gibon Y, Hinch DK. 2015. The intrinsically disordered protein LEA7 from *Arabidopsis thaliana* protects the isolated enzyme lactate dehydrogenase and enzymes in a soluble leaf proteome during freezing and drying. *Biochimica et biophysica acta* **1854**: 1517–1525.

Puhakainen T, Hess MW, Mäkelä P, Svensson J, Heino P, Palva ET. 2004. Overexpression of multiple dehydrin genes enhances tolerance to freezing stress in *Arabidopsis*. *Plant molecular biology* **54**: 743–753.

Ranaweera T, Brown BNI, Wang P, Shiu S-H. 2022. Temporal regulation of cold transcriptional response in switchgrass. *Frontiers in plant science* **13**: 998400.

Roeder AHK, Shi Y, Yang S, Abbas M, Sasidharan R, Yanovsky MJ, Casal JJ, Ruffel S, von Wirén N, Assmann SM, et al. 2025. Translational insights into abiotic interactions: From Arabidopsis to crop plants. *The Plant Cell* **37**: koaf140.

Romero-Pérez PS, Martínez-Castro LV, Linares A, Arroyo-Mosso I, Sánchez-Puig N, Cuevas-Velazquez CL, Sukenik S, Guerrero A, Covarrubias AA. 2024. Self-association and multimer formation in AtLEA4-5, a desiccation-induced intrinsically disordered protein from plants. *Protein science : a publication of the Protein Society* **33**: e5192.

Sage RF, Christin P-A, Edwards EJ. 2011. The C(4) plant lineages of planet Earth. *Journal of experimental botany* **62**: 3155–3169.

Sapay N, Guermeur Y, Deléage G. 2006. Prediction of amphipathic in-plane membrane anchors in monotopic proteins using a SVM classifier. *BMC bioinformatics* **7**: 255.

Schat L, Schubert M, Fjellheim S, Humphreys AM. 2025. Drought tolerance as an evolutionary precursor to frost and winter tolerance in grasses. *Evolution; international journal of organic evolution*: qpaf006.

Schubert M, Grønvold L, Sandve SR, Hvidsten TR, Fjellheim S. 2019. Evolution of cold acclimation and its role in niche transition in the temperate grass subfamily Pooideae. *Plant physiology* **180**: 404–419.

Schubert M, Humphreys AM, Lindberg CL, Preston JC, Fjellheim S. 2020. To coldly go where no grass has gone before: A multidisciplinary review of cold adaptation in Poaceae. *Annual Plant Reviews online*: 523–562.

Singh KK, Graether SP. 2021. The in vitro structure and functions of the disordered late embryogenesis abundant three proteins. *Protein science: a publication of the Protein Society* **30**: 678–692.

Stitzer MC, Seetharam AS, Scheben A, Hsu S-K, Schulz AJ, AuBuchon-Elder T, El-Walid M, Ferebee TH, Hale CO, La T, et al. 2025. Extensive genome evolution distinguishes maize within a stable tribe of grasses. *bioRxiv*: 2025.01. 22.633974.

Sun W, Van Montagu M, Verbruggen N. 2002. Small heat shock proteins and stress tolerance in plants. *Biochimica et biophysica acta* **1577**: 1–9.

Suzuki N, Mittler R. 2006. Reactive oxygen species and temperature stresses: A delicate balance between signaling and destruction. *Physiologia plantarum* **126**: 45–51.

Takamori LM, Pereira AVC, Maia Souza G, Vieira LGE, Ferreira Ribas A. 2017. Identification of endogenous reference genes for RT-qPCR expression analysis in Urochloa brizantha under abiotic stresses. *Scientific reports* **7**: 8502.

Thomashow MF. 1999. PLANT COLD ACCLIMATION: Freezing tolerance genes and regulatory mechanisms. *Annual review of plant physiology and plant molecular biology* **50**: 571–599.

Törönen P, Holm L. 2022. PANNZER-A practical tool for protein function prediction. *Protein science : a publication of the Protein Society* **31**: 118–128.

Vera-Hernández P, Ortega-Ramírez MA, Martínez Nuñez M, Ruiz-Rivas M, Rosas-Cárdenas F de F. 2018. Proline as a probable biomarker of cold stress tolerance in Sorghum (Sorghum bicolor). *Mexican journal of biotechnology* **3**: 77–86.

Woodhouse MR, Cannon EK, Portwood JL 2nd, Harper LC, Gardiner JM, Schaeffer ML, Andorf CM. 2021. A pan-genomic approach to genome databases using maize as a model system.

Wu B, Luo D, Yue Y, Yan H, He M, Ma X, Zhao B, Xu B, Zhu J, Wang J, et al. 2024. Haplotype-resolved genome assembly of upland switchgrass provides insights into cold tolerance. *bioRxiv*: 2024.08.26.609807.

Xie Z, Lin W, Yu G, Cheng Q, Xu B, Huang B. 2019. Improved cold tolerance in switchgrass by a novel CCCH-type zinc finger transcription factor gene, PvC3H72, associated with ICE1-CBF-COR regulon and ABA-responsive genes. *Biotechnology for biofuels* 12: 224.

Xue C, Jiang Y, Wang Z, Shan X, Yuan Y, Hua J. 2021. Tissue-level transcriptomic responses to local and distal chilling reveal potential chilling survival mechanisms in maize. *Journal of experimental botany* 72: 7610–7625.

Xu K, Zhao Y, Gu J, Zhou M, Gao L, Sun R-X, Wang W-W, Zhang S-H, Yang X-J. 2022. Proteomic analysis reveals the molecular mechanism underlying the cold acclimation and freezing tolerance of wheat (*Triticum aestivum* L.). *Plant science: an international journal of experimental plant biology* 318: 111242.


Zhang Y, Ngu DW, Carvalho D, Liang Z, Qiu Y, Roston RL, Schnable JC. 2017. Differentially regulated orthologs in sorghum and the subgenomes of maize. *The plant cell* 29: 1938–1951.

Tables


Table 1. Summary of rhizome proteomics datasets, including number of samples per season, total proteins quantified, proteins shared across seasons, and differentially accumulated proteins (DAPs) accumulated (up) or depleted (down) in winter.

Species	Year	N samples	Genotype / Cultivar	Entry code	Clonal reps	Proteins quantified		DAP up	DAP down
						Total	Shared winter/summer		
<i>Tripsacum</i> spp.	2019	8	<i>T. dactyloides</i> 'LA_Pete_2' × <i>T. floridanum</i>	C002	1	4,451	4,451	123	175
			<i>T. dactyloides</i> 'KS_C9_9' × <i>T. floridanum</i>	C004	1				
			<i>T. dactyloides</i> 'KS_B6_1' × <i>T. floridanum</i>	C022	3				
			<i>T. dactyloides</i> 'KS_C9_10' × <i>T. floridanum</i>	C031	1				
			<i>T. dactyloides</i> 'LA_Pete_1' × <i>T. dactyloides</i> 'FL'	C071	1				
			<i>T. dactyloides</i> 'TX_HC_3' × <i>T. dactyloides</i> 'MO_TP'	C076	1				
<i>Tripsacum</i> spp.	2022	24	<i>T. dactyloides</i> 'LA_Pete_2' × <i>T. floridanum</i>	C002	3	3,468	3,054	100	141
			<i>T. dactyloides</i> 'KS_C9_9' × <i>T. floridanum</i>	C004	3				
			<i>T. dactyloides</i> 'KS_B6_1' × <i>T. floridanum</i>	C022	3				
			<i>T. dactyloides</i> 'KS_C7_3' × <i>T. floridanum</i>	C027	3				
			<i>T. dactyloides</i> 'IL_10_1' × <i>T. floridanum</i>	C043	3				
			<i>T. dactyloides</i> 'IL_14' × <i>T. dactyloides</i> 'FL'	C065	3				
			<i>T. dactyloides</i> 'LA_Pete_1' × <i>T. dactyloides</i> 'FL'	C071	3				
			<i>T. dactyloides</i> 'TX_HC_3' × <i>T. dactyloides</i> 'MO_TP'	C076	3				
<i>Angropogon gerardii</i>	2022	3	Sentinel	01-266*A	3	3,734	3,733	109	76
<i>Miscanthus giganteus</i>	2022	3	-	88-363*A	3	2,372	2,273	58	83
<i>Panicum virgatum</i>	2022	3	Dust Devil	01-277*B	3	4,071	4,068	174	218
<i>Sorghastrum nutans</i>	2022	2	Golden Sunset	22-095*A	2	2,641	2,464	66	56


Supplemental Figures




Supplemental Figure S1. Principal component analysis (PCA) of log-transformed protein abundance data. (A) PCA of 745 orthogroups from five species *Tripsacum* (Td-2019, Td-2022), *A. gerardii* (Ag), *M. giganteus* (Mg), *P. virgatum* (Pv), and *S. nutans* (Sn). **(B)** PCA of 1,889 proteins from *Tripsacum* hybrids. Points are colored by hybrid type—intraspecific *T. dactyloides* hybrids (Td × Td) and interspecific *T. dactyloides* × *T. floridanum* hybrids (Td × Tf). Shapes represent season (circle =summer, triangle = winter). The percentage of total variance explained by each principal component is shown on the axes.



Supplemental Figure S2. Volcano plots showing differentially accumulated proteins (DAPs) in rhizomes of *Tripsacum* spp., *Andropogon gerardii*, *Miscanthus giganteus*, *Panicum virgatum*, and *Sorghastrum nutans* under winter vs. summer conditions, with total number of proteins identified listed. Proteins with significantly elevated (blue) and reduced (pink) abundance in winter relative to summer are highlighted, with non-significant proteins in gray. The y-axis represents the significance as $-\log_{10}(\text{p-value})$, and the x-axis shows the $\log_2$ fold-change ($\log_2\text{FC}$) of the DAPs between winter and summer.




Supplemental Figure S3. Orthogroup sharing of significantly DAPs among five PACMAD grass species. (A) Winter-elevated. (B) Winter-reduced. Set sizes (horizontal bars) indicate the total number of differentially abundant orthogroups identified per species. Intersection sizes (vertical bars) represent orthogroups unique to individual species or shared among multiple species. Bar shading indicates the number of species sharing each orthogroup: unique to one species (white), shared between two species (gray), or shared among more than two species (black). Dots below the plot illustrate species-specific orthogroup intersections.




Supplemental Figure S4. (A) Enriched biological processes (BP) among differentially accumulated proteins (DAPs) in 475 winter-elevated and 609 winter-reduced proteins (proteins with unknown BP term were excluded). Dot size represents gene count per BP category, while dot color indicates the gene ratio (proportion of proteins annotated to each BP category). Only the top significantly enriched BP terms (lowest P-values) are displayed for clarity. **(B)** Mosaic plot of biological processes enriched among winter-elevated DAPs across five Andropogoneae species. The plot summarizes the distribution of functionally annotated proteins (n = 483 of 630; unannotated proteins were excluded) across 139 biological process

(BP) terms. These terms were consolidated into broader functional categories to aid interpretability. Each bar shows the relative abundance of functional categories within a species, highlighting both conserved and species-specific patterns. A Chi-square test for independence ($\chi^2[70]=97.5$, $p=0.017$) was performed to evaluate variation across species. Category labels are shown in the legend.




Supplemental Figure S5. Scatterplots showing the relationship between *Tripsacum* rhizome proteins (x-axis, log₂FC Winter/Summer, 2-year average) and cold-responsive

RNA expression in different seedling tissues (y-axis). (A) *Tripsacum* leaf mRNA from a freezing tolerant F₂ bulk before and after 7 days cold acclimation (tol_5°C/tol_22°C) descended from the same *Tripsacum* hybrids used for rhizome proteomic analysis. (B-E) *Tripsacum* 'Pete' tissues (5°C/22°C): leaf, top-crown, bottom-crown, and root. (F-I) *Zea mays* tissues (5°C/22°C): leaf, top-crown, bottom-crown, and root. Points are colored by shared differential expression status: blue for upregulated, pink for downregulated, and gray for non-significant changes. Dashed horizontal lines indicate the $\pm 1 \log_2FC$ thresholds. Spearman's correlation coefficient (ρ) and P-value are shown in the upper left of each panel, along with the number of genes analyzed (n). The LEA3 gene is labeled in each panel.




Supplemental Figure S6. Hydropathy profiles of LEA3 orthologs from eight species. Kyte-Doolittle plots were generated using ProtScale (ExPASy) with a window size of 11

amino acids. The x-axis indicates amino acid position, and the y-axis represents hydropathy score. Red boxes highlight regions corresponding to conserved 11-mer LEA motifs.




Supplemental Figure S7. Stacked bar chart illustrating the composition of BPs within curated cold-related functional categories. The plot includes 230 proteins from the top 50 DAPs accumulated in winter across species, after excluding those assigned to the cold response category “Other” (i.e., proteins with annotations deemed less relevant to cold tolerance interpretation). Cold-related categories are shown on the x-axis and sorted by total protein count. Each bar is color-coded by BP, highlighting the functional composition of each category. Less informative or poorly annotated BPs were grouped under “Other” in the legend. This visualization supports the manual consolidation of specific biological processes into broader cold-response categories used throughout the study.



Supplemental Figure S8. AlphaFold 3 prediction of LEA3 homo-tetramer assemblies.

Predicted quaternary structures of LEA3 homotetramers for (A) *Zea mays* and (B) *Tripsacum dactyloides*. Both orthologs are predicted to form tetrameric assemblies. Colors indicate per-residue confidence (pLDDT): dark blue (very high, >90), light blue (confident, 70–90), yellow (low, 50–70), orange (very low, <50). Predicted Aligned Error (PAE) matrices (right) show positional uncertainty between residue pairs; dark green indicates low error.



Supplemental Figure S9. Rhizome morphology at sampling. Paired images display representative rhizome tissues (white arrows) collected during summer (August, left) and winter (January, right) for **(A)** *Andropogon gerardii*, **(B)** *Miscanthus giganteus*, **(C)** *Sorghastrum nutans*, **(D)** *Tripsacum dactyloides*, **(E)** and *Panicum virgatum*. Summer samples exhibit active bud growth and pale storage tissue. Winter samples exhibit dormancy and epidermal hardening. Proteomic analysis was restricted to mid-section of these storage organs. Scale bars represent approx. 1 cm.