1. The probability of a leap year selected at random contain 53						
Sunday is:						
(a) 53/366	(b) 1/7	(c) 2/7	(d) 53/365			
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at						
random. The probab	ility of draw	ing a black ball	is:			
(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5			
3. The probability the	nat it will rain	n tomorrow is 0).85. What is the			
probability that it wi	II not rain to	morrow				
(a) 0.25 (l	o) 0.145	(c) 3/20	(d) none of these			
4. What is the proba	ability that a	number selecte	ed from the numbers			
(1, 2, 3,,15) is	a multiple o	f 4?				
(a) 1/5 (l	b) 4/5	(c) 2/15	(d) 1/3			
5. What are the total	al outcomes	when we throw	three coins?			
(a) 4		1 /	, ,			
		number selecte	ed at random from the			
numbers (1,2,3,						
• •	, ,	` '	(d) none of these			
7. The sum of the p						
(a) 2 (b)		` '				
• .		re given; choos	e the correct answer			
for that which is not	•					
	15		(d) none of these.			
9. If three coins are		ultaneously, tha	an the probability of			
getting at least two	•		4.15 - 4.5			
(a) 1/4 (b)		` '				
10. A letter is chos						
*ASSASSINATION						
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.			
	_		ting an even number.			
(A) 2/3 (B) 1 ((C) 5/6 ((D) 1/2			
12 Two coine are th	rown of the	como timo. Ein	d the probability of			
12. Two coins are the	nown at the	Saine unie. Fin	u tile probability of			
getting both heads. (A) 3/4 (B) 1/4	(C) 1/2	(D) 0				
(A) $3/4$ (D) $1/4$	(0) 1/2	(D) 0				
13. Two dice are thrown simultaneously. The probability of getting a						

1

sum of 9 is:

	ime number.	red from 1 to 1 (C) 1/4		probability of 29/100	
of drawing a blue balls in	a blue ball is d n a bag is:	louble that of a	red ball, the	f the probability en the number of	
(A) 5	(B) 10	(C) 15	(D) 20		
	random from	ntains 12 defe			
	(B) 147	7/150 (C)	1/25	(D) 1/50	
mixed thoro	oughly. One ca lity that the nu	mbers 2 to 10 rd is drawn fro mber on card i (C) 3/10	m this box r s a perfect s	andomly, then square.	
18. What is (A) 1/7	the probabilit (B) 53/366	(C) 2/7			
probability	of getting a ki	a well shuffled ng of red suit. C) 7/52 (D)		cards. Find the	
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) $1/6$ (B) $1/12$ (C) $7/12$ (D) $5/12$					
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$					

2

(A) 1/10 (B) 3/10 (C) 1/9 (D) 4/9

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225				
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that x^2 < 2 is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$							
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7							
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$							
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$				
27. Runs scored by batsman in 5 one day matches are 50, 70, 82, 93, and 20. The standard deviation is							
a) 25.79	b) 25.49	c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $\frac{1}{2}$ 30. X is a variate	b) $^1\!/_3$ te between 0 and	c) $\frac{1}{4}$ 3. The value of I 27	$\frac{d}{1}$ d) $\frac{1}{6}$ E(X ²) is				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of the probability?		wing value	es, which	one is no	t possib	ole in
a) P(x) = 1 c) P(x) = 0.5		b) $\sum x P(x) =$				
33.If E(x) =		(z) = 4, th		x) =?		
a) 2	b) 6		c) 0		d) Insu	fficient data
34.The cova	ariance	of two inc	lependen	t random	variable	e is
a) 1	b) 0		c) - 1		d) Und	efined
35.lf Σ P(x)	$= k^2 - 8$	3 then, the	e value of	f k is?		
a) 0	b) 1		c) 3		d) Insu	ıfficient data
36.If P(x) = a) 1	0.5 and b) 0.5	x = 4, the	en E(x) = ' c) 4	?	d) 2	
37.In a disc is always?	rete pro	bability d	istributio	n, the sun	ո of all բ	orobabilities
a) 0	b) Infin	ite	c) 1		d) Unde	efined
38.If the probability of hitting the target is 0.4, find mean and variance.						
a) 0.4, 0.24	ŀ	0) 0.6, 0.2	4	c) 0.4, 0	.16	d) 0.6, 0.16
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6						
40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution?						

c) 5

b) 4

a) 3

4


d) 7

\ 1. 0						
a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0						
c) Mean is 0	and variance is	∞ d) Mean is ∞	and variance	e is 0		
42 Variance	of a random va	ariable X is given b	v			
a) E(X)	D) E(XZ)	c) $E(X2) - (E(X))$	())2	d) (E(X))2		
43.Mean of	a random varia	ble X is given by _				
a) E(X)		c) $E(X2) - (E(X))$		d) (E(X))2		
-, -(,	~ / _(* !_/	0) =(*!=) (=(*!)	/-	-) (-(**))-		
11 Moon of a	a constant 's' is					
	constant 'a' is		• • •			
a) 0	b) a	c) a/2	d) 1			
45.Variance	of a constant 'a	a' is				
a) 0	b) a	c) a/2	d) 1			
4) 0	D) u	0) 4/2	u) 1			
		6340				
16 Find the r	noon and varia	noo of V2				

40.Find the mean and variance of X?

X	0	1	2	3	4
f(x)	1/9	2/9	3/9	2/9	1/9

47. Find the expectation of a random variable X?



48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

5

- a) np
- b) npq
- c) np2q
- d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}