1. The probability of a leap year selected at random contain 53				
Sunday is:				
(a) 53/ 366 (b) 1/7 (c) 2/7				
2. A bag contains 3 red and 2 blue marbles				
random. The probability of drawing a black				
(a) 3/5 (b) 2/5 (c) 0/5				
3. The probability that it will rain tomorrow	v is 0.85. What is the			
probability that it will not rain tomorrow				
(a) 0.25 (b) 0.145 (c) 3/20	•			
4. What is the probability that a number se	elected from the numbers			
(1, 2, 3,,15) is a multiple of 4?				
(a) 1/5 (b) 4/5 (c) 2/15				
5. What are the total outcomes when we t				
(a) 4 (b) 5 (c) 8	• •			
6. The probability that a prime number se	lected at random from the			
numbers (1,2,3,35) is :				
(a) 12/35 (b) 11/35 (c) 13	3/35 (d) none of these			
7. The sum of the probability of an event a				
(a) 2 (b) 1 (c) 0 (d)				
8. The following probabilities are given; c	hoose the correct answer			
for that which is not possible.				
(a) 0.15 (b) 2/7 (c) 7/5				
9. If three coins are tossed simultaneously	, than the probability of			
getting at least two heads, is:				
(a) 1/4 (b) 3/8 (c) ½				
10. A letter is chosen at random from the	letters of the word			
♦ ASSASSINATION ♦ . The probability that	the letter chosen has:			
(a) 6/13 (b) 7/13 (c)) 1 (d) none of these.			
11. A dice is thrown. Find the probability of	f getting an even number.			
(A) 2/3 (B) 1 (C) 5/6	(D) 1/2			
12. Two coins are thrown at the same time	. Find the probability of			
getting both heads.				
(A) 3/4 (B) 1/4 (C) 1/2 (D)	0			
13. Two dice are thrown simultaneously. The probability of getting a				

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	/9			
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4	(B) 27/50	(C) 1/4	(D)	29/100			
_	a blue ball is			If the probability en the number of			
(A) 5	•	(C) 15	(D) 20				
taken out at non-defecti	t random fron ve bulb is:	ontains 12 defo n this box. The	n the probab	ility that it is			
(A) 143/150	(B) 14	7/150 (C)) 1/25	(D) 1/50			
mixed thoro	oughly. One callity that the n	umbers 2 to 10 ard is drawn fr umber on card (C) 3/10	om this box i is a perfect	randomly, then square.			
18. What is the probability of getting 53 Mondays in a leap year? (A) 1/7 (B) 53/366 (C) 2/7 (D) 7/366							
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13							
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12 (E) 1/2							
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4							

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
	•	(C) 1/365	(D) 1/133225				
2. Then the p	er x is chosen at r robability that x ²) 2/5 (C) 3/5	< 2 is?	umbers -2, -1, 0 , 1,				
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7							
Then the prol		a multiple of 3 and	0 natural numbers. l 4 is:				
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 4/21				
			nes are 50, 70, 82,				
	he standard devi	ation is c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
(a) $\frac{1}{2}$	b) $^{1}/_{3}$	c) $^{1}/_{4}$	d) $^{1}/_{6}$				
		nd 3. The value of c) 27	E(X ²) is				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of the following values, which one is not possible in probability?					
a) $P(x) = 1$	b)∑x d) P(x				
	2 and E(z) = 4 b) 6	t , then E(z – c) 0	•	ufficient data	
34.The cov	ariance of two	independer	nt random variab	le is	
a) 1	b) 0	c) - 1	d) Un	defined	
• •) = k² – 8 then, b) 1			sufficient data	
• •	0.5 and x = 4, b) 0.5	, ,	? d) 2		
37.In a disc is always?	rete probabilit	y distributio	on, the sum of all	probabilities	
•	b) Infinite	c) 1	d) Unc	lefined	
38.If the probability of hitting the target is 0.4, find mean and					
variance. a) 0.4, 0.24	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16	
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6					
 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution? 					

c) 5

d) 7

a) 3

b) 4

		and varia and varia		_ ,			
	Variance (X)	e of a rand b) E(X			given by (2) - (E(X))		. d) (E(X))2
		a random b) E(X2					d) (E(X))2
44.N a) 0	Mean of	a constan b) a	t 'a' is	c) a/2	_ ·	d) 1	
45. Variance of a constant 'a' is a) 0							
46.Find the mean and variance of X?							
г 	Х	0	1	2	3	4	
	f(x)	1/9	2/9	3/9	2/9	1/9	
	, 4/3 Find the	b) 3 expectatio			c) 2, 2/3 ariable X?	,	d) 3, 2/3

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2