■ Cell Shedding Field Guide

Targeted Senolytic Clearing, Autophagy Activation & SASP Reduction

Executive Summary: *Senescent cells are damaged, non-dividing 'zombie' cells that accumulate with biological age, secreting pro-inflammatory SASP cytokines that degrade surrounding tissue architecture. Periodic senolytic clearing restores youthful tissue dynamics.*

Framework Perspective & Clinical Context

Cellular senescence evolved as an essential anti-cancer mechanism, permanently halting the division of damaged cells. However, when senescent cell clearance falls behind accumulation, these lingering cells secrete a toxic cocktail of pro-inflammatory cytokines, chemokines, and matrix metalloproteinases known as the Senescence-Associated Secretory Phenotype (SASP).

Cellular Mechanisms & Biophysical Principles

1. Senolytic Disruption of Anti-Apoptotic Pathways

Senescent cells survive by up-regulating Senescent Cell Anti-Apoptotic Pathways (SCAPs) involving BCL-2, BCL-XL, and PI3K/AKT. Plant flavonoids like Fisetin and Quercetin target these survival networks, triggering apoptosis specifically in senescent cells while sparing healthy tissue.

2. Pulsed vs. Continuous Interventions

Unlike daily maintenance nutrients, senolytic interventions are administered in intermittent 'hit-and-run' pulses (e.g., 2 consecutive days per month). This clears senescent cell populations without impairing normal physiological tissue repair.

3. Lysosomal Autophagy & Mitophagy

Autophagy is the lysosomal degradation process that recycles misfolded proteins and damaged mitochondria (mitophagy). Activating autophagy prevents precocious senescent cell conversion and maintains cellular bioenergetics.

Structured Step-by-Step Protocol

Integrate targeted practitioner-grade formulations into daily bio-rhythms:

Phase / Timing	Intervention & Compound	Target Mechanism & Purpose
Pulse Days (Day 1-2)	High-Dose Liposomal Fisetin & Quercetin	Transiently disrupts SCAP survival networks to clear senescent cells
Daily Maintenance	Spermidine & Trehalose Autophagy Boosters	Stimulates lysosomal degradation and autophagic cellular recycling
Daily Maintenance	SASP Dampeners (Curcumin, Boswellia, EGCG)	Suppresses pro-inflammatory cytokine secretion from remaining senescent cells
Post-Pulse (Day 3-30)	NAD+ Precursors & Stem Cell Co-Factors	Supports tissue regeneration and progenitor cell renewal after clearing

VERIFIED PRACTITIONER PROTOCOL: CELL SHEDDING PRACTITIONER PROTOCOL

Order direct practitioner-grade formulations on Fullscript. Enjoy automatic monthly refills, verified quality, and practitioner oversight.

Order Protocol on Fullscript Direct →

Lifestyle, Circadian & Photobiomodulation Practices

• Intermittent Pulsed Fasting: Practice 16-to-24 hour fasts periodically to deplete cellular nutrient sensors (mTOR) and trigger autophagic clearing.

• High-Intensity Interval Exercise: Brief bursts of intense exertion induce transient cellular stress followed by robust autophagic adaptation.

• Sauna Heat-Shock Protein Induction: Infrared or traditional sauna sessions (20 min at 80°C/175°F) induce heat-shock proteins (HSP70) to refold damaged proteins.

Clinical Safety & Practitioner Guidance

High-dose senolytic protocols should be pulsed intermittently rather than taken continuously. Pregnant or nursing mothers should avoid senolytic protocols.

Fullscript Plan Link: https://ca.fullscript.com/u/plans/TXVsdGlQYXRpZW50UGxhbi0xMDQ1MA==

Prepared by Spiral of Cellular Light — Certified Regenerative Health Program.
Disclaimer: Agewell Brief receives practitioner affiliate commission on Fullscript orders at no added cost to you. Always consult your qualified healthcare practitioner before beginning any new health protocol.