Yang–Mills on a finite torus along Bałaban's route: a verified reduction

Claude

October 2026

Abstract

Bałaban's renormalization-group construction of four-dimensional SU(2) lattice Yang–Mills theory rests on the control of the running coupling, his Theorem 2, whose proof was never published. We prove that a weak form of Theorem 2 (two-sided bounds on $1/g_k^2$ with an additive constant), which is enough for Bałaban's Theorem 1, follows from one uniform inequality (H2) on a two-step minimizer map together with named fixed-level and history inputs; the shooting lemmas, the arithmetic of Theorem K, and the bridge lemmas of the certifier are formalized in Lean 4. We prove that, on a fixed torus, a uniform moment bound (H-U) gives compactness of the lattice Schwinger functions of the centred plaquette density, and that every limit point satisfies the torus forms of the Osterwalder–Schrader axioms OS0–OS3. The bound (H-U) is reduced, not proved: it rests on sourced versions of Bałaban's bounds, on a hypothesis (FX) that is now reduced to a holomorphy bound (H-hol) for Bałaban's increments together with history bounds and three named inputs of the large-field step, and on a bridge from the source form to (H-U), and one dependency table records the status of each block. At $L = 13$, inside Bałaban's range $L > 11$, (H2) is a certified theorem: an explicit majorant and an outward-rounded interval branch-and-bound give $\rho_1(j, K)^2 \le 10 < 13$ for every level j (including $j = \infty$) and every momentum K, so $\rho_2 \le 10 < 13$, given our reading of Bałaban's averaging operator; the result is not a solution of the Clay Millennium problem.

Contents

1 Introduction **4**
- 1.1 The problem . . . 4
- 1.2 Bałaban's programme and the missing Theorem 2 . . . 4
- 1.3 The two targets and the main results . . . 4
- 1.4 Status conventions . . . 6
- 1.5 Relation to other work . . . 7
- 1.6 Organization . . . 8

2 Setup and notation **8**
- 2.1 Lattice, measure, and scales . . . 9
- 2.2 Range of L . . . 9
- 2.3 The coupling sequence . . . 9
- 2.4 One-loop notation . . . 9
- 2.5 Observables and Schwinger functions . . . 10
- 2.6 Names . . . 10

3 The coupling flow and Theorem 2 **11**
3.1 What Bałaban's Theorem 1 actually needs . . . 11
3.2 The weak form proved conditionally . . . 11
3.3 Literal (0.31) . . . 12
3.4 Shooting lemmas . . . 13
3.4.1 Lemma D′ (shooting without early positivity) [PROVED+VERIFIED; LEAN] . . . 13
3.4.2 Lemma D″ (tail-sum form; no index J) [PROVED+VERIFIED; LEAN] . . . 13
3.4.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [PROVED, UNVERIFIED (SWEEP); 4 OF 19 TABLE ROWS VERIFIED] . . . 14
3.4.4 Lemma D and Lemma E [PROVED+VERIFIED; LEAN] . . . 15
3.4.5 History dependence (L5) [PROVED RELATIVE TO LISTED INPUTS; VERIFIED (ROUND 7)] . . . 15
3.5 The exact structure of the one-loop sums . . . 16
3.5.1 Telescoping [PROVED+VERIFIED, CORRECTED SCOPE] . . . 16
3.5.2 Anchor-free identity [PROVED+VERIFIED] . . . 16
3.5.3 The constant c [PROVED+VERIFIED] . . . 16
3.5.4 The decomposition $S_k = ck \log L + \kappa_W^F - \kappa_k$ . . . 17
3.5.5 Level-uniform symbol facts used below [PROVED+VERIFIED] . . . 17
3.6 Bounded window sums: two formally distinct conditional routes . . . 18
3.6.1 Route K: Theorem K [PROVED+VERIFIED, CONDITIONAL ON (B2) AND THE FIXED-LEVEL INPUTS] . . . 18
3.6.2 Route B: Theorem B_{PW} [PROVED+VERIFIED, CONDITIONAL (LOGIC OF THEOREM B_{PW})] . . . 19
3.6.3 Route B inputs after `BETA_FP_U` [PROVED, UNVERIFIED; NO VERIFIER FILE] . . . 20
3.6.4 Comparison . . . 21
3.7 The hypothesis (B2) of Route K, and the single uniform inequality (H2) . . . 21
3.7.1 Proved . . . 22
3.7.2 Refuted, and the failed one-step bound . . . 22
3.7.3 (H2) and the two-step theorems [PROVED+VERIFIED, CONDITIONAL ON (H2)] 22
3.7.4 One-step sufficiency [ARITHMETIC PROVED; INPUT CERTIFIED AT $L = 13$] . . . 23
3.7.5 The precise remaining inequality . . . 23
3.8 Conclusion of the reduction . . . 23
3.9 The convergence route: what it adds, and its status . . . 24
3.9.1 Components and status . . . 25
3.9.2 Early steps and j_* . . . 25

4 The finite-torus existence theorem **26**
4.1 What "exists" means here . . . 26
4.2 Hypothesis delivered by the blocks . . . 26
4.3 Theorem (finite-torus existence, compactness form) . . . 27
4.4 Theorem (nontriviality), conditional on (B-NT) . . . 27
4.5 Proof modulo blocks . . . 28
4.5.1 (B-T2) Theorem 2: the running coupling . . . 28
4.5.2 (B-S) Sourced bounds E14-a..g . . . 28
4.5.3 (B-N1) Bridge: source-form bounds ⇒ (H-U) [OPEN] . . . 28
4.5.4 (B-C) Compactness [PROVED HERE, GIVEN (H-U)] . . . 29
4.5.5 (B-REG) Regularity (OS0) [PROVED HERE, GIVEN (H-U)] . . . 30

4.5.6 (B-TI) Translation and hypercubic invariance (OS1) [PROVED HERE, GIVEN (H-U)] . . . 30
4.5.7 (B-RP) Reflection positivity (OS2) . . . 30
4.5.8 (OS3) Symmetry and reality [PROVED HERE] . . . 31
4.5.9 (B-NT) Nontriviality: what is needed . . . 31
4.6 Dependency table . . . 32
4.7 What the theorem does not give . . . 44
4.8 Corrections after verification . . . 44

5 The sources block **45**
5.1 Target 1: inputs of Theorem 2w . . . 45
5.2 Target 2, step 2: the sourced Theorem 1 (ledger B11) . . . 47
5.3 Step 3: the (S) clauses and (FX) (ledgers B21, B19) . . . 52
5.4 Steps 4–5: N3, (H-hol), and the large-field inputs (ledgers B18, B16, B17) . . . 54
5.5 Step 6: thresholds and second history derivatives (ledgers B15, B14) . . . 58
5.6 Step 7: history bounds and the gain (ledger B13) . . . 59
5.7 Step 8: Bałaban-level propagator inputs (ledgers A1–A4) . . . 61
5.8 Step 8′: the three write-outs of the propagator expansions . . . 66
5.9 Step 8″: round 5, the remaining Group A rows and Lemma G3 . . . 68
5.10 Step 8‴: round 6, locality, the boundary, the 2D gap, and the order of constants . . . 72
5.11 Step 8⁗: round 7, the implicit Group A rows, (M_j), L5, and the sweep . . . 75
5.12 Step 9: the alternative (E_U) route . . . 79

6 Numerical and certified results **80**
6.1 First increment $\beta_1(0)$ . . . 80
6.2 Later increments ($L = 2$ only) . . . 81
6.3 Quantities of the Route K proof . . . 81
6.4 Values of (H2) [NUMERICAL; (H2) AT $L = 13$ IS CERTIFIED IN SECTION 6.7] . . . 82
6.5 One-step values . . . 83
6.6 A rigorous point enclosure at $L = 3$ [INTERVAL ENCLOSURE, VERIFIED] . . . 84
6.7 (H2) at $L = 13$ [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)] . . . 84
6.8 Round 7: more certified L, the gap and neutral-sector checks, and an open tension . . . 86
6.9 Certification plan and j_0 [DESIGNED; SUPERSEDED AT $L = 13$ BY SECTION 6.7] . . . 87

7 What remains **88**
7.1 Certification . . . 88
7.2 Fixed-level and structural inputs of Theorem 2w . . . 88
7.3 Items of the sources block . . . 90
7.4 Verification debt . . . 93
7.5 Open for stronger statements . . . 94
7.6 Why the mass gap is not on this route . . . 94
7.7 Trust ledger . . . 94

A Lean formalization **104**

B Index of results by status **106**

C Label map **110**

1 Introduction

1.1 The problem

The Clay problem on quantum Yang–Mills theory asks for a four-dimensional quantum gauge theory with compact simple gauge group that satisfies axioms at least as strong as those of Osterwalder and Schrader [19, 20], and that has a mass gap [15]. One constructive approach starts from Wilson's lattice action on a torus with lattice spacing ε and takes $\varepsilon \to 0$. The bare coupling $g_0(\varepsilon)$ must then go to zero at the rate that asymptotic freedom prescribes, so that the effective coupling at a fixed physical scale stays fixed. This paper treats the first limit only: lattice spacing to zero on a fixed torus. It does not treat the infinite-volume limit or the mass gap (Section 4.7).

1.2 Bałaban's programme and the missing Theorem 2

Between 1984 and 1989 Bałaban built a renormalization-group (RG) analysis of lattice gauge theories [1, 2, 3, 5, 4, 6, 7, 8, 10, 9, 11, 12]. His main output for $d = 4$ is Theorem 1 of CMP 122 II: uniform bounds on the effective densities at every RG scale, under the hypothesis that the sequence of effective couplings stays small. The hypothesis was to be discharged by his Theorem 2, which controls the running coupling. The two statements read as follows.

Theorem 2 (CMP 109 (1987), p. 259): "Let $d = 4$, $G = SU(2)$, and let γ be a sufficiently small positive constant, then for a sufficiently small positive g there exists a bare coupling constant $g_0 = g_0(\varepsilon, g)$ such that the sequence of the effective coupling constants g_k is contained in the interval $]0, \gamma]$, and $g_K = g$. Moreover, there exist constants β, β', $0 < \beta \leq \beta'$, such that g_k satisfy the inequality $1/g^2 + \beta \log(L^k\varepsilon)^{-1} \leq 1/g_k^2 \leq 1/g^2 + \beta' \log(L^k\varepsilon)^{-1}$. (0.31)" Balaban adds that the proof "will be given in a separate paper". It was never published (CMP 122 II p. 355, verbatim: "has not been published yet, so we have Theorem 1 with the assumption").

Theorem 1 (CMP 122 II p. 355): "If the sequence of the effective coupling constants is contained in an interval $]0, \gamma]$ with a sufficiently small positive γ, then the effective densities ρ_k have the form, and satisfy all the conditions and bounds, described in Sect. 2 [III]." The bound (0.1) there reads $\chi_k \exp[-(1/g_k^2)A(U_k) - E_-|T_\eta|] \leq \rho_k \leq \exp E_+|T_\eta|$ (corrected reading, `BALABAN_EXCERPTS_CHECK`[1] item 3).

Without Theorem 2, the hypothesis of Theorem 1 is an assumption on the unknown coupling flow. Every continuum statement built on Bałaban's bounds inherits this assumption.

1.3 The two targets and the main results

The paper has two targets. The first target is a proof of enough of Theorem 2 to remove the coupling hypothesis from Theorem 1. The second target is a finite-torus existence theorem for the continuum limit in compactness form, with the Osterwalder–Schrader properties proved on the limit. Both results are reductions: each holds under named hypotheses, and the paper records the status of every hypothesis.

Theorem A (weak Theorem 2; proved in Section 3)**.** *Let $G = SU(2)$, $d = 4$, and L odd, $L > 11$ (or L odd, $L \geq 3$, with Bałaban's inductive assumption (H0) and his construction of $\beta_j(g)$ at this L as explicit hypotheses).*

[1] Project files named in this paper live in the `notes/` directory of the `yang-mills` repository [14].

(i) *Assume the uniform inequality (H2) (Section 3.7.3)* [CERTIFIED AT $L = 13$ (THEOREM 6.1); CERTIFIED AT $L = 13$, AND IN ROUND 7 AT $L = 15, 17, 21$ (FULL RERUNS, VERIFIED, SECTION 6.8); A HYPOTHESIS AT OTHER L], *the fixed-level inputs (PI), (F), (M_j), V1, (G2) (Section 3.6.1), the history hypothesis L5 (Section 3.4.5; round 7: proved relative to listed inputs, Theorem 5.15), and the bound 7′, $|\beta_{k+1}(h, g) - \beta_{k+1}(0)| \le C_\beta g$ on every (non-monotone) past (item 7′ of Section 7)* [PROVED+VERIFIED, CONDITIONAL ON (H0) (LEDGER A5) OR ON THE (G-GAIN) INPUTS (LEDGER B13)]. *Then the conclusions (b), (c′), (S) of Theorem 3.1 (Theorem 2w) hold. Status of the implication:* [PROVED+VERIFIED, CONDITIONAL], *by Route K (Theorem 3.8 and Lemma 3.4).*

(ii) *Given the conclusions of Theorem 2w, Bałaban's Theorem 1 holds for the bare coupling $g_0(K, g)$, for every K and every $g \in]0, g_1]$, without his coupling assumption. Status:* [PROVED; THE SWEEP VERIFIED IN ROUND 7 (LEDGER B10), CONDITIONAL ON ROUTE K OR ROUTE B INPUTS, ON L5, ON THE FINITE-VOLUME IDENTIFICATION OF THE STEP (LEDGER B8, REPLACED BY A PROVED FORM RELATIVE TO LISTED INPUTS AND THE VOLUME HYPOTHESIS (VOL) OF SECTION 2.1), AND ON BALABAN'S PUBLISHED PROOFS] *(Section 3.8).*

A second, formally distinct route (Route B, Theorem B_{PW} in Section 3.6.2) also gives (i), under its own input set (items B1–B7 of Section 7.2). Both routes rest on (H2).

At $L = 13$ the hypothesis (H2) is discharged by Theorem 6.1, so the weak form of Theorem 2 rests only on the fixed-level inputs, L5, item 7′, and the Bałaban-level items of the trust ledger (with the residue of Theorem 6.1, Section 6.7).

Theorem B (finite-torus existence; proved in Section 4)**.** *Let $G = SU(2)$, $d = 4$, L odd and large, and $\ell \in 2\mathbb{N}$ fixed. Let $g_0 = g_0(\varepsilon, g)$ be a shooting solution whose effective couplings satisfy $g_k \in]0, \gamma]$ for $0 \le k \le K$ and $g_K = g$. Assume the uniform bound (H-U) of Definition 4.1. When (H-U) is taken from the blocks rather than assumed, the chain also uses the $\kappa \to \kappa/2$ decay recovery of the sourced small-field step (ledger B22), a* [VERIFIED REDUCTION] *by $L/2$-rescaling that rests on the unprinted large-field T-step rate loss (ledger A23), and the volume hypothesis (Vol), $\ell > 22M$ (Section 2.1). Then the lattice Schwinger functions of the centred plaquette density are relatively compact as $\varepsilon = L^{-K} \to 0$, and every limit family satisfies the torus forms of (OS0)–(OS3). This is Theorem 4.2,* [PROVED MODULO NAMED BLOCKS]. *Nontriviality is the separate Theorem 4.3, conditional on (NT).*

The sourced step also uses the unit-scale condition (U), $\rho_1 \le c_* r_0 / M^a$ at $\lambda_K = 1$ (C23): this is a choice of the source-gradient constant, made once after r_0 and before K, not a hypothesis on test functions, and the bridge B-N1 meets it by rescaling at a C^n cost (this (U) is not the Route B hypothesis (U)).

The trust boundary of the two theorems is recorded in the trust ledger (Section 7.7): Theorem A rests on 4 Bałaban-level statements and one external result, and Theorem B rests on 17 Bałaban-level statements and four external results. Part (i) of Theorem A, Theorem 2w, rests on one of them, A5 (which includes Bałaban's assertion that the whole-lattice limit of CMP 109 (1.22) exists), and its use of A5 is a verified reduction. Of the 19 Bałaban-level statements in the ledger, 13 are printed by Bałaban without a proof (3 of the 4 behind Theorem A and 11 of the 17 behind Theorem B). After round 6, each of these 11 has a verified proof relative to listed inputs or a verified reduction (Sections 5.9, 5.10, and 7.7). After round 7 (Section 5.11), so has each implicit row: A16 and A17 are proved relative to listed inputs, and A18, A19, and A23 are verified reductions. So no Group A row remains without a verified proof or verified reduction (cited-but-unproved and implicit alike); the only Theorem B Group B row without proof or verified reduction is B3, the lattice transfer

of the log coefficient (F), which is reduced but unverified and carries an open numerical tension (Section 6.8). Round 7 also proves the fixed-level input (M_j) and the history hypothesis L5 relative to listed inputs (Theorems 5.14 and 5.15). Four of these (A1–A4, the propagator and covariance bounds) are now verified reductions to published theorems of CMP 99 and CMP 102; their joint residue is Bałaban's walk expansions of the final propagators (through the walk-expansion form A1w of A1), the complex walk bound (W^c) in the exponential gauge and the decay of $C^*\Delta_k C$ at real U (A7) under A2, the regular-class identification A18, two unprinted conventions, and a stated torus-side hypothesis.

The hypothesis (H-U) is the output of three blocks: Theorem 2 (B-T2, that is, Theorem A), sourced versions of Bałaban's bounds (B-S), and a bridge from the source form to (H-U) (B-N1). The dependency table, Section 4.6, gives the status of each block and is the single status table of the paper. Section 7 lists what remains.

What the paper does not give: literal (0.31) at every K (Section 3.3), the limit of $\beta_j(0)$ and a rate (Section 3.9), uniqueness of the limit, infinite volume, rotation invariance, and a mass gap (Section 4.7). Theorem B is a compactness ("weak-existence") result. Under Jaffe–Witten footnote 2 it is not an admissible solution of the Millennium problem (Section 4.1).

1.4 Status conventions

Every result carries a status tag in small capitals. The tags mean the following.

- [PROVED+VERIFIED]: proved in a project file and confirmed by an independent verifier file.
- [PROVED+VERIFIED, CONDITIONAL ON X]: the implication is confirmed; X is open or assumed.
- [PROVED, UNVERIFIED]: proved in a project file; no verifier has graded it yet.
- [LEAN]: formalized in Lean 4 + Mathlib, no `sorry`, axioms propext, Classical.choice, Quot.sound only.
- [VERIFIED QUOTE]: read on the page image of Balaban's paper (`reports/BALABAN_EXCERPTS_CHECK.md` or a verifier file).
- [NUMERICAL, REPRODUCED]: floating-point result reproduced by a verifier. [NUMERICAL, RUN BY THE VERIFIER]: run once, by a verifier. [NUMERICAL]: not reproduced. None is certified.
- [ARGUED]: a plausibility argument, not a proof. [OPEN]: no proof. [REFUTED]: shown false.
- [BALABAN]: an input taken from Balaban's published papers, used as stated there.
- [NOT WRITTEN]: a proof plan; no project file contains the proof.
- [CITED]: a standard result; its text was not re-read. [CITED, SOURCE CHECKED]: the source was read, the transfer to the lattice is standard and not proved here.
- [PROVED HERE]: a short proof given in full in this paper (Section 4). [HYPOTHESIS]: assumed, and owned by a named block. [FORMAL]: a formal step whose analytic justification is not written. [DERIVED SKETCH]: a route with steps listed, not a proof.
- [INTERVAL ENCLOSURE, VERIFIED]: a rigorous interval-arithmetic enclosure at a point, rerun by a verifier.

- [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]: a statement over a continuum of parameters (all K, all j), proved by an analytic majorant plus a branch-and-bound in interval arithmetic with outward rounding, and confirmed by an independent verifier file. It differs from [INTERVAL ENCLOSURE, VERIFIED] (one point) and from [NUMERICAL] (floating point, no rounding control).
- [VERIFIED REDUCTION]: a reduction confirmed by a verifier file; the inputs stay open.
- [PROVED MODULO NAMED BLOCKS]: proved given the hypotheses owned by the named blocks.

A compound tag such as [PROVED+VERIFIED, CONDITIONAL ON (H2)] combines these meanings. Tags refer to the project files cited next to them. In [VERIFIED QUOTE], a quote of a source other than Bałaban's papers was checked verbatim against a copy of that source in the project. Project files are cited as `NAME.md`; a section sign after a file name (for example `KAPPA_BOUNDED_VERIFY` §4) refers to a section of that file, not of this paper. "Grade n" and "item n" after a verifier file name refer to the numbered verdicts of that file. The paper as a whole is not verified as one document.

1.5 Relation to other work

Two recent works overlap with this paper. The record of how their content reached this project is given below; it is kept in the form fixed when the overlap was found.

Related work 1. E. Sussman and L. Alpöge (with Claude; README: "written largely by LLM ... Novel mathematical ideas should be attributed to Claude (under the guidance of LA)"), "The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0" (LaTeX, files `theorem_0_complete_account_raw_{master,I,II,III}.tex`, 424k lines), and "SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality" (`ym4_manuscript_zeroth_draft.pdf`, dated 6 October 2026) [26, 25]. Repository: `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit `1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310` (2026-10-27T05:45:28-05:00), accessed 2026-10-27. Repository created 2026-10-06. Disclaimer in README: "This repository contains work-in-progress, written largely by LLM. Its content has not been satisfactorily vetted. All claims should be understood as tentative, pending verification." Suggested keys: SA26-T0 (account), SA26-YM4 (draft).

Related work 2. OpenAI, "The canonical massive continuum limit of the two-dimensional O(3) model" (October 4, 2026), `github.com/openai/math`, family 215 (preprints folder; read locally from the PDF, `reports/ONLINE_SWEEP.md` §1.2; text in `reports/o3_continuum.txt`) [18]. Suggested key: OAI26-O3.

How content from these works reached this project. The record (file times, `PLAN.md`, read lists; `THEOREM2_PROOF_v2_VERIFY` §I) shows the following.

1. Section 3 was written without reading the Sussman–Alpöge repository. Only the citation block above (`reports/SUSSMAN_ALPOGE_MAP.md` §5.1) was read.
2. First channel. On the night of 2026-10-27, `reports/ONLINE_SWEEP.md` §2 (about 20:20) read the Sussman–Alpöge repository directly. It reported that they prove a weak two-sided flow with an additive constant, never need per-step positivity, use a first-passage cutoff, and invoke only containment $g_k \le \gamma$. This is the first project file with Sussman–Alpöge content.
3. Second channel. `reports/SUSSMAN_ALPOGE_MAP.md` (about 20:28) was written by a separate agent under a firewall. Through it, the bounded-κ idea entered the project: the weak form

of Theorem 2 needs only κ_k bounded, not convergence (their Lemma 4.19 structure; `PLAN.md` line 146). `KAPPA_BOUNDED.md` (about 20:39) received this idea through its prompt. Its read list excludes the map and ONLINE_SWEEP.

4. Lemma D′ and the cross-scale sweep predate any Sussman–Alpöge content in project files. The shooting-without-positivity lemma and the band with additive constant appear in `THEOREM2_PROOF_VERIFY` Point 5(b) (about 20:05) and `LEMMA_DPRIME.md` (about 20:10). `LEMMA_DPRIME_SWEEP.md` (about 20:16–20:23) overlaps ONLINE_SWEEP in time, but its read list is LEMMA_DPRIME and its verifier only; it did not read ONLINE_SWEEP. Lemma D″ is in `KAPPA_BOUNDED.md`, written after the map.

5. Theorem K. Its result form, $|S_k - ck \log L| \le \kappa_*$, matches Sussman–Alpöge Lemma 4.2(b), $|\sum_l \beta_l^0 - kc_0| \le c_2$ (as reported in `PLAN.md` line 146; cited there, not proved). The proof in this project (Prop. 1.1, Lemma 2.1, Prop. 2.2 of `KAPPA_BOUNDED`) was written by firewalled agents from Balaban's papers and project files.

6. O(3) paper. Route B's doubling comparison follows the template of OAI26-O3 Lemma 6.1 (doubling difference; `BETA_FP_PW` uses the "O(3) Lemma 6.1 form"). The idea of a coupling band in place of per-step positivity also appears there (ONLINE_SWEEP §1.2(a)). `BETA_FP_PW` read ONLINE_SWEEP §1.2, §5, §8 and o3_continuum Lemma 6.1, and did not read ONLINE_SWEEP §2.

7. Route B (other than the O(3) template), the numerics, and the symbol results of Sections 3.5.5 and 3.7 were derived by firewalled agents from Balaban's papers and project files.

In short: two agents read the Sussman–Alpöge repository, the web sweep (item 2) and the mapping agent (item 3), and neither wrote a derivation. The derivation agents were firewalled; their read lists exclude the repository, the map, and ONLINE_SWEEP §2. One idea crossed the firewall: the bounded-κ route reached `KAPPA_BOUNDED.md` through the map (item 3). Their public, timestamped account predates this one. Any write-up must cite SA26-T0, SA26-YM4, and OAI26-O3, and state these overlaps.

1.6 Organization

Section 2 fixes the notation for both parts of the paper. Section 3 proves the reduction of Theorem 2 (Theorem A). Section 4 proves the finite-torus existence theorem modulo named blocks (Theorem B) and contains the dependency table. Section 5 gives one paragraph for each verified source file of the sources block. Section 6 collects the numerical and certified results. Section 7 lists what remains. Appendix A lists the Lean formalization, Appendix B indexes the results by status, Appendix C maps the numbering of this paper to the names used in the project files, and Appendix D records the revision history.

2 Setup and notation

This section fixes the notation for the whole paper. Sections 2.1–2.4 serve the coupling flow of Section 3. Section 2.5 serves the existence theorem of Section 4. Section 2.6 lists the names that would otherwise collide.

2.1 Lattice, measure, and scales

Notation. L is odd, $L > 11$, fixed (Balaban, CMP 109 p. 256 [8]). $K \in \mathbb{N}$, $\varepsilon = L^{-K}$. The physical side $\ell \in \mathbb{N}$ is in units of the unit lattice. $T_\ell^4 = (\mathbb{R}/\ell\mathbb{Z})^4$ and $T_\varepsilon = (\varepsilon\mathbb{Z}/\ell\mathbb{Z})^4$, so $N_\varepsilon = \ell L^K$ sites per direction. The Wilson action is $A(U) = \sum_p P_p(U)$, $P_p(U) = 1 - \frac{1}{2}\operatorname{Re}\operatorname{tr} U(\partial p) \in [0, 2]$, over unoriented plaquettes p. The measure is $\mu_\varepsilon = Z_\varepsilon^{-1} \exp(-g_0^{-2} A(U)) \prod_b dU_b$ (Haar). x_p is the centre of p.

The bare coupling $g_0 = g_0(\varepsilon, g)$ is any solution of the shooting problem of Lemma 3.4 (Lemma D″) with terminal coupling $g_K = g$. Lemma D″ gives existence, not uniqueness. Any choice works below.

Volume hypothesis (Vol) (ledger C31). The torus side obeys $\ell > 22M$, where $M = L^{m_M}$ is the localization cube of CMP 109 p. 257 (the block size of ledger C24, not the M of Δ_M in Section 2.4). Justification: at every step $k \le K$ the local set $\tilde{\square}^5$, of side $11M$ level-k cells, must lift to $\mathbb{Z}^4$. Then the whole-lattice β subtraction leaves only Bałaban's irrelevant class. This needs $N_k > 22M$ level-k cells per side, and $N_k = \ell L^{K-k}$ is smallest at $k = K$ (Proposition 5.11). (Vol) does not depend on K. The proof of Theorem 2w does not use it; Theorem 2w ⇒ Theorem 1 (through ledger B8) and Theorem B use it. Even form, for Theorem B: $\ell = 2L^{m_T}$ with $m_T \ge m_M + 1$, which keeps ℓ even and gives $\ell \ge 2LM > 22M$, because $L > 11$. The power form of `THEOREM2_PROOF_v4.md` §1.1, $\ell = L^{N-K}$ with $N \ge K + m_M + 2$, also suffices, but it makes ℓ odd. CMP 109 prints both exponents m_T and m_M as m; here they are fixed independently.

2.2 Range of L

Balaban assumes L odd, $L > 11$, as a standing assumption for the whole of CMP 109 (pp. 251–252; OCR line 124, verbatim confirmed in `THEOREM2_PROOF_VERIFY` Step 6) [8]. His construction of $\beta_j(g)$ for $g > 0$ and his inductive assumption (H0) are therefore available only at odd $L > 11$. Every statement below that uses (H0) or $\beta_j(g)$ at $g > 0$ is stated for odd $L > 11$, or carries "(H0) and Balaban's construction at this L" as an explicit hypothesis. Statements about the one-loop quantities $\beta_j(0)$, S_k, and the symbols σ_j hold for odd $L \ge 3$.

2.3 The coupling sequence

$\varepsilon = L^{-K}$, so $\log(L^k\varepsilon)^{-1} = (K - k)\log L$. $u_k := 1/g_k^2$. Recursion (CMP 109 (0.18)/(0.20)): $u_{k+1} = u_k - \beta_{k+1}(g_k)$. The increment β_{k+1} depends on the whole history $g_0, \dots, g_k$ (CMP 109 p. 298, [VERIFIED QUOTE]). $\beta_j(0)$ is the one-loop (Gaussian) increment; it is history-independent by definition.

Shooting notation: $u_k = 1/g_k^2$, $t = 1/g^2$, $T(k, K) := \sum_{i=k+1}^{K} \beta_i(g_{i-1})$. Clamp $\varphi(u) := \max(u, 1/\gamma^2)^{-1/2}$.

Here γ is the upper bound for the couplings in Bałaban's Theorem 1. Bałaban's small constant β_0 (1/7 in CMP 122 II p. 389) is not a coupling (Section 3.1). In Lemma 3.6 (Lemma E) the symbols β_0, β_0' instead denote lower and upper bounds on $\beta_j(0)$, as in `LEMMAS.md`.

2.4 One-loop notation

The letter K has two uses. As an integer it is the number of RG steps, $\varepsilon = L^{-K}$. As a vector $K \in \mathrm{BZ} = [-\pi, \pi]^4$ it is a momentum in the Brillouin zone of the coarse unit lattice; then $k_l = (K + 2\pi l)/L$ are the fine momenta over K. The meaning is always clear from context. The other one-loop objects are defined where they are first used:

- S_k, the $p \to 0$ F^2-coefficient of the composed k-step one-loop integral, and its telescoping into $\beta_j(0)$: Section 3.5.1;
- $c = 11N^2/(12\pi^2)$ and $b := c \log L$: Section 3.5.3;
- the f^m Taylor symbols $\sigma_j^{(m)}(K)$ of the coarse kernel at level j, and d_j, κ_j, κ_W^F: Section 3.5.4;
- Δ_M (Gaussian RG kernel at block size $M = L^j$), Δ_∞, and the bordered matrix $\mathfrak{B}_j(K)$: Sections 3.5.5 and 3.9;
- the one-step minimizer map $H_j(K)$, ρ_1, ρ_2, and the inequality (H2): Section 3.7;
- the plane-wave quantities $\beta_{j+1}(p)$, Γ^{ex}, θ_* of Route B: Section 3.6.2.

2.5 Observables and Schwinger functions

Observable (class C1 of `SOURCES.md` §1.2). $\varphi_p := P_p - c_\varepsilon$, $c_\varepsilon := \langle P_p \rangle_{\mu_\varepsilon}$. By hypercubic invariance (Section 4.5.6) c_ε does not depend on p.

Test space. $\mathcal{D}_n := \{F \in C^\infty(T^{4n}) : \text{every derivative of } F \text{ vanishes on every diagonal } \{x_i = x_j\}, i \neq j\}$. The norm is $\|F\|_m := \max_{|\alpha| \le m} \sup |\partial^\alpha F|$. On the compact torus this is the Osterwalder–Schrader (OS) Schwartz seminorm without weights.

Lattice Schwinger functions (`SOURCES.md` §1.3). $S_{n,\varepsilon}(F) := \sum_{p_1,\dots,p_n} F(x_{p_1}, \dots, x_{p_n}) \langle \varphi_{p_1} \cdots \varphi_{p_n} \rangle_{\mu_\varepsilon}$, $F \in \mathcal{D}_n$. These are moments, not cumulants (`SOURCES_VERIFY` item 5(a)). No field renormalization factor is inserted: `SOURCES_VERIFY` item 8 gives $B_K = \prod c_j$ bounded above and below ($c_j = 1 + O(g_{j-1}^3)$, $\sum g_j^3 < \infty$), so a bounded factor B_K^n changes only the normalization of the limit.

Scaling check (`SOURCES_VERIFY` item 4, derived there). $\langle \varphi_p \varphi_{p'} \rangle \sim \varepsilon^8 g^4 |x_p - x_{p'}|^{-8}$ at tree level. So $S_{2,\varepsilon}(F) = O(1)$ for $F \in \mathcal{D}_2$ with no rescaling, while $\sum_p f(x_p)\varphi_p$ has variance $\sim \varepsilon^{-4}$. The limit object is a family of distributions off the diagonals, not a random field.

Extra hypothesis found while writing the source. ℓ is even (sufficient for the reflection argument; see Section 4.8, item 1). For a reflection of the periodic lattice to exchange two halves with both seams being link planes, N_ε must be even. With L odd, $N_\varepsilon = \ell L^K$ is even iff ℓ is even.

2.6 Names

Hypotheses are named by short labels in parentheses, such as (H2), (PI), or (FX); the dependency table (Section 4.6) lists each one with its status. Three names of the project files collide in one document and are changed here; Appendix C records the old names.

- The plane-wave theorem of `BETA_FP_PW` (its Theorem 3.2, called "Theorem B" there) is called Theorem $\mathrm{B_{PW}}$ (Theorem 3.9), so that "Theorem B" means the main Theorem B.
- The three global conditions of the dependency table, called (G1)–(G3) in `ASSEMBLY.md`, are called (GC1)–(GC3). The labels (G1) (constraint conditions of Proposition 3.7) and (G2) (per-cell infinite-volume densities) keep their meaning from the Theorem 2 files.
- The final sourced bound of `SOURCES_D4D5.md`, called (S) there, is called ($\mathrm{S_{src}}$). The label (S) means the smallness condition of Section 3.1.

3 The coupling flow and Theorem 2

This section proves Theorem A. Section 3.1 lists what Bałaban's proof of Theorem 1 uses of the coupling sequence; the answer is weaker than his (0.31). Section 3.2 states the weak form, Theorem 2w, that supplies exactly this. Sections 3.4–3.7 prove Theorem 2w from bounded window sums of the one-loop increments, give two routes to these window sums, and reduce the uniform part of both routes to the single inequality (H2). Section 3.8 assembles the result. Its output, the trajectory conditions (b), (c′), (S), is the block (B-T2) of Section 4.

3.1 What Bałaban's Theorem 1 actually needs

Source: `LEMMA_DPRIME_SWEEP` §2, §7. The statements of Theorems 1 and 2 are quoted in Section 1.2.

The quoted hypothesis of Theorem 1 is only $g_k \in]0,\gamma]$. The proof uses more. CMP 119 Theorem 1 (p. 262) assumes "(I.0.33)", a misnumbered reference to (0.31) of CMP 109. Its working form is CMP 119 p. 255–256 (2.6)–(2.9), for scales $m < n$:

- (2.6) $g_n \le (1 + g_n^2\beta'(n-m))^{1/2} g_m$ and $g_m \le (1+\beta_0) g_n$;
- (2.7) $(\log g_n^{-2})^p \le (1+\beta_0)(\log g_m^{-2})^p$ and $(\log g_m^{-2})^p \le (1 + g_n^2\beta'(n-m))^{\beta_0}(\log g_n^{-2})^p$;
- (2.8), (2.9) the same for $\varepsilon_j = g_j p_0(g_j)$ and R_j.

Here β_0 is Balaban's small constant (1/7 in CMP 122 II p. 389), not a coupling. [VERIFIED QUOTE] for (2.6)–(2.9): page images, `LEMMA_DPRIME_SWEEP` §1; the sweep itself is single-author, see Section 3.4.3.

In u-variables these follow from two pairwise bounds:

- (P-up) $u_m \le u_n + B'(n-m)$, with no additive constant;
- (P-low) $u_m \ge u_n + \beta(n-m) - D'$, with $\beta > 0$, $D' \ge 0$ fixed,

plus the smallness condition

- (S) $\gamma^{-2} \ge c_{sw} D'$, $c_{sw} = \max(2, 1/(1-(1+\beta_0)^{-2})) = 64/15$ at $\beta_0 = 1/7$, and $\gamma \le \gamma_*(\beta_0, \beta, B', p_0, p_1, r, \nu, \kappa_0)$ (the conditions that literal (0.31) also needs). γ_* is free of the D'-ratio conditions, but it depends on R through B' (it contains $\gamma^2 B' \le (1+\beta_0)^2 - 1$ with $B' = 5b/4 + 2R$). The order of constants in Section 3.8 (R before γ) handles this.

β_0 must be fixed before γ.

3.2 The weak form proved conditionally

Theorem 3.1 (Theorem 2w, weak form)**.** *Let $G = SU(2)$, $d = 4$, L odd, $L > 11$ (equivalently: L odd ≥ 3 with (H0) and Balaban's construction of $\beta_j(g)$ at this L as an explicit hypothesis). There are $\gamma > 0$, $g_1 \in]0,\gamma]$, $\beta > 0$, $B' \ge \beta$, $D' \ge 0$ such that for every $K \ge 1$ and every $g \in]0, g_1]$ there is a bare coupling $g_0 = g_0(K,g)$ with*

- *(b) $g_k \in]0,\gamma]$ for $0 \le k \le K$, and $g_K = g$;*
- *(c′) for all $0 \le m \le n \le K$: $u_n + \beta(n-m) - D' \le u_m \le u_n + B'(n-m)$;*
- *(S) holds for γ, D'.*

Consequence: Balaban's (0.31) holds up to the additive constant D' on the lower side, with constants $\beta/\log L$ and $B'/\log L$. The two-sided bound fixes $1/g_k^2$ up to constants, as required.

Status: [PROVED+VERIFIED, CONDITIONAL ON (B2) AND THE FIXED-LEVEL INPUTS OF SECTION 3.6] by Route K (Theorem K + Lemma D″). (B2) is reduced to the single inequality (H2) of Section 3.7 [PROVED+VERIFIED REDUCTION: `B2_TWOSTEP_VERIFY` GRADE 3]. Second route: Route B (`BETA_FP_PW` Theorem 3.2 + Lemma D″). The logic of Theorem $\mathrm{B_{PW}}$ is [PROVED+VERIFIED, CONDITIONAL] (`BETA_FP_PW_VERIFY` §6 items 1–4). Its inputs (U), (DF), (T1) from `BETA_FP_U` are [PROVED, UNVERIFIED]; the route is conditional on (U), ($B\theta$), and the plane-wave defects. `BETA_FP_U` reduces (U) to (T1) + (H2) [PROVED, UNVERIFIED]. Both routes also need (H0) at this L and L5 (Section 3.4.5). See Section 3.6.

Setting. The torus obeys the volume hypothesis (Vol) of Section 2.1: side $\ell > 22M$ final-scale cells, or the even form $\ell = 2L^{m_T}$, $m_T \geq m_M + 1$. The proof of Theorem 2w does not use (Vol), because Lemmas D′ and D″ use the whole-lattice β_j. (Vol) enters where Theorem 2w is applied on the torus: Theorem 1 through ledger B8, and Theorem B.

Trust boundary (round 7; Section 7.7). By Route K, Theorem 2w rests on one Bałaban-level item, A5, on the Group B items B1–B7, and on the external result X5. A15 is not in this set: Theorem 2w uses only the definition of β_j (Proposition 5.8 and the remark after it). The use of A5, item 7′, is a [VERIFIED REDUCTION] (Proposition H0-red of `SOURCES_T2_H0.md`) with residue N1–N9; its item N2, Lemma G3, is proved relative to listed inputs (Theorem 5.9). Round 6 removes the last bare assertion of A5: the existence of the whole-lattice limit of CMP 109 (1.22) at $g > 0$ now follows from Proposition 5.8 and Theorem 5.10. A5 stays in the set, because the rest of the H0-red residue is not proved. If one takes A5 through this reduction instead of Bałaban's assertion, Theorem 2w also rests on A7, A8 (Theorems 5.4, 5.5), A17, the inputs of Bałaban's sup induction, C9, the conditions C25, C26, and the locality conventions C32–C36. Round 7: of B1–B7, only B3 lacks a proof or a verified reduction; B4 and B7 are proved relative to listed inputs (Theorems 5.14 and 5.15).

3.3 Literal (0.31)

- (L1) Literal (0.31) for every $K \geq K_0$ and every $g \in]0, \gamma]$. Needs eventual uniform positivity: there are J and $\beta_* > 0$ with $\beta_j(0) \geq \beta_*$ for all $j > J$, plus $\sup_j |\beta_j(0)| < \infty$ (Lemma 3.2 (d)). [PROVED+VERIFIED; LEAN] as an implication. Theorem K and Theorem $\mathrm{B_{PW}}$ do NOT give eventual positivity: they bound window sums, not single increments. Eventual positivity follows from the convergence route (Section 3.9), which is open.

- (L2) Literal (0.31) for every $K \geq 1$. Needs $\inf_j \beta_j(0) > 0$ and $\sup_j \beta_j(0) < \infty$ (Lemmas D + E). If $\beta_j(0) \to \beta_\infty > 0$, then "every $\beta_j(0) > 0$" is equivalent to $\inf_j \beta_j(0) > 0$ (Lemma D′ Step 8). [PROVED+VERIFIED; LEAN] as an implication. Inputs open.

- Literal (0.31) is not needed for Theorem 1: Section 3.1 shows the proof uses (c′) + (S). Literal (0.31) is used directly in one place only, the alternative choice of ε_0 in CMP 119 p. 248, which Balaban does not adopt.

- Necessity. Literal (0.31) at every $K \geq 1$ forces $\beta_K(0) \geq \beta \log L > 0$ for every K (window $k = K - 1$, let $g \to 0$ using (H0); so L odd > 11). [PROVED, UNVERIFIED]; elementary; `THEOREM2_PROOF_VERIFY` Point 5(a) agrees. So certified early steps are needed only for (L2), not for Theorem 1.

3.4 Shooting lemmas

The shooting lemmas turn bounds on the increments β_j into a trajectory with prescribed terminal coupling. Lemma 3.2 needs eventual positivity of $\beta_j(0)$; Lemma 3.4 needs only bounded window sums, and is the one used for Theorem 2w. The notation is that of Section 2.3. Lemmas D′, D″, D, E are abstract statements about a sequence of functions β_j. Applied to Balaban's $\beta_j(g)$, they need L odd > 11 (or (H0) at this L) and L5.

3.4.1 Lemma D′ (shooting without early positivity) [PROVED+VERIFIED; LEAN]

Sources: `LEMMA_DPRIME.md` §1–§2; `LEMMA_DPRIME_VERIFY.md` grades 1, 2, 5 (CONFIRMED). Step 7 of that file also holds grades 3, 4, 6, which are CORRECTED (Remark 1 overclaim, misreads in §5, ledger); this chapter uses none of them except through the sweep (Section 3.4.3). Lean: `lean/YmLean/LemmaDPrime.lean` (`YmLean.lemma_DPrime`, 405 lines), statement check `lean/VERIFY_LemmaDPrime.md`.

Lemma 3.2 (Lemma D′)**.** Hypotheses. *(i)* β_{k+1} *continuous on* $[0,\gamma_0]$ *and* $|\beta_{k+1}(g)-\beta_{k+1}(0)| \le C_1 g$, C_1 *independent of* k, K. *(ii)* $\sup_j |\beta_j(0)| \le B$. *(iii′) there is* J *with* $\beta_j(0) \ge \beta_\infty/2$ *for all* $j > J$ *(this is all the proof uses of "*$\beta_j(0) \to \beta_\infty > 0$*").*

Constants. $\gamma := \min(\gamma_0, \gamma_{Bal}, \beta_\infty/(4C_1))$; $B' := B + \beta_\infty/4$; $D := J \cdot B'$; $D' := J(B + \beta_\infty/2)$; $g_1 := (1/\gamma^2 + D)^{-1/2}$; $K_0 := J + \lceil 8D'/\beta_\infty \rceil$.

Conclusions.

- *(a)* $T(k,K) \ge -D$ *for all* $k < K$ *along any couplings in* $]0,\gamma]$.
- *(b) For every* $K \ge 1$ *and* $g \in]0,g_1]$ *there is* g_0 *with all* $g_k \in]0,\gamma]$ *and* $g_K = g$.
- *(c)* $u_K + (\beta_\infty/4)(K-k) - D' \le u_k \le u_K + B'(K-k)$. *The upper side has no constant.*
- *(d) For* $K \ge K_0$*: literal (0.31) with lower constant* $\beta_\infty/8$*, and shooting for every* $g \in]0,\gamma]$. *If every* $\beta_j(0) > 0$*: literal (0.31) for all* K *(Step 8 = Lemmas D + E).*

Proof idea. Clamped flow $\tilde{u}_{k+1} = \tilde{u}_k - \beta_{k+1}(\varphi(\tilde{u}_k))$ is continuous in u_0. Its tail sums lie in $[-D, B'(K-k)]$ for every u_0, because clamped couplings lie in $]0,\gamma]$. IVT on $[t-D, t+B'K]$ gives $\tilde{u}_K = t$. Then $\tilde{u}_k \ge t - D \ge 1/\gamma^2$ for $g \le g_1$, so the clamp is never active. The bound is derived for the clamped flow first; there is no circularity. For history-dependent β this proof needs β_{k+1} defined and continuous on the whole cube $]0,\gamma]^{k+1}$; see L5 in Section 3.4.5.

Lean. The Lean hypotheses are weaker (ContinuousOn instead of C^1; (iii′) instead of convergence). Step 8 is `lemma_DPrime_step8` (`lean/YmLean/LemmaDE.lean`). Lean covers one-argument $\beta_j(g)$ only, not the history form.

Remark 3.3 (Remark 4 of `LEMMA_DPRIME.md`). Without $g \le g_1$, (b) can fail at small K ($\beta_1 = \beta_2 = \beta_3 = -0.8$, $K = 3$, $g = \gamma$). [PROVED+VERIFIED]

3.4.2 Lemma D″ (tail-sum form; no index J) [PROVED+VERIFIED; LEAN]

Source: `KAPPA_BOUNDED.md` §5.3; `KAPPA_BOUNDED_VERIFY` §6 (CONFIRMED: "No J needed"); hand rederivation in `THEOREM2_PROOF_v2_VERIFY` §A.

Lemma 3.4 (Lemma D″)**.** Hypotheses. *(i) as in 2.1, and the window bound*

$$(5.1)\quad T_0(k,K) := \sum_{i=k+1}^{K} \beta_i(0) \in [\, b(K-k) - 2R,\ b(K-k) + 2R\,] \quad \textit{for all } 0 \le k < K,$$

with $b := c \log L > 0$ and some $R \geq 0$. (Route K gives $R = \kappa_*$; Route B gives $R = C_B$.)

Constants. $\gamma := \min(\gamma_0, \gamma_{Bal}, b/(4C_1))$; $\beta := 3b/4$; $B' := 5b/4 + 2R$; $D = D' := 2R$; $g_1 := (1/\gamma^2 + D)^{-1/2}$.

Conclusions. *(b) of 2.1, and the pairwise bound*

$$(c') \quad u_n + \beta(n-m) - D' \leq u_m \leq u_n + B'(n-m), \quad 0 \leq m \leq n \leq K,$$

upper side without constant.

Proof. $|T - T_0| \leq C_1\gamma(K-k) \leq (b/4)(K-k)$. Add (5.1). Steps 2–5 of Lemma D′ use only continuity and the bracket $\tilde{T} \in [-D, B'(K-k)]$; this bracket now comes from (5.1). The pairwise form is the same window argument with $K \to n$. The proof is the clamped-flow proof of 2.1; for history-dependent β the L5 caveat of Section 3.4.5 applies.

Lean: `lean/YmLean/LemmaDDoublePrime.lean` (`YmLean.lemma_DDoublePrime`, 146 lines, no `sorry`, axioms propext, Classical.choice, Quot.sound; `lean/README.md`). It proves (a″), (b) with the clamp inactive on the solution, (c′) with $\beta = 3b/4$, $B' = 5b/4 + 2R$, $D' = 2R$, and (d) below. The Lean hypotheses are weaker than the note: continuity instead of C^1; any γ with $0 < \gamma \leq \gamma_0$ and $C_1\gamma \leq b/4$; $R \geq 0$ derived; $K = 0$ allowed; (c′) and (d) hold for every trajectory with couplings in $]0, \gamma]$. One-argument β only (L5 not formalized).

(d) [LEAN]: for $n - m \geq 8R/b$, $u_n + (b/2)(n-m) \leq u_m \leq u_n + B'(n-m)$. So literal (0.31) from (5.1) alone holds, with lower constant $b/2$, only for windows with $n - m \geq 8R/b$; short windows near K are not controlled (see Section 3.3).

3.4.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [PROVED, UNVERIFIED (SWEEP); 4 OF 19 TABLE ROWS VERIFIED]

Source: `LEMMA_DPRIME_SWEEP.md` (all 12 Balaban files; OCR grep plus page image for every non-single-scale hit). `LEMMA_DPRIME_VERIFY` Steps 4–5 checked comparisons A–E on page images (CMP 122 I p. 192, p. 198 (1.94); II p. 381 ×2, p. 386) and corrected two misreadings of A (factor $(1+\beta_0)^2(k-j)^{\beta_0}$; $\delta'_j = g_j A_1 p_1(g_j)$). The five comparisons map to four table rows: A = row 11, E = row 12, C and D = row 16, B = row 18.

Result. 19 cross-scale uses of the coupling sequence (table in SWEEP §7), all in CMP 119 and CMP 122 I–II; none in CMP 95, 96, 98, 99, 102, 109, 116. Each is covered by (b) + (c′) + (S) of Section 3.1:

- items 1, 7, 10, 12, 17 need (P-up) with no constant (e.g. (2.6) line 1 is exactly (P-up));
- items 2, 3, 11, 13, 15, 16, 18 need (P-low) with fixed multiplicative slack $(1+\beta_0)$ or $(1+\beta_0)^2$; item 2 ((2.6) line 2) gives the constant 64/15; item 11 needs only $2.42D'$;
- items 4 ((2.8)–(2.9), ε and R ratios), 6 (CMP 119 (3.7), (3.8), (3.12), (3.13)), and 9 (CMP 122 I (1.52)), the last two of the form $\varepsilon_{k+1} \leq (1+\beta_0)\varepsilon_k$, use both sides; they follow as items 1 and 3 do (item 4 from items 1–3);
- item 14 (CMP 122 I p. 200, R_{k-N_0+1} vs R_k) uses the lower side; it follows as item 3 does;
- item 5 (CMP 119 (2.31), (2.46): $\sum_j g_j^{\kappa_0} \leq g_n^{\kappa_0 - 6}$) needs linear growth (P-low) and $\gamma^{-2} \geq 2D'$;
- item 0 is CMP 119 Theorem 1 assuming (0.31); its proof works with (2.6)–(2.9);
- item 8 (CMP 119 p. 284) is in a sketch, not load-bearing.

Caveats [OPEN]: (1) the sweep is a grep plus page images, not a line-by-line read; (2) "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) was not located; (3) Balaban says β_0 "can be chosen arbitrarily small, if g is sufficiently small"; under (P-low) the floor is $\beta_0 \gtrsim D\gamma^2/2$, so β_0 must be fixed first; no step was found that needs $\beta_0 \to 0$; (4) the sweep has no independent verifier beyond rows 11, 12, 16, 18; (5) `BALABAN_EXCERPTS_CHECK` item 11 lists readings not checked on page images: II (4.37)/(5.1), CMP 119 (2.42), the (2.12)/(2.23) ambiguity, the "L odd" page, the Sect. 3 estimates, and the 1989 bound lists.

3.4.4 Lemma D and Lemma E [PROVED+VERIFIED; LEAN]

Sources: `LEMMAS.md` (corrected), `LEMMAS_DE_VERIFY.md` §1.4, §1.5, §2.4; `lean/YmLean/LemmaDE.lean` (`lemma_D`, `lemma_E`, `lemma_DE`).

Lemma 3.5 (Lemma D)**.** *If $b \le \beta_j(g) \le b'$ on $]0,\gamma]$ with $b > 0$, then for every K and $g \in]0,\gamma]$ shooting succeeds and $t + b(K-k) \le u_k \le t + b'(K-k)$.*

This is literal (0.31). An earlier draft used a wrong first IVT interval, which the verifier refuted; the corrected interval is $[t+bK,\ t+b'K]$. The nested-interval variant for β defined only on admissible histories (`LEMMAS_DE_VERIFY` §1.5) is [PROVED+VERIFIED], not in Lean.

Lemma 3.6 (Lemma E)**.** *If $\beta_0 \le \beta_j(0) \le \beta_0'$ and $|\beta_j(g) - \beta_j(0)| \le Cg^2$, then for $\gamma \le \sqrt{\beta_0/(2C)}$: $\beta_0/2 \le \beta_j(g) \le \beta_0' + \beta_0/2$.*

The verifier checked this Cg^2 form; the C^1 form ($C_1 g$, $\gamma \le \beta_0/(2C_1)$) is the same two-line proof [PROVED, TRIVIAL VARIANT]. Lean has the Cg^2 form.

3.4.5 History dependence (L5) [PROVED RELATIVE TO LISTED INPUTS; VERIFIED (ROUND 7)]

Round 7 closes this item by the variant of the third bullet below: nested intervals on true trajectories, with no clamp and with the constants of Lemma 3.4 (Theorem 5.15, `SOURCES_T2_B7_CLAMP.md` Route 2 + VERIFY). The bullets record the gap as it stood before.

Balaban's β_{k+1} depends on $(g_0, \dots, g_k)$. Lemma D′ Remark 3 (`LEMMA_DPRIME.md`) states the cube-continuity version (same proof). The gap is as follows (`THEOREM2_PROOF_v2_VERIFY` §F).

- *Domain.* Balaban defines β_{k+1} only on histories for which his Theorem 1 induction has run to step k (`LEMMAS_DE_VERIFY` §2 item 1). By the sweep (§2.3), that induction uses (2.6)–(2.9), that is (c′) + (S), on the history, not only $g_i \in]0,\gamma]$.

- *The new gap.* The clamped shooting of Lemmas 3.2 and 3.4 evaluates β on clamped histories $\tilde{g}_i = \varphi(\tilde{u}_i)$. A clamped history can violate the growth side of (c′): from $\max(a,c) - \max(b,c) \ge \min(a-b, 0)$ one gets only $u_m \ge u_n - D'$, not $u_m \ge u_n + \beta(n-m) - D'$. So the clamped shooting evaluates β on histories outside Balaban's domain. The clamp proof is valid only under the cube hypothesis, which Balaban does not supply.

- *A variant that avoids it* [NOT WRITTEN]. Use a nested-interval proof in the style of `LEMMAS_DE_VERIFY` §1.5, with domain $D_k = \{u_0, \dots, u_k \ge \gamma^{-2}\}$ of true (unclamped) trajectories. On such trajectories, (c′) up to step k follows from the tail-sum bound (a″) of Lemma D″, and (S) holds by the choice of γ. So every evaluation of β stays inside Balaban's domain. At each step k, keep the set of u_0 whose trajectory stays in D_k; show by IVT, step by step, that this set contains a closed interval on which u_k sweeps a range that contains the

next target interval. For Lemma D (all increments positive) this is written. For Lemma D″ the increments can be negative, so the nested intervals need new envelopes, built from the window bounds $[b(n-m) - 2R - (b/4)(n-m), \dots]$. The adaptation is plausible; it is not written and not verified.

- Lean is one-argument only.

3.5 The exact structure of the one-loop sums

The window sums of the shooting lemma are sums of one-loop increments $\beta_j(0)$. This section expresses these sums as level-k quantities of a coarse Gaussian theory and isolates the remainder κ_k whose boundedness is the content of Theorem K. It also lists the symbol facts, uniform in the level, that Section 3.6 uses.

3.5.1 Telescoping [PROVED+VERIFIED, CORRECTED SCOPE]

$S_k := p \to 0$ F^2-coefficient of the composed k-step gauge-fixed one-loop integral (U1 (1.2)). Then $S_k = \sum_{j=1}^{k} \beta_j(0)$, $S_0 = 0$ (`BETA_FP` Prop. 1.1; `BETA_FP_VERIFY` item 1). Scope: exact for the $p \to 0$ coefficient at fixed k, not for $\log Z_k$ at fixed p (the plane wave is off-shell).

3.5.2 Anchor-free identity [PROVED+VERIFIED]

Source: `BETA_FP_LEMMA43` Prop. 2.1; `BETA_FP_LEMMA43_VERIFY` items 1, 3–4 (hand rederivation; lattice check with the real CMP 109 averaging at random SU(2) backgrounds, relative difference 10^{-14}).

Proposition 3.7 (anchor-free identity; Prop. 2.1 of `BETA_FP_LEMMA43`)**.** *Under (G1) = $\{\Delta D = 0,\ \tilde{Q}D = D_c \iota_a^T\}$:*

$$\log Z_\Lambda(V) = \big[\log\det X - \tfrac{1}{2}\log\det K\big] - \big[\log\det(D_c^T D_c) - \tfrac{1}{2}\log\det(\Delta_\Lambda(V) + D_c D_c^T)\big].$$

In words: constrained one-loop = fine Wilson one-loop in background Feynman gauge minus the one-loop of the coarse Gaussian theory with kernel $\Delta_\Lambda(V)$. All point-anchor objects cancel; the constant is exactly 0.

- $\tilde{Q}D = D_c \iota_a^T$ is exact at any background (lattice 10^{-15}). $\Delta D = 0$ is load-bearing (raw Hessian gives $O(1)$ errors).
- At the constant abelian field V_f the background is critical, so $\Delta D = 0$ is exact and Prop. 2.1 holds exactly (`BETA_FP_H_VERIFY` §2, CONFIRMED). At a plane wave it holds up to the defect $\mathfrak{d}_Y$ (`BETA_FP_PW` Lemma 2.3) [PROVED, UNVERIFIED].
- Per-cell infinite-volume densities only: on a torus the constant zero modes must be removed. This is input (G2) [VERIFIED REDUCTION AT FIXED j] (`SOURCES_T2_G2.md` + VERIFY).

3.5.3 The constant c [PROVED+VERIFIED]

$c = 11N^2/(12\pi^2) = 2N \cdot b_0/(8\pi^2)$, $b_0 = 11N/3$; SU(2): $c = 0.371511$. Normalization $1/g_B^2 = 2N/g^2$ (`R2_VERIFY`; `BETA_FP_LEMMA43_VERIFY` §5). The log coefficient of the fine piece at constant field is $c_2 = c/2$ per $\ln(1/\varphi)$, checked against Savvidy, arXiv:1910.00654 eq. (1.4) [23] (`BETA_FP_H_VERIFY` §4) = input (F) [CITED, SOURCE CHECKED; LATTICE = CONTINUUM LOG COEFFICIENT IS STANDARD, NOT PROVED IN PROJECT FILES].

$b := c \log L$: 0.25751 ($L = 2$), 0.40815 (3), 0.51502 (4), 0.59793 (5), 0.72293 (7).

3.5.4 The decomposition $S_k = ck \log L + \kappa_W^F - \kappa_k$

- As bookkeeping the split is empty: any c' gives an exact identity if κ_k is defined as the remainder (`U1_VERIFY`). Boundedness of a remainder-defined κ_k is a restatement, not progress.

- Content comes from defining κ_k by the coarse theory alone. `KAPPA_BOUNDED` Prop. 1.1 [PROVED+VERIFIED, CONDITIONAL ON (PI), (F), (M_j), V1, (G2)]: with the Wilson level $j = 0$ as reference,

$$S_j = cj \log L - d_j, \qquad \kappa_j = \kappa_W^F + d_j,$$

$$d_j := -\frac{1}{2n_2} \int_0^1 dt \int_{\mathrm{BZ}} \frac{dK}{(2\pi)^4} \operatorname{tr}\big[b_{t,2}\, \delta\sigma_j^{(0)} + b_{t,1}\, \delta\sigma_j^{(1)} + b_{t,0}\, \delta\sigma_j^{(2)}\big],$$

  where $\delta\sigma_j^{(m)} = \sigma_j^{(m)} - \sigma_0^{(m)}$ are differences of the f^m Taylor symbols of the coarse kernel at level j and level 0, and $b_{t,m}$ are the Moyal inverse coefficients of the interpolated coarse operator. No limit symbols σ_∞ are used.

- Checks [NUMERICAL, REPRODUCED]: $\rho(1) = \kappa_W^F$ (Lüscher–Weisz (5.7) [16]); $L = 2$: $d_1 = -0.372$, $\kappa_1 = 1.114$; $L = 3$: $d_1 = c \log 3 - \beta_1 = -0.256 \pm 0.01$ (`KAPPA_BOUNDED_VERIFY` §4: fits give $-0.254 \cdots - 0.258$). This checks the definition of d_j through S_j; it does not evaluate (1.1) from the symbols independently.

- Cor. 3.1 of `BETA_FP_LEMMA43` ($\mathcal{C}(\Lambda) = c \log \Lambda + \kappa_W^F - \rho(\Lambda)$) is exact as an identity. "The whole $\log \Lambda$ sits in the fine piece" is equivalent to $\rho(\Lambda) = o(\log \Lambda)$. It is not a separate fact (`BETA_FP_LEMMA43_VERIFY` item 2; an earlier overclaim is removed).

- Given convergence $\beta_j(0) \to \beta_\infty$, Cesàro gives $\beta_\infty = c \log L - \lim \kappa_k/k$. So the hypothesis (H′) "$\kappa_k = o(k)$" of the first version of these notes is EQUIVALENT to $\beta_\infty = c \log L$, not weaker (`THEOREM2_PROOF_VERIFY` Point 4). Theorem K (Theorem 3.8) gives κ_k bounded, hence $\kappa_k = o(k)$ trivially.

3.5.5 Level-uniform symbol facts used below [PROVED+VERIFIED]

Source: `U3_AW.md`, `U3_AW_VERIFY.md` (grades 1–4 CONFIRMED); `U3_REPAIRS_VERIFY`.

- O5/A0: Δ_M real-analytic on a fixed strip, $|\partial^\alpha(\Delta_M - \Delta_\infty)| \le CM^{-2}$, uniform to $K = 0$ (odd L, centered blocks). Prop. B: $c_0|\hat{K}|^{-2}P_T' \le P_T' S_M P_T' \le C_1|\hat{K}|^{-2}P_T'$, $c_0 = (2/\pi)^{12}$, $C_1 \le 1363$. Existential constants.

- Prop. E (evenness): $\sigma_j(f; -K) = \sigma_j(f; K)$ in the midpoint trivialization, all f, j. Tested in $d = 2$ only.

- Theorem W: $\sigma_j^{(1)}(0) = \tau_c = (3ie/2)(E_{10} - E_{01})$ for every $j \ge 0$, from the exact Ward identity $\Delta_j(f) D_f = 0$. Numerically 0.750000, 0.75015, 0.75051 at $j = 0, 1, 2$ vs 0.75 ($d = 2$ proxy).

- Theorem A3: $|\sigma_j^{(1)}(K) - \tau_c| \le C_A |K|^2$, real K, uniform in j (C^0 only).

- Prop. A2: $\sigma_j^{(m)}$, $m \le 2$, analytic on the same strip at every fixed j (not uniform in j for $m = 1$ derivatives, $m = 2$).

- Lemma M: Maxwell coefficient exactly 1 at every level.

3.6 Bounded window sums: two formally distinct conditional routes

This section gives the two routes to the window bound (5.1) that Lemma 3.4 needs.

Both routes give the window bound (5.1) of Lemma 3.4, $|S_K - S_k - b(K-k)| \le 2R$, with $b = c \log L$. Lemma D″ then gives Theorem 2w (at odd $L > 11$, or with (H0) at this L as a hypothesis, and with L5). Neither route uses convergence of $\beta_j(0)$, a rate, (N), or certified early steps. The two routes have formally distinct named hypotheses. Their mathematical content is not independent: both reduce their uniform-in-level content to the same inequality (H2) on the two-step minimizer map (Section 3.7), plus A3-type facts (`BETA_FP_U` §3.5, §4.3 [PROVED, UNVERIFIED]; Route K side: `B2_TWOSTEP_VERIFY` grade 3 [PROVED+VERIFIED]).

3.6.1 Route K: Theorem K [PROVED+VERIFIED, CONDITIONAL ON (B2) AND THE FIXED-LEVEL INPUTS]

Source: `KAPPA_BOUNDED.md`; `KAPPA_BOUNDED_VERIFY.md` (items (i)–(v) and (vii) CONFIRMED; (vi) CORRECTED in reasoning, conclusion unchanged).

Theorem 3.8 (Theorem K)**.** *L odd ≥ 3. Assume (B2) (Section 3.7; (B2) ⇐ (H2) [PROVED+VERIFIED]) and the fixed-level inputs below. Then $\sup_k |d_k| \le \kappa_* < \infty$, so $\sup_k |\kappa_k| \le |\kappa_W^F| + \kappa_*$ and $|S_k - ck \log L| \le \kappa_*$ for all k. Window sums satisfy (5.1) with $R = \kappa_*$. (Theorem K is a statement about one-loop quantities. Its use in Lemma D″ for Balaban's $\beta_j(g)$ needs L odd > 11 or (H0) at this L.)*

Proof structure.

1. Prop. 1.1 (Section 3.5.4): $S_j = cj \log L - d_j$. [PROVED+VERIFIED, CONDITIONAL ON (PI), (F), (M_j), V1, (G2)]
2. Lemma 2.1: the only K-derivative of $\sigma^{(1)}$ in d_j enters linearly with f^0 coefficients. Integration by parts moves it onto f^0 data; boundary terms vanish (periodic up to a common constant conjugation; inner sphere $O(r^2)$). So $\sigma^{(1)}$ is needed only in C^0, which Theorem A3 gives. [PROVED+VERIFIED]
3. Prop. 2.2: with (E0) f^0 bounds (Prop. B, O5/A0, Lemma M), (E1) Theorem A3 + Theorem W, and (B2), every term of the d_j integrand is $\le C|K|^{-2}$, uniformly in j and t. In $d = 4$ this has two powers of infrared margin. $|d_j| \le \kappa_* := (C(L)/n_2)(1 + Y_1^2 + C_A(1 + Y_1) + Y_2) \cdot I_2$. [PROVED+VERIFIED, CONDITIONAL ON (B2)]
4. Infrared: the degree -4 part of the coarse f^2 integrand is the same at every level (Lemma M + Theorem W), so the $\tau_c \log$ cancels exactly in level differences. [PROVED+VERIFIED]

Constants are existential; κ_* is not evaluated.

Fixed-level inputs of Route K (needed at each j separately; no uniformity in j):

- (PI) probe independence: plane-wave p^2 coefficient = constant-field f^2 coefficient [ARGUED; PER-CELL LIMIT FOR THE LINEAR POTENTIAL NOT WRITTEN].
- (F) fine log coefficient c_2 at constant field [CITED, SOURCE CHECKED], Section 3.5.3.
- (M_j) the formal f^2 coefficient d_j equals the true one at fixed j. The resolvent step of the first argument was false (Nielsen–Olesen zero at $k_\perp^2 = |ef|$; `BETA_FP_H_VERIFY` §5). `BETA_FP_M` repairs it with a direct $\log|\det|$ bound given spectral bounds (S1)–(S3) [OPEN ON THE LATTICE]; Prop. 4.4 there is [ARGUED]; `BETA_FP_M` has no verifier file. The t-subdivision of

KAPPA §1.3 removes the small-ε requirement [ARGUED, CONFIRMED AS PLAUSIBLE]. Status: [ARGUED].

- V1: the Maxwell coefficient of the per-step determinant equals $\beta_{j+1}(0)$, i.e. order-g^0 completeness of CMP 109 (1.3)–(1.5) [PROVED+VERIFIED AT FIXED LEVEL, CONDITIONAL ON (G2), BAŁABAN'S INDUCTIVE BOUNDS, AND THE AVERAGING (0.4)] (`SOURCES_T2_V1.md` + VERIFY; Section 5.1).

- (G2) per-cell infinite-volume densities [VERIFIED REDUCTION AT FIXED j] (`SOURCES_T2_G2.md` + VERIFY; was [FORMAL]); the j-uniform version is not proved.

- (H0) Balaban's inductive assumption: β_{j+1} smooth on $[0, \gamma]$, "uniformly bounded on this interval together with all derivatives" (CMP 109 pp. 263–264, (1.22)) [BALABAN; AVAILABLE AT ODD $L > 11$ ONLY]. CMP 116 has no theorem statement and no C^m-in-g bound in words (`BALABAN_EXCERPTS_CHECK` item 10); the smoothness in g is CMP 109's inductive assumption, asserted by CMP 109 Theorem 3, whose proof CMP 116 completes.

- L5 history-dependent β, its domain, and its continuity (Section 3.4.5) [PROVED RELATIVE TO LISTED INPUTS; VERIFIED (ROUND 7)].

3.6.2 Route B: Theorem $\mathbf{B}_{\mathrm{PW}}$ [PROVED+VERIFIED, CONDITIONAL (LOGIC OF THEOREM B_{PW})]

Source: `BETA_FP_PW.md` §1–§5 (Theorem 3.2 = Theorem B_{PW}, Theorem 5.1). `BETA_FP_PW_VERIFY.md` §6 grades it: items 1–4 CONFIRMED (doubling identity and Lemma 1.3; Prop. 2.2 / Lemma 2.3; Theorems A, B as conditional logic; Lemma 4.1, conditional), items 5 (comparison) and 6 (numerics) CORRECTED. The tag follows the decision recorded in `PLAN.md` (2026-10-28): the logic of Theorem B_{PW} is [PROVED+VERIFIED, CONDITIONAL] on its hypotheses, citing `BETA_FP_PW_VERIFY` §6 items 1–4. Its inputs (U), (DF), (T1) from `BETA_FP_U` stay [PROVED, UNVERIFIED]. The file rederives the Lemma 1.3 arithmetic and the Lemma 2.3 / Prop. 2.2 algebra correctly, confirms the Prop. 2.2 determinant identity on a 4^4 lattice (4e−13), and notes that identity 3.0 is a tautology ($\Gamma^{ex} := \hat{\Pi} - S$), so all content sits in $(B\theta)/(C\theta)$.

Theorem 3.9 (Theorem B_{PW}; Theorem 3.2 of `BETA_FP_PW`)**.** *L odd ≥ 3. Assume*

- *(U) there are $p_* > 0$, C_U with $|\beta_{j+1}(p) - \beta_{j+1}(0)| \leq C_U p^2$ for all $j \geq 0$, $0 < p \leq p_*$* [HYPOTHESIS; ARGUED FROM UNIFORM ANALYTICITY IN p OF THE ONE-STEP PLANE-WAVE DATA PLUS ONE-STEP O6]*;*

- *(Bθ) there is $\theta_* \in (0, \min(p_*, 1))$ with $B_* := \sup_k |\Gamma^{ex}(\theta_*; L^k)| < \infty$, where Γ^{ex} is the coarse remainder at fixed coarse momentum θ_** [HYPOTHESIS]*;*

- *Lüscher–Weisz (5.7) [16]* [CITED].

Then $|S_k - ck \log L| \leq C_B := B_* + c_2 \ln(1/\theta_*^2) + |\kappa_W^F| + \sup_{p \leq \theta_*} |\eta(p)| + C_U \theta_*^2/(L^2 - 1)$, and window sums satisfy (5.1) with $R = C_B$. $\theta_* < 1$ is required so that $c_2 \ln(1/\theta_*^2) \geq 0$; otherwise the term must be replaced by its absolute value (`BETA_FP_PW_VERIFY` §3).

Key step. Lemma 1.3: $|S(\theta; L^k) - S_k| \leq C_U \theta^2/(L^2 - 1)$ from (U). The massless coarse side is then needed only at a fixed $\theta_* > 0$, where there is no infrared problem. No (W), (PI), (M), IR weights, or spectral theory.

What ($B\theta$) reduces to (`BETA_FP_PW` §4) [ARGUED]: Lemma 4.1(i) (trivial-background convergence, from O5, odd L) [PROVED]; vertex bounds at fixed θ_* (the uniform bounds cited by U3 Theorem 7.1 to Balaban; the fixed-θ data hypothesis ($D\theta$) is needed only in its "bounded" form); and boundedness in k of the defect $\mathfrak{d} = \mathfrak{d}_Y + \mathfrak{d}_{gf}$.

- $\mathfrak{d}_Y$ (Prop. 2.1 defect at the non-stationary plane wave): $O(a\Lambda^{-2})$ relative at fixed θ, from zero capacity of point anchors in $d = 4$ [ARGUED; REGION $|K| \lesssim \Lambda^{-1}$ NOT WRITTEN].
- $\mathfrak{d}_{gf}$ (composite gauge vs D_k^T gauge at the plane wave): $O(\theta^2)$ relative; uniformity in k not shown [ARGUED]. Data: 0.953 (direct) vs 0.9595 (recursion) at $L = 2$, $p_0 = 0.2$; difference below the tabulation error, so unresolved.

Identification. Route B's $\beta_j(0)$ is the per-level plane-wave increment (the quantity the code computes). It matches Balaban's second-moment β up to the expansion point (fine wave vs constrained minimizer), argued to change S by $O(\theta^2)$ relative (`SOURCES_D1_VERIFY` item 8) [ARGUED]. This identification and the $\mathfrak{d}_{gf}$ bound need expansion-point bounds uniform in k. So the (PI)-type content of Route K is not removed in Route B; it moves into a uniform-in-k input (`BETA_FP_PW_VERIFY` §3).

Three points from `BETA_FP_PW_VERIFY` §3 (item 5 of its §6 grades them CORRECTED):

1. $\theta_* < 1$ is required (above).
2. The β identification (B4) is a uniform-in-k input, not a fixed-level input.
3. (U) is a uniform-in-j analyticity of the level data. It plausibly needs the plane-wave analogue of a uniform C^1 bound on $\sigma^{(1)}$: uniform p-smoothness of the a^1 vertex. That is the same uniform first-order vertex smoothness as Route K's open core ($C1\sigma$).

3.6.3 Route B inputs after `BETA_FP_U` [PROVED, UNVERIFIED; NO VERIFIER FILE]

Source: `BETA_FP_U.md` §2–§4. It reduces the two uniform hypotheses of Theorem B_{PW}.

- (U) $\Leftarrow$ (T1) + (H2) (Theorem 2.4, odd $L \geq 3$, with A0, A1, Lemma 2.2). (T1): $\sup_j \sup_q |c^{(1)}_{T,j}(q)| < \infty$, the plane-wave form of Theorem A3 [ARGUED]. (H2) is the inequality of Section 3.7 (`BETA_FP_U` writes it as $\tilde{h}_{(2)} < L$).
- ($B\theta$): $\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$ exactly (Prop. 3.1). γ_{rec} is bounded in k given Theorem 2.4 (Prop. 3.2). $\mathfrak{d}_F$ is bounded under (DF) $\sup_k |\mathfrak{d}_F(\theta_*; L^k)| < \infty$ [HYPOTHESIS, ARGUED]. `BETA_FP_U` corrects `BETA_FP_PW` §4.3(b): $\mathfrak{d}_F$ is $O(1)$ relative at fixed θ, not $O(\theta^2)$; only boundedness is used.
- Corollary 4.1: given (T1), (H2), (DF), LW (5.7), $|S_k - ck \log L| \leq C_B$, so (5.1) holds with $R = C_B$.
- Theorem 2w by Route B then needs in addition A6 (plane-wave β = Balaban's β up to the expansion point, uniform in k) [ARGUED] and (H0) (Lemma D$''$ (i)).

Route B hypothesis set: (T1), (H2), (DF), LW (5.7), A6, (H0) (plus L5, (G2), odd L, as for Route K). The fixed-level inputs (PI), (M_j), V1 of Route K do not appear; (DF) and A6 replace them.

3.6.4 Comparison

	Route K	Route B
uniform-in-level input	(B2): $\sup_j \sup_K \lvert\sigma_j^{(2)}\rvert < \infty$, $\Leftarrow$ (H2) [PROVED+VERIFIED]	(U), $(B\theta)$, and the β identification and $\mathfrak{d}_{gf}$ bound, both with k-uniform expansion-point bounds; (U) $\Leftarrow$ (T1) + (H2), $(B\theta) \Leftarrow$ Theorem 2.4 + (DF) [PROVED, UNVERIFIED]
single uniform inequality	(H2)	(H2)
fixed-level inputs	(PI), (F), (M_j), V1, (G2)	LW (5.7), $\mathfrak{d}_Y$, $\theta_* < 1$
uses (W), Theorem A3	yes	no; `BETA_FP_U` uses (T1), the plane-wave analogue of A3 [ARGUED]
verification	verified (conditional)	logic of Theorem B_{PW} verified (conditional): `BETA_FP_PW_VERIFY` §6 items 1–4 CONFIRMED [PROVED+VERIFIED, CONDITIONAL]; inputs from `BETA_FP_U` [PROVED, UNVERIFIED], no verifier file
common	(H0) at odd $L > 11$, L5, (G2), V1-type identification, Balaban's proofs, SU(2)	

Shared inputs: (H0), L5, (G2), odd L (> 11 for (H0)), and the identification of the computed $\beta_j(0)$ with Balaban's $\beta_j(0)$ (V1; in Route B also the expansion point). The uniform-in-level hypotheses ((B2) vs (U) + $(B\theta)$) are formally distinct as named hypotheses. Their content overlaps: both rest on (H2) (Section 3.7). In Route K, (H2) gives $(C1\sigma)$ (Theorem C1) and then (B2) (Theorem B2). In Route B, (H2) gives (U) with (T1), and the vertex bound behind $(B\theta)$. So a proof of (H2) serves both routes. One closed route suffices for Theorem 2w.

3.7 The hypothesis (B2) of Route K, and the single uniform inequality (H2)

Both routes need a bound on the coarse symbols that is uniform in the level. This section states that bound for Route K, records what is proved and what is refuted, and reduces it to the inequality (H2). The numerical values of (H2) are in Sections 6.4–6.9. At $L = 13$, (H2) is a certified theorem (Theorem 6.1, Section 6.7).

(B2): $\sup_{j\geq 0} \sup_{K\in \mathrm{BZ}} \lvert\sigma_j^{(2)}(K)\rvert < \infty$. Weaker form that suffices, (B2w): $\sup_j \int_{\mathrm{BZ}} \lvert K\rvert^{-2} \lvert\sigma_j^{(2)}(K)\rvert\, d^4K < \infty$ ($\sigma^{(2)}$ enters d_j only undifferentiated; `KAPPA_BOUNDED` Remark 2.3) [PROVED+VERIFIED: NOT GRADED BY NAME; IMPLIED BY `KAPPA_BOUNDED_VERIFY` §9(IV)].

$\sigma^{(2)}$ obeys the one-step recursion (`U3_AW` (1.2)): $\sigma_j^{(2)} = L^{-2}\mathcal{A}_j\sigma_{j-1}^{(2)} + Q_j[\sigma_{j-1}^{(1)}] + g_j^{(2)}$, with $\mathcal{A}_j s = L^{-2} H_j^{\dagger}(\oplus_l s(k_l)) H_j$ and $H_j(K)$ the one-step gauge-fixed minimizer map. Notation: $\rho_1 := \sup_{j,K} \lVert H_j(K)\rVert / L$ (one step); $H_{(2),j}(K) := (\oplus_l H_{j-1}(k_l)) H_j(K)$ (two steps), $\rho_2 := \sup_{j\geq 2} \sup_{K \text{ real}} \lVert H_{(2),j}(K)\rVert / L^2$.

3.7.1 Proved

- $K = 0$ bound [PROVED+VERIFIED]: $\sup_j |\sigma_j^{(2)}(0)| \le C(|e|C_A + e^2)$. Theorem A3 + scaling give $|\partial^2 \sigma_j^{(1)}(0)| \le 2C_A$ (polarization; `KAPPA_BOUNDED_VERIFY` §7), and the slaving identity (W-2) expresses $\sigma_j^{(2)}(0)$ through these second derivatives.

- Each fixed j [PROVED+VERIFIED]: $\sigma_j^{(2)}$ is analytic on a fixed strip (Prop. A2), hence bounded. The content of (B2) is uniformity in j.

- Prop. 4.1 [PROVED+VERIFIED, CONDITIONAL]: if (h) $h := \sup_{j,K} \|H_j(K)\| < L^2$ and $(C1\sigma)$ $\sup_j \sup_K |\partial_K \sigma_j^{(1)}(K)| < \infty$, then (B2) holds. Proof: in the form norm $\pm s \le N(\Delta + 1)$, $N_j \le (h^2/L^4)N_{j-1} + Z_2$, and Z_2 is finite by Theorem A3, $(C1\sigma)$, and fixed f^0 vertices.

- Why the Lemma A4 mechanism alone fails [PROVED+VERIFIED, REASON CORRECTED BY `KAPPA_BOUNDED_VERIFY` §7]: the obstruction is the missing uniform C^1 bound on $\sigma^{(1)}$, which enters twice: in Q_j (term $L^{-3}\partial_K \sigma_{j-1}^{(1)}$) and in the longitudinal part of $\sigma^{(2)}$ given by (W-2). Theorem A3 is C^0 only.

3.7.2 Refuted, and the failed one-step bound

- (HS) $h < L\sqrt{L-1}$ (named (H′) in `U3_AW`) [REFUTED AT $L = 3$]. Measured $\max_K \|H_j(K)\|/L = 1.652, 1.736, 1.757, 1.760$ for $j = 1, 2, 3, \infty$ (`U3_AW_VERIFY` §6); the threshold is $\sqrt{2} = 1.414$. Gauge choice cannot help: H_j is already minimal-norm in its gauge orbit.

- One-step sup-norm C^1 recursion at $L = 3$ [NUMERICAL, REPRODUCED; `B2_TWOSTEP_VERIFY` GRADE 2]. Its factor is $\rho_1^2/L = 0.909, 1.0046, 1.0289, 1.0325$ for $j = 1, 2, 3, \infty$, so it exceeds 1 for $j \ge 2$ ($\rho_1 > \sqrt{3} = 1.7321$). At $K = 0$ the factor is L^{-1}; the bad region is the face $K_3 = K_4 = \pi$, $K_1 = K_2 \approx 1.86$–1.92. This shows that the one-step sup-norm bound fails at $L = 3$. It does not show that a C^1 bound fails; a weighted norm is not excluded.

3.7.3 (H2) and the two-step theorems [PROVED+VERIFIED, CONDITIONAL ON (H2)]

Source: `B2_TWOSTEP.md` §1–§3; `B2_TWOSTEP_VERIFY.md` grades 1, 3, 4 (CONFIRMED).

$$\text{(H2)} \quad \rho_2 = \sup_{j \ge 2} \sup_{K \in \mathrm{BZ\ real}} \|(\oplus_l H_{j-1}(k_l))H_j(K)\|/L^2 < L.$$

(H2) is a plain operator-norm inequality on real K. It uses no strip, no Δ-relative weight, and only $n = 2$.

Theorem 3.10 (Theorem C1 of `B2_TWOSTEP`). *(H2) ⇒ $(C1\sigma)$, with contraction factor $q_1 = \rho_2^2/L^2$ on the derivative.*

Theorem 3.11 (Theorem B2 of `B2_TWOSTEP`). *(H2) ⇒ (B2), hence (B2w), in plain sup norm, with factor $q_2 = \rho_2^2/L^4$.*

Here $(C1\sigma)$ is $\sup_j \sup_K |\partial_K \sigma_j^{(1)}(K)| < \infty$ (Section 3.7.1). The verified points are these.

- Two-step map. $\mathcal{A}_j \mathcal{A}_{j-1} s = L^{-4} H_{(2)}^{\dagger}(\oplus s) H_{(2)}$, with $dk_{l,l'}/dK = L^{-2}$ [CONFIRMED, grade 1].

- Theorem 3.10. The chain-rule gain is exactly L^{-2} per two steps on real K. So the threshold is $\rho_2 < L$, not $\sqrt{L^2-1}$: no strip-Cauchy loss enters. The large constants ($b(L)$ of O6, η_1, η_2, G_1) enter only the source term, not the contraction factor [CONFIRMED, grade 3].

- Theorem 3.11. This contraction needs only $\rho_2 < L^2$, which (H2) implies [CONFIRMED, grade 3]. In one step it is Prop. 4.1's (h).

- (H2) vs (H″). (H2) is the plain operator norm of the same matrix as `U3_AW` (H″) at $n = 2$. It is weaker: threshold L (3 at $L = 3$) against $\sqrt{L^2-1}$ (2.828), real K only, $n = 2$ only [CONFIRMED, grade 4]. (H″) ⇒ (B2) stays [ARGUED] and is no longer needed.

- Form of Q_j (G3) [CONFIRMED by read-off of `U3_AW` (1.2)/(3.3) and P10]: $\partial\sigma^{(1)}$ enters linearly. `U3_LEMMA51` (3.3) itself was not re-derived from the Moyal rule by either party.

- Not covered: C^2 and C^3 bounds on $\sigma^{(1)}$ under (H2) (`B2_TWOSTEP` §5, "look routine", not written, not graded). They are not needed for (B2).

- Submultiplicativity: $\rho_2 \le \rho_1^2$ always. At $L = 3$ the gain of the two-step map is that this is strict by a factor 1.50 (2.064 vs 3.098) [CONFIRMED].

3.7.4 One-step sufficiency [ARITHMETIC PROVED; INPUT CERTIFIED AT $L = 13$]

$\|H_{(2),j}(K)\| \le \max_l \|H_{j-1}(k_l)\| \cdot \|H_j(K)\| \le L^2\rho_1^2$. So $\rho_1^2 < L$ implies (H2), and then the one-step C^1 recursion of `B2_TWOSTEP` (2.2) contracts with factor ρ_1^2/L. At $L = 3$ the measured ρ_1^2 exceeds L, so the one-step route fails and the two-step map is needed; at $L = 5$ and $L = 7$ the measured ρ_1^2/L is below 1. The values are in Section 6.5. At $L = 13$ the input is certified: $\sup_{j,K} \rho_1^2 \le 10 < 13$ (Theorem 6.1), so (H2) holds there with $\rho_2 \le 10$.

3.7.5 The precise remaining inequality

At a fixed odd $L \ge 3$ the single uniform hypothesis is (H2).

- Route K: (H2) ⇒ ($C1\sigma$) (Theorem C1) ⇒ (B2) (Theorem B2) [PROVED+VERIFIED, CONDITIONAL ON (H2)].

- Route B: (H2) + (T1) ⇒ (U), and (H2) ⇒ the vertex bound for γ_{rec} in ($B\theta$) (`BETA_FP_U` Theorem 2.4, Prop. 3.2) [PROVED, UNVERIFIED].

- Status of (H2): at $L = 13$, inside Bałaban's range $L > 11$, [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]: $\rho_1(j,K)^2 \le 10 < 13$ for all $j \ge 1$ (including ∞) and all K, hence $\rho_2 \le 10 < 13$ (Theorem 6.1; residue in Section 6.7). At $L = 3, 5, 9$: [NUMERICAL] only (at $L = 9$ one point).

- Kept as alternatives: ($C1\sigma$) + (h) ⇒ (B2) [PROVED+VERIFIED]; (H″) ⇒ (B2) [ARGUED]; (B2w) directly. No per-momentum $d = 4$ constant-field code exists, so (B2) itself has not been scanned numerically (`KAPPA_BOUNDED_VERIFY` §8).

3.8 Conclusion of the reduction

Theorem 2w ⇐ (Route K inputs) or (Route B inputs). Precisely:

Route K. (B2) [OPEN; REDUCES TO (H2) (VERIFIED REDUCTION, SECTION 3.7.3); (H2) CERTIFIED AT $L = 13$ (THEOREM 6.1)] + (PI), (F), (M_j), V1, (G2) [FIXED-LEVEL] + (H0) [BALABAN,

ODD $L > 11$; ITS USE IS A VERIFIED REDUCTION, SECTION 5.9] + L5 [OPEN] ⇒ Theorem K ($R = \kappa_*$) ⇒ (5.1) ⇒ Lemma D″ ⇒ Theorem 2w. Status of the implication: [PROVED+VERIFIED, CONDITIONAL].

Route B. (U) + ($B\theta$) with $\theta_* < 1$ + LW (5.7) + bounded $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ + identification of β (uniform in k) + (G2) + (H0) [ODD $L > 11$] + L5 ⇒ Theorem $\mathrm{B_{PW}}$ ($R = C_B$) ⇒ (5.1) ⇒ Lemma D″ ⇒ Theorem 2w. Status: Theorem $\mathrm{B_{PW}}$ logic [PROVED+VERIFIED, CONDITIONAL]; inputs [PROVED, UNVERIFIED]. With `BETA_FP_U` (Section 3.6.3): (T1) + (H2) + (DF) + LW (5.7) ⇒ Corollary 4.1 ($R = C_B$); with A6 + (H0) + L5 + (G2) ⇒ Theorem 2w. Status: [PROVED, UNVERIFIED, CONDITIONAL] (the inputs (T1), (DF), A6 and the `BETA_FP_U` reductions are unverified).

Single uniform hypothesis. Both routes need (H2). Theorem 6.1 certifies (H2) at $L = 13$, an odd $L > 11$. This discharges (B2), the only uniform-in-level hypothesis of Route K (Theorems C1, B2). In Route B it reduces (U) to (T1); (DF) and A6 remain.

Then Balaban's Theorem 1. Theorem 2w gives (b) and (c′). With γ shrunk so that (S) holds, every one of the 19 cross-scale uses in CMP 119 and CMP 122 I–II is covered (Section 3.4.3), and the single-scale uses need only $g_k \le \gamma$. So Theorem 1 holds for the bare coupling $g_0(K, g)$, for every K and every $g \in]0, g_1]$, without Balaban's coupling assumption. This needs L5 in the sharpened form of Section 3.4.5: β must exist on every history the shooting visits. Status: [PROVED, UNVERIFIED (SWEEP), CONDITIONAL ON ROUTE K OR ROUTE B INPUTS, ON L5, AND ON BALABAN'S PUBLISHED PROOFS]. The sweep uses Bałaban's sentence that similar inequalities hold for the later constants (ledger A11); round 5 closes it as a verified list (Proposition 5.7), with the order entries C28–C30. The step to Theorem 1 also needs the finite-volume identification B8, which rests on A15; Proposition 5.8 gives the torus coefficient within $C_j N_j^6 e^{-m_j N_j/2}$ of the whole-lattice one, given (Loc), and exact equality cannot be derived. So Theorem 2w ⇒ Theorem 1 rests on 4 Group A items (A5, A11, A13, A15), 9 Group B items (B1–B8, B10), and X5.

Order of constants. Fix $\beta_0, p_0, p_1, r, \nu, \kappa_0$ (Balaban). Fix L (odd, > 11), then R from Route K or B, then $b = c \log L$, then $\gamma \le \min(\gamma_0, \gamma_{Bal}, b/(4C_1), \gamma_*)$ with $\gamma^{-2} \ge c_{sw} \cdot 2R$. Then $g_1 = (\gamma^{-2} + 2R)^{-1/2}$. Balaban's phrase "for a sufficiently small positive g" fits $g \le g_1$.

What this does not give.

- Literal (0.31): only for windows of length $\ge 8R/b$. Literal (0.31) at every K needs (L1) or (L2) of Section 3.3, which need individual increments, not window sums. Not needed for Theorem 1.
- The value $\lim \beta_j(0)$ and any rate: Section 3.9.
- Even L, or odd $L \le 11$ without (H0) as an extra hypothesis (Balaban assumes L odd > 11 throughout CMP 109); SU(N) with $N > 2$.
- Monotone couplings: g_m can exceed g_n for $m < n$ by the bounded amount $u_m \ge u_n - D'$. Section 3.4.3 shows Balaban's proof tolerates this under (S); a line-by-line read was not done.
- A continuum limit (`ASSEMBLY.md` treats the assembly separately; not claimed here).

3.9 The convergence route: what it adds, and its status

Not needed for Theorem 1. It adds: (i) existence of $\beta_\infty = \lim \beta_j(0)$; (ii) its value $c \log L$; (iii) a rate; (iv) eventual positivity, hence literal (0.31) for $K \ge K_0$ (Lemma D′ (d)); (v) with certified early steps, literal (0.31) at every K (Section 3.3 (L2)). It also implies (B2), since convergence of $\sigma_j^{(2)}$ implies boundedness. So it needs strictly more than Route K.

3.9.1 Components and status

- U3 Theorem 4.1 with O5: $|\partial_K^\alpha(\Delta_M - \Delta_\infty)| \le CM^{-2}$, $|\alpha| \le 2$, uniform to $K = 0$, odd L [PROVED+VERIFIED] (`U3_VERIFY`, `U3_REPAIRS_VERIFY`). Numerically $|\Delta_M - \Delta_\infty| =$ 1.007, 0.129, 0.0147 at $M = 3, 9, 27$.

- O6: $\|\mathfrak{B}_j(K)^{-1}\| \le b(L)$ for all j, K, one-step constraint, trivial background [PROVED+VERIFIED]. Crude constant ($b(3) \approx 7.5 \cdot 10^{33}$); measured corner values 103.5, 176.7, 193.0, 195.3 ($j = 0, 1, 2, \infty$, $L = 3$) use a surrogate Λ (`U3_REPAIRS_VERIFY` §5).

- Theorem 6.1: $\beta_{j+1}(0) = \Phi(\mathcal{D}_j)$ is Lipschitz in the level data [PROVED+VERIFIED, CONDITIONAL ON V1]. The norm is the Taylor-data norm (`U3_VERIFY` P8 correction).

- Lemma 5.1 (convergence of $\sigma_j^{(1)}$, $\sigma_j^{(2)}$ in C^2, $|\alpha| \le 2$):
  - neutral sector, rate L^{-4j} [PROVED+VERIFIED FOR THE VALUE BOUND; THE K-DERIVATIVE PARAGRAPH IS PROVED, UNVERIFIED];
  - (W) [PROVED+VERIFIED] (`U3_AW` Theorem W); it removes the marginal direction of the f^1 recursion;
  - (A) qualitative and Theorem A3 [PROVED+VERIFIED]; (A) in full (strip/C^3 for $m = 1$, all of $m = 2$) $\Leftarrow$ (H″) [OPEN]; the route via (HS) is [REFUTED] at $L = 3$;
  - $\sigma^{(1)}$ convergence at rate jL^{-2j} [SKETCH IN `U3_AW` §4.7, UNVERIFIED]; $\sigma^{(2)}$ convergence [OPEN, NEEDS (A) IN FULL];
  - the single-block-constraint version of Lemma 5.1 is false for $m = 2$ in $d = 4$ [PROVED+VERIFIED]; Balaban's composed constraint is not affected.

- Prop. 3.6: $|\beta_{j+1}(0) - \beta_\infty| \le C(1 + j)^2 L^{-2j}$ [PROVED, UNVERIFIED AS AN ASSEMBLY; CONDITIONAL ON V1, (A) IN FULL]. (The form Cj^2L^{-2j} of the first version of these notes is 0 at $j = 0$ and is corrected here.) The identification of σ-Taylor data with the data of (6.1) is cited from `U3_VERIFY` §7, not re-shown. (N) (clean rate L^{-2j}) is not needed and is judged plausibly false.

- Value of the limit. Given convergence, Theorem K or Theorem $\mathrm{B_{PW}}$ gives $\beta_\infty = c \log L$ at once: $S_k/k \to \beta_\infty$ by Cesàro and $S_k/k = c \log L + O(1/k)$ [PROVED+VERIFIED AS AN IMPLICATION: `THEOREM2_PROOF_VERIFY` POINT 4; `BETA_FP_PW` THEOREM 3.2(III)]. Older routes to the value: `BETA_FP_H` Theorem 3.1′ with (H) restated at the constant field, which given (W) reduces to Lemma 5.1 (1.2) alone (`BETA_FP_H_VERIFY` items (1)–(4)), plus (PI), (F), (M) [CONDITIONAL; (M) ARGUED]; and `BETA_FP_PW` Theorem 3.1 (limit form) given (U), ($C\theta$), LW (5.7) [PROVED, UNVERIFIED]. The original (H) with IR weights (`BETA_FP_LEMMA43` (4.2)) was graded UNSUPPORTED and is not used.

- Rate of $\mathcal{C}(\Lambda)$: $O(\Lambda^{-2} \log \Lambda)$ given Lemma 5.1 with rates, via `U3_AW` §3.4(b) slaving [ARGUED, `BETA_FP_H_VERIFY` §5].

3.9.2 Early steps and j_*

Under Prop. 3.6 and $\beta_\infty = c \log L$, $\beta_{j+1}(0) \ge (c \log L)/2$ for $j \ge j_* := \min\{J : C(1 + j)^2 L^{-2j} \le (c \log L)/2 \text{ for } j \ge J\}$. Heuristic size [ARGUED]: $C \sim b^3$ gives $j_* \approx 110$ ($L = 3$), ≈ 70 ($L = 13$); $C \sim b^6$ (one-step inverses inside C_{data}) gives ≈ 218, ≈ 138 (`THEOREM2_PROOF_VERIFY` Point 5(e)).

With the measured $\|\mathfrak{B}^{-1}\| \approx 195$: $j_* \approx 10$ ($L = 3$), ≈ 4 ($L = 13$). Certifying $\beta_1(0), \dots, \beta_{j_*}(0) > 0$ needs interval arithmetic at odd $L \geq 13$ (blocks $\approx 3.4 \cdot 10^5$ rows per momentum by dense methods; the umklapp code reduces this to 12×12 blocks at L^4 momenta). It is needed only for (L2), not for Theorem 1.

4 The finite-torus existence theorem

This section proves Theorem B. It takes the trajectory of Section 3 as the input block (B-T2). Section 4.1 fixes what "exists" means here. Section 4.2 states the uniform bound (H-U) that the blocks must deliver, and Sections 4.3–4.4 state the theorems. Section 4.5 proves them modulo the named blocks, and Section 4.6 gives the dependency table, which is the single status table of the paper. Section 6 then collects the numerical support for the hypotheses.

Source: `ASSEMBLY.md` (2026-10-27), verified in `ASSEMBLY_VERIFY.md`. The statements carry the verifier's corrections (Section 4.8). The theorem is a reduction to named blocks. It is not unconditional. The notation is that of Sections 2.1 and 2.5.

4.1 What "exists" means here

Jaffe–Witten footnote 2 [15] [VERIFIED QUOTE] (`reports/clay_ym.txt` line 619; anchor: the sentence "Such a result presumably would enable the study of the limit as $T^4 \to \mathbb{R}^4$", line 582): "We specifically exclude weak-existence (compactness) as the solution to the existence part of the Millennium problem, unless one also uses other techniques to establish properties of the limit (such as the existence of a mass gap and the axioms)."

Reading used here. "Exists" means: (i) the family $\{S_{n,\varepsilon}\}$ is relatively compact (uniform bounds); (ii) at least one limit point exists along a sequence $\varepsilon_k \to 0$; (iii) every limit point has the properties (OS0)–(OS3) below, proved on the limit, not assumed. No uniqueness is claimed. Full-sequence convergence needs E13 (comparison of two trajectories; `SOURCES.md` (E14-d), D6), which is not in this paper.

This reading is weaker than the footnote's parenthesis. The theorem below gives no gap and only the torus versions of four axioms. It is NOT an admissible Clay solution. It is the finite-torus step of Jaffe–Witten §6.5 [VERIFIED QUOTE]: "One must then verify the existence of limits of appropriate expectations of gauge-invariant observables as the lattice spacing tends to zero and as the volume tends to infinity" (only the first limit is done here).

The OS axioms are not stated in `reports/clay_ym.txt`. It cites them as OS (CMP 31 (1973), CMP 42 (1975)) [19, 20] and requires "axiomatic properties at least as strong as those cited" (lines 279–280) [VERIFIED QUOTE]. The forms used follow the template's application of OS II (`reports/o3_continuum.txt` lines 4987–5003) [VERIFIED QUOTE]: "The linear growth condition requires a bound $|S_n(F)| \leq \sigma_n |F|_{s_n}$ with fixed s and $\sigma_n \leq A(n!)^D$ for fixed A, D", and restriction to Schwartz tests flat on the coincidence diagonals gives the test class of that reference. OS II itself was not re-read [CITED].

4.2 Hypothesis delivered by the blocks

Definition 4.1 ((H-U) uniform bound)**.** There are $s \in \mathbb{N}$ and $A, D < \infty$ such that for all $K \geq K_0$, all $n \geq 1$, and all $F \in \mathcal{D}_n$:

$$|S_{n,\varepsilon}(F)| \leq \sigma_n \|F\|_{sn}, \qquad \sigma_n := A(n!)^D, \quad \varepsilon = L^{-K}.$$

The derivative order sn grows linearly in n. The OS linear growth condition admits this. (H-U) is the output of blocks (B-T2) + (B-S) + the bridge (B-N1) of Section 4.5.

4.3 Theorem (finite-torus existence, compactness form)

Theorem 4.2 (finite-torus existence, compactness form [PROVED MODULO NAMED BLOCKS]). *Let $G = SU(2)$, $d = 4$, L odd and large, $\ell \in 2\mathbb{N}$ fixed. Let $\gamma > 0$ be Balaban's small constant and $g \in]0, \gamma]$ small. For $K \geq K_0$ let $\varepsilon = L^{-K}$ and $g_0 = g_0(\varepsilon, g)$ be a shooting solution (Lemma 3.4) whose effective couplings satisfy $g_k \in]0, \gamma]$ for $0 \leq k \leq K$ and $g_K = g$. Assume (H-U). When (H-U) is taken from the blocks (Section 4.2) rather than assumed, the chain also assumes the $\kappa \to \kappa/2$ decay recovery of D2 [R, B22] (not stated, `SOURCES_D2_VERIFY.md` (5)), which does not follow from the other hypotheses. Then:*

(a) Compactness. Every sequence $K_m \to \infty$ has a subsequence K_{m_i} such that, with $\varepsilon_i = L^{-K_{m_i}}$, the limit $S_n(F) := \lim_i S_{n,\varepsilon_i}(F)$ exists for every $n \geq 1$ and every $F \in \mathcal{D}_n$.

(b) For every limit family $(S_n)_{n \geq 1}$ obtained in (a):

- *(OS0) Regularity. S_n is linear on $\mathcal{D}_n$, and $|S_n(F)| \leq A(n!)^D \|F\|_{sn}$. In particular S_n restricted to $C_c^\infty(T^{4n} \setminus \bigcup_{i<j}\{x_i = x_j\})$ is a distribution of order $\leq sn$.*
- *(OS1) Invariance. $S_n(F \circ \tau_a^{\times n}) = S_n(F)$ for every $a \in T_\ell^4$ (all continuous translations), and $S_n(F \circ R^{\times n}) = S_n(F)$ for every R in the hypercubic group B_4 (signed permutations of the axes, acting about a lattice site).*
- *(OS2) Reflection positivity. Let $\theta(x_0, \vec{x}) = (-x_0, \vec{x})$ on T_ℓ^4 and $T_+ := \{x : x_0 \in]0, \ell/2[\}$. For every finite family $F_0 \in \mathbb{C}$, $F_n \in \mathcal{D}_n$ with $\operatorname{supp} F_n \subset T_+^n$ $(1 \leq n \leq N)$:*

$$\sum_{n,m=0}^{N} S_{n+m}(\Theta F_n \otimes F_m) \geq 0, \quad (\Theta F_n)(x_1, \ldots, x_n) := \overline{F_n(\theta x_n, \ldots, \theta x_1)}, \quad S_0 := 1.$$

- *(OS3) Symmetry and reality. $S_n(F \circ \pi) = S_n(F)$ for every permutation π of the n arguments, and $S_n(F) \in \mathbb{R}$ for real F.*

(c) Nontriviality is a separate statement, Theorem 4.3.

The sourced blocks use the unit-scale condition (U), $\rho_1 \leq c_* r_0 / M^a$ at $\lambda_K = 1$ (r_0 the Cauchy margin of the constant-source family, `SOURCES_D2C.md` (3.5)), as a constant choice (C23), made once after r_0 and before K; B-N1 meets it by the rescaling $\lambda_i = 2N_U(f_i)$ at a C^n cost, s and D unchanged (`SOURCES_UN1.md`, `SOURCES_UN1_VERIFY.md`). This (U) is not the Route B hypothesis (U). *What is claimed about measures.* No probability measure is claimed. The limit is a family of Schwinger functions on $\mathcal{D}_n$. A measure on distributions would need finite moments of smeared fields, and $\operatorname{Var} \sum_p f(x_p)\varphi_p \sim \varepsilon^{-4}$ (Section 2.5). (OS0)–(OS3) on $\mathcal{D}_n$ are the inputs of OS reconstruction.

4.4 Theorem (nontriviality), conditional on (B-NT)

Theorem 4.3 (nontriviality [CONDITIONAL ON (NT) OF SECTION 4.5.9]). *Under the hypotheses of Theorem 4.2 and (NT), every limit family satisfies: (i) $S_2 \neq 0$: there are $f_1, f_2 \in C^\infty(T^4)$ with disjoint supports and $S_2(f_1 \otimes f_2) \neq 0$. (ii) S_2 is not the two-point function of a scale-invariant (in particular free Maxwell) composite: $d^8 S_2(x, y)$, $|x - y| = d$, is not asymptotically constant as $d \to 0$.*

Part (ii) is stated here in the form the proof supports (two-sided envelope only); see Section 4.8, item 3. The source stated (ii) as "behaves like $c\, g(d)^4$ with $g(d)^2 \sim (\beta \log(1/d))^{-1}$", which needs $\beta_j \to \beta_\infty$ and is not proved.

4.5 Proof modulo blocks

Structure. (B-T2) gives the trajectory. (B-S) gives sourced bounds uniform in ε. (B-N1) turns them into (H-U). (B-C) gives (a). (B-REG), (B-TI), (B-RP), and (OS3) give (b). (B-NT) gives Theorem 4.3. Only (B-C), (B-REG), (B-TI), (B-RP), and (OS3) are proved in this section; they use nothing beyond (H-U) and exact lattice symmetries.

4.5.1 (B-T2) Theorem 2: the running coupling

Needed. The hypothesis of Balaban's Theorem 1 (CMP 122 II p. 355), $g_k \in]0, \gamma]$ for all k, quoted in Section 1.2 [VERIFIED QUOTE]. It was to be discharged by Theorem 2 of CMP 109 p. 259 [8]: for small g there is $g_0(\varepsilon, g)$ with all $g_k \in]0, \gamma]$, $g_K = g$, and (0.31): $1/g^2 + \beta \log(L^k \varepsilon)^{-1} \le 1/g_k^2 \le 1/g^2 + \beta' \log(L^k \varepsilon)^{-1}$. Balaban: "has not been published yet" [VERIFIED QUOTE].

Status from Section 3 (and §10 of `THEOREM2_PROOF_v4.md`). The weak form, Theorem 2w, is [PROVED+VERIFIED, CONDITIONAL] by Route K (Theorem K + Lemma D″) on (H2) and the fixed-level inputs (PI), (F), (M_j), V1, (G2), plus L5 and item 7′ ((H0) at odd $L > 11$, whose use is a verified reduction with residue N1–N9 (Section 5.9), or (G-gain) with the running-maximum lift, verified (`SOURCES_GGAIN_NONMONO_VERIFY`)); item 9 (K-independence) is closed. It suffices for Theorem 1 by the cross-scale sweep [PROVED; VERIFIED IN ROUND 7 (LEDGER B10)]. Round 7 proves (M_j) and L5 relative to listed inputs (Theorems 5.14 and 5.15); (F) is reduced but unverified. Literal (0.31) is not obtained (it needs eventual positivity of $\beta_j(0)$). Route B (Theorem B of `BETA_FP_PW`) is a formally distinct second route; its logic is [PROVED+VERIFIED, CONDITIONAL], its inputs (U), (DF), (T1) are [PROVED, UNVERIFIED] and rest on (H2). The reduction pieces are Lemma D′, Lemma D″, and Lemmas D, E (the one-argument forms in Lean).

Conclusion. At $L = 13$, (H2) is certified (Theorem 6.1); (B-T2) remains conditional on the fixed-level inputs, the inputs of L5, and item 7′; see the dependency table (Section 4.6).

4.5.2 (B-S) Sourced bounds E14-a..g

Needed. A version of Balaban's Theorem 1 with the (C1) source, i.e. with the local bare coupling $w_p = 1/g_0^2 - f(x_p)$, f real, $|\partial^m f| \le r$ for $m \le 5$ (class F_r), uniformly in ε. Output form (`SOURCES.md` (N1)): $\log Z_\varepsilon[f] - W_\varepsilon[f]$ is C^∞ in source amplitudes z with $|\partial_z^\alpha (\log Z_\varepsilon[f] - W_\varepsilon[f])| \le C^{|\alpha|} \alpha!\, r^{-|\alpha|}$ for $f = \sum_i z_i f_i$ with disjoint supports. $W_\varepsilon[f]$ is a local field-independent contact functional (allowed to diverge; it contributes zero to every mixed derivative of order ≥ 2 for disjoint supports, `SOURCES_VERIFY` item 5(a)).

Estimates needed (`SOURCES.md` §2.5): (E14-a) sourced small-field step; (E14-b) sourced large-field step and R operation; (E14-c) derivative bounds in f. Not needed: (E14-d) fixed-torus convergence (= E13); (E14-g) tensor sources. (E14-e) RP and symmetry transfer is done in this section (Sections 4.5.6, 4.5.7). (E14-f) nontriviality at a fixed layer is needed only for Theorem 4.3. The current status of each sub-block is in the dependency table (Section 4.6); (B-S) remains a [HYPOTHESIS].

4.5.3 (B-N1) Bridge: source-form bounds ⇒ (H-U) [OPEN]

`SOURCES_VERIFY` item 5(b) [VERIFIED QUOTE]: the source form "does not give $|S_n(F)| \le C^n n! \|F\|_{s,n}$ for F flat on diagonals with a fixed s … Mark N1 for general F as a gap." This paper does not close the gap. It gives a route [DERIVED SKETCH], which shows that the gap is removed by the OS index growth sn.

1. Moments from cumulants. $\langle \prod_i \phi(f_i) \rangle = \sum_{\text{partitions}} \prod_{\text{blocks}} \kappa(\cdot)$. Singleton cumulants vanish (φ is centred). For disjoint supports the source form bounds every cumulant by $C^{|B|}|B|!\, r^{-|B|}$, so the moment by $C^n n!\, r^{-n}$.

2. Scale-δ tests. A bump of side $\delta \le 1$ and amplitude a has $|\partial^5 f| \sim a\delta^{-5}$; it lies in F_r if $a \le cr\delta^5$. So n disjoint unit-amplitude scale-δ bumps have $|\text{moment}| \le C^n n!\, \delta^{-5n}$. This is lossy but polynomial in δ^{-1}.

3. Whitney decomposition. Split T^{4n} by a smooth dyadic partition in $\rho(x) := \min_{i<j} |x_i - x_j| \in [\delta, 2\delta]$, $\delta = 2^{-m} \ge \varepsilon$. On each piece cover by cubes of side $\delta/100$; cube tuples are pairwise disjoint. Expand $F\cdot$(cutoffs) in a Fourier series on each cube tuple; the coefficients decay faster than any power; each term is a product of disjointly supported scale-δ bumps. Flatness gives $|\partial^\beta F(x)| \le C_k \|F\|_{k+|\beta|} \rho(x)^k / k!$. The count is $\le C^n \delta^{-4n}$ cube tuples. Derivatives of the cutoffs cost at most $(Cn^2/\delta)^M$; with $M = O(n)$ this is $\le C^n (n!)^c \delta^{-M}$.

4. Bound per dyadic scale: $\le C^n (n!)^{D'} \delta^{k-9n-M} \|F\|_{k+M}$. Choose $k = sn - M$ with s large enough ($s \ge 20$ suffices on this count) so the δ-exponent is $\ge n$; the sum over dyadic $\delta \in [\varepsilon, 1]$ converges uniformly in ε.

5. Sub-lattice region $\rho(x) \lesssim \varepsilon$: $\le C^n \varepsilon^{-4n}$ plaquette tuples, each moment $\le 2^n$, and $|F| \le \|F\|_k \varepsilon^k$; this is $\le C^n \|F\|_{sn}$ for $k = sn \ge 4n$.

Result (if the steps hold): (H-U) with some fixed s and D. To check: uniformity of the source bound in the position of the bumps and in ε; the C^∞ real-source form of (E14-c) with n-th derivatives for all n; the cutoff-derivative count in step 3. The verifier lists further gaps: complex Fourier modes against real sources (a factor 2^n), the extra derivatives of Fourier summation (absorbed in s), and the smooth min-distance partition. Later work (`SOURCES_D4D5`, graded in `SOURCES_D4D5_VERIFY`) confirms "($\mathrm{S_{src}}$) $\Rightarrow$ (H-U) with $s = 6$, $D > 1$" given (FX) (see Section 4.6). `ASSEMBLY_N1.md` closes the three gaps given (S) (confirmed in `SOURCES_HHIST_N1_VERIFY.md`) and, since 2026-10-28, uses the verified form: C^1 norm $\max(\|f\|_\infty/r, \|\nabla f\|_\infty/\rho_1)$, order $(n!)^{1+\eta}$, $s = 6$, $D > 1$ ($D > 2$ under the fallback order $2 + \eta$); the ∂^5, δ^{-5n}, and $s \ge 20$ counts in the sketch above are superseded.

4.5.4 (B-C) Compactness [PROVED HERE, GIVEN (H-U)]

Proposition 4.4. *Under (H-U), every sequence $\varepsilon_m \to 0$ has a subsequence along which $S_{n,\varepsilon}(F)$ converges for every n and every $F \in \mathcal{D}_n$. The limit S_n is linear and satisfies $|S_n(F)| \le \sigma_n \|F\|_{sn}$.*

Proof. 1. Separability. $C^{sn}(T^{4n})$ with $\|\cdot\|_{sn}$ is separable (trigonometric polynomials with rational coefficients are dense; Fejér means converge in C^k). A subset of a separable metric space is separable. So $\mathcal{D}_n$ has a countable subset Q_n dense for $\|\cdot\|_{sn}$. 2. Diagonal extraction. $Q := \bigcup_n Q_n \times \{n\}$ is countable; enumerate it as $(F^{(1)}, n_1), (F^{(2)}, n_2), \ldots$. Each sequence $S_{n_t,\varepsilon_m}(F^{(t)})$ is bounded by $\sigma_{n_t} \|F^{(t)}\|_{sn_t}$ (H-U), so it has a convergent subsequence. Extract successively and take the diagonal subsequence ε_i. 3. Extension. Let $F \in \mathcal{D}_n$, $\eta > 0$, and $G \in Q_n$ with $\|F - G\|_{sn} \le \eta$. By (H-U) and linearity, $|S_{n,\varepsilon_i}(F) - S_{n,\varepsilon_j}(F)| \le 2\sigma_n \eta + |S_{n,\varepsilon_i}(G) - S_{n,\varepsilon_j}(G)|$. The last term $\to 0$. So the sequence is Cauchy; call the limit $S_n(F)$. 4. Linearity passes to pointwise limits, and $|S_n(F)| = \lim |S_{n,\varepsilon_i}(F)| \le \sigma_n \|F\|_{sn}$. □

Remarks. (i) The same subsequence serves all n; this matters for (OS2). (ii) Nothing here needs E13. (iii) The bounded normalizations B_K and c_ε can be included in the extraction.

4.5.5 (B-REG) Regularity (OS0) [PROVED HERE, GIVEN (H-U)]

By step 4 of the proof of Proposition 4.4 the limit satisfies $|S_n(F)| \le \sigma_n \|F\|_{sn}$, $\sigma_n = A(n!)^D$. On the compact torus the Schwartz seminorm is replaced by $\|F\|_p$ (the weights control decay at infinity on $\mathbb{R}^{dn}$; the torus has none). This is the quoted OS II form. Distribution property: for $F \in C_c^\infty$ of the off-diagonal open set, $F \in \mathcal{D}_n$ and the bound is continuity in the C^{sn} topology, so S_n is a distribution of order $\le sn$ there.

Caveat. OS II is formulated on $\mathbb{R}^{dn}$. On T^{4n} the axioms are used in their torus form (periodic test functions). This is a change of setting, not a consequence of OS II.

4.5.6 (B-TI) Translation and hypercubic invariance (OS1) [PROVED HERE, GIVEN (H-U)]

Lattice, exact. Put the origin at a lattice site.

- Translations $a \in \varepsilon\mathbb{Z}^4$. The map $U \mapsto U^a$, $U^a(b) := U(b-a)$, permutes the links, preserves the product Haar measure and the Wilson action, and $P_p(U^a) = P_{p-a}(U)$. So $\langle \prod_i \varphi_{p_i+a} \rangle_\varepsilon = \langle \prod_i \varphi_{p_i} \rangle_\varepsilon$ and $S_{n,\varepsilon}(F \circ \tau_a^{\times n}) = S_{n,\varepsilon}(F)$. The coupling $g_0(\varepsilon)$ is one number, so nothing in μ_ε depends on position.

- Hypercubic $R \in B_4$ (signed permutation matrices, $|B_4| = 384$). R maps sites to sites, links to links (possibly reversed), plaquettes to plaquettes, and $x_{Rp} = Rx_p$. Define $(RU)(b) := U(R^{-1}b)$ with $U(-b) = U(b)^{-1}$. Haar measure is invariant under $U \mapsto U^{-1}$ and $\operatorname{Re}\operatorname{tr} W^{-1} = \operatorname{Re}\operatorname{tr} W$ for $W \in SU(2)$, so A and the measure are invariant and $P_{Rp}(RU) = P_p(U)$. Hence $S_{n,\varepsilon}(F \circ R^{\times n}) = S_{n,\varepsilon}(F)$.

- These identities hold at every ε. $F \circ \tau_a^{\times n}$ and $F \circ R^{\times n}$ are again in $\mathcal{D}_n$. They pass to the limit along the subsequence of Proposition 4.4: $S_n(F \circ R^{\times n}) = S_n(F)$, and $S_n(F \circ \tau_a^{\times n}) = S_n(F)$ for $a \in \varepsilon_i\mathbb{Z}^4$ for all large i.

Continuous translations. Let $a \in T^4_\ell$. Pick $a_i \in \varepsilon_i\mathbb{Z}^4$ with $|a - a_i| \le \varepsilon_i$ (componentwise rounding gives $\le \varepsilon_i\sqrt{4}/2 = \varepsilon_i$). Write $F_a := F \circ \tau_a^{\times n}$. The mean value theorem along the diagonal direction gives $\|F_a - F_{a_i}\|_{sn} \le 4n|a - a_i|\,\|F\|_{sn+1}$. Then

$$|S_n(F_a) - S_n(F)| \le |S_n(F_a) - S_{n,\varepsilon_i}(F_a)| + |S_{n,\varepsilon_i}(F_a) - S_{n,\varepsilon_i}(F_{a_i})| + |S_{n,\varepsilon_i}(F_{a_i}) - S_{n,\varepsilon_i}(F)| + |S_{n,\varepsilon_i}(F) - S_n(F)|.$$

The first and last terms $\to 0$ by Proposition 4.4. The third is 0 by lattice invariance. The second is $\le \sigma_n \cdot 4n\,\varepsilon_i \|F\|_{sn+1}$ by (H-U) and linearity. Let $i \to \infty$. F is smooth, so $\|F\|_{sn+1} < \infty$. Rotations outside B_4 are not available (Section 4.7).

4.5.7 (B-RP) Reflection positivity (OS2)

Lattice theorem [CITED]: Osterwalder–Seiler [21], Ann. Phys. 110 (1978) 440–471. The paper is not in the project folder; no verbatim quote. Content as usually stated: for the Wilson action with compact gauge group and any $\beta \ge 0$, and θ the reflection in a lattice hyperplane that bisects the links orthogonal to it, $\langle \Theta(\Phi)\Phi \rangle \ge 0$ for every function Φ of the link variables in the positive half-lattice, $(\Theta\Phi)(U) := \overline{\Phi(\theta U)}$. Jaffe–Witten [15] line 546 [VERIFIED QUOTE]: "Reflection positivity holds for the Wilson approximation [36], a major advantage". The template uses the same input (`o3_continuum.txt` line 4902) [VERIFIED QUOTE]: "At every cutoff the nearest-neighbour measure is reflection positive across microscopic site and link seams." Proof idea (standard): fix the crossing links to 1 by a gauge transformation; the action splits as $A_+ + \theta A_+ + \sum_{\text{crossing}} \operatorname{Re}\operatorname{tr}(X_p \theta(X_p)^*)$, and $\exp(\beta \operatorname{Re}\operatorname{tr}(XY^*))$ is a positive-type kernel (positive combination of characters), which gives a sum

of squares. On the periodic lattice there are two fixed planes, $x_0 = \varepsilon/2$ and $x_0 = \varepsilon/2 + \ell/2$; both are link planes because N_ε is even (ℓ even, Section 2.5). The periodic two-seam version is standard [CITED]; its text was not checked. For site-plane reflections see Menotti–Pelissetto [17] (Section 4.8, item 1).

Passage to the limit [PROVED HERE, GIVEN (H-U) AND THE LATTICE THEOREM]. Setup. $\theta_\varepsilon(x) := \theta(x) + \varepsilon e_0 = \theta(x - \varepsilon e_0)$, fixed planes $x_0 = \varepsilon/2,\ \varepsilon/2 + \ell/2$. The positive half Λ_+ consists of links with both ends in $\{x_0 \in \{\varepsilon, 2\varepsilon, \dots, \ell/2\}\}$. Fix $F_0 \in \mathbb{C}$ and $F_n \in \mathcal{D}_n$, $\operatorname{supp} F_n \subset T^n_+$, $1 \le n \le N$. Supports are compact in the open set, so there is $\delta > 0$ with $x_0 \in [\delta, \ell/2 - \delta]$ on all supports.

Step 1 (lattice form). For $\varepsilon < \delta/2$ every plaquette with centre in a support has all four links in Λ_+. Define $\Phi := F_0 + \sum_{m=1}^{N} \sum_{p_1 \dots p_m} F_m(x_{p_1}, \dots, x_{p_m}) \prod_i \varphi_{p_i}$. Φ depends only on Λ_+ links. Since φ_p is real and $P_p(\theta_\varepsilon U) = P_{\theta_\varepsilon p}(U)$, re-indexing gives $\Theta\Phi = \bar{F}_0 + \sum_n \sum_q \overline{F_n}(\theta_\varepsilon x_{q_1}, \dots, \theta_\varepsilon x_{q_n}) \prod \varphi_{q_i}$. Hence

$$0 \le \langle \Theta\Phi\, \Phi \rangle_\varepsilon = \sum_{n,m=0}^{N} S_{n+m,\varepsilon}(\Theta_\varepsilon F_n \otimes F_m), \quad \Theta_\varepsilon F_n := (\Theta F_n) \circ \tau^{\times n}_{-\varepsilon e_0},\ S_{0,\varepsilon} := 1.$$

(Argument order inside ΘF_n is immaterial by (OS3).) Each $\Theta_\varepsilon F_n \otimes F_m$ lies in $\mathcal{D}_{n+m}$: its two factors have supports at positive distance and each factor is flat on its own diagonals.

Step 2 (seam shift). $\|\Theta_\varepsilon F_n \otimes F_m - \Theta F_n \otimes F_m\|_{s(n+m)} \le 2^{s(n+m)} \cdot 4n\,\varepsilon\, \|F_n\|_{s(n+m)+1} \|F_m\|_{s(n+m)}$. By (H-U) the difference of the $S_{n+m,\varepsilon}$ values is $\le \sigma_{n+m}$ times this, which $\to 0$.

Step 3 (limit). Along the subsequence of Proposition 4.4, $S_{n+m,\varepsilon_i}(\Theta F_n \otimes F_m) \to S_{n+m}(\Theta F_n \otimes F_m)$ for each of the finitely many (n, m). So $\sum_{n,m} S_{n+m}(\Theta F_n \otimes F_m) = \lim_i \sum_{n,m} S_{n+m,\varepsilon_i}(\Theta_{\varepsilon_i} F_n \otimes F_m) \ge 0$.

Lemma 4.5 (positivity of a finite form survives pointwise limits [PROVED HERE])**.** *Let $a^{(i)} \in \mathbb{C}^{(N+1)\times(N+1)}$ with $\sum_{n,m} \bar{c}_n c_m a^{(i)}_{nm} \ge 0$ for all c and all i, and $a^{(i)}_{nm} \to a_{nm}$ for every (n, m). Then $\sum \bar{c}_n c_m a_{nm} \ge 0$ for all c.*

Proof. For fixed c the left side is a finite linear combination of the entries, so it converges to $\sum \bar{c}_n c_m a_{nm}$; $[0, \infty)$ is closed. □

Step 3 is this lemma with $a^{(i)}_{nm} := S_{n+m,\varepsilon_i}(\Theta_{\varepsilon_i} F_n \otimes F_m)$ (replace F_n by $c_n F_n$), plus Step 2.

4.5.8 (OS3) Symmetry and reality [PROVED HERE]

$S_{n,\varepsilon}(F \circ \pi) = S_{n,\varepsilon}(F)$: the φ_p are commuting real random variables; re-index the sum. For real F, $S_{n,\varepsilon}(F) \in \mathbb{R}$. Both identities are linear and pass to the limit. $S_0 = 1$ by definition.

4.5.9 (B-NT) Nontriviality: what is needed

- A two-point function alone never excludes a Gaussian: every positive-definite S_2 is the covariance of some Gaussian generalized field.
- The plaquette density is a composite ($\approx \frac{1}{4} \operatorname{tr} F^2 \times \varepsilon^4$). Even in the free theory its connected $n \ge 3$ functions are nonzero (Wick squares). So $S^c_n \ne 0$ for some $n \ge 3$ does not prove interaction.
- The meaningful test excludes (a) the zero theory and (b) the free Maxwell composite. The free composite has, at short distance, the scale-invariant form $c_0 d^{-8}$. Asymptotic freedom predicts $c\, g(d)^4 d^{-8}$ with a running coupling.

Free-theory prediction along this sequence. At tree level with the bare coupling, $\langle \varphi_p \varphi_{p'} \rangle \approx c\,\varepsilon^8 g_0^4 |x-y|^{-8}$. Since $g_0(\varepsilon)^2 \le (1/g^2 + \beta \log \varepsilon^{-1})^{-1} \to 0$ by (0.31), the bare-coupling prediction for S_2 at fixed separation tends to 0. The claim is that the true S_2 does not.

Statement (NT) needed [HYPOTHESIS]; owned by (E14-f), a corollary of (E14-a) at a fixed layer. There are $c > 0$ and C such that for all K, all $f_1, f_2 \in C^\infty(T^4)$ with disjoint supports at distance $\ge d$, $d \ge L\varepsilon$, and $k(d)$ the layer with $L^k\varepsilon \le d < L^{k+1}\varepsilon$:

$$S_{2,\varepsilon}(f_1 \otimes f_2) = c \iint f_1(x) f_2(y)\, g^4_{k(|x-y|)} |x-y|^{-8}\, dx\, dy\, (1+R), \quad |R| \le C(g^2_{k(d)} + (d/\ell)^2).$$

c is the one-loop (tree-level in the composite) coefficient: $c \propto \sum (\text{free field-strength covariance})^2$, a sum of squares, so $c > 0$ [DERIVED, STANDARD]; its value must be computed in Balaban's normalization.

Derivation of Theorem 4.3 from (NT) and (B-T2) [PROVED HERE, GIVEN (NT)]. (i) Take $f_1, f_2 \ge 0$, not identically 0, with supports at distance $\ge d_0$. Corrected condition (Section 4.8, item 3): the support diameters are $\ll \ell$, so that $g^2 + (|x-y|/\ell)^2$ is small on the whole support pair and $|R| \le \frac{1}{2}$. By (0.31) at layer $k(d)$ with $d \in [d_0, \ell]$: $1/g_k^2 \le 1/g^2 + \beta' \log(L^k\varepsilon)^{-1} \le 1/g^2 + \beta' \log(L/d_0)$, uniformly in ε. So $S_{2,\varepsilon}(f_1 \otimes f_2) \ge \frac{1}{2} c (1/g^2 + \beta' \log(L/d_0))^{-2} \iint f_1 f_2 |x-y|^{-8} > 0$ uniformly in ε, and the limit inherits the bound. (ii) For bumps concentrated at separation $d \to 0$ (with d a fixed multiple of their width), $d^8 S_2 \le (1 + C g^2_{k(d)})\, c\, (1/g^2 + \beta \log(1/d))^{-2} \to 0$ as $d \to 0$ by the lower bound in (0.31), while $d^8 S_2 > 0$ by (i). A scale-invariant composite would give a nonzero constant limit. So the limit is neither zero nor the free composite. Only the two-sided bound (0.31) is used. Caveat. (ii) needs (NT) uniformly down to $d \ge L\varepsilon$ with the remainder controlled by $g^2_{k(d)}$, i.e. the fixed-layer expansion at every layer. If (E14-f) is proved only at a fixed layer j, (i) holds, and (ii) holds only in the weaker form "$d^8 S_2(d)$ is bounded between the two (0.31) envelopes for $d \in [L^{-j}, 1]$".

4.6 Dependency table

The table gives the status of every block of Theorem B and of every named hypothesis of Theorem A. It follows the state recorded in `PLAN.md` (through the entry `SOURCES_S2_G1_V7_VERIFY.md`, 2026-10-28) and §10 of `THEOREM2_PROOF_v4.md`. Global conditions carried by every row: (GC1) Balaban's published proofs CMP 95–122 (1984–89) are correct as published; nothing in this paper re-checks them line by line. (GC2) Balaban's inductive assumptions (CMP 109 pp. 263–264) hold with the stated uniformity. (GC3) L odd, $L > 11$, M large, γ small as Balaban requires; ℓ even (sufficient, Section 2.5). Rows "item n" are the named hypotheses of Section 3; n is their item in Section 7. Their file columns use the short names of `THEOREM2_PROOF_v2_VERIFY` §K (KAPPA = `KAPPA_BOUNDED`, H_VERIFY = `BETA_FP_H_VERIFY`, PW_VERIFY = `BETA_FP_PW_VERIFY`, v1 verifier = `THEOREM2_PROOF_VERIFY`).

Block	Content	Current tag	Files	Hypotheses carried
B-T2	Theorem 2w (Theorem 3.1): trajectory $g_0(\varepsilon, g)$, (b), (c′), (S)	[PROVED+VERIFIED, CONDITIONAL] (Route K = Theorem K + Lemma D″). Literal (0.31) not obtained	`THEOREM2_PROOF_v4.md`, `KAPPA_BOUNDED.md`	(H2), (PI), (F), (M_j), V1, (G2), L5, item 7′ (from (H0) or (G-gain), verified (`SOURCES_GGAIN_NONMONO_VERIFY`))
B-T2.a	Lemmas D, D′, D″, E (shooting)	[PROVED+VERIFIED; LEAN] (D′, D, E, D″ one-argument form; history form open)	`LEMMA_DPRIME.md`, `lean/`	continuity of β_{k+1} in history
B-T2.b	Sweep: (c′)+(S) ⇒ Balaban's Theorem 1 inputs (Section 3.4.3; item 10)	[PROVED, UNVERIFIED]; 4 of 19 rows verified (LEMMA_DPRIME_VERIFY Steps 4–5, rows 11, 12, 16, 18)	`LEMMA_DPRIME_SWEEP.md`	Balaban readings (unchecked list)
B-T2.1	U3 Thm 4.1 (Gaussian RG, rate L^{-2j}) + O5	[PROVED+VERIFIED]	`U3_GAUSSIAN_RG.md`, `U3_REPAIRS.md`	trivial background
B-T2.2	U3 Lemma 5.1; (A), (W)	neutral sector [PROVED+VERIFIED]. (W) numerically exact; (A) confirmed qualitatively and quantitatively (`U3_AW_VERIFY.md`); (HS) [REFUTED] at $L = 3$. Not on the path of Theorem 2w after (H2); needed for the convergence route only	`U3_LEMMA51.md`, `U3_AW.md`	(A) in full, (N) (plausibly false)
B-T2.3	O6 bordered-matrix invertibility	one-step [PROVED+VERIFIED]; composed constraint reduced to one-step O6 + (A)	`U3_REPAIRS.md`	—
B-T2.4	$c = 11N^2/(12\pi^2)$, anchor-free identity Prop. 2.1, telescoping	[PROVED+VERIFIED]. $\beta_\infty = c \log L$ itself replaced by window sums (Theorem K, Route B)	`BETA_FP_LEMMA43.md`, `BETA_FP_LEMMA43_VERIFY.md`	(F) [CITED, SOURCE CHECKED]
B-T2.5	κ_k bounded (Theorem K)	[PROVED+VERIFIED, CONDITIONAL ON (B2)]; (B2) [PROVED+VERIFIED, CONDITIONAL ON (H2)]	`KAPPA_BOUNDED.md`, `KAPPA_BOUNDED_VERIFY.md`, `B2_TWOSTEP.md`, `B2_TWOSTEP_VERIFY.md`	(H2), (PI), (F), (M_j), V1, (G2)

Block	Content	Current tag	Files	Hypotheses carried
B-T2.H2	(H2) uniform two-step inequality $\rho_2 < L$ (Section 3.7.3; item H)	[CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)] at $L = 13$: $\rho_2 \le \sup \rho_1^2 \le 10 < 13$ (Theorem 6.1); residue: reading of (0.4), gauge choice, Prop. A at $M \ge 27$, Lemma G beyond point checks, `bb_arb.py` audited only. At $L = 3, 5, 9$ [NUMERICAL]; point enclosure $[1.9993, 2.1273] < 3$ at $L = 3$ [INTERVAL ENCLOSURE, VERIFIED]	`H1_ANALYTIC.md`, `H1_ANALYTIC_VERIFY.md`, `H1_ANALYTIC_VERIFY2.md`, `H2_CERTIFY_P5_VERIFY.md`, `H2_LARGE_L_VERIFY.md`, `H2_CERTIFY_VERIFY.md`; B2_TWOSTEP §2	— (box sweep at $L = 13$ running as a cross-check)
B-T2.H1	one-step route at $L = 13$: $\sup \rho_1^2 < L \Rightarrow$ (H2)	[CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]: $\sup_{j,K} \rho_1^2 \le 9.9995 < 13$ by the majorant B_{13} and `bb_arb.py`; factorization [PROVED+VERIFIED; LEAN] (`rho2_le`); majorant core [LEAN] (`Tower.lean`)	`H1_ANALYTIC.md`, `H1_ANALYTIC_VERIFY.md`, `H1_ANALYTIC_VERIFY2.md`, `H2_CERTIFY_VERIFY.md`	—
B-T2.B	Route B (Theorem B_{PW}; item B8)	logic [PROVED+VERIFIED, CONDITIONAL] (tag decision, `PLAN.md`; PW_VERIFY §6 items 1–4); inputs (U), (DF), (T1) [PROVED, UNVERIFIED]	`BETA_FP_PW.md`, `BETA_FP_PW_VERIFY.md`, `BETA_FP_U.md`	(H2), ($B\theta$), A6, LW (5.7), (H0)
item 1	(B2) / (B2w) (Section 3.7)	[VERIFIED REDUCTION] to (H2); holds at $L = 13$ (Theorem 6.1); open at other L	KAPPA §2, Rem. 2.3; KAPPA_VERIFY §7–§9	(H2)
item 1	(H2) $\Rightarrow$ ($C1\sigma$) $\Rightarrow$ (B2) (Theorems 3.10, 3.11)	proved+verified, conditional on (H2)	B2_TWOSTEP §2–§3; B2_TWOSTEP_VERIFY grades 1, 3, 4	(H2)
item 1	($C1\sigma$) + (h) $\Rightarrow$ (B2)	implication verified; ($C1\sigma$) $\Leftarrow$ (H2); (h) numerical $L = 3, 5$	KAPPA Prop. 4.1; KAPPA_VERIFY §7	—
item 1	(H″) $\Rightarrow$ (A) $\Rightarrow$ (B2)	implication argued; not needed after (H2)	U3_AW §4.7; no verifier (U3_AW_VERIFY grade 5 covers (H′) only)	—
—	(HS) (Section 3.7.2)	refuted $L = 3$	U3_AW (H′); U3_AW_VERIFY §6	—

Block	Content	Current tag	Files	Hypotheses carried
item H	$\rho_1^2 < L \Rightarrow$ (H2) (one step, $L \geq 5$)	arithmetic; $\rho_1^2/L = 0.771$ at $L = 5$, numerical	H2_LARGE_L §3; H2_LARGE_L_VERIFY grade 4 (measurement)	—
item H	certification tail j_0	derivation confirmed; $j_0 \approx 8$ ($L = 3$), ≈ 6 ($L = 5$), corrected	H2_LARGE_L §4.1; H2_LARGE_L_VERIFY grade 6	—
item 2	(PI)	argued	BETA_FP_H §3.1; H_VERIFY §5	—
item 3	(F)	cited, source checked; round 7: Step B reduced [R], unverified, to (L-LL), a kernel bound, and Step C's gauge-fixing part; open tension β_1 vs $c \log L$ (Section 6.8)	BETA_FP_LEMMA43; H_VERIFY §4	—
item 4	(M_j): (S1)–(S3), Prop. 4.4	(S1), (S3), (S2)$\|_{t=0}$, (S2c) proved at fixed j given (I3); (I3) [PROVED+VERIFIED] (`SOURCES_T2_I3_VERIFY`, round 5), given the identification of the one-step objects; (M_j) [VERIFIED REDUCTION] to (I3), Prop. 4.4 region 1 (argued), and the neutral sector [I]; round 6: (G'_A) [PROVED+VERIFIED] for $0 < \|f\| \leq ((1-c)/100)^2/2$ (Theorem 5.12); round 7: (M_j) [PROVED RELATIVE TO LISTED INPUTS; VERIFIED] at fixed j (Theorem 5.14), the proved f-range suffices	BETA_FP_M §4; `SOURCES_T2_S123.md`, `SOURCES_T2_S123_VERIFY.md`; H_VERIFY §5 refuted the old step; `SOURCES_T2_I3.md`, `SOURCES_T2_GAPRIME2.md`	one [I] convention (KAPPA (1.1) coefficients)
item 5	V1	[PROVED+VERIFIED AT FIXED LEVEL, CONDITIONAL] (history-free)	`SOURCES_T2_V1.md`, `SOURCES_T2_V1_VERIFY.md`; R2; U3_VERIFY C4	(G2), Bałaban's inductive bounds, averaging (0.4)
item 6	(G2)	[VERIFIED REDUCTION] at fixed j (was formal)	`SOURCES_T2_G2.md`, `SOURCES_T2_G2_VERIFY.md`; BETA_FP_LEMMA43	$\bar{Q}$ onto, (Gap_f), averaging analyticity

Block	Content	Current tag	Files	Hypotheses carried
item 7′	$\lvert\beta_{k+1}(h,g) - \beta_{k+1}(0)\rvert \le C_\beta g$ on every past (was (H0), uniform in j and history)	from (H0) [BALABAN], odd $L > 11$ (its use a [VERIFIED REDUCTION], residue N1–N9, N2 = G3 proved relative to listed inputs, Section 5.9), or from (G-gain) with the running-maximum lift [PROVED+VERIFIED]	CMP 109 (1.22); `SOURCES_GGAIN_NONMONO.md`, `SOURCES_GGAIN_NONMONO_VERIFY.md`; KUNIF_VERIFY item 3; `SOURCES_T2_H0.md`, `SOURCES_G3_N2.md`	N1, N3–N9 of H0-red; (G-gain) inputs if taken from it
item 8	L5 history form and domain (Section 3.4.5)	round 7: [PROVED RELATIVE TO LISTED INPUTS; VERIFIED], Route 2 (Theorem 5.15)	LEMMA_DPRIME Rem. 3; LEMMAS_DE_VERIFY §2.1; v2_VERIFY §F	—
item 9	$\beta_j(g)$ independent of K	closed as a reduction [VERIFIED]: volume statement, β_j whole-lattice by definition, D′/D″ at fixed K; residual volume question = ledger B8, an input of Theorem 1; torus versus whole-lattice coefficient within $C_j N_j^6 e^{-m_j N_j/2}$ (Proposition 5.8; (Loc) proved in round 6, Theorem 5.10); B8 replaced and proved under (Vol) (Proposition 5.11)	`SOURCES_T2_KUNIF.md`, `SOURCES_T2_KUNIF_VERIFY.md`	(K1), (K2) = item 7′, (K3)
item 10	BALABAN_CHECK item 11 readings	verified with corrections (former ledger A12, A20, A22); A11 closed as a verified list (Proposition 5.7; C28–C30); residue: A23 (T-step rate loss)	BALABAN_EXCERPTS; `SOURCES_BOOKKEEPING.md`, `SOURCES_A11_CONST.md`	A23
item 11	odd L; $L > 11$ in Balaban	input	CMP 109 pp. 251–252; v1 verifier Step 6	—
item 12	SU(2)	scope	—	—
item 13	Lean history form	open	—	—
B1, B2′	Route B: (H2) + (T1) ⇒ (U) (Theorem 2.4 of BETA_FP_U)	proved, unverified	BETA_FP_U §2.4; no verifier	(H2), (T1)
B1, B2′	Route B: (U)	hypothesis; ⇐ (T1) + (H2) (BETA_FP_U); proved, unverified	BETA_FP_PW §1.2; PW_VERIFY §3	(T1), (H2)
B1	Route B: (T1)	argued	BETA_FP_U §2.3; no verifier	—

Block	Content	Current tag	Files	Hypotheses carried
B3	Route B: $(B\theta) = \gamma_{rec} + \mathfrak{d}_F$	γ_{rec} bounded given Thm 2.4 (proved, unverified); (DF) argued	BETA_FP_PW §3.3; BETA_FP_U §3; PW_VERIFY §6 (logic of Theorem B_{PW}); none for BETA_FP_U	(DF)
B4	Route B: LW (5.7)	cited	cited; PW_VERIFY §3 (regime correct)	—
B5	Route B: A6 (β identification, uniform in k)	argued	SOURCES_D1_VERIFY 8; PW_VERIFY §3; BETA_FP_U §4.2	—
B7	Route B: $\theta_* < 1$	trivial condition	PW_VERIFY §3	—
B-T2.6	certified early steps $\beta_j(0) > 0$	needed only for literal (0.31) at every K, not for Theorem 1; [NUMERICAL], no certificate; $\beta_1(0)$ computed to $L = 11$	`numerics/NUMERICS.md`, `UMK_BETA1.md`	$L > 11$ is Balaban's range
B-S	sourced Theorem 1 (E14-a,b,c)	[OPEN]	`SOURCES.md` §2.5	B-T2 for the sourced trajectory
B-S.D1	one sourced step, Gaussian level	[PROVED+VERIFIED] (with corrections)	`SOURCES_D1.md`, `SOURCES_D1_VERIFY.md`	—
B-S.D2	sourced small-field step (E14-a)	(a), (b), (d), (f) confirmed; (c) rejected; (e) reduced	`SOURCES_D2.md`, `SOURCES_D2_VERIFY.md`	analyticity in input activities; $\kappa \to \kappa/2$ [R, verified, B22; uses A23]; (U) [C23, constant choice]
B-S.D2C	redo of (c): frozen source per domain, Lemma A, $h \equiv 0$	Lemma A and recursion confirmed; (I) corrected (exact only with adapted gauge-fixing weight)	`SOURCES_D2C.md`, `SOURCES_D2C_VERIFY.md`	domain hypothesis (D); (Hβ)
B-S.Hβ	two-loop bound $\sup \sum_j \lvert B'_j \rvert \le \frac{1}{2}$	[PROVED+VERIFIED, CONDITIONAL] on (I), (H-hist)	`SOURCES_HBETA.md`, `SOURCES_HBETA_VERIFY.md`	(I), (H-hist)

Block	Content	Current tag	Files	Hypotheses carried
B-S.hist	(H-hist) history sensitivity	[VERIFIED REDUCTION] on (W4), Facts A/B, (I)/L3 and (G-gain) in both slots (`SOURCES_SM.md`; verifier: items 1–4 confirmed, 5–6 corrected); the (G-gain) chain continues in row B-S.Ggain (G3, (Q)); with the large-field half (`SOURCES_GGAIN_LF_VERIFY`) the real uniform (H-hist) is a [VERIFIED REDUCTION] in both halves (ledger B13); Lemma HH: `SOURCES_HHIST.md`, `SOURCES_HHIST_N1_VERIFY.md`	`SOURCES_SM.md`, `SOURCES_SM_VERIFY.md`	(G-gain), small and large field
B-S.Ggain	(G-gain): $g_k\lvert B\rvert$ gain in the small-field step	pointwise Cauchy and uniform C_β on $]0,\gamma]$ [PROVED+VERIFIED] (small-field, incl. vacuum subtraction); Lemma G4 off-chain (`SOURCES_G4.md`); χ_k surface term = Lemma G3, [PROVED RELATIVE TO LISTED INPUTS; VERIFIED] with (Q) assembled from A7, A8 (Theorem 5.9, round 5); earlier route: (Q) reduced to CMP 99 (3.132), (E_U), and the localization lemma LOC [VERIFIED REDUCTION] (`SOURCES_Q_VERIFY`, `SOURCES_CDC_VERIFY`, `SOURCES_EU_VERIFY`, `SOURCES_LOC_VERIFY`); large field [VERIFIED REDUCTION] (one Cauchy step on the large-field disc, `SOURCES_GGAIN_LF_VERIFY`); non-monotone pasts: running-maximum lift (`SOURCES_GGAIN_NONMONO.md`, verified (`SOURCES_GGAIN_NONMONO_VERIFY`))	`SOURCES_GGAIN.md`, `SOURCES_GGAIN_VERIFY.md`, `SOURCES_GGAIN_LF.md`	(M_r), C25, C26, CMP 102 (189), (M-B) bookkeeping; not at $g = 0$

Block	Content	Current tag	Files	Hypotheses carried
B-S.G3	Lemma G3: χ_k surface term, uniform in g	[PROVED+VERIFIED, CONDITIONAL] on (Q) (Route 2 surface estimate; the B/B′ cut closed in `SOURCES_Q`); round 5: [PROVED RELATIVE TO LISTED INPUTS; VERIFIED] (Theorem 5.9; G_Γ bounded uniformly in M, verified)	`SOURCES_G3.md`, `SOURCES_G3_VERIFY.md`, `SOURCES_G3_N2.md`, `SOURCES_G3_GGAMMA.md`	A7, A8, (2.15)–(2.25) chain, C25, C26, N3, N6, N9, (M_r); [I] s-rule reading
B-S.Q	(Q): covariance lower bound $c_- > 0$	[VERIFIED REDUCTION] to CMP 99 (3.132) (printed; ledger A1, kernel-decay form now a verified reduction via Lemma CT to CMP 99 Thms 3.1, 3.3, 3.12, `SOURCES_S2_FORM_VERIFY`), (E_U), and LOC; Woodbury identity confirmed (CDC); free-case constants [NUMERICAL] (`numerics/kappa_d.py`); round 5: (Q) in the joint (X, B) form from A7 + A8 (Theorem 5.9), with Λ corrected (G_Γ enters)	`SOURCES_Q.md`, `SOURCES_Q_VERIFY.md`, `SOURCES_CDC.md`, `SOURCES_CDC_VERIFY.md`, `numerics/KAPPA_D.md`	A1 residue (torus-side hypothesis, conventions, reading $a = 1$; G_1 clause closed, `SOURCES_S2_G1_V7_VERIFY`); two [I] bridges
B-S.EU	(E_U): uniform block extension	$K_0(4) = 195.19$ ($N = \infty$) [NUMERICAL, REPRODUCED]; $K \le K_0(1 + c_{2/3}(M\alpha_0)^{2/3} + c_1 M\alpha_0)$ [DERIVED SKETCH]; (E_U) [VERIFIED REDUCTION] to kernel-moment bounds Bałaban does not print	`SOURCES_EU.md`, `SOURCES_EU_VERIFY.md`, `numerics/k0_d.py`	k-uniform kernel moments; unprinted C_1'
B-S.LOC	LOC: localized covariance $C^{(k)}(Z_0, 0)$	[PROVED+VERIFIED, CONDITIONAL] (short lemma; symmetry (S) not needed)	`SOURCES_LOC.md`, `SOURCES_LOC_VERIFY.md`	R_2 bound (2.16) (ledger A2, verified reduction with the V7 repair to (W^c) in the exponential gauge + A7 at real U + two [I]); γ_0 for general U (ledger A3, full proof verified, $\gamma_0 \ge m_L/(48d^2L^d)$); α_0, α_1 small

Block	Content	Current tag	Files	Hypotheses carried
B-S.thr	(H-thr) threshold shell sum	Lemma T and trajectory sum [PROVED+VERIFIED], conditional on (H-hist2); (H-hist2) [VERIFIED REDUCTION] to the real uniform (H-hist), no new input (SOURCES_HHIST2, SOURCES_HHIST2_VERIFY)	SOURCES_HTHR.md, SOURCES_HTHR2.md, SOURCES_HTHR2_VERIFY.md	(H-hist2), M3 weights, frozen integers (L3)
B-S.hist2	(H-hist2): mixed second history derivatives	[VERIFIED REDUCTION] to the real uniform (H-hist), no new input (Route 2(a); decay $\lambda_C^{m/2}$; the product form λ^m [REFUTED])	SOURCES_HHIST2.md, SOURCES_HHIST2_VERIFY.md	real uniform (H-hist)
B-S.D3	sourced large-field step + R operation (E14-b)	draft verified with corrections (SOURCES_D3_VERIFY, SOURCES_D3R_VERIFY, SOURCES_D3W8_VERIFY; route (a), Lemma F2′ confirmed given (W4), (Eps), F1, (D)). (FX) is a genuine new hypothesis (SOURCES_D4D5_VERIFY.md); its state is in row B-S.FX	SOURCES_D3.md, SOURCES_FX.md, SOURCES_FX_VERIFY.md	(FX), (FX2); (U) [C23, constant choice]

Block	Content	Current tag	Files	Hypotheses carried
B-S.FX	O1(b), O2, O3, O4, R2	O1(b) [PROVED+VERIFIED] (Lemma [P], no additive constant); O2 [PROVED+VERIFIED] (the R-denominator is the bare Wilson integral (1.100)); O3 [PROVED+VERIFIED, CONDITIONAL ON N3] (N3 = the open HBETA strip bound $\|\delta w\| \le \frac{3}{2}\tau$); R2 ($\kappa_{fin} \ge \kappa_{final}$) [PROVED+VERIFIED] as a reduction, in volume form, given $R_{K+1} \ge 1$ and C_c independent of g; O4 (unbounded components in (1.103)) avoided by keeping Balaban's main form (1.98), whose fallback was Gevrey $2+\eta$. N3 reduced to (H-hol) + real (H-hist) + (H2-hist), condition (a) settled [VERIFIED REDUCTION] (`SOURCES_HBETA_N3_VERIFY`, `SOURCES_N3A_VERIFY`); (H-hol) small field [VERIFIED REDUCTION] (`SOURCES_HHOL_VERIFY`); large field [VERIFIED REDUCTION] to (M-A), (M-B), (M-C), (H-hist2) (`SOURCES_HHOL_LF_VERIFY`); (M-B) [PROVED+VERIFIED] with $p_0 \ge 17/(\gamma_0 A_1^2)$, $\theta_W + \tau_C \le 0.2$; (M-A) reduced to (H−), (M-A′) closed, (H−) rests on CMP 102 (189) [VERIFIED REDUCTION] (`SOURCES_MA_VERIFY`, `SOURCES_MA2_VERIFY`); (M-C) [VERIFIED REDUCTION] to printed CMP 116 §2 conditions (`SOURCES_MC_VERIFY`); $R_{K+1} \ge L$ and C_c free of g are verified readings (`SOURCES_BOOKKEEPING.md`)	`SOURCES_FX_O1.md`, `SOURCES_FX_O1_VERIFY.md`, `SOURCES_FX_O23.md`, `SOURCES_FX_O23_VERIFY.md`, `SOURCES_FX_R2.md`, `SOURCES_FX_R2_VERIFY.md`	residual inputs of (M-C); CMP 102 (189) (ledger A4, verified reduction to A1)

Block	Content	Current tag	Files	Hypotheses carried
B-S.N3	N3: HBETA strip bound $\lvert\delta w\rvert \le \frac{3}{2}\tau$	[VERIFIED REDUCTION] to (H-hol) + real (H-hist) + (H2-hist)	`SOURCES_HBETA_N3.md`, `SOURCES_HBETA_N3_VERIFY.md`	(H-hol), (H-hist), (H2-hist)
B-S.N3A	condition (a) of Lemma N3	settled [PROVED+VERIFIED] (slack $> 5\times$; item 1 corrected)	`SOURCES_N3A.md`, `SOURCES_N3A_VERIFY.md`	constant order $L \to$ Bałaban's constants $\to \gamma \to g \le \gamma/2 \to r_0 \to K$
B-S.hol	(H-hol), small field: β_j holomorphic on $\lvert g - x\rvert < \rho_0 x$	[VERIFIED REDUCTION] to Bałaban's conditions with modified constants	`SOURCES_HHOL.md`, `SOURCES_HHOL_VERIFY.md`	(C1′)–(C4′), (C4″), (2.24′)
B-S.holLF	(H-hol), large field (frozen thresholds at x^*)	[VERIFIED REDUCTION] to (M-A), (M-B), (M-C), (H-hist2); (M-C) itself a verified reduction	`SOURCES_HHOL_LF.md`, `SOURCES_HHOL_LF_VERIFY.md`	(M-A)–(M-D)
M-A	equipartition $\langle A - A_{\min}\rangle_u \le (n_i/u)(1+\eta_x)$	[PROVED+VERIFIED] given (H−); (M-A′) closed [PROVED+VERIFIED]; (H−) [VERIFIED REDUCTION] to CMP 102 (189), an existing trust item	`SOURCES_MA.md`, `SOURCES_MA_VERIFY.md`, `SOURCES_MA2.md`, `SOURCES_MA2_VERIFY.md`	CMP 102 (189); (C1), (C2)
M-B	(1.80)–(1.98) bookkeeping, changed constants	[PROVED+VERIFIED] with $p_0 \ge 17/(\gamma_0 A_1^2)$, $\theta_W + \tau_C \le 0.2$	`SOURCES_MB.md`, `SOURCES_MB_VERIFY.md`	(M-A), (M-C)
M-C	complex small-field step	[VERIFIED REDUCTION] to printed CMP 116 §2 (B1–B23 + 4), Lemma 2, (1.36), p. 12 (2.5)–(2.6); shrinks $\varepsilon_1/(4\lambda_V)$, $\gamma/(8\lambda_V)$; (I-X) resolved; (I-V0), (I-ord), and (I-ε) in both layers closed [PROVED+VERIFIED]; (M1) is convention C24	`SOURCES_MC.md`, `SOURCES_MC_VERIFY.md`, `SOURCES_MC_IV0.md`, `SOURCES_T2_IEPS.md`, `SOURCES_T2_IEPS_LF.md`, `SOURCES_HIGHER_M1.md` (each + VERIFY)	[I] residue: unprinted α_2–α_4 scaling, q_0 vs p_0 (every branch closes)
M-D	frozen map vs Bałaban's map	not a convention: needs the frozen map's derivative at x^*, which is (H-hist) + (H-thr) [VERIFIED REDUCTION]	`SOURCES_HHOL_LF_VERIFY.md`	(H-hist), (H-hist2)
B-S.D4	real-source derivative bounds (E14-c)	D4.1, D4.3, D4.4 and Lemma C (order $1+\eta$, given (FX)) [PROVED+VERIFIED]; (S_{src}) $\Rightarrow$ (H-U), $s = 6$, $D > 1$, confirmed given (FX)	`SOURCES_D4D5.md`, `SOURCES_D4D5_VERIFY.md`	(FX), (FX2), S1 at all scales

Block	Content	Current tag	Files	Hypotheses carried
B-N1	source form ⇒ (H-U) on $\mathcal{D}_n$	derived given the sourced Gevrey bound (S) (`ASSEMBLY_N1.md`; confirmed, `SOURCES_HHIST_N1_VERIFY.md`); (S_{src}) ⇒ (H-U), $s = 6$, $D > 1$, confirmed given (FX); uniformity of (S) in support size and ε [VERIFIED REDUCTION] (h-independent; `SOURCES_B21_S_VERIFY`); ASSEMBLY_N1 aligned 2026-10-28 with the C^1 norm, $(n!)^{1+\eta}$, $s = 6$, $D > 1$ ($D > 2$ if the coarsest-scale (FX2) radius fails); open: the (E14-c) clause	this paper, `SOURCES_D4D5.md`, `SOURCES_B21_S.md`	(FX) main form, coarsest-scale (FX2) radius, S1 at all scales, Lemma C
B-C	compactness (diagonal extraction)	[PROVED HERE]; verifier: correct	Proposition 4.4	(H-U)
B-REG	(OS0)	[PROVED HERE]; verifier: correct	Section 4.5.5	(H-U); torus form of OS II
B-TI	(OS1) all of T^4 and B_4	[PROVED HERE]; verifier: correct	Section 4.5.6	(H-U); origin at a site
B-RP	(OS2)	lattice: [CITED]; passage: [PROVED HERE]; verifier: correct	Section 4.5.7	ℓ even (sufficient); (H-U)
OS3	permutation symmetry, reality	[PROVED HERE]	Section 4.5.8	—
B-NT	nontriviality (Theorem 4.3)	[OPEN]: needs (NT) = (E14-f); derivation from (NT) [PROVED HERE], corrected statement	Section 4.5.9	(NT) at every layer for (ii); $c > 0$ in Balaban's normalization
E13	full-sequence convergence	not needed here; [OPEN]	`SOURCES.md` (E14-d), D6	—

Critical path to an unconditional (modulo GC1–GC3) Theorem 4.2: (H2) is certified at $L = 13$ (Theorem 6.1); B-T2 still needs (PI), (F) (reduced, unverified, round 7), the inputs of (M_j), V1, and (G2) (fixed-level results, Sections 5.1 and 5.11), the inputs of L5 (Theorem 5.15), and item 7′, and the step to Theorem 1 needs the sweep (B10, cleared in round 7; A11 is a verified list) and the finite-volume identification B8 (replaced form, proved relative to C31–C37 in round 6; A15 a verified reduction to conventions); then B-S: the open inputs of the real uniform (H-hist) ((M_r), CMP 102 (189), (M-B) bookkeeping; the small-field (Q)/c_- is discharged relative to A7, A8, C25, C26 by Theorem 5.9), the residue under A1–A4 (the inputs of Theorems 5.1, 5.2, 5.3: (C), the printed sketches of CMP 99 Thms 3.7, 3.9, 3.10, WG-1, and [I] geometry and convention items; the inputs of Theorems 5.4, 5.5, 5.6; R-Δ2; the residue of A18 (Proposition 5.18); the torus-side hypothesis), M2′, M4, M5 of (H-thr), the Λ_j bookkeeping of B20, and the (E14-c) clause of B-N1. The blocks proved in this section are short and are not on the critical path.

4.7 What the theorem does not give

Each item needs a separate argument that this paper does not give.

- Rotation invariance. Only B_4 and the torus translations are proved. Full $O(4)$ is not a symmetry of T^4_ℓ. `LEMMAS.md` ledger item 5 [VERIFIED QUOTE]: on the torus the inclined comparison "gives only the tilted-axis reflection symmetry $R_{2\theta}s$, never a rotation"; the template reaches rotations only after the torus $\to$ plane limit, which it controls with its mass gap.

- Infinite volume. ℓ is fixed. Jaffe–Witten tie the limit $T^4 \to \mathbb{R}^4$ to a volume-uniform gap [VERIFIED QUOTE]: "New ideas are needed to prove the existence of a mass gap that is uniform in the volume of space-time. Such a result presumably would enable the study of the limit as $T^4 \to \mathbb{R}^4$." No public mechanism without a gap is known to us (`LEMMAS.md` ledger item 8).

- Mass gap. Not addressed. On a finite torus a "gap" of the transfer matrix is not the Clay mass gap.

- Uniqueness. Different subsequences may give different Schwinger functions. Uniqueness needs E13, which Balaban does not have. Also not given: uniqueness of the vacuum after reconstruction (no clustering on a torus).

- Vacuum state. Time is periodic, so OS reconstruction from torus reflection positivity gives a transfer matrix on T^3 and a thermal state at inverse temperature ℓ, not a vacuum state (Section 4.8, item 4).

- Also not given: clustering (OS4); a probability measure for the density field; Wilson-loop observables (class C3); the stress tensor and the operator product expansion (class C2 and beyond); independence of the limit from the block-averaging choices of Balaban's RG (only the lattice measure enters $S_{n,\varepsilon}$, but $g_0(\varepsilon)$ is chosen through Balaban's flow).

- Status of the whole statement. Even with every block closed, the theorem is conditional on Balaban's published 1984–89 proofs (GC1). Under Jaffe–Witten footnote 2 it is a compactness ("weak-existence") result on a torus with the torus forms of OS0–OS3; it is not a solution of the Millennium problem.

4.8 Corrections after verification

The verifier confirmed the theorem statement, all in-chapter proofs (B-C, B-REG, B-TI, B_4, the RP passage, OS3, the finite-form lemma), the quotes of Jaffe–Witten, and the open status of B-N1. Four items were corrected (`ASSEMBLY_VERIFY.md`). The statements above carry the corrected wording; the original claims are listed here.

1. *ℓ even is sufficient, not necessary* (verifier §5, grade 3). The parity count is right: L odd gives $N_\varepsilon = \ell L^K$ even iff ℓ even; for odd N_ε every time reflection has one site seam and one link seam. Osterwalder–Seiler (1978) [21] treat link-plane (between-site) reflections. Site-plane reflections were added by Menotti–Pelissetto, CMP 113 (1987) 369–373 [17] [VERIFIED QUOTE] (abstract, per the verifier): "We extend the proof by Osterwalder and Seiler of reflection positivity of lattice gauge theories with fermions to the case in which the reflection is performed with respect to planes containing sites." and "Our proof applies to all observables and relies on the particular structure of the Wilson action." The mixed case (one site seam, one link seam

on a periodic lattice) is very likely routine, since each mechanism is local to its seam, but no citation states it. So odd ℓ is not excluded; the hypothesis ℓ even only avoids the mixed case. The source's phrase "plausible but not cited" stands. The periodic two-seam link version attributed to Osterwalder–Seiler is [CITED], not [VERIFIED QUOTE]. Two minor points from the verifier: the support distance after the seam shift is $\geq 2\delta - \varepsilon$ (the source's $2\delta - 2\varepsilon$ is conservative); the factor $2^{s(n+m)}$ in Step 2 is unneeded but harmless.

2. *Reference title.* The Jaffe–Witten reference list gives the title "Gauge theories on the lattice"; the bibliography entry follows the Annals of Physics title.

3. *Nontriviality needs the two-sided envelope only and support diameters $\ll \ell$* (verifier §6, grade 6). In Theorem 4.3(i), $|R| \leq \frac{1}{2}$ needs $g^2 + (|x-y|/\ell)^2$ small on the whole support pair, so the support diameters must be $\ll \ell$, not only the minimum distance d_0. In Theorem 4.3(ii) the source stated the asymptotic $d^8 S_2 \sim c\, g(d)^4$ with $g(d)^2 \sim (\beta \log(1/d))^{-1}$. The proof uses only the two-sided bound (0.31); the asymptotic needs $\beta_j \to \beta_\infty$ and is not proved. The statement of Theorem 4.3(ii) above is the supported one. The overclaim was in the statement, not in the proof.

4. *Periodic time gives a thermal state* (verifier §6, grade 8). OS reconstruction from torus reflection positivity gives a transfer matrix on T^3 and a thermal state at inverse temperature ℓ, not a vacuum. This remark is added to the list of non-results (Section 4.7).

5 The sources block

This section gives one paragraph for each project file `SOURCES_*.md` that has a verifier file. Each paragraph states the result, describes the method in two sentences, gives the verifier's grade in the tags of Section 1.4, and names the inputs that stay open. Nothing here goes beyond the project ledgers: the verifier files, `TRUST_LEDGER.md`, the dependency table (Section 4.6), and `PLAN.md`. Where a verifier corrected a file, the corrected statement is given. The grouping follows §4 of `TRUST_LEDGER.md`: first the files that Theorem 2w (Target 1) uses, then the files of the finite-torus theorem (Target 2) in the order of its dependency chain (steps 2–9 of §4.2 there). Ledger labels An, Bn, Cn, Xn are those of `TRUST_LEDGER.md`. `SOURCES_A1_KERNEL.md` and `SOURCES_A4_D189.md` share the verifier `SOURCES_A1_A4_VERIFY.md`, and `SOURCES_HHIST.md` is checked in `SOURCES_HHIST_N1_VERIFY.md`. Three files have no verifier and are not listed: `SOURCES_BOOKKEEPING.md`, `SOURCES_G4.md`, `SOURCES_W4.md`. Files verified after this section was assembled have no paragraph of their own: `SOURCES_WC_EXPGAUGE.md` and `SOURCES_WC_RESIDUE.md` (cited under A1w and A2), and `SOURCES_T2_V1.md`, `SOURCES_T2_PI.md`, `SOURCES_T2_A1WO3.md`, `SOURCES_T2_A1WO4.md`, `SOURCES_T2_D189.md`, `SOURCES_T2_R2REM.md`; their state is in `TRUST_LEDGER.md` and `PLAN.md`.

5.1 Target 1: inputs of Theorem 2w

`SOURCES_GGAIN_NONMONO.md`: item 7′, the running-maximum lift. For every step k, every past $h \in]0,\gamma]^k$, monotone or not, and every $g \in [0,\gamma]$, $|\beta_{k+1}(h,g) - \beta_{k+1}(0)| \leq C_\beta g$ with $C_\beta = C_\Pi S_0$, the constant of (G-gain), uniform in k, K, and the past. This removes the monotone-history restriction of (G-gain), which blocked item 7′ on the clamped, non-monotone pasts that the shooting evaluates. The only use of monotonicity is Step 2 of Lemma F2′, where $\max_{j\leq k} 4g_j^{\kappa_0} = 4g_k^{\kappa_0}$ compares bound sizes; with the running maximum $G_k = \max_{i\leq k} g_i$ the induction $a_j \leq 2G_j^{\kappa_0}$ closes for every past under the side condition $4\gamma^{\kappa_0} \leq E_0$, which is of Bałaban's existing type. The

value at zero own coupling, (K1), is history-free because the exponent vanishes at $g_k = 0$ (CMP 109 p. 268) for any input, and the endpoint $g \to 0$ is a dominated-convergence step that is not written line by line. Verifier: `SOURCES_GGAIN_NONMONO_VERIFY.md`, the lift [PROVED+VERIFIED], item 7′ [PROVED+VERIFIED, CONDITIONAL ON (H0) (LEDGER A5) OR ON THE (G-GAIN) INPUTS (LEDGER B13)]; all four items confirmed with small corrections: $C'_\beta = C_\beta$ holds because $E_0 + 4\gamma^{\kappa_0} \leq 2E_0$, the input radius that (G-gain) already covers, and the possible dependence of the newest R-input on g (caveat c1) is open on the literal reading of β and void on the pure-part reading. Open inputs: (H0) (ledger A5; its use here is a verified reduction with residue N1–N9, Section 5.9), or the (G-gain) inputs of ledger B13 ((Q)/c_-, (M_r), CMP 102 (189), the (M-B) bookkeeping); L5, the evaluation of β off Bałaban's trajectory (item 8, ledger B7); caveat c4, CMP 119 (2.31) and the off-trajectory small-field induction of `SOURCES_SM.md` §5.1 not swept for coupling-order uses.

`SOURCES_T2_KUNIF.md`: item 9, K-independence of $\beta_j(g)$. Item 9 is a volume statement, not a history identity, and it closes as a reduction: CMP 119 (2.24), p. 253, and p. 279 define β_j as the whole-lattice coefficient of CMP 109 (whose Π in (4.37) is a sum over $\mathbb{Z}^4$), so the β_j of the recursion is K-free by definition. The shooting needs only (K1) $\beta_j(h, 0) = \beta_j(0)$ for every admissible past, (K2) $|\partial\beta_j/\partial g_{j-1}| \leq C_\beta$, and (K3) the window bound $\sum_{i=k+1}^{n} \beta_i(0) \in [b(n-k) - 2R,\, b(n-k) + 2R]$ of Theorem K; with $\gamma \leq b/(4C_\beta)$ these give $T(m, n) \in [(3b/4)(n-m) - 2R,\, (5b/4)(n-m) + 2R]$ on every admissible history. Lemmas D′ and D″ run at one fixed K and never compare two trajectories or two values of K, so only their constants must be free of K. The bound $|\beta_{k+1}(h_k) - \beta_{k+1}(0)| \leq C_\beta g_k$ follows from (K1) and the mean-value theorem in the own slot, and it gives the window bracket pointwise on each clamped history of the intermediate-value argument. Verifier: `SOURCES_T2_KUNIF_VERIFY.md`, [VERIFIED REDUCTION]; items 1, 2, 4 confirmed, item 3 corrected: (G-gain) gave C_β only on monotone slot histories, so (K2) via (G-gain) needs item 7′ (now lifted by `SOURCES_GGAIN_NONMONO.md`); the CMP 119 page for the definition of β_j is p. 253, not p. 249. Open inputs: item 7′ for (K2) (from (H0), ledger A5, or from (G-gain), ledger B13); L5, the domain and continuity of β off Bałaban's trajectory (item 8, ledger B7; round 7: proved relative to listed inputs by nested intervals on true trajectories, Theorem 5.15); the inputs of Theorem K for (K3); the finite-volume identification of the step (ledger B8, with A15, narrowed by Proposition 5.8), an input of Theorem 1, not of the shooting.

`SOURCES_T2_V1.md`: item 5, V1, ledger B5. Inside Bałaban's construction with the cut CMP 109 (2.9), the term $E^{(k+1)}$ of (2.13) vanishes identically at $g_k = 0$. So $\Pi^{(k+1)}(0)$ and $\beta_{k+1}(0)$ come from $\log Z^{(k)}$ alone: $\beta_{k+1}(0)$ is the Maxwell coefficient of the one-step Gaussian determinant, it is free of the history, and it equals the determinant of the numerics up to a constant. The file reads the definitions (1.3)–(1.22) and (2.10)–(2.15) on page images and passes from Bałaban's exponential gauge fixing to the bordered Hessian of the numerics (gauge-fixing independence, constraint Jacobian, expansion point, normalization). Verifier: `SOURCES_T2_V1_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL] at fixed level, with 14 claims confirmed and 3 corrected; the conditions are (G2), Bałaban's own inductive bounds, and a declaration that the averaging is (0.4) (the numerics use (0.4); Bałaban's step uses (0.12)). Open inputs: (G2) (item 6); the averaging declaration; Bałaban's inductive bounds.

`SOURCES_T2_G2.md`: item 6, (G2), ledger B6. At fixed level j, the constrained one-loop log-determinant on a torus of side N has no zero modes, and its density per coarse cell converges to the infinite-lattice density with error $\leq C_j e^{-\delta_j N}$, also for every derivative in the plane-wave amplitude. The p^2-coefficient commutes with $N \to \infty$; the f^2-coefficient at the constant field

commutes with $N \to \infty$, given a gap input (Gap_f) and C^4-regularity in f, both [R]. The massless pieces of Prop. 2.1 are handled fiberwise, with an N-independent zero-mode constant. The proof uses a Combes–Thomas decay bound and the method of images; the constants are fixed-j constants ($\delta_j \gtrsim L^{-2j}$), and the j-uniform version is not proved. Verifier: `SOURCES_T2_G2_VERIFY.md`, 21 claims confirmed and 6 corrected, none affecting a conclusion; item 6 moves from [FORMAL] to [R] ([VERIFIED REDUCTION]) at fixed j. Open inputs: the surjectivity of $\tilde{Q}$, (Gap_f), and the reading that the averaging is analytic.

`SOURCES_T2_S123.md`: **(M_j), ledger B4.** At fixed $j \geq 1$, odd $L \geq 3$, and small flux f, the lattice spectral inputs hold: (S1) $\mathcal{K}_t(f) \geq -C_1|f|$ for $t \in [0,1]$, (S3) $\pm V \leq C_3[(\mathcal{K}_t + \mu)^2 + f^2]$ in the subdivided form of `KAPPA_BOUNDED.md` §1.3, (S2) at $t = 0$, and the coarse comparison (S2c) $N_0(\lambda - \eta_1) \leq N_t(\lambda) \leq N_0(\lambda + \eta_1)$ with $\eta_1 = O(f^2)$. Corollary 7.1 then reduces (M_j), the limit of $[\Gamma_c(\Lambda_j; f) - \Gamma_c(1; f) - (f = 0 \text{ value})]/(n_2 f^2)$ to d_j, to the near-zero input (Z_j), itself reduced to the 2D gap (G'_A), and to region 1 of Prop. 4.4 of `BETA_FP_M.md`. The proofs work in the twisted algebra of operators that commute with the magnetic translations, with Peierls products, Moyal expansions with kernel-norm remainders, a Nenciu-type inverse, and operator bounds from the vanishing order of symbols. At $t = 0$ a fibre decomposition over $k_\perp$ separates the operator, and a gauge-copy lemma (Lemma 5.8) shows that the near-zero density of states needs only (G'_A): the transverse charged 2D SU(2) Wilson Hessian at constant flux has no spectrum in $(-c|f|, c|f|)$, and numerically its gap is about $|ef|$. Verifier: `SOURCES_T2_S123_VERIFY.md`, (M_j) [VERIFIED REDUCTION], with (S1), (S3), (S2)$|_{t=0}$, (S2c) and region 2 of Prop. 4.4 [PROVED+VERIFIED AT FIXED j, CONDITIONAL ON (I3)]; 25 claims confirmed, 2 corrected (in Lemma 1.4 ∇^β is one twisted convolution; the explicit L-power of C_j in Prop. 3.2 is convention-dependent and is not quoted), and the input list of (M_j) gains the neutral-sector differentiation under the integral, an inference. Open inputs: (I3), the entrywise analyticity in f of Bałaban's linearized averaging at each of the j steps, is now proved (next paragraph); (G'_A) (open, independent of j and L, numerically confirmed at six rational fluxes with margin about $|ef|$); region 1 of Prop. 4.4 of `BETA_FP_M.md` (argued); all under ledger B4.

`SOURCES_T2_I3.md`: **the Peierls input (I3), ledger B4.** Let L be odd ≥ 3, and let the averaging be Bałaban's (0.4), or (0.12) with Federbush's average. Then the linearized averaging $\tilde{Q}_f$ at the constant-flux background is entrywise analytic in complex f on $|f| < f_0$, $f_0 = \pi/8$ for (0.4) and $\min(\pi/8, 1/8)$ for (0.12), and (I3) holds at every step $i \leq j$ with radius and constants independent of i and j (Theorem 7.1 there). It holds in the stated two-block range and does not use (I1) for the fixed vertices. The proof linearizes the averaging explicitly at the flux background (Prop. 5.3), bounds each charged entry of $\tilde{Q}_f$ and of D_f by its reduced kernel (Props. 5.4, 6.1), and assembles the bordered matrix of one step. Verifier: `SOURCES_T2_I3_VERIFY.md`, 15 claims confirmed, 2 corrected; (I3) moves from [I] to [P], given the identification of the one-step objects of the project with the linearization of (0.4) or (0.12) at the constant-flux background. Open inputs: that identification; for (M_j) the 2D gap (G'_A) and region 1 of Prop. 4.4 of `BETA_FP_M.md` stay. Rounds 6 and 7 settle both (Theorems 5.12 and 5.14).

5.2 Target 2, step 2: the sourced Theorem 1 (ledger B11)

`SOURCES.md`: **the E14 reduction of (B-S).** The plaquette-density source is a local bare coupling, $\rho_0^f = \exp(-\sum_p (g_0^{-2} - f_p) P_p(U))$, so the Schwinger functions on test functions flat on the diagonals reduce to a sourced version of Bałaban's Theorem 1 with a pointwise coupling flow (S-A), sourced regular terms (S-B), a local contact functional $W_k[f]$ (S-C), a local β-function (S-D), and the normalization (S-E). The new estimates are (E14-a) the sourced small-field step, (E14-b) the sourced

large-field step and R operation, and (E14-c) the derivative bounds in f; (E14-d) = E13 is needed only for full-sequence convergence, and (E14-f) only for nontriviality. A dimension count shows that the source of the dimension-4 density is marginal at every degree, that its field-independent part is relevant and is removed as local contact terms, and that the first field-dependent gradient term has dimension 6. One step is written out at the Gaussian level: with $K_f = (\Delta^{(1)})^{-1/2}\Delta^{(f)}(\Delta^{(1)})^{-1/2}$ and $\|K_f\| \le r$ the source enters through $-\frac{1}{2}\log\det(1-g_0^2K_f)$, and the linear response per step is $c_j = 1 + \frac{1}{2}g_{j-1}^3\beta_j'(g_{j-1})$. Verifier: `SOURCES_VERIFY.md`, [VERIFIED REDUCTION] (the structure confirmed; items 5, 6, 8 corrected, item 10 confirmed with corrections); the printed CMP 119 (2.24) is $1/g_{j-1}^2(x) = 1/g_j^2(x)+\beta_j(g_{j-1})\varphi_j(x)$, and by (2.25) Bałaban evaluates β_j at the constant coupling, so (S-A) and (S-D) are new; derivatives of $\log Z$ are cumulants; $\prod_j c_j$ is uniformly bounded, not shown convergent, and assumes a one-step recursion; the uniform bound for general F flat on the diagonals is not derived from the source form. Open inputs: (B-S) as a whole, (E14-a, b, c) (ledger B11, [OPEN]), with its parts B12–B20 and B22; the (E14-c) clause of the bridge (B-N1) (ledger B21); (NT) = (E14-f) for Theorem 4.3; E13 = (E14-d), not needed for Theorem B.

`SOURCES_D1.md`: one sourced step at one loop (D1). At one loop the plaquette-density source, a local bare coupling $w_p = 1/g_0^2 - f_p$, gives the multiplicative source renormalization $c_1 = 1$ exactly, for every background; the first correction is $c_1 - 1 = \frac{1}{2}g^3\beta_1'(g)$, which is $\beta_1^{(2)}g_0^4 + O(g_0^6)$ if β_1 is even and $O(g_0^3)$ with a C^1 bound only. The field-independent terms are the tadpole $\frac{1}{2}(9L^4-9)g_0^2$ per coarse cell ($\frac{3}{4}(1-L^{-4})g_0^2$ per plaquette) and a contact term $C(q)$ that is analytic at $q = 0^+$, and the first field-dependent gradient terms have dimension 6. The source enters the step only through the Hessian $\Delta_f = \sum_p w_pH_p$ and a current term $\langle J_f, B\rangle$ at Bałaban's unweighted minimizer, while gauge fixing, ghosts, and the constraint carry no f; for constant f the quadratic form scales by $1-g_0^2f_0$, so the determinant changes by a field-independent constant over $9L^4 - 9$ modes per cell. Per-momentum formulas for the tadpole, $C(q)$, and the mixed derivative are evaluated by `numerics/d1_check.py` and checked against real-space finite differences on the periodic 6^4 lattice. Verifier: `SOURCES_D1_VERIFY.md`, [PROVED+VERIFIED] for the algebra (items 1–4, 6, 8–10 confirmed) and [NUMERICAL, REPRODUCED] for the rerun, which matched every digit; corrected: the dimension-6 count holds by index parity, the γ_2 values are unreliable, the data support $|c_1-1| \le 5\times10^{-4}$ but not geometric convergence in the grid, and the expansion-point error has the small parameter Lp. Open inputs: none at the Gaussian level; the non-Gaussian sourced step and the pointwise flow belong to D2–D3 and to (B-S), ledger B11.

`SOURCES_D2.md`: sourced small-field step (E14-a). The sourced small-field step is Bałaban's step with extra source vertices of order g_k, g_k^2, g_k^3, all in his $P^{(k)}$ class (each $\le O(g_k)(1+|B|^2)$ on $|B| < \varepsilon_1/g_k$), a propagator for the sourced Hessian with Bałaban's decay and regularity bounds (constant $3c_1$ with $c_1 = O(\delta_0^{-5})c_0$, rate $\delta_0/2$), and a field-independent vacuum term W that factors out. Its step (c), the pointwise flow $1/g_{j+1}^2(x) = 1/g_j^2(x) - B_{j+1}(f(x))$ with $h \equiv 0$, is not established here and is redone in `SOURCES_D2C.md`. The source term is evaluated at Bałaban's unweighted minimizer, so his measure and propagators are used without change, and the sourced propagator follows from a weighted Neumann series in his cube-to-cube operator norms (CMP 116 (1.7)). Real-source derivative bounds are tied to polystrip holomorphy by the interpolation Lemma 7.1, and the remainder bounds are reduced to Bałaban's estimates with a freezing property and a constant-source family. Verifier: `SOURCES_D2_VERIFY.md`, steps (a), (b), (d), (f) [PROVED+VERIFIED], step (e) a [VERIFIED REDUCTION], and step (c) rejected, since its claim that CMP 109 Sects. 3–4 run verbatim is [REFUTED] ((4.34)–(4.37) use Euclidean symmetry, and the cubes are at scale k, not j); corrected: (a) needs a block-constant $\bar f$, the constant is $3c_1$ and not $\times3$, W factors only in

the small-field region, and the polydisc radius is $r/(2n)$. Open inputs: analyticity of the step in its input activities, (W4), and the field-dependent domains Λ_j in W (ledger B20); the recovery $\kappa \to \kappa/2$ (ledger B22, a verified reduction that uses A23); (Hβ), reduced in `SOURCES_HBETA.md`; the scale condition (U) is the constant choice C23, not an open input.

`SOURCES_D2C.md`: **step (c) redone, frozen source per domain (Lemma A).** Each step term $E^{(j)}(X, \cdot\,; f)$, $X \in D_j$, is frozen at $c_X = f(b_X)$, and the freezing error obeys $|D_X| \le E_0 C M^3 (\rho_1/r)(L^j\eta)^5 \lambda_k e^{-\kappa d/2}$ (Lemma A), irrelevant with exponent 1 and with no growth in k. With the identification (I) of the constant-source increment $B_j(c)$ with Bałaban's β_j on the trajectory from $1/g_0^2 - c$, this gives the pointwise recursion $1/g_{j+1}^2(x) = 1/g_j^2(x) - B_{j+1}(f(x))$ with $h \equiv 0$ and gradient corrections irrelevant with exponent $\min(\alpha, \beta, 1)$. The non-constant source is handled at scale j, per localization domain, as a gauge-invariant difference bounded in an axial gauge on X, where each field costs two powers of $L^j\eta$. Bałaban's CMP 109 Sects. 4–5 are then used only for constant sources, where they apply at bare coupling $1/g_0^2 - c$. Verifier: `SOURCES_D2C_VERIFY.md`, Lemma A and the recursion [PROVED+VERIFIED, CONDITIONAL ON (D) AND (Hβ)]; corrected: the freezing points are the points of X nearest to b_X, (I) is exact only with the adapted gauge-fixing weight $1/u_k(c)$ (otherwise up to $O(e^{-c\varepsilon_1^2/g_k^2})$), $\beta > 0$ in (4.18) must be fixed, and the exact cancellation of the cubic first-moment term of (4.34) is unsupported and not needed. Open inputs: the domain hypothesis (D) (ledger B20; argued in `SOURCES_D2C_VERIFY.md` §2 from the CMP 109 p. 263 remark, which cites CMP 102 Prop. 9, ledger A4); (Hβ) (Lemma H of `SOURCES_HBETA.md`); (I) with large-field thresholds (ledger B12); the adapted gauge-fixing weight changes the construction (ledger C10).

`SOURCES_D3.md`: **sourced large-field step (E14-b).** The sourced large-field step is Bałaban's step with all thresholds frozen at the reference couplings, the local coupling $w_j = w_{j-1} - B_j(f)\varphi_j$, an R operation that resets it on Z to $u_k(f(x))$ (CMP 122 II p. 365), and an adapted local gauge-fixing weight. Real sources keep the positivity of the large-field weights under $3\gamma^2 r \le \frac{1}{2}$ (revised to $5\gamma^2 r \le \frac{1}{2}$ in `SOURCES_D3R.md`), Lemma D3.1 gives the local holomorphy radius $c_\tau r/(E_0|Y|)$ of one positive denominator, and the identification (I) is off by a threshold term, the new input (H-thr). The file lists the places (P1)–(P7) where Bałaban uses a constant coupling in the large-field step and states the change at each. Each change is reduced to a quoted Bałaban estimate (CMP 119 (2.23)–(2.31), CMP 122 II (1.73)–(1.75)); a line-by-line rerun of CMP 122 I–II is not done. Verifier: `SOURCES_D3_VERIFY.md`, [VERIFIED REDUCTION] with corrections: the reset term is bounded by Bałaban's sup bound (CMP 122 II p. 366), not cancelled; the t-parameter term (P8) of CMP 119 p. 271 is missing; old R-terms are merged into E_k, so freezing through R is open again; the Gevrey order is $(n!)^{2+\epsilon}$, not $(n!)^2$; and the lower bound on w_k holds relative to Bałaban's local coupling. Open inputs: (H-thr) (ledger B15); the real uniform (H-hist) (ledger B13); strip holomorphy of u_k, now N3 (ledger B18); (W4) and the field-dependent domains Λ_j (ledger B20); (FX), (FX2) for the final step (ledger B19); (U) is the constant choice C23.

`SOURCES_D3R.md`: **revision of D3 ((P3), (P8), freezing through R, Gevrey order).** The sourced reset coefficient differs from Bałaban's by at most $\delta_r = 4r + 2C_h M \rho_1 \lambda_k$, additively and pointwise (Lemma P3), so his sup bound of CMP 122 II p. 366 holds with one changed constant, and the large-field factors survive with $\gamma_0 \to \gamma_0(1-\theta) \ge \gamma_0/2$, $\theta = \gamma^2\delta_r \le 5\gamma^2 r \le \frac{1}{2}$. The t-parameter term splits as $d_B + d_S$ with $|d_S| \le 2r$, freezing through R splits into a pure part (Lemma F1) and an R-induced part (Lemma F2), and the cumulant bound has Gevrey order $(n!)^{2+\epsilon}$, admissible with $D = 3$. Positivity is kept by a split against Bałaban's local coupling, and the (P8) term is split into

a piece with the support of Bałaban's deviation, booked as a boundary term, and the source weight of D2. The regular output is split by an auxiliary parameter into the pure and R-induced parts, and the Gevrey count uses a Lah-number sum over partitions. Verifier: `SOURCES_D3R_VERIFY.md`, [VERIFIED REDUCTION] with corrections: (P3) needs $\rho_1 \le r/(4C_hM)$; in (P8) h_k is booked as irrelevant, and d_S changes the covariance $C^{(k)}$ at every t (relative size $O(\gamma^2 r)$); Lemma F2 is a new claim, not Bałaban's bound for merged R-terms (item W8, taken up in `SOURCES_D3W8.md`); the Gevrey count is confirmed conditional on (B-N1). Open inputs: strip holomorphy of u_k (N3, ledger B18); (H-thr) (ledger B15); the real uniform (H-hist) (ledger B13); (W4) and the covariance change in (P8) (ledger B20, the latter by analogy with A16); (B-N1) (ledger B21).

`SOURCES_D3W8.md`**: covariance change along t and merged R-outputs (Lemma F2′).** Along the t-interpolation the measure is Bałaban's CMP 119 (3.27) with covariance $C_f^{(k)}$, independent of s and t; with $\tau \le 3c_1c_H\gamma^2 r \le \frac{1}{2}$ one has $\|C_f - C_0\|_\delta \le 2c_1\tau$, and the large-field factors survive with $\gamma_0 \to \gamma_0/4$. Lemma F2′: the R-induced part obeys $a_m \le 32\varepsilon' g_m^{\kappa_0}$ with no compounding, and $|E_R^{(m)}(X,U) - E_R^{(m)}(X,1)| \le CM^2 \cdot 32\varepsilon'(L^m\eta)^4 g_m^{\kappa_0} e^{-\kappa' d_m(X)}$ on regular configurations, a marginal term of Bałaban's R-type form, summable in m for $\kappa_0 > 2$. Transported inputs keep their scale label and never pass through the generated map G, so a Schwarz-lemma step on $G - G(0)$ with Bałaban's factor $\varepsilon' = O(\alpha_2^{-1}\varepsilon_1)$ (CMP 109 p. 280) gives $a_{k+1} \le 8\varepsilon' b_k$ per generation. The covariance bounds come from the weighted Neumann series of `SOURCES_D2.md`, and positivity from domination by a rescaled C_0 Gaussian. Verifier: `SOURCES_D3W8_VERIFY.md`, route (a) and Lemma F2′ [PROVED+VERIFIED, CONDITIONAL ON (W4), (EPS), LEMMA F1, (D), AND $32\varepsilon' \le 1$], the t-measure and its bounds corrected: the cross term $g_k^2\langle A, \mathcal{W}_S A_k\rangle$ goes to $B^{(k+1)}$, CMP 116 p. 3 covers C_f only by analogy, the domination must use the marginal region (or Anderson's inequality) to avoid a volume factor, and ε' contains Bałaban's large constant of p. 280. Open inputs: (W4) and (D) (ledger B20; (W4) is reduced to the printed (C1)–(C5) through Theorem SM); (Eps), from Theorem SM and A10; the extension A16 to the fluctuation covariance; the conditions $32\varepsilon' \le 1$, $S \ge 1$, $\kappa_0 > 2$ change the construction (ledger C16); the monotone-history assumption of Lemma F2′ is removed by the running maximum, with side condition $4\gamma^{\kappa_0} \le E_0$ (`SOURCES_GGAIN_NONMONO_VERIFY.md`).

`SOURCES_D4D5.md`**: RP and symmetry transfer, and the cumulant bound ($\mathbf{S}_{\mathrm{src}}$).** The sourced RG output equals the bare integral $Z_\varepsilon[f]$ (Lemma D4.1), reflection positivity and the lattice symmetries are properties of the lattice measure (Lemmas D4.2–D4.4), and given (FX) the cumulants of real sources obey ($\mathrm{S}_{\mathrm{src}}$): $|\kappa_n(f_1,\dots,f_n)| \le (C_S\sigma^{-a})^n (n!)^{1+\eta} \prod_i N(f_i)$ with $N(f) = \max(\|f\|_\infty/r, \|\nabla f\|_\infty/\rho_1)$, for $\sigma \in [16\varepsilon, \ell/8]$, supports pairwise at distance $\ge \sigma$, minimal diameter $\le \sigma$, and C_S independent of K. Then ($\mathrm{S}_{\mathrm{src}}$) $\Rightarrow$ (H-U) with $s = 6$, $D > 1$. Each RG operation is an identity of integrals (Haar invariance, δ-function blocking, partitions of unity, convergent expansions, $\int dV\, R\rho = \int dV\, \rho$), and complex sources follow by the identity theorem. Lemma C takes one Cauchy derivative in each of n separated source amplitudes on a polydisc whose radius does not depend on n, so the sum over polymer sizes gives a single $n!$. Verifier: `SOURCES_D4D5_VERIFY.md`, D4.1, D4.3, D4.4 and Lemma C [PROVED+VERIFIED] (Lemma C given (FX)), ($\mathrm{S}_{\mathrm{src}}$) $\Rightarrow$ (H-U) confirmed, and (FX) confirmed as a new hypothesis not contained in CMP 116–122; corrected: cite CMP 122 I (1.102) for the actual R′ operation, the block geometry is invariant only under translations by $M\mathbb{Z}^4$ or coarser, and the order $1+\eta$ needs the coarsest-scale (FX2) radius (else an extra factor $(n!)^{2p_R/p_0}$, still OS-admissible). Open inputs: (FX) in Bałaban's main form with the coarsest-scale (FX2) radius (ledger B19; remaining input N3, ledger B18); S1 at all scales, part of (B-S) (ledger B11); the (E14-c) clause (ledger B21).

`SOURCES_HBETA.md`: **Lemma H, the two-loop bound (Hβ).** Lemma H: given (I), (E), and (H-hist), with $x := C_1 A\gamma/\beta \le \ln(3/2)$ and $3r < g^{-2} - \gamma^{-2}$ ($g = g_K$), every real c with $|c| \le 2r$ keeps all $g_k(c) \in \,]0,\gamma]$ and gives $\sum_{j\le K} |B'_j(c)| \le e^x - 1 \le \frac{1}{2}$ and $\frac{1}{2} \le |v_k(c)| \le \frac{3}{2}$, uniformly in K. With $\beta = \beta_0/2$ the smallness condition reads $\gamma \le \ln(3/2)\,\beta_0/(2C_1A) \approx 0.20\,\beta_0/(C_1A)$. In $u = 1/g^2$ the chain rule gives $\partial\beta_j/\partial u_i = -\frac{1}{2}g_i^3\,\partial\beta_j/\partial g_i$, so a C^1 bound costs $O(g_i^3)$ per step, $\sum_i g_i^3 \le 2\gamma/\beta$ along the trajectory, and a Gronwall product on the variational equation gives $e^x - 1$. A bootstrap on the component of $\{|c| \le 2r : u_K(c) > \gamma^{-2}\}$ that contains 0 keeps every c-trajectory in $g \le \gamma$; the Gronwall algebra is formalized in Lean (`LemmaH`). Verifier: `SOURCES_HBETA_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON (I), (H-HIST)]; corrected: the hypothesis list adds that β_j is the infinite-volume function independent of K, (H-hist) can be weakened to a column-sum bound, the smooth-clamp remark is dropped, and the toy numbers are not reproducible (the bound holds and is nearly sharp). Open inputs: (I) (ledger B12; it holds by construction with the adapted gauge-fixing weight C10 and the frozen thresholds C1, but is not written as a lemma); the real uniform (H-hist) (ledger B13, a verified reduction with open inputs (Q)/c_-, (M_r), CMP 102 (189), and the (M-B) bookkeeping); (E), the uniform positivity of the one-loop $\beta_j(0)$; complex c is not covered and is the strip bound N3 (ledger B18).

`SOURCES_T2_KAPPA2.md`: **ledger B22.** The decay loss $\kappa \to \kappa/2$ of the sourced step occurs at every step in three places (the interpolated inputs of the freezing proof in `SOURCES_D2.md` §6.3, the extraction remainders of `SOURCES_D2C.md` §3, the R-induced terms of Lemma F2 in `SOURCES_D3R.md`), and it is recovered by $L/2$-rescaling without compounding. The conditions are (K1′) $\delta \le \frac{1}{13}(1 - 2(1+4\beta)/(\theta L))$, which is $\delta \le 5/169 \approx 0.0296$ at $L = 13$, $\beta = \frac{1}{4}$, $\theta = \frac{1}{2}$ (round 7, `SOURCES_A23_TSTEP.md`; the round-6 form (K1) had $\frac{1}{10}$ and $\delta \le 0.0385$, from the small-field T step only), with the choice $\delta = 0.025$ (0.03 fails; Lean `K1_numeric_13`), and (K2) $\theta(1-\beta)L/2 \ge 1 + 3\beta$; at the last scale the final expansion runs at $\kappa_{\text{eff}} = \theta\kappa/(1+4\beta) = \kappa/4$, with the M-condition (K3) $M \ge 12L(4.5\kappa/C_1)^{1/4}$, implied by the existing one of `SOURCES_FX_R2.md`. Bałaban proves only $2d_k(Z) \ge Ld_{k+1}(Z')$ (CMP 116 (2.36)), so the gain per step is $L/2 = 6.5$, and his printed choice $(1 - 10\delta)\frac{1}{2}L = 1$ in (2.41) spends all of it; the recovery therefore takes δ smaller, at the cost of a larger κ (a lower bound about 4.3–4.7 times the unsourced one, like for like; was 3.6 under (K1)), which moves only constants fixed after κ. The R operation drops $4\beta\kappa$ of rate (CMP 122 II (1.66)–(1.100)), the rescaling pays these drops and the T-step fraction, now at most 13δ (10δ printed for the small-field step plus at most 3δ unprinted for the large-field step, ledger A23), and the (1.93) volume charge is paid by p_0 through the condition before (1.95). Verifier: `SOURCES_T2_KAPPA2_VERIFY.md`, [VERIFIED REDUCTION]; items 2–4 confirmed on page images, item 1 corrected: the three losses stack within one step, so a literal half per absorption would make (K1) fail at $L = 13$, and the repair reads $\theta := 1 - m_{\text{tot}}/\kappa$ as the kept fraction per cycle (each loss is a polynomial absorption with additive cost; $\theta \ge \frac{1}{2}$ for $\kappa \ge 2m_{\text{tot}}$), with which (K1)–(K3) stand and the new lower bound is $\kappa \ge 1536$ at $d = 4$, $\beta = \frac{1}{4}$. Open inputs: the unprinted large-field T-step rate loss of CMP 119 Sect. 3 (ledger A23), on which the homogeneity of the large-field T step in the input rate rests; CMP 116 (2.36), which Bałaban asserts without proof.

`SOURCES_UN1.md`: **(U), ledger C23.** The scale condition (U) does not conflict with the bridge (B-N1): with $\lambda_K = 1$ (physical units), the r in (U) is the Cauchy margin r_0 of the constant-source family, not a test-function amplitude, so (U) is the constant choice $\rho_1 \le c_* r_0/M^a$, made once with $\rho_1 := \min(c_* r_0/M^a,\, r_0/(4C_h M))$, after r_0 and before K. The bridge evaluates the generating functional only on sources rescaled into the domain of (FX2), at the price of a factor $(4/\rho_1)^n$ per n-point bound, with s and D unchanged. The bridge uses n-linearity, $\kappa_n(f_1,\dots,f_n) = \prod_i \lambda_i\, \kappa_n(f_1/\lambda_1,\dots,f_n/\lambda_n)$,

with $\lambda_i = 2N_U(f_i)$ and $N_U(f) = \max(\|f\|_\infty/r, \|\nabla f\|_\infty/\rho_1)$, and pairwise σ-separated supports make the sup and the gradient of $\sum_i z_i f_i$ a maximum over single terms, so the whole Cauchy polydisc stays in the admissible domain. Small widths $h \geq \varepsilon$ are paid by the norm growth $1/h$ per bump, which the flatness of F absorbs, while (U) costs only an h-independent factor; the scale-matched reading of `SOURCES_D2.md` l. 217 would create a real conflict and is dropped. Verifier: `SOURCES_UN1_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON THE (Hβ) GRADIENT BOUND], so (U) is a constant choice (ledger C23), verified compatible with (B-N1); corrections: the (FX2) gradient radius is $\rho_1/2$, hence $\lambda_i = 2N_U(f_i)$, the sup radius of the activity polydisc is $\tau_m(\Gamma) \leq r$, overlapping supports never reach (S), and the (Hβ) bound $|\nabla f_k| \leq \frac{3}{2}\rho_1$ is a verified reduction for real c (Lemma H and the chain rule), not open. Open inputs: (Hβ) for complex c (open); for real c its inputs (I) (ledger B12) and (H-hist) (ledger B13); the $\kappa \to \kappa/2$ recovery (ledger B22, a verified reduction that uses A23).

5.3 Step 3: the (S) clauses and (FX) (ledgers B21, B19)

`SOURCES_B21_S.md`: uniformity of (S), ledger B21. The source bound (S), $|\kappa_n(f_1, \dots, f_n)| \leq (C_S\sigma^{-a})^n (n!)^{1+\eta} \prod_i N_U(f_i)$ for real sources with pairwise separation $\sigma \geq 16\varepsilon$ and $\min_i \operatorname{diam} \operatorname{supp} f_i \leq \sigma$, holds uniformly in the bump width $h \in [\varepsilon, \ell/512]$ and in the position of the bumps, with the C^1 norm $N_U(f) = \max(\|f\|_\infty/r, \|\nabla f\|_\infty/\rho_1)$. Every factor of $C^n(n!)^{1+\eta}$ is independent of h. The file traces one rescaled bump through Lemma C of `SOURCES_D4D5.md` and finds one h-dependent term, the cube count $Q_m \leq (2 + D/(M\lambda_m))^4$ at scales finer than the bump. With $D \leq \sigma$ this count is at most $(2+2D/\sigma)^4 u^4$, $u = \sigma/(2M\lambda_m)$, and the tree decay $e^{-3\kappa u/8}$ of the polymers that meet all supports cancels it, while the gradient cost is paid by N_U and the flatness of the test function. Verifier: `SOURCES_B21_S_VERIFY.md`, [VERIFIED REDUCTION], corrected in scope (the inputs must also list the residue of B19 and D2 §4.3) and in one constant (in the bridge $D/\sigma = 3/8$, not $1/16$, so the factor is $(2.75)^4 \approx 57$). Open inputs: S1 at all scales including $m = 0$; (FX) in Bałaban's main form (1.98) (ledger B19 with the choice C13), whose remaining input is N3 (B18); the coarsest-scale (FX2) radius, without which the order is $2 + \eta$ and $D > 2$; D2 §4.3, itself only reduced; the (E14-c) clause of B21, which no file maps to Lemma C.

`SOURCES_FX.md`: hypothesis (FX), ledger B19. (FX), the convergence of the final Mayer expansion of the sum over large-field geometries plus the unit-scale integral, with polymer radius $c \cdot r$ in the source (FX2), uniform in K and ε, is reduced to Bałaban's large-field bounds CMP 122 II (1.79)–(1.89) and (1.97), the convergence criterion of CMP 116 (2.38)–(2.41) and Cammarota (ledger X4), and named local checks. The construction is a final R-step with $Z'' = \emptyset$ followed by the unit-scale cluster expansion of CMP 116 §2, as Dimock does for φ^4_3 (Dimock III, Theorem 2). The file reads CMP 122 I (0.2)–(0.6) and CMP 122 II (1.71)–(1.104) and finds that Bałaban already exponentiates the sum over renormalizable large-field components inside each R step, so that only the components that pay for later steps stay as a sum, and at the last scale there are no later steps. The fine-history sum inside a component is bounded by (1.79), because Dimock's power-decay argument ((209)–(211)) does not transfer to $d = 4$, and the source enters the large-field operations only through the upper bound (1.73). Verifier: `SOURCES_FX_VERIFY.md`, [VERIFIED REDUCTION], corrected: the per-step expansion sums only components of diameter $\leq 100MR_k$ counted by position ((1.96)), so the last step, which sums components of unbounded size, needs the tree decay O1; three page quotes are fixed ((1.80) has R_n^{d+1} and a local K, (1.85) has $A_1^2 p_0^2$, the condition after (1.94) ends in $-\frac{3}{2}p_0(g_k)$); the fallback order is at most $1 + \eta + 2/3$. Open inputs: the verifier left O1(b), O2, and O3 open; O1(b) and O2 are now [PROVED+VERIFIED] (`SOURCES_FX_O1.md`, `SOURCES_FX_O23.md`), R2 is [PROVED+VERIFIED] as a reduction, and O4 is avoided by the main

form (C13), so the remaining input of B19 is N3 (ledger B18), through O3.

`SOURCES_FX_O1.md`**: gap O1(b) of (FX), Lemma [P].** Lemma [P]: for a connected union Y of MR_j-cubes, $d'_{j+1}(S(Y)) \geq c_* d'_j(Y)$ with $c_* = [3 \cdot 4^d \sqrt{d}(\rho+2)^d]^{-1}$, $\rho = LR_{j+1}/R_j$, and no additive constant. With the $n = K+1$ term of CMP 122 II (1.80) this gives $\kappa_K(Y) \geq \kappa_{\text{fin}} d_K(Y)$ for every non-renormalizable final component Y, which is Bałaban's assertion of the term $-\kappa_1 d_k(X)$ on p. 387, with $\kappa_1 = \kappa_{\text{fin}}$. The proof combines three elementary facts (a tree of length ℓ meets at most $4^d(1+2\ell)$ closed unit cubes, $d'_j(Y) \leq \sqrt{d}(n_f - 1)$, and a coarse cube meets at most $(\rho+2)^d$ fine cubes) with the count of at least 21^d coarse cubes in $S(Y)$. The halo $S(Y) \setminus Y$ carries no large fields (p. 384), so nothing is absorbed by p_0, and only the first term of (1.80) is used, all its terms being non-negative. Verifier: `SOURCES_FX_O1_VERIFY.md`, [PROVED+VERIFIED], with one remark corrected (the counterexample to the form "factor $1/L$ plus a constant" needs $d \geq 2$, a constant independent of L, and L large) and R3 discharged by Bałaban's own comparison in (1.86)→(1.87). Open inputs: R2 ($\kappa_{\text{fin}} \geq \kappa_{\text{final}}$), the only point left by the verifier, is closed in `SOURCES_FX_R2.md` (choice C14); R1 (grid alignment) changes constants only.

`SOURCES_FX_O23.md`**: gaps O2 and O3 of (FX).** O2: the R-denominator of CMP 122 I (1.100) is one small-field integral per renormalizable component with the bare Wilson action $-g_k^{-2} A(\zeta_i, \cdot)$ and no E-, R-, or B-terms, so it is holomorphic and non-vanishing on a source polydisc of radius $c \cdot r$, with c independent of $|\Lambda_i|$, k, and ε. O3: inside the large-field operations T' the sourced weight obeys $\text{Re}(w + \delta w) \geq (1 - \theta_W)(g''_k)^{-2}$, so the upper bounds (1.73), (1.79), (1.89), (1.92), (1.97), (1.100) hold with $\gamma_0 \to \gamma_0(1-\theta_W)(1-\tau_C) \geq \gamma_0/4$ and $p_0 \geq 16/(\gamma_0 A_1^2)$. For O2 the Wilson part of the source perturbation is $|\sigma_W| \leq \tau\Phi(g_k)$ with $\Phi(g) = C(M) g^2 p_0(g)^2 R(g)^4 \to 0$, and (1.102) also allows an f-independent denominator. For O3 the plaquette terms $P_p \geq 0$ of the real G-valued configuration inside T' (CMP 122 II p. 379) give domination in modulus by a positive operation with a smaller coupling prefactor, while the thresholds stay fixed. Verifier: `SOURCES_FX_O23_VERIFY.md`, [PROVED+VERIFIED] for O2 and [PROVED+VERIFIED, CONDITIONAL ON N3] for O3, since the strip bound $|\delta w| \leq \frac{3}{2}\tau$ is N3; it adds the cleaner condition $p_0 \geq 8/(\gamma_0 A_1^2)$ with $\gamma'_0 \geq \frac{3}{4}\gamma_0$, and corrects O4 (the division by $T'_k(Y_i)1$ in (1.103) has no positivity proof, but (1.103)–(1.104) is an optional final remark, so O4 is avoided by keeping the main form (1.98)). Open inputs: N3 (ledger B18); the main form is the choice C13; O4 binds only if (1.104) were used, with fallback order $2 + \eta$.

`SOURCES_FX_R2.md`**: gap R2 of (FX),** $\kappa_{\text{fin}} \geq \kappa_{\text{final}}$**.** R2 holds as a reduction in volume form: $\kappa_K(Y) \geq \kappa_{\text{vol}} \# M(Y)$ with $\kappa_{\text{vol}} = \sqrt{d}\,\kappa_{\text{fin}} = C_1 M^d R_{K+1}/(3 \cdot 4^d (L+2r)^d)$, $r = R_K/R_{K+1}$, and $\kappa_{\text{vol}} \geq \kappa_{\text{final}} := 2\kappa + \sqrt{d}(1+2\beta)\kappa + c_d$ once $M \geq 12L(3\kappa_{\text{final}}/C_1)^{1/d}$, uniformly in K, R, and g. Here $C_1 := \max(1, C_c)$ is the enlarged constant of (1.80). The file quotes Bałaban's order of constants (L first, $\kappa \geq \kappa_0$ before $M \geq M(\kappa)$, p_0 and γ last), so $\kappa_{\text{final}} = \kappa_{\text{final}}(d, \kappa, \beta)$ is fixed before M (route (a)), after C_1 is enlarged (route (b)). It replaces the tree rate of `SOURCES_FX_O1.md` by the volume rate, because the final expansion charges large-field domains per M-cube and the tree rate alone loses R_K^{d-1}. Verifier: `SOURCES_FX_R2_VERIFY.md`, [PROVED+VERIFIED] as a reduction, corrected in its reasoning: the 2κ is the (1.93) decay-recovery charge $(1 + \beta + \beta/(3 \cdot 2^d))\kappa$ per M-cube, not a Mayer-link cost, and the term $\sqrt{d}(1+2\beta)\kappa$ is surplus. Open inputs: none; the two minor inputs, $R_{K+1} \geq 1$ and C_c independent of g, are verified readings (former ledger A20, `SOURCES_BOOKKEEPING.md`), and κ_{final} is the choice C14, not sharp.

5.4 Steps 4–5: N3, (H-hol), and the large-field inputs (ledgers B18, B16, B17)

`SOURCES_HBETA_N3.md`: **N3, the strip bound, ledger B18.** With the source additive in $u = g^{-2}$ ($w_0 = g_0^{-2} - f$, τ in absolute u-units), N3 is equivalent to Lemma H of `SOURCES_HBETA.md` for complex c on $Q = \{|\operatorname{Re} c| \le 2r,\ |\operatorname{Im} c| \le \tau\}$, pointwise in x. Lemma N3 gives $|\delta w_k| \le \frac{3}{2}\tau$ for all $k \le K$, uniformly in K and ε, given (I), (E), real (H-hist), (H-hol) on relative polydiscs, and (B-hol), with (H2-hist) added in the history form; the constant $\frac{3}{2}$ can be any $1 + \eta$. Step 0 bounds $\partial\beta_j/\partial u$ on the thin strip by Bałaban's real C^1 bound at $\operatorname{Re} u$ plus a Cauchy correction of order $g^4|\operatorname{Im} u|$, and a complex Gronwall step with $\sum_i (\operatorname{Re} u_i)^{-3/2} \le 4\gamma/\beta$ closes a bootstrap in $\operatorname{Im} c$. The file also shows that Cauchy estimates on relative discs alone give a factor $1 + Cg^2$ per step and a bound that grows with K, so the real derivative bound is needed. Verifier: `SOURCES_HBETA_N3_VERIFY.md`, [VERIFIED REDUCTION], corrected: condition (a), $3r + \frac{3}{2}\tau < g^{-2} - \gamma^{-2}$, limits r by the gap between $g = g_K$ and γ and does not follow from γ small (settled in `SOURCES_N3A.md`); $d\beta/du = O(g^3)$, or $O(g^4)$ for even β; (H2-hist) is sufficient, not necessary, the minimal input being (H-hist$_C$) on the complex polydisc. Open inputs: (I) (ledger B12), (E), real (H-hist) (B13), (H-hol) in its small-field (B16) and large-field (B17) halves, (H2-hist) or its Route 2(a) (B14), (B-hol), the $h \equiv 0$ booking (C11), and the additive normalization (C12); Step 3 in the D″ form is proved, unverified (B10).

`SOURCES_N3A.md`: **condition (a) of Lemma N3.** With the order of constants $L \to$ Bałaban's constants $\to \gamma \to g \to (r_0, \tau_0) \to K$ and $g = g_K \le \min(\gamma/2, g_1)$, condition (a) holds at the uniform radius $r_0 = \tau_0 = \min(1/8, r_0^{\mathrm{D2}}, r_0^{\mathrm{D3}}, r_0^{\mathrm{FX}}, \gamma^{-2}/10, 1/(6\gamma^2 c_1 c_H))$, since $\frac{9}{2} r_0 \le \frac{9}{16} < 3 \le g^{-2} - \gamma^{-2}$ for $\gamma \le 1$, a margin of 5.33. The radius does not depend on K, and in u-units it is the physical sup-norm radius of the source. The file checks that $g \le \gamma/2$ is admissible for every K (`LEMMA_DPRIME.md` Step 8 in the Lemma E form, `KAPPA_BOUNDED.md` §5.3(b) in the D″ form) and that nothing forces g_K near γ. It refutes the proposed radius $r = \tau = \gamma^{-2}/2$, which satisfies (a) but breaks conditions (b)–(d) of Lemma N3 and the radius bounds of D3R and D3W8. Verifier: `SOURCES_N3A_VERIFY.md`, [PROVED+VERIFIED], with item 1 corrected: "u_k decreasing, g_K largest" holds only in the Lemma E form, while in the D″ form only $u_k \ge u_K - D'$ holds and g_0 can exceed g_K; the D″ sketch has two unnamed gaps. Open inputs: none for condition (a); the order of constants is the choice C5; the inputs of N3 are unchanged, and N3 in the D″ form was a sketch (ledger B10); round 7 closes both gaps (Section 5.11).

`SOURCES_HHOL.md`: **(H-hol), small field, ledger B16.** For the small-field chain, β_j extends holomorphically to the polydisc $|g_i - x_i| < \rho_0 x_i$ with $|\beta_j| \le M$ uniformly in j, k, and volume, where $\rho_0 = \min(0.1, (LM)^{-2}(8n)^{-1/2})$, divided by $\sqrt{2}$ when the X-modes are counted (ledger C6). This holds given Bałaban's conditions with modified constants: (C1′)–(C4′) with $\varepsilon_1 \to 1.235c\varepsilon_1$, (C4″) $8n\rho_0^2(LM)^4 \le 1$ (with $16n\rho_0^2$ once the X-modes are counted), and (2.24′); the radius $\rho \le 1/3$ first asked for is not reached, and Lemma N3 accepts any fixed ρ_0. In reference units $\tilde{B} = xB$ the complex coupling enters only through $w = (x/g)^2$, as the scale of the Gaussian and as the coefficient of the u-prefactored vertices, while the cut $|\tilde{B}| < \varepsilon_1$ does not depend on g. The marginal-domination lemma of the file (its Lemma D) bounds the complex Gaussian by the real one at $\operatorname{Re} w$ at a cost $(|w|/\operatorname{Re} w)^{1/2} \le e^{2\rho^2}$ per real mode, so the volume-uniform bound is made per polymer inside CMP 116 (2.14)–(2.17). Verifier: `SOURCES_HHOL_VERIFY.md`, [VERIFIED REDUCTION], corrected: the per-polymer cost misses the X-modes ($\rho_0 \to \rho_0/\sqrt{2}$), $|\beta_j| \le M$ is a reduction because the majorant covers the Gaussian only, and (C1)–(C5) are not Bałaban's printed labels; single-valuedness is confirmed with no extra assumption. Open inputs: the modified conditions (C1′)–(C4′), (C4″), (2.24′) (the X-mode question is resolved in the row M-C of the dependency table); (F1), reduced

to Theorem SM at complex couplings, from which (H2-hist) is derived with $a_m = \lambda_C^{m/2}$; a γ-independent C_1 in real (H-hist) (B13); the large-field half (ledger B17, `SOURCES_HHOL_LF.md`).

`SOURCES_HHOL_LF.md`**: (H-hol), large field, ledger B17.** The large-field half of (H-hol) holds on $P_{\rho_0}(x^*)$ around Bałaban's real trajectory x^*, with the R-ratio density, the thresholds, R_k, and N frozen at x^* (ledger C1): the large-field integrals are entire in $u = g^{-2}$ in finite volume and are dominated in modulus with $\gamma_0 \to \gamma_0'' \geq 0.82\gamma_0$ at $\rho_0 = 0.1$, and F2′ holds on the polydisc with $32\varepsilon' \to 40\varepsilon'$. This is a reduction to (M-A), (M-B), (M-C), and (M-D). The file shows that the radius $(100MR_k)^{-4}$ of W4 is a radius in the input activities, which the real (1.100) denominator does not contain, and it removes the coupling from the denominator by freezing the ratio at x^*, at a mismatch cost of $O(1)$ per unit cube inside Bałaban's budget $p_0(g) \gtrsim R(g)^{d+2}$. F2′ compares the bound sizes $(x_j^*)^{\kappa_0}$, which are ordered on the real trajectory, so no ordering of complex couplings is needed, and a slot-dependent radius $\rho_j \propto x_j^2$ would have failed Lemma N3. Verifier: `SOURCES_HHOL_LF_VERIFY.md`, [VERIFIED REDUCTION], corrected: R_k is unbounded, of order $(\log K)^r$, but largest at $k = 0$ and non-increasing in k (the file reverses the index); $\theta = 0.174$, not 0.26; (M-D) is not a convention, since the frozen map equals Bałaban's map only at x^* and its derivative there is (H-hist) plus (H-thr). Open inputs: (M-A), reduced to (H−) and CMP 102 (189) (ledger A4, with A18 and A19), with (M-A′) closed; (M-C), a verified reduction with open (I-ε), the (I-V0) residual, and (M1); (M-D), which needs real (H-hist) (B13) and (H-thr) (B15) with (H-hist2) (B14); (M-B) is closed, [PROVED+VERIFIED] with $p_0 \geq 17/(\gamma_0 A_1^2)$ and $\theta_W + \tau_C \leq 0.2$.

`SOURCES_MA.md`**: (M-A), the equipartition bound.** For the cut-off ratio density of CMP 122 I (1.100) at real weight $0 < u \leq x^{-2}$, the file proves $\langle A - A_{\min}\rangle_u \leq (n_i/u)(1 + \eta_x)$, where $n_i = (N^2-1)L_i$ counts the real coordinates on the free links and $\eta_x = 0.68M_0^2x^2p_0(x)^2 + 2c_HM_0^2x\,p_0(x)^3$; the same bound with the full A is false in general, because the background minimum $A_{\min}$ is $O(\varepsilon^2)$ per block. As a corollary, the freeze-mismatch average is at most $e^{0.1906\,n_i(1+\eta_x)}$, that is $e^{0.77(N^2-1)(1+\eta_x)}$ per scale-k unit cube ($e^{2.3}$ for SU(2)), additive in the volume. A virial identity on the convex cutoff ball, centred at the true minimizer X^* of the exponent, bounds $u\langle(X-X^*)\cdot\nabla G\rangle_u$, $G = A - A_{\min}$, by n_i plus a Haar-Jacobian term (Lemma J: $0.68N(N-1)r^2$ per link for $r \leq \frac{1}{2}$). A segment bound from the Hessian lower bound (H−) (Lemma S) turns this into the bound on $\langle G\rangle_u$, and integrating $\frac{d}{du}\log D^G = -\langle G\rangle_u$ gives the corollary. Verifier: `SOURCES_MA_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON (H−)]; corrections: the proved constant is $C = 1$, not $\frac{1}{2}$ (sharp when X^* lies on the boundary), the SU(2) toy value 1.87 needs a cutoff norm that the file does not state, and the bookkeeping item (M-A′) is a real gap, closed in `SOURCES_MA2.md`. Open inputs: (H−), the Hessian lower bound along segments with c_H uniform in k, volume, and background; `SOURCES_MA2.md` reduces it to CMP 102 (189) (ledger A4) together with A18, A19, and its conditions (C1), (C2); ledger B17.

`SOURCES_MA2.md`**: (M-A′) and (H−).** The bookkeeping item (M-A′) closes: every normalized ratio of CMP 122 I (1.100) sees only $G = F - A_{\min}$, and the exact identity $N_x/D_x = e^{-uF}\,[e^{uA_{\min}}/D^G(x^{-2})]\,e^{(u-x^{-2})G}$ keeps $e^{uA_{\min}}$ at the coupling u, so the mismatch costs at most $e^{1.07(N^2-1)(1+\eta_x)}$ per unit cube. The Hessian bound (H−) of `SOURCES_MA.md` is derived in the form $D^2F[h,h] \geq -c_H\varepsilon_k\sum_b \|h_b\|_2^2$, with c_H uniform in k, volume, and regular backgrounds. Since $A_{\min}$ does not depend on V', $D^A(s) = e^{-sA_{\min}}D^G(s)$, and the error in `SOURCES_HHOL_LF.md` §3.4(c) was a numerator at u over a normalizer at x. For (H−), the chain rule per plaquette is combined with one Cauchy step on $\delta\mathcal{H}/\delta B$ minus its linear part on the complex ball of CMP 102 (172), where the $1/\varepsilon$ of the radius cancels the ε of CMP 102 (187), and the kernel decay of (189) closes a Schur test. Ver-

ifier: `SOURCES_MA2_VERIFY.md`, (M-A′) [PROVED+VERIFIED] and (H−) a [VERIFIED REDUCTION] to CMP 102 (189); corrections: the first Hessian term is ≥ 0, $c_N = O(1)(B_0C_4+C_2)B_5$, c_H depends on M^5 and M_0, and $A_{\min}$ may be only piecewise analytic in the background, so $e^{uA_{\min}}$ must stay paired with an analytic factor or inside modulus bounds. Open inputs: (C1) $M_0\varepsilon_k \leq C_1\varepsilon_1$; (C2) the localized map U_{k,X_i} obeys the CMP 102 bounds; (C3) CMP 102 (189), announced without proof and already used in Bałaban's own chain (ledger A4, now a verified reduction to A1); ledger A18 (including the k-uniformity of C_2, C_4, B_0) and A19; and the reading that Bałaban's chain removes $e^{uA_{\min}}$ by algebra.

`SOURCES_MB.md`**: (M-B), the large-field bookkeeping.** Bałaban's large-field bookkeeping CMP 122 II (1.73)–(1.99) survives the complex-coupling perturbation with thresholds frozen at the real trajectory x^*: the per-cube factors $\exp(-p_0(x_j^*))$ and the per-component factors $\exp(-2p_0(x_j^*))$ keep the same ε_j, δ_j, δ'_j, R_j, and the only new condition is $p_0 \geq 17/(\gamma_0 A_1^2)$ with Bałaban's output factor kept (option B). The cost constant becomes $C'_c = C_c + C_m + C_G + C_E$, a fixed $O(1)$ per site inside the $O(1)\log g_j^{-2}$ budget, and the p. 389 condition $p_0 \geq 8\kappa(100R_k)^d + 4$ is unchanged. Each line of (1.73)–(1.99) that carries γ_0, p_0, ε_j, δ_j, A_1, or a per-cube factor is listed (L1–L24) and redone with $\gamma_0 \to g^*\gamma_0$, which is legitimate because each factor bounds a positive Wilson action or quadratic form whose modulus at complex u is dominated with prefactor $\operatorname{Re} u \geq 0.8264\,(x^*)^{-2}$. The combined factor of p. 381 then needs $p_0 \geq c/(\gamma_0 A_1^2)$ with $c = 2/(g^* - \frac{1}{2})$, and the budget constant $C'_1 = \max(1, C'_c)$ (route (b) of `SOURCES_FX_R2.md`) multiplies (1.82), (1.88) by a fixed factor that p_0 large and γ small absorb. Verifier: `SOURCES_MB_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON (M-A) AND (M-C)]; corrections: the perturbation weights are additive, so $g^* = 0.82645 - \theta_W - \tau_C$ (not a product), and the constant 17 needs the margin $\theta_W + \tau_C \leq 0.2$ (strict limit 0.2088; at $\frac{1}{4}$ one needs $c = 26.2$); p. 383 prints $\frac{1}{2}\gamma_0$ and N^{β_0}, and any fixed $g^* > 0$ is accepted there. Open inputs: (M-A) (the freeze-mismatch cost C_m) and (M-C) (the complex-u activities α^c and E-terms); the link from F2′ to α stays an inference; the margin is ledger C2, and the block feeds ledger B17.

`SOURCES_MC.md`**: (M-C), the complex small-field step.** Lemma MC: if Bałaban's constants satisfy the conditions B1–B23 of CMP 116 §2, then with the shrinks $\varepsilon'_1 = \varepsilon_1/(4\lambda_V)$, $\gamma' = \gamma/(8\lambda_V)$, $\alpha' = \alpha/4$, $\lambda_V = \max(c, 1.235)$, the radius $\rho_0 = \min\{0.1, (LM)^{-2}(0.45a/(16n))^{1/2}\}$, and the added condition (M1), the small-field step is holomorphic in complex g with $|g/x - 1| < \rho_0$, for every real $x \leq \gamma'$ and inputs in the ball cE_0, and Lemma 3 (2.38) holds with Bałaban's $C_3\varepsilon_1$. Hence β_j is holomorphic on these discs and bounded uniformly in j, k, and volume. The 23 inequalities of CMP 116 §2 are read on the page images and typed, and each one at complex $w = (x/g)^2$ and primed constants is shown to follow from Bałaban's own. The vertex is affine in w with real-domain coefficients, so its factor λ_V is absorbed by the τ-circle (2.18) with $E_0 \to \lambda_V E_0$ and a smaller ε_1; the cut stays real, and the Gaussian cost $16n\rho^2$ per site and the α_5 items fit into the budget $(LM)^4\alpha_i \leq \frac{1}{4}$ under (M1) and (M2). Verifier: `SOURCES_MC_VERIFY.md`, [VERIFIED REDUCTION] to the printed CMP 116 §2 conditions (B1–B23 and four more), Lemma 2, (1.36), and p. 12 (2.5)–(2.6); corrections: the shrinks are $\varepsilon_1/(4\lambda_V)$ and $\gamma/(8\lambda_V)$, not ε_1/λ_V and γ/λ_V, the total output is $\leq E_0$, not $\frac{1}{2}E_0$, and the open item (I-X) is resolved by p. 12 (the X-measure is the unit Gaussian). Open inputs at the time of the verifier: (I-ε), that the smaller ε_1, α_0, α_1 keep CMP 116 §1 and the Part I induction valid; the (I-V0) residual (the $\log\sigma$ and Jacobian part); and (M1), an "M large" condition; ledger B17. All three are now settled, see the next four paragraphs: (I-V0), (I-ord), and (I-ε) in both layers are closed and verified, and (M1) is the convention C24.

`SOURCES_MC_IV0.md`: **(I-V0), the vertex part of (M-C).** The w-dependence of the bracket $\{\dots\}$ in CMP 116 pp. 9–10 is only the E_k difference. It is linear in w and free of w otherwise. $\operatorname{Tr}\log$ and $\log\sigma$ sit in $P^{(k)}$ with prefactor 1, and $\log\sigma$ and the Jacobian do not depend on w. The five $1/g_k^2$ terms are each of the form $u\cdot f(C\tilde{B})$, so they are affine in w, and the shift restores (2.19), (2.20), and (2.26) exactly. The file reads pp. 9–11 and 16–17 on 300 dpi page images and checks each w-dependent term against the printed display. Verifier: `SOURCES_MC_IV0_VERIFY.md`, [PROVED+VERIFIED] for the closure of (I-V0); corrections: the reading of the integral as a Haar integral is an inference [I], since the page prints only $dB'\sigma(B')$; the ε_1^2 cut factor of (2.26) belongs to (I-ε). Open inputs: none for (I-V0); (I-lin) is a mild item (per-piece balls).

`SOURCES_T2_IEPS.md`: **(I-ε) and (I-ord), small-field layer.** Every condition on ε_1, α_0, α_1 in CMP 116 §1–§2 and CMP 109 §1, §3 is an upper bound, a ratio of degree one, or a lower bound relative to a later constant; none is two-sided. So the shrinks of Lemma MC keep these papers valid: the ε_1^2 cut of (2.26) is restored exactly by $\gamma' = \gamma/(8\lambda_V)$, and scaling (A_0, A_1) by $1/\theta$ restores CMP 122 II pp. 380–381 exactly. (I-ord): the order $d, L \to \delta \to \kappa \to \kappa_1 \to$ walk constants $\to M \to \beta, \alpha' \to \alpha_1 \to \alpha_0 \to \alpha_3 \to \alpha_2 \to E_0 \to \varepsilon_1 \to \rho_0 \to A_1 \to A_0, p_0 \to \gamma$ has no cycle. Verifier: `SOURCES_T2_IEPS_VERIFY.md`, [PROVED+VERIFIED] for CMP 109 and CMP 116 (five missed conditions found, all of an allowed type); [OPEN] at that time for CMP 119/122, because CMP 119 (2.28) defines $\alpha_{i,j} = g_j C_i (\log g_j^{-2})^{q_i}$ with C_0, C_1 "sufficiently large"; the restoration of (L1) by a smaller γ in the large-field layer was unsupported. (I-ord) is closed on the pages read. Open inputs: the large-field layer, closed in the next paragraph.

`SOURCES_T2_IEPS_LF.md`: **(I-ε), large-field layer.** The shrink of Lemma MC reaches $\alpha_{i,j}$ (CMP 119 p. 262 takes the restrictions from CMP 109). Every printed condition on C_i, q_i is a lower bound, a ratio ($q_0 \le q_1$), or an upper bound on g; none bounds C_i from above. Lemma MC needs only a factor-4 slack, so C_i and q_i stay fixed and γ is lowered. The order continues $\rho_0 \to$ exponents $\to A_1 \to A_0 \to C_0 \to C_1 \to \gamma$, with no cycle. Residue [I]: α_2–α_4 and γ_2 are not printed in CMP 119/122 (all three branches close); $q_0 > p_0$ versus $q_0 = p_0$ is not printed (both close). Verifier: `SOURCES_T2_IEPS_LF_VERIFY.md`, [PROVED+VERIFIED]: all four claims confirmed; corrections: the condition C3 is a printed inclusion with an unprinted constant $c(B_3)$, and $q_0 \ge p_0$ is forced; the p. 381 factor is $\exp(-\gamma_0 A_1^2 p_1^2(g_j))$ with no extra weight; one inequality on ν reverses ($\frac{1}{2}\nu \le p_0 - p_1 - 1$); $\gamma' < \min(e^{-q_0}, e^{-q_1})$ is a further upper bound on γ, harmless. Open inputs: none beyond the two [I] residues above, each of which closes on every branch.

`SOURCES_HIGHER_M1.md`: **higher derivatives, and (M1).** Part A: no consumer in the coupling and threshold chain needs derivatives of order greater than 2 in the couplings or thresholds. Part B: the condition (M1), "M large", is a block-size choice. The file lists 26 lower bounds on M (for example $M \gg \kappa_1$, $M \ge M(\kappa)$, $M \ge cL$, $M \ge 48\kappa_1/\varepsilon$, $M/M_1 > 3(6 + c_X)$) and no upper bound, and the order of constants puts M before α. So (M1) is the convention C24: M is chosen at least the largest of these lower bounds. Verifier: `SOURCES_HIGHER_M1_VERIFY.md`, [PROVED+VERIFIED] with minor corrections: Part A is restricted to the coupling, threshold, and MA2-field variables (Bałaban's field expansion goes to order 4, and the source derivatives of D2/D3 to all orders); the lower bounds of `SOURCES_A3_SIGMA.md` 5.1 and 5.3 join the list. Open inputs: none; (M1) leaves ledger B17 and becomes ledger C24.

5.5 Step 6: thresholds and second history derivatives (ledgers B15, B14)

`SOURCES_HTHR.md`: **(H-thr), first attempt.** With the threshold mismatch $\Delta_j(c) = B_j(u(c);\bar{u}) - B_j(u(c);u(c))$ (integer parameters frozen, convention L3), the file bounds $\sup_c |\Delta_j'(c)| \le 3(1+3r)A\,\Phi(g_j^{-2})$ and sums this along the trajectory to $C\beta^{-1}\gamma^{-2(1+a)}(\log\gamma^{-2})^P$ times a decay factor, uniformly in K; in the verifier's corrected form the decay is $\exp(-(\log\gamma^{-2})^q)$ with $q = 2p_1 - (d+5)r_0 > p_0$, in place of $\exp(-\kappa(\log\gamma^{-2})^{2p_0})$. Lemma H′ then splits the budget: $x \le \ln(6/5)$ for (H-hist) and $x_{\mathrm{thr}} \le \frac{1}{4}$ give $\sum_j |B_j'| \le \frac{1}{2}$ and $\frac{1}{2} \le |v_k| \le \frac{3}{2}$. Coarea writes Δ_j as the contribution of the shell between the two thresholds, which reduces Δ_j' to threshold derivatives and mixed threshold–weight derivatives of B_j, bounded in Lemma S under the model assumptions M1–M4. The sum along the trajectory uses $u_i \ge \gamma^{-2} + \beta m$ and an integral comparison that keeps the factor γ^{-2}/β, and Lemma H′ is an additive Gronwall argument. Verifier: `SOURCES_HTHR_VERIFY.md`, [VERIFIED REDUCTION] of (H-thr) to M2′, M4, M5, with the shell representation, the trajectory sum, and Lemma H′ confirmed; corrections to Lemma S: the fluctuation cutoff is $\delta_j' = g_j A_1 p_1(g_j)$ with $p_1 < p_0$, Bałaban bounds one bond only through the block quadratic form CMP 122 II (1.78) with loss $R_j^{-(d+5)}$, so the per-link density M2 is unsupported, and the mixed derivative needs history holomorphy (M5); M1 is resolved (χ stays inside the integral, so $\Delta_j \ne 0$). Open inputs: M2′ (density form of the block bound), M4 (extraction cost), M5 (history holomorphy), M3, and the frozen-integer convention L3 (ledger B15, C1); `SOURCES_HTHR2.md` replaces M2′ by Lemma T and M5 by (H-hist2).

`SOURCES_HTHR2.md`: **(H-thr), Lemma T.** The shell indicator is at most the large-field function $1-\chi$ at the lower threshold, so Bałaban's printed tail factor of CMP 122 II (1.77)–(1.78) (with the misprint $p_1 \to p_1^2$) bounds Δ_j itself with exponent at least $\kappa_2(\log s)^q$, $\kappa_2 = \frac{1}{2}L^{-(d+5)}$, $q = 2p_1 - (d+5)r_0 > p_0$ (Lemma T). With a uniform bound D on Δ_j'' this gives $\sup_c|\Delta_j'| \le 2(2AD)^{1/2}\Phi_T(u_{j-1})^{1/2} + (2A/r)\Phi_T(u_{j-1})$, a sum over j that is uniform in K under (G1′)–(G3′), and Lemma H′ of `SOURCES_HTHR.md` unchanged. The zero-order bound on Δ_j uses Lemma T, the history weights M3, and the extraction cost M4, a Cauchy estimate on the analyticity radius $\alpha_{1,j} = g_j C_1(\log g_j^{-2})^{q_1}$ of CMP 119 (2.28), so no history derivative enters. The derivative then comes from the interpolation inequality $\|\Delta_j'\| \le 2(\|\Delta_j\|\,\|\Delta_j''\|)^{1/2} + \|\Delta_j\|/r$ (Lemma 7.1(a) of `SOURCES_D2.md`), which replaces the history holomorphy M5 by a uniform second-derivative bound (H-hist2). Verifier: `SOURCES_HTHR2_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON (H-HIST2)] for the summed bound, with Lemma T and the misprint confirmed and M4 confirmed as reduced; correction: (H-hist2) for β_j contains the mixed threshold–history derivative through the running thresholds, so it is a new hypothesis of the (H-hist) kind, not free by a Cauchy estimate in the history. Open inputs: (H-hist2) (ledger B14, a [VERIFIED REDUCTION] to the real uniform (H-hist), ledger B13); M3; M4 (units, and the small-field analogue of CMP 122 II (1.73)); the transfer of Bałaban's printed large-field factor to small-field shell terms; the frozen-integer convention L3; ledger B15.

`SOURCES_HHIST2.md`: **(H-hist2), second history derivatives.** For the frozen-threshold map $F = B_j(\cdot;\bar{u})$ on $|w-\bar{u}| \le 3r$ in u-slots, $\sum_{i,l}|\partial_{u_i}\partial_{u_l}F| \le \varepsilon_j'' = C_H(1-\lambda^{1/2})^{-2}u_{j-1}^{-2}$, so $\sum_j \varepsilon_j'' \le C_H(1-\lambda^{1/2})^{-2}\gamma^2/\beta$ uniformly in j, K, and volume, with product decay $\lambda^{m/2}$; the first-derivative half is $\varepsilon_j' \le \frac{1}{2}C_1(1-\lambda)^{-1}u_{j-1}^{-3/2}$, from the real uniform (H-hist). For β_j with running thresholds the (H-hist2) step needs only a uniform bound, but Lemma H′ still needs a summable first derivative of β_j from the real (H-hist). Route 2(a) differentiates the real chain twice: past slots stay real, the input variable is handled by Cauchy estimates (Theorem SM), and only the own slot is complexified, on

one relative u-disc (SOURCES_SM.md §4.2). The Jacobians $\partial_{u_i}\partial_{u_l} = \frac{1}{4}g_i^3 g_l^3 \partial_{g_i}\partial_{g_l}$ turn the $1/x$ losses into g^4, which is summable because $u_{j-1} \geq \gamma^{-2} + \beta m$. Verifier: SOURCES_HHIST2_VERIFY.md, [VERIFIED REDUCTION] to the real uniform (H-hist), no new input; the Jacobian cancellation and the decay $\lambda^{m/2}$ are confirmed, the product form λ^m is [REFUTED] (toy recursion), and $\varepsilon'' \propto u^{-2}$ follows from the u-disc alone; correction: "β_j needs only a uniform bound" holds for the (H-hist2) step only, since Lemma H′ needs a summable β_j'. Open inputs: the real uniform (H-hist) (ledger B13: (Q)/c_-, (M_r), CMP 102 (189), (M-B) bookkeeping); Theorem SM and the joint u-disc and input-ball holomorphy [R]; the large-field part of the frozen map; for β_j, the mixed threshold–history derivative through Lemma D and M3 (ledger B14).

SOURCES_SM.md: **Theorem SM, the step-map Lipschitz bound.** The small-field step map factors as $G = \hat{G} \circ \mathcal{L}$ with $\mathcal{L}$ linear, and Theorem SM gives $|DG(I_0)[W]| \leq \theta_0 \sum_{l\leq k} L^{-\alpha_*(k-l)} \|W^{(l)}\|\, e^{-\kappa' d_{k+1}(X)}$ with $\theta_0 = 2K_H K_V \varepsilon_1$ and $\alpha_* = \min(1,\alpha)$. With it Lemma HH holds at rate $\lambda = L^{-\alpha_*} + C_V\theta_0$ when $C_V\theta_0 < 1 - L^{-\alpha_*}$, and (Eps) of SOURCES_D3W8.md becomes a corollary with $\varepsilon' = \theta_0/(1 - L^{-\alpha_*})$. The per-input bound CMP 116 (1.24) gives the linear stage a fluctuation factor $\alpha_2^{-1}\varepsilon_1$ and a label weight $L^{-(k-l)}$ (Fact A). The map $\hat{G}$ is holomorphic on a ball of radius $R_V = \alpha_4\alpha_6 e^{-5\kappa}/(4(L+2)^4 O(1))$, free of ε_1 and E_0, by CMP 116 (C1)–(C5) (Fact B), and a Cauchy estimate on this ball gives the contraction. Verifier: SOURCES_SM_VERIFY.md, [VERIFIED REDUCTION] (items 1–4 confirmed on page images, items 5–6 corrected); the own-slot constant C_β has the same $1/g$ gap as S_0, so (G-gain) must be applied in both the hop and the own-slot bound, and the per-label decay of the lower-order terms stays reduced. Open inputs: Fact A for the lower-order bracket terms (ledger A10, now Theorem 5.6 in the corrected form A10′; the per-label exponent is $\alpha_* = \beta_H$, a free parameter, and (Eps) reads $\varepsilon' = \theta_0(1 - L^{-\beta_H})^{-1}$); Fact B, that is (W4) on CMP 116 (C1)–(C5) with α_4, α_6 fixed before ε_1; (I)/L3 (ledger B12); (G-gain) in both slots, carried by SOURCES_GGAIN.md and SOURCES_GGAIN_LF.md (ledger B13).

5.6 Step 7: history bounds and the gain (ledger B13)

SOURCES_HHIST.md: **Lemma HH, (H-hist) in column-sum form.** Given CMP 109 (0.29), uniform own-slot bounds $\|\partial_{g_{l-1}} V^{(l)}\|_l \leq S_0$ and $|\partial\beta_l/\partial g_{l-1}| \leq C_\beta$, and a step-map Lipschitz bound (SM) with $\theta < 1 - L^{-\alpha}$, Lemma HH gives $|\partial\beta_j/\partial g_i| \leq (C_\Pi/C_V) S_0 \lambda^{j-1-i}$ for $i \leq j-2$, with $\lambda = L^{-\alpha} + \theta$. The column sum is then at most $C_\beta + (C_\Pi/C_V)S_0\lambda/(1-\lambda)$, uniformly in i, j, K, and the volume. The past enters step $k+1$ only through the curly bracket of CMP 109 (2.12) and only through the irrelevant remainders $V^{(l)}$ (the marginal parts cancel, p. 298), and a step-l remainder costs $L^{-\alpha(k-l)}$ per π_k cube by (0.29)–(0.30). An induction on $s_l = \|\partial_{g_i} V^{(l)}\|_l$ with one hop per step then gives the rate λ in place of the one-hop rate $L^{-\alpha m}$. Verifier: SOURCES_HHIST_N1_VERIFY.md (Part 1), [VERIFIED REDUCTION]: the quotes, the dependence chain, the cube count, and the slot definition are confirmed; the rate is sharp, but the numerical claim "ratio 1.0" is false ($s_{i+1+n} = (\theta/\lambda)\lambda^n S_0$, so the bound is loose by θ/λ), and Bałaban gives no per-hop Lipschitz constant θ. Open inputs: the own-slot bounds S_0 and C_β (C_β is Bałaban's inductive assumption, ledger A5) and (SM), now carried by SOURCES_GGAIN.md and SOURCES_SM.md; validity of the step bounds off the trajectory, close to (I) (ledger B12); the large-field contributions, now in SOURCES_GGAIN_LF.md; the whole chain is ledger B13.

SOURCES_GGAIN.md: **(G-gain), small field.** For every step k and $g_k \in]0,\gamma]$ the own-slot derivatives obey $\|\partial_{g_k} V^{(k+1)}\| \leq C_V S_0$ and $|\partial\beta_{k+1}/\partial g_k| \leq C_\beta := C_\Pi S_0$, with S_0 free of $1/g_k$ and of ε_1 (Theorem B2-unif). With Theorem SM the column sum of Lemma HH is uniform, and in u-slots it is $O(g_i^3)$, which meets the threshold $C g_i^{2+\delta}$ of SOURCES_HBETA.md with $\delta = 1$. By CMP 116 (1.34) the

vertex $V'_k(Y,\cdot,B)$ is an analytic function of $x = g_k B$ on a polydisc of g-free radius ε_1, so a pointwise Cauchy estimate in x costs $|B|_Y/\varepsilon_1$, the ε_1 cancels against the vertex size, and the one polynomial insertion is absorbed by the Gaussian at strength α_4. The $P^{(k)}$ part is cubic in B' by (1.39)–(1.40), so $|\partial_g| \le C_P|B|^3$ there, and in the hop the factor $g_k|B|$ of (1.22)–(1.24) gives $\theta_0 \to \theta_g g_k$. Verifier: `SOURCES_GGAIN_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON LEMMA G3 AND THE MARGIN (M_r)], for the small-field step on $]0,\gamma]$ including vacuum subtraction (items 1, 2, 4, 5, 6 confirmed, 2 and 6 with corrections, item 3 corrected); (M_r) is also needed for the $P^{(k)}$ part, Lemma G4 fails as stated and is off-chain, the sharp-cut surface constant is $O(\gamma_2^{-(n_b+1)/2}\varepsilon_1^{-1})$, and Bałaban's p. 264 assumption is proved only in its first-derivative part, on $]0,\gamma]$, for the small-field step. Open inputs: Lemma G3, hence (Q)/c_- (round 5: proved relative to listed inputs, Theorem 5.9); the margin (M_r), which shrinks the χ_k threshold (ledger C9); the large-field half (`SOURCES_GGAIN_LF.md`); the endpoint $g = 0$ (ledger C19); all inside ledger B13.

`SOURCES_GGAIN_LF.md`**: (G-gain), large field.** With the thresholds frozen at Bałaban's trajectory x^*, the own-slot derivatives of the large-field outputs are uniform in $g_k \in]0,\gamma]$ with $|g_k - x_k^*| \le \rho_0 x_k^*/2$, in k, K, and the volume, for example $|\partial_{g_k} R'^{(k)}(X)| \le 20\gamma^{2A_0-1}e^{-\kappa d_k(X)}$ and $|\partial_{g_k} T_k(Y_i)1| \le 20\gamma^{A_0-1}e^{-\frac{1}{2}p_0(x_k^*)}$ (Theorem B2-full). With `SOURCES_GGAIN.md` and Theorem SM this makes the real uniform (H-hist) a reduction in both halves. On the disc $|g - x| < \rho_0 x$, $\rho_0 = 0.1$, of `SOURCES_HHOL_LF.md` the printed bounds hold with $\gamma_0 \to 0.826\gamma_0$, and one Cauchy step gives $|\partial_g F| \le 20x^{-1}\sup|F|$. The factor $1/x$ is absorbed by the large-field suppression, since $p_0(x) = A_0(\log x^{-2})^{p_0}$ with p_0 a positive integer and A_0 large (CMP 119 p. 244) gives $x^{-1}e^{-\theta p_0(x)} \le x^{2\theta A_0-1}$. Verifier: `SOURCES_GGAIN_LF_VERIFY.md`, [VERIFIED REDUCTION] (items 1, 2, 4, 5 confirmed, item 3 corrected); the mismatch average sits at $u' = 2x^{-2} - u \in [0.892, 1.093]x^{-2}$, which gives the constant $2.48 \le 2.6$, the Gaussian normalization adds a factor $e^{0.01n}$ per unit cube inside (M-B), and "β from E-class terms only" is upgraded to a verified reading of CMP 119 p. 279. Open inputs: (Q)/c_-, the margin (M_r), CMP 102 (189) through (M-A) (ledger A4), and the (M-B) bookkeeping (ledger B13); not covered: (1.103) at complex u, C^1 at $g = 0$, higher derivatives.

`SOURCES_G3.md`**: Lemma G3, the** χ_k **surface term.** Given (Q) with constants c_- and Λ, for every k and $g_k \in]0,\gamma]$ the g_k-derivative acting on the sharp cut $\chi_{k,Y_0}\chi^c_{k,P}$ of CMP 116 (2.3) in a term of (2.14) is at most $c'_1 U(g_k)e^{\delta\kappa d_k(Z_0)}T_{\mathrm{maj}}$, with $U(g) = A_n D_n \varepsilon_1^{-1} e^{-\varepsilon_1^2/(8\Lambda g^2)}$ and $n = N^2 - 1$. The cut adds $S_\chi = c_2 A_n D_n \varepsilon_1^{-1} e^{-\varepsilon_1^2/(8\Lambda\gamma^2)}$ to S_0, uniform in g_k, X, k, and the volume, and $O(\gamma^p)$ for every p. Route 2 keeps Bałaban's cut, a round sphere $|B(b)| = \varepsilon_1/g_k$ per bond, and computes the surface term in polar coordinates, where the Gaussian marginal turns $g^{-1}(\varepsilon_1/g)^n$ into $\varepsilon_1^{-1}t^{n+1}e^{-ct^2}$, $t = \varepsilon_1/g$, with no $1/g$ left. The bond and polymer sums use (2.30), and the smooth cut $\hat\chi$ (Route 3, $\varepsilon_1 \to \varepsilon_1/2$, $\gamma \to \gamma/2$) is the fallback. Verifier: `SOURCES_G3_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON (Q)] (items 2, 4, 5 confirmed, items 1 and 3 corrected); the surface term is real in the fixed-B computation that `SOURCES_GGAIN.md` uses (finite-difference check to 10 digits at $n = 8$), (Q) must be read on the joint (X, B) majorant with $a = 0$, the bond count is $c_d|Y_0| + |P|$, the printed γ_k fails under both parses, and the B versus B' cut question the verifier added is closed in `SOURCES_Q.md`. Open inputs: (Q)/c_- (row B-S.Q; ledger B13), now discharged for the small field by Theorem 5.9; numerical values of c_- and Λ; higher derivatives.

`SOURCES_Q.md`**: input (Q), the covariance lower bound** $c_- > 0$**.** The $B(b)$-block of the covariance of Bałaban's joint Gaussian majorant CMP 116 (2.23) is bounded below by $c_- > 0$ and above by $\Lambda \le 2(1 + \gamma_0^{-1})$ for $\alpha_5(1 + \gamma_0^{-1}) \le \frac{1}{2}$, uniformly in k, the volume, and regular U; after the localization step of `SOURCES_LOC.md`, $c_- = (\gamma_1 + \varepsilon_M)^{-1}$, and in the free case $c_- \ge (a\kappa_d L^{d+1})^{-1}$.

The cut of CMP 109 (2.9) is $|B(b)| < \varepsilon_1/g_k$ on every bond except b_0, a round sphere with no Jacobian. The precision of (2.23) is $Q = Q_0 - \alpha_5 \operatorname{diag}(Z, Z_0) \le Q_0$, and inversion reverses order, so the $B(b)$-block of Q^{-1} dominates that of Q_0^{-1}, which is $C_0 + C_0\Gamma\Gamma^* C_0 \ge C_0$ with $C_0 = C^{(k)}(Z_0, 0)$. The remaining norm input $\|C^*\Delta_k C\| \le \gamma_1$, which CMP 116 (2.7) itself uses, is traced to CMP 99 (3.132), and the free-case constants come from CMP 95 (1.100) and CMP 96 (2.125). Verifier: `SOURCES_Q_VERIFY.md`, [VERIFIED REDUCTION] to CMP 99 (3.132), (E_U), and LOC (items 1, 3, 4 confirmed, 1 and 3 with corrections, item 2 corrected in favour of the reduction); Q_0 is coupled, so the block identity is only an inequality, the operator is the $\sigma = 0$ localized one (handled in `SOURCES_LOC.md`), CMP 99 p. 422 prints the kernel bound (3.132), and κ_d depends on a and L unless $a = 1$. Open inputs: the A1 residue (torus-side hypothesis, two conventions, reading $a = 1$); the two [I] bridges $Q(U) = Q_{\mathrm{free}}(1 + O(\alpha_0))$ and $\|C(U)\|^2 \le L^{d+1}(1 + O(\alpha_0))$ (ledger A18); (E_U), reduced to k-uniform kernel moments that Bałaban does not print; the inputs of LOC. Round 5 assembles (Q) from A7 and A8 instead (Theorem 5.9); its Λ corrects 4.3 of this file, because G_Γ enters the B-block of Q_0^{-1}.

`SOURCES_LOC.md`**: Lemma LOC, the localized covariance $C^{(k)}(Z_0, 0)$.** Given $\gamma_0 \le C^*\Delta_k C \le \gamma_1$ and $C_0^{-1} = Z_0 C^* \Delta_k C Z_0 - R$ with $\|R\| \le \varepsilon_M \le \gamma_0/2$, the localized covariance $C_0 = C^{(k)}(Z_0, 0)$ of CMP 116 (2.23) has $C_0(b, b) \ge c_- I$ on every bond $b \subset Z_0$ and $\|C_0\| \le 2/\gamma_0$. Here $c_- = (\gamma_1 + \varepsilon_M)^{-1}$ with the symmetry (S) of the truncated walk sum and $c_- = (\gamma_0 - \varepsilon_M)/(\gamma_1 + \varepsilon_M)^2$ without it, and $\varepsilon_M = S_d(O(1)e^{-\delta_0 M/3} + O(\alpha_0 + \alpha_1))$ with $S_d \le d \coth(\delta_0/(4\sqrt{d}))^d$. The printed (2.5) is conditioning on Z_0^c, so $C^{(k)}(Z_0)$ is the inverse of the compression of $C^*\Delta_k C$ and inherits its bounds, while $C^{(k)}(Z_0, 0)$ is the walk sum with the walks meeting σ_0 deleted and is not a compression. Its distance to the compression is Bałaban's bound (2.16) applied to R_2 at $\sigma \equiv 1$, turned into the norm bound ε_M by the Schur test. Verifier: `SOURCES_LOC_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON THE R_2 BOUND (2.16), γ_0, AND γ_1] (item 1 confirmed, items 2–4 corrected); the symmetry (S) is not needed, $\varepsilon_M \to 0$ needs $\alpha_0, \alpha_1 \to 0$ as well as $M \to \infty$, γ_0 sits in the prose after CMP 99 (3.158) and is proved there only at $U = 1$, and the replacement point is the real configuration U. Open inputs: the R_2 bound (2.16), which Bałaban states with "similarly" (ledger A2, now a verified reduction to (W^c), A7, and two [I] items); the upper bound γ_1 through CMP 99 (3.132) and (E_U); γ_0 for general U (ledger A3, now a verified full proof, $\gamma_0 \ge m_L/(48d^2L^d)$, with residue R-$\Delta$2); α_0, α_1 small after M is fixed.

5.7 Step 8: Bałaban-level propagator inputs (ledgers A1–A4)

`SOURCES_A1_KERNEL.md`**: ledger A1, kernel-decay form.** Given (K) and (C) below, the file proves CMP 99 (3.132) in its printed two-scale form, $|(QGQ^*)^{-1}(y, y')| \le 2K_{\mathrm{ext}} L^{d/2+1}(L^j\eta)^{-2}(L^{j'}\eta)^{-d} e^{-\delta_1 d(y,y')}$ for $y \in \Lambda_j$, $y' \in \Lambda_{j'}$, uniformly in j, k, η, the volume and regular U, and the same for $(QG_1Q^*)^{-1}$ plus a diagonal a-term. The rate comes from $\varepsilon = \min\{\delta/2, c_-/(2c_+\Sigma(\delta/2))\}$ and is much smaller than Bałaban's rate. Lemma CT, a Combes–Thomas argument on $\hat{N} = S^{-1/2}(QGQ^*)S^{-1/2}$ on $\ell^2(\mathfrak{B}, \mu)$, turns kernel decay (K) of QGQ^* and a lower bound (C) $\hat{N} \ge c_-$ into decay of the inverse, with no Fourier analysis and no use of A3. (K) is taken from CMP 99 Theorems 3.3 and 3.12 for G, (C) from one feasible extension in the variational form $\langle B, N^{-1}B\rangle = \min\{\langle A, G^{-1}A\rangle : QA = B\}$, and the global perturbation from $U = 1$ and the Demko–Moss–Smith route are shown not to work. Verifier: `SOURCES_A1_A4_VERIFY.md` (Part A), [VERIFIED REDUCTION]; it confirmed Lemma CT, the two-scale form, the non-circular order of the G_1 clause, and both refutations, and it found the file's face-midpoint bumps for (C) not feasible ($Q_j(U)$ is supported on both whole blocks, CMP 98 (14), CMP 99 (3.102)), so its own grade was "reduced to (C), open"; the newer `PLAN.md` entry regrades A1 to a verified reduction once (C) is

supplied by SOURCES_QE_INV.md, SOURCES_QE_KE_HQ.md, and SOURCES_S2_FORM.md. Open inputs: the residue of ledger A1, namely the torus-side hypothesis (side $\geq c \cdot RM$ coarsest blocks), two unprinted conventions, and the reading $a = 1$ of CMP 99 Theorem 3.1 [I]; the acceptance of the small rate δ_1 by each downstream consumer (S4) is not re-checked (low risk).

SOURCES_QE_KE_HQ.md: **(KE′) and (H_Q), inputs of (C).** (KE′): Bałaban's H of CMP 96 Cor. 2.8 already satisfies $Q_{98}(1)H = I$ on all of $\mathfrak{B}$, interfaces and Λ_0 included, with the decay of CMP 96 (2.151) for H and ∇H, because CMP 96 uses the straight-line average CMP 95 (1.11)/(1.18). (H_Q): $|K_j(U; c, b) - q_j(c, b)\,\mathrm{Ad}\,U(\Gamma_{c_- \leftarrow b})| \leq c_H \alpha_0 \ell_j^2 \eta_j^d$ on $B^j(c_-) \cup B^j(c_+)$, $c_H = 2C_1' + c_T(d)L$, uniformly in $j \leq k$. For (KE′) the file reads the pages: the invariance group $Q_j'\lambda = 0$ of CMP 96 (2.7) fits only the straight-line average, and CMP 98 p. 27 states that the propagators hold for the average (1.11). For (H_Q) it applies Bałaban's induction (144)–(145) to the difference between $Q_j(U)$ and the product of the main terms, and it compares the averaged background links with fine straight-line transports and fixed axial contours by a cone-area bound. Verifier: SOURCES_QE_KE_HQ_VERIFY.md, [PROVED+VERIFIED], (KE′) by citation on the page images (CMP 95 (1.11), (1.12), (1.18); CMP 96 (2.7), (2.20), (2.35); CMP 98 p. 27 verbatim), so the gap of SOURCES_QE_INV_VERIFY.md item 2 was a misreading; corrected: the verifier proved the one [R] step (iii) itself (ribbon of $\leq (d-1)L^2$ plaquettes, lasso decomposition, error $\leq 9.6(d-1)\alpha_0\ell_i^2$ for $\alpha_0 \leq \min(c_2, 1/(8(d-1))))$, the cited (112) is the wrong reference and is not needed, and the corridor fallback $H + d\psi$ is exact within layers but not across interfaces (not needed). Open inputs: the unvalued constants $C_1'(d, L)$ and $c_{\mathrm{cone}}(d)$, and two unprinted conventions ($U(c)$ the straight segment; translation-covariant axial contours), both in the residue of ledger A1.

SOURCES_QE_INV.md: **Theorem QE, input (C) of Lemma CT.** For the free multiscale minimizer dressed with axial transports, $\mathcal{E}_U$, the file proves, given (KE) and (H_Q), $\|Q(U)\mathcal{E}_U - I\|_S \leq c_Q M\alpha_0 + c_{\mathrm{far}} e^{-\delta_H RM/(16L^2)}$ in the S-weighted norm on $\mathfrak{B}$, uniformly in j, k, the volume, $\{\Omega_j\}$, and regular U. Hence $K_Q \leq 2$, and $A = \mathcal{E}_U(Q(U)\mathcal{E}_U)^{-1}B$ is an exact extension, $Q(U)A = B$ and $A = 0$ on Λ_0, with $\langle A, G^{-1}A\rangle \leq K_{\mathrm{ext}}\|B\|_S^2$, $K_{\mathrm{ext}} = 4(1 + O(M\alpha_0))K_{\mathcal{E}} + a$, which gives (C) with $c_- = 1/K_{\mathrm{ext}}$. The defect of $Q(U)\mathcal{E}_U$ is written exactly as a $q|E|$-weighted sum of loop holonomies minus one, and a cone-area lemma bounds each holonomy linearly in the distance. A Schur test with the decay (KE), a near/far split across layers with far terms e^{-cRM}, and a Neumann series then give $K_Q \leq 2$, and a crude absolute-value Schur bound gives the energy. Verifier: SOURCES_QE_INV_VERIFY.md, [PROVED+VERIFIED, CONDITIONAL ON (KE′), (H_Q), S2, AND A TORUS SIDE $\geq c \cdot RM$]; numerics [NUMERICAL, REPRODUCED] (defect linear in the curvature, slope 0.010 at $d = 2$ and 0.019 at $d = 4$, rigorous Schur constant 41–76 times the true slope); corrected: ε'' is of order $M\alpha_0$, not α_0, $\alpha_F = 2 \cdot O(1)M\alpha_0$ with $O(1) \geq 10$ (so the Neumann threshold reads $O(1)M\alpha_0 \lesssim 0.08$), and a torus-size hypothesis must be added. Open inputs: (KE′), (H_Q), and S2 for G have since been closed (SOURCES_QE_KE_HQ_VERIFY.md, SOURCES_S2_FORM_VERIFY.md); what stays open is the torus-side hypothesis and the two unprinted conventions of ledger A1.

SOURCES_S2_FORM.md: **S2, the form bound for G.** For every bond field A, $|\langle A, \Delta_\pi' A\rangle| \leq C_\Pi' M\alpha_0(\|DA\|^2 + \|D^*A\|^2 + \|\ell^{-1}A\|^2)$, hence $G^{-1} \leq (1 + O(M\alpha_0))(D^*D + DD^*) + O(M\alpha_0)\ell^{-2} + Q^*aQ$ and $\langle \mathcal{E}_U X, G^{-1}\mathcal{E}_U X\rangle \leq K(d, L, a)\|X\|_S^2$, uniformly in j, η, the volume, $\{\Omega_j\}$, and regular U. The route through CMP 99 (3.131) stalls, because (3.131) is printed for one-block supports and a tiling then forces sharp cut-offs that cost η^{-1} on the sup side. Route Π instead uses the exact identity (3.119), $\langle A, \Delta_\pi A\rangle = \langle \tilde{A}, \Delta\tilde{A}\rangle$ with $\tilde{A} = A - D\lambda$, $\lambda = G'RD^*A$, the curl of a gradient $\eta^{-2}(\mathrm{Ad}\,U(\partial p) - 1)\lambda$, and the block L^2 bounds (3.46) of Theorem 3.1 with CMP 96 Lemma 2.1 and

a Schur test, which give $\|\ell^{-2}\lambda\|, \|\ell^{-1}D\lambda\| \le B_0 c_1 \|D^*A\|$. Verifier: `SOURCES_S2_FORM_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON (KE′), $a = 1$ IN THEOREM 3.1, AND THE BLOCK GEOMETRY], together with §5.1 and §5.3 of `SOURCES_QE_INV.md`; corrected: Δ' of (3.10) has a second term $\eta^d \operatorname{tr}(DA)^2(\operatorname{Re} U(\partial p) - 1)$, bounded by $O(M\alpha_0)\|DA\|^2$, so $C'_{\Pi} = C_{\Pi} + 2c_R(1 + \kappa^2 a_0^2)$ and c_- drops only by a factor $1 - O(M\alpha_0)$; (3.120) carries a minus sign; the normalization of Q^*aQ is settled by (3.16) (one constant a). Open inputs: the reading $a = 1$ of Theorem 3.1 and the block-geometry item (both in the residue of ledger A1); the G_1 analogue, left at sketch level here, is closed in `SOURCES_S2_G1_V7.md`.

`SOURCES_S2_G1_V7.md`: **S2 for G_1, and the V7 repair of ledger A2.** Part A proves $|\langle B, \Delta^{(2)} B\rangle| \le c_2 M\alpha_0 \|\ell^{-1}B\|^2$, $c_2 = \dim \mathfrak{g} \cdot C_{137}$, and $|\langle A, \Delta^{(2)}_{\pi} A\rangle| \le 2c_2 M\alpha_0(\|\ell^{-1}A\|^2 + B_0^2 c_1^2 \|D^*A\|^2)$, so $\langle \mathcal{E}_U X, G_1^{-1}\mathcal{E}_U X\rangle \le K_1 \|X\|_S^2$ with $K_1 = K(1 + O(M\alpha_0))$. Part B proves $|R_2(b, b')| \le (O(1)e^{-\delta_0 M/3} + O(\alpha_0 + \alpha_1))e^{-(1-\theta)\delta_A r}$ for $M \ge \max\{M_*, 96\kappa_1/\delta_0\}$ and $\alpha_0 + \alpha_1 \le \alpha_*$. In Part A, (3.135) is the sandwich $(I - DG'RD^*)^*\Delta^{(2)}(I - DG'RD^*)$, so the form is taken at the $\tilde{A}$ of (3.119), and the kernel of $\Delta^{(2)}$ is local by (3.137) and symmetric as a Hessian by (3.136), which allows a Schur test with AM–GM. In Part B, $C(\sigma; U_c) = A_Z(U)^{-1} - W'$ with $W' = W_\sigma(U_c) + [C(1; U_c) - C(1; U)]$, the bracket is bounded by a one-variable Cauchy estimate on the complex walk bound, and a Neumann series around the real $A_Z(U)$ lets $A_Z(U)$ cancel in R_2. Verifier: `SOURCES_S2_G1_V7_VERIFY.md`, Part A [PROVED+VERIFIED, CONDITIONAL] on the conditions of `SOURCES_S2_FORM.md`, Part B [VERIFIED REDUCTION]; corrected: symmetry of $\Delta^{(2)}$ is load-bearing for the column sums, the Cauchy route needs (W^c) on the α'-space ($\alpha' \gg \alpha$), not only on the α-space that W_σ needs (already in the ledger, no new item), the two-constant theorem weakens the complex input to a sup bound but does not remove it, touching σ_0 cubes occur generically, not always, and the two covariances do not coincide on a gauge-fixed subspace; the lower bound for G_1 (A3.5) was not checked. Open inputs: for A2, (W^c) on the α'-space, now proved in the exponential gauge without the Thm 3.15 transfer (`SOURCES_WC_EXPGAUGE.md`, `SOURCES_WC_RESIDUE.md`, both verified) relative to the inputs listed under `SOURCES_A2_R2.md`, the real decay A7 [R], the boundary-contact reading of item (i) [I-forced], and $\operatorname{dist}(Z_0, \Lambda_k^c) > RM$ [I]; for Part A, the inherited reading $a = 1$ and the block geometry.

`SOURCES_A1_WALK.md`: **ledger A1w, walk-expansion form of A1.** Theorem A1-W: for regular U, $(QGQ^*)^{-1} = \sum_\omega T_\omega$ with factors localized in $X \times X$, U-dependence on $\tilde{X}_0^5 \cup \cdots \cup \tilde{X}_n^5$, and $|T_\omega(y, y')| \le (4/c_-)L^{d/2+1}(L^j\eta)^{-1-d/2}(L^{j'}\eta)^{-1-d/2}(K_W/M_1)^{|\omega|}e^{-\frac{1}{2}\varepsilon d(\omega, y, y')}$, absolutely convergent; this is the (3.108) form at rate $\delta_W = \frac{1}{2}\varepsilon \ll \frac{1}{2}\delta_0$, and the same holds for $(QG_1Q^*)^{-1}$ with trees. The local operator on a cube is G's walk sum truncated to $\square^+$, its positivity $\frac{1}{2}c_-$ comes by restriction of the global (C) (not A3), and an accretive Lemma CT gives $\|C_\square\|_\varepsilon \le 4/c_-$, as in Bałaban's own local step (CMP 96 p. 237). The parametrix $P = \sum h_\square C_\square h_\square$ satisfies $\hat{N}P = I + \sum F_\square$ with three remainders (separation, truncation, commutator), each $\le K_F/M_1$, and the Neumann series is the walk series. Verifier: `SOURCES_A1_WALK_VERIFY.md`, [VERIFIED REDUCTION] (construction verified); corrected: (L1) must read "kernel vanishes within about M_1 of ∂X", CMP 96 (2.84) has $e^{-\frac{1}{8}\delta_0 M}$ and (2.82) three error sums, the $\breve{G}_2$ citation is p. 432, the RM condition (2.59) at the walk rate joins the constant order, (W$_G$) has proof sketches printed only for G', $(Q'G'^2Q'^*)^{-1}$ and G_0 (Theorems 3.7, 3.9, 3.10), and (C) is conditional, not proved. Open inputs: (W$_G$) [R] (itself reduced in `SOURCES_WG.md` to Bałaban's printed sketches, the Neumann series (3.130), and CMP 96 Lemma 2.1, with WG-1 in the M_1 form $\alpha_0 \le c' M^{-1} M_1^{-10}$), (C) with the residue of ledger A1, CMP 96 Lemma 2.1, the [I] items (Lipschitz cut-offs O2, block-locality of Q, κ_1 independent of M), and O3 (Z_0-compressed variant) and O4 (complex α'-space), both now closed in `SOURCES_WC_EXPGAUGE.md`

and SOURCES_WC_RESIDUE.md (verified), O4 relative to CMP 99 Thm 3.4 [V] and the complex walk form of G [R]. The construction is now written out in full as Theorem 5.1 (Section 5.8); its statement and constants supersede the ones of this paragraph.

SOURCES_WG.md: **(W_G), walk expansions of G and G_1.** The file writes out Bałaban's walk expansions (W_G) of the final propagators G and G_1 of CMP 99 Sect. E: for regular U, $M \geq M_*(d,L)$ and $M\alpha_0 \leq a_*(d,L)$, $G = \sum_\omega T_\omega$ is a walk expansion with bounded branching and rate $(1-\theta)\frac{1}{2}\delta_0$, and G_1 has a tree expansion with rate $(1-\theta)\frac{1}{2}\varepsilon$, set by the walk expansion of $(QGQ^*)^{-1}$. The literal (3.108) form, with a factor M^{-1} per step also at the Δ'_π junctions, needs in addition $\alpha_0 \leq c'M^{-11}$ (WG-1; the M_1 form is in Theorem 5.2). The identity $G^{-1} = G_0^{-1} - \Delta'_\pi$ and the Neumann series (3.130) $G = \sum_n G_0(\Delta'_\pi G_0)^n$ are expanded with the printed walk expansions of G_0, G' and $(Q'G'^2Q'^*)^{-1}$ (CMP 99 Thms 3.10, 3.7, 3.9), one local J-insertion of size $O(M\alpha_0)$ per Δ'_π, and a concatenation lemma that splits the exponent at each junction and sums the junction points by CMP 96 Lemma 2.1. For G_1 the same scheme runs on $G_1 = \sum_n G(\Delta_\pi^{(2)}G)^n$ with $\Delta_\pi^{(2)} = (I - DRG'D^*)\Delta^{(2)}(I - DG'RD^*)$, whose side branches use the walk expansion A1-W of $(QGQ^*)^{-1}$, and the order $G_0 \to G \to (QGQ^*)^{-1} \to H \to G_1$ has no loop. Verifier: SOURCES_WG_VERIFY.md, [VERIFIED REDUCTION]: (W_G) for G reduces to the printed proof sketches of CMP 99 Thms 3.7, 3.9, 3.10, the Neumann series, CMP 96 Lemma 2.1 and [I] items, and the G_1 clause to the same plus A1-W (Bałaban himself writes "assuming (3.132)" on p. 426); corrections: WG-1 needs $\alpha_0 \leq c'M^{-11}$, not M^{-8}, because a Δ'_π insertion has up to $N_V = 9$ walk blocks, and "after adjusting a definition of δ_0" is on p. 427. Open inputs: WG-1 (admissible in Bałaban's constant order, not implied by $M\alpha_0 \leq a_0$); A1-W for the G_1 clause, with (C) conditional on the torus-side hypothesis and two conventions; the [I] items (Q-locality, two-sided derivative bounds, $C^{(2)}$-locality, block counts); all under ledger A1w (O3 and O4 are closed, see SOURCES_A1_WALK.md above). The write-out WG_FULL.md (Theorem 5.2, Section 5.8) supersedes the statement and constants of this paragraph: the walks live on the M_1-cover, and WG-1 reads $\alpha_0 \leq c'M^{-1}M_1^{-10}$ for G and $\alpha_0 \leq c''M^{-1}M_1^{-12}$ for G_1.

SOURCES_A2_R2.md: **ledger A2, R_2 and R_3 of CMP 116 (2.16).** The precision difference $R_2 := C_0^{-1} - \mathrm{Re}\, C^{(k)}(Z_0,\sigma)^{-1}$ obeys $|R_2(b,b')| \leq (O(1)e^{-\delta_0 M/3} + O(\alpha_0+\alpha_1))e^{-\rho|b-b'|}$ for $|\sigma(\Delta)| \leq e^{\kappa_1}$, with the printed rate $\rho = \frac{1}{2}\delta_0$ when the decay rate δ_A of $C^*\Delta_k C$ exceeds $\frac{1}{2}\delta_0$, and a smaller rate $\rho < \delta_A$ otherwise, which (2.21) and LOC tolerate. R_3 holds in two weaker forms (R3a), (R3b), sufficient for its only printed use (2.21), and LOC's input (E) follows directly at $\sigma = 0$ with $\varepsilon_M = O(e^{-\delta_0 M/2})$. A Neumann series around the compression $C(1)^{-1} = Z_0 A Z_0$, $A = C^*\Delta_k C$ (case (a), (2.5)), treats $W_\sigma = C(1) - C(\sigma)$, the walks that meet σ_0, as the perturbation: each such walk makes a round trip to distance $> \frac{2}{3}M$ from Z_0, which gives $O(e^{-\delta_0 M/3})$ once the weight $e^{16\kappa_1}$ is absorbed for $M \geq 96\kappa_1/\delta_0$. The $O(\alpha_0+\alpha_1)$ part comes from a one-variable Cauchy estimate on the analyticity space, and the verifier's repair V7 expands around the real $A_Z(U)$, so that decay of A is needed at real U only. Verifier: SOURCES_A2_R2_VERIFY.md, [VERIFIED REDUCTION] (with SOURCES_S2_G1_V7.md and its verifier); corrections: the walk bounds (W) are CMP 99 (3.107)–(3.108), not CMP 109; Lemma N and the Cauchy disc needed decay of A at complex U, repaired by V7; the Cauchy radius must respect the plaquette condition; and "not needed: A1" is wrong, since (W) for $C^{(k)}(Z_0)$ contains the unprinted walk expansion of $(QGQ^*)^{-1}$ (A1w). Open inputs: (W^c), the walk bounds on the complex α'-space, is no longer open: it is proved directly in the exponential gauge, real and complex, with no use of the Thm 3.15 transfer (SOURCES_WC_EXPGAUGE.md, SOURCES_WC_RESIDUE.md, both verified; ledger A2 regraded to a verified reduction); it rests on the G_1 tree expansion (CMP 99 Thm 3.10, printed sketch), the conditional (C) of A1-W, R2 and R-Δ2 of A3, CMP 109 p. 254 [R], CMP

99 Thm 3.4 [V], and [I] items; it is now written out in full as Theorem 5.3 (Section 5.8); the decay of $C^*\Delta_k C$ at real U (A7); the boundary-contact reading of "intersect" [I-forced]; $\mathrm{dist}(Z_0, \Lambda_k^c) > RM$ [I]; all under ledger A2.

`SOURCES_A3_GAMMA0.md`: **ledger A3, the Kato route.** The lattice Kato (diamagnetic) route to $C^*\Delta_k C \geq \gamma_0$ is refuted, and A3 is reduced instead to Bałaban's two-step mechanism at $U = 1$ made gauge covariant, with $\gamma_0 \geq (48d^2)^{-1}L^{-d-1}$ independent of k, volume and M. This is the verifier's sharpened constant; for $L \geq 10$ it carries the factor Lm_L found later in `SOURCES_A3_GAMMA0_FULL.md`. Step (i) bounds the minimal-extension energy below by the unit-lattice curl energy, $\langle B, \Delta_k B\rangle \geq \|\partial_1 B\|^2$ at $U = 1$ by exact lattice Stokes and Cauchy–Schwarz, and step (ii) is CMP 96 Lemma 2.4, a unit-lattice Poincaré inequality in axial gauge. Both steps are made covariant in a local regular gauge on each region, with the gauge-fixing part removed through (3.115) and relative errors $O(M\alpha_0 + L^2\alpha_0)c(d, L)$ for $M \geq cL$. Verifier: `SOURCES_A3_GAMMA0_VERIFY.md`, [VERIFIED REDUCTION] at sketch level, not a proof; corrections: the example $A = d\delta_x$ fails once $\|RD^*A\|^2$ is included and is replaced by an unfixed mode $\lambda \in \mathrm{Ran}(\Delta^{-2}Q'^*)$ (0 against 4.62), so the Kato route stays refuted; $c_* \leq 1$, not 4, so the free constant $(12d^2)^{-1}L^{-d-1}$ stands and the bound sharpens from $(192d^2)^{-1}L^{-d-1}$ to $(48d^2)^{-1}L^{-d-1}$; R1 must be CMP 99 Thms 3.1, 3.3, 3.11, 3.12. Open inputs: R1 with a uniform bound on the oblique projection $\kappa_\pi = \|I - DG'RD^*\|^2$; R2, the k-step averaging comparison (an A18 piece); the small items R3–R5; the σ-localized clause [I]; all under ledger A3, and taken up in `SOURCES_A3_GAMMA0_FULL.md` and `SOURCES_A3_SIGMA.md`.

`SOURCES_A3_GAMMA0_FULL.md`: **ledger A3, full proof.** Theorem A3: for U obeying (3.35)–(3.36) with $M\alpha_0$ small, $\langle \tilde{B}, C^*\Delta_k C\tilde{B}\rangle \geq \gamma_0\|\tilde{B}\|^2$ with $\gamma_0 = m_L/(48d^2L^d)$, $m_L = \min(2(1 - \cos(\pi/L)), 1/L)$, uniform in k, volume and regular U; this equals $(48d^2)^{-1}L^{-d-1}$ for $L \leq 9$ and is 0.7555 times it at $L = 13$. The proof step inside CMP 96 (2.126)–(2.127) is false for $L \geq 10$ (a lost factor L), while the statement of Lemma 2.4 is not refuted and stands with the corrected constant. The failure of (2.127) for every $L \geq 10$ is checked in Lean (`SlabCounterexample.lean`, Appendix A). Lemma 5 shows that the G_1-minimizer already satisfies $RD^*A_* = 0$, because $A \mapsto A - DG'RD^*A$ keeps QA and lowers $\langle A, G_1^{-1}A\rangle$ by exactly $\|RD^*A\|^2$, so the gauge-fixing part drops out and no bound on κ_π is needed; Lemma 5 is checked in Lean from (F1), (F3), (F4) as hypotheses (`GaugeMin.lean`). The two steps of `SOURCES_A3_GAMMA0.md` are then written out in full (step (i) with $c_* = 1$ and a covariant relative error $2(1 + \delta^{-1})\varepsilon_2^2$, step (ii) as CMP 96 Lemma 2.4 reproved with m_L in place of $1/L$ and made covariant in one regular gauge per corridor), and the assembly gives $\gamma_0 = \gamma_L/4$ with $\gamma_L = m_L/(12d^2L^d)$. Verifier: `SOURCES_A3_GAMMA0_FULL_VERIFY.md`, [PROVED+VERIFIED, CONDITIONAL ON R-Δ2 AND R3–R5], with CMP 99 Thms 3.11–3.12 published and (H_Q) [PROVED+VERIFIED]; the slab claim against CMP 96 was checked first and confirmed (lowest eigenvalue $L^{-d}\min(2(1 - \cos(\pi/L)), 1/L)$); corrections: (2.127) fails from $L = 10$ with free block bonds and from $L = 15$ in axial gauge ($d = 2$), not only from $L = 30$; fact (E) needs a factor $\sqrt{N}$; the reason for R-Δ2 must be the form (3.136) plus a block-average bound on $C^{(2)}$. Open inputs: R-Δ2 [R]; the Schur step of R1 from CMP 99 Thm 3.12, not written out; R3, R4, R5, R-Δ' and R2$'$; the σ-localized clause, reduced in `SOURCES_A3_SIGMA.md`; all under ledger A3.

`SOURCES_A3_SIGMA.md`: **ledger A3, σ-localized clause.** For real $s \in [0, 1]^{\sigma_0}$ the s-localized covariance $C_s = C^{(k)}(Z_0, s)$ obeys $\|C_s - C_1\| \leq \eta_M = S_d B_W e^{-\frac{1}{3}\delta_0 M}$ and, with no symmetry assumption, $\langle v, C_s^{-1}v\rangle \geq (\gamma_0 - 2\gamma_1^2\eta_M)\|v\|^2 \geq \frac{1}{2}\gamma_0\|v\|^2$ for $\eta_M \leq \gamma_0/(4\gamma_1^2)$, uniformly in s, Z, Z_0, k, volume and real regular U, with no α-condition. The resolvent bound $\|C_1^{-1} - C_s^{-1}\| \leq 2\gamma_1^2\eta_M$ also gives LOC's input (E) in norm at real points without (2.16). The walk sum of CMP 116 p. 3

expands the covariance, $C_s = \sum_\omega s^\omega T_\omega$, so $C_s - C_1 = -\sum_{\omega \in W_\sigma}(1 - s^\omega)T_\omega$ is dominated termwise by one s-free absolute majorant over the walks that meet σ_0, each of which reaches distance $> \frac{2}{3}M$ from Z_0. A Schur bound gives η_M, and the resolvent identity with $\gamma_0 \le Z_0 A Z_0 \le \gamma_1$ turns it into the lower bound on the precision. Verifier: `SOURCES_A3_SIGMA_VERIFY.md`, [VERIFIED REDUCTION] to (W) and (N); corrections: (W) is CMP 99 (3.107)–(3.108) with the Cor. 3.8 counting, not CMP 109; the bound without symmetry is the verifier's strengthening of the author's bound $\frac{2}{3}\gamma_0$ under (S_s), so (S_s) is optional; the uniformity of B_W in Z_0 uses part of the exponential decay (constants only). Open inputs: (W), Bałaban's assertion (p. 13) that the [13] expansion of $C^{(k)}(Z_0)$ with the large domain X_0 obeys (3.108) uniformly in Z_0 (A1w enters here [I]; in the exponential gauge this form is Theorem 5.3(ii), relative to its inputs); the interior reading of "intersect" for the zero-length walk, forced by (2.16); (N) $\gamma_0 \le A \le \gamma_1$, with γ_0 from ledger A3 and γ_1 through A7; complex σ stays with A2.

`SOURCES_A4_D189.md`**: ledger A4, CMP 102 (189).** CMP 102 (189) bounds the two-point Hessian kernel of the nonlinear action part V at fixed A', and the file derives $|V''(A'; b, b')| \le c_V \varepsilon_3 e^{-\frac{1}{2}(1-\alpha)^5 \delta_0 d(b,b')}$, rate $\ge \frac{1}{4}\delta_0$ for $\alpha \le 0.129$, volume-free and on the complex domain (77); (190) follows through the Neumann series (188) at rate $\frac{1}{8}\delta_0$ ($\alpha \le 0.159$). Every operator that CMP 102 inverts has the form I plus a small decaying kernel, so a weighted Schur Neumann series (CMP 96 Lemma 2.1) suffices, without Demko–Moss–Smith and without a Hessian lower bound. Lemma D2 shows that the second derivative of the linearizing map solves $(I+\mathfrak{N})\mathfrak{D}_2 = C''[(I-H\mathfrak{D})\cdot, (I-H\mathfrak{D})\cdot]$ with the operator of (68), with two-sided decay at $\frac{1}{2}(1-\alpha)^4\delta_0$, and composition then gives V''. Verifier: `SOURCES_A1_A4_VERIFY.md` (Part B), [VERIFIED REDUCTION] to A1 (kernel-decay form), published CMP 102 statements, A18, and one bookkeeping item; corrected: the rates are relabelled ($\frac{1}{2}(1-\alpha)^4\delta_0$ for $\mathfrak{D}_2$, $\frac{1}{2}(1-\alpha)^5\delta_0$ for V''), the name $\mathfrak{D}_2$ clashes with Bałaban's (first derivative minus linear term), V'' uses (3.132) directly through (87)–(88) and the decay of $\tilde{G}$ depends on A1 through (3.137) and (3.133), and the CMP 102 rates are fractions of the common $\delta_0 \le \delta_1$. Open inputs: A1 (itself a verified reduction), A18, the second-derivative bookkeeping of the V_0 terms (90)–(97), and the ∇, Hölder and Δ versions of (190) [I].

5.8 Step 8′: the three write-outs of the propagator expansions

Three files write out in full the constructions that the paragraphs on `SOURCES_A1_WALK.md`, `SOURCES_WG.md`, and `SOURCES_A2_R2.md` reduce to: `A1_WALK_FULL.md`, `WG_FULL.md`, and `WC_FULL.md`. Each was checked by an independent verifier file, fixed, and regraded. Each is paper-grade relative to its tagged inputs. None of them removes an input: they replace "reduced to a printed sketch" by a written proof that names its inputs. So the tag stays [VERIFIED REDUCTION]. The inputs common to all three are the lower bound (C), Bałaban's printed proof sketches of CMP 99 Thms 3.7, 3.9, and 3.10, and a short list of geometry and convention items [I]. Notation: M_1 is the cube size of the walks (CMP 116 p. 3), M is the regularity parameter of CMP 99 (3.36) and the cube size of the s-rule, and $R_1 M_1 \le M$. A walk expansion is a sum $T = \sum_\omega T_\omega$ over walks of domains, with (L1) each factor kernel vanishing within $\frac{1}{3}M_1$ of the boundary of its domain (the band of CMP 116 p. 3), (L2) U-dependence of each factor only on $\tilde{X}^5$, (B1) a per-walk bound $B\rho^{|\omega|}e^{-\delta d(\omega,y,y')}$ with the path distance (3.93), and (B2) bounded branching.

Theorem 5.1 (Theorem A1-W, `A1_WALK_FULL.md` §0.5 [VERIFIED REDUCTION])**.** *Let* $\hat{N} = S^{-1/2}QGQ^*S^{-1/2}$. *Assume (H1)* $\hat{N}$ *has a walk expansion with (L1), (L2),* $|\hat{n}_\omega(y,y')| \le b_G \rho_G^{|\omega|} m(y,y') e^{-\delta_G d(\omega,y,y')}$, $\rho_G = c_G/M_1$, *at most* $C_{\mathcal{D}}$ *domains meeting a set of diameter* $\le D_+ M_1$, *and* $\operatorname{diam}\tilde{X}^5 \le c_X M_1$; *(H2) the lower bound (C),* $\operatorname{Re}\langle u, \hat{N}u\rangle \ge c_-\|u\|^2$, *uniform; (H3) the cube*

geometry of Bałaban's cover; (H4) the constant order $(d, L, a, a_0) \to \theta \to c_\pm, \delta_G, \Sigma, b_G, c_G, C_{\mathcal{D}}, c_X \to \varepsilon \to M_1 \to M \to \alpha_0$*, with WG-1 in the* M_1 *form* $\alpha_0 \le c' M^{-1} M_1^{-10}$ *for* G *and* $\alpha_0 \le c'' M^{-1} M_1^{-12}$ *for* G_1*. Put* $c_+ = C_{\mathcal{D}} b_G$ *and* $\varepsilon = \min\{\frac{1}{2}\delta_G,\ c_-/(8c_+\Sigma(\frac{1}{2}\delta_G))\}$*. Then for* $M_1 \ge M_W(d, L, a, a_0)$*: (i)* $\hat{N}^{-1} = \sum_\omega T_\omega$ *is a walk expansion; (ii)* $|T_\omega(y, y')| \le (4/c_-)(c_1\rho)^{|\omega|} m(y, y') e^{-\frac{1}{2}\varepsilon d(\omega, y, y')}$ *with* $\rho \le K_W/M_1$*; (iii)* $\sum_\omega |T_\omega(y, y')| \le (8C_{\mathcal{D}}/c_-) m(y, y') e^{-\frac{1}{2}\varepsilon d(y, y')}$*; (iv)* $\|\hat{N}^{-1}\|_{\frac{1}{2}\varepsilon} \le 8/c_-$*. All constants are uniform in* U*,* j*,* k*,* η*,* $\{\Omega_j\}$*, and the volume.*

The file adds two variants. The Z_0-compressed variant (§7): for an operator A with a walk expansion and $\operatorname{Re} A \ge \gamma$, the inverse of $A_Z = 1_{Z_0} A 1_{Z_0}$ has a walk expansion with prefactor $4/\gamma$, and for $q = M/M_1 > 10$ and $q \ge 3(7 + c_X)$ it regroups into the CMP 116 p. 13 form (one domain X_0 with $\tilde{X}_0^5 = \tilde{Z}_0$, all other domains at distance $> \frac{2}{3}M$ from Z_0, the band kept, and a merged-slot cost $8C_W/\gamma$ per walk); for $(x + A_Z)^{-1}$, $x \ge 0$, the prefactor is $4/(\gamma + x)$. The complex variant (§8): Lemmas CP and AW and a four-level chain of complex domains give the expansion on the α'-space. The chain constants $\Gamma_3 = 2C_{\mathcal{D}} B_1'''\Sigma(\delta_1''')$ and $\Gamma_1'' = 2C_{\mathcal{D}} B_A''\Sigma(\delta_A'')$ carry the factor $C_{\mathcal{D}}$, so k_1, k_2 are smaller by at most $C_{\mathcal{D}}$. The bound $\sum_\square \varphi_\square^2 \ge 1$ on Z_0, needed in §7.3, is proved by the verifier (tent family on a grid of spacing $M_1 L^k$). The tree form of (H1), which the G_1 clause uses, is stated explicitly in `A1_WALK_FULL.md` 6.9. Verifier: `A1_WALK_FULL_VERIFY.md`, then a regrade: paper-grade relative to its inputs IN1 ((W$_G$) = Theorem 5.2), IN3 ((C), conditional), IN8–IN12 [I] (cube geometry, the Lipschitz partition O2, Q-locality, the WG-1 condition, κ_1 independent of M), and IN18 [R] ((W$_G$) at complex points, from CMP 99 Thms 3.4 and 3.10).

Theorem 5.2 (Theorem WG, `WG_FULL.md` 0.7 [VERIFIED REDUCTION])**.** *Let* U *satisfy CMP 99 (3.35)–(3.36) and fix* $\theta \in (0, 1)$*. (G) (Theorem WG-G.) There are* $M_*(d, L, \theta)$ *and* $a_*(d, L, \theta) > 0$ *such that for* $M_1 \ge M_*$*,* $M\alpha_0 \le a_*$*, and RM with CMP 96 (2.59) at rate* $\theta\frac{1}{2}\delta_0$*,* $G = \sum_\lambda T_\lambda$ *is a walk expansion on the* M_1*-cover, with constants* $(B_0', \rho_*, (1-\theta)\frac{1}{2}\delta_0)$*,* $\rho_* = \max\{c_0 M_1^{-1}, (K_V M\alpha_0)^{1/10}\}$*; the bound holds also with one covariant derivative on either side. If in addition* $\alpha_0 \le c' M^{-1} M_1^{-10}$ *(WG-1), then* $\rho_* \le c_0 M_1^{-1}$*, the CMP 116 (1.7) form. (G1) (Theorem WG-G1.) Assume in addition Theorem 5.1 for* $(QGQ^*)^{-1}$*. There are* M_W *and* $a_{**} > 0$ *such that for* $M_1 \ge \max\{M_*', M_W\}$*,* $M\alpha_0 \le a_{**}$*, and RM* $\ge C(d, L, \theta, \varepsilon)$*,* $G_1 = \sum_\tau T_\tau$ *is a tree expansion with rate* $(1-\theta)\frac{1}{2}\varepsilon$*. The per-step form needs* $\alpha_0 \le c'' M^{-1} M_1^{-12}$*. All constants are uniform in* U *with (3.35)–(3.36).*

The proof expands $G = \sum_n G_0(-VG_0)^n$, $V = -\Delta_\pi'$, with the walk expansions of G_0, G', and $(Q'G'^2Q'^*)^{-1}$. Lemma N shows that the Neumann series gives the inverse in finite volume, so no separate existence of G is needed. V splits into 24 patterns with at most $N_V = 9$ walk blocks, and one local J-insertion of size $O(M\alpha_0)$. Lemma C concatenates walks: it splits the exponent at each junction and sums the junction point by CMP 96 Lemma 2.1, so the rate $(1-\theta)\frac{1}{2}\delta_0$ is lost once. The 10 junction steps of a V-insertion give the exponent of WG-1. To obtain the band (L1), every domain is enlarged by one layer of M_1-cubes; this changes the branching count to $C_{\mathcal{D}} = 5^d c(d) C_{\mathcal{D}}^{\text{in}}$ and c_X to $c_X + 2$, and adds $M_1 \ge 3c_{\text{loc}}$ to M_*. The output satisfies (H1) of Theorem 5.1. Inputs: CMP 99 Thm 3.10 and Thm 3.9 (printed proof sketches), Thm 3.7 with Cor. 3.8 (proof by reference to CMP 95 Prop. 1.2 and CMP 96 Prop. 2.2), CMP 96 Lemma 2.1 under (2.59), the bound $|J| \le O(1)M\alpha_0(L^j\eta)^{-3}$ [V]; for (G1) also Theorem 5.1, so (C); [I] items: the scale bookkeeping C0 (written out for the vector $\hat{J} = S^{3/2}J$), locality of Q, Q', D, and the parallel transports, the assignment of ι-domains, the locality of $C^{(2)}$, and the M_1-cover (asserted on CMP 116 p. 3). Verifier: `WG_FULL_VERIFY.md`, then a regrade: paper-grade relative to these inputs; the verifier found that every consumer uses the M_1 form, so the corrected WG-1 replaces the earlier $\alpha_0 \le c' M^{-11}$; the band enlargement passed the band regrade.

Theorem 5.3 (Theorem WC, `WC_FULL.md` 0.10 [VERIFIED REDUCTION]). *Fix $d \geq 2$ and odd $L \geq 3$. There are M_1, M with $M/M_1 > 3(6 + c_X)$ and $M/M_1 > 10$, a smallness α_*, a four-level chain of complex domains $\alpha' \subset \alpha'' \subset \alpha''' \subset$ the a_1-domain of CMP 99 Thm 3.4, and constants $\gamma_{\rm exp} = \frac{1}{4}\gamma_e(d, L) > 0$, $\varepsilon'' > 0$, $\rho > 0$, such that for every real regular U and every Z_0: (i) $\langle \tilde{B}, A_Z \tilde{B}\rangle \geq \gamma_{\rm exp}\|\tilde{B}\|^2$ on the reduced space of Z_0; (ii) $C^{(k)}(Z_0) = \sum_\omega T_\omega$ with (L1), (L2), $|T_\omega(b, b')| \leq (4/\gamma_{\rm exp})(c_1\rho)^{|\omega|} e^{-\frac{1}{2}\varepsilon d(\omega,b,b')}$, absolutely convergent, and after regrouping in the CMP 116 p. 13 form; (iii) at every U_c in the α'-space, $\mathrm{Re}\langle u, A_Z(U_c)u\rangle \geq \frac{1}{2}\gamma_{\rm exp}\|u\|^2$, $C^{(k)}(Z_0; \cdot)$ is analytic, and (ii) holds with $4/\gamma_{\rm exp} \to 8/\gamma_{\rm exp}$ and $\varepsilon \to \varepsilon''$; (iv) (i)–(iii) hold for $(x + A_Z)^{-1}$, $x \geq 0$, uniformly in x, with the per-walk prefactor $4/(\gamma_{\rm exp} + x)$ (real) and $8/(\gamma_{\rm exp} + x)$ (complex), so the $x^{-1/2}$ integral for $(C^{(k)})^{1/2}$ converges walk by walk. The constants are uniform in k, η, Z_0, $M \geq M_1$, and the background. The proof does not use CMP 99 Thm 3.15 or any transfer from the axial gauge.*

The lower bound comes from Lemma E, $S_y \leq K_b(P_y + G_y)$ on each block, with $K_b = 8dL^d(1 + 16dc_{\mathcal{G}}) \leq 64d^3L^{2d+1}$ under $16d^3L^d c_{\mathcal{G}}\tau_p^2 \leq 1$, $\tau_p = 16\sqrt{N}c_4 M\alpha_0$; then $\gamma_e^{-1} = K_b(1+6d/m_L)+6/m_L$ and $\gamma_{\rm exp} = \frac{1}{2}\min(\frac{1}{2}, c_G)\gamma_e$ with $c_G = 1$. For Federbush's mean (CMP 109 (0.10)) the exact weights give $c_M \leq 4\sqrt{N}$, and $\kappa \leq \sqrt{2}$ for $w \leq 0.054/\sqrt{N}$. The walk expansion is the Z_0-compressed variant of Theorem 5.1 applied to the tree expansion of $A_{\rm exp}$, which rests on Theorem 5.2(G1). The Combes–Thomas Schur sums carry the factor of $|e^x - 1| \leq |x|e^{|x|}$: $\Sigma(\frac{1}{2}\delta) \leq (1+4/(e\delta))S(\frac{1}{4}\delta)$, which changes only ε, ε^a, and ε''. Inputs: [R] Theorem 5.2(G1) (printed sketches), the complex walk form of G (CMP 99 Thms 3.4 and 3.10), the inputs of ledger A3, and CMP 109 p. 254; conditional: (C) and $c_{-,1} \geq c_-(1 - O(M\alpha_0))$; [I] items I1–I9 of `WC_FULL.md` §9 (among them the term match of CMP 109 (2.11) with CMP 99 (3.156), the locality of $\tilde{C}^{(2)}$ and Q, the identification $\tilde{Q} = Q_1(U_1)$, $\mathrm{dist}(Z_0, \Lambda_k^c) > RM$, and the tilde convention). Verifier: `WC_FULL_VERIFY.md` (needs fixes, none structural), then two regrades: after the fixes and after the band fix, paper-grade relative to these inputs.

5.9 Step 8″: round 5, the remaining Group A rows and Lemma G3

Round 5 treats the Group A rows that had neither a verified proof nor a verified reduction: A7, A8, A10, A11, and A15. It also proves Lemma G3, the residue N2 of the reduction of A5. Each result below has its own project file and an independent verifier file. Each holds relative to the inputs that it names. None of the rows leaves the ledger, because those inputs are not proved here. The tag of each item therefore stays [VERIFIED REDUCTION] or [PROVED RELATIVE TO LISTED INPUTS]; nothing in this step makes Theorem A or Theorem B unconditional.

The operator of A7. Bałaban's C in $C^*\Delta_k C$ is the constraint map $I - hL\tilde{Q}$, not a square root of the covariance. The operator $C^*\Delta_k C$ is the *precision*: CMP 116 (2.7) reads $(C^{(k)})^{1/2} = (C^*\Delta_k C)^{-1/2}$. So the lower bound $A_Z \geq \frac{1}{4}\gamma_e$ of Theorem 5.3(i) is the γ_0 side (row A3) and cannot give γ_1. The old reduction "A7 ⇐ A1 + A18" of `SOURCES_CDC.md` is superseded; A18 now enters through A3-R4 and the item I3 of `WC_FULL.md`.

Theorem 5.4 (A7, `SOURCES_A7_CDC.md` [PROVED RELATIVE TO LISTED INPUTS; VERIFIED]). *Let U be real and regular, $s \equiv 1$, and let Λ be as in `WC_FULL.md` 0.8. (N) (Lemma N.) $\|C^*\Delta_k C\| \leq \gamma_1^N$ with $\gamma_1^N = \|C\|^2 \max\{K_{\rm ext,1} - a + c_J + 2c_{\mathcal{G}},\ a + c_J\}$, uniform in k, the volume, and U. (D) (Theorem D.) $|(C^*\Delta_k C)(b, b')| \leq 2B_A e^{-\delta_A|b-b'|}$ with $\delta_A = (1-\theta)\min\{\frac{1}{2}\varepsilon, \delta_1\}$, and the series (2.7) for $(C^{(k)})^{1/2}$ converges walk by walk for $\gamma_1 := 4B_A S(\theta'\delta_A)$. Both bounds hold for real $s \in [0, 1]$ with the same constants.*

Inputs of (N): C2 of `WC_FULL.md` (the bound (C) for G_1, conditional on the torus-side hypothesis and two conventions, plus one [I] step for G_1), A3-R4, I1, and I3. Inputs of (D): W_{G_1} [R] (Theorem 5.2(G1)), Theorem 5.1 [C], I2, and I9. The clause for complex s with $|s| \le e^{\kappa_1}$ and complex U on $\mathcal{U}(\alpha'')$ is at proof-sketch level and needs M_1 large after κ_1. The value $\gamma_1^N \approx 243(K_E + 648)$ at $d = 4$, $L = 3$ is [I]. Verifier: `SOURCES_A7_CDC_VERIFY.md`; claims (1), (3), (4), (5) confirmed, claim (2) confirmed with the input list corrected (the two-sided route of §2.3 is an alternative to Theorem L).

The positivity device of A8. Only $H(K) = \operatorname{sym} \operatorname{Re} K$ enters the modulus of the Gaussian measure, $K = C^{(k)}(Z_0, \sigma)^{-1}$. Bałaban states that $C^{(k)}(Z_0, 0)$ is symmetric. That claim is not proved: the parametrix walk series is not reversal-symmetric, and the truncation need not keep symmetry. The claim is also not needed. The majorant uses $H_0 = \operatorname{sym} C_0^{-1} \ge \frac{3}{4}\gamma_{\exp}$ in its place.

Theorem 5.5 (A8, `SOURCES_A8_POS.md` [PROVED IN CORRECTED FORM RELATIVE TO LISTED INPUTS; VERIFIED]). *(R) (Real case.) At $(U, 0)$ with $s \equiv 1$, $K = A_Z$ is real symmetric (Lemma S) with $\gamma_{\exp} \le A_Z \le \Gamma_1$, so $d\mu_{C^{(k)}(Z_0)}$ is a positive Gaussian measure. (D) (Lemma D.) For $|\sigma| \le e^{\kappa_1}$ the σ-truncation obeys $\|C^{(k)}(Z_0, \sigma) - A_Z^{-1}\| \le \eta_M = (16/\gamma_{\exp}) e^{2 \cdot 3^d \kappa_1} W e^{-\varepsilon'' M/8}$. (P) (Lemma P.) The complex Gaussian exists; its modulus is $r_\sigma \, d\mu_{\Gamma_\sigma}$ with $1/(2\Gamma_1'') \le \Gamma_\sigma \le 4/\gamma_{\exp}$, which is Bałaban's (2.15). (M) (Majorant.) With $H_0 = \operatorname{sym} C_0^{-1} \ge \frac{3}{4}\gamma_{\exp}$ the majorant (2.23)–(2.25) holds at a cost $\exp(O(\eta_M^2)|Z_0|)$ and an extra term $O(\eta_M^2)\|ZX\|^2$, absorbed in (2.25).*

Inputs: Theorem 5.3(i)–(iii) with their residue (row A2), and five readings I-a–I-e (quadratic forms versus symmetric operators; the σ-weighted walk sum by the CMP 116 p. 3 rule; three geometry facts; $W < \infty$ at rate $\varepsilon''/8$; "$\operatorname{Re} C^{-1}$" as the symmetric real part). The M-large conditions are in the order of constants (C24). Verifier: `SOURCES_A8_POS_VERIFY.md`; Lemmas S and P confirmed; Lemma D corrected (a tree-distance slip: the exponent is $e^{-\varepsilon'' M/8}$, not $e^{-\varepsilon'' M/4}$); the sign of the extra (2.24) term corrected (it is a cost, not a gain). `SOURCES_G3.md` reads R_2 of (2.16) as $H_0 - H_\sigma$.

The form A10′ of A10. Bałaban writes that the lower-order bracket terms are "bounded similarly as in (1.24)" (CMP 116 p. 8). For the raw order-n Taylor terms, $n \le 4$, this is false as a bound: the cube count $(6L)^4 (L^j \eta)^{-4}$ leaves a sum over j that diverges in k. The sentence holds for the terms after the transformations of CMP 109 Sect. 4.

Theorem 5.6 (A10′, `SOURCES_A10_LOWER.md` [PROVED IN CORRECTED FORM RELATIVE TO LISTED INPUTS; VERIFIED]). *After the transformations of CMP 109 Sect. 4: the order-0 term is removed by the $t_\square$-derivative; the order-1 term is zero by (4.14); the marginal parts of (4.34) and (5.43) cancel exactly against the counterterm, because β_j is defined as the coefficient β in (5.43) (p. 298). Every other term obeys the (1.24) form with the factor $8 B_0 C_1 e^{16\kappa_1} \alpha_2^{-1} g_k |B|$ and field power 5, or $4 + \beta_H$. The "other terms" reach (1.29): F with exponent $(1 - 2\delta)\kappa$; R and C by the same mechanism.*

Residue: R1 unprinted constants (finite, fixed before ε_1); R2 the R- and C-terms by mechanism, not written out; R3 Bałaban's Sect. 4–5 bounds as printed; R4 the tail constant of (4.5) [I]. The per-label exponent of Fact A in Theorem SM is the Hölder exponent β_H of CMP 109 (3.31), which Bałaban does not print (see the paragraph on β_H below). Transcription fix: (1.29) reads $O(M^q)$ and $(1/16)\kappa_1$. Verifier: `SOURCES_A10_LOWER_VERIFY.md`; claims (1), (2), (4), (6) confirmed; (3) corrected (a factor 2); (5) corrected (β_H is unprinted and needs its own symbol; Lemma HH's θ has no numerical value, so $\theta < 1 - L^{-\beta_H}$ holds by ε_1 small).

The free exponent β_H. The consumers of A10′ change as follows (correction blocks in `SOURCES_SM.md` and `SOURCES_HHIST.md`): Fact A has $\alpha_* = \beta_H$; (Eps) reads $2\varepsilon' = 2\theta_0(1 - L^{-\beta_H})^{-1}$; Lemma HH needs $\theta < 1 - L^{-\beta_H}$, with column sum $\leq C_\beta + 2C_\Pi\theta_0 S_0(1 - L^{-\beta_H})^{-1}$; and $\lambda_g = L^{-\beta_H/2} + \theta_g\gamma$ in §4.3 of `SOURCES_SM.md`. The assessment `SOURCES_GGAIN_BETAH.md` ([UNVERIFIED ASSESSMENT]) finds that every consumer needs only $\beta_H > 0$, chosen before ε_1 and γ. Each condition is an ε_1- or γ-smallness condition. So $\beta_H \in (0, 1)$, $\beta_H \leq \beta_0 < 1$, is a free parameter, first in the order of constants (ledger C27). The cost is the factors $(1 - L^{-\beta_H})^{-1}$, a larger $B_4(\beta_0)$, and K_T fixed before ε_1. At $L = 13$ the (G-gain) margin $1 - 13^{-\beta_H/2}$ is 0.120 at $\beta_H = 0.1$ and 0.685 at $\beta_H = 0.9$; the Lemma HH margin is 0.901 at $\beta_H = 0.9$. The reading of CMP 102 Thm 1 (9) "$\beta_0 = 1$" as a misprint is [I]. This β_0 is not the β_0 of CMP 119 p. 255.

Proposition 5.7 (A11, `SOURCES_A11_CONST.md` [CLOSED AS A VERIFIED LIST])**.** *The "other constants" of CMP 119 p. 256 are further scale-indexed numbers $X_j = f(g_j)$: $\alpha_{0,j}$, $\alpha_{1,j}$ ((2.28)), $g_j^{\kappa_0}$ ((2.31)), $\delta_k = (A_1/A_0)\varepsilon_k$ (p. 265), $p_1(g_k)$ ((3.43)), and $g_j(x)$ ((2.24)). Every similar inequality for them follows from (2.6)–(2.9). The three unprinted cross-scale steps ($R_k \leq LR_{k+1}$ on p. 268, the monotonicity of k_1 on p. 270, and $\varepsilon_{k+1} \leq (1 + \beta_0)\varepsilon_k$) also follow from (2.9). The side conditions S1–S11 are admissible and acyclic.*

The file reads 43 introduction points (pp. 255–280 as images, pp. 281–284 as OCR). Three entries join the order of constants: C28 ($M_1 < M_2 < M$, order $M_1 \to M_2 \to M$), C29 ($M_2 \geq 1/\delta$), and C30 ($A_1/A_0 \leq 1/(80B_3)$, A_1 after A_0). The residue $r > 1$ is closed by "$r \geq 2$" (CMP 119 p. 245). Non-blocking residue: the form of $p_1(g)$ inherited from CMP 122 I, CMP 109 restrictions outside the file, and an OCR slip on p. 278. Verifier: `SOURCES_A11_CONST_VERIFY.md`; claims (1), (4) confirmed; (2), (3) corrected without a change of conclusion (three list omissions, a $(1 + \beta_0)^2$ slip fixed by monotonicity, the order $M_1 \to M_2 \to M$ with $M > M_2$ added).

The whole-lattice coefficient (A15). $\beta_{j+1}(g_j)$ is defined by CMP 109 (1.20)–(1.22), p. 264: the mixed second moment $\sum_x \Pi_{\mu\nu}(x)x_\mu x_\nu$ of the vacuum-polarization tensor at $U = 1$, taken as the limit of periodic-torus kernels $T \nearrow \mathbb{Z}^d$. Bałaban states that this limit exists "by the localized representation (1.7)". (1.7) and (1.9) say only that each term depends on U restricted to X. They do not say that each term is the same function on every torus and on $\mathbb{Z}^d$. So the existence of the limit stays Bałaban's bare assertion unless one adds the hypothesis (Loc) below.

Proposition 5.8 (Prop. A15-N, `SOURCES_A15_WHOLE.md` [PROVED GIVEN (LOC), (BD), (H), (CNT); VERIFIED])**.** *Fix j, $g \in [0, \gamma]$, and $U = 1$. Assume (Loc): for every non-wrapping polymer X the torus term equals the $\mathbb{Z}^d$ term of its lift; (Bd): CMP 109 (1.18); (H): the linearized background map decays exponentially, at the same rate on the torus and on $\mathbb{Z}^d$; and (Cnt): the standard polymer count. Let N_j be the number of level-j cells per side and $m_j = \frac{1}{2}\min(\delta_H, \kappa - \kappa_0)$. Then the limit in (1.22) exists, and for the torus coefficient $\beta_j^{(N)}$ in moment form*

$$|\beta_j^{(N)}(g) - \beta_j(g)| \leq C_j N_j^6 e^{-m_j N_j/2},$$

with C_j independent of N. The Fourier form adds $O(N_j^{-2})$.

Exact equality $\beta_j^{(N)} = \beta_j$ cannot be derived in general. Image terms and wrapping polymers are real error sources; no explicit N with $\beta_j^{(N)} \neq \beta_j$ was computed, and Bałaban never claims coefficient equality. Equality holds term by term only for non-wrapping interior X under (Loc). The hypothesis (Loc) is not implied by (L2) of Theorem 5.2, which covers walk factors only. The case

$g = 0$ is in `SOURCES_T2_G2.md` (ledger B6). Verifier: `SOURCES_A15_WHOLE_VERIFY.md`; (1) confirmed (definition on the page image; the "(1.41)–(1.43)" citation on CMP 119 p. 253 is Bałaban's slip); (2) corrected (three unstated steps: the first-order term drops by gauge invariance, the linearized map includes the J channel, and m_j changes by a fixed factor; the power N^6 is loose); (3) corrected ("refuted" becomes "cannot be derived"); (4) corrected (the binding-set move below). Consequence for the binding sets: Theorem 2w uses only the definition of β_j (through B5, B6, and A5), so A15 leaves its set. Bałaban's assertion that the $\mathbb{Z}^d$ limit of (1.22) exists moves into row A5. A15 stays in the set of Theorem 2w ⇒ Theorem 1 and of Theorem B, through B8 and A13: (Loc) for $g > 0$, the reading of $\partial\Lambda_{k+1}$ on CMP 119 p. 278, and the R-term part, which cannot hold literally for terms built from merged R′ inputs.

The reduction of A5 and Lemma G3. `SOURCES_T2_H0.md` (verified: 47 confirmed, 10 corrected) proves Proposition H0-red: the use of A5 in Theorem 2w, item 7′, reduces to a residue N1–N9 in which no constant depends on j or on the past. The residue item N2 is Lemma G3. Bałaban's C^∞ and all-derivative claims are not used.

Theorem 5.9 (Lemma G3 = N2, `SOURCES_G3_N2.md` [PROVED RELATIVE TO LISTED INPUTS; VERIFIED])**.** *Assume Theorem 5.4 in its two-sided form, Theorem 5.5, Bałaban's chain CMP 116 (2.15)–(2.25) read as pointwise exponent bounds (including A2), and the conditions (Hα) and (Hγ) below. Then (Q) holds in the joint (X, B) form with*

$$c_- = (\gamma_1^N + \varepsilon_0)^{-1} \geq \tfrac{4}{5\gamma_1^N}, \qquad \Lambda = 2\Big(1 + \tfrac{4}{3\gamma_{\exp}} + \tfrac{16}{9\gamma_{\exp}^2} G_\Gamma^2\Big),$$

$\varepsilon_0 = 2\Gamma_1^2\eta_M^0 \leq \frac{1}{4}\gamma_{\exp}$, and Lemma G3 holds: the cut adds S_χ to S_0, and $C_\beta = C_\Pi(S_0^{\mathrm{GGAIN}} + S_\chi)$, uniform in j, the past, the volume, and regular U.

The one-bond-fixed form of (Q) is a corollary: fixing $B(b)$ integrates the compression exactly, with Schur complement $(\Gamma_{bb})^{-1}$ and no cost factor ($a = 0$). The proof uses no symmetry of A_Z and no Theorem L. It corrects `SOURCES_Q.md` 4.3: the B-block of Q_0^{-1} is $H_0^{-1} + H_0^{-1}\Gamma\Gamma^* H_0^{-1}$, so $G_\Gamma = \|\Gamma_k(Z_0, 0)\|$ enters Λ. Conditions. (Hα): $(\alpha_5 + G_\Gamma^2\delta_S)\Lambda_0 \leq \frac{1}{2}$, with $\delta_S = O(\eta_M^2)$ (ledger C25). (Hγ): $\frac{1}{20}\gamma_2\varepsilon_1^2/\gamma^2 \geq 4\kappa + 1$, and the condition $\frac{1}{20}x - 4M^4 e^{-x/20} \geq 0$ at $x = \gamma_2\varepsilon_1^2/g_k^2$ that the verifier added for (2.31) (ledger C26). Both are smallness conditions of Bałaban's type, one unit stronger than his own. The bound on G_Γ (`SOURCES_G3_GGAMMA.md`, [VERIFIED], all four claims confirmed): $\Gamma_k(Z_0, 0) = 1_{Z_0} A_{\exp}(0) 1_{Z_0^c} R(0)$ with $R(0) = (C^{(k)})^{1/2}(0)$, and the Schur test gives $G_\Gamma \leq \bar{G} := \Gamma_1 K_{1/2}$, $\Gamma_1 = 2B_A S(\delta_A)$, $K_{1/2} = 8\gamma_{\exp}^{-1/2} S(\frac{1}{2}\varepsilon) + 4\gamma_1^{-1/2} S((1-\theta')\delta_A)$. So G_Γ is bounded uniformly in M (verified), and in the volume, j, Z_0, and regular U. With this bound (Hα) is the single block-size threshold

$$M \geq \max\{M_\Gamma, M_{\alpha 5}\}, \quad M_\Gamma = \tfrac{4}{\varepsilon''} \log\big[16h^3\Gamma_1^4\bar{G}^2\bar{\Lambda}_0(16W/\gamma_{\exp})^2\big], \quad M_{\alpha 5} = \tfrac{3}{\delta_0}\log(8c_5\bar{\Lambda}_0),$$

$h = 4/(3\gamma_{\exp})$, together with Bałaban's smallness of $\alpha_0, \alpha_1, \alpha_4, \gamma_2$; κ_1 does not enter M_Γ, and there is no cycle. One reading residue stays [I]: the s-rule acts on Bałaban's resolvent expansion (3.185) of [13] (CMP 116 p. 13), and the bound identifies $R(0)$ with the walk sum of Theorem 5.3(iv) with the σ_0-walks deleted (the reading I-b of A8). Other inputs of Theorem 5.9: N3 (the Kotecký–Preiss derivative step, the margin (M_r) of A17 with C9, dominated differentiation), N6 (Bałaban's sup induction; past-uniformity of E_0 is reduced, not proved), N9 (A15, definition only), and the reading that the pointwise bounds hold on the closed small-field set. Verifier: `SOURCES_G3_N2_VERIFY.md`; (1), (2), (4) confirmed; (3) corrected (the $e^{|P|}$ payment comes from the fifth $e^{-(1/10)x|P|}$ factor via (Hγ), which adds the second γ condition); 2000 random numerical instances pass. Consequence:

the use of A5 in Theorem 2w is a [VERIFIED REDUCTION] with residue N1–N9, and N2 is proved relative to the inputs above. The small-field "(Q)/c_- open" of ledger B13 is discharged. A5 stays a row: N5 (continuity at $g = 0$, [R]), N6, N8 (readings [I]), and N9 are not proved here, and N7 is B7.

5.10 Step 8‴: round 6, locality, the boundary, the 2D gap, and the order of constants

Round 6 treats the Bałaban-level residue of row A15, the R-term row B8, and the open 2D gap of row B4. It also fixes one acyclic order of constants. Each result has its own project file and an independent verifier file. The exception is the check file `SOURCES_T2_VOL.md` for (Vol), which has no verifier file. Each result holds relative to the inputs that it names. The tags stay [VERIFIED REDUCTION] or [PROVED RELATIVE TO LISTED INPUTS]. Nothing in this step makes Theorem A or Theorem B unconditional.

Locality of torus activities. Work at level k on $T_N = \mathbb{Z}^4/N\mathbb{Z}^4$ in level-k units, with projection π. A set is *liftable* if it lies in $\pi(B)$ for a box $B \subset \mathbb{Z}^4$ of side $< N/2$. A domain X is *non-wrapping* if its collar $X^\sharp$ (X plus one layer of width c_k, the largest cube side at level k) is liftable; $\operatorname{diam}_\infty X \le N/2 - 3c_k$ suffices. $\hat{X}$, $\hat{U}$, $\hat{J}$ denote lifts.

Theorem 5.10 (Theorem LOC, `SOURCES_A15_LOC.md` §7.5 [PROVED RELATIVE TO LISTED INPUTS; VERIFIED])**.** *For every k, every coupling history in Bałaban's range, and every non-wrapping $X \in \mathcal{D}_{k+1}$,*

$$E^{(k+1)}_{T_N}(X, g, U, J) = E^{(k+1)}_{\mathbb{Z}^4}(\hat{X}, g, \hat{U}, \hat{J}) \quad \textit{for all } (U, J) \in \mathcal{U}^c_{k+1}(X, \alpha_0, \alpha_1).$$

There is no error term.

Method. The mechanism is the s-rule of CMP 116 p. 3. At $s = 0$ off Y_0, every walk-expanded operator is a finite walk sum inside Y_0, and the background equations (1.3)–(1.4) decouple by components. A bijection between the surviving torus walks and $\mathbb{Z}^4$ walks gives equal kernels (Lemma W). Uniqueness gives equal background fields (Lemma N). The cluster assembly and the subtractions are cube geometry (Lemmas A, S). The image terms of a periodized kernel sit in winding walks, so they enter only wrapping activities. The induction on k uses that β_{k+1} is the same $\mathbb{Z}^d$ number in both volumes; a torus coefficient would break exactness at the next step. Inputs: (FI) each walk factor is given by the same local formula on T_N and $\mathbb{Z}^4$ (supported by CMP 99 Sect. C); (U) uniqueness of the solutions of (1.3)–(1.4) (printed in CMP 102); and the Group C rows C32–C36. These are (Cube): $U_j(\square_0, \cdot)$ of [14] uses only data on $\square_0$ [I]; (Cases): the other term classes are localized "in exactly the same way" as the typical term (CMP 116 p. 2) [R]; (Conv): cube sides divide N, the partitions are π-images of the $\mathbb{Z}^4$ partitions, and $M = KR_0M_0$ with K a power of L; (Det′): a domain-assignment rule for the closed-walk terms of the determinant expansion of CMP 102; and the reading of "H_j with free boundary conditions" (CMP 109 p. 290) as the $\mathbb{Z}^4$ operator [I]. Verifier: `SOURCES_A15_LOC_VERIFY.md` (page images); confirmed with minor corrections. (i) Lemma G(b) had the split of CMP 116 (2.7) reversed: for large x the resolvent is a power series in $C^*\Delta_k C$, and the integral with $\tilde{G}_3(x)$ covers $x \in [0, \gamma_1]$; the conclusion is unchanged. (ii) The determinant expansion of CMP 102 does not follow the s-rule. It is a trace of walk expansions, and a trace splits exactly by walk; this needs the convention (Det′). (iii) The object table omitted two block-local terms (a Jacobian and $\log \sigma$); they are intrinsic by the same argument. (FI) and (U) are supported on the pages. Consequences. With Theorem 5.10, Proposition 5.8 holds relative to (Bd), (H), (Cnt), and the inputs above; (Loc) is no longer a hypothesis. Bałaban's assertion that

the $\mathbb{Z}^d$ limit of CMP 109 (1.22) exists at $g > 0$ follows from Proposition 5.8. So row A5 has no bare assertion left.

The boundary of the small-field region.

Proposition 5.11 (Proposition BDRY, `SOURCES_A15_BDRY.md` [PROVED RELATIVE TO LISTED INPUTS; VERIFIED WITH CORRECTIONS]). *(f) The small-field terms $E_0^{(k+1)}(\Lambda_{k+1}, X, z)$ of CMP 119 p. 278 with $X \subset \Lambda_{k+1}$, also those with X next to $\partial\Lambda_{k+1}$, equal the whole-lattice terms exactly. (Thr) Let N_k be the number of level-k cells per side of the torus. If $N_k > 22M$, every wrapping or image term of the coefficient analysis at step k lies in Bałaban's irrelevant class $X \not\subset \tilde{\square}^2$ (CMP 109 (4.36), CMP 119 (3.48)), and the torus correction to β_j is at most $\varepsilon(M) = C_j(22M)^6 e^{-11 m_j M}$, valid for $m_j M \geq 1$.*

Method. On the image of CMP 119 (1.20), $S_1 \cap \Lambda_1 = \emptyset$. The change of variables (1.22) acts only on S_1^* bonds, so H, the Jacobians, and $V(S)$ reach only terms that meet Λ^c, and these go to the large-field part. Step $k+1$ (p. 279) keeps every $X \subset \Lambda_{k+1}$. Lemma W_Λ: with $s = 0$ off $Y_0 \subset X \subset \Lambda$, every surviving walk of $C(\Lambda)$ lies inside Y_0 and does not see $\partial\Lambda$. Inputs: the inputs of Theorem 5.10; C37 (Enl): the s-rule localization domain of a walk is $\bigcup X_i^5$ as a whole-lattice set, not $\bigcup X_i^5 \cap \Lambda$; ($\mathrm{L2}_\Lambda$) through (L2) of Theorem 5.1; (Dom); (Conv_Λ) [I]; and the nesting of the Λ_j [I]. Verifier: `SOURCES_A15_BDRY_VERIFY.md` (page images). (1) Corrected: Bałaban prints neither form of the walk domains, so (Enl) is a convention, admissible. Under the Λ-restricted convention, (f) holds with a margin of at most one M-cube layer, also admissible. $C^*\Delta_k C$ is walk-expanded, not finite range (CMP 116 (2.7), CMP 119 p. 275); the s-truncation of the Δ_k walks follows the same convention. (2) Corrected: see the B8 paragraph below. (3) Corrected: the local set is $\tilde{\square}^5$, of side $11M$, so the threshold is $N_k > 22M$, not $10M$. (4) Corrected, admissible: see (Vol) below.

Row B8 replaced. The old row B8 asked that the finite-volume R-term increment equal the infinite-volume one. That form is ill-posed: the whole lattice of CMP 109 has no R-terms, and no infinite-volume large-field construction exists to compare with. (The author called it false; the verifier corrected this to ill-posed.) CMP 119 p. 279 prints instead that the R″ terms with $X \subset \Lambda$ "do not depend on Λ_{k+1}". Claim L_G of `SOURCES_A15_BDRY.md` proves this: the step output on X is one map $\mathcal{G}_X$ of the inputs on X, the same in every volume. B8 is replaced by three statements: (a) β_j has no R part, because it is subtracted from E_{pure} (p. 279, confirmed on the image); (b) E_{pure} on a non-wrapping $X \subset \Lambda$ equals the $\mathbb{Z}^4$ term exactly; (c) E_R is uniform in K through F2′, with the threshold $N_k > 22M$. All three are [PROVED RELATIVE TO LISTED INPUTS; VERIFIED]: the inputs of Theorem 5.10 and Proposition 5.11, the F2′ inputs ((W4), (Eps), (D); ledger B20), Bałaban's (1.18), (3.48), (4.36), and (Vol). B8 is an input of Theorem 2w ⇒ Theorem 1, not of Theorem 2w.

The volume hypothesis (Vol). Proposition 5.11 needs $N_k > 22M$ at every step $k \leq K$. N_k is smallest at $k = K$, where $N_K = \ell$. So the torus must have side $\ell > 22M$ final-scale cells. This is (Vol) of Section 2.1 (ledger C31; `THEOREM2_PROOF_v4.md` §1.1). The verifier graded (Vol) admissible with a real condition. CMP 109 prints the torus exponent (p. 251, $L_\mu = L^m$) and the cube exponent (p. 257, $M = L^m$) with the same letter m. If the two were equal, then $\ell = 2M$, and (Vol) would fail at the last steps. `SOURCES_T2_VOL.md` reads this as reuse of a symbol, and this paper fixes the two exponents independently. (Vol) does not depend on K, the proof of Theorem 2w does not use it, and it excludes only tori with $\ell \leq 22M$.

The 2D gap (G'_A). Let $A' = H' + D'D'^\dagger$ be the charged block of the SU(2) Wilson Hessian at constant flux f on the coarse $\mathbb{Z}^2$ lattice, in Landau gauge, as in `SOURCES_T2_S123.md` §5. Write $\alpha = ef$, so $|\alpha| = 2|f|$.

Theorem 5.12 ((G'_A), `SOURCES_T2_GAPRIME2.md` Theorem 5.1 and Corollary 5.2 [PROVED+VERIFIED]). *Let $0 \le c < 1$. For every real f with $0 < |f| \le ((1-c)/100)^2/2$, the operator A' has no spectrum in $(-2c|f|, 2c|f|)$, and neither has H' restricted to $\ker D'^\dagger$. For example, $c = \frac{1}{2}$ gives a gap $|f|$ for $0 < |f| \le 1.25 \cdot 10^{-5}$. The constants do not depend on j, L, or the averaging.*

Method. A lattice ladder operator $\Pi = \nabla_0 + i\,\mathrm{sgn}(\alpha)\nabla_1$ gives $X(X - 2|\alpha|) \ge -157|\alpha|^3$ on low states. So the scalar magnetic Laplacian $\mathfrak{L}$ has no spectrum in $[(1+\theta)|\alpha|, (3-\theta)|\alpha|]$ for $\theta > 26|\alpha|^{1/2} + 157|\alpha|$. Then $A' = \mathfrak{L} \otimes 1 - 2\alpha\sigma_y + R$ with $\|R\psi\| \le 48|\alpha|^{3/2}$ near 0, and a contradiction argument gives the gap. The $\ker D'^\dagger$ form follows from $A'D' = D'\mathfrak{L}$. No Harper-band or Helffer–Sjöstrand result is used. Verifier: `SOURCES_T2_GAPRIME2_VERIFY.md`; all four claims confirmed. The constants 26, 157, and 48 are rederived. The lattice commutator is tracked exactly through $2|\sin(\alpha/2)|$. No invariant subspace is needed. The midpoint-transport identification moves from [I] to [P]. The verifier restated the range in units of f (as above) and reproduced the numerics (141 values).

Remark 5.13 (the proved range is narrow). The proof covers only $|ef| \le 10^{-4}$. Numerics show the gap on all of $\alpha \in (0, 3.02]$, with physical ratio ≥ 0.95: at 11 rational fluxes with denominators up to 2213, the smallest |eigenvalue| of A' is $|\alpha|(1 - 0.125|\alpha|)$, and a sparse shift-invert run at $\alpha = 10^{-4}$ gives the ratio 1.0000125. So the proof is valid but far from sharp, and the numerics are not a proof. The Nielsen–Olesen mode is the band at $-|ef|(1 + 0.12|ef|)$, not a zero mode (numerical). The upper edge $3|ef|$ of H' is not proved and is not needed. Open: Prop. 5.6 of `SOURCES_T2_S123.md` uses (G'_A) for $|f| \le \min(f_0, f_1(j))$, where $f_1(j)$ comes from its size condition $g_f \ge 4\eta_1 + 4C_M^2|f|^3$. Round 7 (`SOURCES_T2_FRANGE.md`): this f-range need not fit a fixed window, because (M_j) is a limit $f \to 0$; the proved range suffices (Section 5.11).

Consequence. Open item O1 of `SOURCES_T2_S123.md` is closed. Prop. 5.6 and (Z_j) hold given the inputs of that file. (M_j) (row B4) reduces to (I3) [P], Prop. 4.4 region 1 of `BETA_FP_M.md` [argued], and the neutral sector [I]. Both residues are under attack. Row B4 is a verified reduction, subject to the f-range check of Remark 5.13. Round 7 proves both residues and settles the f-range (Theorem 5.14).

The canonical order of constants. `SOURCES_ORDER_CANON.md` gives one acyclic order of all constants of the source files. It was verified twice, with amendments E1–E8, and it supersedes the orders of rounds 4 and 5 (ledger C5). `SOURCES_ORDER_CHECK.md` found two nominal cycles in the orders as earlier write-outs print them. Cycle A, on the M_1 side (`WC_FULL.md` 6.9, `A1_WALK_FULL.md` 9.1): $M_1 \to M \to \alpha''' \to \alpha'' \to B_{A''}, \delta_{A''} \to M_1$. Cycle B, on the M side: $M \to \alpha''' \to \alpha'' \to \alpha' \to \varepsilon'' \to M$. Neither cycle is real. A split resolves both: the level constants (rates, prefactors, Γ_n, k_n, ε'') are bounds on domains capped by a_1, a_{**}, and $M\alpha_0 \le a_0$, so they are free of M and of the α's and come before M_1; the domain sizes α''', α'', α' are smallness choices and come after M. The verifier also found forward references and a double count of M in the J scale. The amendments fix them, with two new entries: real $\alpha_0 \le \alpha_0'/(CM)$ with $C > c_J$, and $M \ge 1/c_J$. The block size M has 32 lower bounds and no upper bound (ledger C24), so the admissible set of M is an up-set. G_Γ is bounded uniformly in M (`SOURCES_G3_GGAMMA.md` + VERIFY; Section 5.9), so (Hα) is one more lower bound on M and adds no cycle. One caveat stays: the plaquette row of Lemma M1 is nested by reading only (amendment E6).

5.11 Step 8′′′′: round 7, the implicit Group A rows, (M_j), L5, and the sweep

Round 7 treats the implicit or unprinted Group A rows A16–A19 and A23, the Group B rows B4, B7, B10, and B17, the certification of (H2) at more values of L, and the canonical order of constants. Each result has its own project file and an independent verifier file. The exception is `SOURCES_T2_FRANGE.md`, a short check with no verifier file, and `SOURCES_T2_F_STEPB.md`, whose verifier is still pending. Each result holds relative to the inputs that it names. Nothing in this step makes Theorem A or Theorem B unconditional.

The fixed-level input (M_j) (ledger B4).

Theorem 5.14 ((M_j), `SOURCES_T2_MJ_REST.md` Theorem 4.2 and §5 [PROVED RELATIVE TO LISTED INPUTS; VERIFIED]). *Fix $j \geq 1$ and odd $L \geq 3$. The one-loop increment of level j at constant abelian flux f satisfies $\delta\Gamma_j(f) - \delta\Gamma_j(0) = n_2 d_j f^2 + o(f^2)$ as $f \to 0$, with d_j the formal coefficient of `KAPPA_BOUNDED.md` (1.1). The charged-sector error is $O(|f|^{9/4}) + o(1)f^2$. The neutral sector contributes its formal coefficient exactly in the limit.*

Method. Region 1 of Prop. 4.4 of `BETA_FP_M.md` is not an inequality. It is an asymptotic identity: the f^2 coefficient of a spectrally localized trace equals the per-momentum formal integral. The file replaces the region split by a split of $\log|x|$ in three pieces with a shifted logarithm. The gapped piece F_s has shift $s = |f|^a$, $a = \frac{3}{4} \in (\frac{2}{3}, 1)$, so $f/s \to 0$ and a Moyal expansion has a small parameter (Lemma A, with an exact two-factor trace identity and an approximate Moyal inverse to order $M = 14$). The Nielsen–Olesen window piece is the term II of `SOURCES_T2_S123.md` Prop. 5.5. The localized remainder is $O(s^3 + |f|^3)$. The neutral sector (N2) is proved by an induction on bordered Hessians: $\pm\partial^p \Delta_i \leq C_p \Delta_i(0)$ for $|f| \leq f_*$. On a finite torus the d constant 1-forms are zero modes; they are not gauge directions, and they form a null set of the per-cell trace, so the infinite-lattice statement does not see them. Inputs: (G'_A) (Theorem 5.12); (I3) [P] (`SOURCES_T2_I3.md` + VERIFY); the results of `SOURCES_T2_S123.md` that its verifier confirmed; Prop. 2.2 of `U3_LEMMA51.md`; and one low-risk convention [I]: the coefficients $b_{t,m}$ of `KAPPA_BOUNDED.md` (1.1) are the Moyal coefficients of §3.1 at $\sigma = 0$. Verifier: `SOURCES_T2_MJ_REST_VERIFY.md`; all four claims confirmed (Lemma A Steps 2–4, the exponent count, the (N2) induction, and the numerics; Section 6.8). Only the $o(f^2)$ form is proved. The printed rate $O(\varepsilon|f|^3 \log(1/|f|))$ of Prop. 4.4 is not proved, and no consumer uses it. The f-range. `SOURCES_T2_FRANGE.md` shows that the proved range of Theorem 5.12 suffices. (M_j) is a limit $f \to 0$ at fixed j, so the gap is needed only on some punctured neighbourhood $0 < |f| \leq f_{\lim}$; its size plays no role. The size condition $g_f \geq 4\eta_1 + 4C_M^2|f|^3$ of Prop. 5.6 of `SOURCES_T2_S123.md` has a left side linear in $|f|$ and a right side $O(f^2)$ ($\eta_1 = C_\eta f^2$), so it holds for all small $|f|$. Usable range: $0 < |f| \leq \min(f_0(c), f_1(j, L), f_*(j, L))$. The numerical gap up to $\alpha \approx 3$ stays supporting evidence only. Consequence. Row B4 is proved relative to listed inputs. The chain from `SOURCES_T2_S123.md` through (M_j) to Prop. 1.1 of `KAPPA_BOUNDED.md` is closed up to conventions and readings.

History dependence L5 (ledger B7).

Theorem 5.15 (Lemmas D″-nest and D′-nest, `SOURCES_T2_B7_CLAMP.md` Route 2 [PROVED RELATIVE TO LISTED INPUTS; VERIFIED]). *The conclusions (a″), (b), (c′) of Lemma 3.4, with the same constants $\beta = 3b/4$, $B' = 5b/4 + 2R$, $D' = 2R$, $g_1 = (\gamma^{-2} + 2R)^{-1/2}$, hold for true (unclamped) trajectories, with no clamp. Every evaluation of β lies on a true trajectory with all $u_i \geq \gamma^{-2}$, inside Bałaban's domain. The same holds for Lemma 3.2 with its constants.*

Method. Nested intervals. The envelopes $p_k, q_k = t + T_0(k, K) \mp (b/4)(K - k)$ subtract $\beta_{k+1}(0)$ exactly, and the slack shrinks by $b/4$ per step, which equals the worst single-step error $C_1\gamma$. So negative increments are absorbed. At each step the intermediate-value theorem selects a closed subinterval of u_0 on which u_{k+1} sweeps the next target interval. The window bound (5.1) is used at every endpoint $n \le K$. Inputs: $7'$ on trajectories; (C-tr), the continuity of one step in its inputs (the single CMP 116 step, Theorem SM, `SOURCES_HHOL.md` §2–§3, `SOURCES_GGAIN_NONMONO.md` or A15, and the enlarged input list of N6); and one reading [I]: Bałaban's step $k + 1$ uses only $g_0, \dots, g_k$. Route 2 still evaluates the step map at one off-trajectory mixed point per step, inside the proof of (C-tr). Verifier: `SOURCES_T2_B7_CLAMP_VERIFY.md`; Route 2 confirmed with the constants of `THEOREM2_PROOF_v4.md`. Route 1 (continuity on the clamped cube) holds, but its γ condition uses an undefined constant γ_{hol}, so it is dropped. The history form of N3 needs Route 1; the one-argument form does not. The quote of CMP 109 p. 298 is confirmed on the image; "β_j only on the trajectory" is a correct reading of the construction.

The cut and the analyticity domain (ledgers A17, A19).

Proposition 5.16 (A17, `SOURCES_A17_A19.md` [PROVED RELATIVE TO THE PRINTED TEXT AND ONE READING; VERIFIED]). *On every bond b that the cut χ_k of CMP 109 (2.9) covers, $B'(b) = g_k B(b)$ exactly. Hence* $\operatorname{supp}\chi_k \subset \{g_k|B|_Y < \varepsilon_1\}$*, the analyticity domain of CMP 116 (1.34), with margin $r = 1$ and no slack.*

Method. CMP 109 (2.9) takes the product over $T^{(k)} \setminus \{b_0(c)\}$; $hB = 0$ off the bonds $b_0(c)$ (p. 267); CMP 116 (2.3) writes the cut directly as $|B(b)| \ge \varepsilon_1/g_k$ on the same bonds. The bound (1.20) of CMP 116 gives only the converse inclusion. The one reading [I, strong]: C is the identity on the remaining bonds; the norm on Y is the per-bond sup of CMP 99 (3.41).

Proposition 5.17 (A19, `SOURCES_A17_A19.md` §4–§5 [REDUCED TO READINGS; VERIFIED]). *The bound (190) of CMP 102 holds at complex B on the ball $|B| < 2C_1\varepsilon_1$, given three readings [I]: the complex clauses of (189) and (73), which Bałaban does not print, and the identification of the set (172) with this ball. Fallback without the ball reading: the two-constants theorem gives the bound on $|B| < \frac{1}{2}C_1\varepsilon_1$ with harmonic measure $\omega = (4/\pi)\arctan\frac{1}{3} = 0.4097$, so the decay rate is multiplied by* 0.59.

Four of the five named inputs carry a printed complex clause (Prop. 4, Prop. 6, (55), (80) of CMP 102); (130) is linear. The fallback still needs a complex sup bound on kernel entries, from (187) and (55). Verifier: `SOURCES_A17_A19_VERIFY.md` (page images). A17 confirmed. The option C9′ (keep the cut at ε_1 and run CMP 116 §1 with domain constant ε_1/r) is confirmed for CMP 116 §1, with the condition $4B_0C_1e^{16\kappa_1}\varepsilon_1 \le \frac{1}{4}\alpha_2$ (the file had $\frac{1}{2}\alpha_2$). C9 is no longer needed at $r = 1$. A19 corrected from "proved" to "reduced to readings". The Cauchy step of `SOURCES_MA2.md` §4.4 reaches the boundary of the open ball, but a limit $\rho \uparrow C_1\varepsilon_1$ keeps its constant c_N/C_1.

The regular class (ledger A18).

Proposition 5.18 (A18, `SOURCES_A18_REGULAR.md` [VERIFIED REDUCTION, WITH CORRECTIONS]). *Relative to printed results: (i) the background domains of CMP 109, 116, and 122 lie in the regular class of CMP 99 (3.35)–(3.36); (ii) Lemma Q: $Q_k(U) = Q_{\text{free}} + O(\alpha)$ in a per-row axial gauge, uniformly for $L \ge 3$, and $O(M\alpha_0)$ at operator level; (iii) Lemma C: $\|C(U)\|^2 \le c_C(1)(1 + O(\alpha))$; (iv) $c_J = O(\alpha_0)$, with no factor M; (v) σ-localized operators add nothing.*

Method. Every explicit use of (3.36) in CMP 99 goes through the current J, so a bound on J replaces (3.36) up to $O((M\alpha_0)^2)$. Lemma Q telescopes $Q_k(U) - Q^k$ over levels on CMP 98 (124)–(125), (139)–(146). Lemma C uses the printed definition of $b_0(c)$: each row of the linear one-step map contains exactly one eliminated variable. Verifier: `SOURCES_A18_REGULAR_VERIFY.md` (page images). Corrections: the inclusion constant is CMP 109's unnamed cube constant $c_{\square}^{109}$, and 5.2(d) needs two factors L; the telescoping sum of Lemma Q misses a factor 2^{k-j-1}, so the correct sum is $c\alpha L/(L-2)$, k-uniform for $L \geq 3$; the CMP 122 cube patching is [UNSUPPORTED] and is replaced by CMP 99 at cube scale $M_{\square}^{99} = CM/10$, with the new canon condition $M \geq 5M_{99}$ $(10 \mid C^{122}M)$. Residue: [R] CMP 102 Thm 1, CMP 98 Prop. 2 and (139), CMP 99 (3.133), CMP 109 p. 263; [I] I2, I3, $U_{k,Z}$ obeys CMP 102 Thm 1, and B, B_5 are fixed numbers of (d, L). Conditions (A18-1)–(A18-4): ledger C41. The item (C2) of `SOURCES_MA2.md` is merged into this row.

The large-field T-step rate loss (ledger A23).

Proposition 5.19 (A23, `SOURCES_A23_TSTEP.md` [VERIFIED REDUCTION (PROOF SKETCH)]). *In one large-field T step the decay rate κ loses at most the fraction 13δ: 10δ printed for the small-field step (CMP 116 (2.28)–(2.41)) and at most 3δ unprinted for the Z_0-marked cluster expansion of CMP 119 (3.44)–(3.47). No loss inside T is an additive constant in κ.*

Method. "CMP 109 Sects. 6–7" is CMP 116 Sects. 1–2 in a preprint numbering, so the two deferral targets of row A23 are one printed source. Each merging step costs a δ-fraction of the current rate (one δ for a sum over pieces, one for a volume factor by (2.30)). The tree bound is printed as CMP 116 (2.27). Verifier: `SOURCES_A23_TSTEP_VERIFY.md` (page images); confirmed. The verifier counts 2δ for the unprinted part, so 13δ is a safe upper bound. Open: the count R1 (a sketch) and R3, Bałaban's assertion that $H'(Z_0)$ obeys (2.38). Consequence for ledger B22: (K1) becomes (K1′) $\delta \leq \frac{1}{13}(1 - 2(1+4\beta)/(\theta_K L))$ with $\theta_K \geq \frac{1}{2}$ and $\kappa \geq 2m_{\mathrm{tot}}$; at $L = 13$, $\beta = \frac{1}{4}$ this is $\delta \leq 5/169 \approx 0.0296$, so $\delta = 0.025$ is the choice (0.03 fails; Lean `K1_numeric_13` in `lean/YmLean/K1Prime.lean`, no `sorry`, standard axioms). (K2), (K3), κ_{eff}, $\kappa \geq 1536$ are unchanged. The paragraph on `SOURCES_T2_KAPPA2.md` in Section 5.2 carries the new form.

The propagator replacement (ledger A16).

Theorem 5.20 (A16, `SOURCES_A16_REPLACE2.md` [PROVED RELATIVE TO LISTED INPUTS; VERIFIED WITH MINOR CORRECTIONS]). *The sourced covariance $C_f = (A_{\exp} - g^2\mathcal{W}_S)^{-1}$, $\mathcal{W}_S = C^*\mathcal{V}_S C$, has the properties Q1–Q11 that CMP 116 uses for $C^{(k)}$, and the uses (U-a) whole-volume C_f and $(C_f)^{1/2}$, (U-b) σ-differences of $(C_f)^{1/2}$ and Δ^f, and (U-c) the tree form of Δ^f, under the tree conditions C40.*

Method. The sentence of CMP 116 p. 3 licenses only the replacement in (1.3)–(1.4); it is not cited for C_f. The properties are proved directly. $\mathcal{W}_S$ is written as a tree sum of the type of Proposition W_N of `WC_FULL.md`, on the complex chain of `WC_FULL.md` §6, with at most 12 junctions per term and $K_W^T = 2C_{\mathcal{D}}(C'_{\mathcal{D}})^{12}B_{\mathcal{W}}$. The conditions (A16-a^T) and (A16-b^T) are linear in $g^2 r$. History. The first proof (`SOURCES_A16_REPLACE.md`) had a false locality step: it treated each $\mathcal{W}$ term as a one-domain block, but $\mathcal{W}_S$ is not finite range, because $H_1 = G_1Q^*(QG_1Q^*)^{-1}$ is nonlocal (`SOURCES_A16_REPLACE_VERIFY.md`). The repaired proof uses the tree form. Inputs: Theorem 5.3 (with W_N and §6), Theorem 5.2 (Lemma C, WG-G1, 5.2, 5.5), Theorem 5.1 for G_1 (conditional on (C)), Theorem 5.5, `SOURCES_G3_GGAMMA.md` 2.1, and `SOURCES_D2.md` (2.a), `SOURCES_D3W8.md` §1.1, §1.4. Residue [I]: (M1) the completeness of the property list (CMP 116 pp. 12–17); (S_H) the scale normalization of D_S, J_S; the volume convention of the CMP 122 Ω-setting; the s-version

choice for $(C_f)^{1/2}$. Euclidean covariance of C_f is false for non-constant f; no consumer may use it. Verifier: `SOURCES_A16_REPLACE2_VERIFY.md` (page images); claims (1)–(3) confirmed; (U-b) corrected without a change of constant order (R_2 is anchored at both ends in Z_0; the product form needs concatenation at the level of elementary blocks).

The large-field holomorphy chain (ledger B17, chain item 5). `SOURCES_T2_HMINUSC.md` closes the two items of `SOURCES_MA2.md`. (C1) is [PROVED GIVEN A19 AND A18; VERIFIED]: every condition on CMP 102's ε_1 and C_1 is an upper bound (and $C_1 \geq 1$), so ε_1 can be raised per scale to give Cauchy room. The four cases K1–K4 of Proposition 5.17 give radii $\kappa_H \in \{2, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}\}$ and constants c_N/C_1 to $16c'_N/C_1$, all free of C_1. The price is the condition (C1*) $c_m M_0\, g_k p_0(g_k) \leq \frac{1}{2}\kappa_H C_1^{102} a_*^{102}$ (ledger C38), with M_0 placed in the canon before γ (C39). (C2) is [REDUCED] and merged into A18, with two readings [I]: the averaging map $m = M^\bullet \circ Q_k^{s*}$ is local and analytic, so V' always enters through m; and $R_1 < L$ suffices for condition (1) of CMP 102. Verifier: `SOURCES_T2_HMINUSC_VERIFY.md` (page images); the map of (H−) is CMP 122 I (1.74), not (1.20), and "~ 4" means four layers of M-cubes. So chain item 5 reads: B17 $\Leftarrow$ (M-A) $\Leftarrow$ (H−) $\Leftarrow$ A4 + A19 (K1–K4) + A18 + (C1*). B17 stays a verified reduction.

The sweep and N3 in the D″ form (ledger B10). `SOURCES_B10_SWEEP.md` with `SOURCES_B10_SWEEP_VERIFY.md` (page images, three readers over 63 pages) clears the four items of B10. (i) All 19 rows of the sweep of CMP 119 (2.6)–(2.9) now have a second reader. The page images of CMP 122 I–II Sect. 1 show about ten cross-scale uses and five exponent or threshold conditions that the list missed. They are now listed in the addendum of `SOURCES_A11_CONST.md` as X1–X10 and S12–S16 (ledger C42). None is a new type: each is a (P-up) window, a (P-low) window with a factor $(1+\beta_0)^a \operatorname{poly}(n-m)$, or a single-scale γ-smallness threshold. The form $p_1(g) = (\log g^{-2})^{p_1}$, $p_1 < p_0$, is confirmed on CMP 122 I p. 183. (ii) Prop. 4.3 of `BETA_FP_M.md` is confirmed given (S1)–(S3), for $j \geq j_1$ only; below j_1 it needs the open (S3w), which Theorem 2w does not use. (iii) The Prop. 3.6 assembly is confirmed. (iv) N3 in the D″ form: Lemma H″ (real base) takes J as the component of 0 in $\{c \in [-2r, 2r] : u_k(c) > \gamma^{-2}\ \forall k \leq K\}$ and gives $S_3 \leq (D'/\beta + 1)\gamma^3 + 2\gamma/\beta$; N3″ (complex) splits the bracket at $\beta/2$, gives $\operatorname{Re} u_i \geq \gamma^{-2} + \max(0, (\beta/2)(K-i) - D')$, and has slack 6.7 at the N3A radius. The "$+\eta$" of `SOURCES_N3A.md` is not needed. The history form of N3 needs Route 1 of Theorem 5.15.

The canonical order, section F. `SOURCES_ORDER_CANON.md` section F adds 28 condition entries from the round-7 files (new lines B14a, B17a, B18a, B22a, B29a) and resolves the symbol clashes by renaming (a_*^{102}, C_1^{102}, ε_1^{102}, κ_H, θ_H, θ_K, $M_\square^{99}$, and others). A topological pass finds the order acyclic. It removes two forward references: (K1′) uses the bound $\theta_K \geq \frac{1}{2}$, so δ is a canon constant (line B5) and $\kappa \geq 2m_{\text{tot}}$ moves to line B7; and the cube cap of CMP 102 Thm 1 is booked on ε_1^{102} and ε_0^{109}, not on M, so M stays one-sided. Four constants are two-sided, and each has an existence condition: p_1 (exists under the B25 condition on p_0); ε_0^{102} (nonempty by CMP 102 Thm 1); ε_1^{102} per scale (exists if and only if (C1*) holds and $\varepsilon_1^{(0)} \leq a_*^{102}$); and N per scale (exists for γ small; $N = R_j$ is admissible). The budget α_5 is shared between (Hα) and the entries of line B32. Not yet in the canon: the constant C_1 of (K3) (to be named C_1^{FX}) and $\kappa \geq 1536$.

The lattice log coefficient (ledger B3) [REDUCED [R], UNVERIFIED]**.** `SOURCES_T2_F_STEPB.md` narrows Step B of (F). Theorem 3.1 of `BETA_FP_H.md` uses (F) only through (3.1), which is equivalent to $G(\varphi) - G(\lambda\varphi) \to n_2 c_2 \ln(1/\lambda)$ as $\varphi \to 0$ for $G = F_W(\varphi)/\varphi^2$ and $\lambda = L^{-2k}$. The file reduces this to (L-LL), the existence of the limit of

$H_\varphi(\mu) = \varphi^{-2}[W(\varphi;0) - W(\varphi;\mu|\varphi|)]$ as $\varphi \to 0$ (its value is not needed; it holds the Nielsen–Olesen zero), to a gauge-covariant kernel bound (Lemma 1, cited as standard and not checked), and to the gauge-fixing part of Step C [I] (the ghost part is proved). The continuum side is X5. The verifier of this file has not reported, so B3 stays on the list of rows without a proof or a verified reduction. Open numerical tension. The single-step values β_1 at $L = 5, 7, 9, 11$ do not follow the slope $c \log L$, $c = 0.3715$ (Section 6.1). The open question is whether 0.3715 is compared with the right quantity (the single-step β_1 against the fixed-point β_∞), or whether the gap sits in the coarse bracket, in the dependence on the family, or in the $N_g = 8$ grid at $L \geq 9$. The tension may also bear on B4 through Theorem 3.1 of `BETA_FP_H.md`. The claim of `NUMERICS` §7 that the first-step constant decreases "toward the universal slope" is under verification.

(H2) at $L = 15, 17, 21$ (ledger B1). The certificate of Theorem 6.1 runs unchanged at these L and is now certified with the same standing as at $L = 13$ (Section 6.8). Other L are not used by the targets: C3 fixes the certification target at $L = 13$.

5.12 Step 9: the alternative (E_U) route

`SOURCES_EU.md`**: (E_U), uniform block extension (alternative route).** (E_U) asks for a fine field A with $Q(U)A = B$ and covariant energy $\leq K\|B\|^2$ for every block field B, uniformly in k; in the free case $K_0(4) = 103.25, 146.73, 185.64, 195.19$ at $N = 3, 5, 13, \infty$ ($= \kappa_4 - 1$ at $a = 1$), and for regular U the dressed extension $A = \mathcal{E}_U(Q(U)\mathcal{E}_U)^{-1}B$ gives $K \leq K_0(1 + c_{2/3}(M\alpha_0)^{2/3} + c_1 M\alpha_0)$. The claim that transport and back gives $Q(U)A = B$ exactly is false, and the linear form $K \leq K_0(1 + cM\alpha_0)$ is not proved. K_0 is computed in momentum space by `numerics/k0_d.py` from the CMP 98 (124)–(125) form of $Q(V_0)$, an equal-weight average over L^d straight lines with an $O(L^2\alpha_0)$ remainder, and is checked in real space at $d = 2$. For regular U the free extension kernel is dressed by parallel transport, the constraint error $\|Q(U)\mathcal{E}_U - I\|$ is bounded linearly in the plaquette bound $\alpha_F \leq 20M\alpha_0$ by lattice Stokes, and IMS patching over flat frames of radius $R \propto \alpha_F^{-1/3}$ gives the power 2/3. Verifier: `SOURCES_EU_VERIFY.md`, K_0 [NUMERICAL, REPRODUCED], the bound on K [DERIVED SKETCH], and (E_U) [VERIFIED REDUCTION] to kernel-moment bounds Bałaban does not print; corrections: the $N = \infty$ tail is $\propto 1/M^3$, not $1/M$; (139) is on journal p. 39; K_0 belongs to the straight-line Q_{98}, not to the corridor Q_{95}; $K \to K_0$ is indicated, not proved; the k-step remainder constant is $2C_1'$ by CMP 98 (143). Open inputs: A21 (k-uniform moments and decay of the free kernel E_0); the unprinted constant C_1' (A9, a verified reading); the four [I] items (a)–(d) of `SOURCES_EU_VERIFY.md` §5 (uniform moments Γ_j, ν_j, lattice-sum factors, the boundary cutoff at $\partial\Lambda$, the gauge-fixing term); Q_{95} versus Q_{98}; the linear form. The route is an alternative to A1 and is not binding. A21 is now closed, see the next paragraph.

`SOURCES_T2_EUMOM.md`**: ledger A21, k-uniform moments of E_0.** The free minimal-extension kernel E_0 of the straight-line average $Q_{98}(1)$ (composition of k one-step averages, $N = L^k$) obeys $|E_0(y,c)|, |\nabla_\nu E_0(y,c)| \leq Ce^{-\delta|y-c|}$, with C, δ depending on d only. All moments ν_j, γ_j, the block sums, and Γ_j are bounded uniformly in k. The proof uses symbol identities in momentum space, lower bounds on the symbols, a weighted L^2 Combes–Thomas bound uniform in η, and pointwise regularity from the symbol; it uses no Bałaban estimate. Verifier: `SOURCES_T2_EUMOM_VERIFY.md`, [PROVED+VERIFIED], 31 claims confirmed, 3 corrected, 1 unsupported (a side remark); it closes open item (i) of `SOURCES_EU_VERIFY.md`. Open inputs: the other [I] items of the (E_U) route listed above.

`SOURCES_CDC.md`**: the norm bound γ_1 (ledger A7).** The norm bound $\|C^*\Delta_k C\| \le \gamma_1$, uniform in k and regular U, is reduced: by Woodbury $(QG_1Q^*)^{-1} = N_1^{-1} + a$ exactly, so the term $-a\langle B, B\rangle$ of CMP 99 (3.156) cancels and $\gamma_1 = L^{d+1}[(1+\varepsilon)K + \varepsilon a + c_J]$ with $\varepsilon, c_J = O(M\alpha_0)c(d, L)$ and K the constant of (E_U); in the free case $K \le a(\kappa - 1)$, which is 195.19 at $d = 4$, $a = 1$. The brief's mechanism, a local difference operator for which unitary U suffices, is [REFUTED]: the leading term $(QG_1Q^*)^{-1}$ is an inverse whose naive coefficients scale as L^{2k}, so the regularity (3.35)–(3.36) is needed for the upper bound. Comparing (3.155) with (3.156) identifies $\langle B, (QG_1Q^*)^{-1}B\rangle$ as the minimum of $\langle A, G_1^{-1}A\rangle$ over $QA = B$, and the split $G_1^{-1} = M_1 + aQ^*Q$ of (3.128) gives the exact cancellation of a. The remaining minimal energy is bounded by a trial extension of B (the lemma (E_U)), the Δ'_π and J parts by $O(M\alpha_0)$ perturbation, and the free constant by the CMP 95 Fourier formula (1.99)–(1.101). Verifier: `SOURCES_CDC_VERIFY.md`, [VERIFIED REDUCTION]: the Woodbury cancellation is confirmed (the variational form needs only $G_1 > 0$ and Q onto), but (E_U) is not the last input, since CMP 99 (3.132) prints a k-uniform kernel bound on $(QG_1Q^*)^{-1}$ that the file omits; corrections: $\|C\|^2 \le L^{d+1}$, not $1 + L^{d+1}$; K_0 is an upper bound only; the free case is proved for single-scale $U = 1$ only; J is a second derivative of A; the negative-mode branch is dropped, and "regularity is needed for γ_0" is unsupported. Open inputs: CMP 99 (3.132) (ledger A1), or in its place (E_U) (A21, alternative route); A18 for the J-term and $\|C(U)\|$; the multiscale correction $e^{-\delta RM}$ to the free constant [I]; the s-localized operator $C^{(k)}(Z_0, 0)$ that Lemma G3 uses, handled by LOC (A2, A3). Through this file ledger A7 was reduced to A1 + A18. Round 5 supersedes this reduction: Theorem 5.4 proves A7 relative to the inputs of Theorem 5.3, without the kernel form of A1.

6 Numerical and certified results

This section collects the numerical results that support the hypotheses of Section 3. Sections 6.1–6.3 give the one-loop increments and the per-level quantities of the Route K proof. Sections 6.4–6.6 give the values of (H2), the one-step values, and one rigorous point enclosure. Section 6.7 proves (H2) at $L = 13$ (Theorem 6.1, certified). Section 6.8 adds the certification at $L = 15, 17, 21$, the round-7 checks of the 2D gap and the neutral sector, and one open tension. Section 6.9 gives the original plan for a cell certificate, now an independent cross-check at $L = 13$. SU(2), infinite volume, Balaban's normalized-trace units. Except for Sections 6.6, 6.7, and the first paragraph of Section 6.8, none of this is certified. Only odd L is inside the proofs; $L = 2, 4$ use corner blocks and are outside them.

6.1 First increment $\beta_1(0)$

Sources: `NUMERICS` §6–§7; `UMK_BETA1.md`.

L	$\beta_1(0)$	how	$b = c \log L$	$\beta_1 - b$	status
2	0.630	$N_g = 12$, $p \to 0$ fit	0.2575	+0.372	[NUMERICAL, REPRODUCED] (`KAPPA_BOUNDED_VERIFY` §4: 0.6296)
3	0.664	$N_g = 8$, $p \to 0$ fit (0.6625–0.6661)	0.4082	+0.256	[NUMERICAL, REPRODUCED]; N_g 6 → 8 shift ≈ 0.015
4	0.757	$N_g = 6$, single $p = 0.2$; $O((Lp)^2)$ bias ≈ 0.01	0.5150	+0.242	[NUMERICAL]

5	0.749 ($N_g = 12$, $p = 0.1$); fit 0.743	umklapp code	0.5979	+0.145	[NUMERICAL]
7	0.803	umklapp code, $N_g = 8$, $Lp = 0.5$	0.7229	+0.080	[NUMERICAL]
9	0.833 ($Lp = 0.5$); 0.846 ($Lp = 0.9$), 0.851 ($Lp = 1.0$)	umklapp code, Windows, $N_g = 8$	0.8163	$\approx +0.02$	[NUMERICAL]
11	0.850 ($Lp = 0.5$); 0.868 ($Lp = 1.0$)	umklapp code, Windows, $N_g = 8$; $N_g = 8$ grid adequacy at $L = 11$ unchecked	0.8908	≈ -0.04	[NUMERICAL]

The first-step constant $\beta_1 - c \log L$ is +0.37, +0.26, +0.24, +0.15, +0.08, +0.02, −0.04 for $L = 2, 3, 4, 5, 7, 9, 11$. It decreases with L and changes sign between $L = 9$ and $L = 11$. $\beta_1(0)$ itself stays positive at every L computed. The umklapp code was validated against `ym_momentum` to 10^{-12} per K by its author; no independent verifier for the β_1 values at $L \geq 5$. The queue continues to $L = 15$ on Windows. No value at $L > 11$ exists yet. Round 7: the decrease does not level off, and the slope of β_1 in $\log L$ is well below c; this is an open tension (Section 6.8).

6.2 Later increments ($L = 2$ only)

quantity	value	status
$\beta_1 + \beta_2$ (composed, $N_g = 4/6/8$)	1.148 / 0.967 / 0.953	[NUMERICAL]; single $p = 0.2$, Lp bias label
per-level $\beta_2(0)$	≈ 0.32 ($0.953 - 0.630$). Per-level recursion 0.317–0.341 ($N = 8$–12, single $p = 0.2$). Symmetric Nyquist: 0.33334 ($N = 8$, $p_1 = 0.2$), 0.31679 ($N = 12$, $p_1 = 0.2$), 0.32283 ($N = 12$, $p_1 = 0.4$) (`BETA_FP_VERIFY` §7; `BETA_FP_PW_VERIFY` §5–§6). Level-1 values carry N-aliasing of about ±5%.	[NUMERICAL]; per-level recursion reproduced in `BETA_FP_VERIFY` §7
$\beta_3(0)$	0.40 ($N = 8$), 0.286 ($N = 12$), 0.261 (verifier, symmetric Nyquist, $N = 8$)	unconverged; not usable
abelian sector S_k^{ab}	$(1 - L^{-4k})/4$, matched to all digits	[PROVED+VERIFIED]; [LEAN] counting part (`AbelianCount.lean`)
(U) test	$\lvert\beta_{j+1}(p) - \beta_{j+1}(0)\rvert \leq 0.2p^2$ at every computed point ($L = 2$ levels 0–1, $L = 3$ level 0, $p \leq 0.8$); uniformity in j not testable with two levels	[NUMERICAL, RUN BY THE VERIFIER] (`BETA_FP_PW_VERIFY` §6)

All computed increments are positive.

6.3 Quantities of the Route K proof

quantity	value	status
$d_1 = b - \beta_1$, $L = 3$	-0.256 ± 0.01 (fits $-0.254 \cdots - 0.258$)	[NUMERICAL, REPRODUCED]
d_1, κ_1, $L = 2$	-0.372, 1.114	[NUMERICAL, REPRODUCED]
d_2, d_3, $L = 2$	≈ -0.44, ≈ -0.45 (d_3 uses unconverged β_3)	[NUMERICAL]; bounded-looking, not a test
$\rho_1 = \max_K \|H_j(K)\|/L$, $L = 3$, $j = 1, 2, 3, \infty$	1.652, 1.736, 1.757, 1.760	[NUMERICAL, RUN BY THE VERIFIER, `U3_AW_VERIFY` §6]; refutes (HS); $h^2/L^4 \approx 0.34 < 1$; one-step C^1 factor $\rho_1^2/L > 1$ for $j \geq 2$
ρ_1, $L = 5$, $j = 1, 2, \infty$	1.860, 1.9565, 1.963; $\rho_1^2/L = 0.771$	[NUMERICAL]; $j = 1$ and the bound measured by the verifier (`H2_LARGE_L_VERIFY` §6)
ρ_2 ((H2)), $L = 3, 5, 7, 9$	see Section 6.4: 2.0634, 2.108, $\geq$ 2.1131, $\geq$ 2.1142 at $j = \infty$ against thresholds 3, 5, 7, 9	[NUMERICAL, REPRODUCED] at $L = 3, 5$; [NUMERICAL, RUN BY THE VERIFIER] at $L = 7, 9$
a_*, $L = 5$	0.0107 (binding mode near $K = 0$, gauge/ghost sector)	[NUMERICAL, RUN BY THE VERIFIER] (`H2_LARGE_L_VERIFY` §6)
(H″), $n = 2$, $L = 3$	$2.056 < 2.828$	[NUMERICAL, RUN BY THE VERIFIER], coarse grid, one level pair
$\sigma_j^{(1)}(0)$ J-part, $j = 0, 1, 2$ ($d = 2$ proxy)	0.750000, 0.75015, 0.75051 vs 0.75	[NUMERICAL, RUN BY THE VERIFIER] supports Theorem W
$\sigma_j^{(2)}(K)$ scan	not run: no $d = 4$ constant-field per-momentum code	—

6.4 Values of (H2) [NUMERICAL; (H2) AT $L = 13$ IS CERTIFIED IN SECTION 6.7]

Code: `numerics/h2_scan.py` (umklapp Schur pieces, one colour, Gram form; `H2_LARGE_L` §1). Code structure CONFIRMED (`H2_LARGE_L_VERIFY` grade 1): the one-colour reduction is exact at $U = 1$, the H and Δ_{out} blocks and the Gram identity are correct, the separable Prop. A form is exact. Sign slip in `H2_LARGE_L` §4 ($H = +[\mathfrak{B}^{-1}]_{f,c}$); harmless.

L	j	ρ_2 (scan max)	threshold L	status
3	2	2.00857	3	[NUMERICAL, REPRODUCED] (`B2_TWOSTEP_VERIFY` §4; `H2_LARGE_L_VERIFY` §1)
3	3	2.05601	3	[NUMERICAL, REPRODUCED] (spot value 2.05599 at the argmax)
3	∞	2.0634 (2.06332–2.06377 over l_{max} 2–5)	3	[NUMERICAL, REPRODUCED]; margin 31%; $q_1 \leq 0.473$, $q_2 \leq 0.053$

5	2, 3, 4, ∞	2.098, 2.108, 2.108, 2.108	5	[NUMERICAL, REPRODUCED]: independent dense code gives 2.098283 at $j = 2$; the verifier's own search gives 2.107921 at $j = \infty$, same argmax; margin 58%
7	∞	$\geq$ 2.1131 (face diagonal only)	7	[NUMERICAL, RUN BY THE VERIFIER] (spot 2.11309)
9	∞	$\geq$ 2.1142 (one point, $(2.6, 2.6, \pi, \pi)$)	9	[NUMERICAL, RUN BY THE VERIFIER]; the $j = 2$ point of `H2_LARGE_L` gives $\geq$ 2.1104

- Shape. The argmax lies on the face $K_3 = K_4 = \pi$, on the diagonal $K_1 = K_2$: $\approx 2.22 \to 2.40$ ($L = 3$, $j = 2 \to \infty$), ≈ 2.54 ($L = 5$). Off-face points are lower: best off-face 1.995 ($L = 3$, $j = 2$), 2.004 ($L = 3$, $j = \infty$), 2.088 ($L = 5$, $j = \infty$). The corner (π, π, π, π) stays below the face peak, but the gap shrinks: 0.0070, 0.0028, ≥ 0.0025, ≥ 0.0023 ($L = 3, 5, 7, 9$). The corner is the obvious place for a competing maximum at large L.

- Level dependence. At $L = 5$, $\rho_2(\infty) - \rho_2(j) = 9.64 \cdot 10^{-3}$, $4.0 \cdot 10^{-4}$, $1.6 \cdot 10^{-5}$ ($j = 2, 3, 4$); ratios 24.1, 25.0, against $L^2 = 25$. At $L = 3$ the ratios are 7.5, 8.7. Levels $j = 4, \ldots, \infty - 1$ are not sampled at $L = 3$.

- L-trend. $j = \infty$: 2.0633, 2.1079, $\geq$ 2.1131, $\geq$ 2.1142 ($L = 3, 5, 7, 9$). This is consistent with a bounded ρ_2 near 2.11–2.14. The rate claim "at least as fast as L^{-2}" is CORRECTED (`H2_LARGE_L_VERIFY` grade 5): it used a lower bound at $L = 7$ as if it were the sup. "(H2) gets easier with L" is a measured trend for $L \leq 9$, not a result for $L > 11$.

- All values are lower estimates of a sup (grid plus local search).

6.5 One-step values

The arithmetic of Section 3.7.4 shows that $\rho_1^2 < L$ implies (H2). The measured one-step values are as follows.

- $L = 3$: $\rho_1^2 = 3.10 > 3$. The one-step route fails (Section 3.7.2); the two-step map is needed.

- $L = 5$: $\rho_1 = 1.860$ ($j = 1$, Wilson fine kernel, dense, measured by the verifier), 1.9565 ($j = 2$), 1.963 ($j = \infty$). So $\rho_1^2 \leq 3.853 < 5$ and $\rho_1^2/L = 0.771$ [NUMERICAL, CONFIRMED AS A MEASUREMENT, UNCERTIFIED; `H2_LARGE_L_VERIFY` GRADE 4].

- $L = 7$: $\rho_1 \geq 2.032$ (diagonal only), $\rho_1^2/L \approx 0.59$ [NUMERICAL].

- $L = 13$: measured $\sup \rho_1^2 = 4.37$; certified bound $\sup_{j,K} \rho_1^2 \leq 10 < 13$ (Theorem 6.1).

So at $L \geq 5$ the one-step map suffices for Theorem C1, as far as measured. For certification this replaces L^8 blocks per K (two-step map) by L^4 blocks per K (625× fewer at $L = 5$).

6.6 A rigorous point enclosure at $L = 3$ [INTERVAL ENCLOSURE, VERIFIED]

Source: `H2_CERTIFY.md` with the prototype `numerics/h2cert_proto.py`; verifier `H2_CERTIFY_VERIFY.md`. At $L = 3$, $j = \infty$, and the face point $K^* = (2.4035, 2.4035, \pi, \pi)$ near the argmax of Section 6.4, ball arithmetic (python-flint) gives

$$\rho_2(\infty, K^*) \in [1.9993,\ 2.1273] < 3.$$

The width comes from the umklapp tail and scales as N^{-4} in the truncation N. The verifier reran the code at $N = 12$ and obtained [1.5475, 2.5792], which contains the floating-point value 2.063361 and confirms the N^{-4} scaling; at $N = 6$ the code stops with an assertion instead of returning a wrong interval. The verdict is CONFIRMED given the premises P1–P3 of `H2_CERTIFY_VERIFY`; P1 (the loop-family coefficients of the (0.4) row are exact rationals $k/648$, k even) is proved there, and the fidelity of the loop family to Bałaban's paper stays inside P3. Further results of the same two files:

- One-step factorization: $\rho_2(j, K) \le \rho_1(j, K) \cdot \max_l \rho_1(j-1, k_l)$, exact [PROVED+VERIFIED]. So two-step work is needed only where the right side is $\ge L$, near the face diagonal.
- Oishi–Rump bound $\|\mathfrak{B}^{-1}\| \in [42.7,\ 43.6]$ at K^* (bounds CONFIRMED). The values 177 and 195 of Section 3.9 use the full Δ instead of $\frac{1}{4}\Delta$; this factor reconciles them. No $\|\mathfrak{B}^{-1}\|$ enters ρ_2.
- The umklapp tail enclosure is CONFIRMED for $M = \infty$. For finite M it is CORRECTED: its lower bound must be 0, with class-wise factors $(\pi/2)^{12}$ for two or more large components.
- A uniform Lipschitz grid is refuted as a route: $\Lambda_K \approx 2.8 \cdot 10^4$ (CONFIRMED) would need about 10^{13} cells. The replacement is a per-cell invertibility test with a mean-value enclosure.
- Cost: `H2_CERTIFY` estimates about $1.5 \cdot 10^4$ core-hours for a full certification at $L = 3$. The verifier grades this estimate UNSUPPORTED: at the tested sizes the arb overhead dominates, and the box-sum speedup is overstated.

This is one point at one level. It does not certify the supremum over the Brillouin zone or over j. Section 6.7 does this at $L = 13$.

6.7 (H2) at $L = 13$ [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]

Sources: `H1_ANALYTIC.md` §0–§8 (statement, lemmas, majorant); verifier `H1_ANALYTIC_VERIFY.md` V0–V9 with `numerics/h1analytic/bb_arb.py`; operator check `H1_ANALYTIC_VERIFY2.md` with `numerics/h1analytic/indep_H.py`; symmetry check `H2_CERTIFY_P5_VERIFY.md`. Here $\rho_1(j, K) = \|H_j(K)\|/L$ with $H_j(K)$ the one-step gauge-fixed minimizer map of Section 3.7.

Theorem 6.1 ((H2) at $L = 13$)**.** *For $L = 13$, every level $j \ge 1$ including $j = \infty$, and every coarse momentum $K \in \mathrm{BZ}$,*

$$\rho_1(j, K)^2 \le 10 < 13.$$

Hence $\rho_2 \le \sup_{j,K} \rho_1(j, K)^2 \le 10 < L$ by the factorization $\rho_2 \le \rho_1(j) \max_l \rho_1(j-1, k_l)$ (Lean `rho2_le`, Section 3.7.4). That is, (H2) holds at $L = 13$.

Proof sketch. Write $s = L^{-2}$, $k_l = (K + 2\pi l)/L$, $M = L^{j-1}$, d^M for the Prop. A symbol at level M, $P_\perp$ for the projection onto $\hat{K}^\perp$, and T, T_2, σ, $Z = (P_\perp T P_\perp)^+$ as in `H1_ANALYTIC` §1.

1. Exact split. $\rho_1(j,K)^2L^2 = \lambda_{\max}\big(ZT_2Z + cc^\dagger/(\sigma|\hat{K}|^4)\big)$, $c^\dagger = \hat{K}^\dagger(1-TZ)$: a transverse part and an orthogonal pure-gauge part.

2. Three exact lemmas. Lemma S: the averaging symbol is $E_q(k_l) = E_l^0 - \hat{K}g_l^\dagger$, with E_l^0 diagonal (block average times straight path), $|E^0_{l,\mu\mu}|$ built from $a_{l,\rho} = \sin(K_\rho/2)/(L\sin(k_{l,\rho}/2))$, and $\sum_l \prod_\rho a_{l,\rho}^2 = 1$ (Fejér). Tower identity: $d_\mu^{ML}(K) = L^{-2}\sum_l w_{l,\mu} d_\mu^M(k_l)$, hence $P_\perp T P_\perp = P_\perp \operatorname{diag}(d^{ML}(K))P_\perp$ and Z is the next-level kernel of Prop. A. Lemma G: a closed form of the gauge row g_l from the path average, so the averaging operator is explicit.

3. Majorant. Lemma T (transverse): $\lambda_{\max}(ZT_2Z)/L^2 \le \gamma_L F(K)$, $F(K) = 1/(|K|^2\underline{d}(K))$, $\gamma_L = \operatorname{sinc}(\pi/2L)^{-2}$, from $T_2 \le d_{\max}T$, $ZTZ = Z$, $d_{\max} \le L^2\gamma_L/|K|^2$, and $\|Z\| \le 1/\underline{d}$. Lemma Gauge: the gauge part is $\le (1+\tau)^2/(|\hat{K}|^2 S_L)$ with explicit sums V, U and $\tau = (V+|\hat{K}|U)/\underline{d}$. So $\rho_1(j,K)^2 \le B_L(K) := \gamma_L F(K) + (1+\tau)^2/(|\hat{K}|^2 S_L)$. By Cauchy–Schwarz $F \le F_{CS}$, a closed form made of one-dimensional factors in each K_ρ.

4. Uniformity in j. B_L does not depend on j. For every $M \in \{1, L, L^2, \dots, \infty\}$ the fine symbol satisfies $d^M \le 1/|\tilde{q}_0^M|^2 \le 1/|\hat{k}|^2$, because the Fejér weights sum to 1. The coarse symbol d^{ML} dominates the truncated $M = \infty$ sum termwise. No monotonicity in j and no O5 rate is used.

5. Branch-and-bound. B_L is invariant under the 384 signed permutations of K by construction, so the sup over the sorted chamber $0 \le K_1 \le \cdots \le K_4 \le \pi$ is the sup over BZ. The program `numerics/h1analytic/bb_arb.py` encloses $\gamma_L F_{CS} + \bar{G}_L$ on boxes $K = \pi u$ with exact dyadic u, evaluates every one-dimensional value in arb (python-flint), and rounds outward by one ulp after every floating-point operation, with Higham bounds on every sum. Result: $\sup_{\text{chamber}} B_{13} \le 9.9995277$, 25,169 boxes, 8,075 gauge evaluations, 0 failed leaves, 58 s. Boxes that contain $K = 0$ are enclosed directly; $\rho_1(j,0) = 1$.

Margins. The measured $\sup \rho_1^2$ is 4.37. The majorant is tight to 2.6% at $|K| \approx 1.5$ ($\min B_{13}/\rho_1^2 = 1.026$ over 896 cases and a local search). The certificate refines boxes only until each is ≤ 10, so its margin to 10 is $5 \cdot 10^{-5}$; its margin to the threshold 13 is 23%.

Verification. `H1_ANALYTIC_VERIFY` rederives the split, Lemmas S, G, the tower identity, Lemma T, and Lemma Gauge on paper and checks them against the production objects to $\le 5.3 \cdot 10^{-13}$ (V1–V2); it checks $B_{13} \ge \rho_1^2$ at 896 (j,K) cases (V3–V4); it grades the original float program CORRECTED (no directed rounding) and replaces it by `bb_arb.py` (V5–V8). `H1_ANALYTIC_VERIFY2` rebuilds $H_j(K)$ from the definitions (position-space (0.4) paths, sparse KKT, ghost-domain gauge) and reproduces the production ρ_1 to $5.8 \cdot 10^{-9}$ (10^{-14} with the same d symbol); the two parts of the split agree to $3 \cdot 10^{-8}$, and the Lemma S conventions are derived by hand and confirmed to $1.2 \cdot 10^{-13}$. `H2_CERTIFY_P5_VERIFY` proves and checks the invariance of ρ_1 under the 384 signed permutations (P5a, P5b); the analytic certificate does not need it, the box sweep does. Lean: `lean/YmLean/Tower.lean` formalizes the algebraic core of the majorant (tower identity in weighted form, Rayleigh and rank-one bounds, `majorant_core`, `lemmaT_core`); `BlockNorm.rho2_le` is the one-step ⇒ two-step factorization.

Residue. The theorem is certified given the following, which no second source has checked.

- The reading of Bałaban's (0.4) (the path family that defines the averaging operator) and the gauge choice (`NUMERICS` §5). The certificate is a statement about H_j so defined; both implementations use the same reading.

- Prop. A at $M \ge 27$ and $M = \infty$: checked only by a second implementation of the same formula (independent recursion at $M = 3, 9$).

- Lemma G and the bound chain: proved on paper and rederived by the verifier, checked numerically at points only. The Fourier/umklapp derivation, Lemmas S and G, and the identification of the split are not in Lean.
- `bb_arb.py`: audited line by line and checked against the float program on 30 boxes; not formally verified. It trusts python-flint arb and IEEE correctly rounded $+, -, \times, \div, \sqrt{}$.
- Independent cross-check: the box sweep of the cell certificate (Section 6.9) is running; its result is not in yet.

6.8 Round 7: more certified L, the gap and neutral-sector checks, and an open tension

(H2) at $L = 15, 17, 21$ [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]. The program `numerics/h1analytic/bb_arb.py` of Theorem 6.1 was run with target $L - 1$ on the initial grid $n_0 = 16$ of the sorted chamber (3,876 boxes, no refinement). The lemmas behind the majorant B_L are stated for odd $L \geq 3$, and B_L does not depend on j. A full rerun on the Mac covered all boxes, with 0 failed leaves.

L	$\sup_{j,K} \rho_1^2 \leq$	target	$F_{\max}$	$G_{\max}$	rerun time
15	13.737723	14	10.020	4.005	33 s
17	12.826432	16	10.020	3.061	51 s
21	11.776985	20	10.020	1.951	113 s

So $\rho_2 \leq \sup \rho_1^2 < L$ by `rho2_le`, and (H2) holds at $L = 15, 17, 21$, with the same standing and the same residue as at $L = 13$ (Section 6.7). Sources: `numerics/H1_OTHER_L.md`; verifier `H2_CERTIFY_L15_17_21_VERIFY.md`. The verifier reproduced the worst cell, $F_{\max}$, and $G_{\max}$ at each L to the printed digits, checked 9 spot cells, and found $B_L \geq \rho_1^2$ at 5 points of an independent H_j (ratio ≥ 1.63). The full sweep of 896 cases $B_L \geq \rho_1^2$ and the production cross-checks were done at $L = 13$ only; neither affects the proof logic. These values of L are not used by the targets (ledger C3).

The 2D gap ($\mathbf{G'_A}$) [NUMERICAL]. The proof of Theorem 5.12 covers only $|ef| \leq 10^{-4}$. Numerics (`numerics/t2_gaprime2_check.py`; reproduced by the verifier, 141 values) show the gap on all of $\alpha = ef \in (0, 3.02]$: at 11 rational fluxes with denominators up to 2213, the smallest |eigenvalue| of A' is $|\alpha|(1 - 0.125|\alpha|)$, so the ratio to $|\alpha|$ is ≥ 0.95; a sparse shift-invert run at $\alpha = 10^{-4}$ gives the ratio 1.0000125. Round 7 (`SOURCES_T2_FRANGE.md`) shows that (M_j) needs the gap only on a punctured neighbourhood of $f = 0$. So these values are supporting evidence, not proof inputs.

The neutral sector of (M_j) [NUMERICAL, REPRODUCED]. Model: the 2D $N \times N$ torus of `SOURCES_T2_S123.md` §6, Landau gauge, flux $\varphi = 2\pi m/N$ per plaquette, the operator $4\,\mathrm{Hess} + D_r D_r^T$ restricted to the neutral (colour-3) block. Prediction: the spectrum is $\bigcup_K \{l(K), \cos\varphi\, l(K)\}$, with exactly $d = 2$ zero modes (the constant 1-forms). Author's run (`numerics/t2_mjrest_neutral.py`): $N = 12$–48, $m = 0$–2, maximal deviation from the prediction $\leq 4.9 \cdot 10^{-8}$ (the finite-difference floor), two zero modes at every flux, gap $\cos\varphi \cdot 4\sin^2(\pi/N)$ to relative error $\leq 8 \cdot 10^{-8}$. Verifier's run (`numerics/t2_mjrest_verify_neutral.py`): the author's JSON is reproduced bit for bit, and at new sizes

N	m	φ	zero modes	first positive	$\cos\varphi \cdot 4\sin^2(\pi/N)$
36	3	0.5236	2	0.02631374	0.02631374
20	3	0.9425	2	0.05753652	0.05753652
64	2	0.1963	2	0.00944548	0.00944550

The zero modes span the two constant 1-forms (residual $\leq 5.6 \cdot 10^{-7}$), and the charged–neutral leak is 0. Limits: 2D, Wilson level ($t = 0$, no Bałaban averaging), rational fluxes. The level-j neutral symbol is not checked numerically; Theorem 5.14 uses only the bounds (N1), (N2).

Open: the single-step β_1 against $c \log L$. `SOURCES_T2_F_STEPB.md` §3.3 takes the extrapolated single-step values $\beta_1(Lp \to 0) = 0.743, 0.797, 0.827, 0.844$ at $L = 5, 7, 9, 11$. The constants $\beta_1 - c \log L$ are 0.145, 0.074, 0.010, -0.047. The local slopes $d\beta_1/d\log L$ are 0.161 ($5 \to 7$), 0.119 ($7 \to 9$), and 0.084 ($9 \to 11$), and they decrease with L. Fits give slopes 0.129 ($a + s \log L$), 0.052 ($a + s \log L + d/L^2$), and -0.18 ($a + s \log L + d(\log L)^2/L^2$). None is near $c = 0.3715$. The first-step constant falls by about 0.06 per step and does not level off at $L \leq 11$. So the reading of `NUMERICS` §7, "decreasing toward the universal slope", is not supported by these data, and that claim is under verification. Open: whether 0.3715 is compared with the right quantity (a single-step β_1 at each L is a different family from the fixed-L sequence $\Lambda = L^j$ of Theorem 3.1 of `BETA_FP_H.md`), or whether the gap sits in the coarse bracket, in the dependence on the family, or in the $N_g = 8$ grid at $L \geq 9$ (unchecked). This tension is [OPEN]. It bears on ledger B3 and possibly on B4 through Theorem 3.1 of `BETA_FP_H.md`.

6.9 Certification plan and j_0 [DESIGNED; SUPERSEDED AT $L = 13$ BY SECTION 6.7]

At $L = 13$ the analytic route of Section 6.7 replaces this plan: it needs no tail level j_0, no $\|\mathfrak{B}^{-1}\|$ bound, and no Lipschitz constant. The cell certificate below is run at $L = 13$ only as an independent cross-check (box sweep, in progress). `B2_TWOSTEP_VERIFY` grade 6 (CORRECTED in size): certifying (H2) is a computer-assisted proof project, not a check. `H2_LARGE_L` §4 gives the design; `H2_LARGE_L_VERIFY` grades 6–8 check it. Three parts.

(a) Tail $j \geq j_0$. From the resolvent identity, $\rho_2(j) \leq R_\infty + (\rho_1^2/a_*)(\delta_{j-2} + \delta_{j-1})$, where $R_\infty = \rho_2$ at $j = \infty$, $a_* = \inf_K a_j(K)$ with $\|[\mathfrak{B}^{-1}]_{f,f}\| = 1/a_j(K)$ exactly, and $\delta_m = \sup_K \|\Delta_{L^m} - \Delta_\infty\|$ (O5 decay) [derivation CONFIRMED]. Inputs CORRECTED by the verifier: the 2-norm sup of δ_m is at the corner, $\delta = 179, 92.0, 18.7, 2.39, 0.317$ ($M = 1, 3, 9, 27, 81$), 45–90× the assumed value at small m; $a_*(L = 5) = 0.0107$, not 0.05. With these inputs, $j_0 \approx 8$ ($L = 3$) and ≈ 6 ($L = 5$). With the O6 coercivity constant, $j_0 = 25$, which cannot be used in practice (`B2_TWOSTEP_VERIFY` §5 estimated $j \gtrsim 45$ with $\|\mathfrak{B}^{-1}\| \leq b(L) \sim 10^{19}$).

(b) Levels $2 \leq j < j_0$ and $j = \infty$. A certified sup over the BZ of $\|H_{(2),j}(K)\|$ (at $L \geq 5$: of $\|H_j(K)\|$, Section 3.7.4). Plan: interval arithmetic with local mean-value enclosures on boxes in the sorted chamber (1/384 of the BZ), 4×4 and 5×5 verified inverses, one 4×4 λ_{max} per K; about $2.5 \cdot 10^3$ boxes at half-width 0.05 at $L = 3$; the analytic form of Δ near $K = 0$, where $\rho_2(0) = 1$. At $j = \infty$ also a certified tail bound for the Prop. A sum (l_{max} truncation; observed effect $4 \cdot 10^{-4}$). There is no explicit global Lipschitz constant today: the A0/O5 strip width is existential, and the constant that can be made explicit is $\sim 10^{40}$. Monotonicity in j is not proved and is not used.

(c) Inputs to certify first: $\|\mathfrak{B}_j(K)^{-1}\| \leq B_{num}$ for all j, K (the O6 constant $b(L)$ is unusable), the O5 constant C_0 explicitly (or certified δ_m), certified a_*, ρ_1, R_∞.

The run plan (`H2_LARGE_L_VERIFY` grade 8) uses atomic JSON writes and pre-copied profile caches; the profile build time at $L \geq 11$ is not budgeted. At $L = 13$ the cell certificate runs as the

box sweep (`numerics/h1cert_cell2.py`, sorted chamber, valid by P5), an independent cross-check of Theorem 6.1; its result is not in yet.

7 What remains

This section lists every open item of the paper in one place. The items are grouped by the kind of work they need: certification, fixed-level and structural inputs of Theorem 2w, items of the sources block, and verification. The item labels H, 1–13, and B1–B8 are those of the dependency table (Section 4.6). For Theorem 2w ⇒ Theorem 1 the list covers §10 of `THEOREM2_PROOF_v4.md`. The trust ledger (Section 7.7, `TRUST_LEDGER.md`) supersedes this list as the complete record of what the two main theorems take on trust: Bałaban-level statements, external results, our open items, and conventions.

7.1 Certification

Single uniform hypothesis (both routes):

H. (H2) $\rho_2 = \sup_{j\geq 2} \sup_{K \text{ real}} \|(\oplus_l H_{j-1}(k_l))H_j(K)\|/L^2 < L$. [DONE AT $L = 13$: CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)], Theorem 6.1: $\rho_1(j,K)^2 \leq 10 < 13$ for all $j \geq 1$ (including ∞) and all K, hence $\rho_2 \leq 10 < 13$ by `rho2_le`. Route K: (H2) ⇒ ($C1\sigma$) ⇒ (B2) [PROVED+VERIFIED, `B2_TWOSTEP_VERIFY` GRADE 3]. Route B: (H2) + (T1) ⇒ (U) [PROVED, UNVERIFIED, `BETA_FP_U` THEOREM 2.4]. Cross-check in progress: the box sweep of the cell certificate at $L = 13$ (Section 6.9). Round 7: (H2) is also certified at $L = 15, 17, 21$ (full reruns, verified; Section 6.8). At $L = 3, 5, 9$ (H2) stays [NUMERICAL].

H′. Residue of Theorem 6.1 (Section 6.7): the reading of Bałaban's (0.4) and the gauge choice (single source); Prop. A at $M \geq 27$ and $M = \infty$ (second implementation of the same formula only); Lemma G and the bound chain (point checks only beyond the paper proof); `bb_arb.py` audited, not formally verified.

At $L = 3$ one point is enclosed: $j = \infty$, $K^* = (2.4035, 2.4035, \pi, \pi)$, $\rho_2 \in [1.9993, 2.1273]$ [INTERVAL ENCLOSURE, VERIFIED] (Section 6.6). Certified early steps $\beta_1(0), \ldots, \beta_{j_*}(0) > 0$ at odd $L > 11$ are needed only for literal (0.31) at every K (Section 3.9.2).

7.2 Fixed-level and structural inputs of Theorem 2w

Route K hypothesis set (all must hold):

1. (B2) $\sup_j \sup_K |\sigma_j^{(2)}| < \infty$ (or (B2w)). Reduced to H [VERIFIED REDUCTION]. Older reductions kept: ($C1\sigma$) + (h) [PROVED+VERIFIED]; (H″) [ARGUED]. (HS) is refuted.

2. (PI) probe independence, at each fixed level [VERIFIED REDUCTION] (`SOURCES_T2_PI.md` + VERIFY), given (I1) massiveness at $U = 1$, (I2) gauge covariance, and (I3).

3. (F) lattice = continuum log coefficient at constant field [CITED SOURCE CHECKED (SAVVIDY [22], LEDGER X5); LATTICE TRANSFER NOT PROVED]. Round 7: Step B is [REDUCED [R], UNVERIFIED] to (L-LL), a kernel bound, and the gauge-fixing part of Step C (Section 5.11). Open numerical tension: the single-step β_1 at $L = 5$–11 does not follow $c \log L$ (Section 6.8). This is the only Group B row of Theorem B without a proof or a verified reduction.

4. (M_j) formal = true f^2 coefficient at each fixed level: (S1) $\mathcal{K}_t(f) \geq -C_1|f|$, (S2) density of states $\leq C_2|f|$ near 0, (S3) relative bound of the level difference, all on the lattice: [VERIFIED REDUCTION] (`SOURCES_T2_S123.md` + VERIFY, 25 confirmed, 2 corrected): (S1), (S3), (S2)$|_{t=0}$, (S2c) are proved at fixed j given the Peierls input (I3). Round 5 proves (I3) at every step, with constants independent of the step (`SOURCES_T2_I3.md` + VERIFY, 15 confirmed, 2 corrected; given the identification of the one-step objects with the linearized averaging). Round 6 proves (G'_A) for $0 < |f| \leq ((1-c)/100)^2/2$, gap $2c|f|$ (Theorem 5.12) [PROVED+VERIFIED]. So (M_j) reduces to (I3), the region-1 step of Prop. 4.4 [ARGUED], and the neutral sector [I]. The proved f-range is narrow. Round 7: (M_j) is [PROVED RELATIVE TO LISTED INPUTS; VERIFIED] at fixed j (Theorem 5.14); region 1 and the neutral sector are proved, only the $o(f^2)$ form, and the proved f-range suffices (`SOURCES_T2_FRANGE.md`).

5. V1 order-g^0 completeness of CMP 109 (1.3)–(1.5): $\beta_{j+1}(0)$ is the Maxwell coefficient of the per-step Gaussian determinant [PROVED+VERIFIED AT FIXED LEVEL, CONDITIONAL ON (G2), BAŁABAN'S INDUCTIVE BOUNDS, AND THE DECLARATION THAT THE AVERAGING IS (0.4)] (`SOURCES_T2_V1.md` + VERIFY). It is history-free.

6. (G2) per-cell infinite-volume densities [VERIFIED REDUCTION] at fixed j (`SOURCES_T2_G2.md` + VERIFY; was [FORMAL]): the torus densities converge exponentially, and the p^2- and f^2-coefficients commute with $N \to \infty$, modulo the surjectivity of $\tilde{Q}$, (Gap_f), and the analyticity reading of the averaging. The j-uniform version is not proved.

Route B hypothesis set (alternative to 1–5; per `BETA_FP_U` §4; all [PROVED, UNVERIFIED] as implications, no verifier file for `BETA_FP_U`):

B1. (T1) $\sup_j \sup_{q \text{ real}} |c^{(1)}_{T,j}(q)| < \infty$, the plane-wave form of Theorem A3 [ARGUED; THE LANDAU-TO-SYMMETRIC GAUGE CONVERSION AND STEPS 2–5 OF A3 FOR (2.1) ARE NOT WRITTEN].

B2′. (H2) = item H.

B3. (DF) $\sup_k |\mathfrak{d}_F(\theta_*; L^k)| < \infty$, the gauge-fixing difference at the plane wave [HYPOTHESIS, ARGUED]. With the exact split $\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$ (`BETA_FP_U` Prop. 3.1) there is no Y-defect; all non-stationarity sits in $\mathfrak{d}_F$. This replaces the earlier $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ items for this route (Section 3.6.2).

B4. Lüscher–Weisz (5.7) [CITED].

B5. A6 (the label of v4; not ledger A6): the plane-wave increment equals Balaban's β up to the expansion point, uniformly in k [ARGUED]. A uniform-in-k input, of the same kind as (DF).

B6. (H0) = item 7′ (Lemma D″ (i)).

B7. $\theta_* < 1$, so that $c_2 \ln(1/\theta_*^2) \geq 0$ in C_B [TRIVIAL CONDITION ON THE CHOICE OF θ_*].

Route B also needs 6 ((G2)), item 7′, and the items 8 and 10–13 (item 9 is closed). It does not need (PI), (M_j), V1; it needs (DF) and A6 instead.

Common to both routes:

7′. The own-slot bound $|\beta_{k+1}(h,g) - \beta_{k+1}(0)| \leq C_\beta g$ on $]0,\gamma]^k \times [0,\gamma]$, for every past h, monotone or not; the lift [PROVED+VERIFIED] (`SOURCES_GGAIN_NONMONO_VERIFY`), 7′ itself conditional on (H0) or on the (G-gain) inputs. Item 7 asked for (H0), Bałaban's smoothness in g with bounds uniform in j and history (CMP 109 inductive assumption at odd $L > 11$; "uniform in j" is our reading of p. 264, not printed, `R2_VERIFY` row 12; uniformity in the

history is not printed; ledger A5). The shooting evaluates β on clamped, non-monotone pasts, and `SOURCES_T2_KUNIF_VERIFY` found that (G-gain) proves C_β only on monotone pasts. `SOURCES_GGAIN_NONMONO.md` lifts the restriction with no change of constant: the only use of monotonicity is a comparison of bound sizes in Lemma F2′, and the running maximum $G_k = \max_{i \le k} g_i$ closes the induction for every past. So 7′ follows from (H0) or from (G-gain) with this lift. The proof uses the Lipschitz form on $]0, \gamma]$ only. Round 5: the use of (H0) here is a [VERIFIED REDUCTION] (Proposition H0-red, `SOURCES_T2_H0.md` + VERIFY) with residue N1–N9; N2 is Lemma G3, proved relative to listed inputs (Theorem 5.9); the rest of the residue (N5 [R], N6, N8 [I], N9 definition only, N7 = B7) is not proved, so A5 stays a ledger row.

8. L5 history dependence. Round 7: [PROVED RELATIVE TO LISTED INPUTS; VERIFIED] by Route 2, nested intervals on true trajectories with no clamp (Theorem 5.15); (a)–(c) below are kept as the history of the item. (a) The clamp proof of Lemmas D′/D″ is valid only if $\beta_{k+1}(g_0, \dots, g_k)$ is defined and continuous on the whole cube $]0, \gamma]^{k+1}$; Balaban does not supply this. (b) Balaban's β_{k+1} exists only on histories where his induction has run; that domain must carry (c′) + (S), not only $g_i \in]0, \gamma]$. Clamped histories can violate the growth side of (c′). (c) True trajectories with all $u_i \ge \gamma^{-2}$ inherit (c′) from (a″). A nested-interval variant of Lemma D″ on these trajectories would avoid the gap [NOT WRITTEN]; the variant is written only for Lemma D (`LEMMAS_DE_VERIFY` §1.5). Lean covers one-argument β only (Lemmas D′, D″, D, E).

9. K-independence of $\beta_j(g)$ [CLOSED AS A REDUCTION; VERIFIED] (`SOURCES_T2_KUNIF.md` + VERIFY). Item 9 is a volume statement, not a history identity: K enters β_j only through the torus size. CMP 119 (2.24), p. 249, and p. 279 define β_j as the whole-lattice coefficient, so the β_j of the recursion is K-free by definition. Lemmas D′/D″ run at one fixed K and never compare trajectories. The window bound needs only (K1) (history-free at zero own coupling), (K2) C_β (item 7′), and (K3) the window bound (5.1) of Theorem K. The residual volume question (the finite-volume step equals the whole-lattice one, ledger A15, B8) is an input of Theorem 1, not of the shooting. Round 5 narrows it: given (Loc), the torus coefficient is within $C_j N_j^6 e^{-m_j N_j/2}$ of the whole-lattice one (Proposition 5.8); exact equality cannot be derived in general. Round 6 proves (Loc) (Theorem 5.10), so this bound no longer assumes it, and proves the boundary reading and the replaced B8 under (Vol) (Proposition 5.11).

11. Odd L. All one-loop proofs need odd $L \ge 3$ (centered blocks: U3 Lemma 1.2, O5/Prop. A, evenness, Lemma 4.1 of LEMMA43). Balaban assumes L odd > 11 throughout CMP 109, so (H0) and $\beta_j(g)$ at $g > 0$ exist only there. Theorem 2w is stated for odd $L > 11$, or with (H0) at this L as a hypothesis. Numerics: $\beta_1(0)$ up to $L = 11$ (Section 6.1); (H2) at $L = 3, 5$ (scans), 7 (face diagonal), 9 (one point); ρ_1 at $L = 3, \dots, 13$. Only $L = 13$ is inside Bałaban's range; there (H2) is certified (Theorem 6.1).

12. SU(2) only. The charged/neutral split of Lemma 5.1 and (W) are SU(2) statements; $c = 11N^2/(12\pi^2)$ is general but the chain is not checked for $N > 2$.

7.3 Items of the sources block

These items stand between the trajectory of Section 3 and the bound (H-U) of Section 4.2. Tags and files are those of the dependency table.

- (B-S) The sourced version of Bałaban's Theorem 1, (E14-a, b, c), is [OPEN] as a whole.

- (H-hist) via (G-gain). The small-field step is done: pointwise Cauchy and uniform C_β on $]0,\gamma]$ [PROVED+VERIFIED]. Lemma G3 (the χ_k surface term) holds given (Q). Round 5 proves G3 relative to listed inputs, with (Q) assembled from A7 and A8 in the joint (X, B) form (Theorem 5.9); its conditions are C25 and C26, and G_Γ is bounded uniformly in M (verified). The large-field half is a [VERIFIED REDUCTION] (`SOURCES_GGAIN_LF.md` + VERIFY): one Cauchy step on the large-field disc, with the factor $1/g$ absorbed by the exponent of p_0 (CMP 119 p. 244). The real uniform (H-hist) is therefore reduced [R] in both halves; open inputs: the margin (M_r), CMP 102 (189), and the (M-B) bookkeeping; the small-field (Q)/c_- is discharged relative to the inputs of Theorem 5.9. Lemma G4 is off-chain (`SOURCES_G4.md`). Second derivatives: (H-hist2).

- Bałaban-level core (round 3). The propagator and covariance inputs A1–A4 of the ledger reduce to published theorems:

  - A1, kernel-decay form: [VERIFIED REDUCTION] via Lemma CT to the printed CMP 99 Theorems 3.1, 3.3, and the G part of 3.12, with (KE′) proved by citation (CMP 95 (1.11)/(1.18), CMP 96 (2.35)), (H_Q) proved, and S2 closed for G. Residue: a stated torus-side hypothesis (side $\geq c \cdot RM$ coarsest blocks), two unprinted conventions, and the reading $a = 1$ of Theorem 3.1 ([I]); S2 for G_1 is closed (`SOURCES_S2_G1_V7.md` + VERIFY).
  - A1w, walk-expansion form of A1 (CMP 99 Theorem 3.15, "analyzed in the same way"): [VERIFIED REDUCTION], now written out in full. Theorem 5.1 (`A1_WALK_FULL.md`) gives the walk expansion of $(QGQ^*)^{-1}$ with (3.108)-form bounds at rate $\frac{1}{2}\varepsilon$, the Z_0-compressed variant (O3), and the complex α'-space (O4). Its input (W_G) for the final propagators is Theorem 5.2 (`WG_FULL.md`): the walk expansion of G and the tree expansion of G_1, built from Bałaban's printed proof sketches of CMP 99 Thms 3.7, 3.9, 3.10, the Neumann series (3.130), and CMP 96 Lemma 2.1. Both are paper-grade relative to their tagged inputs. What remains: the lower bound (C) (the residue of A1), the three printed sketches [R], (W_G) at complex points [R], the condition WG-1 in the M_1 form $\alpha_0 \leq c' M^{-1} M_1^{-10}$ ($c'' M^{-1} M_1^{-12}$ for G_1), admissible in Bałaban's order, and geometry and convention items [I] (cube geometry, Lipschitz partition, Q-locality, $C^{(2)}$-locality, the scale bookkeeping C0, the M_1-cover, κ_1 independent of M).
  - A2 (R_2, R_3 of CMP 116 (2.16)): [VERIFIED REDUCTION] (`SOURCES_A2_R2.md`, `SOURCES_S2_G1_V7.md` + VERIFY): R_2 by a Neumann series around the compression with the complex-U repair V7, to the complex walk bounds (W^c) on the α'-space, now Theorem 5.3 (`WC_FULL.md`, paper-grade relative to its tagged inputs): in the exponential gauge, $A_Z \geq \frac{1}{4}\gamma_e$ at real U, the walk expansion of $C^{(k)}(Z_0)$ in the CMP 116 p. 13 form, $\operatorname{Re} A_Z \geq \frac{1}{2}\gamma_{\exp}$ on the α'-space, and the resolvent bound with per-walk prefactor $4/(\gamma_{\exp} + x)$, with no use of Thm 3.15; its inputs are Theorem 5.2(G1), (C), the inputs of A3, CMP 109 p. 254 [R], and [I] items; the decay of $C^* \Delta_k C$ at real U (A7), the boundary-contact reading (item (i), [I-forced]), and $\operatorname{dist}(Z_0, \Lambda_k^c) > RM$ [I]; R_3 in weaker forms sufficient for (2.21).
  - A3 ($C^* \Delta_k C \geq \gamma_0$): full proof, verified (`SOURCES_A3_GAMMA0_FULL.md` + VERIFY), from the printed CMP 99 Theorems 3.11–3.12, (H_Q) [PROVED+VERIFIED], R-Δ2 [R], and items R3–R5: $\gamma_0 \geq m_L/(48 d^2 L^d)$ with $m_L = \min(2(1 - \cos(\pi/L)), 1/L)$, uniform in k, volume, and regular U, for $M \geq L/2$ (this equals $(48d^2)^{-1} L^{-d-1}$ for $L \leq 9$ and is 0.7555 times it at $L = 13$); the σ-localized clause reduces to (W) + (N), without symmetry.

The proof step inside CMP 96 (2.126)–(2.127) is false for $L \geq 10$ (a lost factor L); the statement of Lemma 2.4 stands with the corrected constant.

- A4 (CMP 102 (189)): [VERIFIED REDUCTION] to A1 + published CMP 102 statements + A18 + one bookkeeping item.
- Round 5 (Section 5.9). A7: proved relative to listed inputs (Theorem 5.4); C is the constraint map and $C^*\Delta_k C$ the precision; the norm bound by Lemma N from C2 of `WC_FULL.md`, A3-R4, I1, I3; the decay from W_{G_1}, Theorem 5.1, I2, I9; complex s at proof-sketch level. A8: proved in corrected form (Theorem 5.5); Bałaban's symmetry claim is not proved and not needed; the majorant uses $H_0 = \operatorname{sym} C_0^{-1}$; Lemma D with $e^{-\varepsilon'' M/8}$; readings I-a–I-e. A10: proved in the corrected form A10′ (Theorem 5.6); the raw Taylor terms do not obey (1.24); the transformed terms do, after the exact marginal cancellation; residue R1–R4; per-label exponent β_H. A11: closed as a verified list (Proposition 5.7). A15: narrowed (Proposition 5.8); (Loc) for $g > 0$, the $\partial\Lambda_{k+1}$ reading, and the R-term part (B8) stay asserted. A5: its use in Theorem 2w is a verified reduction, with N2 = Lemma G3 proved (Theorem 5.9).
- Round 6 (Section 5.10). A15: (Loc) proved exact for non-wrapping polymers relative to C32–C36 (Theorem 5.10); the $\partial\Lambda_{k+1}$ reading (f) proved exact relative to C37 (Proposition 5.11); threshold $N_k > 22M$, $\varepsilon(M) = C_j(22M)^6 e^{-11 m_j M}$. A15 is a verified reduction to conventions. A5: the $\mathbb{Z}^d$-limit assertion now follows, so A5 has no bare assertion left. B8: ill-posed as stated; replaced by (a)–(c), proved relative to listed inputs and (Vol).
- Round 7 (Section 5.11). A16 proved relative to listed inputs (Theorem 5.20; the first proof's locality step was false). A17 proved relative to the printed text and one reading (Proposition 5.16). A18 (Proposition 5.18), A19 (Proposition 5.17), and A23 (Proposition 5.19) are verified reductions. Conditions C38–C42.

So the Bałaban-level core now rests on published propagator theorems, Bałaban's printed proof sketches of CMP 99 Thms 3.7, 3.9, 3.10 (through Theorems 5.1, 5.2, 5.3), the lower bound (C), the inputs of Theorems 5.4, 5.5, 5.6, A18, R-Δ2, two [I] items of A2, the geometry and convention items of the three write-outs, the two unprinted conventions, the torus-side hypothesis, and, since round 6, the locality conventions C32–C37 and (Vol) (C31). This is a verified reduction, not a proof.

- (H-hist2), on which the threshold-shell bound (H-thr) depends: [VERIFIED REDUCTION] to the real uniform (H-hist), no new input; with it the M3 history weights and the frozen-integer convention (L3).

- (FX). O1(b), O2, and R2 are [PROVED+VERIFIED]; O3 is [PROVED+VERIFIED, CONDITIONAL ON N3]; O4 is avoided by Bałaban's main form (1.98). N3, the HBETA strip bound $|\delta w| \leq \frac{3}{2}\tau$, is reduced to (H-hol) + real (H-hist) + (H2-hist) (condition (a) settled); (H-hol) is a verified reduction to (M-A) (via (H−) and CMP 102 (189)), (M-B) (verified), and (M-C). (M-C) is a [VERIFIED REDUCTION] (`SOURCES_MC.md` + VERIFY) to the printed conditions of CMP 116 §2 (B1–B23 and four more), Lemma 2, (1.36), and p. 12 (2.5)–(2.6), with shrinks $\varepsilon_1/(4\lambda_V)$, $\gamma/(8\lambda_V)$. Its residual inputs are now settled: (I-V0) is closed [PROVED+VERIFIED] (`SOURCES_MC_IV0.md`); (I-ε) is closed [PROVED+VERIFIED] for CMP 109/116 (`SOURCES_T2_IEPS.md`) and for the large-field layer of CMP 119/122 (`SOURCES_T2_IEPS_LF.md`; residue [I]: the unprinted scaling of α_2–α_4 and q_0 versus p_0, each branch closes); (I-ord) is closed; (M1) is the convention C24 (`SOURCES_HIGHER_M1.md` + VERIFY). No consumer needs coupling or threshold derivatives of order greater than

2. The two minor inputs of R2 ($R_{K+1} \geq L$, C_c independent of g) are verified readings (`SOURCES_BOOKKEEPING.md`). (FX2) is conditional on O3, so also on N3.

- (B-S.D2C, B-S.Hβ) The domain hypothesis (D), and the conditions (I) and (H-hist) of the two-loop bound.

- (B-S.D2, B-S.D3) The recovery of the decay loss $\kappa \to \kappa/2$ [VERIFIED REDUCTION] by $L/2$-rescaling, $\kappa \geq 1536$ at $d = 4$ (ledger B22; it uses the unprinted large-field T-step rate loss, ledger A23, now a verified reduction with loss $\leq 13\delta$, so (K1′) and $\delta = 0.025$ at $L = 13$). The scale condition (U) ($\rho_1 \leq c_* r_0/M^a$ at the unit scale) is a constant choice (ledger C23), verified compatible with (B-N1) (`SOURCES_UN1.md`), and does not appear in Theorem B.

- (B-N1) The bridge from the source form to (H-U) derived given (S) (`ASSEMBLY_N1.md`, confirmed); the uniformity of (S) in support size and in ε is a [VERIFIED REDUCTION] (`SOURCES_B21_S.md` + VERIFY): every factor of $C^n(n!)^{1+\eta}$ is independent of h, given Lemma C, S1 at all scales, (FX) in the main form, and the coarsest-scale (FX2) radius; open: the (E14-c) clause. Given (FX), the step (S_{src}) $\Rightarrow$ (H-U) with $s = 6$, $D > 1$ is confirmed.

- (NT) = (E14-f), for nontriviality (Theorem 4.3); at every layer for part (ii). The constant $c > 0$ must be computed in Bałaban's normalization.

- E13, full-sequence convergence: not needed for Theorem B; [OPEN].

7.4 Verification debt

10. Balaban readings: the sweep (Section 3.4.3) is grep plus page images, not line-by-line, with 4 of 19 rows checked by a second reader; "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) is closed as a verified list (Proposition 5.7, ledger A11); "β_0 can be chosen arbitrarily small" is on CMP 119 p. 255, and β_0 is fixed before γ, as used (verified reading, `SOURCES_BOOKKEEPING.md`; former ledger A12); the readings of `BALABAN_EXCERPTS_CHECK` item 11 (Π (4.37)/(5.1), CMP 119 (2.42), (2.12)/(2.23), the "L odd" page) are verified, with corrections, in `SOURCES_BOOKKEEPING.md` (former ledger A22); the CMP 119 Sect. 3 residue is the unprinted T-step rate loss (ledger A23); Theorem 1 itself rests on Balaban's published proof, which uses the Group A items of Section 7.7.

B8. Verification: `BETA_FP_U` has no verifier file. `BETA_FP_PW_VERIFY` §6 items 1–4 grade the logic of Theorem B [PROVED+VERIFIED, CONDITIONAL] (tag decision, `PLAN.md`); the inputs (U), (DF), (T1) stay [PROVED, UNVERIFIED].

13. Lean formalization of the history form of Lemmas D′/D″ [OPEN]. Lemma D″ in the one-argument form is done (Section 3.4.2).

- Results tagged [PROVED, UNVERIFIED] need a verifier pass; Appendix B lists them. They include the `BETA_FP_U` reductions and H2_CERTIFY_L13. Round 7 cleared the sweep, `BETA_FP_M` Prop. 4.3 (for $j \geq j_1$), the Prop. 3.6 assembly, and N3 in the D″ form (ledger B10, Section 5.11); this also covers the sweep in the step Theorem 2w $\Rightarrow$ Theorem 1. H2_CERTIFY_L13 (its cell plan is superseded at $L = 13$ by Theorem 6.1; the box sweep runs as a cross-check). H2_CERTIFY_P4 has partial support: identity (D) is proved in Lean (`P4_divergence`), Lemma S is rederived in `H1_ANALYTIC_VERIFY.md` V1, and the rebuilt H_j of `H1_ANALYTIC_VERIFY2.md` confirms it numerically; the Fourier step has no independent verifier.

- The paper is not verified as one document. Its statements carry the tags of the project files.

7.5 Open for stronger statements

Not needed for Theorem 1, open for the stronger statements: (A) in full and Lemma 5.1 for $m = 2$ (convergence, rate); C^2, C^3 of $\sigma^{(1)}$ under (H2) (not written); certified $\beta_1(0), \dots, \beta_{j_*}(0) > 0$ at odd $L > 11$ (literal (0.31) at every K); explicit j_*; (N).

7.6 Why the mass gap is not on this route

Write β_W for the coefficient of $\sum_p (1 - \frac{1}{N} \operatorname{Re} \operatorname{Tr} U_p)$ in the Wilson action. Bałaban's action is $g^{-2} \sum_p (1 - \operatorname{Re} \operatorname{tr} U_p)$ with the normalized trace (CMP 109 (0.2)), so $\beta_W = 1/g_B^2$. Shen, Zhu, and Zhu [24] prove a mass gap for SU(N) lattice Yang–Mills with action $N\beta \operatorname{Re} \sum_p \operatorname{Tr} Q_p$ (their (1.2)) when $|\beta| < 1/(16(d-1))$ (their (1.3)); with $\beta_W = N^2\beta$, $N = 2$, and $d = 4$, this is $\beta_W < 1/12$. Bałaban's construction requires $-2(1+\beta_0)^{-1} p_0(g_k) + 1 + 2\kappa(100 R_k)^d \le -\frac{3}{2} p_0(g_k)$ (CMP 122 II p. 389). With $d = 4$ and $R_K \ge 1$, this gives $p_0(g_K) > 4\kappa \cdot 10^8 + 2$ for every $\beta_0 > 0$ (twice this at Bałaban's $\beta_0 = 1/7$). The unit-scale threshold $g_K p_0(g_K)$ must be below 2, the largest value of $\|U - 1\|$ on SU(2); with $\kappa \ge 1$ this gives $\beta_W = 1/g_K^2 > p_0(g_K)^2/4 > 4 \times 10^{16}$ at the unit scale. The two regimes are therefore separated by at least seventeen orders of magnitude in β_W, and no choice of L narrows this, because L does not enter the condition. A proof of a mass gap needs a mechanism outside the renormalization-group construction, and Bałaban's papers do not address it.

7.7 Trust ledger

The file `TRUST_LEDGER.md`, corrected by `TRUST_LEDGER_VERIFY.md` (Steps 1–28), lists every statement that the two main theorems use and that this project does not prove with a verifier's confirmation. The three tables below condense it to one row per item. Group A holds Bałaban-level statements and external results (X), Group B holds our own open items, and Group C holds conventions and choices. The IDs are those of the ledger. Ledger B1–B22 are not the Route B items B1–B7 or item B8 of this section, and ledger A6 is not the item "A6" of Route B (that label comes from v4 §4.2a). Sources are cited by paper and page; "[I]" marks an inference step of ours that no verifier proved.

Group A: Bałaban-level statements and external results

ID	statement	source	status
A1	Kernel bounds on $(QGQ^*)^{-1}$, $(QG_1Q^*)^{-1}$, uniform in k and regular U	CMP 99 p. 422 (3.132)	asserted, "not repeated"; kernel-decay form a verified reduction via Lemma CT to CMP 99 Thms 3.1, 3.3, 3.12 + (KE′) + S2; residue: torus-side hypothesis, two conventions, the reading $a = 1$ (G_1 clause closed)

Group A: Bałaban-level statements and external results

ID	statement	source	status
A1w	Walk-expansion form of A1 (builds the walk expansion of $C^{(k)}(Z_0)$, bounds (3.107)–(3.108))	CMP 99 Thm 3.15, p. 416	asserted, "analyzed in the same way"; verified reduction, written out as Theorems 5.1 and 5.2 (paper-grade): rests on (C), CMP 99 Thms 3.7, 3.9, 3.10 (printed sketches), CMP 96 Lemma 2.1, WG-1 ($\alpha_0 \le c' M^{-1} M_1^{-10}$), and [I] geometry/convention items; O3, O4 closed
A2	R_2 and R_3 obey (2.16)	CMP 116 p. 16	asserted, "similarly"; verified reduction (V7 repair) to (W^c) = Theorem 5.3 (paper-grade, no Thm 3.15 transfer) + A7 at real U + item (i) [I-forced] + RM distance [I]; R_3 in weaker forms
A3	$C^* \Delta_k C \ge \gamma_0 > 0$, uniform in k and U	CMP 99 p. 428	asserted, "we can prove"; full proof, verified, from CMP 99 Thms 3.11–3.12 + (H_Q) + R-Δ2 [R]; $\gamma_0 \ge m_L/(48 d^2 L^d)$; σ-localized clause to (W) + (N); CMP 96 (2.126)–(2.127) proof step false for $L \ge 10$
A4	Kernel decay (189) of $\delta^2 V / \delta A'^2$ and its consequence (190)	CMP 102 p. 308, Prop. 9	announced; verified reduction to A1 + published CMP 102 + A18 + one bookkeeping item
A5	(H0): $E^{(j)}$, β_{j+1} smooth in g with bounded derivatives; includes the assertion that the whole-lattice limit of (1.22) exists	CMP 109 pp. 263–264	inductive assumption; Theorem 2w uses only item $7'$ (Lipschitz form on $]0, \gamma]$, every past), which also follows from (G-gain) with the running-maximum lift [verified]; round 5: that use is a verified reduction (H0-red), residue N1–N9, N2 = Lemma G3 proved relative to listed inputs (Theorem 5.9); round 6: the $\mathbb{Z}^d$-limit assertion follows from Proposition 5.8 and Theorem 5.10, so no bare assertion is left; for Theorem 2w only the H0-red residue remains
A6	$E^{(j)}$, β_j analytic in the couplings	CMP 109 pp. 264, 266	asserted; not used

Group A: Bałaban-level statements and external results

ID	statement	source	status
A7	$\|C^*\Delta_k C\| \le \gamma_1$ and uniform decay of $C^*\Delta_k C$	CMP 116 p. 13 (2.7); CMP 99 p. 428	premise; round 5: proved relative to listed inputs, verified (Theorem 5.4): C = constraint map, $C^*\Delta_k C$ = precision; inputs C2 [C + one I step], A3-R4, I1, I3 (norm), W_{G_1} [R], A1-W [C], I2, I9 (decay); complex s at sketch level; old A1 + A18 route superseded
A8	Positive Gaussian measure for the real pair $(U, 0)$; complex case "perturbative"	CMP 116 p. 15	asserted; round 5: proved in corrected form relative to Theorem 5.3 and readings I-a–I-e, verified (Theorem 5.5); symmetry of $C^{(k)}(Z_0, 0)$ not proved and not needed ($H_0 = \operatorname{sym} C_0^{-1}$); Lemma D at $e^{-\varepsilon'' M/8}$
A10	Lower-order bracket terms "bounded similarly as in (1.24)"	CMP 116 pp. 8–9	asserted, "similarly"; round 5: proved in corrected form A10′ relative to CMP 109 Sect. 3–5, verified (Theorem 5.6); raw Taylor terms do not obey (1.24) (the j-sum diverges); transformed terms do after the marginal cancellation; residue R1–R4; exponent β_H (C27)
A11	"Similar inequalities hold for other constants"	CMP 119 p. 256	asserted; round 5: closed as a verified list (Proposition 5.7); all follow from (2.6)–(2.9); order entries C28–C30
A13	Theorem 1	CMP 122 II p. 355; CMP 116, 119, 122 I–II	proved, conditional on $g_k \in]0, \gamma]$; uses A1–A4, A1w, A7, A8, A10, A14, A15, A17, A18
A14	Induction closure recovers E_0	CMP 109 p. 280	asserted, constant unspecified; inside A13 only

Group A: Bałaban-level statements and external results

ID	statement	source	status
A15	Small-field terms and β_j are whole-lattice quantities, independent of Λ_{k+1}	CMP 119 p. 278, (2.24), p. 249; CMP 109 (4.35)–(4.37)	asserted, small-field part only; round 6: verified reduction to conventions: (Loc) proved exact relative to C32–C36 (Theorem 5.10), so Proposition 5.8 needs only (Bd), (H), (Cnt); (f) proved exact relative to C37 (Proposition 5.11); threshold $N_k > 22M$, $\varepsilon(M) = C_j(22M)^6 e^{-11 m_j M}$; exact equality only term by term for non-wrapping X; R-term part is B8; not in the set of Theorem 2w
A16	Propagators may be replaced by operators with the same bounds	CMP 116 p. 3	asserted for (1.3)–(1.4); round 7: for C_f proved relative to listed inputs, verified (Theorem 5.20; tree form of $\mathcal{W}_S$; the first proof's locality step was false); residue four [I]: (M1) completeness, (S_H), the Ω-setting volume convention, the s-version of $(C_f)^{1/2}$; conditions C40
A17	$\operatorname{supp}\chi_k$ lies in the analyticity domain of (1.34)	used in CMP 116 Lemma 1	implicit; inside A13; round 7: proved relative to the printed text and one reading, verified (Proposition 5.16): $B' = g_k B$ on every cut bond, $r = 1$; C9 no longer needed
A18	Backgrounds lie in the regular class of CMP 99/102; C-maps agree	used in CMP 109, 116, 122 I–II	implicit; round 7: verified reduction with corrections (Proposition 5.18): inclusions, Lemma Q, Lemma C, $c_J = O(\alpha_0)$; cube patching replaced, $M \geq 5M_{99}$ (C41); residue [R] CMP 102 Thm 1, CMP 98, CMP 99 (3.133), [I] I2, I3; absorbs (C2) of MA2
A19	(190) at complex B	CMP 102 Prop. 9 (silent)	implicit; round 7: reduced to three readings [I], verified (Proposition 5.17): complex (189), (73), and the (172) ball; fallback without the ball reading: two constants, $\omega = 0.4097$, half ball, rate $\times 0.59$

Group A: Bałaban-level statements and external results

ID	statement	source	status
A21	Uniform-in-k moments of the free extension kernel E_0	not printed	proved+verified (`SOURCES_T2_EUMOM`; constants depend on d only); left the ledger in round 4; row kept for reference
A23	Large-field T-step rate loss	CMP 119 Sect. 3 (defers to CMP 109 Sects. 6–7 = CMP 116 Sects. 1–2)	not printed; round 7: verified reduction (proof sketch, Proposition 5.19): loss $\leq 13\delta$; open R1 (count sketch), R3 ($H'(Z_0)$ obeys (2.38)); used by B22 through (K1′)
X1	Lüscher–Weisz (5.7) one-loop coefficient	Lüscher–Weisz [16]	cited; Route B only
X2	OS II growth condition and reconstruction from torus data	OS II [20]	cited
X3	Reflection positivity of the Wilson action, link planes	Osterwalder–Seiler [21]	cited, text not fetched
X4	Convergence criterion of the final cluster expansion	Cammarota, CMP 85 (1982)	cited
X5	(F): continuum log coefficient at constant field	Savvidy [22]	cited, source checked

Group B: our own open items

ID	statement	source	status
B1	(H2) $\rho_2 < L$	none	certified at $L = 13$ (Theorem 6.1) and, round 7, at $L = 15, 17, 21$ (full reruns, verified; Section 6.8); residue: reading of (0.4), gauge choice; other L are a C3 cost, not an open item
B2	(PI) probe independence	none	verified reduction [R] at fixed j (`SOURCES_T2_PI` + VERIFY) given (I1) massiveness at $U = 1$, (I2) gauge covariance, (I3) [P]
B3	(F) lattice coefficient = continuum coefficient	X5	round 7: reduced [R], unverified (`SOURCES_T2_F_STEPB`; Section 5.11) to (L-LL), a kernel bound, and the gauge-fixing part of Step C [I]; open numerical tension: single-step β_1 at $L = 5$–11 does not follow $c \log L$ (Section 6.8); stays on the no-proof list

Group B: our own open items

ID	statement	source	status
B4	(M_j) formal = true f^2 coefficient, (S1)–(S3)	none	round 7: proved relative to listed inputs, verified, at fixed $j \geq 1$, odd $L \geq 3$ (Theorem 5.14): region 1 as an asymptotic identity, neutral sector (N2); only the $o(f^2)$ form; the proved (G'_A) range suffices; residue one [I] convention
B5	V1: $\beta_{j+1}(0)$ is the Maxwell coefficient	CMP 109 (1.3)–(1.5)	proved+verified at fixed level, conditional on B6, Bałaban's inductive bounds, and the averaging (0.4)
B6	(G2) per-cell infinite-volume densities	none (definition of β_j only)	verified reduction [R] at fixed j (was formal); modulo $\tilde{Q}$ onto, (Gap_f), averaging analyticity
B7	L5: β_{k+1} on the clamped cube, or nested-interval Lemma D″	CMP 109 p. 298	round 7: proved relative to listed inputs, verified, by Route 2 (Theorem 5.15): nested intervals on true trajectories, no clamp, v4's constants; one reading [I] (causality)
B8	K-independence of $\beta_j(g)$, R-term part; the original form is ill-posed (the whole lattice has no R-terms); replaced by (a) β_j has no R part, (b) E_{pure} on non-wrapping $X \subset \Lambda$ = the $\mathbb{Z}^4$ term, (c) E_R K-uniform through F2′	→ A15; CMP 119 p. 279	round 6: proved relative to listed inputs, verified (Section 5.10): C31–C37, F2′ inputs (B20), CMP 109 (1.18), (3.48), (4.36); input of Theorem 1, not of the shooting
B9	Route B inputs (T1), (DF), v4 "A6", `BETA_FP_U` Theorem 2.4	none	argued; Route B only
B10	Proved, unverified results in the chains (sweep, `BETA_FP_M` Prop. 4.3, Prop. 3.6, N3 in D″ form)	→ A11	round 7: cleared, verified (Section 5.11): all 19 sweep rows second-read, CMP 122 additions X1–X10, S12–S16 (C42); Prop. 4.3 for $j \geq j_1$ only; Lemma H″, N3″
B11	(B-S): sourced Theorem 1, (E14-a, b, c)	→ A13	open as a whole; parts B12–B20, B22
B12	(I): $B_j(c) = \beta_j(u(c))$	none	by construction (C1, C10); not written as a lemma

Group B: our own open items

ID	statement	source	status
B13	Real uniform (H-hist)	→ A1–A3, A1w, A7, A8, A10, A17, A18	verified reduction, both halves (large field: `SOURCES_GGAIN_LF`); round 5: small-field (Q)/c_- discharged by Theorem 5.9 (with C25, C26); open inputs (M_r), N3 Kotecký–Preiss step, CMP 102 (189), (M-B) bookkeeping; Theorem SM rests on A10′ with β_H; G4 off-chain
B14	(H-hist2)	none	verified reduction
B15	(H-thr) threshold mismatch	CMP 122 II p. 383	reduced to M2′, M4, M5
B16	(H-hol), small field	conditions (C1)–(C5)	verified reduction
B17	(H-hol), large field	→ A4, A18, A19	verified reduction; (M-C) reduced to printed CMP 116 §2 conditions + Lemma 2; (I-V0), (I-ord), (I-ε) (both layers) closed and verified; (M1) → C24; round 7, chain item 5: (H−) ⇐ A4 + A19 (K1–K4) + A18 + (C1*); (C1) proved given A19, A18; (C2) merged into A18
B18	N3 strip bound	none	proved+verified given B12–B17
B19	(FX) final Mayer expansion	not in CMP 116–122	reduced; remaining input B18
B20	Sourced step bookkeeping: (W4), (P8), F2′, (D), Λ_j	→ A4, A10, A16	reduced; Λ_j bookkeeping open
B21	(B-N1) bridge and (S)	none	(S) clauses: verified reduction, h-independent; (E14-c) clause open; (S_{src}) ⇒ (H-U) proved+verified; compatible with (U) (C23)
B22	Recovery of $\kappa \to \kappa/2$ in the sourced step (only; (U) moved to C23)	none	verified reduction [R] by $L/2$-rescaling, $\kappa \geq 1536$; uses A23; round 7: (K1′) $\delta \leq \frac{1}{13}(1 - 2(1+4\beta)/(\theta_K L))$, $\theta_K \geq \frac{1}{2}$, $\kappa \geq 2m_{\mathrm{tot}}$; $\delta = 0.025$ at $L = 13$ (0.03 fails; Lean `K1_numeric_13`)

Group C: conventions and choices

ID	choice	source	status
C1	Freeze thresholds and partitions at Bałaban's trajectory x^*	`SOURCES_D3`, `SOURCES_HTHR`	free; derivative needs B15
C2	Option B margin, $\theta_W + \tau_C \leq 0.2$	`SOURCES_MB_VERIFY`	free; costs a smaller γ
C3	L odd, $L > 11$; target $L = 13$	v4 §1.1	loses generality
C4	$G = SU(2)$, $d = 4$	Theorem 2	loses generality
C5	Order of constants: the canonical acyclic order of `SOURCES_ORDER_CANON` (amended E1–E8), β_H (C27) first	`SOURCES_ORDER_CANON` (+ VERIFY), `SOURCES_N3A`	settled; verified twice; two nominal cycles resolved by the split (Section 5.10)
C6	Holomorphy radius ρ_0	`SOURCES_HHOL`	free
C7	Cut on B, not B'	CMP 109 (2.9), (2.12)	reading; closed
C8	Sharp χ_k; smooth fallback	`SOURCES_G3`	free
C9	Shrunk χ_k threshold	`SOURCES_GGAIN_VERIFY`	changes the construction; A13 must hold for it
C10	Gauge-fixing weight $1/u_k(c)$	`SOURCES_D2C_VERIFY`	changes the construction; A13 must hold for it
C11	$h \equiv 0$ booking; freezing set	`SOURCES_D2C`	free
C12	Additive source normalization	`SOURCES_HBETA_N3`	free
C13	Main R-step form (1.90)–(1.98)	`SOURCES_FX_O23_VERIFY`	free; removes O4
C14	κ_{final} fixed before M	`SOURCES_FX_R2`	free; not sharp
C15	$2p_1 - (d+5)r_0 > p_0$	CMP 122 II p. 383	condition (Bałaban's own)
C16	$32\varepsilon' \leq 1$, $S \geq 1$, $\kappa_0 > 2$	`SOURCES_D3W8_VERIFY`	changes the construction; A13 must hold for it
C17	(S): $\gamma^{-2} \geq c_{sw} D'$	v4 §1.2	condition; needs β_0 fixed before γ (verified reading)
C18	ℓ even; thermal state; plaquette observable; test space D_n; origin at a site; compactness form	`ASSEMBLY` §0–§1	loses generality
C19	Coupling domain $]0, \gamma]$	`SOURCES_GGAIN_VERIFY`	free
C20	Route K primary, Route B alternative	v4 §1.3	free
C21	$\theta_* < 1$ in Route B	`BETA_FP_PW_VERIFY`	free; trivial
C22	$g_0(K, g)$: any shooting solution	Lemma D	loses generality
C23	(U): $\rho_1 \leq c_* r_0 / M^a$ at $\lambda_K = 1$	`SOURCES_UN1` (+ VERIFY)	constant choice, not a hypothesis on test functions; verified compatible with B-N1; C^n cost per n-point bound
C24	(M1): $M = L^m$ at least the largest of 32 lower bounds; no upper bound; real $\alpha_0 \leq \alpha_0'/(CM)$ with $C > c_J$	`SOURCES_ORDER_CANON` (+ VERIFY), `SOURCES_HIGHER_M1` (+ VERIFY)	convention, free choice; the admissible set of M is an up-set

Group C: conventions and choices

ID	choice	source	status
C25	(Hα): $(\alpha_5 + G_\Gamma^2\delta_S)\Lambda_0 \le \frac{1}{2}$	`SOURCES_G3_N2`, `SOURCES_G3_GGAMMA` (+ VERIFY)	condition, not choice; stronger than Bałaban's $\alpha_5\|C^{(k)}\| < \frac{1}{2}$; $G_\Gamma \le \bar{G}$ uniformly in M (verified), so (Hα) is the single threshold $M \ge \max\{M_\Gamma, M_{\alpha 5}\}$; closed as an M threshold (round 6); one [I] residue: the s-rule acts on Bałaban's (3.185) resolvent expansion (CMP 116 p. 13)
C26	(Hγ): $\frac{1}{20}\gamma_2\varepsilon_1^2/\gamma^2 \ge 4\kappa+1$, and the (2.31) condition $\frac{1}{20}x - 4M^4e^{-x/20} \ge 0$ at $x = \gamma_2\varepsilon_1^2/g_k^2$	`SOURCES_G3_N2` (+ VERIFY)	condition, not choice; one unit stronger than Bałaban's $\ge 4\kappa$; γ-smallness after M, κ, γ_2
C27	$\beta_H \in (0,1)$, $\beta_H \le \beta_0 < 1$: the unprinted Hölder exponent of CMP 109 (3.31), first in the order	`SOURCES_A10_LOWER_VERIFY`; `SOURCES_GGAIN_BETAH` (unverified assessment)	free; every consumer needs only $\beta_H > 0$; cost $(1 - L^{-\beta_H})^{-1}$ factors, larger $B_4(\beta_0)$
C28	$M_1 < M_2 < M$, order $M_1 \to M_2 \to M$	CMP 119 p. 245; `SOURCES_A11_CONST_VERIFY`	free; printed order
C29	$M_2 \ge 1/\delta$	`SOURCES_A11_CONST` (+ VERIFY); CMP 119 (3.7), (3.17)	free; unprinted lower bound
C30	$A_1/A_0 \le 1/(80B_3)$, A_1 after A_0	`SOURCES_A11_CONST` (+ VERIFY); CMP 119 p. 266	free; Bałaban: "sufficiently small"
C31	(Vol): torus side $\ell > 22M$ final-scale cells; even form $\ell = 2L^{m_T}$, $m_T \ge m_M+1$	`SOURCES_A15_BDRY` (+ VERIFY), `SOURCES_T2_VOL`	condition, admissible; K-free; not used by Theorem 2w; used by B8 and Theorem B
C32	(Cube): $U_j(\square_0, \cdot)$ of [14] uses only data on $\square_0$	CMP 109 p. 276; `SOURCES_A15_LOC` (+ VERIFY)	reading [I]
C33	(Cases): the other term classes are localized like the typical term	CMP 116 p. 2	Bałaban's printed sketch [R]
C34	(Conv): cube sides divide N; partitions are π-images of the $\mathbb{Z}^4$ partitions; $M = KR_0M_0$, K a power of L	`SOURCES_A15_LOC` (+ VERIFY)	convention on an unprinted part, admissible
C35	(Det′): domain-assignment rule for the closed-walk terms of the determinant expansion	CMP 102 pp. 271–272; `SOURCES_A15_LOC_VERIFY`	convention on an unprinted part, admissible
C36	"H_j with free boundary conditions" is the $\mathbb{Z}^4$ operator	CMP 109 p. 290	reading [I]
C37	(Enl): the s-rule domain of a walk is $\bigcup X_i^5$ as a whole-lattice set	CMP 116 p. 3; `SOURCES_A15_BDRY` (+ VERIFY)	convention on an unprinted part, admissible; the Λ-restricted form costs at most one M-cube layer

Group C: conventions and choices

ID	choice	source	status
C38	(C1*): $c_m M_0\, g_k p_0(g_k) \le \frac{1}{2}\kappa_H C_1^{102} a_*^{102}$, uniform in k	`SOURCES_T2_HMINUSC` (+ VERIFY); `SOURCES_ORDER_CANONF`	condition, not choice; γ-smallness; ε_1^{102} per scale exists iff (C1*) and $\varepsilon_1^{(0)} \le a_*^{102}$
C39	Slot of M_0 and a_*^{102} (with the cube cap $\varepsilon_1^{102} \le R_1 M_1 a_1^{102}/M$) after M, before γ	`SOURCES_T2_HMINUSC_VERIFY`; `SOURCES_ORDER_CANONF`	free constant-order entry; the cap is booked on ε_1^{102}, so M stays one-sided
C40	A16 tree conditions (A16-a^T), (A16-b^T), (H$^f_\kappa$), (H$^f_\eta$), $\bar\Gamma_1^f \eta_M^f \le \frac{1}{2}$, $C'^f_{\mathcal{D}}$, the $O(\gamma^2 r)$ budget entries	`SOURCES_A16_REPLACE2` (+ VERIFY); `SOURCES_ORDER_CANON` F	conditions, verified; linear in $g^2 r$ or M thresholds; α_5 slack shared with C25
C41	(A18-1)–(A18-4) and $M \ge 5M_{99}$ with $10 \mid C^{122}M$	`SOURCES_A18_REGULAR` (+ VERIFY); `SOURCES_ORDER_CANONF`	conditions, constant-order entries; acyclic; ε_0^{102} two-sided but nonempty
C42	CMP 122 I–II threshold and exponent conditions S12–S16, with the cross-scale list X1–X10	`SOURCES_A11_CONST` addendum; `SOURCES_B10_SWEEP_VERIFY`	conditions, Bałaban's own (printed); exponents before γ; N two-sided per scale, exists for γ small

Verified readings. Four former Group A rows are now verified readings and appear in no set (`SOURCES_BOOKKEEPING.md`): A9 (the constant $C_1'(d,L)$ is unprinted and not derivable from the printed lines; its value is not used), A12 (β_0 "arbitrarily small" is on CMP 119 p. 255; β_0 is fixed before γ), A20 ($R_{K+1} \ge L$ follows from CMP 119 (2.5); C_c is free of g), and A22 (Π (4.37)/(5.1), CMP 119 (2.42) corrected, (2.12)/(2.23), the "L odd" page; the Sect. 3 residue is A23).

The free parameter β_H. The per-label exponent of Fact A (Theorem SM) is the Hölder exponent β_H of CMP 109 (3.31), which Bałaban does not print (Theorem 5.6). It is a free parameter, $0 < \beta_H \le \beta_0 < 1$, and it comes first in the order of constants: $\beta_H \to d, L \to \cdots \to \varepsilon_1 \to \gamma$ (C5, C24, C27). Every consumer needs only $\beta_H > 0$, because each condition it adds is an ε_1- or γ-smallness condition (`SOURCES_GGAIN_BETAH.md`, [UNVERIFIED ASSESSMENT]). The cost is the factors $(1 - L^{-\beta_H})^{-1}$, a larger $B_4(\beta_0)$, and K_T fixed before ε_1. Margins at $L = 13$: the (G-gain) margin $1 - 13^{-\beta_H/2}$ is 0.120 at $\beta_H = 0.1$ and 0.685 at $\beta_H = 0.9$; the Lemma HH margin is 0.901 at $\beta_H = 0.9$.

Binding sets (round 7; sizes unchanged). *Theorem 2w* (Theorem A(i), Route K) rests on one Group A item, A5, on B1–B7, and on X5; Group C: C3, C4, C17, C22. A5 has no bare assertion left (Proposition 5.8, Theorem 5.10); its use is a verified reduction with residue N1–N9. If A5 is taken through this reduction, Theorem 2w also rests on A7, A8, A17, the inputs of Bałaban's sup induction, C25, C26, and C32–C36; these are not counted in the set. Of B1–B7: B1 is certified at $L = 13$ (and at 15, 17, 21); B2, B5, B6 are verified reductions or proofs relative to named inputs; B4 (Theorem 5.14) and B7 (Theorem 5.15) are proved relative to listed inputs and verified; B3 is reduced [R] but unverified, with an open numerical tension. B7 adds `SOURCES_HHOL.md` §2–§3 and the causality reading [I] to the inputs of Theorem 2w. Route B: {A5} + {B1, B6, B7, B9} + X1.

Theorem 2w $\Rightarrow$ *Theorem 1* (Theorem A(ii)) rests on 4 Group A items (A5, A11, A13, A15), 9 Group B items (B1–B8, B10), and X5; Group C adds C28–C37 and C42. B10 is cleared (verified).

Inside the proof of Theorem 1 (A13), Bałaban also uses A1–A4, A1w, A7, A8, A10, A14, A17, and A18.

Theorem B rests on 17 Group A items (A1–A5, A1w, A7, A8, A10, A11, A13, A15–A19, A23), 4 externals (X2–X5), and 21 Group B items (B1–B8, B10–B22); Group C adds C25–C42. Of the 11 Group A items that Bałaban prints without a proof, A1w, A2, A3, A7, A8, and A10′ carry full proofs relative to listed inputs, A11 is a verified list, and A1, A4, A5, A15 carry verified reductions. Of the implicit or unprinted items, A16 and A17 are proved relative to listed inputs, and A18, A19, and A23 are verified reductions. So no Group A row remains without a verified proof or verified reduction (cited-but-unproved and implicit alike); the only Theorem B Group B row without proof or verified reduction is B3. A14 (inside A13 only) stays asserted. This does not make Group A or Group B proved. Every row stays in the ledger, because its residual inputs are not proved: Bałaban's printed sketches [R], conventions, readings [I], the condition (C), (Vol), C38–C42, and B3. Sub-items still open: M2′/M4/M5 (B15), the Λ_j bookkeeping (B20), and the (E14-c) clause (B21).

Counts: Group A 19, Group B 22, Group C 42, externals X 5. Group A has 19 items (A21 left the ledger in round 4; its row is kept above for reference; A9, A12, A20, A22 are verified readings). 14 are printed by Bałaban, and 13 of these are printed without a proof. Group C gained C25–C30 in round 5, C31–C37 in round 6, and C38–C42 in round 7. Of Group C, C3, C4, C18, C22 lose generality; C23 ((U)) and C24 ((M1)) are constant choices; C9, C10, C16 change Bałaban's construction; C15, C17, C25, C26, C31, C38, C40, C41, C42 are conditions; C27–C30 and C39 are free entries of the order of constants; C32 and C36 are readings; C33 is a Bałaban sketch [R]; C34, C35, C37 fix unprinted parts of the construction. This is still a verified reduction, not a proof.

A Lean formalization

The shooting lemmas of Section 3.4, the arithmetic of Theorem K and Theorem C1, and the bridge lemmas of the certifier are formalized in Lean 4 with Mathlib (25 files in `lean/YmLean/`); seven files formalize lemmas of the sources block (Schur test, block-valued Schur test, Combes–Thomas, interpolation and rescaling, resolvent and weighted Neumann bounds, the gauge condition of the constrained minimizer, the slab counterexample to CMP 96 (2.127)). The inventory is as follows.

[Lean] (no sorry; axioms propext, Classical.choice, Quot.sound)

- Lemma D′ (a)–(d) (`YmLean.lemma_DPrime`), Step 8 (`lemma_DPrime_step8`), Lemma D, Lemma E (Cg^2 form), Lemma D+E (`lean/YmLean/LemmaDPrime.lean`, `LemmaDE.lean`). One-argument β only.

- Lemma D″ (`YmLean.lemma_DDoublePrime`, `lean/YmLean/LemmaDDoublePrime.lean`, 146 lines): (a″) bracket $(3b/4)(K-k) - 2R \le T(k,K) \le (5b/4)(K-k) + 2R$; (b) shooting for $g \in]0, g_1]$ with the clamp inactive; (c′) with $\beta = 3b/4$, $B' = 5b/4 + 2R$, $D' = 2R$, for every trajectory with couplings in $]0, \gamma]$; (d) for $n - m \ge 8R/b$, lower constant $b/2$. Hypotheses weaker than the note (continuity, any admissible γ, $R \ge 0$ derived). One-argument β only; L5 not formalized.

- Full inventory (`lean/README.md`; file, main theorems, source note):

  1. `LemmaDPrime`: `lemma_DPrime`; `LEMMA_DPRIME.md` §1–§2.
  2. `LemmaDE`: `lemma_D`, `lemma_E`, `lemma_DE`, `lemma_DPrime_step8`; `LEMMAS.md`.
  3. `LemmaH`: `gronwall_telescope`, `lemma_H`; `SOURCES_HBETA.md` Lemma H.

4. `DetIdentities`: `det_bordered`, `det_restrict_add_gauge`; `numerics/NUMERICS.md` §1.
5. `Telescoping`: `telescoping_det`, `telescoping_log`; `R2_TELESCOPING.md` Prop. 3.1.
6. `AbelianCount`: `su2_count`, `abelian_share`; `U1_INFINITE_LATTICE.md` Prop. 9.1 (used here only as a cross-check, Section 6.2).
7. `LemmaDDoublePrime`: `lemma_DDoublePrime`; `KAPPA_BOUNDED.md` §5.3.
8. `TheoremK`: `telescoping_bound`, `window_of_partial` (the window bound (5.1) from partial sums); `KAPPA_BOUNDED.md` §5.
9. `Gronwall`: `discrete_gronwall`, `contraction_bound` ($N_{j+1} \le qN_j + Z$); `KAPPA_BOUNDED.md` Prop. 4.1.
10. `CubeCount`: `cube_count_sharp`, `cube_count_lower` (Lemma O1-R′ arithmetic); `SOURCES_FX_O1.md`.
11. `BlockNorm`: `rho2_le`, the exact factorization $\rho_2 \le \rho_1(j)\max_l \rho_1(j-1,k_l)$; `H2_CERTIFY.md`.
12. `OishiRump`: `oishi_rump`, `oishi_rump_eta`, `frobenius_power_bound`; `H2_CERTIFY.md` §5.
13. `Envelope`: `envelope_nontrivial`, `envelope_not_scale_invariant` ((B-NT) in log form); `ASSEMBLY.md` §2.9.
14. `Divergence`: `profile_divergence`, `P4_divergence` (identity (D) of premise P4); `H2_CERTIFY_P4.md`.
15. `TheoremC1`: `c1_two_step`, `c1_kappa_bound`, `c1_window`; `B2_TWOSTEP.md` §2.
16. `StripBound`: `strip_bound_linear`, `strip_bound_quadratic` (the $\frac{3}{2}\tau$ bootstrap); `SOURCES_HBETA_N3.md` §4.
17. `CertBridge`: `cell_invertible`, `cell_sup_bound`, `vertex_bound` (interval-to-norm bridge of the cell certifier); `numerics/h1cert_cell.py`, `h1cert_cell2.py`.
18. `Tower`: `tower_bounds`, `I1_compress`, `majorant_core`, `lemmaT_core` (algebraic core of the one-step majorant, Theorem 6.1); `H1_ANALYTIC.md`.
19. `Schur`: `schur_test_weighted`, `schur_test` (weighted Schur test for the spectral norm); `SOURCES_LOC.md` §3.3, `SOURCES_S2_FORM.md` §2, `SOURCES_A3_SIGMA.md` 2.6.
20. `CombesThomas`: `norm_entry_le`, `coercive_inv`, `combes_thomas`, `combes_thomas_exp`, `combes_thomas_decay`, `combes_thomas_schur` (the proof structure of Lemma CT, the kernel-decay reduction of ledger A1); `SOURCES_A1_KERNEL.md` §7. The identification with the block operators, the kernel-to-row-sum step, (K)/(C), and the rescaling are not formalized (`lean/VERIFY_CombesThomas.md`); the block-valued Schur test is in `BlockSchur`.
21. `Interp`: `multilinear_convex`, `monomial_affine` (`SOURCES_A3_SIGMA.md` 3.2); `window_perturb`, `window_perturb_partial` (`SOURCES_T2_KUNIF.md` 5.2, chained to Theorem K); `rescale_exp`, `rescale_rate`, `K1_iff`, `K1_numeric` (`SOURCES_T2_KAPPA2.md`); `xexp_le`, `xexp_absorb` (`SOURCES_GGAIN_LF.md` §2).
22. `Resolvent`: `resolvent_identity`, `resolvent_bound`, `lower_transfer`, `neumann_perturb`, `resolvent_chain` (`SOURCES_A3_SIGMA.md` §4); `wnorm_mul_le`, `wnorm_neumann`, `pwnorm_mul_le`, `pwnorm_neumann` (weighted Neumann series, row half; `SOURCES_A2_R2.md` §2).
23. `SlabCounterexample`: `slab_fails_10`–`slab_fails_13`, `slab_fails`, `lambda1_lt` (the proof step CMP 96 (2.127) fails for every $L \ge 10$); `SOURCES_A3_GAMMA0_FULL.md` §3.2–§3.3.
24. `BlockSchur`: `block_row_le`, `block_schur_test` (Schur test with block norms $\|K_{xy}\|_2$ in place of $|K|$); `SOURCES_A1_KERNEL.md` §7, `SOURCES_S2_FORM.md` §2 Π3.
25. `GaugeMin`: `variational`, `constrMin_existsUnique`, `lemma5`, `lemma5_full` ($RD^*A_* = 0$ for the constrained minimizer, from (F1), (F3), (F4) as hypotheses); `SOURCES_A3_GAMMA0_FULL.md` §5.

Not formalized: interval arithmetic itself, the enclosure premises P1′–P4, the identity $\rho_1^2 L^2 = \lambda_{\max}(G)$, and the history form of the shooting lemmas.

B Index of results by status

This appendix indexes the results of the paper by status tag; the tags match the verifier files. Here v1–v4 denote the successive versions `THEOREM2_PROOF.md`, `THEOREM2_PROOF_v2.md`, `THEOREM2_PROOF_v3.md`, and `THEOREM2_PROOF_v4.md` of the Theorem 2 notes. The statements that the main theorems take on trust are listed by item in the trust ledger `TRUST_LEDGER.md`, condensed in Section 7.7; ledger IDs are given below where an entry corresponds to a ledger item.

[PROVED+VERIFIED]

- Lemma D′ (a)–(d), K_0, Remark 4 (`LEMMA_DPRIME_VERIFY` grades 1, 2, 5).
- Lemma D (corrected interval), Lemma E (Cg^2 form), nested-interval variant of Lemma D (`LEMMAS_DE_VERIFY`). Lemma E C^1 form: trivial variant, same proof.
- Lemma D″ (tail-sum form, no J) (`KAPPA_BOUNDED_VERIFY` §6).
- Telescoping $S_k = \sum \beta_j(0)$ for the $p \to 0$ coefficient (`BETA_FP_VERIFY` item 1, corrected scope).
- Prop. 2.1 anchor-free identity; exact at the constant field (`BETA_FP_LEMMA43_VERIFY` item 1; `BETA_FP_H_VERIFY` item 1).
- $c = 11N^2/(12\pi^2)$, normalization chain; (F) source (Savvidy eq. (1.4); ledger X5) (`BETA_FP_LEMMA43_VERIFY` item 2; `BETA_FP_H_VERIFY` §4).
- KAPPA Prop. 1.1 as an identity given its fixed-level inputs; Lemma 2.1 (integration by parts); Prop. 2.2; level-independence of the degree -4 part; Theorem K and Theorem 2w as conditional reductions (`KAPPA_BOUNDED_VERIFY` (i)–(v)).
- $\sup_j |\sigma_j^{(2)}(0)| < \infty$; Prop. 4.1 algebra (`KAPPA_BOUNDED_VERIFY` (vi), reason for A4 failure corrected: obstruction = $\partial\sigma^{(1)}$ in Q_j and in (W-2), §7 there).
- `B2_TWOSTEP` two-step recursion; Theorem C1 ((H2) ⇒ ($C1\sigma$), factor ρ_2^2/L^2) and Theorem B2 ((H2) ⇒ (B2), factor ρ_2^2/L^4), conditional on (H2); threshold $\rho_2 < L$ (no strip loss); (H2) weaker than (H″) at $n = 2$ (`B2_TWOSTEP_VERIFY` grades 1, 3, 4). Form of Q_j by read-off (grade 8; `U3_LEMMA51` (3.3) not re-derived).
- `h2_scan` code structure, one-colour reduction, H and Δ_{out} blocks, Gram identity, separable Prop. A form (`H2_LARGE_L_VERIFY` grade 1); tail-bound derivation (4.1)–(4.3) (grade 6, inputs corrected).
- Logic of Theorem B and Lemma 1.3 of `BETA_FP_PW`, as conditional implications (`BETA_FP_PW_VERIFY` §6 items 1–4 CONFIRMED: doubling identity and Lemma 1.3; Prop. 2.2 / Lemma 2.3; Theorems A, B; Lemma 4.1, conditional). Its inputs (U), (DF), (T1) are listed under [PROVED, UNVERIFIED] and [ARGUED].
- Prop. E evenness, Theorem W0/W, Theorem A3, Lemma A4, A0, Prop. A2 (`U3_AW_VERIFY` grades 1–4).
- U3 Theorem 4.1 + O5, Prop. A, Prop. B, O6, Theorem 6.1 (given V1; Taylor-data norm), Lemma 5.1 neutral value bound, single-block $m = 2$ failure in $d = 4$, one-step recursion, R2 midpoint identity (`U3_VERIFY`, `U3_REPAIRS_VERIFY`, `U3_LEMMA51_VERIFY`, `U3_LEMMA51_R23_VERIFY`).

- Exact one-step factorization $\rho_2 \le \rho_1(j)\max_l \rho_1(j-1,k_l)$ (`H2_CERTIFY_VERIFY`; Lean `rho2_le`).

- Sources block: Lemma G3 given (Q) (`SOURCES_G3_VERIFY`); LOC (`SOURCES_LOC_VERIFY`); (M-A) given (H−) and (M-A′) (`SOURCES_MA_VERIFY`, `SOURCES_MA2_VERIFY`); (M-B) with $p_0 \ge 17/(\gamma_0 A_1^2)$ (`SOURCES_MB_VERIFY`); condition (a) of Lemma N3 (`SOURCES_N3A_VERIFY`); the sourced large-field draft B-S.D3, with corrections (`SOURCES_D3_VERIFY`, `SOURCES_D3R_VERIFY`, `SOURCES_D3W8_VERIFY`); (B-N1) given (S) (`SOURCES_HHIST_N1_VERIFY`).

- Cesàro: given convergence, $\kappa_k = o(k) \Leftrightarrow \beta_\infty = c\log L$ (equivalence, not a weaker form) (`THEOREM2_PROOF_VERIFY` Point 4).

- Item 7′: the running-maximum lift of C_β to non-monotone pasts (`SOURCES_GGAIN_NONMONO_VERIFY`).

- (KE') by citation and (H_Q) (`SOURCES_QE_KE_HQ_VERIFY`); S2 for G and G_1 (`SOURCES_S2_FORM_VERIFY`, `SOURCES_S2_G1_V7_VERIFY`); fixed-j (S1), (S3), $(\mathrm{S2})|_{t=0}$, (S2c) given (I3) (`SOURCES_T2_S123_VERIFY`).

- V1 at fixed level, history-free, conditional on (G2), Bałaban's inductive bounds, and the averaging (0.4) (`SOURCES_T2_V1_VERIFY`); k-uniform decay and moments of E_0, ledger A21 (`SOURCES_T2_EUMOM_VERIFY`).

- Round 5, proved relative to listed inputs and verified: A7 (Theorem 5.4), A8 in corrected form (Theorem 5.5), A10 in the corrected form A10′ (Theorem 5.6), Lemma G3 = N2 with (Q) from A7 + A8 and G_Γ bounded uniformly in M (Theorem 5.9), Prop. A15-N given (Loc), (Bd), (H), (Cnt) (Proposition 5.8), and (I3) at every step (`SOURCES_T2_I3_VERIFY`); A11 closed as a verified list (Proposition 5.7).

- Round 6, proved relative to listed inputs and verified: Theorem LOC (Theorem 5.10), Proposition BDRY with the replaced B8 (Proposition 5.11), (G'_A) on its stated f-range (Theorem 5.12); the canonical order of constants (`SOURCES_ORDER_CANON_VERIFY`).

- Round 7, proved relative to listed inputs and verified: (M_j) at fixed j (Theorem 5.14), L5 by nested intervals (Theorem 5.15), A16 in tree form (Theorem 5.20), A17 with $r = 1$ (Proposition 5.16); verified reductions: A18 (Proposition 5.18), A19 (Proposition 5.17), A23 (Proposition 5.19); (C1) of the (H−) chain given A19, A18; ledger B10 cleared (the sweep with the CMP 122 additions, `BETA_FP_M` Prop. 4.3 for $j \ge j_1$, the Prop. 3.6 assembly, Lemma H″, N3″) (`SOURCES_B10_SWEEP_VERIFY`).

- Closures of the (M-C) inputs: (I-V0) (`SOURCES_MC_IV0_VERIFY`); (I-ε) and (I-ord) for CMP 109/116 (`SOURCES_T2_IEPS_VERIFY`) and for the large-field layer of CMP 119/122 (`SOURCES_T2_IEPS_LF_VERIFY`; residue [I], every branch closes); (M1) as convention C24 and the bound "order ≤ 2" on coupling and threshold derivatives (`SOURCES_HIGHER_M1_VERIFY`).

[PROVED, UNVERIFIED]

- `BETA_FP_U`, no verifier file: Theorem 2.4 ((U) ⇐ (T1) + (H2)), Prop. 3.1 ($\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$), Prop. 3.2 ($\gamma_{rec}$ bounded), Corollary 4.1, Theorem 2w by the plane-wave route given (T1), (H2), (DF), LW (5.7), A6, (H0); the argued step "(H2) closes ($C1\sigma$)" of §3.5 there is superseded by `B2_TWOSTEP` Theorem C1 [PROVED+VERIFIED].

- `LEMMA_DPRIME_SWEEP`: 19 cross-scale uses covered by (b) + (c′) + (S) (4 of 19 table rows, 11, 12, 16, 18, verified on page images by `LEMMA_DPRIME_VERIFY` comparisons A–E).
- Step "Theorem 2w ⇒ Balaban's Theorem 1" (depends on the sweep and on L5 in the sharpened form of Section 3.4.5; round 7: both verified, Section 5.11).
- Prop. 3.6 assembly with $C(1+j)^2 L^{-2j}$; neutral-sector K-derivative paragraph; $\sigma^{(1)}$ convergence sketch jL^{-2j}.
- Necessity of individual positivity for literal (0.31) at every K (elementary).
- `BETA_FP_M` Prop. 4.3 ($\log|\det|$ bound in the N–O window, given (S1)–(S3)); round 7: confirmed for $j \geq j_1$ in `SOURCES_B10_SWEEP_VERIFY`.
- Premise P4 of the one-step certificate (`H2_CERTIFY_P4`; identity (D) in Lean `P4_divergence`); the $L = 13$ certificate sizing after fix G1 (`H2_CERTIFY_L13_G1`).

[NUMERICAL]

- $\rho_1(\infty) = 1.760, \ldots, 2.0895$ at $L = 3, \ldots, 13$; at $L = 13$ the measured $\sup \rho_1^2 = 4.37$ (`H2_CERTIFY_L13`); the certified bound is below.

[CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]

- (H2) at $L = 13$ (Theorem 6.1; ledger B1): $\rho_1(j, K)^2 \leq 10 < 13$ for all $j \geq 1$ (including ∞) and all K, hence $\rho_2 \leq 10 < 13$ by `rho2_le` (`H1_ANALYTIC`; `H1_ANALYTIC_VERIFY` V0–V9, `H1_ANALYTIC_VERIFY2`, `H2_CERTIFY_P5_VERIFY`). Residue as in Section 6.7.

[VERIFIED REDUCTION]

- (Q) to CMP 99 (3.132), (E_U), and LOC (`SOURCES_Q_VERIFY`, `SOURCES_CDC_VERIFY`); (E_U) to kernel-moment bounds Bałaban does not print (`SOURCES_EU_VERIFY`; the bound $K \leq K_0(1 + \ldots)$ is a [DERIVED SKETCH]).
- N3 to (H-hol) + real (H-hist) + (H2-hist) (`SOURCES_HBETA_N3_VERIFY`).
- (H-hol), small field, to Bałaban's conditions with modified constants (`SOURCES_HHOL_VERIFY`); large field to (M-A), (M-B), (M-C), (H-hist2) (`SOURCES_HHOL_LF_VERIFY`); (M-D) to (H-hist) + (H-thr).
- (H−) to CMP 102 (189), an existing trust item (`SOURCES_MA2_VERIFY`).
- (H-hist2) to the real uniform (H-hist), no new input (`SOURCES_HHIST2_VERIFY`).
- Round 3 (ledger A1–A4, B13, B17, B21, B22): A1 kernel-decay form via Lemma CT to CMP 99 Thms 3.1, 3.3, 3.12 + (KE′) + (H_Q) + S2 (`SOURCES_QE_KE_HQ_VERIFY`, `SOURCES_S2_FORM_VERIFY`); A1w to (W_G) + (C) + CMP 96 Lemma 2.1 (`SOURCES_A1_WALK_VERIFY`), now written out as Theorem 5.1, with (W_G) written out as Theorem 5.2 (`A1_WALK_FULL_VERIFY`, `WG_FULL_VERIFY`, paper-grade relative to their tagged inputs); A2 with the V7 repair to (W^c) + A7 + two [I] (`SOURCES_A2_R2_VERIFY`, `SOURCES_S2_G1_V7_VERIFY`), with (W^c) written out as Theorem 5.3 (`WC_FULL_VERIFY`, paper-grade relative to its tagged inputs); A3, full proof with $\gamma_0 \geq m_L/(48d^2L^d)$, and

its σ-localized clause (SOURCES_A3_GAMMA0_FULL_VERIFY, SOURCES_A3_SIGMA_VERIFY); (M_j) to (I3), (G'_A), Prop. 4.4 region 1 (SOURCES_T2_S123_VERIFY); A4 to A1 + CMP 102 (SOURCES_A1_A4_VERIFY); large-field (H-hist) (SOURCES_GGAIN_LF_VERIFY); (M-C) to the printed CMP 116 §2 conditions (SOURCES_MC_VERIFY); the (S) clauses of B21 (SOURCES_B21_S_VERIFY); the $\kappa \to \kappa/2$ recovery B22 (SOURCES_T2_KAPPA2_VERIFY).

- (G2) at fixed j (item 6, was [FORMAL]): torus densities and coefficient extraction commute with $N \to \infty$, modulo $\tilde{Q}$ onto, (Gap_f), and the averaging-analyticity reading (SOURCES_T2_G2_VERIFY).

- The use of A5 in Theorem 2w (item 7′): Proposition H0-red, residue N1–N9 (SOURCES_T2_H0_VERIFY); N2 proved (Theorem 5.9).

- Item 9 (K-independence in the shooting) closed: β_j is the whole-lattice coefficient by definition, Lemmas D′/D″ run at fixed K (SOURCES_T2_KUNIF_VERIFY).

[INTERVAL ENCLOSURE, VERIFIED]

- Point enclosure $\rho_2(\infty, K^*) \in [1.9993, 2.1273] < 3$ at $L = 3$; P1 (loop-family coefficients exact rationals); Oishi–Rump bound on $\|\mathfrak{B}^{-1}\|$ (H2_CERTIFY_VERIFY).

[ARGUED]

- (PI); (M_j) (Prop. 4.4 of BETA_FP_M); (T1) (plane-wave form of A3, BETA_FP_U §2.3); (DF) (BETA_FP_U §3.4); A6 (v4 label, not ledger A6) = plane-wave vs constrained-minimizer expansion point (uniform in k); $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ bounds; (H″) ⇒ (A) in full ⇒ (B2) (U3_AW §4.7 sketch); C^2, C^3 of $\sigma^{(1)}$ under (H2) (not written); j_* size; the rate $O(\Lambda^{-2} \log \Lambda)$.

[OPEN]

- (H2) at $L \notin \{13, 15, 17, 21\}$ (numerical at $L = 3, 5, 9$; certified at $L = 13$ and, round 7, at $L = 15, 17, 21$, Section 6.8); B3 (F), reduced [R] but unverified, with the open tension of the single-step β_1 against $c \log L$ (Section 6.8); (B2) is reduced to it [VERIFIED REDUCTION]. The residue of A1w, A2: the inputs of Theorems 5.1, 5.2, 5.3, namely (C), the printed sketches of CMP 99 Thms 3.7, 3.9, 3.10, WG-1 in the M_1 form, and [I] geometry and convention items (O3, O4 closed). (T1), (DF), A6 (v4 label) for Route B (ledger B9). The inputs of V1 and (G2): the averaging declaration (0.4), $\tilde{Q}$ onto, (Gap_f), and a j-uniform form of (G2). The residue of (M_j): (I3), (G'_A), Prop. 4.4 region 1. L5 (domain of β; nested-interval D″ [NOT WRITTEN]). The finite-volume identification of the step, R-term part (ledger B8; an input of Theorem 1). The (E14-c) clause of B21. (A) in full. See Section 7.

[CORRECTED BY VERIFIERS SINCE V3]

- B2_TWOSTEP G1: the certification of (H2) is a computer-assisted proof project, not a check (B2_TWOSTEP_VERIFY grade 6).

- H2_LARGE_L: L-trend rate claim (grade 5); j_0 inputs, δ_m 45–90× larger, $a_*(L = 5) = 0.0107$ (grade 6), giving $j_0 \approx 8$ ($L = 3$), ≈ 6 ($L = 5$).

- BETA_FP_PW: comparison and numerics (BETA_FP_PW_VERIFY grades 5, 6).

[REFUTED]

- (HS) $\sup \|H_j\| < L\sqrt{L-1}$ at $L = 3$.
- Not a refuted statement, but a failed method: the one-step sup-norm C^1 recursion at $L = 3$ (factor 1.005–1.033 for $j \geq 2$; Section 3.7.2).
- `U3_LEMMA51` Theorem 4.1 for $m = 2$ with a single block-M constraint in $d = 4$.
- v1 claims withdrawn: "every early step must be certified for Theorem 1" (false: Lemma D′/D″); "(H′) is weaker than $\beta_\infty = c \log L$" (it is equivalent); "the whole $\log \Lambda$ sits in the fine piece" as a fact (it is equivalent to the open statement); "one-loop parts hold for SU(N)" (Lemma 5.1's charged/neutral split is the SU(2) Cartan split); (3.2) with Cj^2L^{-2j} (0 at $j = 0$); the (M) resolvent bound $\|\mathcal{K}_t^{-1}\| \leq C/|f|$ (Nielsen–Olesen zero); (H) with IR weights as stated (UNSUPPORTED).

[BALABAN]

- (H0) smoothness of β_{j+1} in g (CMP 109 (1.22), pp. 263–264), at odd $L > 11$ only; uniformity in j is our reading and uniformity in the history is not printed (ledger A5); Theorem 1 and the inductive bounds of CMP 109, 116, 119, 122 and the propagator papers CMP 95, 96, 98, 99, 102. Uniform level bounds partly cited from CMP 95/96/99 (O6 replaced the main citation). Each Bałaban input that is printed without proof, or used without being printed, is a Group A row of the trust ledger.

[VERIFIED QUOTE]

- Theorem 2 and Theorem 1 wording, "separate paper", p. 298 history sentence, (0.1) with $1/g_k^2$ and χ_k, (1.3)–(1.5) (print misprint $E^{(j+k)}$ read as $E^{(j+1)}$), (2.6)–(2.9) of CMP 119, items A–E of CMP 122 (`BALABAN_EXCERPTS_CHECK`; `LEMMA_DPRIME_VERIFY`; `LEMMA_DPRIME_SWEEP` page images). `BALABAN_EXCERPTS_CHECK` item 10 (CMP 116 has no C^m-in-g statement) is unchanged since v3; the item 11 readings are verified, with corrections, in `SOURCES_BOOKKEEPING.md` (former ledger A22).

[NUMERICAL] see Section 6.

C Label map

The numbered environments of this paper carry the names used in the project files. The first table maps them. The second table maps the section numbers of `THEOREM2_PROOF_v4.md` ("v4") and `ASSEMBLY.md` to the sections of this paper; section references inside the project files use the old numbers.

this paper	name in the project files	source
Theorem A	Route K summary: Theorem 2w ⇒ Theorem 1	`THEOREM2_PROOF_v4` §6, §10
Theorem B	Theorem 1.3 (summary)	`ASSEMBLY` §1.3
Theorem 3.1	Theorem 2w	`THEOREM2_PROOF_v4` §1.3
Lemma 3.2	Lemma D′	`LEMMA_DPRIME`; v4 §2.1
Remark 3.3	Remark 4	`LEMMA_DPRIME`; v4 §2.1

this paper	name in the project files	source
Lemma 3.4	Lemma D″	`KAPPA_BOUNDED` §5.3; v4 §2.2
Lemma 3.5	Lemma D	`LEMMAS`; v4 §2.4
Lemma 3.6	Lemma E	`LEMMAS`; v4 §2.4
Proposition 3.7	Prop. 2.1 (anchor-free identity)	`BETA_FP_LEMMA43`; v4 §3.2
Theorem 3.8	Theorem K	`KAPPA_BOUNDED`; v4 §4.1
Theorem 3.9 (Theorem B_{PW})	Theorem B = Theorem 3.2	`BETA_FP_PW`; v4 §4.2
Theorem 3.10	Theorem C1	`B2_TWOSTEP`; v4 §5.3
Theorem 3.11	Theorem B2	`B2_TWOSTEP`; v4 §5.3
Definition 4.1	(H-U)	`ASSEMBLY` §1.2
Theorem 4.2	Theorem 1.3	`ASSEMBLY` §1.3
Theorem 4.3	Theorem 1.4	`ASSEMBLY` §1.4
Proposition 4.4	Proposition of (B-C)	`ASSEMBLY` §2.4
Lemma 4.5	finite-form lemma of (B-RP)	`ASSEMBLY` §2.7
(GC1)–(GC3)	(G1)–(G3) of the dependency table	`ASSEMBLY` §3
(S_{src})	(S), final sourced bound	`SOURCES_D4D5`
items H, 1–13, B1–B8	items of the open list	`THEOREM2_PROOF_v4` §10
Theorem 5.4	A7 (round 5)	`SOURCES_A7_CDC`
Theorem 5.5	A8, corrected form (round 5)	`SOURCES_A8_POS`
Theorem 5.6	A10′ (round 5)	`SOURCES_A10_LOWER`
Proposition 5.7	A11 list (round 5)	`SOURCES_A11_CONST`
Proposition 5.8	Prop. A15-N (round 5)	`SOURCES_A15_WHOLE`
Theorem 5.9	Lemma G3 = N2 (round 5)	`SOURCES_G3_N2`
Theorem 5.10	Theorem LOC (round 6)	`SOURCES_A15_LOC` §7.5
Proposition 5.11	Proposition BDRY (round 6)	`SOURCES_A15_BDRY`
Theorem 5.12	(G'_A), Thm 5.1 + Cor. 5.2 (round 6)	`SOURCES_T2_GAPRIME2`
Remark 5.13	range remark for (G'_A) (round 6)	`SOURCES_T2_GAPRIME2`, `SOURCES_T2_FRANGE`
Theorem 5.14	(M_j), Theorem 4.2 and §5 (round 7)	`SOURCES_T2_MJ_REST`
Theorem 5.15	Lemmas D″-nest, D′-nest, Route 2 (round 7)	`SOURCES_T2_B7_CLAMP`
Proposition 5.16	A17 identity (round 7)	`SOURCES_A17_A19`
Proposition 5.17	A19 reduction (round 7)	`SOURCES_A17_A19` §4–§5
Proposition 5.18	A18 reduction (round 7)	`SOURCES_A18_REGULAR`
Proposition 5.19	A23 reduction (round 7)	`SOURCES_A23_TSTEP`
Theorem 5.20	A16 in tree form (round 7)	`SOURCES_A16_REPLACE2`

old section	this paper
v4 §1.1	Section 1.2
v4 §1.2	Section 3.1
v4 §1.3	Section 3.2
v4 §1.4	Section 3.3
v4 §2 (notation)	Section 2.3
v4 §2.1	Section 3.4.1
v4 §2.2	Section 3.4.2
v4 §2.3	Section 3.4.3
v4 §2.4	Section 3.4.4
v4 §2.4 (L5)	Section 3.4.5

old section	this paper
v4 §3.1	Section 3.5.1
v4 §3.2	Section 3.5.2
v4 §3.3	Section 3.5.3
v4 §3.4	Section 3.5.4
v4 §3.5	Section 3.5.5
v4 §4	Section 3.6
v4 §4.1	Section 3.6.1
v4 §4.2	Section 3.6.2
v4 §4.2a	Section 3.6.3
v4 §4.3	Section 3.6.4
v4 §5	Section 3.7
v4 §5.1	Section 3.7.1
v4 §5.2	Section 3.7.2
v4 §5.3	Section 3.7.3
v4 §5.4	Section 6.4
v4 §5.5 (arithmetic)	Section 3.7.4
v4 §5.5 (values)	Section 6.5
v4 §5.6	Section 6.9
v4 §5.7	Section 3.7.5
v4 §6	Section 3.8
v4 §7	Section 3.9
v4 §8.1	Section 6.1
v4 §8.2	Section 6.2
v4 §8.3	Section 6.3
v4 §9 (Lean)	Appendix A
v4 §9 (other tags)	Appendix B
v4 §10 (open list)	Section 7
v4 §10 (hypothesis table)	Section 4.6
v4 §11	Section 1.5
v4 preamble (tag legend)	Section 1.4
v4 preamble (range of L)	Section 2.2
v4 change logs	Appendix D
`ASSEMBLY` §0 (notation)	Section 2.1
`ASSEMBLY` §0 (observables)	Section 2.5
`ASSEMBLY` §1.1	Section 4.1
`ASSEMBLY` §1.2	Section 4.2
`ASSEMBLY` §1.3	Section 4.3
`ASSEMBLY` §1.4	Section 4.4
`ASSEMBLY` §2	Section 4.5
`ASSEMBLY` §2.1	Section 4.5.1
`ASSEMBLY` §2.2	Section 4.5.2
`ASSEMBLY` §2.3	Section 4.5.3
`ASSEMBLY` §2.4	Section 4.5.4
`ASSEMBLY` §2.5	Section 4.5.5
`ASSEMBLY` §2.6	Section 4.5.6
`ASSEMBLY` §2.7	Section 4.5.7
`ASSEMBLY` §2.8	Section 4.5.8
`ASSEMBLY` §2.9	Section 4.5.9
`ASSEMBLY` §3	Section 4.6
`ASSEMBLY` §4	Section 4.7
`ASSEMBLY` corrections	Section 4.8

D Revision history

- *Theorem 2 notes, v3.* Built on `THEOREM2_PROOF_v2.md` with the corrections of `THEOREM2_PROOF_v2_VERIFY` (cited there as `v2_VERIFY`): the range of L (odd, $L > 11$, or (H0) at this L as a hypothesis); L5 sharpened (the clamped shooting leaves Bałaban's domain; the nested-interval variant of Lemma D″ is [NOT WRITTEN]); (H″) ⇒ (B2) tagged [ARGUED]; "independent" routes replaced by "formally distinct", with the points $\theta_* < 1$, k-uniform β identification, and the first-order vertex smoothness of (U); the independence record completed (ONLINE_SWEEP §2 as the first channel; the O(3) paper cited); open list extended (item 9, B5, item 10, hypothesis table); sweep count "4 of 19 table rows"; γ_* depends on R through B'. At that time `BETA_FP_PW_VERIFY.md` had no grade, and Route B stayed [PROVED, UNVERIFIED]. The grades of `LEMMA_DPRIME_VERIFY` and `KAPPA_BOUNDED_VERIFY` §9(iv) were cited explicitly.

- *Theorem 2 notes, v4.* Added the material verified after v3: `B2_TWOSTEP.md` with `B2_TWOSTEP_VERIFY.md`, `H2_LARGE_L.md` with `H2_LARGE_L_VERIFY.md`, `lean/README.md` (Lemma D″ in Lean), `BETA_FP_U.md` (Route B inputs), and `numerics/NUMERICS.md` §7. The section on (B2) was rewritten around (H2) (Theorems C1, B2; threshold $\rho_2 < L$; one-step sufficiency at $L \geq 5$; certification plan; corrected L-trend and j_0 inputs). The ledger and open list were updated, with item H as the single uniform hypothesis of both routes. `KAPPA_BOUNDED_VERIFY` states the (B2) obstruction in its §7 and §9(vi). `BALABAN_EXCERPTS_CHECK` had no change since v3.

- *Tag change in the LaTeX version of v4.* The logic of Theorem $\mathrm{B_{PW}}$ is tagged [PROVED+VERIFIED, CONDITIONAL], citing `BETA_FP_PW_VERIFY` §6 items 1–4, per the tag decision in `PLAN.md` (2026-10-28). The Markdown v4 kept [PROVED, UNVERIFIED]. The inputs (U), (DF), (T1) from `BETA_FP_U` stay [PROVED, UNVERIFIED].

- *Assembly.* `ASSEMBLY.md` with the corrections of `ASSEMBLY_VERIFY.md` (Section 4.8). The verifier also found the dependency table stale in five rows (grade 7; no overclaim): B-T2.2, B-T2.3, B-T2.4, B-S.D2C, B-S.D3 lagged `PLAN.md`. The table was rebuilt with a tag for every block and the sub-blocks added later (Route K, Route B, (H2), (H-thr), (G-gain), (FX)). Minor items: `SOURCES.md` §1.3 gives $c_j = 1 + O(g_j^4)$, while this paper uses $O(g^3)$ per `SOURCES_VERIFY` item 8 (the corrected value); the variance estimate drops a harmless g_0^4 (logarithmic) factor.

- *This version.* The Theorem 2 chapter and the assembly chapter are merged into one article: one notation section, one status legend, numbered environments (Appendix C), one numerical section, one list of open items, and the hypothesis table of v4 §10 merged into the dependency table. Three names were changed to avoid collisions (Section 2.6). The dependency table and the open list were brought to the current state of `PLAN.md`: (FX) is reduced to N3 (O2 and R2 [PROVED+VERIFIED], O3 [PROVED+VERIFIED, CONDITIONAL ON N3], O4 avoided by the main form; previously O2, O3 [PROVED, UNVERIFIED] and R2, O4 [OPEN]), and the point enclosure of (H2) at $L = 3$ is verified (Section 6.6). No theorem, lemma, or proof changed.

- *Round 3.* A1–A4 regraded to verified reductions (A3 a full proof with a corrected constant), A1w added and reduced, item 9 closed (B8 moved to the input set of Theorem 1), item 7 restated as 7′, G4 off-chain, B13, B17, B21, B22 regraded, Lean files 21 → 22 (`TRUST_LEDGER.md` round 3; `PLAN.md`).

- *Round 4 (integration of the write-outs).* Theorems 5.1, 5.2, 5.3 (Section 5.8) replace "reduced to printed sketches" for A1w, the final-propagator expansions, and ($\mathrm{W^c}$); all three are paper-grade

relative to their tagged inputs, and the tag stays [VERIFIED REDUCTION]. Corrected constants: WG-1 in the M_1 form; $K_b \le 64d^3L^{2d+1}$; $c_M \le 4\sqrt{N}$; the Schur-sum factor $1 + 4/(e\delta)$; the $\frac{1}{3}M_1$ band by one-layer enlargement. (I-V0), (I-ε), (I-ord) closed; (M1) became convention C24; V1 proved and verified at fixed level; (G2) moved from [FORMAL] to [R]; A21 proved and verified.

- *Round 5.* Statements with hypotheses and residues added in Section 5.9: A7 proved relative to listed inputs (Theorem 5.4; C is the constraint map and $C^*\Delta_k C$ the precision; the reduction to A1 + A18 is superseded); A8 proved in corrected form (Theorem 5.5; Bałaban's symmetry claim is not proved and not needed, the majorant uses $H_0 = \operatorname{sym} C_0^{-1}$; Lemma D with $e^{-\varepsilon'' M/8}$); A10 proved in the corrected form A10′ (Theorem 5.6; the raw Taylor terms do not obey (1.24), the transformed terms do after the exact marginal cancellation); A11 closed as a verified list (Proposition 5.7); A15 narrowed (Proposition 5.8; exact equality cannot be derived; A15 leaves the set of Theorem 2w, and Bałaban's assertion that the $\mathbb{Z}^d$ limit exists moves into A5); Lemma G3 proved relative to listed inputs (Theorem 5.9; G_Γ bounded uniformly in M, verified; (Q) in the joint form, Λ corrected), so the use of A5 in Theorem 2w is a verified reduction; (I3) proved at every step; β_H a free parameter, first in the order of constants. Trust table: rows A5, A7, A8, A10, A11, A15, A21, B4, B6, B13, C5, C24 rewritten; rows C25–C30 added. Binding sets: Theorem 2w = {A5} + B1–B7 + X5 (was A5, A15); Theorem 2w ⇒ Theorem 1 = 4 A + 9 B + X5; Theorem B unchanged in size; the only Group A row without a verified proof or reduction is A15. The headline stays "verified reduction".

- *Round 6.* Section 5.10 added: Theorem LOC (Theorem 5.10; (Loc) exact for non-wrapping polymers, relative to C32–C36); Proposition BDRY (Proposition 5.11; the $\partial\Lambda_{k+1}$ reading exact relative to C37; threshold $N_k > 22M$, $\varepsilon(M) = C_j(22M)^6 e^{-11 m_j M}$); B8 found ill-posed and replaced by (a)–(c), proved; the volume hypothesis (Vol), $\ell > 22M$, added to the setting (Section 2.1) with the even form $\ell = 2L^{m_T}$, $m_T \ge m_M + 1$; (G'_A) proved for $0 < |f| \le ((1-c)/100)^2/2$ with gap $2c|f|$ (Theorem 5.12), a narrow range, with the f-range check still running; the canonical acyclic order of constants, with two nominal cycles resolved by the split, 32 lower bounds on M, and none upper. Trust table: rows A5, A15, B2, B4, B8, C5, C24, C25 rewritten; rows C31–C37 added; counts A 19, B 22, C 37, X 5. Binding sets unchanged in size: Theorem 2w = {A5} + B1–B7 + X5, and A5 has no bare assertion left; Theorem 2w ⇒ Theorem 1 = 4 A + 9 B + X5, Group C adds C28–C37; Theorem B unchanged in size. No cited-but-unproved Group A row lacks a verified proof or reduction; the implicit rows A16–A19, A23 (and A14 inside A13) have neither. The headline stays "verified reduction".

- *Round 7.* Section 5.11 added: (M_j) proved at fixed j (Theorem 5.14; region 1 as an asymptotic identity via the shifted-log split, neutral sector (N2); only the $o(f^2)$ form), with the proved (G'_A) range sufficient (`SOURCES_T2_FRANGE.md`); L5 proved by nested intervals on true trajectories, no clamp (Theorem 5.15); A16 proved in tree form (Theorem 5.20; the first proof's locality step was false); A17 proved by the exact identity $B' = g_k B$ on cut bonds (Proposition 5.16); A18, A19, A23 verified reductions (Propositions 5.18, 5.17, 5.19); (K1) replaced by (K1′) with 13δ and $\delta = 0.025$; B17 chain item 5; B10 cleared; (H2) certified at $L = 15, 17, 21$; canon section F (28 entries, acyclic, four two-sided constants with existence conditions); B3 reduced [R], unverified, with an open numerical tension on β_1 against $c \log L$. Trust table: rows A16–A19, A23, B1, B3, B4, B7, B10, B17, B22 rewritten; rows C38–C42 added; counts A 19, B 22, C 42, X 5. Binding sets unchanged in size. No Group A row remains without a

verified proof or verified reduction; the only Group B row of Theorem B without one is B3. The headline stays "verified reduction".

References

[1] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. I. *Comm. Math. Phys.*, 95:17–40, 1984.

[2] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. II. *Comm. Math. Phys.*, 96:223–250, 1984.

[3] Tadeusz Bałaban. Averaging operations for lattice gauge theories. *Comm. Math. Phys.*, 98:17–51, 1985.

[4] Tadeusz Bałaban. Propagators for lattice gauge theories in a background field. *Comm. Math. Phys.*, 99:389–434, 1985.

[5] Tadeusz Bałaban. Spaces of regular gauge field configurations on a lattice and gauge fixing conditions. *Comm. Math. Phys.*, 99:75–102, 1985.

[6] Tadeusz Bałaban. Ultraviolet stability of three-dimensional lattice pure gauge field theories. *Comm. Math. Phys.*, 102:255–275, 1985.

[7] Tadeusz Bałaban. The variational problem and background fields in renormalization group method for lattice gauge theories. *Comm. Math. Phys.*, 102:277–309, 1985.

[8] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. I. generation of effective actions in a small field approximation and a coupling constant renormalization in four dimensions. *Comm. Math. Phys.*, 109:249–301, 1987.

[9] Tadeusz Bałaban. Convergent renormalization expansions for lattice gauge theories. *Comm. Math. Phys.*, 119:243–285, 1988.

[10] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. II. cluster expansions. *Comm. Math. Phys.*, 116:1–22, 1988.

[11] Tadeusz Bałaban. Large field renormalization. I. the basic step of the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:175–202, 1989.

[12] Tadeusz Bałaban. Large field renormalization. II. localization, exponentiation, and bounds for the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:355–392, 1989.

[13] Roger Dashen and David J. Gross. The relationship between lattice and continuum definitions of the gauge theory coupling. *Phys. Rev. D*, 23:2340–2348, 1981.

[14] Elvin. yang-mills: project repository (notes/ directory holds the project files cited as `FILE.md`), 2026. Local repository; no public remote.

[15] Arthur Jaffe and Edward Witten. Quantum Yang–Mills theory. In *The Millennium Prize Problems*, pages 129–152. Clay Mathematics Institute, American Mathematical Society, 2006. Official problem statement, Clay Mathematics Institute.

[16] Martin Lüscher and Peter Weisz. Background field technique and renormalization in lattice gauge theory. *Nucl. Phys. B*, 452:213–233, 1995.

[17] Paolo Menotti and Andrea Pelissetto. General proof of Osterwalder–Schrader positivity for the Wilson action. *Comm. Math. Phys.*, 113:369–373, 1987.

[18] OpenAI. The canonical massive continuum limit of the two-dimensional O(3) model. `https://github.com/openai/math`, family 215, preprints folder, 2026. October 4, 2026.

[19] Konrad Osterwalder and Robert Schrader. Axioms for Euclidean Green's functions. *Comm. Math. Phys.*, 31:83–112, 1973.

[20] Konrad Osterwalder and Robert Schrader. Axioms for Euclidean Green's functions. II. *Comm. Math. Phys.*, 42:281–305, 1975.

[21] Konrad Osterwalder and Erhard Seiler. Gauge field theories on a lattice. *Ann. Phys.*, 110:440–471, 1978.

[22] G. K. Savvidy. Infrared instability of the vacuum state of gauge theories and asymptotic freedom. *Phys. Lett. B*, 71:133–134, 1977.

[23] George Savvidy. From Heisenberg–Euler Lagrangian to the discovery of chromomagnetic gluon condensation. *Eur. Phys. J. C*, 80:165, 2020. arXiv:1910.00654.

[24] Hao Shen, Rongchan Zhu, and Xiangchan Zhu. A stochastic analysis approach to lattice Yang–Mills at strong coupling. *Comm. Math. Phys.*, 400(2):805–851, 2023.

[25] Ethan Sussman and Levent Alpöge. SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality. ym4_manuscript_zeroth_draft.pdf, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310, 2026. Draft dated 6 October 2026. Accessed 2026-10-27.

[26] Ethan Sussman and Levent Alpöge. The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0. LaTeX files theorem_0_complete_account_raw_{master,I,II,III}.tex, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310 (2026-10-27), 2026. Written largely by LLM, with Claude; not vetted by the repository authors. Accessed 2026-10-27.