Theorem 2 for lattice Yang–Mills theory: proof chapter, version 4

(shortest route; Route K reduction verified)

Written 2026-10-27/28

Contents

1 Statement **3**
- 1.1 Balaban's statements [verified quote] . . . 3
- 1.2 What Balaban's Theorem 1 actually needs (`LEMMA_DPRIME_SWEEP` §2, §7) . . . 3
- 1.3 The weak form proved conditionally in this chapter . . . 4
- 1.4 Literal (0.31) . . . 4

2 Shooting lemmas **5**
- 2.1 Lemma D′ (shooting without early positivity) [proved+verified; Lean] . . . 5
- 2.2 Lemma D″ (tail-sum form; no index J) [proved+verified; Lean] . . . 5
- 2.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [proved, unverified (sweep); 4 of 19 table rows verified] . . . 6
- 2.4 Lemma D and Lemma E [proved+verified; Lean] . . . 7

3 The exact structure of the one-loop sums **8**
- 3.1 Telescoping [proved+verified, corrected scope] . . . 8
- 3.2 Prop. 2.1, anchor-free identity [proved+verified] . . . 8
- 3.3 The constant c [proved+verified] . . . 8
- 3.4 The decomposition $S_k = ck \log L + \kappa_W^F - \kappa_k$ . . . 8
- 3.5 Level-uniform symbol facts used below [proved+verified] . . . 9

4 Bounded window sums: two formally distinct conditional routes **9**
- 4.1 Route K: Theorem K [proved+verified, conditional on (B2) and the fixed-level inputs] 10
- 4.2 Route B: Theorem B of `BETA_FP_PW` [proved+verified, conditional (logic of Theorem B)] . . . 11
- 4.3 Route B inputs after BETA_FP_U [proved, unverified; no verifier file] . . . 12
- 4.4 Comparison . . . 12

5 The hypothesis (B2) of Route K, and the single uniform inequality (H2) **13**
- 5.1 Proved . . . 13
- 5.2 Refuted, and the failed one-step bound . . . 13
- 5.3 (H2) and the two-step theorems [proved+verified, conditional on (H2)] . . . 14
- 5.4 Numerics for (H2) [numerical; none certified] . . . 14
- 5.5 One-step sufficiency at $L \geq 5$ [arithmetic proved; inputs numerical] . . . 15
- 5.6 Certification plan and j_0 [designed, not done] . . . 15
- 5.7 The precise remaining inequality . . . 16

6 Conclusion **16**

7 The convergence route: what it adds, and its status **17**
7.1 Components and status 17
7.2 Early steps and j_* 18

8 Numerics **18**
8.1 First increment $\beta_1(0)$ (NUMERICS §6–§7; UMK_BETA1.md) 18
8.2 Later increments ($L = 2$ only) 19
8.3 Quantities of the Route K proof 20

9 Ledger (tags match the verifier files) **20**

10 Open list (complete for Theorem 2w ⇒ Theorem 1) **23**

11 Citation and independence **27**

A Change log (v2 → v3) **28**

B Change log (v3 → v4) **29**

Assembly chapter: the finite-torus existence theorem **31**

Preamble

Written 2026-10-28 (Opus). v4 = v3 (THEOREM2_PROOF_v3.md[1]) updated with the verified material that appeared after v3: B2_TWOSTEP.md + B2_TWOSTEP_VERIFY.md, H2_LARGE_L.md + H2_LARGE_L_VERIFY.md, lean/README.md (Lemma D″ in Lean), BETA_FP_U.md (Route B inputs), numerics/NUMERICS.md §7. Main changes: §5 is rewritten around (H2); §8, §9, §10 are updated; small consistency edits elsewhere. It does not edit v1, v2, or v3. It is not itself verified as a whole document. It adds no claim beyond the cited verifier verdicts. One deliberate tag change: the logic of Theorem B (Route B) is tagged [proved+verified, conditional] per BETA_FP_PW_VERIFY §6 items 1–4; see the change log. The change log is at the end (Appendix).

Tag legend.

- [proved+verified]: proved in a project file and confirmed by an independent verifier file.
- [proved+verified, conditional on X]: the implication is confirmed; X is open or assumed.
- [proved, unverified]: proved in a project file; no verifier has graded it yet.
- [Lean]: formalized in Lean 4 + Mathlib, no sorry, axioms propext, Classical.choice, Quot.sound only.
- [verified quote]: read on the page image of Balaban's paper (reports/BALABAN_EXCERPTS_CHECK.md or a verifier file).
- [numerical, reproduced]: floating-point result reproduced by a verifier. [numerical, run by the verifier]: run once, by a verifier. [numerical]: not reproduced. None is certified.

[1] Project files named in this chapter live in the notes/ directory of the yang-mills repository.

- [argued]: a plausibility argument, not a proof. [open]: no proof. [refuted]: shown false.
- [Balaban]: an input taken from Balaban's published papers, used as stated there.
- [not written]: a proof plan; no project file contains the proof.

Range of L. Balaban assumes L odd, $L > 11$, as a standing assumption for the whole of CMP 109 (pp. 251–252) [8]. His construction of $\beta_j(g)$ for $g > 0$ and his inductive assumption (H0) are therefore available only at odd $L > 11$. Every statement below that uses (H0) or $\beta_j(g)$ at $g > 0$ is stated for odd $L > 11$, or carries "(H0) and Balaban's construction at this L" as an explicit hypothesis. Statements about the one-loop quantities $\beta_j(0)$, S_k, and the symbols σ_j hold for odd $L \geq 3$.

1 Statement

1.1 Balaban's statements [verified quote]

Theorem 2 (CMP 109 (1987), p. 259): "Let $d = 4$, $G = SU(2)$, and let γ be a sufficiently small positive constant, then for a sufficiently small positive g there exists a bare coupling constant $g_0 = g_0(\varepsilon, g)$ such that the sequence of the effective coupling constants g_k is contained in the interval $]0, \gamma]$, and $g_K = g$. Moreover, there exist constants β, β', $0 < \beta \leq \beta'$, such that g_k satisfy the inequality $1/g^2 + \beta \log(L^k\varepsilon)^{-1} \leq 1/g_k^2 \leq 1/g^2 + \beta' \log(L^k\varepsilon)^{-1}$. (0.31)" Balaban adds that the proof "will be given in a separate paper". It was never published (CMP 122 II p. 355, verbatim: "has not been published yet, so we have Theorem 1 with the assumption").

Theorem 1 (CMP 122 II p. 355): "If the sequence of the effective coupling constants is contained in an interval $]0, \gamma]$ with a sufficiently small positive γ, then the effective densities ρ_k have the form, and satisfy all the conditions and bounds, described in Sect. 2 [III]." The bound (0.1) there reads $\chi_k \exp[-(1/g_k^2)A(U_k) - E_-|T_\eta|] \leq \rho_k \leq \exp E_+|T_\eta|$ (corrected reading, `BALABAN_EXCERPTS_CHECK` item 3).

Setting. $\varepsilon = L^{-K}$, so $\log(L^k\varepsilon)^{-1} = (K-k)\log L$. $u_k := 1/g_k^2$. Recursion (CMP 109 (0.18)/(0.20)): $u_{k+1} = u_k - \beta_{k+1}(g_k)$. The increment β_{k+1} depends on the whole history $g_0, \ldots, g_k$ (CMP 109 p. 298, [verified quote]). $\beta_j(0)$ is the one-loop (Gaussian) increment; it is history-independent by definition. Balaban assumes L odd, $L > 11$, for the whole paper (CMP 109 pp. 251–252, OCR line 124, verbatim confirmed in `THEOREM2_PROOF_VERIFY` Step 6).

1.2 What Balaban's Theorem 1 actually needs (`LEMMA_DPRIME_SWEEP` §2, §7)

The quoted hypothesis of Theorem 1 is only $g_k \in]0, \gamma]$. The proof uses more. CMP 119 Theorem 1 (p. 262) assumes "(I.0.33)", a misnumbered reference to (0.31) of CMP 109. Its working form is CMP 119 p. 255–256 (2.6)–(2.9), for scales $m < n$:

- (2.6) $g_n \leq (1 + g_n^2\beta'(n-m))^{1/2} g_m$ and $g_m \leq (1+\beta_0) g_n$;
- (2.7) $(\log g_n^{-2})^p \leq (1+\beta_0)(\log g_m^{-2})^p$ and $(\log g_m^{-2})^p \leq (1 + g_n^2\beta'(n-m))^{\beta_0}(\log g_n^{-2})^p$;
- (2.8), (2.9) the same for $\varepsilon_j = g_j p_0(g_j)$ and R_j.

Here β_0 is Balaban's small constant (1/7 in CMP 122 II p. 389), not a coupling. [verified quote] for (2.6)–(2.9): page images, `LEMMA_DPRIME_SWEEP` §1; the sweep itself is single-author, see §2.3.

In u-variables these follow from two pairwise bounds:

- (P-up) $u_m \le u_n + B'(n-m)$, with no additive constant;
- (P-low) $u_m \ge u_n + \beta(n-m) - D'$, with $\beta > 0$, $D' \ge 0$ fixed,

plus the smallness condition

- (S) $\gamma^{-2} \ge c_{sw} D'$, $c_{sw} = \max(2, 1/(1-(1+\beta_0)^{-2})) = 64/15$ at $\beta_0 = 1/7$, and $\gamma \le \gamma_*(\beta_0, \beta, B', p_0, p_1, r, \nu, \kappa_0)$ (the conditions that literal (0.31) also needs). γ_* is free of the D'-ratio conditions, but it depends on R through B' (it contains $\gamma^2 B' \le (1+\beta_0)^2 - 1$ with $B' = 5b/4 + 2R$). The order of constants in §6 (R before γ) handles this.

β_0 must be fixed before γ.

1.3 The weak form proved conditionally in this chapter

Theorem (Theorem 2w (weak form))**.** *Let $G = SU(2)$, $d = 4$, L odd, $L > 11$ (equivalently: L odd ≥ 3 with (H0) and Balaban's construction of $\beta_j(g)$ at this L as an explicit hypothesis). There are $\gamma > 0$, $g_1 \in]0, \gamma]$, $\beta > 0$, $B' \ge \beta$, $D' \ge 0$ such that for every $K \ge 1$ and every $g \in]0, g_1]$ there is a bare coupling $g_0 = g_0(K, g)$ with*

- *(b) $g_k \in]0, \gamma]$ for $0 \le k \le K$, and $g_K = g$;*
- *(c′) for all $0 \le m \le n \le K$: $u_n + \beta(n-m) - D' \le u_m \le u_n + B'(n-m)$;*
- *(S) holds for γ, D'.*

Consequence: Balaban's (0.31) holds up to the additive constant D' on the lower side, with constants $\beta/\log L$ and $B'/\log L$. The two-sided bound fixes $1/g_k^2$ up to constants, as required.

Status: [proved+verified, conditional on (B2) and the fixed-level inputs of §4] by Route K (Theorem K + Lemma D″). (B2) is reduced to the single inequality (H2) of §5 [proved+verified reduction: `B2_TWOSTEP_VERIFY` grade 3]. Second route: Route B (`BETA_FP_PW` Theorem 3.2 + Lemma D″). The logic of Theorem B is [proved+verified, conditional] (`BETA_FP_PW_VERIFY` §6 items 1–4). Its inputs (U), (DF), (T1) from `BETA_FP_U` are [proved, unverified]; the route is conditional on (U), ($B\theta$), and the plane-wave defects. `BETA_FP_U` reduces (U) to (T1) + (H2) [proved, unverified]. Both routes also need (H0) at this L and L5 (§2.4). See §4.

1.4 Literal (0.31)

- (L1) Literal (0.31) for every $K \ge K_0$ and every $g \in]0, \gamma]$. Needs eventual uniform positivity: there are J and $\beta_* > 0$ with $\beta_j(0) \ge \beta_*$ for all $j > J$, plus $\sup_j |\beta_j(0)| < \infty$ (Lemma D′ (d), §2.1). [proved+verified; Lean] as an implication. Theorem K and Theorem B do NOT give eventual positivity: they bound window sums, not single increments. Eventual positivity follows from the convergence route (§7), which is open.
- (L2) Literal (0.31) for every $K \ge 1$. Needs $\inf_j \beta_j(0) > 0$ and $\sup_j \beta_j(0) < \infty$ (Lemmas D + E). If $\beta_j(0) \to \beta_\infty > 0$, then "every $\beta_j(0) > 0$" is equivalent to $\inf_j \beta_j(0) > 0$ (Lemma D′ Step 8). [proved+verified; Lean] as an implication. Inputs open.
- Literal (0.31) is not needed for Theorem 1: §1.2 shows the proof uses (c′) + (S). Literal (0.31) is used directly in one place only, the alternative choice of ε_0 in CMP 119 p. 248, which Balaban does not adopt.

- Necessity. Literal (0.31) at every $K \geq 1$ forces $\beta_K(0) \geq \beta \log L > 0$ for every K (window $k = K-1$, let $g \to 0$ using (H0); so L odd > 11). [proved, unverified]; elementary; `THEOREM2_PROOF_VERIFY` Point 5(a) agrees. So certified early steps are needed only for (L2), not for Theorem 1.

2 Shooting lemmas

Notation: $u_k = 1/g_k^2$, $t = 1/g^2$, $T(k,K) := \sum_{i=k+1}^{K} \beta_i(g_{i-1})$. Clamp $\varphi(u) := \max(u, 1/\gamma^2)^{-1/2}$. Lemmas D′, D″, D, E are abstract statements about a sequence of functions β_j. Applied to Balaban's $\beta_j(g)$, they need L odd > 11 (or (H0) at this L) and L5.

2.1 Lemma D′ (shooting without early positivity) [proved+verified; Lean]

Sources: `LEMMA_DPRIME.md` §1–§2; `LEMMA_DPRIME_VERIFY.md` grades 1, 2, 5 (CONFIRMED). Step 7 of that file also holds grades 3, 4, 6, which are CORRECTED (Remark 1 overclaim, misreads in §5, ledger); this chapter uses none of them except through the sweep (§2.3). Lean: `lean/YmLean/LemmaDPrime.lean` (`YmLean.lemma_DPrime`, 405 lines), statement check `lean/VERIFY_LemmaDPrime.md`.

Lemma (Lemma D′)**.** Hypotheses. *(i) β_{k+1} continuous on $[0,\gamma_0]$ and $|\beta_{k+1}(g) - \beta_{k+1}(0)| \leq C_1 g$, C_1 independent of k, K. (ii) $\sup_j |\beta_j(0)| \leq B$. (iii′) there is J with $\beta_j(0) \geq \beta_\infty/2$ for all $j > J$ (this is all the proof uses of "$\beta_j(0) \to \beta_\infty > 0$").*

Constants. $\gamma := \min(\gamma_0, \gamma_{Bal}, \beta_\infty/(4C_1))$; $B' := B + \beta_\infty/4$; $D := J \cdot B'$; $D' := J(B + \beta_\infty/2)$; $g_1 := (1/\gamma^2 + D)^{-1/2}$; $K_0 := J + \lceil 8D'/\beta_\infty \rceil$.

Conclusions.

- *(a) $T(k,K) \geq -D$ for all $k < K$ along any couplings in $]0,\gamma]$.*
- *(b) For every $K \geq 1$ and $g \in]0,g_1]$ there is g_0 with all $g_k \in]0,\gamma]$ and $g_K = g$.*
- *(c) $u_K + (\beta_\infty/4)(K-k) - D' \leq u_k \leq u_K + B'(K-k)$. The upper side has no constant.*
- *(d) For $K \geq K_0$: literal (0.31) with lower constant $\beta_\infty/8$, and shooting for every $g \in]0,\gamma]$. If every $\beta_j(0) > 0$: literal (0.31) for all K (Step 8 = Lemmas D + E).*

Proof idea. Clamped flow $\tilde{u}_{k+1} = \tilde{u}_k - \beta_{k+1}(\varphi(\tilde{u}_k))$ is continuous in u_0. Its tail sums lie in $[-D, B'(K-k)]$ for every u_0, because clamped couplings lie in $]0,\gamma]$. IVT on $[t-D, t+B'K]$ gives $\tilde{u}_K = t$. Then $\tilde{u}_k \geq t - D \geq 1/\gamma^2$ for $g \leq g_1$, so the clamp is never active. The bound is derived for the clamped flow first; there is no circularity. For history-dependent β this proof needs β_{k+1} defined and continuous on the whole cube $]0,\gamma]^{k+1}$; see L5 in §2.4.

Lean. The Lean hypotheses are weaker (ContinuousOn instead of C^1; (iii′) instead of convergence). Step 8 is `lemma_DPrime_step8` (`lean/YmLean/LemmaDE.lean`). Lean covers one-argument $\beta_j(g)$ only, not the history form.

Remark (Remark 4). Without $g \leq g_1$, (b) can fail at small K ($\beta_1 = \beta_2 = \beta_3 = -0.8$, $K = 3$, $g = \gamma$). [proved+verified]

2.2 Lemma D″ (tail-sum form; no index J) [proved+verified; Lean]

Source: `KAPPA_BOUNDED.md` §5.3; `KAPPA_BOUNDED_VERIFY` §6 (CONFIRMED: "No J needed"); hand rederivation in `THEOREM2_PROOF_v2_VERIFY` §A.

Lemma (Lemma D″)**.** Hypotheses. *(i) as in 2.1, and the window bound*

$$\textit{(5.1)}\quad T_0(k,K) := \sum_{i=k+1}^{K} \beta_i(0) \in [\, b(K-k) - 2R,\ b(K-k) + 2R\,] \quad \textit{for all } 0 \le k < K,$$

with $b := c \log L > 0$ *and some* $R \ge 0$*. (Route K gives* $R = \kappa_*$*; Route B gives* $R = C_B$*.)*

Constants. $\gamma := \min(\gamma_0, \gamma_{Bal}, b/(4C_1))$*;* $\beta := 3b/4$*;* $B' := 5b/4 + 2R$*;* $D = D' := 2R$*;* $g_1 := (1/\gamma^2 + D)^{-1/2}$.

Conclusions. *(b) of 2.1, and the pairwise bound*

$$(c')\quad u_n + \beta(n-m) - D' \le u_m \le u_n + B'(n-m), \quad 0 \le m \le n \le K,$$

upper side without constant.

Proof. $|T - T_0| \le C_1\gamma(K-k) \le (b/4)(K-k)$. Add (5.1). Steps 2–5 of Lemma D′ use only continuity and the bracket $\tilde{T} \in [-D, B'(K-k)]$; this bracket now comes from (5.1). The pairwise form is the same window argument with $K \to n$. The proof is the clamped-flow proof of 2.1; for history-dependent β the L5 caveat of §2.4 applies.

Lean: `lean/YmLean/LemmaDDoublePrime.lean` (`YmLean.lemma_DDoublePrime`, 146 lines, no `sorry`, axioms propext, Classical.choice, Quot.sound; `lean/README.md`). It proves (a″), (b) with the clamp inactive on the solution, (c′) with $\beta = 3b/4$, $B' = 5b/4 + 2R$, $D' = 2R$, and (d) below. The Lean hypotheses are weaker than the note: continuity instead of C^1; any γ with $0 < \gamma \le \gamma_0$ and $C_1\gamma \le b/4$; $R \ge 0$ derived; $K = 0$ allowed; (c′) and (d) hold for every trajectory with couplings in $]0, \gamma]$. One-argument β only (L5 not formalized).

(d) [Lean]: for $n - m \ge 8R/b$, $u_n + (b/2)(n-m) \le u_m \le u_n + B'(n-m)$. So literal (0.31) from (5.1) alone holds, with lower constant $b/2$, only for windows with $n - m \ge 8R/b$; short windows near K are not controlled (see §1.4).

2.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [proved, unverified (sweep); 4 of 19 table rows verified]

Source: `LEMMA_DPRIME_SWEEP.md` (all 12 Balaban files; OCR grep plus page image for every non-single-scale hit). `LEMMA_DPRIME_VERIFY` Steps 4–5 checked comparisons A–E on page images (CMP 122 I p. 192, p. 198 (1.94); II p. 381 ×2, p. 386) and corrected two misreadings of A (factor $(1+\beta_0)^2(k-j)^{\beta_0}$; $\delta'_j = g_j A_1 p_1(g_j)$). The five comparisons map to four table rows: A = row 11, E = row 12, C and D = row 16, B = row 18.

Result. 19 cross-scale uses of the coupling sequence (table in SWEEP §7), all in CMP 119 and CMP 122 I–II; none in CMP 95, 96, 98, 99, 102, 109, 116. Each is covered by (b) + (c′) + (S) of §1.2:

- items 1, 7, 10, 12, 17 need (P-up) with no constant (e.g. (2.6) line 1 is exactly (P-up));
- items 2, 3, 11, 13, 15, 16, 18 need (P-low) with fixed multiplicative slack $(1+\beta_0)$ or $(1+\beta_0)^2$; item 2 ((2.6) line 2) gives the constant 64/15; item 11 needs only $2.42D'$;
- items 4 ((2.8)–(2.9), ε and R ratios), 6 (CMP 119 (3.7), (3.8), (3.12), (3.13)), and 9 (CMP 122 I (1.52)), the last two of the form $\varepsilon_{k+1} \le (1+\beta_0)\varepsilon_k$, use both sides; they follow as items 1 and 3 do (item 4 from items 1–3);
- item 14 (CMP 122 I p. 200, R_{k-N_0+1} vs R_k) uses the lower side; it follows as item 3 does;

- item 5 (CMP 119 (2.31), (2.46): $\sum_j g_j^{\kappa_0} \le g_n^{\kappa_0-6}$) needs linear growth (P-low) and $\gamma^{-2} \ge 2D'$;
- item 0 is CMP 119 Theorem 1 assuming (0.31); its proof works with (2.6)–(2.9);
- item 8 (CMP 119 p. 284) is in a sketch, not load-bearing.

Caveats [open]: (1) the sweep is a grep plus page images, not a line-by-line read; (2) "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) was not located; (3) Balaban says β_0 "can be chosen arbitrarily small, if g is sufficiently small"; under (P-low) the floor is $\beta_0 \gtrsim D\gamma^2/2$, so β_0 must be fixed first; no step was found that needs $\beta_0 \to 0$; (4) the sweep has no independent verifier beyond rows 11, 12, 16, 18; (5) `BALABAN_EXCERPTS_CHECK` item 11 lists readings not checked on page images: Π (4.37)/(5.1), CMP 119 (2.42), the (2.12)/(2.23) ambiguity, the "L odd" page, the Sect. 3 estimates, and the 1989 bound lists.

2.4 Lemma D and Lemma E [proved+verified; Lean]

Sources: `LEMMAS.md` (corrected), `LEMMAS_DE_VERIFY.md` §1.4, §1.5, §2.4; `lean/YmLean/LemmaDE.lean` (`lemma_D`, `lemma_E`, `lemma_DE`).

- **Lemma D.** If $b \le \beta_j(g) \le b'$ on $]0,\gamma]$ with $b > 0$, then for every K and $g \in]0,\gamma]$ shooting succeeds and $t+b(K-k) \le u_k \le t+b'(K-k)$. This is literal (0.31). The first IVT interval in the draft was wrong and was refuted by the verifier; the corrected interval is $[t+bK,\, t+b'K]$. The nested-interval variant for β defined only on admissible histories (`LEMMAS_DE_VERIFY` §1.5) is [proved+verified], not in Lean.
- **Lemma E.** If $\beta_0 \le \beta_j(0) \le \beta_0'$ and $|\beta_j(g) - \beta_j(0)| \le Cg^2$, then for $\gamma \le \sqrt{\beta_0/(2C)}$: $\beta_0/2 \le \beta_j(g) \le \beta_0' + \beta_0/2$. The verifier checked this Cg^2 form; the C^1 form ($C_1 g$, $\gamma \le \beta_0/(2C_1)$) is the same two-line proof [proved, trivial variant]. Lean has the Cg^2 form.
- **History dependence (ledger item L5)** [open]. Balaban's β_{k+1} depends on $(g_0, \dots, g_k)$. Lemma D′ Remark 3 states the cube-continuity version (same proof). The gap is sharper than v2 stated (`THEOREM2_PROOF_v2_VERIFY` §F):
  - *Domain.* Balaban defines β_{k+1} only on histories for which his Theorem 1 induction has run to step k (`LEMMAS_DE_VERIFY` §2 item 1). By the sweep (§2.3), that induction uses (2.6)–(2.9), that is (c′) + (S), on the history, not only $g_i \in]0,\gamma]$.
  - *The new gap.* The clamped shooting of §2.1–§2.2 evaluates β on clamped histories $\tilde{g}_i = \varphi(\tilde{u}_i)$. A clamped history can violate the growth side of (c′): from $\max(a,c) - \max(b,c) \ge \min(a-b, 0)$ one gets only $u_m \ge u_n - D'$, not $u_m \ge u_n + \beta(n-m) - D'$. So the clamped shooting evaluates β on histories outside Balaban's domain. The clamp proof is valid only under the cube hypothesis, which Balaban does not supply.
  - *A variant that avoids it* [not written]. Use a nested-interval proof in the style of `LEMMAS_DE_VERIFY` §1.5, with domain $D_k = \{u_0, \dots, u_k \ge \gamma^{-2}\}$ of true (unclamped) trajectories. On such trajectories, (c′) up to step k follows from the tail-sum bound (a″) of Lemma D″, and (S) holds by the choice of γ. So every evaluation of β stays inside Balaban's domain. At each step k, keep the set of u_0 whose trajectory stays in D_k; show by IVT, step by step, that this set contains a closed interval on which u_k sweeps a range that contains the next target interval. For Lemma D (all increments positive) this is written. For Lemma D″ the increments can be negative, so the nested intervals need new envelopes, built from the window bounds $[b(n-m) - 2R - (b/4)(n-m), \dots]$. The adaptation is plausible; it is not written and not verified.

– Lean is one-argument only.

3 The exact structure of the one-loop sums

3.1 Telescoping [proved+verified, corrected scope]

$S_k := p \to 0$ F^2-coefficient of the composed k-step gauge-fixed one-loop integral (U1 (1.2)). Then $S_k = \sum_{j=1}^{k} \beta_j(0)$, $S_0 = 0$ (`BETA_FP` Prop. 1.1; `BETA_FP_VERIFY` item 1). Scope: exact for the $p \to 0$ coefficient at fixed k, not for $\log Z_k$ at fixed p (the plane wave is off-shell).

3.2 Prop. 2.1, anchor-free identity [proved+verified]

Source: `BETA_FP_LEMMA43` Prop. 2.1; `BETA_FP_LEMMA43_VERIFY` items 1, 3–4 (hand rederivation; lattice check with the real CMP 109 averaging at random SU(2) backgrounds, relative difference 10^{-14}).

Under (G1) $= \{\Delta D = 0,\ \tilde{Q}D = D_c \iota_a^T\}$:

$$\log Z_\Lambda(V) = \big[\log\det X - \tfrac{1}{2}\log\det K\big] - \big[\log\det(D_c^T D_c) - \tfrac{1}{2}\log\det(\Delta_\Lambda(V) + D_c D_c^T)\big].$$

In words: constrained one-loop = fine Wilson one-loop in background Feynman gauge minus the one-loop of the coarse Gaussian theory with kernel $\Delta_\Lambda(V)$. All point-anchor objects cancel; the constant is exactly 0.

- $\tilde{Q}D = D_c\iota_a^T$ is exact at any background (lattice 10^{-15}). $\Delta D = 0$ is load-bearing (raw Hessian gives $O(1)$ errors).
- At the constant abelian field V_f the background is critical, so $\Delta D = 0$ is exact and Prop. 2.1 holds exactly (`BETA_FP_H_VERIFY` §2, CONFIRMED). At a plane wave it holds up to the defect $\mathfrak{d}_Y$ (`BETA_FP_PW` Lemma 2.3) [proved, unverified].
- Per-cell infinite-volume densities only: on a torus the constant zero modes must be removed. This is input (G2) [formal].

3.3 The constant c [proved+verified]

$c = 11N^2/(12\pi^2) = 2N \cdot b_0/(8\pi^2)$, $b_0 = 11N/3$; SU(2): $c = 0.371511$. Normalization $1/g_B^2 = 2N/g^2$ (`R2_VERIFY`; `BETA_FP_LEMMA43_VERIFY` §5). The log coefficient of the fine piece at constant field is $c_2 = c/2$ per $\ln(1/\varphi)$, checked against Savvidy, arXiv:1910.00654 eq. (1.4) (`BETA_FP_H_VERIFY` §4) = input (F) [cited, source checked; lattice = continuum log coefficient is standard, not proved in project files].

$b := c\log L$: 0.25751 ($L = 2$), 0.40815 (3), 0.51502 (4), 0.59793 (5), 0.72293 (7).

3.4 The decomposition $S_k = ck\log L + \kappa_W^F - \kappa_k$

- As bookkeeping the split is empty: any c' gives an exact identity if κ_k is defined as the remainder (`U1_VERIFY`). Boundedness of a remainder-defined κ_k is a restatement, not progress.
- Content comes from defining κ_k by the coarse theory alone. `KAPPA_BOUNDED` Prop. 1.1 [proved+verified, conditional on (PI), (F), (M_j), V1, (G2)]: with the Wilson level $j = 0$ as reference,

$$S_j = cj\log L - d_j, \qquad \kappa_j = \kappa_W^F + d_j,$$

$$d_j := -\frac{1}{2n_2}\int_0^1 dt \int_{\mathrm{BZ}} \frac{dK}{(2\pi)^4}\ \mathrm{tr}\big[b_{t,2}\,\delta\sigma_j^{(0)} + b_{t,1}\,\delta\sigma_j^{(1)} + b_{t,0}\,\delta\sigma_j^{(2)}\big],$$

where $\delta\sigma_j^{(m)} = \sigma_j^{(m)} - \sigma_0^{(m)}$ are differences of the f^m Taylor symbols of the coarse kernel at level j and level 0, and $b_{t,m}$ are the Moyal inverse coefficients of the interpolated coarse operator. No limit symbols σ_∞ are used.

- Checks [numerical, reproduced]: $\rho(1) = \kappa_W^F$ (Lüscher–Weisz (5.7) [16]); $L = 2$: $d_1 = -0.372$, $\kappa_1 = 1.114$; $L = 3$: $d_1 = c\log 3 - \beta_1 = -0.256 \pm 0.01$ (`KAPPA_BOUNDED_VERIFY` §4: fits give $-0.254 \cdots - 0.258$). This checks the definition of d_j through S_j; it does not evaluate (1.1) from the symbols independently.

- Cor. 3.1 of `BETA_FP_LEMMA43` ($\mathcal{C}(\Lambda) = c\log\Lambda + \kappa_W^F - \rho(\Lambda)$) is exact as an identity. "The whole $\log\Lambda$ sits in the fine piece" is equivalent to $\rho(\Lambda) = o(\log\Lambda)$. It is not a separate fact (`BETA_FP_LEMMA43_VERIFY` item 2; v1 overclaim removed).

- Given convergence $\beta_j(0) \to \beta_\infty$, Cesàro gives $\beta_\infty = c\log L - \lim \kappa_k/k$. So v1's (H′) "$\kappa_k = o(k)$" is EQUIVALENT to $\beta_\infty = c\log L$, not weaker (`THEOREM2_PROOF_VERIFY` Point 4). Theorem K (§4) gives κ_k bounded, hence $\kappa_k = o(k)$ trivially.

3.5 Level-uniform symbol facts used below [proved+verified]

Source: `U3_AW.md`, `U3_AW_VERIFY.md` (grades 1–4 CONFIRMED); `U3_REPAIRS_VERIFY`.

- O5/A0: Δ_M real-analytic on a fixed strip, $|\partial^\alpha(\Delta_M - \Delta_\infty)| \le CM^{-2}$, uniform to $K = 0$ (odd L, centered blocks). Prop. B: $c_0|\hat{K}|^{-2}P_T' \le P_T' S_M P_T' \le C_1|\hat{K}|^{-2}P_T'$, $c_0 = (2/\pi)^{12}$, $C_1 \le 1363$. Existential constants.

- Prop. E (evenness): $\sigma_j(f;-K) = \sigma_j(f;K)$ in the midpoint trivialization, all f, j. Tested in $d = 2$ only.

- Theorem W: $\sigma_j^{(1)}(0) = \tau_c = (3ie/2)(E_{10} - E_{01})$ for every $j \ge 0$, from the exact Ward identity $\Delta_j(f)D_f = 0$. Numerically 0.750000, 0.75015, 0.75051 at $j = 0, 1, 2$ vs 0.75 ($d = 2$ proxy).

- Theorem A3: $|\sigma_j^{(1)}(K) - \tau_c| \le C_A|K|^2$, real K, uniform in j (C^0 only).

- Prop. A2: $\sigma_j^{(m)}$, $m \le 2$, analytic on the same strip at every fixed j (not uniform in j for $m = 1$ derivatives, $m = 2$).

- Lemma M: Maxwell coefficient exactly 1 at every level.

4 Bounded window sums: two formally distinct conditional routes

Both routes give the window bound (5.1) of §2.2, $|S_K - S_k - b(K-k)| \le 2R$, with $b = c\log L$. Lemma D″ then gives Theorem 2w (at odd $L > 11$, or with (H0) at this L as a hypothesis, and with L5). Neither route uses convergence of $\beta_j(0)$, a rate, (N), or certified early steps. The two routes have formally distinct named hypotheses. Their mathematical content is not independent: both reduce their uniform-in-level content to the same inequality (H2) on the two-step minimizer map (§5), plus A3-type facts (`BETA_FP_U` §3.5, §4.3 [proved, unverified]; Route K side: `B2_TWOSTEP_VERIFY` grade 3 [proved+verified]).

4.1 Route K: Theorem K [proved+verified, conditional on (B2) and the fixed-level inputs]

Source: `KAPPA_BOUNDED.md`; `KAPPA_BOUNDED_VERIFY.md` (items (i)–(v) and (vii) CONFIRMED; (vi) CORRECTED in reasoning, conclusion unchanged).

Theorem (Theorem K)**.** *L odd ≥ 3. Assume (B2) (§5; (B2) $\Leftarrow$ (H2)* [PROVED+VERIFIED]*) and the fixed-level inputs below. Then $\sup_k |d_k| \leq \kappa_* < \infty$, so $\sup_k |\kappa_k| \leq |\kappa_W^F| + \kappa_*$ and $|S_k - ck \log L| \leq \kappa_*$ for all k. Window sums satisfy (5.1) with $R = \kappa_*$. (Theorem K is a statement about one-loop quantities. Its use in Lemma D′ for Balaban's $\beta_j(g)$ needs L odd > 11 or (H0) at this L.)*

Proof structure.

1. Prop. 1.1 (§3.4): $S_j = cj \log L - d_j$. [proved+verified, conditional on (PI), (F), (M_j), V1, (G2)]
2. Lemma 2.1: the only K-derivative of $\sigma^{(1)}$ in d_j enters linearly with f^0 coefficients. Integration by parts moves it onto f^0 data; boundary terms vanish (periodic up to a common constant conjugation; inner sphere $O(r^2)$). So $\sigma^{(1)}$ is needed only in C^0, which Theorem A3 gives. [proved+verified]
3. Prop. 2.2: with (E0) f^0 bounds (Prop. B, O5/A0, Lemma M), (E1) Theorem A3 + Theorem W, and (B2), every term of the d_j integrand is $\leq C|K|^{-2}$, uniformly in j and t. In $d = 4$ this has two powers of infrared margin. $|d_j| \leq \kappa_* := (C(L)/n_2)(1 + Y_1^2 + C_A(1 + Y_1) + Y_2) \cdot I_2$. [proved+verified, conditional on (B2)]
4. Infrared: the degree -4 part of the coarse f^2 integrand is the same at every level (Lemma M + Theorem W), so the $\tau_c \log$ cancels exactly in level differences. [proved+verified]

Constants are existential; κ_* is not evaluated.

Fixed-level inputs of Route K (needed at each j separately; no uniformity in j):

- (PI) probe independence: plane-wave p^2 coefficient = constant-field f^2 coefficient [argued; per-cell limit for the linear potential not written].
- (F) fine log coefficient c_2 at constant field [cited, source checked, §3.3].
- (M_j) the formal f^2 coefficient d_j equals the true one at fixed j. The resolvent step of the first argument was false (Nielsen–Olesen zero at $k_\perp^2 = |ef|$; `BETA_FP_H_VERIFY` §5). `BETA_FP_M` repairs it with a direct $\log|\det|$ bound given spectral bounds (S1)–(S3) [open on the lattice]; Prop. 4.4 there is [argued]; `BETA_FP_M` has no verifier file. The t-subdivision of KAPPA §1.3 removes the small-ε requirement [argued, CONFIRMED as plausible]. Status: [argued].
- V1: the Maxwell coefficient of the per-step determinant equals $\beta_{j+1}(0)$, i.e. order-g^0 completeness of CMP 109 (1.3)–(1.5) [open; supported at order p^2 by `U3_VERIFY` §6].
- (G2) per-cell infinite-volume densities [formal].
- (H0) Balaban's inductive assumption: β_{j+1} smooth on $[0, \gamma]$, "uniformly bounded on this interval together with all derivatives" (CMP 109 pp. 263–264, (1.22)) [Balaban; available at odd $L > 11$ only]. CMP 116 has no theorem statement and no C^m-in-g bound in words (`BALABAN_EXCERPTS_CHECK` item 10); the smoothness in g is CMP 109's inductive assumption, asserted by CMP 109 Theorem 3, whose proof CMP 116 completes.
- L5 history-dependent β, its domain, and its continuity (§2.4) [open].

4.2 Route B: Theorem B of BETA_FP_PW [proved+verified, conditional (logic of Theorem B)]

Source: BETA_FP_PW.md §1–§5 (Theorem 3.2 = Theorem B, Theorem 5.1). BETA_FP_PW_VERIFY.md now has a §6 with grades: items 1–4 CONFIRMED (doubling identity and Lemma 1.3; Prop. 2.2 / Lemma 2.3; Theorems A, B as conditional logic; Lemma 4.1, conditional), items 5 (comparison) and 6 (numerics) CORRECTED. v3 stated that the file had no grade; that is outdated. Tag decision (PLAN.md, 2026-10-28): the logic of Theorem B is [proved+verified, conditional] on its hypotheses, citing BETA_FP_PW_VERIFY §6 items 1–4. Its inputs (U), (DF), (T1) from BETA_FP_U stay [proved, unverified]. The file rederives the Lemma 1.3 arithmetic and the Lemma 2.3 / Prop. 2.2 algebra correctly, confirms the Prop. 2.2 determinant identity on a 4^4 lattice (4e−13), and notes that identity 3.0 is a tautology ($\Gamma^{ex} := \hat{\Pi} - S$), so all content sits in $(B\theta)/(C\theta)$.

Theorem (Theorem B)**.** *L odd ≥ 3. Assume*

- *(U) there are $p_* > 0$, C_U with $|\beta_{j+1}(p) - \beta_{j+1}(0)| \leq C_U p^2$ for all $j \geq 0$, $0 < p \leq p_*$* [HYPOTHESIS; ARGUED FROM UNIFORM ANALYTICITY IN p OF THE ONE-STEP PLANE-WAVE DATA PLUS ONE-STEP O6]*;*
- *(Bθ) there is $\theta_* \in (0, \min(p_*, 1))$ with $B_* := \sup_k |\Gamma^{ex}(\theta_*; L^k)| < \infty$, where Γ^{ex} is the coarse remainder at fixed coarse momentum θ_** [HYPOTHESIS]*;*
- *Lüscher–Weisz (5.7) [16]* [CITED].

Then $|S_k - ck \log L| \leq C_B := B_* + c_2 \ln(1/\theta_*^2) + |\kappa_W^F| + \sup_{p \leq \theta_*} |\eta(p)| + C_U \theta_*^2/(L^2 - 1)$, and window sums satisfy (5.1) with $R = C_B$. $\theta_* < 1$ is required so that $c_2 \ln(1/\theta_*^2) \geq 0$; otherwise the term must be replaced by its absolute value (BETA_FP_PW_VERIFY §3).

Key step. Lemma 1.3: $|S(\theta; L^k) - S_k| \leq C_U \theta^2/(L^2 - 1)$ from (U). The massless coarse side is then needed only at a fixed $\theta_* > 0$, where there is no infrared problem. No (W), (PI), (M), IR weights, or spectral theory.

What $(B\theta)$ reduces to (BETA_FP_PW §4) [argued]: Lemma 4.1(i) (trivial-background convergence, from O5, odd L) [proved]; vertex bounds at fixed θ_* (the uniform bounds cited by U3 Theorem 7.1 to Balaban; the fixed-θ data hypothesis $(D\theta)$ is needed only in its "bounded" form); and boundedness in k of the defect $\mathfrak{d} = \mathfrak{d}_Y + \mathfrak{d}_{gf}$.

- $\mathfrak{d}_Y$ (Prop. 2.1 defect at the non-stationary plane wave): $O(a\Lambda^{-2})$ relative at fixed θ, from zero capacity of point anchors in $d = 4$ [argued; region $|K| \lesssim \Lambda^{-1}$ not written].
- $\mathfrak{d}_{gf}$ (composite gauge vs D_k^T gauge at the plane wave): $O(\theta^2)$ relative; uniformity in k not shown [argued]. Data: 0.953 (direct) vs 0.9595 (recursion) at $L = 2$, $p_0 = 0.2$; difference below the tabulation error, so unresolved.

Identification. Route B's $\beta_j(0)$ is the per-level plane-wave increment (the quantity the code computes). It matches Balaban's second-moment β up to the expansion point (fine wave vs constrained minimizer), argued to change S by $O(\theta^2)$ relative (SOURCES_D1_VERIFY item 8) [argued]. This identification and the $\mathfrak{d}_{gf}$ bound need expansion-point bounds uniform in k. So the (PI)-type content of Route K is not removed in Route B; it moves into a uniform-in-k input (BETA_FP_PW_VERIFY §3).

Three points from BETA_FP_PW_VERIFY §3 (item 5 of its §6 grades them CORRECTED):

1. $\theta_* < 1$ is required (above).
2. The β identification (B4) is a uniform-in-k input, not a fixed-level input.

3. (U) is a uniform-in-j analyticity of the level data. It plausibly needs the plane-wave analogue of a uniform C^1 bound on $\sigma^{(1)}$: uniform p-smoothness of the a^1 vertex. That is the same uniform first-order vertex smoothness as Route K's open core ($C1\sigma$).

4.3 Route B inputs after BETA_FP_U [proved, unverified; no verifier file]

Source: `BETA_FP_U.md` §2–§4. It reduces the two uniform hypotheses of Theorem B.

- (U) $\Leftarrow$ (T1) + (H2) (Theorem 2.4, odd $L \geq 3$, with A0, A1, Lemma 2.2). (T1): $\sup_j \sup_q |c^{(1)}_{T,j}(q)| < \infty$, the plane-wave form of Theorem A3 [argued]. (H2) is the inequality of §5 (`BETA_FP_U` writes it as $\tilde{h}_{(2)} < L$).
- ($B\theta$): $\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$ exactly (Prop. 3.1). γ_{rec} is bounded in k given Theorem 2.4 (Prop. 3.2). $\mathfrak{d}_F$ is bounded under (DF) $\sup_k |\mathfrak{d}_F(\theta_*; L^k)| < \infty$ [hypothesis, argued]. `BETA_FP_U` corrects `BETA_FP_PW` §4.3(b): $\mathfrak{d}_F$ is $O(1)$ relative at fixed θ, not $O(\theta^2)$; only boundedness is used.
- Corollary 4.1: given (T1), (H2), (DF), LW (5.7), $|S_k - ck \log L| \leq C_B$, so (5.1) holds with $R = C_B$.
- Theorem 2w by Route B then needs in addition A6 (plane-wave β = Balaban's β up to the expansion point, uniform in k) [argued] and (H0) (Lemma D″ (i)).

Route B hypothesis set in v4: (T1), (H2), (DF), LW (5.7), A6, (H0) (plus L5, (G2), odd L, as for Route K). The fixed-level inputs (PI), (M_j), V1 of Route K do not appear; (DF) and A6 replace them.

4.4 Comparison

	Route K	Route B
uniform-in-level input	(B2): $\sup_j \sup_K \|\sigma^{(2)}_j\| < \infty$, $\Leftarrow$ (H2) [proved+verified]	(U), ($B\theta$), and the β identification and $\mathfrak{d}_{gf}$ bound, both with k-uniform expansion-point bounds; (U) $\Leftarrow$ (T1) + (H2), ($B\theta$) $\Leftarrow$ Theorem 2.4 + (DF) [proved, unverified]
single uniform inequality	(H2)	(H2)
fixed-level inputs	(PI), (F), (M_j), V1, (G2)	LW (5.7), $\mathfrak{d}_Y$, $\theta_* < 1$
uses (W), Theorem A3	yes	no; `BETA_FP_U` uses (T1), the plane-wave analogue of A3 [argued]
verification	verified (conditional)	logic of Theorem B verified (conditional): `BETA_FP_PW_VERIFY` §6 items 1–4 CONFIRMED [proved+verified, conditional]; inputs from `BETA_FP_U` [proved, unverified], no verifier file
common	(H0) at odd $L > 11$, L5, (G2), V1-type identification, Balaban's proofs, SU(2)	

Shared inputs: (H0), L5, (G2), odd L (> 11 for (H0)), and the identification of the computed $\beta_j(0)$ with Balaban's $\beta_j(0)$ (V1; in Route B also the expansion point). The uniform-in-level hypotheses ((B2) vs (U) + ($B\theta$)) are formally distinct as named hypotheses. Their content overlaps:

both rest on (H2) (§5). In Route K, (H2) gives ($C1\sigma$) (Theorem C1) and then (B2) (Theorem B2). In Route B, (H2) gives (U) with (T1), and the vertex bound behind ($B\theta$). So a proof of (H2) serves both routes. One closed route suffices for Theorem 2w.

5 The hypothesis (B2) of Route K, and the single uniform inequality (H2)

(B2): $\sup_{j\geq 0}\sup_{K\in \mathrm{BZ}}|\sigma_j^{(2)}(K)| < \infty$. Weaker form that suffices, (B2w): $\sup_j \int_{\mathrm{BZ}}|K|^{-2}|\sigma_j^{(2)}(K)|\,d^4K < \infty$ ($\sigma^{(2)}$ enters d_j only undifferentiated; `KAPPA_BOUNDED` Remark 2.3) [proved+verified: not graded by name; implied by `KAPPA_BOUNDED_VERIFY` §9(iv)].

$\sigma^{(2)}$ obeys the one-step recursion (`U3_AW` (1.2)): $\sigma_j^{(2)} = L^{-2}\mathcal{A}_j\sigma_{j-1}^{(2)} + Q_j[\sigma_{j-1}^{(1)}] + g_j^{(2)}$, with $\mathcal{A}_j s = L^{-2}H_j^{\dagger}(\oplus_l s(k_l))H_j$ and $H_j(K)$ the one-step gauge-fixed minimizer map. Notation: $\rho_1 := \sup_{j,K}\|H_j(K)\|/L$ (one step); $H_{(2),j}(K) := (\oplus_l H_{j-1}(k_l))H_j(K)$ (two steps), $\rho_2 := \sup_{j\geq 2}\sup_{K\ \mathrm{real}}\|H_{(2),j}(K)\|/L^2$.

5.1 Proved

- $K=0$ bound [proved+verified]: $\sup_j|\sigma_j^{(2)}(0)| \leq C(|e|C_A + e^2)$. Theorem A3 + scaling give $|\partial^2\sigma_j^{(1)}(0)| \leq 2C_A$ (polarization; `KAPPA_BOUNDED_VERIFY` §7), and the slaving identity (W-2) expresses $\sigma_j^{(2)}(0)$ through these second derivatives.

- Each fixed j [proved+verified]: $\sigma_j^{(2)}$ is analytic on a fixed strip (Prop. A2), hence bounded. The content of (B2) is uniformity in j.

- Prop. 4.1 [proved+verified, conditional]: if (h) $h := \sup_{j,K}\|H_j(K)\| < L^2$ and ($C1\sigma$) $\sup_j\sup_K|\partial_K\sigma_j^{(1)}(K)| < \infty$, then (B2) holds. Proof: in the form norm $\pm s \leq N(\Delta+1)$, $N_j \leq (h^2/L^4)N_{j-1} + Z_2$, and Z_2 is finite by Theorem A3, ($C1\sigma$), and fixed f^0 vertices.

- Why the Lemma A4 mechanism alone fails [proved+verified, reason corrected by `KAPPA_BOUNDED_VERIFY` §7]: the obstruction is the missing uniform C^1 bound on $\sigma^{(1)}$, which enters twice: in Q_j (term $L^{-3}\partial_K\sigma_{j-1}^{(1)}$) and in the longitudinal part of $\sigma^{(2)}$ given by (W-2). Theorem A3 is C^0 only.

5.2 Refuted, and the failed one-step bound

- (HS) $h < L\sqrt{L-1}$ (named (H′) in `U3_AW`) [refuted at $L=3$]. Measured $\max_K\|H_j(K)\|/L =$ 1.652, 1.736, 1.757, 1.760 for $j=1,2,3,\infty$ (`U3_AW_VERIFY` §6); the threshold is $\sqrt{2}=1.414$. Gauge choice cannot help: H_j is already minimal-norm in its gauge orbit.

- One-step sup-norm C^1 recursion at $L=3$ [numerical, reproduced; `B2_TWOSTEP_VERIFY` grade 2]. Its factor is $\rho_1^2/L = 0.909$, 1.0046, 1.0289, 1.0325 for $j=1,2,3,\infty$, so it exceeds 1 for $j\geq 2$ ($\rho_1 > \sqrt{3} = 1.7321$). At $K=0$ the factor is L^{-1}; the bad region is the face $K_3 = K_4 = \pi$, $K_1 = K_2 \approx 1.86$–1.92. This shows that the one-step sup-norm bound fails at $L=3$. It does not show that a C^1 bound fails; a weighted norm is not excluded.

5.3 (H2) and the two-step theorems [proved+verified, conditional on (H2)]

Source: `B2_TWOSTEP.md` §1–§3; `B2_TWOSTEP_VERIFY.md` grades 1, 3, 4 (CONFIRMED).

$$\text{(H2)} \quad \rho_2 = \sup_{j\geq 2} \sup_{K\in \mathrm{BZ\ real}} \|(\oplus_l H_{j-1}(k_l))H_j(K)\|/L^2 < L.$$

(H2) is a plain operator-norm inequality on real K. It uses no strip, no Δ-relative weight, and only $n = 2$.

- Two-step map. $\mathcal{A}_j\mathcal{A}_{j-1}s = L^{-4}H^{\dagger}_{(2)}(\oplus s)H_{(2)}$, with $dk_{l,l'}/dK = L^{-2}$ [CONFIRMED, grade 1].
- Theorem C1: (H2) $\Rightarrow$ ($C1\sigma$), with contraction factor $q_1 = \rho_2^2/L^2$ on the derivative. The chain-rule gain is exactly L^{-2} per two steps on real K. So the threshold is $\rho_2 < L$, not $\sqrt{L^2-1}$: no strip-Cauchy loss enters. The large constants ($b(L)$ of O6, η_1, η_2, G_1) enter only the source term, not the contraction factor [CONFIRMED, grade 3].
- Theorem B2: (H2) $\Rightarrow$ (B2), hence (B2w), in plain sup norm, with factor $q_2 = \rho_2^2/L^4$. This contraction needs only $\rho_2 < L^2$, which (H2) implies [CONFIRMED, grade 3]. In one step it is Prop. 4.1's (h).
- (H2) vs (H″). (H2) is the plain operator norm of the same matrix as `U3_AW` (H″) at $n = 2$. It is weaker: threshold L (3 at $L = 3$) against $\sqrt{L^2-1}$ (2.828), real K only, $n = 2$ only [CONFIRMED, grade 4]. (H″) $\Rightarrow$ (B2) stays [argued] and is no longer needed.
- Form of Q_j (G3) [CONFIRMED by read-off of `U3_AW` (1.2)/(3.3) and P10]: $\partial\sigma^{(1)}$ enters linearly. `U3_LEMMA51` (3.3) itself was not re-derived from the Moyal rule by either party.
- Not covered: C^2 and C^3 bounds on $\sigma^{(1)}$ under (H2) (`B2_TWOSTEP` §5, "look routine", not written, not graded). They are not needed for (B2).
- Submultiplicativity: $\rho_2 \leq \rho_1^2$ always. At $L = 3$ the gain of the two-step map is that this is strict by a factor 1.50 (2.064 vs 3.098) [CONFIRMED].

5.4 Numerics for (H2) [numerical; none certified]

Code: `numerics/h2_scan.py` (umklapp Schur pieces, one colour, Gram form; `H2_LARGE_L` §1). Code structure CONFIRMED (`H2_LARGE_L_VERIFY` grade 1): the one-colour reduction is exact at $U = 1$, the H and Δ_{out} blocks and the Gram identity are correct, the separable Prop. A form is exact. Sign slip in `H2_LARGE_L` §4 ($H = +[\mathfrak{B}^{-1}]_{f,c}$); harmless.

L	j	ρ_2 (scan max)	threshold L	status
3	2	2.00857	3	[numerical, reproduced] (`B2_TWOSTEP_VERIFY` §4; `H2_LARGE_L_VERIFY` §1)
3	3	2.05601	3	[numerical, reproduced] (spot value 2.05599 at the argmax)
3	∞	2.0634 (2.06332–2.06377 over l_{max} 2–5)	3	[numerical, reproduced]; margin 31%; $q_1 \leq 0.473$, $q_2 \leq 0.053$

5	2, 3, 4, ∞	2.098, 2.108, 2.108, 2.108	5	[numerical, reproduced]: independent dense code gives 2.098283 at $j = 2$; the verifier's own search gives 2.107921 at $j = \infty$, same argmax; margin 58%
7	∞	$\geq$ 2.1131 (face diagonal only)	7	[numerical, run by the verifier] (spot 2.11309)
9	∞	$\geq$ 2.1142 (one point, $(2.6, 2.6, \pi, \pi)$)	9	[numerical, run by the verifier]; the author's $j = 2$ point gives ≥ 2.1104

- Shape. The argmax lies on the face $K_3 = K_4 = \pi$, on the diagonal $K_1 = K_2$: $\approx 2.22 \to 2.40$ ($L = 3$, $j = 2 \to \infty$), ≈ 2.54 ($L = 5$). Off-face points are lower: best off-face 1.995 ($L = 3$, $j = 2$), 2.004 ($L = 3$, $j = \infty$), 2.088 ($L = 5$, $j = \infty$). The corner (π, π, π, π) stays below the face peak, but the gap shrinks: 0.0070, 0.0028, ≥ 0.0025, ≥ 0.0023 ($L = 3, 5, 7, 9$). The corner is the obvious place for a competing maximum at large L.

- Level dependence. At $L = 5$, $\rho_2(\infty) - \rho_2(j) = 9.64 \cdot 10^{-3}$, $4.0 \cdot 10^{-4}$, $1.6 \cdot 10^{-5}$ ($j = 2, 3, 4$); ratios 24.1, 25.0, against $L^2 = 25$. At $L = 3$ the ratios are 7.5, 8.7. Levels $j = 4, \ldots, \infty - 1$ are not sampled at $L = 3$.

- L-trend. $j = \infty$: 2.0633, 2.1079, ≥ 2.1131, ≥ 2.1142 ($L = 3, 5, 7, 9$). This is consistent with a bounded ρ_2 near 2.11–2.14. The rate claim "at least as fast as L^{-2}" is CORRECTED (`H2_LARGE_L_VERIFY` grade 5): it used a lower bound at $L = 7$ as if it were the sup. "(H2) gets easier with L" is a measured trend for $L \leq 9$, not a result for $L > 11$.

- All values are lower estimates of a sup (grid plus local search).

5.5 One-step sufficiency at $L \geq 5$ [arithmetic proved; inputs numerical]

$\|H_{(2),j}(K)\| \leq \max_l \|H_{j-1}(k_l)\| \cdot \|H_j(K)\| \leq L^2 \rho_1^2$. So $\rho_1^2 < L$ implies (H2), and then the one-step C^1 recursion of `B2_TWOSTEP` (2.2) contracts with factor ρ_1^2 / L.

- $L = 3$: $\rho_1^2 = 3.10 > 3$. The one-step route fails (§5.2); the two-step map is needed.

- $L = 5$: $\rho_1 = 1.860$ ($j = 1$, Wilson fine kernel, dense, measured by the verifier), 1.9565 ($j = 2$), 1.963 ($j = \infty$). So $\rho_1^2 \leq 3.853 < 5$ and $\rho_1^2 / L = 0.771$ [numerical, CONFIRMED as a measurement, uncertified; `H2_LARGE_L_VERIFY` grade 4].

- $L = 7$: $\rho_1 \geq 2.032$ (diagonal only), $\rho_1^2 / L \approx 0.59$ [numerical].

So at $L \geq 5$ the one-step map suffices for Theorem C1, as far as measured. For certification this replaces L^8 blocks per K (two-step map) by L^4 blocks per K (625× fewer at $L = 5$).

5.6 Certification plan and j_0 [designed, not done]

`B2_TWOSTEP_VERIFY` grade 6 (CORRECTED in size): certifying (H2) is a computer-assisted proof project, not a check. `H2_LARGE_L` §4 gives the design; `H2_LARGE_L_VERIFY` grades 6–8 check it. Three parts.

(a) Tail $j \geq j_0$. From the resolvent identity, $\rho_2(j) \leq R_\infty + (\rho_1^2/a_*)(\delta_{j-2} + \delta_{j-1})$, where $R_\infty = \rho_2$ at $j = \infty$, $a_* = \inf_K a_j(K)$ with $\|[\mathfrak{B}^{-1}]_{f,f}\| = 1/a_j(K)$ exactly, and $\delta_m = \sup_K \|\Delta_{L^m} - \Delta_\infty\|$ (O5 decay) [derivation CONFIRMED]. Inputs CORRECTED by the verifier: the 2-norm sup of δ_m is at the corner, $\delta = 179, 92.0, 18.7, 2.39, 0.317$ ($M = 1, 3, 9, 27, 81$), 45–90× the assumed value at small m; $a_*(L = 5) = 0.0107$, not 0.05. With these inputs, $j_0 \approx 8$ ($L = 3$) and ≈ 6 ($L = 5$). With the O6 coercivity constant, $j_0 = 25$, which cannot be used in practice (`B2_TWOSTEP_VERIFY` §5 estimated $j \gtrsim 45$ with $\|\mathfrak{B}^{-1}\| \leq b(L) \sim 10^{19}$).

(b) Levels $2 \leq j < j_0$ and $j = \infty$. A certified sup over the BZ of $\|H_{(2),j}(K)\|$ (at $L \geq 5$: of $\|H_j(K)\|$, §5.5). Plan: interval arithmetic with local mean-value enclosures on boxes in the sorted chamber (1/384 of the BZ), 4×4 and 5×5 verified inverses, one 4×4 λ_{max} per K; about $2.5 \cdot 10^3$ boxes at half-width 0.05 at $L = 3$; the analytic form of Δ near $K = 0$, where $\rho_2(0) = 1$. At $j = \infty$ also a certified tail bound for the Prop. A sum (l_{max} truncation; observed effect $4 \cdot 10^{-4}$). There is no explicit global Lipschitz constant today: the A0/O5 strip width is existential, and the constant that can be made explicit is $\sim 10^{40}$. Monotonicity in j is not proved and is not used.

(c) Inputs to certify first: $\|\mathfrak{B}_j(K)^{-1}\| \leq B_{num}$ for all j, K (the O6 constant $b(L)$ is unusable), the O5 constant C_0 explicitly (or certified δ_m), certified a_*, ρ_1, R_∞.

Operational fixes for the Windows runs (`H2_LARGE_L_VERIFY` grade 8): atomic JSON writes, pre-copied profile caches, profile build time at $L \geq 11$ not budgeted. Nothing is launched.

5.7 The precise remaining inequality

At a fixed odd $L \geq 3$ the single uniform hypothesis is (H2).

- Route K: (H2) ⇒ ($C1\sigma$) (Theorem C1) ⇒ (B2) (Theorem B2) [proved+verified, conditional on (H2)].
- Route B: (H2) + (T1) ⇒ (U), and (H2) ⇒ the vertex bound for γ_{rec} in ($B\theta$) (`BETA_FP_U` Theorem 2.4, Prop. 3.2) [proved, unverified].
- Status of (H2): [numerical at $L = 3, 5, 9$; certification open]. At $L = 9$ there is one point only. No value exists at $L > 11$ (Balaban's range), so the numbers are supportive only.
- Kept as alternatives: ($C1\sigma$) + (h) ⇒ (B2) [proved+verified]; (H″) ⇒ (B2) [argued]; (B2w) directly. No per-momentum $d = 4$ constant-field code exists, so (B2) itself has not been scanned numerically (`KAPPA_BOUNDED_VERIFY` §8).

6 Conclusion

Theorem 2w ⇐ (Route K inputs) or (Route B inputs). Precisely:

Route K. (B2) [open; reduces to (H2) (verified reduction, §5.3); (H2) numerical at $L = 3, 5, 9$, certification open] + (PI), (F), (M_j), V1, (G2) [fixed-level] + (H0) [Balaban, odd $L > 11$] + L5 [open] ⇒ Theorem K ($R = \kappa_*$) ⇒ (5.1) ⇒ Lemma D″ ⇒ Theorem 2w. Status of the implication: [proved+verified, conditional].

Route B. (U) + ($B\theta$) with $\theta_* < 1$ + LW (5.7) + bounded $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ + identification of β (uniform in k) + (G2) + (H0) [odd $L > 11$] + L5 ⇒ Theorem B ($R = C_B$) ⇒ (5.1) ⇒ Lemma D″ ⇒ Theorem 2w. Status: Theorem B logic [proved+verified, conditional]; inputs [proved, unverified]. With `BETA_FP_U` (§4.2a): (T1) + (H2) + (DF) + LW (5.7) ⇒ Corollary 4.1 ($R = C_B$); with A6 + (H0) + L5 + (G2) ⇒ Theorem 2w. Status: [proved, unverified, conditional] (the inputs (T1), (DF), A6 and the `BETA_FP_U` reductions are unverified).

Single uniform hypothesis. Both routes need (H2). A certified (H2) at one odd $L > 11$ would discharge (B2), the only uniform-in-level hypothesis of Route K (Theorems C1, B2). In Route B it would reduce (U) to (T1); (DF) and A6 would remain.

Then Balaban's Theorem 1. Theorem 2w gives (b) and (c′). With γ shrunk so that (S) holds, every one of the 19 cross-scale uses in CMP 119 and CMP 122 I–II is covered (§2.3), and the single-scale uses need only $g_k \le \gamma$. So Theorem 1 holds for the bare coupling $g_0(K, g)$, for every K and every $g \in]0, g_1]$, without Balaban's coupling assumption. This needs L5 in the sharpened form of §2.4: β must exist on every history the shooting visits. Status: [proved, unverified (sweep), conditional on Route K or Route B inputs, on L5, and on Balaban's published proofs].

Order of constants. Fix $\beta_0, p_0, p_1, r, \nu, \kappa_0$ (Balaban). Fix L (odd, > 11), then R from Route K or B, then $b = c \log L$, then $\gamma \le \min(\gamma_0, \gamma_{Bal}, b/(4C_1), \gamma_*)$ with $\gamma^{-2} \ge c_{sw} \cdot 2R$. Then $g_1 = (\gamma^{-2} + 2R)^{-1/2}$. Balaban's phrase "for a sufficiently small positive g" fits $g \le g_1$.

What this does not give.

- Literal (0.31): only for windows of length $\ge 8R/b$. Literal (0.31) at every K needs (L1) or (L2) of §1.4, which need individual increments, not window sums. Not needed for Theorem 1.
- The value $\lim \beta_j(0)$ and any rate: §7.
- Even L, or odd $L \le 11$ without (H0) as an extra hypothesis (Balaban assumes L odd > 11 throughout CMP 109); SU(N) with $N > 2$.
- Monotone couplings: g_m can exceed g_n for $m < n$ by the bounded amount $u_m \ge u_n - D'$. §2.3 shows Balaban's proof tolerates this under (S); a line-by-line read was not done.
- A continuum limit (`ASSEMBLY.md` treats the assembly separately; not claimed here).

7 The convergence route: what it adds, and its status

Not needed for Theorem 1. It adds: (i) existence of $\beta_\infty = \lim \beta_j(0)$; (ii) its value $c \log L$; (iii) a rate; (iv) eventual positivity, hence literal (0.31) for $K \ge K_0$ (Lemma D′ (d)); (v) with certified early steps, literal (0.31) at every K (§1.4 (L2)). It also implies (B2), since convergence of $\sigma_j^{(2)}$ implies boundedness. So it needs strictly more than Route K.

7.1 Components and status

- U3 Theorem 4.1 with O5: $|\partial_K^\alpha(\Delta_M - \Delta_\infty)| \le CM^{-2}$, $|\alpha| \le 2$, uniform to $K = 0$, odd L [proved+verified] (`U3_VERIFY`, `U3_REPAIRS_VERIFY`). Numerically $|\Delta_M - \Delta_\infty| = 1.007, 0.129, 0.0147$ at $M = 3, 9, 27$.
- O6: $\|\mathfrak{B}_j(K)^{-1}\| \le b(L)$ for all j, K, one-step constraint, trivial background [proved+verified]. Crude constant ($b(3) \approx 7.5 \cdot 10^{33}$); measured corner values 103.5, 176.7, 193.0, 195.3 ($j = 0, 1, 2, \infty$, $L = 3$) use a surrogate Λ (`U3_REPAIRS_VERIFY` §5).
- Theorem 6.1: $\beta_{j+1}(0) = \Phi(\mathcal{D}_j)$ is Lipschitz in the level data [proved+verified, conditional on V1]. The norm is the Taylor-data norm (`U3_VERIFY` P8 correction).
- Lemma 5.1 (convergence of $\sigma_j^{(1)}$, $\sigma_j^{(2)}$ in C^2, $|\alpha| \le 2$):
  - neutral sector, rate L^{-4j} [proved+verified for the value bound; the K-derivative paragraph is proved, unverified];

  - (W) [proved+verified] (`U3_AW` Theorem W); it removes the marginal direction of the f^1 recursion;
  - (A) qualitative and Theorem A3 [proved+verified]; (A) in full (strip/C^3 for $m = 1$, all of $m = 2$) $\Leftarrow$ (H″) [open]; the route via (HS) is [refuted] at $L = 3$;
  - $\sigma^{(1)}$ convergence at rate jL^{-2j} [sketch in `U3_AW` §4.7, unverified]; $\sigma^{(2)}$ convergence [open, needs (A) in full];
  - the single-block-constraint version of Lemma 5.1 is false for $m = 2$ in $d = 4$ [proved+verified]; Balaban's composed constraint is not affected.

- Prop. 3.6: $|\beta_{j+1}(0) - \beta_\infty| \le C(1+j)^2 L^{-2j}$ [proved, unverified as an assembly; conditional on V1, (A) in full]. (v1 wrote Cj^2L^{-2j}, which is 0 at $j = 0$; corrected.) The identification of σ-Taylor data with the data of (6.1) is cited from `U3_VERIFY` §7, not re-shown. (N) (clean rate L^{-2j}) is not needed and is judged plausibly false.

- Value of the limit. Given convergence, Theorem K or Theorem B gives $\beta_\infty = c \log L$ at once: $S_k/k \to \beta_\infty$ by Cesàro and $S_k/k = c \log L + O(1/k)$ [proved+verified as an implication: `THEOREM2_PROOF_VERIFY` Point 4; `BETA_FP_PW` Theorem 3.2(iii)]. Older routes to the value: `BETA_FP_H` Theorem 3.1′ with (H) restated at the constant field, which given (W) reduces to Lemma 5.1 (1.2) alone (`BETA_FP_H_VERIFY` items (1)–(4)), plus (PI), (F), (M) [conditional; (M) argued]; and `BETA_FP_PW` Theorem 3.1 (limit form) given (U), ($C\theta$), LW (5.7) [proved, unverified]. The original (H) with IR weights (`BETA_FP_LEMMA43` (4.2)) was graded UNSUPPORTED and is not used.

- Rate of $\mathcal{C}(\Lambda)$: $O(\Lambda^{-2} \log \Lambda)$ given Lemma 5.1 with rates, via `U3_AW` §3.4(b) slaving [argued, `BETA_FP_H_VERIFY` §5].

7.2 Early steps and j_*

Under Prop. 3.6 and $\beta_\infty = c \log L$, $\beta_{j+1}(0) \ge (c \log L)/2$ for $j \ge j_* := \min\{J : C(1+j)^2 L^{-2j} \le (c \log L)/2 \text{ for } j \ge J\}$. Heuristic size [argued]: $C \sim b^3$ gives $j_* \approx 110$ ($L = 3$), ≈ 70 ($L = 13$); $C \sim b^6$ (one-step inverses inside C_{data}) gives ≈ 218, ≈ 138 (`THEOREM2_PROOF_VERIFY` Point 5(e)). With the measured $\|\mathfrak{B}^{-1}\| \approx 195$: $j_* \approx 10$ ($L = 3$), ≈ 4 ($L = 13$). Certifying $\beta_1(0), \dots, \beta_{j_*}(0) > 0$ needs interval arithmetic at odd $L \ge 13$ (blocks $\approx 3.4 \cdot 10^5$ rows per momentum by dense methods; the umklapp code reduces this to 12×12 blocks at L^4 momenta). It is needed only for (L2), not for Theorem 1.

8 Numerics

SU(2), infinite volume, Balaban's normalized-trace units. None of this is certified. Only odd L is inside the proofs; $L = 2, 4$ use corner blocks and are outside them.

8.1 First increment $\beta_1(0)$ (`NUMERICS` §6–§7; `UMK_BETA1.md`)

L	$\beta_1(0)$	how	$b = c \log L$	$\beta_1 - b$	status

2	0.630	$N_g = 12$, $p \to 0$ fit	0.2575	+0.372	[numerical, reproduced] (KAPPA_BOUNDED_VERIFY §4: 0.6296)
3	0.664	$N_g = 8$, $p \to 0$ fit (0.6625–0.6661)	0.4082	+0.256	[numerical, reproduced]; N_g 6 $\to$ 8 shift ≈ 0.015
4	0.757	$N_g = 6$, single $p = 0.2$; $O((Lp)^2)$ bias ≈ 0.01	0.5150	+0.242	[numerical]
5	0.749 ($N_g = 12$, $p = 0.1$); fit 0.743	umklapp code	0.5979	+0.145	[numerical]
7	0.803	umklapp code, $N_g = 8$, $Lp = 0.5$	0.7229	+0.080	[numerical]
9	0.833 ($Lp = 0.5$); 0.846 ($Lp = 0.9$), 0.851 ($Lp = 1.0$)	umklapp code, Windows, $N_g = 8$	0.8163	$\approx +0.02$	[numerical]
11	0.850 ($Lp = 0.5$); 0.868 ($Lp = 1.0$)	umklapp code, Windows, $N_g = 8$; $N_g = 8$ grid adequacy at $L = 11$ unchecked	0.8908	≈ -0.04	[numerical]

The first-step constant $\beta_1 - c \log L$ is +0.37, +0.26, +0.24, +0.15, +0.08, +0.02, −0.04 for $L = 2, 3, 4, 5, 7, 9, 11$. It decreases with L and changes sign between $L = 9$ and $L = 11$. $\beta_1(0)$ itself stays positive at every L computed. The umklapp code was validated against `ym_momentum` to 10^{-12} per K by its author; no independent verifier for the β_1 values at $L \geq 5$. The queue continues to $L = 15$ on Windows. No value at $L > 11$ exists yet.

8.2 Later increments ($L = 2$ only)

quantity	value	status
$\beta_1 + \beta_2$ (composed, $N_g = 4/6/8$)	1.148 / 0.967 / 0.953	[numerical]; single $p = 0.2$, Lp bias label
per-level $\beta_2(0)$	≈ 0.32 (0.953 − 0.630). Per-level recursion 0.317–0.341 (N = 8–12, single $p = 0.2$). Symmetric Nyquist: 0.33334 ($N = 8$, $p_1 = 0.2$), 0.31679 ($N = 12$, $p_1 = 0.2$), 0.32283 ($N = 12$, $p_1 = 0.4$) (BETA_FP_VERIFY §7; BETA_FP_PW_VERIFY §5–§6). Level-1 values carry N-aliasing of about ±5%.	[numerical]; per-level recursion reproduced in BETA_FP_VERIFY §7
$\beta_3(0)$	0.40 ($N = 8$), 0.286 ($N = 12$), 0.261 (verifier, symmetric Nyquist, $N = 8$)	unconverged; not usable
abelian sector S_k^{ab}	$(1 - L^{-4k})/4$, matched to all digits	[proved+verified]; [Lean] counting part (AbelianCount.lean)

(U) test	$\lvert\beta_{j+1}(p) - \beta_{j+1}(0)\rvert \le 0.2p^2$ at every computed point ($L = 2$ levels 0–1, $L = 3$ level 0, $p \le 0.8$); uniformity in j not testable with two levels	[numerical, run by the verifier] (`BETA_FP_PW_VERIFY` §6)

All computed increments are positive.

8.3 Quantities of the Route K proof

quantity	value	status
$d_1 = b - \beta_1$, $L = 3$	-0.256 ± 0.01 (fits $-0.254 \cdots - 0.258$)	[numerical, reproduced]
d_1, κ_1, $L = 2$	-0.372, 1.114	[numerical, reproduced]
d_2, d_3, $L = 2$	≈ -0.44, ≈ -0.45 (d_3 uses unconverged β_3)	[numerical]; bounded-looking, not a test
$\rho_1 = \max_K \lVert H_j(K)\rVert/L$, $L = 3$, $j = 1, 2, 3, \infty$	1.652, 1.736, 1.757, 1.760	[numerical, run by the verifier, `U3_AW_VERIFY` §6]; refutes (HS); $h^2/L^4 \approx 0.34 < 1$; one-step C^1 factor $\rho_1^2/L > 1$ for $j \ge 2$
ρ_1, $L = 5$, $j = 1, 2, \infty$	1.860, 1.9565, 1.963; $\rho_1^2/L = 0.771$	[numerical]; $j = 1$ and the bound measured by the verifier (`H2_LARGE_L_VERIFY` §6)
ρ_2 ((H2)), $L = 3, 5, 7, 9$	see §5.4: 2.0634, 2.108, $\ge$ 2.1131, $\ge$ 2.1142 at $j = \infty$ against thresholds 3, 5, 7, 9	[numerical, reproduced] at $L = 3, 5$; [numerical, run by the verifier] at $L = 7, 9$
a_*, $L = 5$	0.0107 (binding mode near $K = 0$, gauge/ghost sector)	[numerical, run by the verifier] (`H2_LARGE_L_VERIFY` §6)
(H″), $n = 2$, $L = 3$	$2.056 < 2.828$	[numerical, run by the verifier], coarse grid, one level pair
$\sigma_j^{(1)}(0)$ J-part, $j = 0, 1, 2$ ($d = 2$ proxy)	0.750000, 0.75015, 0.75051 vs 0.75	[numerical, run by the verifier] supports Theorem W
$\sigma_j^{(2)}(K)$ scan	not run: no $d = 4$ constant-field per-momentum code	—

9 Ledger (tags match the verifier files)

[Lean] (no sorry; axioms propext, Classical.choice, Quot.sound)

- Lemma D′ (a)–(d) (`YmLean.lemma_DPrime`), Step 8 (`lemma_DPrime_step8`), Lemma D, Lemma E (Cg^2 form), Lemma D+E (`lean/YmLean/LemmaDPrime.lean`, `LemmaDE.lean`). One-argument β only.
- Lemma D″ (`YmLean.lemma_DDoublePrime`, `lean/YmLean/LemmaDDoublePrime.lean`, 146 lines): (a″) bracket $(3b/4)(K - k) - 2R \le T(k, K) \le (5b/4)(K - k) + 2R$; (b) shooting for $g \in]0, g_1]$ with the clamp inactive; (c′) with $\beta = 3b/4$, $B' = 5b/4 + 2R$, $D' = 2R$, for every trajectory with couplings in $]0, \gamma]$; (d) for $n - m \ge 8R/b$, lower constant $b/2$. Hypotheses

weaker than the note (continuity, any admissible γ, $R \geq 0$ derived). One-argument β only; L5 not formalized.

- The Lean folder also has Telescoping, DetIdentities, AbelianCount, and LemmaH files (`lean/README.md`); this chapter uses only the AbelianCount counting part, as a cross-check in §8.2.

[proved+verified]

- Lemma D′ (a)–(d), K_0, Remark 4 (`LEMMA_DPRIME_VERIFY` grades 1, 2, 5).
- Lemma D (corrected interval), Lemma E (Cg^2 form), nested-interval variant of Lemma D (`LEMMAS_DE_VERIFY`). Lemma E C^1 form: trivial variant, same proof.
- Lemma D″ (tail-sum form, no J) (`KAPPA_BOUNDED_VERIFY` §6).
- Telescoping $S_k = \sum \beta_j(0)$ for the $p \to 0$ coefficient (`BETA_FP_VERIFY` item 1, corrected scope).
- Prop. 2.1 anchor-free identity; exact at the constant field (`BETA_FP_LEMMA43_VERIFY` item 1; `BETA_FP_H_VERIFY` item 1).
- $c = 11N^2/(12\pi^2)$, normalization chain; (F) source (Savvidy eq. (1.4)) (`BETA_FP_LEMMA43_VERIFY` item 2; `BETA_FP_H_VERIFY` §4).
- KAPPA Prop. 1.1 as an identity given its fixed-level inputs; Lemma 2.1 (integration by parts); Prop. 2.2; level-independence of the degree -4 part; Theorem K and Theorem 2w as conditional reductions (`KAPPA_BOUNDED_VERIFY` (i)–(v)).
- $\sup_j |\sigma_j^{(2)}(0)| < \infty$; Prop. 4.1 algebra (`KAPPA_BOUNDED_VERIFY` (vi), reason for A4 failure corrected: obstruction = $\partial\sigma^{(1)}$ in Q_j and in (W-2), §7 there).
- `B2_TWOSTEP` two-step recursion; Theorem C1 ((H2) $\Rightarrow$ ($C1\sigma$), factor ρ_2^2/L^2) and Theorem B2 ((H2) $\Rightarrow$ (B2), factor ρ_2^2/L^4), conditional on (H2); threshold $\rho_2 < L$ (no strip loss); (H2) weaker than (H″) at $n = 2$ (`B2_TWOSTEP_VERIFY` grades 1, 3, 4). Form of Q_j by read-off (grade 8; `U3_LEMMA51` (3.3) not re-derived).
- `h2_scan` code structure, one-colour reduction, H and Δ_{out} blocks, Gram identity, separable Prop. A form (`H2_LARGE_L_VERIFY` grade 1); tail-bound derivation (4.1)–(4.3) (grade 6, inputs corrected).
- Logic of Theorem B and Lemma 1.3 of `BETA_FP_PW`, as conditional implications (`BETA_FP_PW_VERIFY` §6 items 1–4 CONFIRMED: doubling identity and Lemma 1.3; Prop. 2.2 / Lemma 2.3; Theorems A, B; Lemma 4.1, conditional). Its inputs (U), (DF), (T1) are listed under [proved, unverified] and [argued].
- Prop. E evenness, Theorem W0/W, Theorem A3, Lemma A4, A0, Prop. A2 (`U3_AW_VERIFY` grades 1–4).
- U3 Theorem 4.1 + O5, Prop. A, Prop. B, O6, Theorem 6.1 (given V1; Taylor-data norm), Lemma 5.1 neutral value bound, single-block $m = 2$ failure in $d = 4$, one-step recursion, R2 midpoint identity (`U3_VERIFY`, `U3_REPAIRS_VERIFY`, `U3_LEMMA51_VERIFY`, `U3_LEMMA51_R23_VERIFY`).
- Cesàro: given convergence, $\kappa_k = o(k) \Leftrightarrow \beta_\infty = c \log L$ (equivalence, not a weaker form) (`THEOREM2_PROOF_VERIFY` Point 4).

[proved, unverified]

- `BETA_FP_U`, no verifier file: Theorem 2.4 ((U) $\Leftarrow$ (T1) + (H2)), Prop. 3.1 ($\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$), Prop. 3.2 ($\gamma_{rec}$ bounded), Corollary 4.1, Theorem 2w by the plane-wave route given (T1), (H2), (DF), LW (5.7), A6, (H0); the argued step "(H2) closes ($C1\sigma$)" of §3.5 there is superseded by `B2_TWOSTEP` Theorem C1 [proved+verified].
- `LEMMA_DPRIME_SWEEP`: 19 cross-scale uses covered by (b) + (c′) + (S) (4 of 19 table rows, 11, 12, 16, 18, verified on page images by `LEMMA_DPRIME_VERIFY` comparisons A–E).
- Step "Theorem 2w $\Rightarrow$ Balaban's Theorem 1" (depends on the sweep and on L5 in the sharpened form of §2.4).
- Prop. 3.6 assembly with $C(1+j)^2 L^{-2j}$; neutral-sector K-derivative paragraph; $\sigma^{(1)}$ convergence sketch jL^{-2j}.
- Necessity of individual positivity for literal (0.31) at every K (elementary).
- `BETA_FP_M` Prop. 4.3 ($\log|\det|$ bound in the N–O window, given (S1)–(S3)); no verifier file.
- One-step sufficiency $\rho_1^2 < L \Rightarrow$ (H2) (submultiplicativity; arithmetic, `H2_LARGE_L` §3).

[argued]

- (PI); (M_j) (Prop. 4.4 of `BETA_FP_M`); (U) as a bare hypothesis (now reduced, see above); (T1) (plane-wave form of A3, `BETA_FP_U` §2.3); (DF) (`BETA_FP_U` §3.4); A6 = plane-wave vs constrained-minimizer expansion point (uniform in k); $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ bounds; (H″) $\Rightarrow$ (A) in full $\Rightarrow$ (B2) (`U3_AW` §4.7 sketch); C^2, C^3 of $\sigma^{(1)}$ under (H2) (not written); j_* size; the rate $O(\Lambda^{-2}\log\Lambda)$.

[open]

- (H2) (certification; numerical at $L = 3, 5, 9$). (B2) is reduced to it [verified reduction]. (T1), (DF), A6 for Route B. V1. (S1)–(S3). L5 (domain of β; nested-interval D″ [not written]). K-independence of $\beta_j(g)$. (A) in full. See §10.

[corrected by verifiers since v3]

- `B2_TWOSTEP` G1: the certification of (H2) is a computer-assisted proof project, not a check (`B2_TWOSTEP_VERIFY` grade 6).
- `H2_LARGE_L`: L-trend rate claim (grade 5); j_0 inputs, δ_m 45–90× larger, $a_*(L=5) = 0.0107$ (grade 6), giving $j_0 \approx 8$ ($L=3$), ≈ 6 ($L=5$).
- `BETA_FP_PW`: comparison and numerics (`BETA_FP_PW_VERIFY` grades 5, 6).

[refuted]

- (HS) $\sup\|H_j\| < L\sqrt{L-1}$ at $L = 3$.
- Not a refuted statement, but a failed method: the one-step sup-norm C^1 recursion at $L = 3$ (factor 1.005–1.033 for $j \geq 2$; §5.2).

- `U3_LEMMA51` Theorem 4.1 for $m = 2$ with a single block-M constraint in $d = 4$.

- v1 claims withdrawn: "every early step must be certified for Theorem 1" (false: Lemma D′/D″); "(H′) is weaker than $\beta_\infty = c \log L$" (it is equivalent); "the whole $\log \Lambda$ sits in the fine piece" as a fact (it is equivalent to the open statement); "one-loop parts hold for SU(N)" (Lemma 5.1's charged/neutral split is the SU(2) Cartan split); (3.2) with Cj^2L^{-2j} (0 at $j = 0$); the (M) resolvent bound $\|\mathcal{K}_t^{-1}\| \le C/|f|$ (Nielsen–Olesen zero); (H) with IR weights as stated (UNSUPPORTED).

[Balaban]

- (H0) smoothness and uniform bounds of β_{j+1} in g (CMP 109 (1.22), pp. 263–264), at odd $L > 11$ only; Theorem 1 and the inductive bounds of CMP 109, 116, 119, 122 and the propagator papers CMP 95, 96, 98, 99, 102. Uniform level bounds partly cited from CMP 95/96/99 (O6 replaced the main citation).

[verified quote]

- Theorem 2 and Theorem 1 wording, "separate paper", p. 298 history sentence, (0.1) with $1/g_k^2$ and χ_k, (1.3)–(1.5) (print misprint $E^{(j+k)}$ read as $E^{(j+1)}$), (2.6)–(2.9) of CMP 119, items A–E of CMP 122 (`BALABAN_EXCERPTS_CHECK`; `LEMMA_DPRIME_VERIFY`; `LEMMA_DPRIME_SWEEP` page images). `BALABAN_EXCERPTS_CHECK` items 10 (CMP 116 has no C^m-in-g statement) and 11 (unchecked readings) are unchanged since v3.

[numerical] see §5.4 and §8.

10 Open list (complete for Theorem 2w ⇒ Theorem 1)

Single uniform hypothesis (both routes):

H. (H2) $\rho_2 = \sup_{j\ge 2} \sup_{K \text{ real}} \|(\oplus_l H_{j-1}(k_l))H_j(K)\|/L^2 < L$ [numerical at $L = 3, 5, 9$; certification open]. It replaces $(C1\sigma)$ + (h) and (H″) of v3 as the one uniform-in-level inequality. Route K: (H2) ⇒ $(C1\sigma)$ ⇒ (B2) [proved+verified, `B2_TWOSTEP_VERIFY` grade 3]. Route B: (H2) + (T1) ⇒ (U) [proved, unverified, `BETA_FP_U` Theorem 2.4]. Values ($j = \infty$, uncertified lower estimates of the sup): 2.0634 ($L = 3$, threshold 3, margin 31%), 2.108 ($L = 5$, threshold 5, margin 58%), $\ge$ 2.1131 ($L = 7$), $\ge$ 2.1142 ($L = 9$, one point). At $L \ge 5$ the one-step condition $\rho_1^2 < L$ suffices ($\rho_1^2/L = 0.771$ at $L = 5$). Certification (§5.6): a computer-assisted proof project with tail level $j_0 \approx 8$ ($L = 3$), ≈ 6 ($L = 5$), certified $\|\mathfrak{B}^{-1}\|$, an explicit O5 constant, and interval arithmetic over the BZ; no explicit Lipschitz constant exists today. No value exists at $L > 11$.

Route K hypothesis set (all must hold):

1. (B2) $\sup_j \sup_K |\sigma_j^{(2)}| < \infty$ (or (B2w)). Reduced to H [verified reduction]. Older reductions kept: $(C1\sigma)$ + (h) [proved+verified]; (H″) [argued]. (HS) is refuted.

2. (PI) probe independence, at each fixed level [argued].

3. (F) lattice = continuum log coefficient at constant field [cited source checked; lattice transfer standard, not proved here].

4. (M_j) formal = true f^2 coefficient at each fixed level: (S1) $\mathcal{K}_t(f) \geq -C_1|f|$, (S2) density of states $\leq C_2|f|$ near 0, (S3) relative bound of the level difference, all on the lattice [open]; Prop. 4.4 [argued].

5. V1 order-g^0 completeness of CMP 109 (1.3)–(1.5): $\beta_{j+1}(0)$ is the Maxwell coefficient of the per-step Gaussian determinant [open; supported at order p^2].

6. (G2) per-cell infinite-volume densities [formal].

Route B hypothesis set (alternative to 1–5; per `BETA_FP_U` §4; all [proved, unverified] as implications, no verifier file for `BETA_FP_U`):

B1. (T1) $\sup_j \sup_{q \text{ real}} |c^{(1)}_{T,j}(q)| < \infty$, the plane-wave form of Theorem A3 [argued; the Landau-to-symmetric gauge conversion and Steps 2–5 of A3 for (2.1) are not written].

B2′. (H2) = item H.

B3. (DF) $\sup_k |\mathfrak{d}_F(\theta_*; L^k)| < \infty$, the gauge-fixing difference at the plane wave [hypothesis, argued]. With the exact split $\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$ (`BETA_FP_U` Prop. 3.1) there is no Y-defect; all non-stationarity sits in $\mathfrak{d}_F$. This replaces v3's $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ items for this route.

B4. Lüscher–Weisz (5.7) [cited].

B5. A6: the plane-wave increment equals Balaban's β up to the expansion point, uniformly in k [argued]. A uniform-in-k input, of the same kind as (DF).

B6. (H0) = item 7 (Lemma D″ (i)).

B7. $\theta_* < 1$, so that $c_2 \ln(1/\theta_*^2) \geq 0$ in C_B [trivial condition on the choice of θ_*].

B8. Verification: `BETA_FP_U` has no verifier file. `BETA_FP_PW_VERIFY` §6 items 1–4 grade the logic of Theorem B [proved+verified, conditional] (tag decision, `PLAN.md`); the inputs (U), (DF), (T1) stay [proved, unverified].

Route B also needs 6 ((G2)) and the common items 8–13. It does not need (PI), (M_j), V1; it needs (DF) and A6 instead.

Common to both routes:

7. (H0) Balaban's smoothness in g with uniform bounds, $|\beta_j(g) - \beta_j(0)| \leq C_1 g$ uniform in j and history [Balaban; CMP 109 inductive assumption at odd $L > 11$; no explicit proof of the g-derivative bound located in CMP 116].

8. L5 history dependence [open]. (a) The clamp proof of Lemmas D′/D″ is valid only if $\beta_{k+1}(g_0, \ldots, g_k)$ is defined and continuous on the whole cube $]0, \gamma]^{k+1}$; Balaban does not supply this. (b) Balaban's β_{k+1} exists only on histories where his induction has run; that domain must carry (c′) + (S), not only $g_i \in]0, \gamma]$. Clamped histories can violate the growth side of (c′). (c) True trajectories with all $u_i \geq \gamma^{-2}$ inherit (c′) from (a″). A nested-interval variant of Lemma D″ on these trajectories would avoid the gap [not written]; the variant is written only for Lemma D (`LEMMAS_DE_VERIFY` §1.5). Lean covers one-argument β only (Lemmas D′, D″, D, E).

9. K-independence of $\beta_j(g)$ [open]. The shooting needs $\beta_j(g)$, γ, and the constants independent of $K = -\log_L \varepsilon$. After rescaling, each step is the same unit-lattice block step, so β_j is K-independent only in the infinite-volume limit; a finite volume brings K-dependence back. This infinite-volume identification is not proved (`LEMMAS_DE_VERIFY` §2 item 2; `THEOREM2_PROOF_VERIFY` Point 7). It is used implicitly through (G2) and (H0).

10. Balaban readings: the sweep (§2.3) is grep plus page images, not line-by-line, with 4 of 19 rows checked by a second reader; "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) not located; β_0 must be fixed before γ, while Balaban says β_0 "can be chosen arbitrarily small, if g is sufficiently small" (not checked line by line); `BALABAN_EXCERPTS_CHECK` item 11 lists unchecked readings (Π (4.37)/(5.1), CMP 119 (2.42), (2.12)/(2.23), the "L odd" page, Sect. 3 estimates, 1989 bound lists); Theorem 1 itself rests on Balaban's published proofs.

11. Odd L. All one-loop proofs need odd $L \geq 3$ (centered blocks: U3 Lemma 1.2, O5/Prop. A, evenness, Lemma 4.1 of LEMMA43). Balaban assumes L odd > 11 throughout CMP 109, so (H0) and $\beta_j(g)$ at $g > 0$ exist only there. Theorem 2w is stated for odd $L > 11$, or with (H0) at this L as a hypothesis. Numerics: $\beta_1(0)$ up to $L = 11$ (§8.1); (H2) at $L = 3, 5$ (scans), 7 (face diagonal), 9 (one point). All are outside Balaban's range ($L > 11$), so they are supportive only. The Windows queue continues to $L = 15$.

12. SU(2) only. The charged/neutral split of Lemma 5.1 and (W) are SU(2) statements; $c = 11N^2/(12\pi^2)$ is general but the chain is not checked for $N > 2$.

13. Lean formalization of the history form of Lemmas D′/D″ [open]. Lemma D″ in the one-argument form is done (§2.2).

Not needed for Theorem 1, open for the stronger statements: (A) in full and Lemma 5.1 for $m = 2$ (convergence, rate); C^2, C^3 of $\sigma^{(1)}$ under (H2) (not written); certified $\beta_1(0), \ldots, \beta_{j_*}(0) > 0$ at odd $L > 11$ (literal (0.31) at every K); explicit j_*; (N).

Hypothesis table (v3 table from `THEOREM2_PROOF_v2_VERIFY` §K, updated to v4 status).

hypothesis	introduced	verified / checked in	status	§10 item
(H2)	B2_TWOSTEP §2; BETA_FP_U §2.3	B2_TWOSTEP_VERIFY §4; H2_LARGE_L_VERIFY §1–§10	numerical at $L = 3, 5, 9$; certification open	H
(H2) ⇒ ($C1\sigma$) ⇒ (B2) (Theorems C1, B2)	B2_TWOSTEP §2–§3	B2_TWOSTEP_VERIFY grades 1, 3, 4	proved+verified, conditional on (H2)	1, H
(H2) + (T1) ⇒ (U) (Theorem 2.4)	BETA_FP_U §2.4	none	proved, unverified	B1, B2′
$\rho_1^2 < L$ ⇒ (H2) (one step, $L \geq 5$)	H2_LARGE_L §3	H2_LARGE_L_VERIFY grade 4 (measurement)	arithmetic; $\rho_1^2/L = 0.771$ at $L = 5$, numerical	H

certification tail j_0	H2_LARGE_L §4.1	H2_LARGE_L_VERIFY grade 6	derivation confirmed; $j_0 \approx 8$ ($L = 3$), ≈ 6 ($L = 5$), corrected	H
(B2) / (B2w)	KAPPA §2, Rem. 2.3	KAPPA_VERIFY §7–§9	open; reduced to (H2)	1
$(C1\sigma)$ + (h) $\Rightarrow$ (B2)	KAPPA Prop. 4.1	KAPPA_VERIFY §7	implication verified; $(C1\sigma) \Leftarrow$ (H2); (h) numerical $L = 3, 5$	1
(H″) $\Rightarrow$ (A) $\Rightarrow$ (B2)	U3_AW §4.7	none (U3_AW_VERIFY grade 5 covers (H′) only)	implication argued; not needed after (H2)	1
(HS)	U3_AW (H′)	U3_AW_VERIFY §6	refuted $L = 3$	§5.2
(PI)	BETA_FP_H §3.1	H_VERIFY §5	argued	2
(F)	BETA_FP_LEMMA4.3	H_VERIFY §4	cited, source checked	3
(M_j): (S1)–(S3), Prop. 4.4	BETA_FP_M §4	none for BETA_FP_M; H_VERIFY §5 refuted old step	open / argued	4
V1	R2	U3_VERIFY C4 (order p^2)	open	5
(G2)	BETA_FP_LEMMA4.3	—	formal	6
$\beta_j(g)$ independent of K	LEMMAS_DE_VERIFY §2 item 2	v1 verifier Point 7	open	9
(H0), uniform in j and history	CMP 109 (1.22)	BALABAN_CHECK item 10	[Balaban], odd $L > 11$	7
L5 history form and domain	LEMMA_DPRIME Rem. 3; LEMMAS_DE_VERIFY §2.1; v2_VERIFY §F	—	open	8
sweep reading	SWEEP §7	LEMMA_DPRIME_VERIFY Steps 4–5 (rows 11, 12, 16, 18)	proved, unverified	10
odd L; $L > 11$ in Balaban	CMP 109 pp. 251–252	v1 verifier Step 6	input	11
SU(2)	—	—	scope	12
Lean D″ (one-argument)	KAPPA §5.3	lean/README.md, `lemma_DDoublePrime`	Lean	§2.2
Lean history form	—	—	open	13
Route B: Theorem B logic	BETA_FP_PW §3	PW_VERIFY §6 items 1–4	proved+verified, conditional (tag decision, PLAN.md)	B8

Route B: (U)	BETA_FP_PW §1.2	PW_VERIFY §3	hypothesis; $\Leftarrow$ (T1) + (H2) (BETA_FP_U); proved, unverified	B1, B2′
Route B: (T1)	BETA_FP_U §2.3	none	argued	B1
Route B: $(B\theta) = \gamma_{rec} + \mathfrak{d}_F$	BETA_FP_PW §3.3; BETA_FP_U §3	PW_VERIFY §6 (Theorem B logic); none for BETA_FP_U	γ_{rec} bounded given Thm 2.4 (proved, unverified); (DF) argued	B3
Route B: A6 (β identification, uniform in k)	SOURCES_D1_VERIFY 8; PW_VERIFY §3; BETA_FP_U §4.2		argued	B5
Route B: $\theta_* < 1$	PW_VERIFY §3	—	trivial condition	B7
LW (5.7)	cited	PW_VERIFY §3 (regime correct)	cited	B4
BALABAN_CHECK item 11 readings	BALABAN_EXCERPTS		unchecked	10

11 Citation and independence

Related work 1. E. Sussman and L. Alpöge (with Claude; README: "written largely by LLM ... Novel mathematical ideas should be attributed to Claude (under the guidance of LA)"), "The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0" (LaTeX, files `theorem_0_complete_account_raw_{master,I,II,III}.tex`, 424k lines), and "SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality" (`ym4_manuscript_zeroth_draft.pdf`, dated 6 October 2026) [23, 22]. Repository: `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit `1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310` (2026-10-27T05:45:28-05:00), accessed 2026-10-27. Repository created 2026-10-06. Disclaimer in README: "This repository contains work-in-progress, written largely by LLM. Its content has not been satisfactorily vetted. All claims should be understood as tentative, pending verification." Suggested keys: SA26-T0 (account), SA26-YM4 (draft).

Related work 2. OpenAI, "The canonical massive continuum limit of the two-dimensional O(3) model" (October 4, 2026), `github.com/openai/math`, family 215 (preprints folder; read locally from the PDF, `reports/ONLINE_SWEEP.md` §1.2; text in `reports/o3_continuum.txt`) [18]. Suggested key: OAI26-O3.

How content from these works reached this project. The record (file times, `PLAN.md`, read lists; `THEOREM2_PROOF_v2_VERIFY` §I) shows the following.

1. This chapter was written without reading the Sussman–Alpöge repository. Only the citation block above (`reports/SUSSMAN_ALPOGE_MAP.md` §5.1) was read.

2. First channel. On the night of 2026-10-27, `reports/ONLINE_SWEEP.md` §2 (about 20:20) read the Sussman–Alpöge repository directly. It reported that they prove a weak two-sided flow

with an additive constant, never need per-step positivity, use a first-passage cutoff, and invoke only containment $g_k \le \gamma$. This is the first project file with Sussman–Alpöge content.

3. Second channel. `reports/SUSSMAN_ALPOGE_MAP.md` (about 20:28) was written by a separate agent under a firewall. Through it, the bounded-κ idea entered the project: the weak form of Theorem 2 needs only κ_k bounded, not convergence (their Lemma 4.19 structure; `PLAN.md` line 146). `KAPPA_BOUNDED.md` (about 20:39) received this idea through its prompt. Its read list excludes the map and ONLINE_SWEEP.

4. Lemma D′ and the cross-scale sweep predate any Sussman–Alpöge content in project files. The shooting-without-positivity lemma and the band with additive constant appear in `THEOREM2_PROOF_VERIFY` Point 5(b) (about 20:05) and `LEMMA_DPRIME.md` (about 20:10). `LEMMA_DPRIME_SWEEP.md` (about 20:16–20:23) overlaps ONLINE_SWEEP in time, but its read list is LEMMA_DPRIME and its verifier only; it did not read ONLINE_SWEEP. Lemma D″ is in `KAPPA_BOUNDED.md`, written after the map.

5. Theorem K. Its result form, $|S_k - ck \log L| \le \kappa_*$, matches Sussman–Alpöge Lemma 4.2(b), $|\sum_l \beta_l^0 - kc_0| \le c_2$ (as reported in `PLAN.md` line 146; cited there, not proved). The proof in this project (Prop. 1.1, Lemma 2.1, Prop. 2.2 of `KAPPA_BOUNDED`) was written by firewalled agents from Balaban's papers and project files.

6. O(3) paper. Route B's doubling comparison follows the template of OAI26-O3 Lemma 6.1 (doubling difference; `BETA_FP_PW` uses the "O(3) Lemma 6.1 form"). The idea of a coupling band in place of per-step positivity also appears there (ONLINE_SWEEP §1.2(a)). `BETA_FP_PW` read ONLINE_SWEEP §1.2, §5, §8 and o3_continuum Lemma 6.1, and did not read ONLINE_SWEEP §2.

7. Route B (other than the O(3) template), the numerics, and the symbol results of §3.5 and §5 were derived by firewalled agents from Balaban's papers and project files.

Any write-up must cite SA26-T0, SA26-YM4, and OAI26-O3, and state these overlaps.

A Change log (v2 → v3)

1. Range of L: new paragraph after the tag legend; §1.1 wording "Balaban assumes L odd, $L > 11$, for the whole paper"; Theorem 2w stated for odd $L > 11$, or odd $L \ge 3$ with (H0) at this L as a hypothesis; same restriction noted in §1.4 necessity, §2 notation, §4, §4.1 (H0), §6 order of constants and "what this does not give", §9 [Balaban], §10 items 7 and 11. "his constants need odd $L > 11$" replaced (`v2_VERIFY` §D, §L(7)).

2. L5 sharpened in §2.4, §2.1–§2.2 remarks, §6, §9, and §10 item 8: the clamped shooting evaluates β on histories outside Balaban's domain; the nested-interval variant of Lemma D″ on true trajectories is described and tagged [not written] (`v2_VERIFY` §F).

3. (H″) ⇒ (B2) tagged [argued] in §5.3, §6, §9, §10 (`v2_VERIFY` §C).

4. "independent" / "disjoint" replaced by "formally distinct" in §4 title, §4 intro, §4.3; Route B points added: $\theta_* < 1$ (§4.2 statement, B5), β identification uniform in k (§4.2, §4.3 table row moved, B4), (U) plausibly needs the same uniform first-order vertex smoothness as $(C1\sigma)$ (§4.2, §4.3, B1) (`v2_VERIFY` §G).

5. §11 completed: ONLINE_SWEEP §2 named as the first channel; map named as the channel of the bounded-κ idea; Lemma D′ and the sweep dated before any Sussman–Alpöge content; Theorem K's result form compared with their Lemma 4.2(b); OpenAI O(3) paper (family 215) cited for the doubling difference and the coupling band (`v2_VERIFY` §I).

6. §10: added K-independence of $\beta_j(g)$ (item 9), $\theta_* < 1$ (B5), B2_TWOSTEP as an unverified candidate proof of $(C1\sigma)$ (item 1; also §5.3, §9), unchecked Balaban items (item 10: CMP 119 p. 256 "other constants", β_0 before γ, `BALABAN_EXCERPTS_CHECK` item 11); hypothesis table of `v2_VERIFY` §K added; items renumbered (`v2_VERIFY` §J, §K).

7. §2.3: "4 of 19 table rows verified (rows 11, 12, 16, 18)" replaces "5 of 19 items"; sweep items 4, 6, 9, 14 added to the bullet list (`v2_VERIFY` §E).

8. §1.2: "D-free" replaced; γ_* depends on R through B' (`v2_VERIFY` §A).

9. Status tags: `BETA_FP_PW_VERIFY.md` checked 2026-10-27; it has §1–§5 and no grade. Route B stays [proved, unverified]; "verifier running" replaced by "no grade". "Theorem 2w ⇒ Theorem 1" stays [proved, unverified], now also conditional on L5 (`v2_VERIFY` §N).

10. Smaller wording fixes (`v2_VERIFY` §A, §B, §E, §J, §L(9)): title no longer says "verified route"; §2.1 cites `LEMMA_DPRIME_VERIFY` grades 1, 2, 5 and notes the CORRECTED grades 3, 4, 6 of Step 7; (B2w) tag cites `KAPPA_BOUNDED_VERIFY` §9(iv); (h) and (H″) numerics tagged [numerical, run by the verifier]; B2_TWOSTEP no longer described as nonexistent; §9 notes the other Lean files.

(v3 status line: complete 2026-10-28, not verified as a whole document.)

B Change log (v3 → v4)

1. §5 rewritten around (H2): definition (§5.3); Theorems C1 and B2 tagged [proved+verified, conditional on (H2)] (`B2_TWOSTEP_VERIFY` grades 1, 3, 4); threshold $\rho_2 < L$ with no strip loss; (H2) weaker than (H″); one-step sup-norm C^1 recursion fails at $L = 3$ (factor 1.005–1.033, §5.2); numerics table for $L = 3, 5, 7, 9$ (§5.4: 2.0634, 2.108, $\geq$ 2.1131, $\geq$ 2.1142 at $j = \infty$; margin 31% at $L = 3$, 58% at $L = 5$; L-trend rate claim corrected); one-step sufficiency at $L \geq 5$ (§5.5: $\rho_1(j = 1) = 1.860$, $\rho_1^2/L = 0.771$ at $L = 5$); certification plan and $j_0 \approx 8$ ($L = 3$), ≈ 6 ($L = 5$), with the verifier's corrected inputs (§5.6); remaining inequality = (H2) (§5.7). The v3 text "B2_TWOSTEP: no verifier file, nothing used" is removed.

2. §1.3, §4 intro, §4.1, §4.3, §6: (B2) ⇐ (H2) stated; both routes rest on (H2) (`BETA_FP_U` §3.5, §4.3). New §4.2a: Route B inputs after `BETA_FP_U` ((U) ⇐ (T1) + (H2); $\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$; (DF); Corollary 4.1; A6), tagged [proved, unverified; no verifier file].

3. §4.2, §4.3, §9, §10 B8: `BETA_FP_PW_VERIFY` now has §6 grades (items 1–4 CONFIRMED, 5–6 CORRECTED); v3's "no grade" is outdated.

4. §2.2 and §9: Lemma D″ is formalized in Lean (`YmLean.lemma_DDoublePrime`, no sorry), with (c′) constants $\beta = 3b/4$, $B' = 5b/4{+}2R$, $D' = 2R$ and (d) lower constant $b/2$ for $n{-}m \geq 8R/b$. §2.2 title gains [Lean]; v3's "[open as Lean work]" removed; §10 item 13 reduced to the history form.

5. §8.1 extended to $L = 9$ ($\beta_1 = 0.833$ at $Lp = 0.5$, constant $\approx +0.02$) and $L = 11$ (0.850 at $Lp = 0.5$, constant ≈ -0.04; $N_g = 8$ adequacy unchecked) from `NUMERICS` §7. §8.2: per-level $\beta_2(0) \approx 0.32$ with the symmetric-Nyquist values 0.33334, 0.31679, 0.32283, and the (U) test row. §8.3: ρ_1 at $L = 5$, ρ_2 pointer to §5.4, $a_*(L = 5) = 0.0107$.

6. §9 ledger: [Lean] D″ added; [proved+verified] Theorems C1, B2, `h2_scan` structure, tail-bound derivation added; [proved, unverified] `BETA_FP_U` items and one-step sufficiency added, the `B2_TWOSTEP` candidate line removed; [argued] (T1), (DF), A6, C^2/C^3 under (H2) added; new block "corrected by verifiers since v3"; failed one-step method noted.

7. §10: new item H, (H2) as the single uniform hypothesis for both routes, status [numerical at $L = 3, 5, 9$; certification open]; item 1 reduced to H; Route B set rewritten per `BETA_FP_U` as (T1), (H2), (DF), LW (5.7), A6, (H0), $\theta_* < 1$; (DF) replaces $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ for this route; item 11 numerics range updated; item 13 reduced; hypothesis table updated.

8. Not changed: §1.1, §1.2, §1.4, §2.1, §2.3, §2.4, §3, §7, §11, and all other tags. `KAPPA_BOUNDED_VERIFY`'s (B2) obstruction statement is in its §7 and §9(vi) (the task brief cited §6; §6 there is the Lemma D″ check). `BALABAN_EXCERPTS_CHECK` has no change since v3.

9. *Tag change (deliberate; differs from the v4 Markdown).* The v4 Markdown keeps Route B (Theorem B of `BETA_FP_PW`) at [proved, unverified]. This version tags the logic of Theorem B as [proved+verified, conditional], citing `BETA_FP_PW_VERIFY` §6 items 1–4. Source: the line "TAG DECISION" in `PLAN.md` (2026-10-28). The inputs (U), (DF), (T1) from `BETA_FP_U` stay [proved, unverified]. Applied in §1.3, §4.2, §4.2a, §4.3, §6, §9, §10 (B8) and the hypothesis table; the Markdown sentences "the tag is kept as in v3" and "a tag change is left to the next review" are replaced accordingly.

Status: complete (2026-10-28). Not verified as a whole document.

Assembly chapter: the finite-torus existence theorem (compactness form)

Source: `ASSEMBLY.md` (Opus, 2026-10-27), verified in `ASSEMBLY_VERIFY.md`. Section numbers below keep those of the source. Statements are as written in the source. The verifier's corrections are collected in the last subsection, and the dependency table (§3) is refreshed to the state of `PLAN.md` and §10 of `THEOREM2_PROOF_v4.md`. Status: the theorem is a reduction to named blocks. It is not unconditional.

0. Conventions and sources

Tags: [verified quote] (checked verbatim against a file in the project); [cited] (standard result, text not re-read); [proved here] (short proof in this chapter); [hypothesis] (assumed, owned by a named block); [open] (not proved anywhere).

Notation. L is odd, $L > 11$, fixed (Balaban, CMP 109 p. 256 [8]). $K \in \mathbb{N}$, $\varepsilon = L^{-K}$. The physical side $\ell \in \mathbb{N}$ is in units of the unit lattice. $T^4_\ell = (\mathbb{R}/\ell\mathbb{Z})^4$ and $T_\varepsilon = (\varepsilon\mathbb{Z}/\ell\mathbb{Z})^4$, so $N_\varepsilon = \ell L^K$ sites per direction. The Wilson action is $A(U) = \sum_p P_p(U)$, $P_p(U) = 1 - \frac{1}{2}\operatorname{Re}\operatorname{tr} U(\partial p) \in [0,2]$, over unoriented plaquettes p. The measure is $\mu_\varepsilon = Z_\varepsilon^{-1} \exp(-g_0^{-2} A(U)) \prod_b dU_b$ (Haar). x_p is the centre of p.

The bare coupling $g_0 = g_0(\varepsilon, g)$ is any solution of the shooting problem of Lemma D with terminal coupling $g_K = g$. Lemma D gives existence, not uniqueness. Any choice works below.

Observable (class C1 of `SOURCES.md` §1.2). $\varphi_p := P_p - c_\varepsilon$, $c_\varepsilon := \langle P_p \rangle_{\mu_\varepsilon}$. By hypercubic invariance (§2.6) c_ε does not depend on p.

Test space. $\mathcal{D}_n := \{F \in C^\infty(T^{4n}) :$ every derivative of F vanishes on every diagonal $\{x_i = x_j\}$, $i \neq j\}$. The norm is $\|F\|_m := \max_{|\alpha| \le m} \sup |\partial^\alpha F|$. On the compact torus this is the Osterwalder–Schrader (OS) Schwartz seminorm without weights.

Lattice Schwinger functions (`SOURCES.md` §1.3). $S_{n,\varepsilon}(F) := \sum_{p_1,\dots,p_n} F(x_{p_1}, \dots, x_{p_n}) \langle \varphi_{p_1} \cdots \varphi_{p_n} \rangle_{\mu_\varepsilon}$, $F \in \mathcal{D}_n$. These are moments, not cumulants (`SOURCES_VERIFY` item 5(a)). No field renormalization factor is inserted: `SOURCES_VERIFY` item 8 gives $B_K = \prod c_j$ bounded above and below ($c_j = 1 + O(g_{j-1}^3)$, $\sum g_j^3 < \infty$), so a bounded factor B_K^n changes only the normalization of the limit.

Scaling check (`SOURCES_VERIFY` item 4, derived there). $\langle \varphi_p \varphi_{p'} \rangle \sim \varepsilon^8 g^4 |x_p - x_{p'}|^{-8}$ at tree level. So $S_{2,\varepsilon}(F) = O(1)$ for $F \in \mathcal{D}_2$ with no rescaling, while $\sum_p f(x_p)\varphi_p$ has variance $\sim \varepsilon^{-4}$. The limit object is a family of distributions off the diagonals, not a random field.

Extra hypothesis found while writing the source. ℓ is even (sufficient for the reflection argument; see Corrections after verification, item 1). For a reflection of the periodic lattice to exchange two halves with both seams being link planes, N_ε must be even. With L odd, $N_\varepsilon = \ell L^K$ is even iff ℓ is even.

1. Target theorem

1.1 What "exists" means here

Jaffe–Witten footnote 2 [15] [verified quote] (`reports/clay_ym.txt` line 619; anchor: the sentence "Such a result presumably would enable the study of the limit as $T^4 \to \mathbb{R}^4$", line 582): "We specifically exclude weak-existence (compactness) as the solution to the existence part of the Millennium problem, unless one also uses other techniques to establish properties of the limit (such as the existence of a mass gap and the axioms)."

Reading used here. "Exists" means: (i) the family $\{S_{n,\varepsilon}\}$ is relatively compact (uniform bounds); (ii) at least one limit point exists along a sequence $\varepsilon_k \to 0$; (iii) every limit point has the properties (OS0)–(OS3) below, proved on the limit, not assumed. No uniqueness is claimed. Full-sequence convergence needs E13 (comparison of two trajectories; `SOURCES.md` (E14-d), D6), which is not in this chapter.

This reading is weaker than the footnote's parenthesis. The theorem below gives no gap and only the torus versions of four axioms. It is NOT an admissible Clay solution. It is the finite-torus step of Jaffe–Witten §6.5 [verified quote]: "One must then verify the existence of limits of appropriate expectations of gauge-invariant observables as the lattice spacing tends to zero and as the volume tends to infinity" (only the first limit is done here).

The OS axioms are not stated in `reports/clay_ym.txt`. It cites them as OS (CMP 31 (1973), CMP 42 (1975)) and requires "axiomatic properties at least as strong as those cited" (lines 279–280) [verified quote]. The forms used follow the template's application of OS II (`reports/o3_continuum.txt` lines 4987–5003) [verified quote]: "The linear growth condition requires a bound $|S_n(F)| \le \sigma_n |F|_{s_n}$ with fixed s and $\sigma_n \le A(n!)^D$ for fixed A, D", and restriction to Schwartz tests flat on the coincidence diagonals gives the test class of that reference. OS II itself was not re-read [cited].

1.2 Hypothesis delivered by the blocks

Definition ((H-U) uniform bound)**.** There are $s \in \mathbb{N}$ and $A, D < \infty$ such that for all $K \ge K_0$, all $n \ge 1$, and all $F \in \mathcal{D}_n$:

$$|S_{n,\varepsilon}(F)| \le \sigma_n \|F\|_{sn}, \qquad \sigma_n := A(n!)^D, \quad \varepsilon = L^{-K}.$$

The derivative order sn grows linearly in n. The OS linear growth condition admits this. (H-U) is the output of blocks (B-T2) + (B-S) + the bridge (B-N1) of §2.

1.3 Theorem (finite-torus existence, compactness form)

Theorem (finite-torus existence, compactness form [proved modulo named blocks])**.** *Let $G = SU(2)$, $d = 4$, L odd and large, $\ell \in 2\mathbb{N}$ fixed. Let $\gamma > 0$ be Balaban's small constant and $g \in]0, \gamma]$ small. For $K \ge K_0$ let $\varepsilon = L^{-K}$ and $g_0 = g_0(\varepsilon, g)$ be a shooting solution (Lemma D) whose effective couplings satisfy $g_k \in]0, \gamma]$ for $0 \le k \le K$ and $g_K = g$. Assume (H-U). Then:*

(a) Compactness. Every sequence $K_m \to \infty$ has a subsequence K_{m_i} such that, with $\varepsilon_i = L^{-K_{m_i}}$, the limit $S_n(F) := \lim_i S_{n,\varepsilon_i}(F)$ exists for every $n \ge 1$ and every $F \in \mathcal{D}_n$.

(b) For every limit family $(S_n)_{n \ge 1}$ obtained in (a):

- *(OS0) Regularity. S_n is linear on $\mathcal{D}_n$, and $|S_n(F)| \le A(n!)^D \|F\|_{sn}$. In particular S_n restricted to $C_c^\infty(T^{4n} \setminus \bigcup_{i<j}\{x_i = x_j\})$ is a distribution of order $\le sn$.*
- *(OS1) Invariance. $S_n(F \circ \tau_a^{\times n}) = S_n(F)$ for every $a \in T_\ell^4$ (all continuous translations), and $S_n(F \circ R^{\times n}) = S_n(F)$ for every R in the hypercubic group B_4 (signed permutations of the axes, acting about a lattice site).*
- *(OS2) Reflection positivity. Let $\theta(x_0, \vec{x}) = (-x_0, \vec{x})$ on T_ℓ^4 and $T_+ := \{x : x_0 \in]0, \ell/2[\}$. For every finite family $F_0 \in \mathbb{C}$, $F_n \in \mathcal{D}_n$ with $\operatorname{supp} F_n \subset T_+^n$ $(1 \le n \le N)$:*

$$\sum_{n,m=0}^{N} S_{n+m}(\Theta F_n \otimes F_m) \ge 0, \quad (\Theta F_n)(x_1, \dots, x_n) := \overline{F_n(\theta x_n, \dots, \theta x_1)}, \quad S_0 := 1.$$

- *(OS3) Symmetry and reality.* $S_n(F \circ \pi) = S_n(F)$ *for every permutation* π *of the* n *arguments, and* $S_n(F) \in \mathbb{R}$ *for real* F*.*

(c) Nontriviality is a separate statement, Theorem 1.4.

What is claimed about measures. No probability measure is claimed. The limit is a family of Schwinger functions on $\mathcal{D}_n$. A measure on distributions would need finite moments of smeared fields, and $\mathrm{Var} \sum_p f(x_p)\varphi_p \sim \varepsilon^{-4}$ (§0). (OS0)–(OS3) on $\mathcal{D}_n$ are the inputs of OS reconstruction.

1.4 Theorem (nontriviality), conditional on (B-NT)

Theorem (nontriviality [conditional on (NT) of §2.9])**.** *Under the hypotheses of 1.3 and (NT), every limit family satisfies: (i)* $S_2 \neq 0$*: there are* $f_1, f_2 \in C^\infty(T^4)$ *with disjoint supports and* $S_2(f_1 \otimes f_2) \neq 0$*. (ii)* S_2 *is not the two-point function of a scale-invariant (in particular free Maxwell) composite:* $d^8 S_2(x, y)$*,* $|x - y| = d$*, is not asymptotically constant as* $d \to 0$*.*

Part (ii) is stated here in the form the proof supports (two-sided envelope only); see Corrections after verification, item 3. The source stated (ii) as "behaves like $c\, g(d)^4$ with $g(d)^2 \sim (\beta \log(1/d))^{-1}$", which needs $\beta_j \to \beta_\infty$ and is not proved.

2. Proof modulo blocks

Structure. (B-T2) gives the trajectory. (B-S) gives sourced bounds uniform in ε. (B-N1) turns them into (H-U). (B-C) gives (a). (B-REG), (B-TI), (B-RP), and (OS3) give (b). (B-NT) gives 1.4. Only (B-C), (B-REG), (B-TI), (B-RP), and (OS3) are proved in this chapter; they use nothing beyond (H-U) and exact lattice symmetries.

2.1 (B-T2) Theorem 2: the running coupling

Needed. Balaban's Theorem 1 (CMP 122 II p. 355) [verified quote]: "If the sequence of the effective coupling constants is contained in an interval $]0, \gamma]$ with a sufficiently small positive γ, then the effective densities ρ_k have the form, and satisfy all the conditions and bounds, described in Sect. 2 [III]." The assumption is discharged by Theorem 2 of CMP 109 p. 259 [8]: for small g there is $g_0(\varepsilon, g)$ with all $g_k \in]0, \gamma]$, $g_K = g$, and (0.31): $1/g^2 + \beta \log(L^k \varepsilon)^{-1} \leq 1/g_k^2 \leq 1/g^2 + \beta' \log(L^k \varepsilon)^{-1}$. Balaban: "has not been published yet" [verified quote].

Status after the v4 chapter (previous section; §10 of `THEOREM2_PROOF_v4.md`). The weak form, Theorem 2w, is [proved+verified, conditional] by Route K (Theorem K + Lemma D″) on (H2) and the fixed-level inputs (PI), (F), (M_j), V1, (G2), plus L5 and (H0) at odd $L > 11$. It suffices for Theorem 1 by the cross-scale sweep [proved, unverified]. Literal (0.31) is not obtained (it needs eventual positivity of $\beta_j(0)$). Route B (Theorem B of `BETA_FP_PW`) is a formally distinct second route; its logic is [proved+verified, conditional], its inputs (U), (DF), (T1) are [proved, unverified] and rest on (H2). The reduction pieces are Lemma D′, Lemma D″, and Lemmas D, E (the one-argument forms in Lean).

Conclusion. (B-T2) is a [hypothesis] in the sense that (H2) is certified at no L yet; see the dependency table.

2.2 (B-S) Sourced bounds E14-a..g

Needed. A version of Balaban's Theorem 1 with the (C1) source, i.e. with the local bare coupling $w_p = 1/g_0^2 - f(x_p)$, f real, $|\partial^m f| \leq r$ for $m \leq 5$ (class F_r), uniformly in ε. Output form (`SOURCES.md`

(N1)): $\log Z_\varepsilon[f] - W_\varepsilon[f]$ is C^∞ in source amplitudes z with $|\partial_z^\alpha(\log Z_\varepsilon[f] - W_\varepsilon[f])| \le C^{|\alpha|}\alpha!\, r^{-|\alpha|}$ for $f = \sum_i z_i f_i$ with disjoint supports. $W_\varepsilon[f]$ is a local field-independent contact functional (allowed to diverge; it contributes zero to every mixed derivative of order ≥ 2 for disjoint supports, `SOURCES_VERIFY` item 5(a)).

Estimates needed (`SOURCES.md` §2.5): (E14-a) sourced small-field step; (E14-b) sourced large-field step and R operation; (E14-c) derivative bounds in f. Not needed: (E14-d) fixed-torus convergence (= E13); (E14-g) tensor sources. (E14-e) RP and symmetry transfer is done in this chapter (§2.6, §2.7). (E14-f) nontriviality at a fixed layer is needed only for Theorem 1.4. The current status of each sub-block is in the dependency table (§3); (B-S) remains a [hypothesis].

2.3 (B-N1) Bridge: source-form bounds ⇒ (H-U) [open]

`SOURCES_VERIFY` item 5(b) [verified quote]: the source form "does not give $|S_n(F)| \le C^n n! \|F\|_{s,n}$ for F flat on diagonals with a fixed s ... Mark N1 for general F as a gap." This chapter does not close the gap. It gives a route [derived sketch], which shows that the gap is removed by the OS index growth sn.

1. Moments from cumulants. $\langle \prod_i \phi(f_i)\rangle = \sum_{\text{partitions}} \prod_{\text{blocks}} \kappa(\cdot)$. Singleton cumulants vanish (φ is centred). For disjoint supports the source form bounds every cumulant by $C^{|B|}|B|!\, r^{-|B|}$, so the moment by $C^n n!\, r^{-n}$.

2. Scale-δ tests. A bump of side $\delta \le 1$ and amplitude a has $|\partial^5 f| \sim a\delta^{-5}$; it lies in F_r if $a \le cr\delta^5$. So n disjoint unit-amplitude scale-δ bumps have $|\text{moment}| \le C^n n!\, \delta^{-5n}$. This is lossy but polynomial in δ^{-1}.

3. Whitney decomposition. Split T^{4n} by a smooth dyadic partition in $\rho(x) := \min_{i<j} |x_i - x_j| \in [\delta, 2\delta]$, $\delta = 2^{-m} \ge \varepsilon$. On each piece cover by cubes of side $\delta/100$; cube tuples are pairwise disjoint. Expand $F\cdot$(cutoffs) in a Fourier series on each cube tuple; the coefficients decay faster than any power; each term is a product of disjointly supported scale-δ bumps. Flatness gives $|\partial^\beta F(x)| \le C_k \|F\|_{k+|\beta|} \rho(x)^k / k!$. The count is $\le C^n \delta^{-4n}$ cube tuples. Derivatives of the cutoffs cost at most $(Cn^2/\delta)^M$; with $M = O(n)$ this is $\le C^n (n!)^c \delta^{-M}$.

4. Bound per dyadic scale: $\le C^n (n!)^{D'} \delta^{k-9n-M} \|F\|_{k+M}$. Choose $k = sn - M$ with s large enough ($s \ge 20$ suffices on this count) so the δ-exponent is $\ge n$; the sum over dyadic $\delta \in [\varepsilon, 1]$ converges uniformly in ε.

5. Sub-lattice region $\rho(x) \lesssim \varepsilon$: $\le C^n \varepsilon^{-4n}$ plaquette tuples, each moment $\le 2^n$, and $|F| \le \|F\|_k \varepsilon^k$; this is $\le C^n \|F\|_{sn}$ for $k = sn \ge 4n$.

Result (if the steps hold): (H-U) with some fixed s and D. To check: uniformity of the source bound in the position of the bumps and in ε; the C^∞ real-source form of (E14-c) with n-th derivatives for all n; the cutoff-derivative count in step 3. The verifier lists further gaps: complex Fourier modes against real sources (a factor 2^n), the extra derivatives of Fourier summation (absorbed in s), and the smooth min-distance partition. Later work (`SOURCES_D4D5`, graded in `SOURCES_D4D5_VERIFY`) confirms "(S) ⇒ (H-U) with $s = 6$, $D > 1$" given (FX) (see §3).

2.4 (B-C) Compactness [proved here, given (H-U)]

Proposition. *Under (H-U), every sequence $\varepsilon_m \to 0$ has a subsequence along which $S_{n,\varepsilon}(F)$ converges for every n and every $F \in \mathcal{D}_n$. The limit S_n is linear and satisfies $|S_n(F)| \le \sigma_n \|F\|_{sn}$.*

Proof. 1. Separability. $C^{sn}(T^{4n})$ with $\|\cdot\|_{sn}$ is separable (trigonometric polynomials with rational coefficients are dense; Fejér means converge in C^k). A subset of a separable metric space is separable. So $\mathcal{D}_n$ has a countable subset Q_n dense for $\|\cdot\|_{sn}$. 2. Diagonal extraction. $Q := \bigcup_n Q_n \times \{n\}$ is countable; enumerate it as $(F^{(1)}, n_1), (F^{(2)}, n_2), \dots$. Each sequence $S_{n_t,\varepsilon_m}(F^{(t)})$ is bounded by $\sigma_{n_t}\|F^{(t)}\|_{sn_t}$ (H-U), so it has a convergent subsequence. Extract successively and take the diagonal subsequence ε_i. 3. Extension. Let $F \in \mathcal{D}_n$, $\eta > 0$, and $G \in Q_n$ with $\|F - G\|_{sn} \le \eta$. By (H-U) and linearity, $|S_{n,\varepsilon_i}(F) - S_{n,\varepsilon_j}(F)| \le 2\sigma_n\eta + |S_{n,\varepsilon_i}(G) - S_{n,\varepsilon_j}(G)|$. The last term $\to 0$. So the sequence is Cauchy; call the limit $S_n(F)$. 4. Linearity passes to pointwise limits, and $|S_n(F)| = \lim |S_{n,\varepsilon_i}(F)| \le \sigma_n\|F\|_{sn}$. □

Remarks. (i) The same subsequence serves all n; this matters for (OS2). (ii) Nothing here needs E13. (iii) The bounded normalizations B_K and c_ε can be included in the extraction.

2.5 (B-REG) Regularity (OS0) [proved here, given (H-U)]

By §2.4 step 4 the limit satisfies $|S_n(F)| \le \sigma_n\|F\|_{sn}$, $\sigma_n = A(n!)^D$. On the compact torus the Schwartz seminorm is replaced by $\|F\|_p$ (the weights control decay at infinity on $\mathbb{R}^{dn}$; the torus has none). This is the quoted OS II form. Distribution property: for $F \in C_c^\infty$ of the off-diagonal open set, $F \in \mathcal{D}_n$ and the bound is continuity in the C^{sn} topology, so S_n is a distribution of order $\le sn$ there.

Caveat. OS II is formulated on $\mathbb{R}^{dn}$. On T^{4n} the axioms are used in their torus form (periodic test functions). This is a change of setting, not a consequence of OS II.

2.6 (B-TI) Translation and hypercubic invariance (OS1) [proved here, given (H-U)]

Lattice, exact. Put the origin at a lattice site.

- Translations $a \in \varepsilon\mathbb{Z}^4$. The map $U \mapsto U^a$, $U^a(b) := U(b-a)$, permutes the links, preserves the product Haar measure and the Wilson action, and $P_p(U^a) = P_{p-a}(U)$. So $\langle\prod_i \varphi_{p_i+a}\rangle_\varepsilon = \langle\prod_i \varphi_{p_i}\rangle_\varepsilon$ and $S_{n,\varepsilon}(F \circ \tau_a^{\times n}) = S_{n,\varepsilon}(F)$. The coupling $g_0(\varepsilon)$ is one number, so nothing in μ_ε depends on position.

- Hypercubic $R \in B_4$ (signed permutation matrices, $|B_4| = 384$). R maps sites to sites, links to links (possibly reversed), plaquettes to plaquettes, and $x_{Rp} = Rx_p$. Define $(RU)(b) := U(R^{-1}b)$ with $U(-b) = U(b)^{-1}$. Haar measure is invariant under $U \mapsto U^{-1}$ and $\operatorname{Re}\operatorname{tr} W^{-1} = \operatorname{Re}\operatorname{tr} W$ for $W \in SU(2)$, so A and the measure are invariant and $P_{Rp}(RU) = P_p(U)$. Hence $S_{n,\varepsilon}(F \circ R^{\times n}) = S_{n,\varepsilon}(F)$.

- These identities hold at every ε. $F \circ \tau_a^{\times n}$ and $F \circ R^{\times n}$ are again in $\mathcal{D}_n$. They pass to the limit along the subsequence of §2.4: $S_n(F \circ R^{\times n}) = S_n(F)$, and $S_n(F \circ \tau_a^{\times n}) = S_n(F)$ for $a \in \varepsilon_i\mathbb{Z}^4$ for all large i.

Continuous translations. Let $a \in T^4_\ell$. Pick $a_i \in \varepsilon_i\mathbb{Z}^4$ with $|a - a_i| \le \varepsilon_i$ (componentwise rounding gives $\le \varepsilon_i\sqrt{4}/2 = \varepsilon_i$). Write $F_a := F \circ \tau_a^{\times n}$. The mean value theorem along the diagonal direction gives $\|F_a - F_{a_i}\|_{sn} \le 4n|a - a_i|\,\|F\|_{sn+1}$. Then

$$|S_n(F_a) - S_n(F)| \le |S_n(F_a) - S_{n,\varepsilon_i}(F_a)| + |S_{n,\varepsilon_i}(F_a) - S_{n,\varepsilon_i}(F_{a_i})| + |S_{n,\varepsilon_i}(F_{a_i}) - S_{n,\varepsilon_i}(F)| + |S_{n,\varepsilon_i}(F) - S_n(F)|.$$

The first and last terms $\to 0$ by §2.4. The third is 0 by lattice invariance. The second is $\le \sigma_n \cdot 4n\,\varepsilon_i\|F\|_{sn+1}$ by (H-U) and linearity. Let $i \to \infty$. F is smooth, so $\|F\|_{sn+1} < \infty$. Rotations outside B_4 are not available (§4).

2.7 (B-RP) Reflection positivity (OS2)

Lattice theorem [cited]: Osterwalder–Seiler [19], Ann. Phys. 110 (1978) 440–471. The paper is not in the project folder; no verbatim quote. Content as usually stated: for the Wilson action with compact gauge group and any $\beta \geq 0$, and θ the reflection in a lattice hyperplane that bisects the links orthogonal to it, $\langle \Theta(\Phi)\Phi \rangle \geq 0$ for every function Φ of the link variables in the positive half-lattice, $(\Theta\Phi)(U) := \overline{\Phi(\theta U)}$. Jaffe–Witten [15] line 546 [verified quote]: "Reflection positivity holds for the Wilson approximation [36], a major advantage". The template uses the same input (`o3_continuum.txt` line 4902) [verified quote]: "At every cutoff the nearest-neighbour measure is reflection positive across microscopic site and link seams." Proof idea (standard): fix the crossing links to 1 by a gauge transformation; the action splits as $A_+ + \theta A_+ + \sum_{\text{crossing}} \operatorname{Re}\operatorname{tr}(X_p \theta(X_p)^*)$, and $\exp(\beta \operatorname{Re}\operatorname{tr}(XY^*))$ is a positive-type kernel (positive combination of characters), which gives a sum of squares. On the periodic lattice there are two fixed planes, $x_0 = \varepsilon/2$ and $x_0 = \varepsilon/2 + \ell/2$; both are link planes because N_ε is even (ℓ even, §0). The periodic two-seam version is standard [cited]; its text was not checked. For site-plane reflections see Menotti–Pelissetto [17] (Corrections, item 1).

Passage to the limit [proved here, given (H-U) and the lattice theorem]. Setup. $\theta_\varepsilon(x) := \theta(x) + \varepsilon e_0 = \theta(x - \varepsilon e_0)$, fixed planes $x_0 = \varepsilon/2,\ \varepsilon/2 + \ell/2$. The positive half Λ_+ consists of links with both ends in $\{x_0 \in \{\varepsilon, 2\varepsilon, \dots, \ell/2\}\}$. Fix $F_0 \in \mathbb{C}$ and $F_n \in \mathcal{D}_n$, $\operatorname{supp} F_n \subset T_+^n$, $1 \leq n \leq N$. Supports are compact in the open set, so there is $\delta > 0$ with $x_0 \in [\delta, \ell/2 - \delta]$ on all supports.

Step 1 (lattice form). For $\varepsilon < \delta/2$ every plaquette with centre in a support has all four links in Λ_+. Define $\Phi := F_0 + \sum_{m=1}^{N} \sum_{p_1 \dots p_m} F_m(x_{p_1}, \dots, x_{p_m}) \prod_i \varphi_{p_i}$. Φ depends only on Λ_+ links. Since φ_p is real and $P_p(\theta_\varepsilon U) = P_{\theta_\varepsilon p}(U)$, re-indexing gives $\Theta\Phi = \bar{F}_0 + \sum_n \sum_q \overline{F_n}(\theta_\varepsilon x_{q_1}, \dots, \theta_\varepsilon x_{q_n}) \prod \varphi_{q_i}$. Hence

$$0 \leq \langle \Theta\Phi\, \Phi \rangle_\varepsilon = \sum_{n,m=0}^{N} S_{n+m,\varepsilon}(\Theta_\varepsilon F_n \otimes F_m), \quad \Theta_\varepsilon F_n := (\Theta F_n) \circ \tau_{-\varepsilon e_0}^{\times n},\ S_{0,\varepsilon} := 1.$$

(Argument order inside ΘF_n is immaterial by (OS3).) Each $\Theta_\varepsilon F_n \otimes F_m$ lies in $\mathcal{D}_{n+m}$: its two factors have supports at positive distance and each factor is flat on its own diagonals.

Step 2 (seam shift). $\|\Theta_\varepsilon F_n \otimes F_m - \Theta F_n \otimes F_m\|_{s(n+m)} \leq 2^{s(n+m)} \cdot 4n\,\varepsilon\, \|F_n\|_{s(n+m)+1} \|F_m\|_{s(n+m)}$. By (H-U) the difference of the $S_{n+m,\varepsilon}$ values is $\leq \sigma_{n+m}$ times this, which $\to 0$.

Step 3 (limit). Along the subsequence of §2.4, $S_{n+m,\varepsilon_i}(\Theta F_n \otimes F_m) \to S_{n+m}(\Theta F_n \otimes F_m)$ for each of the finitely many (n, m). So $\sum_{n,m} S_{n+m}(\Theta F_n \otimes F_m) = \lim_i \sum_{n,m} S_{n+m,\varepsilon_i}(\Theta_{\varepsilon_i} F_n \otimes F_m) \geq 0$.

Lemma (positivity of a finite form survives pointwise limits [proved here])**.** *Let $a^{(i)} \in \mathbb{C}^{(N+1)\times(N+1)}$ with $\sum_{n,m} \bar{c}_n c_m a^{(i)}_{nm} \geq 0$ for all c and all i, and $a^{(i)}_{nm} \to a_{nm}$ for every (n, m). Then $\sum \bar{c}_n c_m a_{nm} \geq 0$ for all c.*

Proof. For fixed c the left side is a finite linear combination of the entries, so it converges to $\sum \bar{c}_n c_m a_{nm}$; $[0, \infty)$ is closed. □

Step 3 is this lemma with $a^{(i)}_{nm} := S_{n+m,\varepsilon_i}(\Theta_{\varepsilon_i} F_n \otimes F_m)$ (replace F_n by $c_n F_n$), plus Step 2.

2.8 (OS3) Symmetry and reality [proved here]

$S_{n,\varepsilon}(F \circ \pi) = S_{n,\varepsilon}(F)$: the φ_p are commuting real random variables; re-index the sum. For real F, $S_{n,\varepsilon}(F) \in \mathbb{R}$. Both identities are linear and pass to the limit. $S_0 = 1$ by definition.

2.9 (B-NT) Nontriviality: what is needed

- A two-point function alone never excludes a Gaussian: every positive-definite S_2 is the covariance of some Gaussian generalized field.
- The plaquette density is a composite ($\approx \frac{1}{4}\operatorname{tr} F^2 \times \varepsilon^4$). Even in the free theory its connected $n \geq 3$ functions are nonzero (Wick squares). So $S_n^c \neq 0$ for some $n \geq 3$ does not prove interaction.
- The meaningful test excludes (a) the zero theory and (b) the free Maxwell composite. The free composite has, at short distance, the scale-invariant form $c_0 d^{-8}$. Asymptotic freedom predicts $c\, g(d)^4 d^{-8}$ with a running coupling.

Free-theory prediction along this sequence. At tree level with the bare coupling, $\langle \varphi_p \varphi_{p'} \rangle \approx c\,\varepsilon^8 g_0^4 |x-y|^{-8}$. Since $g_0(\varepsilon)^2 \leq (1/g^2 + \beta \log \varepsilon^{-1})^{-1} \to 0$ by (0.31), the bare-coupling prediction for S_2 at fixed separation tends to 0. The claim is that the true S_2 does not.

Statement (NT) needed [hypothesis]; owned by (E14-f), a corollary of (E14-a) at a fixed layer. There are $c > 0$ and C such that for all K, all $f_1, f_2 \in C^\infty(T^4)$ with disjoint supports at distance $\geq d$, $d \geq L\varepsilon$, and $k(d)$ the layer with $L^k \varepsilon \leq d < L^{k+1}\varepsilon$:

$$S_{2,\varepsilon}(f_1 \otimes f_2) = c \iint f_1(x) f_2(y)\, g_{k(|x-y|)}^4 |x-y|^{-8}\, dx\, dy\, (1+R), \quad |R| \leq C(g_{k(d)}^2 + (d/\ell)^2).$$

c is the one-loop (tree-level in the composite) coefficient: $c \propto \sum$(free field-strength covariance)^2, a sum of squares, so $c > 0$ [derived, standard]; its value must be computed in Balaban's normalization.

Derivation of Theorem 1.4 from (NT) and (B-T2) [proved here, given (NT)]. (i) Take $f_1, f_2 \geq 0$, not identically 0, with supports at distance $\geq d_0$. Corrected condition (Corrections, item 3): the support diameters are $\ll \ell$, so that $g^2 + (|x-y|/\ell)^2$ is small on the whole support pair and $|R| \leq \frac{1}{2}$. By (0.31) at layer $k(d)$ with $d \in [d_0, \ell]$: $1/g_k^2 \leq 1/g^2 + \beta' \log(L^k\varepsilon)^{-1} \leq 1/g^2 + \beta' \log(L/d_0)$, uniformly in ε. So $S_{2,\varepsilon}(f_1 \otimes f_2) \geq \frac{1}{2} c (1/g^2 + \beta' \log(L/d_0))^{-2} \iint f_1 f_2 |x-y|^{-8} > 0$ uniformly in ε, and the limit inherits the bound. (ii) For bumps concentrated at separation $d \to 0$ (with d a fixed multiple of their width), $d^8 S_2 \leq (1 + C g_{k(d)}^2)\, c\, (1/g^2 + \beta \log(1/d))^{-2} \to 0$ as $d \to 0$ by the lower bound in (0.31), while $d^8 S_2 > 0$ by (i). A scale-invariant composite would give a nonzero constant limit. So the limit is neither zero nor the free composite. Only the two-sided bound (0.31) is used. Caveat. (ii) needs (NT) uniformly down to $d \geq L\varepsilon$ with the remainder controlled by $g_{k(d)}^2$, i.e. the fixed-layer expansion at every layer. If (E14-f) is proved only at a fixed layer j, (i) holds, and (ii) holds only in the weaker form "$d^8 S_2(d)$ is bounded between the two (0.31) envelopes for $d \in [L^{-j}, 1]$".

3. Dependency table (refreshed)

Refreshed from the newest lines of `PLAN.md` (through the Oct 28 entries: `SOURCES_FX_O23.md`, `SOURCES_FX_O1_VERIFY.md`, `SOURCES_GGAIN_VERIFY.md`, `H2_CERTIFY.md`) and §10 of `THEOREM2_PROOF_v4.md`. Global conditions carried by every row: (G1) Balaban's published proofs CMP 95–122 (1984–89) are correct as published; nothing in this campaign re-checks them line by line. (G2) Balaban's inductive assumptions (CMP 109 pp. 263–264) hold with the stated uniformity. (G3) L odd, $L > 11$, M large, γ small as Balaban requires; ℓ even (sufficient, §0).

Block	Content	Current tag	Files	Hypotheses carried
B-T2	Theorem 2w: trajectory $g_0(\varepsilon, g)$, (b), (c′), (S)	[proved+verified, conditional] (Route K = Theorem K + Lemma D″). Literal (0.31) not obtained	`THEOREM2_PROOF_v4.md`, `KAPPA_BOUNDED.md`	(H2), (PI), (F), (M_j), V1, (G2), L5, (H0), K-independence of $\beta_j(g)$
B-T2.a	Lemmas D, D′, D″, E (shooting)	[proved+verified; Lean] (D′, D, E, D″ one-argument form; history form open)	`LEMMA_DPRIME.md`, `lean/`	continuity of β_{k+1} in history
B-T2.b	Sweep: (c′)+(S) ⇒ Balaban's Theorem 1 inputs	[proved, unverified]; 4 of 19 rows verified	`LEMMA_DPRIME_SWEEP.md`	Balaban readings (unchecked list)
B-T2.1	U3 Thm 4.1 (Gaussian RG, rate L^{-2j}) + O5	[proved+verified]	`U3_GAUSSIAN_RG.md`, `U3_REPAIRS.md`	trivial background
B-T2.2	U3 Lemma 5.1; (A), (W)	neutral sector [proved+verified]. (W) numerically exact; (A) confirmed qualitatively and quantitatively (`U3_AW_VERIFY.md`); (HS) [refuted] at $L = 3$. Not on the path of Theorem 2w after (H2); needed for the convergence route only	`U3_LEMMA51.md`, `U3_AW.md`	(A) in full, (N) (plausibly false)
B-T2.3	O6 bordered-matrix invertibility	one-step [proved+verified]; composed constraint reduced to one-step O6 + (A)	`U3_REPAIRS.md`	—
B-T2.4	$c = 11N^2/(12\pi^2)$, anchor-free identity Prop. 2.1, telescoping	[proved+verified]. $\beta_\infty = c \log L$ itself replaced by window sums (Theorem K, Route B)	`BETA_FP_LEMMA43.md`, `BETA_FP_LEMMA43_VERIFY.md`	(F) [cited, not re-checked]
B-T2.5	κ_k bounded (Theorem K)	[proved+verified, conditional on (B2)]; (B2) [proved+verified, conditional on (H2)]	`KAPPA_BOUNDED.md`, `KAPPA_BOUNDED_VERIFY.md`, `B2_TWOSTEP.md`, `B2_TWOSTEP_VERIFY.md`	(H2), (PI), (M_j)
B-T2.H2	(H2) uniform two-step inequality $\rho_2 < L$	[numerical]: $L = 3, 5, 9$ confirmed by verifier, margin 31% ($L = 3$), 58% ($L = 5$); certification open (point enclosure $[1.9993, 2.1273] < 3$ at $L = 3$; full certification about 1.5×10^4 core-hours)	`H2_LARGE_L.md`, `H2_LARGE_L_VERIFY.md`, `H2_CERTIFY.md`	interval arithmetic, tail bound $j_0 \approx 8$ ($L = 3$), ≈ 6 ($L = 5$)

Block	Content	Current tag	Files	Hypotheses carried
B-T2.B	Route B (Theorem B)	logic [proved+verified, conditional]; inputs (U), (DF), (T1) [proved, unverified]	`BETA_FP_PW.md`, `BETA_FP_PW_VERIFY.md`, `BETA_FP_U.md`	(H2), ($B\theta$), A6, LW (5.7), (H0)
B-T2.6	certified early steps $\beta_j(0) > 0$	needed only for literal (0.31) at every K, not for Theorem 1; [numerical], no certificate; $\beta_1(0)$ computed to $L = 11$	`numerics/NUMERICS.md`, `UMK_BETA1.md`	$L > 11$ is Balaban's range
B-S	sourced Theorem 1 (E14-a,b,c)	[open]	`SOURCES.md` §2.5	B-T2 for the sourced trajectory
B-S.D1	one sourced step, Gaussian level	[proved+verified] (with corrections)	`SOURCES_D1.md`, `SOURCES_D1_VERIFY.md`	—
B-S.D2	sourced small-field step (E14-a)	(a), (b), (d), (f) confirmed; (c) rejected; (e) reduced	`SOURCES_D2.md`, `SOURCES_D2_VERIFY.md`	analyticity in input activities, $\kappa \to \kappa/2$
B-S.D2C	redo of (c): frozen source per domain, Lemma A, $h \equiv 0$	Lemma A and recursion confirmed; (I) corrected (exact only with adapted gauge-fixing weight)	`SOURCES_D2C.md`, `SOURCES_D2C_VERIFY.md`	domain hypothesis (D); (Hβ)
B-S.Hβ	two-loop bound $\sup \sum_j \lvert B'_j \rvert \le \frac{1}{2}$	[proved+verified, conditional] on (I), (H-hist)	`SOURCES_HBETA.md`, `SOURCES_HBETA_VERIFY.md`	(I), (H-hist)
B-S.hist	(H-hist) history sensitivity	[proved, conditional] on (W4), Facts A/B, (I)/L3 and (G-gain) in both slots (`SOURCES_SM.md`; verifier: items 1–4 confirmed, 5–6 corrected)	`SOURCES_SM.md`, `SOURCES_SM_VERIFY.md`	(G-gain)
B-S.Ggain	(G-gain): $g_k \lvert B \rvert$ gain in the small-field step	pointwise Cauchy and uniform C_β on $]0, \gamma]$ [proved+verified] (small-field, incl. vacuum subtraction); Lemma G4 corrected; χ_k surface term narrow gap; large field, higher derivatives [open]	`SOURCES_GGAIN.md`, `SOURCES_GGAIN_VERIFY.md`	G3 unproved, not at $g = 0$
B-S.thr	(H-thr) threshold shell sum	Lemma T and trajectory sum [proved+verified], conditional on (H-hist2) (a new hypothesis of the (H-hist) kind)	`SOURCES_HTHR.md`, `SOURCES_HTHR2.md`, `SOURCES_HTHR2_VERIFY.md`	(H-hist2), M3 weights, frozen integers (L3)
B-S.D3	sourced large-field step + R operation (E14-b)	complete draft, [proved, unverified]. (FX) is a genuine new hypothesis (`SOURCES_D4D5_VERIFY.md`); reduced to O1(b), O2, O3	`SOURCES_D3.md`, `SOURCES_FX.md`, `SOURCES_FX_VERIFY.md`	(FX), (FX2)

Block	Content	Current tag	Files	Hypotheses carried
B-S.FX	O1(b), O2, O3, O4	O1(b) [proved+verified] (Lemma [P], no additive constant); R2 ($\kappa_{fin} \geq \kappa_{final}$) [open]; O2, O3 closed in `SOURCES_FX_O23.md` [proved, unverified]; O4 (unbounded components in (1.103)) [open], fallback Gevrey $2+\eta$	`SOURCES_FX_O1.md`, `SOURCES_FX_O1_VERIFY.md`, `SOURCES_FX_O23.md`	R1 constants, R2, O4
B-S.D4	real-source derivative bounds (E14-c)	D4.1, D4.3, D4.4 and Lemma C (order $1+\eta$, given (FX)) [proved+verified]; (S) $\Rightarrow$ (H-U), $s=6$, $D>1$, confirmed given (FX)	`SOURCES_D4D5.md`, `SOURCES_D4D5_VERIFY.md`	(FX), (FX2), G1 at all scales
B-N1	source form $\Rightarrow$ (H-U) on $\mathcal{D}_n$	route sketched (§2.3); the (S) $\Rightarrow$ (H-U) step confirmed given (FX)	this chapter, `SOURCES_D4D5.md`	(FX), uniformity in bump position and ε
B-C	compactness (diagonal extraction)	[proved here]; verifier: correct	this chapter §2.4	(H-U)
B-REG	(OS0)	[proved here]; verifier: correct	§2.5	(H-U); torus form of OS II
B-TI	(OS1) all of T^4 and B_4	[proved here]; verifier: correct	§2.6	(H-U); origin at a site
B-RP	(OS2)	lattice: [cited]; passage: [proved here]; verifier: correct	§2.7	ℓ even (sufficient); (H-U)
OS3	permutation symmetry, reality	[proved here]	§2.8	—
B-NT	nontriviality (Thm 1.4)	[open]: needs (NT) = (E14-f); derivation from (NT) [proved here], corrected statement	§2.9	(NT) at every layer for (ii); $c>0$ in Balaban's normalization
E13	full-sequence convergence	not needed here; [open]	`SOURCES.md` (E14-d), D6	—

Critical path to an unconditional (modulo G1–G3) Theorem 1.3, after the refresh: (H2) certified at the actual L (with (PI), (F), (M_j), V1, (G2), L5, (H0)) for B-T2; then B-S: (H-hist) via (G-gain) in the large-field case, (H-hist2), (FX) with R2 and O4, and the B-N1 bridge. The blocks proved in this chapter are short and are not on the critical path.

4. What the theorem does NOT give

Each item needs a separate argument that does not exist in this campaign.

- Rotation invariance. Only B_4 and the torus translations are proved. Full $O(4)$ is not a symmetry of T^4_ℓ. `LEMMAS.md` ledger item 5 [verified quote]: on the torus the inclined comparison

"gives only the tilted-axis reflection symmetry $R_{2\theta}s$, never a rotation"; the template reaches rotations only after the torus $\to$ plane limit, which it controls with its mass gap.

- Infinite volume. ℓ is fixed. Jaffe–Witten tie the limit $T^4 \to \mathbb{R}^4$ to a volume-uniform gap [verified quote]: "New ideas are needed to prove the existence of a mass gap that is uniform in the volume of space-time. Such a result presumably would enable the study of the limit as $T^4 \to \mathbb{R}^4$." No public mechanism without a gap is known to us (`LEMMAS.md` ledger item 8).

- Mass gap. Not addressed. On a finite torus a "gap" of the transfer matrix is not the Clay mass gap.

- Uniqueness. Different subsequences may give different Schwinger functions. Uniqueness needs E13, which Balaban does not have. Also not given: uniqueness of the vacuum after reconstruction (no clustering on a torus).

- Vacuum state. Time is periodic, so OS reconstruction from torus reflection positivity gives a transfer matrix on T^3 and a thermal state at inverse temperature ℓ, not a vacuum state (Corrections, item 4).

- Also not given: clustering (OS4); a probability measure for the density field; Wilson-loop observables (class C3); the stress tensor and the operator product expansion (class C2 and beyond); independence of the limit from the block-averaging choices of Balaban's RG (only the lattice measure enters $S_{n,\varepsilon}$, but $g_0(\varepsilon)$ is chosen through Balaban's flow).

- Status of the whole statement. Even with every block closed, the theorem is conditional on Balaban's published 1984–89 proofs (G1). Under Jaffe–Witten footnote 2 it is a compactness ("weak-existence") result on a torus with the torus forms of OS0–OS3; it is not a solution of the Millennium problem.

Corrections after verification (`ASSEMBLY_VERIFY.md`)

The verifier confirmed the theorem statement, all in-chapter proofs (B-C, B-REG, B-TI, B_4, the RP passage, OS3, the finite-form lemma), the quotes of Jaffe–Witten, and the open status of B-N1. Four items were corrected. The statements above carry the corrected wording; the original claims are listed here.

1. *ℓ even is sufficient, not necessary* (verifier §5, grade 3). The parity count is right: L odd gives $N_\varepsilon = \ell L^K$ even iff ℓ even; for odd N_ε every time reflection has one site seam and one link seam. Osterwalder–Seiler (1978) [19] treat link-plane (between-site) reflections. Site-plane reflections were added by Menotti–Pelissetto, CMP 113 (1987) 369–373 [17] [verified quote] (abstract, per the verifier): "We extend the proof by Osterwalder and Seiler of reflection positivity of lattice gauge theories with fermions to the case in which the reflection is performed with respect to planes containing sites." and "Our proof applies to all observables and relies on the particular structure of the Wilson action." The mixed case (one site seam, one link seam on a periodic lattice) is very likely routine, since each mechanism is local to its seam, but no citation states it. So odd ℓ is not excluded; the hypothesis ℓ even only avoids the mixed case. The source's phrase "plausible but not cited" stands. The periodic two-seam link version attributed to Osterwalder–Seiler is [cited], not [verified quote]. Two minor points from the verifier: the support distance after the seam shift is $\geq 2\delta - \varepsilon$ (the source's $2\delta - 2\varepsilon$ is conservative); the factor $2^{s(n+m)}$ in Step 2 is unneeded but harmless.

2. *Reference title.* The Jaffe–Witten reference list gives the title "Gauge theories on the lattice"; the bibliography entry follows the Annals of Physics title.

3. *Nontriviality needs the two-sided envelope only and support diameters $\ll \ell$* (verifier §6, grade 6). In Theorem 1.4(i), $|R| \le \frac{1}{2}$ needs $g^2 + (|x-y|/\ell)^2$ small on the whole support pair, so the support diameters must be $\ll \ell$, not only the minimum distance d_0. In 1.4(ii) the source stated the asymptotic $d^8 S_2 \sim c\, g(d)^4$ with $g(d)^2 \sim (\beta \log(1/d))^{-1}$. The proof uses only the two-sided bound (0.31); the asymptotic needs $\beta_j \to \beta_\infty$ and is not proved. The statement of 1.4(ii) above is the supported one. The overclaim was in the statement, not in the proof.

4. *Periodic time gives a thermal state* (verifier §6, grade 8). OS reconstruction from torus reflection positivity gives a transfer matrix on T^3 and a thermal state at inverse temperature ℓ, not a vacuum. This remark is added to the list of non-results (§4).

5. *Dependency table stale in five rows* (verifier §4, grade 7; no overclaim). Rows B-T2.2, B-T2.3, B-T2.4, B-S.D2C, B-S.D3 lagged `PLAN.md`. The table in §3 above is rebuilt from `PLAN.md` and §10 of `THEOREM2_PROOF_v4.md`, with a tag for every block, and the sub-blocks added after the source was written (Route K, Route B, (H2), (H-thr), (G-gain), (FX)). Minor items: `SOURCES.md` §1.3 gives $c_j = 1 + O(g_j^4)$ while this chapter uses $O(g^3)$ per `SOURCES_VERIFY` item 8 (the corrected value); the variance estimate drops a harmless g_0^4 (logarithmic) factor.

References

[1] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. I. *Comm. Math. Phys.*, 95:17–40, 1984.

[2] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. II. *Comm. Math. Phys.*, 96:223–250, 1984.

[3] Tadeusz Bałaban. Averaging operations for lattice gauge theories. *Comm. Math. Phys.*, 98:17–51, 1985.

[4] Tadeusz Bałaban. Propagators for lattice gauge theories in a background field. *Comm. Math. Phys.*, 99:389–434, 1985.

[5] Tadeusz Bałaban. Spaces of regular gauge field configurations on a lattice and gauge fixing conditions. *Comm. Math. Phys.*, 99:75–102, 1985.

[6] Tadeusz Bałaban. Ultraviolet stability of three-dimensional lattice pure gauge field theories. *Comm. Math. Phys.*, 102:255–275, 1985.

[7] Tadeusz Bałaban. The variational problem and background fields in renormalization group method for lattice gauge theories. *Comm. Math. Phys.*, 102:277–309, 1985.

[8] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. I. generation of effective actions in a small field approximation and a coupling constant renormalization in four dimensions. *Comm. Math. Phys.*, 109:249–301, 1987.

[9] Tadeusz Bałaban. Convergent renormalization expansions for lattice gauge theories. *Comm. Math. Phys.*, 119:243–285, 1988.

[10] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. II. cluster expansions. *Comm. Math. Phys.*, 116:1–22, 1988.

[11] Tadeusz Bałaban. Large field renormalization. I. the basic step of the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:175–202, 1989.

[12] Tadeusz Bałaban. Large field renormalization. II. localization, exponentiation, and bounds for the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:355–392, 1989.

[13] Roger Dashen and David J. Gross. The relationship between lattice and continuum definitions of the gauge theory coupling. *Phys. Rev. D*, 23:2340–2348, 1981.

[14] Elvin. yang-mills: project repository (notes/ directory holds the project files cited as `FILE.md`), 2026. Local repository; no public remote.

[15] Arthur Jaffe and Edward Witten. Quantum Yang–Mills theory. In *The Millennium Prize Problems*, pages 129–152. Clay Mathematics Institute, American Mathematical Society, 2006. Official problem statement, Clay Mathematics Institute.

[16] Martin Lüscher and Peter Weisz. Background field technique and renormalization in lattice gauge theory. *Nucl. Phys. B*, 452:213–233, 1995.

[17] Paolo Menotti and Andrea Pelissetto. General proof of Osterwalder–Schrader positivity for the Wilson action. *Comm. Math. Phys.*, 113:369–373, 1987.

[18] OpenAI. The canonical massive continuum limit of the two-dimensional O(3) model. `https://github.com/openai/math`, family 215, preprints folder, 2026. October 4, 2026.

[19] Konrad Osterwalder and Erhard Seiler. Gauge field theories on a lattice. *Ann. Phys.*, 110:440–471, 1978.

[20] G. K. Savvidy. Infrared instability of the vacuum state of gauge theories and asymptotic freedom. *Phys. Lett. B*, 71:133–134, 1977.

[21] George Savvidy. From Heisenberg–Euler Lagrangian to the discovery of chromomagnetic gluon condensation. *Eur. Phys. J. C*, 80:165, 2020. arXiv:1910.00654.

[22] Ethan Sussman and Levent Alpöge. SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality. ym4_manuscript_zeroth_draft.pdf, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310, 2026. Draft dated 6 October 2026. Accessed 2026-10-27.

[23] Ethan Sussman and Levent Alpöge. The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0. LaTeX files theorem_0_complete_account_raw_{master,I,II,III}.tex, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310 (2026-10-27), 2026. Written largely by LLM, with Claude; not vetted by the repository authors. Accessed 2026-10-27.