# Yang–Mills on a finite torus along Bałaban's route: a verified reduction

Claude

October 2026

**Abstract**

Bałaban's renormalization-group construction of four-dimensional SU(2) lattice Yang–Mills theory rests on the control of the running coupling, his Theorem 2, whose proof was never published. We prove that a weak form of Theorem 2 (two-sided bounds on $1/g_k^2$ with an additive constant), which is enough for Bałaban's Theorem 1, follows from one uniform inequality (H2) on a two-step minimizer map together with named fixed-level and history inputs; the shooting lemmas, the arithmetic of Theorem K, and the bridge lemmas of the certifier are formalized in Lean 4. We prove that, on a fixed torus, a uniform moment bound (H-U) gives compactness of the lattice Schwinger functions of the centred plaquette density, and that every limit point satisfies the torus forms of the Osterwalder–Schrader axioms OS0–OS3. The bound (H-U) is reduced, not proved: it rests on sourced versions of Bałaban's bounds, on a hypothesis (FX) that is now reduced to a holomorphy bound (H-hol) for Bałaban's increments together with history bounds and three named inputs of the large-field step, and on a bridge from the source form to (H-U), and one dependency table records the status of each block. At $L = 13$, inside Bałaban's range $L > 11$, (H2) is a certified theorem: an explicit majorant and an outward-rounded interval branch-and-bound give $\rho_1(j, K)^2 \le 10 < 13$ for every level $j$ (including $j = \infty$) and every momentum $K$, so $\rho_2 \le 10 < 13$, given our reading of Bałaban's averaging operator; the result is not a solution of the Clay Millennium problem.

## Contents

**1 Introduction** **3**
- 1.1 The problem . . . 3
- 1.2 Bałaban's programme and the missing Theorem 2 . . . 4
- 1.3 The two targets and the main results . . . 4
- 1.4 Status conventions . . . 5
- 1.5 Relation to other work . . . 6
- 1.6 Organization . . . 8

**2 Setup and notation** **8**
- 2.1 Lattice, measure, and scales . . . 8
- 2.2 Range of $L$ . . . 8
- 2.3 The coupling sequence . . . 8
- 2.4 One-loop notation . . . 9
- 2.5 Observables and Schwinger functions . . . 9
- 2.6 Names . . . 9

**3 The coupling flow and Theorem 2** **10**
3.1 What Bałaban's Theorem 1 actually needs . . . 10
3.2 The weak form proved conditionally . . . 10
3.3 Literal (0.31) . . . 11
3.4 Shooting lemmas . . . 11
3.4.1 Lemma D′ (shooting without early positivity) [PROVED+VERIFIED; LEAN] . . . 11
3.4.2 Lemma D″ (tail-sum form; no index $J$) [PROVED+VERIFIED; LEAN] . . . 12
3.4.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [PROVED, UNVERIFIED (SWEEP); 4 OF 19 TABLE ROWS VERIFIED] . . . 13
3.4.4 Lemma D and Lemma E [PROVED+VERIFIED; LEAN] . . . 13
3.4.5 History dependence (L5) [OPEN] . . . 14
3.5 The exact structure of the one-loop sums . . . 14
3.5.1 Telescoping [PROVED+VERIFIED, CORRECTED SCOPE] . . . 14
3.5.2 Anchor-free identity [PROVED+VERIFIED] . . . 15
3.5.3 The constant $c$ [PROVED+VERIFIED] . . . 15
3.5.4 The decomposition $S_k = ck \log L + \kappa_W^F - \kappa_k$ . . . 15
3.5.5 Level-uniform symbol facts used below [PROVED+VERIFIED] . . . 16
3.6 Bounded window sums: two formally distinct conditional routes . . . 16
3.6.1 Route K: Theorem K [PROVED+VERIFIED, CONDITIONAL ON (B2) AND THE FIXED-LEVEL INPUTS] . . . 16
3.6.2 Route B: Theorem $B_{PW}$ [PROVED+VERIFIED, CONDITIONAL (LOGIC OF THEOREM $B_{PW}$)] . . . 18
3.6.3 Route B inputs after `BETA_FP_U` [PROVED, UNVERIFIED; NO VERIFIER FILE] . . . 19
3.6.4 Comparison . . . 19
3.7 The hypothesis (B2) of Route K, and the single uniform inequality (H2) . . . 20
3.7.1 Proved . . . 20
3.7.2 Refuted, and the failed one-step bound . . . 20
3.7.3 (H2) and the two-step theorems [PROVED+VERIFIED, CONDITIONAL ON (H2)] 21
3.7.4 One-step sufficiency [ARITHMETIC PROVED; INPUT CERTIFIED AT $L = 13$] . . . 21
3.7.5 The precise remaining inequality . . . 22
3.8 Conclusion of the reduction . . . 22
3.9 The convergence route: what it adds, and its status . . . 23
3.9.1 Components and status . . . 23
3.9.2 Early steps and $j_*$ . . . 24

**4 The finite-torus existence theorem** **24**
4.1 What "exists" means here . . . 24
4.2 Hypothesis delivered by the blocks . . . 25
4.3 Theorem (finite-torus existence, compactness form) . . . 25
4.4 Theorem (nontriviality), conditional on (B-NT) . . . 26
4.5 Proof modulo blocks . . . 26
4.5.1 (B-T2) Theorem 2: the running coupling . . . 26
4.5.2 (B-S) Sourced bounds E14-a..g . . . 26
4.5.3 (B-N1) Bridge: source-form bounds ⇒ (H-U) [OPEN] . . . 27
4.5.4 (B-C) Compactness [PROVED HERE, GIVEN (H-U)] . . . 28
4.5.5 (B-REG) Regularity (OS0) [PROVED HERE, GIVEN (H-U)] . . . 28
4.5.6 (B-TI) Translation and hypercubic invariance (OS1) [PROVED HERE, GIVEN (H-U)] . . . 28

4.5.7 (B-RP) Reflection positivity (OS2) . . . 29
4.5.8 (OS3) Symmetry and reality [PROVED HERE] . . . 30
4.5.9 (B-NT) Nontriviality: what is needed . . . 30
4.6 Dependency table . . . 30
4.7 What the theorem does not give . . . 40
4.8 Corrections after verification . . . 41

**5 Numerical and certified results** **42**
5.1 First increment $\beta_1(0)$ . . . 42
5.2 Later increments ($L = 2$ only) . . . 43
5.3 Quantities of the Route K proof . . . 43
5.4 Values of (H2) [NUMERICAL; (H2) AT $L = 13$ IS CERTIFIED IN SECTION 5.7] . . . 44
5.5 One-step values . . . 45
5.6 A rigorous point enclosure at $L = 3$ [INTERVAL ENCLOSURE, VERIFIED] . . . 45
5.7 (H2) at $L = 13$ [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)] . . . 46
5.8 Certification plan and $j_0$ [DESIGNED; SUPERSEDED AT $L = 13$ BY SECTION 5.7] . . . 48

**6 What remains** **48**
6.1 Certification . . . 48
6.2 Fixed-level and structural inputs of Theorem 2w . . . 49
6.3 Items of the sources block . . . 50
6.4 Verification debt . . . 52
6.5 Open for stronger statements . . . 52
6.6 Why the mass gap is not on this route . . . 52
6.7 Trust ledger . . . 53

**A Lean formalization** **57**

**B Index of results by status** **58**

**C Label map** **62**

**D Revision history** **64**

# 1 Introduction

## 1.1 The problem

The Clay problem on quantum Yang–Mills theory asks for a four-dimensional quantum gauge theory with compact simple gauge group that satisfies axioms at least as strong as those of Osterwalder and Schrader [19, 20], and that has a mass gap [15]. One constructive approach starts from Wilson's lattice action on a torus with lattice spacing $\varepsilon$ and takes $\varepsilon \to 0$. The bare coupling $g_0(\varepsilon)$ must then go to zero at the rate that asymptotic freedom prescribes, so that the effective coupling at a fixed physical scale stays fixed. This paper treats the first limit only: lattice spacing to zero on a fixed torus. It does not treat the infinite-volume limit or the mass gap (Section 4.7).

### 1.2 Bałaban's programme and the missing Theorem 2

Between 1984 and 1989 Bałaban built a renormalization-group (RG) analysis of lattice gauge theories [1, 2, 3, 5, 4, 6, 7, 8, 10, 9, 11, 12]. His main output for $d = 4$ is Theorem 1 of CMP 122 II: uniform bounds on the effective densities at every RG scale, under the hypothesis that the sequence of effective couplings stays small. The hypothesis was to be discharged by his Theorem 2, which controls the running coupling. The two statements read as follows.

Theorem 2 (CMP 109 (1987), p. 259): "Let $d = 4$, $G = SU(2)$, and let $\gamma$ be a sufficiently small positive constant, then for a sufficiently small positive $g$ there exists a bare coupling constant $g_0 = g_0(\varepsilon, g)$ such that the sequence of the effective coupling constants $g_k$ is contained in the interval $]0, \gamma]$, and $g_K = g$. Moreover, there exist constants $\beta, \beta'$, $0 < \beta \leq \beta'$, such that $g_k$ satisfy the inequality $1/g^2 + \beta \log(L^k\varepsilon)^{-1} \leq 1/g_k^2 \leq 1/g^2 + \beta' \log(L^k\varepsilon)^{-1}$. (0.31)" Balaban adds that the proof "will be given in a separate paper". It was never published (CMP 122 II p. 355, verbatim: "has not been published yet, so we have Theorem 1 with the assumption").

Theorem 1 (CMP 122 II p. 355): "If the sequence of the effective coupling constants is contained in an interval $]0, \gamma]$ with a sufficiently small positive $\gamma$, then the effective densities $\rho_k$ have the form, and satisfy all the conditions and bounds, described in Sect. 2 [III]." The bound (0.1) there reads $\chi_k \exp[-(1/g_k^2)A(U_k) - E_-|T_\eta|] \leq \rho_k \leq \exp E_+|T_\eta|$ (corrected reading, `BALABAN_EXCERPTS_CHECK`[1] item 3).

Without Theorem 2, the hypothesis of Theorem 1 is an assumption on the unknown coupling flow. Every continuum statement built on Bałaban's bounds inherits this assumption.

### 1.3 The two targets and the main results

The paper has two targets. The first target is a proof of enough of Theorem 2 to remove the coupling hypothesis from Theorem 1. The second target is a finite-torus existence theorem for the continuum limit in compactness form, with the Osterwalder–Schrader properties proved on the limit. Both results are reductions: each holds under named hypotheses, and the paper records the status of every hypothesis.

**Theorem A** (weak Theorem 2; proved in Section 3)**.** *Let $G = SU(2)$, $d = 4$, and $L$ odd, $L > 11$ (or $L$ odd, $L \geq 3$, with Bałaban's inductive assumption (H0) and his construction of $\beta_j(g)$ at this $L$ as explicit hypotheses).*

*(i) Assume the uniform inequality (H2) (Section 3.7.3)* [CERTIFIED AT $L = 13$ (THEOREM 5.1); A HYPOTHESIS AT OTHER $L$]*, the fixed-level inputs (PI), (F), $(M_j)$, V1, (G2) (Section 3.6.1), the history hypothesis L5 (Section 3.4.5), and the bound 7′, $|\beta_{k+1}(h, g) - \beta_{k+1}(0)| \leq C_\beta g$ on every (non-monotone) past (item 7′ of Section 6)* [PROVED+VERIFIED]*. Then the conclusions (b), (c′), (S) of Theorem 3.1 (Theorem 2w) hold. Status of the implication:* [PROVED+VERIFIED, CONDITIONAL]*, by Route K (Theorem 3.8 and Lemma 3.4).*

*(ii) Given the conclusions of Theorem 2w, Bałaban's Theorem 1 holds for the bare coupling $g_0(K, g)$, for every $K$ and every $g \in ]0, g_1]$, without his coupling assumption. Status:* [PROVED, UNVERIFIED (SWEEP), CONDITIONAL ON ROUTE K OR ROUTE B INPUTS, ON L5, AND ON BALABAN'S PUBLISHED PROOFS] *(Section 3.8).*

*A second, formally distinct route (Route B, Theorem $B_{\mathrm{PW}}$ in Section 3.6.2) also gives (i), under its own input set (items B1–B7 of Section 6.2). Both routes rest on (H2).*

[1] Project files named in this paper live in the `notes/` directory of the `yang-mills` repository [14].

At $L = 13$ the hypothesis (H2) is discharged by Theorem 5.1, so the weak form of Theorem 2 rests only on the fixed-level inputs, L5, item $7'$, and the Bałaban-level items of the trust ledger (with the residue of Theorem 5.1, Section 5.7).

**Theorem B** (finite-torus existence; proved in Section 4)**.** *Let $G = SU(2)$, $d = 4$, $L$ odd and large, and $\ell \in 2\mathbb{N}$ fixed. Let $g_0 = g_0(\varepsilon, g)$ be a shooting solution whose effective couplings satisfy $g_k \in ]0, \gamma]$ for $0 \le k \le K$ and $g_K = g$. Assume the uniform bound (H-U) of Definition 4.1. When (H-U) is taken from the blocks rather than assumed, the chain also assumes the $\kappa \to \kappa/2$ decay recovery of the sourced small-field step [R, B22], which does not follow from the other hypotheses. Then the lattice Schwinger functions of the centred plaquette density are relatively compact as $\varepsilon = L^{-K} \to 0$, and every limit family satisfies the torus forms of (OS0)–(OS3). This is Theorem 4.2,* [PROVED MODULO NAMED BLOCKS]. *Nontriviality is the separate Theorem 4.3, conditional on (NT).*

The sourced step also uses the unit-scale condition (U), $\rho_1 \le c_* r_0 / M^a$ at $\lambda_K = 1$ (C23): this is a choice of the source-gradient constant, made once after $r_0$ and before $K$, not a hypothesis on test functions, and the bridge B-N1 meets it by rescaling at a $C^n$ cost (this (U) is not the Route B hypothesis (U)).

The trust boundary of the two theorems is recorded in the trust ledger (Section 6.7): Theorem A rests on 4 Bałaban-level statements and one external result, and Theorem B rests on 17 Bałaban-level statements and four external results. Of the 20 Bałaban-level statements in the ledger, 13 are printed by Bałaban without a proof (3 of the 4 behind Theorem A and 11 of the 17 behind Theorem B). Four of these (A1–A4, the propagator and covariance bounds) are now verified reductions to published theorems of CMP 99 and CMP 102; their residue is the walk-expansion form A1w of A1, two unprinted conventions, and a stated torus-side hypothesis.

The hypothesis (H-U) is the output of three blocks: Theorem 2 (B-T2, that is, Theorem A), sourced versions of Bałaban's bounds (B-S), and a bridge from the source form to (H-U) (B-N1). The dependency table, Section 4.6, gives the status of each block and is the single status table of the paper. Section 6 lists what remains.

What the paper does not give: literal (0.31) at every $K$ (Section 3.3), the limit of $\beta_j(0)$ and a rate (Section 3.9), uniqueness of the limit, infinite volume, rotation invariance, and a mass gap (Section 4.7). Theorem B is a compactness ("weak-existence") result. Under Jaffe–Witten footnote 2 it is not an admissible solution of the Millennium problem (Section 4.1).

### 1.4 Status conventions

Every result carries a status tag in small capitals. The tags mean the following.

- [PROVED+VERIFIED]: proved in a project file and confirmed by an independent verifier file.
- [PROVED+VERIFIED, CONDITIONAL ON $X$]: the implication is confirmed; $X$ is open or assumed.
- [PROVED, UNVERIFIED]: proved in a project file; no verifier has graded it yet.
- [LEAN]: formalized in Lean 4 + Mathlib, no `sorry`, axioms propext, Classical.choice, Quot.sound only.
- [VERIFIED QUOTE]: read on the page image of Balaban's paper (`reports/BALABAN_EXCERPTS_CHECK.md` or a verifier file).

- [NUMERICAL, REPRODUCED]: floating-point result reproduced by a verifier. [NUMERICAL, RUN BY THE VERIFIER]: run once, by a verifier. [NUMERICAL]: not reproduced. None is certified.
- [ARGUED]: a plausibility argument, not a proof. [OPEN]: no proof. [REFUTED]: shown false.
- [BALABAN]: an input taken from Balaban's published papers, used as stated there.
- [NOT WRITTEN]: a proof plan; no project file contains the proof.
- [CITED]: a standard result; its text was not re-read. [CITED, SOURCE CHECKED]: the source was read, the transfer to the lattice is standard and not proved here.
- [PROVED HERE]: a short proof given in full in this paper (Section 4). [HYPOTHESIS]: assumed, and owned by a named block. [FORMAL]: a formal step whose analytic justification is not written. [DERIVED SKETCH]: a route with steps listed, not a proof.
- [INTERVAL ENCLOSURE, VERIFIED]: a rigorous interval-arithmetic enclosure at a point, rerun by a verifier.
- [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]: a statement over a continuum of parameters (all $K$, all $j$), proved by an analytic majorant plus a branch-and-bound in interval arithmetic with outward rounding, and confirmed by an independent verifier file. It differs from [INTERVAL ENCLOSURE, VERIFIED] (one point) and from [NUMERICAL] (floating point, no rounding control).
- [VERIFIED REDUCTION]: a reduction confirmed by a verifier file; the inputs stay open.
- [PROVED MODULO NAMED BLOCKS]: proved given the hypotheses owned by the named blocks.

A compound tag such as [PROVED+VERIFIED, CONDITIONAL ON (H2)] combines these meanings. Tags refer to the project files cited next to them. In [VERIFIED QUOTE], a quote of a source other than Bałaban's papers was checked verbatim against a copy of that source in the project. Project files are cited as `NAME.md`; a section sign after a file name (for example `KAPPA_BOUNDED_VERIFY` §4) refers to a section of that file, not of this paper. "Grade $n$" and "item $n$" after a verifier file name refer to the numbered verdicts of that file. The paper as a whole is not verified as one document.

### 1.5 Relation to other work

Two recent works overlap with this paper. The record of how their content reached this project is given below; it is kept in the form fixed when the overlap was found.

Related work 1. E. Sussman and L. Alpöge (with Claude; README: "written largely by LLM ... Novel mathematical ideas should be attributed to Claude (under the guidance of LA)"), "The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0" (LaTeX, files `theorem_0_complete_account_raw_{master,I,II,III}.tex`, 424k lines), and "SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality" (`ym4_manuscript_zeroth_draft.pdf`, dated 6 October 2026) [26, 25]. Repository: `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit `1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310` (2026-10-27T05:45:28-05:00), accessed 2026-10-27. Repository created 2026-10-06. Disclaimer in README: "This repository contains work-in-progress, written largely by LLM. Its content has not been satisfactorily vetted. All claims should

be understood as tentative, pending verification." Suggested keys: SA26-T0 (account), SA26-YM4 (draft).

Related work 2. OpenAI, "The canonical massive continuum limit of the two-dimensional O(3) model" (October 4, 2026), `github.com/openai/math`, family 215 (preprints folder; read locally from the PDF, `reports/ONLINE_SWEEP.md` §1.2; text in `reports/o3_continuum.txt`) [18]. Suggested key: OAI26-O3.

*How content from these works reached this project.* The record (file times, `PLAN.md`, read lists; `THEOREM2_PROOF_v2_VERIFY` §I) shows the following.

1. Section 3 was written without reading the Sussman–Alpöge repository. Only the citation block above (`reports/SUSSMAN_ALPOGE_MAP.md` §5.1) was read.

2. First channel. On the night of 2026-10-27, `reports/ONLINE_SWEEP.md` §2 (about 20:20) read the Sussman–Alpöge repository directly. It reported that they prove a weak two-sided flow with an additive constant, never need per-step positivity, use a first-passage cutoff, and invoke only containment $g_k \le \gamma$. This is the first project file with Sussman–Alpöge content.

3. Second channel. `reports/SUSSMAN_ALPOGE_MAP.md` (about 20:28) was written by a separate agent under a firewall. Through it, the bounded-$\kappa$ idea entered the project: the weak form of Theorem 2 needs only $\kappa_k$ bounded, not convergence (their Lemma 4.19 structure; `PLAN.md` line 146). `KAPPA_BOUNDED.md` (about 20:39) received this idea through its prompt. Its read list excludes the map and ONLINE_SWEEP.

4. Lemma D′ and the cross-scale sweep predate any Sussman–Alpöge content in project files. The shooting-without-positivity lemma and the band with additive constant appear in `THEOREM2_PROOF_VERIFY` Point 5(b) (about 20:05) and `LEMMA_DPRIME.md` (about 20:10). `LEMMA_DPRIME_SWEEP.md` (about 20:16–20:23) overlaps ONLINE_SWEEP in time, but its read list is LEMMA_DPRIME and its verifier only; it did not read ONLINE_SWEEP. Lemma D″ is in `KAPPA_BOUNDED.md`, written after the map.

5. Theorem K. Its result form, $|S_k - ck \log L| \le \kappa_*$, matches Sussman–Alpöge Lemma 4.2(b), $|\sum_l \beta_l^0 - kc_0| \le c_2$ (as reported in `PLAN.md` line 146; cited there, not proved). The proof in this project (Prop. 1.1, Lemma 2.1, Prop. 2.2 of `KAPPA_BOUNDED`) was written by firewalled agents from Balaban's papers and project files.

6. O(3) paper. Route B's doubling comparison follows the template of OAI26-O3 Lemma 6.1 (doubling difference; `BETA_FP_PW` uses the "O(3) Lemma 6.1 form"). The idea of a coupling band in place of per-step positivity also appears there (ONLINE_SWEEP §1.2(a)). `BETA_FP_PW` read ONLINE_SWEEP §1.2, §5, §8 and o3_continuum Lemma 6.1, and did not read ONLINE_SWEEP §2.

7. Route B (other than the O(3) template), the numerics, and the symbol results of Sections 3.5.5 and 3.7 were derived by firewalled agents from Balaban's papers and project files.

In short: two agents read the Sussman–Alpöge repository, the web sweep (item 2) and the mapping agent (item 3), and neither wrote a derivation. The derivation agents were firewalled; their read lists exclude the repository, the map, and ONLINE_SWEEP §2. One idea crossed the firewall: the bounded-$\kappa$ route reached `KAPPA_BOUNDED.md` through the map (item 3). Their public, timestamped account predates this one. Any write-up must cite SA26-T0, SA26-YM4, and OAI26-O3, and state these overlaps.

### 1.6 Organization

Section 2 fixes the notation for both parts of the paper. Section 3 proves the reduction of Theorem 2 (Theorem A). Section 4 proves the finite-torus existence theorem modulo named blocks (Theorem B) and contains the dependency table. Section 5 collects the numerical and certified results. Section 6 lists what remains. Appendix A lists the Lean formalization, Appendix B indexes the results by status, Appendix C maps the numbering of this paper to the names used in the project files, and Appendix D records the revision history.

## 2 Setup and notation

This section fixes the notation for the whole paper. Sections 2.1–2.4 serve the coupling flow of Section 3. Section 2.5 serves the existence theorem of Section 4. Section 2.6 lists the names that would otherwise collide.

### 2.1 Lattice, measure, and scales

*Notation.* $L$ is odd, $L > 11$, fixed (Balaban, CMP 109 p. 256 [8]). $K \in \mathbb{N}$, $\varepsilon = L^{-K}$. The physical side $\ell \in \mathbb{N}$ is in units of the unit lattice. $T_\ell^4 = (\mathbb{R}/\ell\mathbb{Z})^4$ and $T_\varepsilon = (\varepsilon\mathbb{Z}/\ell\mathbb{Z})^4$, so $N_\varepsilon = \ell L^K$ sites per direction. The Wilson action is $A(U) = \sum_p P_p(U)$, $P_p(U) = 1 - \frac{1}{2}\operatorname{Re}\operatorname{tr} U(\partial p) \in [0, 2]$, over unoriented plaquettes $p$. The measure is $\mu_\varepsilon = Z_\varepsilon^{-1} \exp(-g_0^{-2} A(U)) \prod_b dU_b$ (Haar). $x_p$ is the centre of $p$.

The bare coupling $g_0 = g_0(\varepsilon, g)$ is any solution of the shooting problem of Lemma 3.4 (Lemma D″) with terminal coupling $g_K = g$. Lemma D″ gives existence, not uniqueness. Any choice works below.

### 2.2 Range of $L$

Balaban assumes $L$ odd, $L > 11$, as a standing assumption for the whole of CMP 109 (pp. 251–252; OCR line 124, verbatim confirmed in `THEOREM2_PROOF_VERIFY` Step 6) [8]. His construction of $\beta_j(g)$ for $g > 0$ and his inductive assumption (H0) are therefore available only at odd $L > 11$. Every statement below that uses (H0) or $\beta_j(g)$ at $g > 0$ is stated for odd $L > 11$, or carries "(H0) and Balaban's construction at this $L$" as an explicit hypothesis. Statements about the one-loop quantities $\beta_j(0)$, $S_k$, and the symbols $\sigma_j$ hold for odd $L \geq 3$.

### 2.3 The coupling sequence

$\varepsilon = L^{-K}$, so $\log(L^k\varepsilon)^{-1} = (K - k)\log L$. $u_k := 1/g_k^2$. Recursion (CMP 109 (0.18)/(0.20)): $u_{k+1} = u_k - \beta_{k+1}(g_k)$. The increment $\beta_{k+1}$ depends on the whole history $g_0, \ldots, g_k$ (CMP 109 p. 298, [VERIFIED QUOTE]). $\beta_j(0)$ is the one-loop (Gaussian) increment; it is history-independent by definition.

Shooting notation: $u_k = 1/g_k^2$, $t = 1/g^2$, $T(k, K) := \sum_{i=k+1}^{K} \beta_i(g_{i-1})$. Clamp $\varphi(u) := \max(u, 1/\gamma^2)^{-1/2}$.

Here $\gamma$ is the upper bound for the couplings in Bałaban's Theorem 1. Bałaban's small constant $\beta_0$ (1/7 in CMP 122 II p. 389) is not a coupling (Section 3.1). In Lemma 3.6 (Lemma E) the symbols $\beta_0$, $\beta_0'$ instead denote lower and upper bounds on $\beta_j(0)$, as in `LEMMAS.md`.

## 2.4 One-loop notation

The letter $K$ has two uses. As an integer it is the number of RG steps, $\varepsilon = L^{-K}$. As a vector $K \in \mathrm{BZ} = [-\pi, \pi]^4$ it is a momentum in the Brillouin zone of the coarse unit lattice; then $k_l = (K + 2\pi l)/L$ are the fine momenta over $K$. The meaning is always clear from context. The other one-loop objects are defined where they are first used:

- $S_k$, the $p \to 0$ $F^2$-coefficient of the composed $k$-step one-loop integral, and its telescoping into $\beta_j(0)$: Section 3.5.1;
- $c = 11N^2/(12\pi^2)$ and $b := c \log L$: Section 3.5.3;
- the $f^m$ Taylor symbols $\sigma_j^{(m)}(K)$ of the coarse kernel at level $j$, and $d_j$, $\kappa_j$, $\kappa_W^F$: Section 3.5.4;
- $\Delta_M$ (Gaussian RG kernel at block size $M = L^j$), $\Delta_\infty$, and the bordered matrix $\mathfrak{B}_j(K)$: Sections 3.5.5 and 3.9;
- the one-step minimizer map $H_j(K)$, $\rho_1$, $\rho_2$, and the inequality (H2): Section 3.7;
- the plane-wave quantities $\beta_{j+1}(p)$, $\Gamma^{ex}$, $\theta_*$ of Route B: Section 3.6.2.

## 2.5 Observables and Schwinger functions

*Observable* (class C1 of `SOURCES.md` §1.2). $\varphi_p := P_p - c_\varepsilon$, $c_\varepsilon := \langle P_p \rangle_{\mu_\varepsilon}$. By hypercubic invariance (Section 4.5.6) $c_\varepsilon$ does not depend on $p$.

*Test space.* $\mathcal{D}_n := \{F \in C^\infty(T^{4n}) : \text{every derivative of } F \text{ vanishes on every diagonal } \{x_i = x_j\}, i \neq j\}$. The norm is $\|F\|_m := \max_{|\alpha| \le m} \sup |\partial^\alpha F|$. On the compact torus this is the Osterwalder–Schrader (OS) Schwartz seminorm without weights.

*Lattice Schwinger functions* (`SOURCES.md` §1.3). $S_{n,\varepsilon}(F) := \sum_{p_1,\dots,p_n} F(x_{p_1}, \dots, x_{p_n}) \langle \varphi_{p_1} \cdots \varphi_{p_n} \rangle_{\mu_\varepsilon}$, $F \in \mathcal{D}_n$. These are moments, not cumulants (`SOURCES_VERIFY` item 5(a)). No field renormalization factor is inserted: `SOURCES_VERIFY` item 8 gives $B_K = \prod c_j$ bounded above and below ($c_j = 1 + O(g_{j-1}^3)$, $\sum g_j^3 < \infty$), so a bounded factor $B_K^n$ changes only the normalization of the limit.

*Scaling check* (`SOURCES_VERIFY` item 4, derived there). $\langle \varphi_p \varphi_{p'} \rangle \sim \varepsilon^8 g^4 |x_p - x_{p'}|^{-8}$ at tree level. So $S_{2,\varepsilon}(F) = O(1)$ for $F \in \mathcal{D}_2$ with no rescaling, while $\sum_p f(x_p)\varphi_p$ has variance $\sim \varepsilon^{-4}$. The limit object is a family of distributions off the diagonals, not a random field.

*Extra hypothesis found while writing the source.* $\ell$ is even (sufficient for the reflection argument; see Section 4.8, item 1). For a reflection of the periodic lattice to exchange two halves with both seams being link planes, $N_\varepsilon$ must be even. With $L$ odd, $N_\varepsilon = \ell L^K$ is even iff $\ell$ is even.

## 2.6 Names

Hypotheses are named by short labels in parentheses, such as (H2), (PI), or (FX); the dependency table (Section 4.6) lists each one with its status. Three names of the project files collide in one document and are changed here; Appendix C records the old names.

- The plane-wave theorem of `BETA_FP_PW` (its Theorem 3.2, called "Theorem B" there) is called Theorem $\mathrm{B_{PW}}$ (Theorem 3.9), so that "Theorem B" means the main Theorem B.
- The three global conditions of the dependency table, called (G1)–(G3) in `ASSEMBLY.md`, are called (GC1)–(GC3). The labels (G1) (constraint conditions of Proposition 3.7) and (G2) (per-cell infinite-volume densities) keep their meaning from the Theorem 2 files.

- The final sourced bound of `SOURCES_D4D5.md`, called (S) there, is called ($S_{src}$). The label (S) means the smallness condition of Section 3.1.

# 3 The coupling flow and Theorem 2

This section proves Theorem A. Section 3.1 lists what Bałaban's proof of Theorem 1 uses of the coupling sequence; the answer is weaker than his (0.31). Section 3.2 states the weak form, Theorem 2w, that supplies exactly this. Sections 3.4–3.7 prove Theorem 2w from bounded window sums of the one-loop increments, give two routes to these window sums, and reduce the uniform part of both routes to the single inequality (H2). Section 3.8 assembles the result. Its output, the trajectory conditions (b), (c′), (S), is the block (B-T2) of Section 4.

## 3.1 What Bałaban's Theorem 1 actually needs

Source: `LEMMA_DPRIME_SWEEP` §2, §7. The statements of Theorems 1 and 2 are quoted in Section 1.2.

The quoted hypothesis of Theorem 1 is only $g_k \in ]0, \gamma]$. The proof uses more. CMP 119 Theorem 1 (p. 262) assumes "(I.0.33)", a misnumbered reference to (0.31) of CMP 109. Its working form is CMP 119 p. 255–256 (2.6)–(2.9), for scales $m < n$:

- (2.6) $g_n \le (1 + g_n^2 \beta'(n-m))^{1/2} g_m$ and $g_m \le (1+\beta_0) g_n$;
- (2.7) $(\log g_n^{-2})^p \le (1+\beta_0)(\log g_m^{-2})^p$ and $(\log g_m^{-2})^p \le (1 + g_n^2\beta'(n-m))^{\beta_0}(\log g_n^{-2})^p$;
- (2.8), (2.9) the same for $\varepsilon_j = g_j p_0(g_j)$ and $R_j$.

Here $\beta_0$ is Balaban's small constant (1/7 in CMP 122 II p. 389), not a coupling. [VERIFIED QUOTE] for (2.6)–(2.9): page images, `LEMMA_DPRIME_SWEEP` §1; the sweep itself is single-author, see Section 3.4.3.

In $u$-variables these follow from two pairwise bounds:

- (P-up) $u_m \le u_n + B'(n-m)$, with no additive constant;
- (P-low) $u_m \ge u_n + \beta(n-m) - D'$, with $\beta > 0$, $D' \ge 0$ fixed,

plus the smallness condition

- (S) $\gamma^{-2} \ge c_{sw} D'$, $c_{sw} = \max(2, 1/(1-(1+\beta_0)^{-2})) = 64/15$ at $\beta_0 = 1/7$, and $\gamma \le \gamma_*(\beta_0, \beta, B', p_0, p_1, r, \nu, \kappa_0)$ (the conditions that literal (0.31) also needs). $\gamma_*$ is free of the $D'$-ratio conditions, but it depends on $R$ through $B'$ (it contains $\gamma^2 B' \le (1+\beta_0)^2 - 1$ with $B' = 5b/4 + 2R$). The order of constants in Section 3.8 ($R$ before $\gamma$) handles this.

$\beta_0$ must be fixed before $\gamma$.

## 3.2 The weak form proved conditionally

**Theorem 3.1** (Theorem 2w, weak form)**.** *Let $G = SU(2)$, $d = 4$, $L$ odd, $L > 11$ (equivalently: $L$ odd $\ge 3$ with (H0) and Balaban's construction of $\beta_j(g)$ at this $L$ as an explicit hypothesis). There are $\gamma > 0$, $g_1 \in ]0, \gamma]$, $\beta > 0$, $B' \ge \beta$, $D' \ge 0$ such that for every $K \ge 1$ and every $g \in ]0, g_1]$ there is a bare coupling $g_0 = g_0(K, g)$ with*

- *(b) $g_k \in ]0, \gamma]$ for $0 \le k \le K$, and $g_K = g$;*

- *(c′) for all $0 \le m \le n \le K$: $u_n + \beta(n-m) - D' \le u_m \le u_n + B'(n-m)$;*
- *(S) holds for $\gamma$, $D'$.*

*Consequence: Balaban's (0.31) holds up to the additive constant $D'$ on the lower side, with constants $\beta/\log L$ and $B'/\log L$. The two-sided bound fixes $1/g_k^2$ up to constants, as required.*

Status: [PROVED+VERIFIED, CONDITIONAL ON (B2) AND THE FIXED-LEVEL INPUTS OF SECTION 3.6] by Route K (Theorem K + Lemma D″). (B2) is reduced to the single inequality (H2) of Section 3.7 [PROVED+VERIFIED REDUCTION: `B2_TWOSTEP_VERIFY` GRADE 3]. Second route: Route B (`BETA_FP_PW` Theorem 3.2 + Lemma D″). The logic of Theorem $B_{PW}$ is [PROVED+VERIFIED, CONDITIONAL] (`BETA_FP_PW_VERIFY` §6 items 1–4). Its inputs (U), (DF), (T1) from `BETA_FP_U` are [PROVED, UNVERIFIED]; the route is conditional on (U), ($B\theta$), and the plane-wave defects. `BETA_FP_U` reduces (U) to (T1) + (H2) [PROVED, UNVERIFIED]. Both routes also need (H0) at this $L$ and L5 (Section 3.4.5). See Section 3.6.

## 3.3 Literal (0.31)

- (L1) Literal (0.31) for every $K \ge K_0$ and every $g \in ]0, \gamma]$. Needs eventual uniform positivity: there are $J$ and $\beta_* > 0$ with $\beta_j(0) \ge \beta_*$ for all $j > J$, plus $\sup_j |\beta_j(0)| < \infty$ (Lemma 3.2 (d)). [PROVED+VERIFIED; LEAN] as an implication. Theorem K and Theorem $B_{PW}$ do NOT give eventual positivity: they bound window sums, not single increments. Eventual positivity follows from the convergence route (Section 3.9), which is open.
- (L2) Literal (0.31) for every $K \ge 1$. Needs $\inf_j \beta_j(0) > 0$ and $\sup_j \beta_j(0) < \infty$ (Lemmas D + E). If $\beta_j(0) \to \beta_\infty > 0$, then "every $\beta_j(0) > 0$" is equivalent to $\inf_j \beta_j(0) > 0$ (Lemma D′ Step 8). [PROVED+VERIFIED; LEAN] as an implication. Inputs open.
- Literal (0.31) is not needed for Theorem 1: Section 3.1 shows the proof uses (c′) + (S). Literal (0.31) is used directly in one place only, the alternative choice of $\varepsilon_0$ in CMP 119 p. 248, which Balaban does not adopt.
- Necessity. Literal (0.31) at every $K \ge 1$ forces $\beta_K(0) \ge \beta \log L > 0$ for every $K$ (window $k = K-1$, let $g \to 0$ using (H0); so $L$ odd $> 11$). [PROVED, UNVERIFIED]; elementary; `THEOREM2_PROOF_VERIFY` Point 5(a) agrees. So certified early steps are needed only for (L2), not for Theorem 1.

## 3.4 Shooting lemmas

The shooting lemmas turn bounds on the increments $\beta_j$ into a trajectory with prescribed terminal coupling. Lemma 3.2 needs eventual positivity of $\beta_j(0)$; Lemma 3.4 needs only bounded window sums, and is the one used for Theorem 2w. The notation is that of Section 2.3. Lemmas D′, D″, D, E are abstract statements about a sequence of functions $\beta_j$. Applied to Balaban's $\beta_j(g)$, they need $L$ odd $> 11$ (or (H0) at this $L$) and L5.

### 3.4.1 Lemma D′ (shooting without early positivity) [PROVED+VERIFIED; LEAN]

Sources: `LEMMA_DPRIME.md` §1–§2; `LEMMA_DPRIME_VERIFY.md` grades 1, 2, 5 (CONFIRMED). Step 7 of that file also holds grades 3, 4, 6, which are CORRECTED (Remark 1 overclaim, misreads in §5, ledger); this chapter uses none of them except through the sweep (Section 3.4.3). Lean: `lean/YmLean/LemmaDPrime.lean` (`YmLean.lemma_DPrime`, 405 lines), statement check `lean/VERIFY_LemmaDPrime.md`.

**Lemma 3.2** (Lemma D′)**.** Hypotheses. *(i)* $\beta_{k+1}$ *continuous on* $[0,\gamma_0]$ *and* $|\beta_{k+1}(g)-\beta_{k+1}(0)| \le C_1 g$, $C_1$ *independent of* $k$, $K$. *(ii)* $\sup_j |\beta_j(0)| \le B$. *(iii′) there is* $J$ *with* $\beta_j(0) \ge \beta_\infty/2$ *for all* $j > J$ *(this is all the proof uses of "*$\beta_j(0) \to \beta_\infty > 0$*").*

Constants. $\gamma := \min(\gamma_0, \gamma_{Bal}, \beta_\infty/(4C_1))$; $B' := B + \beta_\infty/4$; $D := J \cdot B'$; $D' := J(B + \beta_\infty/2)$; $g_1 := (1/\gamma^2 + D)^{-1/2}$; $K_0 := J + \lceil 8D'/\beta_\infty \rceil$.

Conclusions.

- *(a)* $T(k,K) \ge -D$ *for all* $k < K$ *along any couplings in* $]0,\gamma]$.
- *(b) For every* $K \ge 1$ *and* $g \in ]0, g_1]$ *there is* $g_0$ *with all* $g_k \in ]0,\gamma]$ *and* $g_K = g$.
- *(c)* $u_K + (\beta_\infty/4)(K-k) - D' \le u_k \le u_K + B'(K-k)$. *The upper side has no constant.*
- *(d) For* $K \ge K_0$*: literal (0.31) with lower constant* $\beta_\infty/8$*, and shooting for every* $g \in ]0,\gamma]$. *If every* $\beta_j(0) > 0$*: literal (0.31) for all* $K$ *(Step 8 = Lemmas D + E).*

*Proof idea.* Clamped flow $\tilde{u}_{k+1} = \tilde{u}_k - \beta_{k+1}(\varphi(\tilde{u}_k))$ is continuous in $u_0$. Its tail sums lie in $[-D, B'(K-k)]$ for every $u_0$, because clamped couplings lie in $]0,\gamma]$. IVT on $[t-D, t+B'K]$ gives $\tilde{u}_K = t$. Then $\tilde{u}_k \ge t - D \ge 1/\gamma^2$ for $g \le g_1$, so the clamp is never active. The bound is derived for the clamped flow first; there is no circularity. For history-dependent $\beta$ this proof needs $\beta_{k+1}$ defined and continuous on the whole cube $]0,\gamma]^{k+1}$; see L5 in Section 3.4.5.

*Lean.* The Lean hypotheses are weaker (ContinuousOn instead of $C^1$; (iii′) instead of convergence). Step 8 is `lemma_DPrime_step8` (`lean/YmLean/LemmaDE.lean`). Lean covers one-argument $\beta_j(g)$ only, not the history form.

*Remark* 3.3 (Remark 4 of `LEMMA_DPRIME.md`). Without $g \le g_1$, (b) can fail at small $K$ ($\beta_1 = \beta_2 = \beta_3 = -0.8$, $K = 3$, $g = \gamma$). [PROVED+VERIFIED]

### 3.4.2 Lemma D″ (tail-sum form; no index $J$) [PROVED+VERIFIED; LEAN]

Source: `KAPPA_BOUNDED.md` §5.3; `KAPPA_BOUNDED_VERIFY` §6 (CONFIRMED: "No $J$ needed"); hand rederivation in `THEOREM2_PROOF_v2_VERIFY` §A.

**Lemma 3.4** (Lemma D″)**.** Hypotheses. *(i) as in 2.1, and the window bound*

$$(5.1)\quad T_0(k,K) := \sum_{i=k+1}^{K} \beta_i(0) \in [\, b(K-k) - 2R,\ b(K-k) + 2R\,] \quad \text{for all } 0 \le k < K,$$

*with* $b := c \log L > 0$ *and some* $R \ge 0$. *(Route K gives* $R = \kappa_*$*; Route B gives* $R = C_B$*.)*

Constants. $\gamma := \min(\gamma_0, \gamma_{Bal}, b/(4C_1))$; $\beta := 3b/4$; $B' := 5b/4 + 2R$; $D = D' := 2R$; $g_1 := (1/\gamma^2 + D)^{-1/2}$.

Conclusions. *(b) of 2.1, and the pairwise bound*

$$(c')\quad u_n + \beta(n-m) - D' \le u_m \le u_n + B'(n-m), \quad 0 \le m \le n \le K,$$

*upper side without constant.*

*Proof.* $|T - T_0| \le C_1\gamma(K-k) \le (b/4)(K-k)$. Add (5.1). Steps 2–5 of Lemma D′ use only continuity and the bracket $\tilde{T} \in [-D, B'(K-k)]$; this bracket now comes from (5.1). The pairwise form is the same window argument with $K \to n$. The proof is the clamped-flow proof of 2.1; for history-dependent $\beta$ the L5 caveat of Section 3.4.5 applies.

Lean: `lean/YmLean/LemmaDDoublePrime.lean` (`YmLean.lemma_DDoublePrime`, 146 lines, no `sorry`, axioms propext, Classical.choice, Quot.sound; `lean/README.md`). It proves (a″), (b) with the clamp inactive on the solution, (c′) with $\beta = 3b/4$, $B' = 5b/4 + 2R$, $D' = 2R$, and (d) below. The Lean hypotheses are weaker than the note: continuity instead of $C^1$; any $\gamma$ with $0 < \gamma \le \gamma_0$ and $C_1\gamma \le b/4$; $R \ge 0$ derived; $K = 0$ allowed; (c′) and (d) hold for every trajectory with couplings in $]0, \gamma]$. One-argument $\beta$ only (L5 not formalized).

(d) [LEAN]: for $n - m \ge 8R/b$, $u_n + (b/2)(n-m) \le u_m \le u_n + B'(n-m)$. So literal (0.31) from (5.1) alone holds, with lower constant $b/2$, only for windows with $n - m \ge 8R/b$; short windows near $K$ are not controlled (see Section 3.3).

### 3.4.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [PROVED, UNVERIFIED (SWEEP); 4 OF 19 TABLE ROWS VERIFIED]

Source: `LEMMA_DPRIME_SWEEP.md` (all 12 Balaban files; OCR grep plus page image for every non-single-scale hit). `LEMMA_DPRIME_VERIFY` Steps 4–5 checked comparisons A–E on page images (CMP 122 I p. 192, p. 198 (1.94); II p. 381 ×2, p. 386) and corrected two misreadings of A (factor $(1+\beta_0)^2(k-j)^{\beta_0}$; $\delta'_j = g_j A_1 p_1(g_j)$). The five comparisons map to four table rows: A = row 11, E = row 12, C and D = row 16, B = row 18.

*Result.* 19 cross-scale uses of the coupling sequence (table in SWEEP §7), all in CMP 119 and CMP 122 I–II; none in CMP 95, 96, 98, 99, 102, 109, 116. Each is covered by (b) + (c′) + (S) of Section 3.1:

- items 1, 7, 10, 12, 17 need (P-up) with no constant (e.g. (2.6) line 1 is exactly (P-up));
- items 2, 3, 11, 13, 15, 16, 18 need (P-low) with fixed multiplicative slack $(1+\beta_0)$ or $(1+\beta_0)^2$; item 2 ((2.6) line 2) gives the constant 64/15; item 11 needs only $2.42D'$;
- items 4 ((2.8)–(2.9), $\varepsilon$ and $R$ ratios), 6 (CMP 119 (3.7), (3.8), (3.12), (3.13)), and 9 (CMP 122 I (1.52)), the last two of the form $\varepsilon_{k+1} \le (1+\beta_0)\varepsilon_k$, use both sides; they follow as items 1 and 3 do (item 4 from items 1–3);
- item 14 (CMP 122 I p. 200, $R_{k-N_0+1}$ vs $R_k$) uses the lower side; it follows as item 3 does;
- item 5 (CMP 119 (2.31), (2.46): $\sum_j g_j^{\kappa_0} \le g_n^{\kappa_0 - 6}$) needs linear growth (P-low) and $\gamma^{-2} \ge 2D'$;
- item 0 is CMP 119 Theorem 1 assuming (0.31); its proof works with (2.6)–(2.9);
- item 8 (CMP 119 p. 284) is in a sketch, not load-bearing.

Caveats [OPEN]: (1) the sweep is a grep plus page images, not a line-by-line read; (2) "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) was not located; (3) Balaban says $\beta_0$ "can be chosen arbitrarily small, if $g$ is sufficiently small"; under (P-low) the floor is $\beta_0 \gtrsim D\gamma^2/2$, so $\beta_0$ must be fixed first; no step was found that needs $\beta_0 \to 0$; (4) the sweep has no independent verifier beyond rows 11, 12, 16, 18; (5) `BALABAN_EXCERPTS_CHECK` item 11 lists readings not checked on page images: II (4.37)/(5.1), CMP 119 (2.42), the (2.12)/(2.23) ambiguity, the "$L$ odd" page, the Sect. 3 estimates, and the 1989 bound lists.

### 3.4.4 Lemma D and Lemma E [PROVED+VERIFIED; LEAN]

Sources: `LEMMAS.md` (corrected), `LEMMAS_DE_VERIFY.md` §1.4, §1.5, §2.4; `lean/YmLean/LemmaDE.lean` (`lemma_D`, `lemma_E`, `lemma_DE`).

**Lemma 3.5** (Lemma D)**.** *If $b \le \beta_j(g) \le b'$ on $]0,\gamma]$ with $b > 0$, then for every $K$ and $g \in ]0,\gamma]$ shooting succeeds and $t + b(K-k) \le u_k \le t + b'(K-k)$.*

This is literal (0.31). An earlier draft used a wrong first IVT interval, which the verifier refuted; the corrected interval is $[t+bK,\ t+b'K]$. The nested-interval variant for $\beta$ defined only on admissible histories (`LEMMAS_DE_VERIFY` §1.5) is [PROVED+VERIFIED], not in Lean.

**Lemma 3.6** (Lemma E)**.** *If $\beta_0 \le \beta_j(0) \le \beta_0'$ and $|\beta_j(g) - \beta_j(0)| \le Cg^2$, then for $\gamma \le \sqrt{\beta_0/(2C)}$: $\beta_0/2 \le \beta_j(g) \le \beta_0' + \beta_0/2$.*

The verifier checked this $Cg^2$ form; the $C^1$ form ($C_1 g$, $\gamma \le \beta_0/(2C_1)$) is the same two-line proof [PROVED, TRIVIAL VARIANT]. Lean has the $Cg^2$ form.

#### 3.4.5 History dependence (L5) [OPEN]

Balaban's $\beta_{k+1}$ depends on $(g_0, \dots, g_k)$. Lemma D′ Remark 3 (`LEMMA_DPRIME.md`) states the cube-continuity version (same proof). The gap is as follows (`THEOREM2_PROOF_v2_VERIFY` §F).

- *Domain.* Balaban defines $\beta_{k+1}$ only on histories for which his Theorem 1 induction has run to step $k$ (`LEMMAS_DE_VERIFY` §2 item 1). By the sweep (§2.3), that induction uses (2.6)–(2.9), that is (c′) + (S), on the history, not only $g_i \in ]0,\gamma]$.
- *The new gap.* The clamped shooting of Lemmas 3.2 and 3.4 evaluates $\beta$ on clamped histories $\tilde{g}_i = \varphi(\tilde{u}_i)$. A clamped history can violate the growth side of (c′): from $\max(a,c) - \max(b,c) \ge \min(a-b, 0)$ one gets only $u_m \ge u_n - D'$, not $u_m \ge u_n + \beta(n-m) - D'$. So the clamped shooting evaluates $\beta$ on histories outside Balaban's domain. The clamp proof is valid only under the cube hypothesis, which Balaban does not supply.
- *A variant that avoids it* [NOT WRITTEN]. Use a nested-interval proof in the style of `LEMMAS_DE_VERIFY` §1.5, with domain $D_k = \{u_0, \dots, u_k \ge \gamma^{-2}\}$ of true (unclamped) trajectories. On such trajectories, (c′) up to step $k$ follows from the tail-sum bound (a″) of Lemma D″, and (S) holds by the choice of $\gamma$. So every evaluation of $\beta$ stays inside Balaban's domain. At each step $k$, keep the set of $u_0$ whose trajectory stays in $D_k$; show by IVT, step by step, that this set contains a closed interval on which $u_k$ sweeps a range that contains the next target interval. For Lemma D (all increments positive) this is written. For Lemma D″ the increments can be negative, so the nested intervals need new envelopes, built from the window bounds $[b(n-m) - 2R - (b/4)(n-m), \dots]$. The adaptation is plausible; it is not written and not verified.
- Lean is one-argument only.

### 3.5 The exact structure of the one-loop sums

The window sums of the shooting lemma are sums of one-loop increments $\beta_j(0)$. This section expresses these sums as level-$k$ quantities of a coarse Gaussian theory and isolates the remainder $\kappa_k$ whose boundedness is the content of Theorem K. It also lists the symbol facts, uniform in the level, that Section 3.6 uses.

#### 3.5.1 Telescoping [PROVED+VERIFIED, CORRECTED SCOPE]

$S_k := p \to 0$ $F^2$-coefficient of the composed $k$-step gauge-fixed one-loop integral (U1 (1.2)). Then $S_k = \sum_{j=1}^{k} \beta_j(0)$, $S_0 = 0$ (`BETA_FP` Prop. 1.1; `BETA_FP_VERIFY` item 1). Scope: exact for the $p \to 0$ coefficient at fixed $k$, not for $\log Z_k$ at fixed $p$ (the plane wave is off-shell).

### 3.5.2 Anchor-free identity [PROVED+VERIFIED]

Source: `BETA_FP_LEMMA43` Prop. 2.1; `BETA_FP_LEMMA43_VERIFY` items 1, 3–4 (hand rederivation; lattice check with the real CMP 109 averaging at random SU(2) backgrounds, relative difference $10^{-14}$).

**Proposition 3.7** (anchor-free identity; Prop. 2.1 of `BETA_FP_LEMMA43`)**.** *Under (G1) =* $\{\Delta D = 0,\ \tilde{Q}D = D_c \iota_a^T\}$*:*

$$\log Z_\Lambda(V) = \big[\log\det X - \tfrac{1}{2}\log\det K\big] - \big[\log\det(D_c^T D_c) - \tfrac{1}{2}\log\det(\Delta_\Lambda(V) + D_c D_c^T)\big].$$

In words: constrained one-loop = fine Wilson one-loop in background Feynman gauge minus the one-loop of the coarse Gaussian theory with kernel $\Delta_\Lambda(V)$. All point-anchor objects cancel; the constant is exactly 0.

- $\tilde{Q}D = D_c\iota_a^T$ is exact at any background (lattice $10^{-15}$). $\Delta D = 0$ is load-bearing (raw Hessian gives $O(1)$ errors).
- At the constant abelian field $V_f$ the background is critical, so $\Delta D = 0$ is exact and Prop. 2.1 holds exactly (`BETA_FP_H_VERIFY` §2, CONFIRMED). At a plane wave it holds up to the defect $\mathfrak{d}_Y$ (`BETA_FP_PW` Lemma 2.3) [PROVED, UNVERIFIED].
- Per-cell infinite-volume densities only: on a torus the constant zero modes must be removed. This is input (G2) [FORMAL].

### 3.5.3 The constant $c$ [PROVED+VERIFIED]

$c = 11N^2/(12\pi^2) = 2N\cdot b_0/(8\pi^2)$, $b_0 = 11N/3$; SU(2): $c = 0.371511$. Normalization $1/g_B^2 = 2N/g^2$ (`R2_VERIFY`; `BETA_FP_LEMMA43_VERIFY` §5). The log coefficient of the fine piece at constant field is $c_2 = c/2$ per $\ln(1/\varphi)$, checked against Savvidy, arXiv:1910.00654 eq. (1.4) [23] (`BETA_FP_H_VERIFY` §4) = input (F) [CITED, SOURCE CHECKED; LATTICE = CONTINUUM LOG COEFFICIENT IS STANDARD, NOT PROVED IN PROJECT FILES].

$b := c\log L$: 0.25751 ($L = 2$), 0.40815 (3), 0.51502 (4), 0.59793 (5), 0.72293 (7).

### 3.5.4 The decomposition $S_k = ck\log L + \kappa_W^F - \kappa_k$

- As bookkeeping the split is empty: any $c'$ gives an exact identity if $\kappa_k$ is defined as the remainder (`U1_VERIFY`). Boundedness of a remainder-defined $\kappa_k$ is a restatement, not progress.
- Content comes from defining $\kappa_k$ by the coarse theory alone. `KAPPA_BOUNDED` Prop. 1.1 [PROVED+VERIFIED, CONDITIONAL ON (PI), (F), ($M_j$), V1, (G2)]: with the Wilson level $j = 0$ as reference,

$$S_j = cj\log L - d_j, \qquad \kappa_j = \kappa_W^F + d_j,$$

$$d_j := -\frac{1}{2n_2}\int_0^1 dt \int_{\mathrm{BZ}} \frac{dK}{(2\pi)^4}\ \mathrm{tr}\big[b_{t,2}\,\delta\sigma_j^{(0)} + b_{t,1}\,\delta\sigma_j^{(1)} + b_{t,0}\,\delta\sigma_j^{(2)}\big],$$

  where $\delta\sigma_j^{(m)} = \sigma_j^{(m)} - \sigma_0^{(m)}$ are differences of the $f^m$ Taylor symbols of the coarse kernel at level $j$ and level 0, and $b_{t,m}$ are the Moyal inverse coefficients of the interpolated coarse operator. No limit symbols $\sigma_\infty$ are used.

- Checks [NUMERICAL, REPRODUCED]: $\rho(1) = \kappa_W^F$ (Lüscher–Weisz (5.7) [16]); $L = 2$: $d_1 = -0.372$, $\kappa_1 = 1.114$; $L = 3$: $d_1 = c\log 3 - \beta_1 = -0.256 \pm 0.01$ (`KAPPA_BOUNDED_VERIFY` §4: fits give $-0.254 \cdots - 0.258$). This checks the definition of $d_j$ through $S_j$; it does not evaluate (1.1) from the symbols independently.

- Cor. 3.1 of `BETA_FP_LEMMA43` ($\mathcal{C}(\Lambda) = c\log\Lambda + \kappa_W^F - \rho(\Lambda)$) is exact as an identity. "The whole $\log\Lambda$ sits in the fine piece" is equivalent to $\rho(\Lambda) = o(\log\Lambda)$. It is not a separate fact (`BETA_FP_LEMMA43_VERIFY` item 2; an earlier overclaim is removed).

- Given convergence $\beta_j(0) \to \beta_\infty$, Cesàro gives $\beta_\infty = c\log L - \lim \kappa_k/k$. So the hypothesis (H′) "$\kappa_k = o(k)$" of the first version of these notes is EQUIVALENT to $\beta_\infty = c\log L$, not weaker (`THEOREM2_PROOF_VERIFY` Point 4). Theorem K (Theorem 3.8) gives $\kappa_k$ bounded, hence $\kappa_k = o(k)$ trivially.

#### 3.5.5 Level-uniform symbol facts used below [PROVED+VERIFIED]

Source: `U3_AW.md`, `U3_AW_VERIFY.md` (grades 1–4 CONFIRMED); `U3_REPAIRS_VERIFY`.

- O5/A0: $\Delta_M$ real-analytic on a fixed strip, $|\partial^\alpha(\Delta_M - \Delta_\infty)| \le CM^{-2}$, uniform to $K = 0$ (odd $L$, centered blocks). Prop. B: $c_0|\hat{K}|^{-2}P_T' \le P_T' S_M P_T' \le C_1|\hat{K}|^{-2}P_T'$, $c_0 = (2/\pi)^{12}$, $C_1 \le 1363$. Existential constants.

- Prop. E (evenness): $\sigma_j(f;-K) = \sigma_j(f;K)$ in the midpoint trivialization, all $f$, $j$. Tested in $d = 2$ only.

- Theorem W: $\sigma_j^{(1)}(0) = \tau_c = (3ie/2)(E_{10} - E_{01})$ for every $j \ge 0$, from the exact Ward identity $\Delta_j(f)D_f = 0$. Numerically 0.750000, 0.75015, 0.75051 at $j = 0, 1, 2$ vs 0.75 ($d = 2$ proxy).

- Theorem A3: $|\sigma_j^{(1)}(K) - \tau_c| \le C_A|K|^2$, real $K$, uniform in $j$ ($C^0$ only).

- Prop. A2: $\sigma_j^{(m)}$, $m \le 2$, analytic on the same strip at every fixed $j$ (not uniform in $j$ for $m = 1$ derivatives, $m = 2$).

- Lemma M: Maxwell coefficient exactly 1 at every level.

### 3.6 Bounded window sums: two formally distinct conditional routes

This section gives the two routes to the window bound (5.1) that Lemma 3.4 needs.

Both routes give the window bound (5.1) of Lemma 3.4, $|S_K - S_k - b(K-k)| \le 2R$, with $b = c\log L$. Lemma D″ then gives Theorem 2w (at odd $L > 11$, or with (H0) at this $L$ as a hypothesis, and with L5). Neither route uses convergence of $\beta_j(0)$, a rate, (N), or certified early steps. The two routes have formally distinct named hypotheses. Their mathematical content is not independent: both reduce their uniform-in-level content to the same inequality (H2) on the two-step minimizer map (Section 3.7), plus A3-type facts (`BETA_FP_U` §3.5, §4.3 [PROVED, UNVERIFIED]; Route K side: `B2_TWOSTEP_VERIFY` grade 3 [PROVED+VERIFIED]).

#### 3.6.1 Route K: Theorem K [PROVED+VERIFIED, CONDITIONAL ON (B2) AND THE FIXED-LEVEL INPUTS]

Source: `KAPPA_BOUNDED.md`; `KAPPA_BOUNDED_VERIFY.md` (items (i)–(v) and (vii) CONFIRMED; (vi) CORRECTED in reasoning, conclusion unchanged).

**Theorem 3.8** (Theorem K). *$L$ odd $\geq 3$. Assume (B2) (Section 3.7; (B2) $\Leftarrow$ (H2) [PROVED+VERIFIED]) and the fixed-level inputs below. Then $\sup_k |d_k| \leq \kappa_* < \infty$, so $\sup_k |\kappa_k| \leq |\kappa_W^F| + \kappa_*$ and $|S_k - ck \log L| \leq \kappa_*$ for all $k$. Window sums satisfy (5.1) with $R = \kappa_*$. (Theorem K is a statement about one-loop quantities. Its use in Lemma D′ for Balaban's $\beta_j(g)$ needs $L$ odd $> 11$ or (H0) at this $L$.)*

*Proof structure.*

1. Prop. 1.1 (Section 3.5.4): $S_j = cj \log L - d_j$. [PROVED+VERIFIED, CONDITIONAL ON (PI), (F), ($M_j$), V1, (G2)]
2. Lemma 2.1: the only $K$-derivative of $\sigma^{(1)}$ in $d_j$ enters linearly with $f^0$ coefficients. Integration by parts moves it onto $f^0$ data; boundary terms vanish (periodic up to a common constant conjugation; inner sphere $O(r^2)$). So $\sigma^{(1)}$ is needed only in $C^0$, which Theorem A3 gives. [PROVED+VERIFIED]
3. Prop. 2.2: with (E0) $f^0$ bounds (Prop. B, O5/A0, Lemma M), (E1) Theorem A3 + Theorem W, and (B2), every term of the $d_j$ integrand is $\leq C|K|^{-2}$, uniformly in $j$ and $t$. In $d = 4$ this has two powers of infrared margin. $|d_j| \leq \kappa_* := (C(L)/n_2)(1 + Y_1^2 + C_A(1 + Y_1) + Y_2) \cdot I_2$. [PROVED+VERIFIED, CONDITIONAL ON (B2)]
4. Infrared: the degree $-4$ part of the coarse $f^2$ integrand is the same at every level (Lemma M + Theorem W), so the $\tau_c \log$ cancels exactly in level differences. [PROVED+VERIFIED]

Constants are existential; $\kappa_*$ is not evaluated.

Fixed-level inputs of Route K (needed at each $j$ separately; no uniformity in $j$):

- (PI) probe independence: plane-wave $p^2$ coefficient = constant-field $f^2$ coefficient [ARGUED; PER-CELL LIMIT FOR THE LINEAR POTENTIAL NOT WRITTEN].
- (F) fine log coefficient $c_2$ at constant field [CITED, SOURCE CHECKED], Section 3.5.3.
- ($M_j$) the formal $f^2$ coefficient $d_j$ equals the true one at fixed $j$. The resolvent step of the first argument was false (Nielsen–Olesen zero at $k_\perp^2 = |ef|$; `BETA_FP_H_VERIFY` §5). `BETA_FP_M` repairs it with a direct $\log|\det|$ bound given spectral bounds (S1)–(S3) [OPEN ON THE LATTICE]; Prop. 4.4 there is [ARGUED]; `BETA_FP_M` has no verifier file. The $t$-subdivision of KAPPA §1.3 removes the small-$\varepsilon$ requirement [ARGUED, CONFIRMED AS PLAUSIBLE]. Status: [ARGUED].
- V1: the Maxwell coefficient of the per-step determinant equals $\beta_{j+1}(0)$, i.e. order-$g^0$ completeness of CMP 109 (1.3)–(1.5) [OPEN; SUPPORTED AT ORDER $p^2$ BY `U3_VERIFY` §6].
- (G2) per-cell infinite-volume densities [FORMAL].
- (H0) Balaban's inductive assumption: $\beta_{j+1}$ smooth on $[0, \gamma]$, "uniformly bounded on this interval together with all derivatives" (CMP 109 pp. 263–264, (1.22)) [BALABAN; AVAILABLE AT ODD $L > 11$ ONLY]. CMP 116 has no theorem statement and no $C^m$-in-$g$ bound in words (`BALABAN_EXCERPTS_CHECK` item 10); the smoothness in $g$ is CMP 109's inductive assumption, asserted by CMP 109 Theorem 3, whose proof CMP 116 completes.
- L5 history-dependent $\beta$, its domain, and its continuity (Section 3.4.5) [OPEN].

### 3.6.2 Route B: Theorem $B_{PW}$ [PROVED+VERIFIED, CONDITIONAL (LOGIC OF THEOREM $B_{PW}$)]

Source: `BETA_FP_PW.md` §1–§5 (Theorem 3.2 = Theorem $B_{PW}$, Theorem 5.1). `BETA_FP_PW_VERIFY.md` §6 grades it: items 1–4 CONFIRMED (doubling identity and Lemma 1.3; Prop. 2.2 / Lemma 2.3; Theorems A, B as conditional logic; Lemma 4.1, conditional), items 5 (comparison) and 6 (numerics) CORRECTED. The tag follows the decision recorded in `PLAN.md` (2026-10-28): the logic of Theorem $B_{PW}$ is [PROVED+VERIFIED, CONDITIONAL] on its hypotheses, citing `BETA_FP_PW_VERIFY` §6 items 1–4. Its inputs (U), (DF), (T1) from `BETA_FP_U` stay [PROVED, UNVERIFIED]. The file rederives the Lemma 1.3 arithmetic and the Lemma 2.3 / Prop. 2.2 algebra correctly, confirms the Prop. 2.2 determinant identity on a $4^4$ lattice (4e−13), and notes that identity 3.0 is a tautology ($\Gamma^{ex} := \hat{\Pi} - S$), so all content sits in $(B\theta)/(C\theta)$.

**Theorem 3.9** (Theorem $B_{PW}$; Theorem 3.2 of `BETA_FP_PW`)**.** *$L$ odd $\geq 3$. Assume*

- *(U) there are $p_* > 0$, $C_U$ with $|\beta_{j+1}(p) - \beta_{j+1}(0)| \leq C_U p^2$ for all $j \geq 0$, $0 < p \leq p_*$* [HYPOTHESIS; ARGUED FROM UNIFORM ANALYTICITY IN $p$ OF THE ONE-STEP PLANE-WAVE DATA PLUS ONE-STEP O6]*;*
- *($B\theta$) there is $\theta_* \in (0, \min(p_*, 1))$ with $B_* := \sup_k |\Gamma^{ex}(\theta_*; L^k)| < \infty$, where $\Gamma^{ex}$ is the coarse remainder at fixed coarse momentum $\theta_*$* [HYPOTHESIS]*;*
- *Lüscher–Weisz (5.7) [16]* [CITED].

*Then $|S_k - ck \log L| \leq C_B := B_* + c_2 \ln(1/\theta_*^2) + |\kappa_W^F| + \sup_{p \leq \theta_*} |\eta(p)| + C_U \theta_*^2/(L^2 - 1)$, and window sums satisfy (5.1) with $R = C_B$. $\theta_* < 1$ is required so that $c_2 \ln(1/\theta_*^2) \geq 0$; otherwise the term must be replaced by its absolute value (`BETA_FP_PW_VERIFY` §3).*

*Key step.* Lemma 1.3: $|S(\theta; L^k) - S_k| \leq C_U \theta^2/(L^2 - 1)$ from (U). The massless coarse side is then needed only at a fixed $\theta_* > 0$, where there is no infrared problem. No (W), (PI), (M), IR weights, or spectral theory.

What ($B\theta$) reduces to (`BETA_FP_PW` §4) [ARGUED]: Lemma 4.1(i) (trivial-background convergence, from O5, odd $L$) [PROVED]; vertex bounds at fixed $\theta_*$ (the uniform bounds cited by U3 Theorem 7.1 to Balaban; the fixed-$\theta$ data hypothesis ($D\theta$) is needed only in its "bounded" form); and boundedness in $k$ of the defect $\mathfrak{d} = \mathfrak{d}_Y + \mathfrak{d}_{gf}$.

- $\mathfrak{d}_Y$ (Prop. 2.1 defect at the non-stationary plane wave): $O(a\Lambda^{-2})$ relative at fixed $\theta$, from zero capacity of point anchors in $d = 4$ [ARGUED; REGION $|K| \lesssim \Lambda^{-1}$ NOT WRITTEN].
- $\mathfrak{d}_{gf}$ (composite gauge vs $D_k^T$ gauge at the plane wave): $O(\theta^2)$ relative; uniformity in $k$ not shown [ARGUED]. Data: 0.953 (direct) vs 0.9595 (recursion) at $L = 2$, $p_0 = 0.2$; difference below the tabulation error, so unresolved.

*Identification.* Route B's $\beta_j(0)$ is the per-level plane-wave increment (the quantity the code computes). It matches Balaban's second-moment $\beta$ up to the expansion point (fine wave vs constrained minimizer), argued to change $S$ by $O(\theta^2)$ relative (`SOURCES_D1_VERIFY` item 8) [ARGUED]. This identification and the $\mathfrak{d}_{gf}$ bound need expansion-point bounds uniform in $k$. So the (PI)-type content of Route K is not removed in Route B; it moves into a uniform-in-$k$ input (`BETA_FP_PW_VERIFY` §3).

Three points from `BETA_FP_PW_VERIFY` §3 (item 5 of its §6 grades them CORRECTED):

1. $\theta_* < 1$ is required (above).

2. The $\beta$ identification (B4) is a uniform-in-$k$ input, not a fixed-level input.

3. (U) is a uniform-in-$j$ analyticity of the level data. It plausibly needs the plane-wave analogue of a uniform $C^1$ bound on $\sigma^{(1)}$: uniform $p$-smoothness of the $a^1$ vertex. That is the same uniform first-order vertex smoothness as Route K's open core ($C1\sigma$).

#### 3.6.3 Route B inputs after `BETA_FP_U` [PROVED, UNVERIFIED; NO VERIFIER FILE]

Source: `BETA_FP_U.md` §2–§4. It reduces the two uniform hypotheses of Theorem $\mathrm{B_{PW}}$.

- (U) $\Leftarrow$ (T1) + (H2) (Theorem 2.4, odd $L \geq 3$, with A0, A1, Lemma 2.2). (T1): $\sup_j \sup_q |c^{(1)}_{T,j}(q)| < \infty$, the plane-wave form of Theorem A3 [ARGUED]. (H2) is the inequality of Section 3.7 (`BETA_FP_U` writes it as $\tilde{h}_{(2)} < L$).

- ($B\theta$): $\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$ exactly (Prop. 3.1). $\gamma_{rec}$ is bounded in $k$ given Theorem 2.4 (Prop. 3.2). $\mathfrak{d}_F$ is bounded under (DF) $\sup_k |\mathfrak{d}_F(\theta_*; L^k)| < \infty$ [HYPOTHESIS, ARGUED]. `BETA_FP_U` corrects `BETA_FP_PW` §4.3(b): $\mathfrak{d}_F$ is $O(1)$ relative at fixed $\theta$, not $O(\theta^2)$; only boundedness is used.

- Corollary 4.1: given (T1), (H2), (DF), LW (5.7), $|S_k - ck \log L| \leq C_B$, so (5.1) holds with $R = C_B$.

- Theorem 2w by Route B then needs in addition A6 (plane-wave $\beta$ = Balaban's $\beta$ up to the expansion point, uniform in $k$) [ARGUED] and (H0) (Lemma D″ (i)).

Route B hypothesis set: (T1), (H2), (DF), LW (5.7), A6, (H0) (plus L5, (G2), odd $L$, as for Route K). The fixed-level inputs (PI), ($M_j$), V1 of Route K do not appear; (DF) and A6 replace them.

#### 3.6.4 Comparison

| | Route K | Route B |
|---|---|---|
| uniform-in-level input | (B2): $\sup_j \sup_K \lvert\sigma^{(2)}_j\rvert < \infty$, $\Leftarrow$ (H2) [PROVED+VERIFIED] | (U), ($B\theta$), and the $\beta$ identification and $\mathfrak{d}_{gf}$ bound, both with $k$-uniform expansion-point bounds; (U) $\Leftarrow$ (T1) + (H2), ($B\theta$) $\Leftarrow$ Theorem 2.4 + (DF) [PROVED, UNVERIFIED] |
| single uniform inequality | (H2) | (H2) |
| fixed-level inputs | (PI), (F), ($M_j$), V1, (G2) | LW (5.7), $\mathfrak{d}_Y$, $\theta_* < 1$ |
| uses (W), Theorem A3 | yes | no; `BETA_FP_U` uses (T1), the plane-wave analogue of A3 [ARGUED] |
| verification | verified (conditional) | logic of Theorem $\mathrm{B_{PW}}$ verified (conditional): `BETA_FP_PW_VERIFY` §6 items 1–4 CONFIRMED [PROVED+VERIFIED, CONDITIONAL]; inputs from `BETA_FP_U` [PROVED, UNVERIFIED], no verifier file |

| | |
|---|---|
| common | (H0) at odd $L > 11$, L5, (G2), V1-type identification, Balaban's proofs, SU(2) |

Shared inputs: (H0), L5, (G2), odd $L$ ($> 11$ for (H0)), and the identification of the computed $\beta_j(0)$ with Balaban's $\beta_j(0)$ (V1; in Route B also the expansion point). The uniform-in-level hypotheses ((B2) vs (U) + $(B\theta)$) are formally distinct as named hypotheses. Their content overlaps: both rest on (H2) (Section 3.7). In Route K, (H2) gives $(C1\sigma)$ (Theorem C1) and then (B2) (Theorem B2). In Route B, (H2) gives (U) with (T1), and the vertex bound behind $(B\theta)$. So a proof of (H2) serves both routes. One closed route suffices for Theorem 2w.

## 3.7 The hypothesis (B2) of Route K, and the single uniform inequality (H2)

Both routes need a bound on the coarse symbols that is uniform in the level. This section states that bound for Route K, records what is proved and what is refuted, and reduces it to the inequality (H2). The numerical values of (H2) are in Sections 5.4–5.8. At $L = 13$, (H2) is a certified theorem (Theorem 5.1, Section 5.7).

(B2): $\sup_{j\geq 0}\sup_{K\in \mathrm{BZ}} |\sigma_j^{(2)}(K)| < \infty$. Weaker form that suffices, (B2w): $\sup_j \int_{\mathrm{BZ}} |K|^{-2}|\sigma_j^{(2)}(K)|\, d^4K < \infty$ ($\sigma^{(2)}$ enters $d_j$ only undifferentiated; `KAPPA_BOUNDED` Remark 2.3) [PROVED+VERIFIED: NOT GRADED BY NAME; IMPLIED BY `KAPPA_BOUNDED_VERIFY` §9(IV)].

$\sigma^{(2)}$ obeys the one-step recursion (`U3_AW` (1.2)): $\sigma_j^{(2)} = L^{-2}\mathcal{A}_j\sigma_{j-1}^{(2)} + Q_j[\sigma_{j-1}^{(1)}] + g_j^{(2)}$, with $\mathcal{A}_j s = L^{-2}H_j^\dagger(\oplus_l s(k_l))H_j$ and $H_j(K)$ the one-step gauge-fixed minimizer map. Notation: $\rho_1 := \sup_{j,K}\|H_j(K)\|/L$ (one step); $H_{(2),j}(K) := (\oplus_l H_{j-1}(k_l))H_j(K)$ (two steps), $\rho_2 := \sup_{j\geq 2}\sup_{K \text{ real}}\|H_{(2),j}(K)\|/L^2$.

### 3.7.1 Proved

- $K = 0$ bound [PROVED+VERIFIED]: $\sup_j |\sigma_j^{(2)}(0)| \leq C(|e|C_A + e^2)$. Theorem A3 + scaling give $|\partial^2\sigma_j^{(1)}(0)| \leq 2C_A$ (polarization; `KAPPA_BOUNDED_VERIFY` §7), and the slaving identity (W-2) expresses $\sigma_j^{(2)}(0)$ through these second derivatives.

- Each fixed $j$ [PROVED+VERIFIED]: $\sigma_j^{(2)}$ is analytic on a fixed strip (Prop. A2), hence bounded. The content of (B2) is uniformity in $j$.

- Prop. 4.1 [PROVED+VERIFIED, CONDITIONAL]: if (h) $h := \sup_{j,K}\|H_j(K)\| < L^2$ and $(C1\sigma)$ $\sup_j\sup_K|\partial_K\sigma_j^{(1)}(K)| < \infty$, then (B2) holds. Proof: in the form norm $\pm s \leq N(\Delta + 1)$, $N_j \leq (h^2/L^4)N_{j-1} + Z_2$, and $Z_2$ is finite by Theorem A3, $(C1\sigma)$, and fixed $f^0$ vertices.

- Why the Lemma A4 mechanism alone fails [PROVED+VERIFIED, REASON CORRECTED BY `KAPPA_BOUNDED_VERIFY` §7]: the obstruction is the missing uniform $C^1$ bound on $\sigma^{(1)}$, which enters twice: in $Q_j$ (term $L^{-3}\partial_K\sigma_{j-1}^{(1)}$) and in the longitudinal part of $\sigma^{(2)}$ given by (W-2). Theorem A3 is $C^0$ only.

### 3.7.2 Refuted, and the failed one-step bound

- (HS) $h < L\sqrt{L-1}$ (named (H′) in `U3_AW`) [REFUTED AT $L = 3$]. Measured $\max_K\|H_j(K)\|/L = 1.652, 1.736, 1.757, 1.760$ for $j = 1, 2, 3, \infty$ (`U3_AW_VERIFY` §6); the

threshold is $\sqrt{2} = 1.414$. Gauge choice cannot help: $H_j$ is already minimal-norm in its gauge orbit.

- One-step sup-norm $C^1$ recursion at $L = 3$ [NUMERICAL, REPRODUCED; `B2_TWOSTEP_VERIFY` GRADE 2]. Its factor is $\rho_1^2/L = 0.909, 1.0046, 1.0289, 1.0325$ for $j = 1, 2, 3, \infty$, so it exceeds 1 for $j \geq 2$ ($\rho_1 > \sqrt{3} = 1.7321$). At $K = 0$ the factor is $L^{-1}$; the bad region is the face $K_3 = K_4 = \pi$, $K_1 = K_2 \approx 1.86$–$1.92$. This shows that the one-step sup-norm bound fails at $L = 3$. It does not show that a $C^1$ bound fails; a weighted norm is not excluded.

### 3.7.3 (H2) and the two-step theorems [PROVED+VERIFIED, CONDITIONAL ON (H2)]

Source: `B2_TWOSTEP.md` §1–§3; `B2_TWOSTEP_VERIFY.md` grades 1, 3, 4 (CONFIRMED).

$$\text{(H2)} \quad \rho_2 = \sup_{j \geq 2} \sup_{K \in \mathrm{BZ\ real}} \|(\oplus_l H_{j-1}(k_l))H_j(K)\|/L^2 < L.$$

(H2) is a plain operator-norm inequality on real $K$. It uses no strip, no $\Delta$-relative weight, and only $n = 2$.

**Theorem 3.10** (Theorem C1 of `B2_TWOSTEP`). *(H2) ⇒ (C1σ), with contraction factor $q_1 = \rho_2^2/L^2$ on the derivative.*

**Theorem 3.11** (Theorem B2 of `B2_TWOSTEP`). *(H2) ⇒ (B2), hence (B2w), in plain sup norm, with factor $q_2 = \rho_2^2/L^4$.*

Here $(C1\sigma)$ is $\sup_j \sup_K |\partial_K \sigma_j^{(1)}(K)| < \infty$ (Section 3.7.1). The verified points are these.

- Two-step map. $\mathcal{A}_j \mathcal{A}_{j-1} s = L^{-4} H_{(2)}^{\dagger}(\oplus s) H_{(2)}$, with $dk_{l,l'}/dK = L^{-2}$ [CONFIRMED, grade 1].
- Theorem 3.10. The chain-rule gain is exactly $L^{-2}$ per two steps on real $K$. So the threshold is $\rho_2 < L$, not $\sqrt{L^2 - 1}$: no strip-Cauchy loss enters. The large constants ($b(L)$ of O6, $\eta_1$, $\eta_2$, $G_1$) enter only the source term, not the contraction factor [CONFIRMED, grade 3].
- Theorem 3.11. This contraction needs only $\rho_2 < L^2$, which (H2) implies [CONFIRMED, grade 3]. In one step it is Prop. 4.1's (h).
- (H2) vs (H″). (H2) is the plain operator norm of the same matrix as `U3_AW` (H″) at $n = 2$. It is weaker: threshold $L$ (3 at $L = 3$) against $\sqrt{L^2 - 1}$ (2.828), real $K$ only, $n = 2$ only [CONFIRMED, grade 4]. (H″) ⇒ (B2) stays [ARGUED] and is no longer needed.
- Form of $Q_j$ (G3) [CONFIRMED by read-off of `U3_AW` (1.2)/(3.3) and P10]: $\partial\sigma^{(1)}$ enters linearly. `U3_LEMMA51` (3.3) itself was not re-derived from the Moyal rule by either party.
- Not covered: $C^2$ and $C^3$ bounds on $\sigma^{(1)}$ under (H2) (`B2_TWOSTEP` §5, "look routine", not written, not graded). They are not needed for (B2).
- Submultiplicativity: $\rho_2 \leq \rho_1^2$ always. At $L = 3$ the gain of the two-step map is that this is strict by a factor 1.50 (2.064 vs 3.098) [CONFIRMED].

### 3.7.4 One-step sufficiency [ARITHMETIC PROVED; INPUT CERTIFIED AT $L = 13$]

$\|H_{(2),j}(K)\| \leq \max_l \|H_{j-1}(k_l)\| \cdot \|H_j(K)\| \leq L^2 \rho_1^2$. So $\rho_1^2 < L$ implies (H2), and then the one-step $C^1$ recursion of `B2_TWOSTEP` (2.2) contracts with factor $\rho_1^2/L$. At $L = 3$ the measured $\rho_1^2$ exceeds $L$, so the one-step route fails and the two-step map is needed; at $L = 5$ and $L = 7$ the measured $\rho_1^2/L$ is below 1. The values are in Section 5.5. At $L = 13$ the input is certified: $\sup_{j,K} \rho_1^2 \leq 10 < 13$ (Theorem 5.1), so (H2) holds there with $\rho_2 \leq 10$.

### 3.7.5 The precise remaining inequality

At a fixed odd $L \geq 3$ the single uniform hypothesis is (H2).

- Route K: (H2) $\Rightarrow$ $(C1\sigma)$ (Theorem C1) $\Rightarrow$ (B2) (Theorem B2) [PROVED+VERIFIED, CONDITIONAL ON (H2)].
- Route B: (H2) + (T1) $\Rightarrow$ (U), and (H2) $\Rightarrow$ the vertex bound for $\gamma_{rec}$ in $(B\theta)$ (`BETA_FP_U` Theorem 2.4, Prop. 3.2) [PROVED, UNVERIFIED].
- Status of (H2): at $L = 13$, inside Bałaban's range $L > 11$, [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]: $\rho_1(j, K)^2 \leq 10 < 13$ for all $j \geq 1$ (including $\infty$) and all $K$, hence $\rho_2 \leq 10 < 13$ (Theorem 5.1; residue in Section 5.7). At $L = 3, 5, 9$: [NUMERICAL] only (at $L = 9$ one point).
- Kept as alternatives: $(C1\sigma)$ + (h) $\Rightarrow$ (B2) [PROVED+VERIFIED]; (H″) $\Rightarrow$ (B2) [ARGUED]; (B2w) directly. No per-momentum $d = 4$ constant-field code exists, so (B2) itself has not been scanned numerically (`KAPPA_BOUNDED_VERIFY` §8).

## 3.8 Conclusion of the reduction

Theorem 2w $\Leftarrow$ (Route K inputs) or (Route B inputs). Precisely:

*Route K.* (B2) [OPEN; REDUCES TO (H2) (VERIFIED REDUCTION, SECTION 3.7.3); (H2) CERTIFIED AT $L = 13$ (THEOREM 5.1)] + (PI), (F), $(M_j)$, V1, (G2) [FIXED-LEVEL] + (H0) [BALABAN, ODD $L > 11$] + L5 [OPEN] $\Rightarrow$ Theorem K ($R = \kappa_*$) $\Rightarrow$ (5.1) $\Rightarrow$ Lemma D″ $\Rightarrow$ Theorem 2w. Status of the implication: [PROVED+VERIFIED, CONDITIONAL].

*Route B.* (U) + $(B\theta)$ with $\theta_* < 1$ + LW (5.7) + bounded $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ + identification of $\beta$ (uniform in $k$) + (G2) + (H0) [ODD $L > 11$] + L5 $\Rightarrow$ Theorem $\mathrm{B_{PW}}$ ($R = C_B$) $\Rightarrow$ (5.1) $\Rightarrow$ Lemma D″ $\Rightarrow$ Theorem 2w. Status: Theorem $\mathrm{B_{PW}}$ logic [PROVED+VERIFIED, CONDITIONAL]; inputs [PROVED, UNVERIFIED]. With `BETA_FP_U` (Section 3.6.3): (T1) + (H2) + (DF) + LW (5.7) $\Rightarrow$ Corollary 4.1 ($R = C_B$); with A6 + (H0) + L5 + (G2) $\Rightarrow$ Theorem 2w. Status: [PROVED, UNVERIFIED, CONDITIONAL] (the inputs (T1), (DF), A6 and the `BETA_FP_U` reductions are unverified).

*Single uniform hypothesis.* Both routes need (H2). Theorem 5.1 certifies (H2) at $L = 13$, an odd $L > 11$. This discharges (B2), the only uniform-in-level hypothesis of Route K (Theorems C1, B2). In Route B it reduces (U) to (T1); (DF) and A6 remain.

*Then Balaban's Theorem 1.* Theorem 2w gives (b) and (c′). With $\gamma$ shrunk so that (S) holds, every one of the 19 cross-scale uses in CMP 119 and CMP 122 I–II is covered (Section 3.4.3), and the single-scale uses need only $g_k \leq \gamma$. So Theorem 1 holds for the bare coupling $g_0(K, g)$, for every $K$ and every $g \in ]0, g_1]$, without Balaban's coupling assumption. This needs L5 in the sharpened form of Section 3.4.5: $\beta$ must exist on every history the shooting visits. Status: [PROVED, UNVERIFIED (SWEEP), CONDITIONAL ON ROUTE K OR ROUTE B INPUTS, ON L5, AND ON BALABAN'S PUBLISHED PROOFS].

*Order of constants.* Fix $\beta_0, p_0, p_1, r, \nu, \kappa_0$ (Balaban). Fix $L$ (odd, $> 11$), then $R$ from Route K or B, then $b = c \log L$, then $\gamma \leq \min(\gamma_0, \gamma_{Bal}, b/(4C_1), \gamma_*)$ with $\gamma^{-2} \geq c_{sw} \cdot 2R$. Then $g_1 = (\gamma^{-2} + 2R)^{-1/2}$. Balaban's phrase "for a sufficiently small positive $g$" fits $g \leq g_1$.

*What this does not give.*

- Literal (0.31): only for windows of length $\geq 8R/b$. Literal (0.31) at every $K$ needs (L1) or (L2) of Section 3.3, which need individual increments, not window sums. Not needed for Theorem 1.

- The value $\lim \beta_j(0)$ and any rate: Section 3.9.
- Even $L$, or odd $L \leq 11$ without (H0) as an extra hypothesis (Balaban assumes $L$ odd $> 11$ throughout CMP 109); SU($N$) with $N > 2$.
- Monotone couplings: $g_m$ can exceed $g_n$ for $m < n$ by the bounded amount $u_m \geq u_n - D'$. Section 3.4.3 shows Balaban's proof tolerates this under (S); a line-by-line read was not done.
- A continuum limit (`ASSEMBLY.md` treats the assembly separately; not claimed here).

## 3.9 The convergence route: what it adds, and its status

Not needed for Theorem 1. It adds: (i) existence of $\beta_\infty = \lim \beta_j(0)$; (ii) its value $c \log L$; (iii) a rate; (iv) eventual positivity, hence literal (0.31) for $K \geq K_0$ (Lemma D′ (d)); (v) with certified early steps, literal (0.31) at every $K$ (Section 3.3 (L2)). It also implies (B2), since convergence of $\sigma_j^{(2)}$ implies boundedness. So it needs strictly more than Route K.

### 3.9.1 Components and status

- U3 Theorem 4.1 with O5: $|\partial_K^\alpha(\Delta_M - \Delta_\infty)| \leq CM^{-2}$, $|\alpha| \leq 2$, uniform to $K = 0$, odd $L$ [PROVED+VERIFIED] (`U3_VERIFY`, `U3_REPAIRS_VERIFY`). Numerically $|\Delta_M - \Delta_\infty| =$ 1.007, 0.129, 0.0147 at $M = 3, 9, 27$.
- O6: $\|\mathfrak{B}_j(K)^{-1}\| \leq b(L)$ for all $j$, $K$, one-step constraint, trivial background [PROVED+VERIFIED]. Crude constant ($b(3) \approx 7.5 \cdot 10^{33}$); measured corner values 103.5, 176.7, 193.0, 195.3 ($j = 0, 1, 2, \infty$, $L = 3$) use a surrogate $\Lambda$ (`U3_REPAIRS_VERIFY` §5).
- Theorem 6.1: $\beta_{j+1}(0) = \Phi(\mathcal{D}_j)$ is Lipschitz in the level data [PROVED+VERIFIED, CONDITIONAL ON V1]. The norm is the Taylor-data norm (`U3_VERIFY` P8 correction).
- Lemma 5.1 (convergence of $\sigma_j^{(1)}$, $\sigma_j^{(2)}$ in $C^2$, $|\alpha| \leq 2$):
  - neutral sector, rate $L^{-4j}$ [PROVED+VERIFIED FOR THE VALUE BOUND; THE $K$-DERIVATIVE PARAGRAPH IS PROVED, UNVERIFIED];
  - (W) [PROVED+VERIFIED] (`U3_AW` Theorem W); it removes the marginal direction of the $f^1$ recursion;
  - (A) qualitative and Theorem A3 [PROVED+VERIFIED]; (A) in full (strip/$C^3$ for $m = 1$, all of $m = 2$) $\Leftarrow$ (H″) [OPEN]; the route via (HS) is [REFUTED] at $L = 3$;
  - $\sigma^{(1)}$ convergence at rate $jL^{-2j}$ [SKETCH IN `U3_AW` §4.7, UNVERIFIED]; $\sigma^{(2)}$ convergence [OPEN, NEEDS (A) IN FULL];
  - the single-block-constraint version of Lemma 5.1 is false for $m = 2$ in $d = 4$ [PROVED+VERIFIED]; Balaban's composed constraint is not affected.
- Prop. 3.6: $|\beta_{j+1}(0) - \beta_\infty| \leq C(1 + j)^2 L^{-2j}$ [PROVED, UNVERIFIED AS AN ASSEMBLY; CONDITIONAL ON V1, (A) IN FULL]. (The form $Cj^2L^{-2j}$ of the first version of these notes is 0 at $j = 0$ and is corrected here.) The identification of $\sigma$-Taylor data with the data of (6.1) is cited from `U3_VERIFY` §7, not re-shown. (N) (clean rate $L^{-2j}$) is not needed and is judged plausibly false.

- Value of the limit. Given convergence, Theorem K or Theorem $\mathrm{B_{PW}}$ gives $\beta_\infty = c \log L$ at once: $S_k/k \to \beta_\infty$ by Cesàro and $S_k/k = c \log L + O(1/k)$ [PROVED+VERIFIED AS AN IMPLICATION: `THEOREM2_PROOF_VERIFY` POINT 4; `BETA_FP_PW` THEOREM 3.2(III)]. Older routes to the value: `BETA_FP_H` Theorem 3.1′ with (H) restated at the constant field, which given (W) reduces to Lemma 5.1 (1.2) alone (`BETA_FP_H_VERIFY` items (1)–(4)), plus (PI), (F), (M) [CONDITIONAL; (M) ARGUED]; and `BETA_FP_PW` Theorem 3.1 (limit form) given (U), ($C\theta$), LW (5.7) [PROVED, UNVERIFIED]. The original (H) with IR weights (`BETA_FP_LEMMA43` (4.2)) was graded UNSUPPORTED and is not used.

- Rate of $\mathcal{C}(\Lambda)$: $O(\Lambda^{-2} \log \Lambda)$ given Lemma 5.1 with rates, via `U3_AW` §3.4(b) slaving [ARGUED, `BETA_FP_H_VERIFY` §5].

### 3.9.2 Early steps and $j_*$

Under Prop. 3.6 and $\beta_\infty = c \log L$, $\beta_{j+1}(0) \geq (c \log L)/2$ for $j \geq j_* := \min\{J : C(1+j)^2 L^{-2j} \leq (c \log L)/2 \text{ for } j \geq J\}$. Heuristic size [ARGUED]: $C \sim b^3$ gives $j_* \approx 110$ ($L = 3$), $\approx 70$ ($L = 13$); $C \sim b^6$ (one-step inverses inside $C_{data}$) gives $\approx 218$, $\approx 138$ (`THEOREM2_PROOF_VERIFY` Point 5(e)). With the measured $\|\mathfrak{B}^{-1}\| \approx 195$: $j_* \approx 10$ ($L = 3$), $\approx 4$ ($L = 13$). Certifying $\beta_1(0), \ldots, \beta_{j_*}(0) > 0$ needs interval arithmetic at odd $L \geq 13$ (blocks $\approx 3.4 \cdot 10^5$ rows per momentum by dense methods; the umklapp code reduces this to $12 \times 12$ blocks at $L^4$ momenta). It is needed only for (L2), not for Theorem 1.

# 4 The finite-torus existence theorem

This section proves Theorem B. It takes the trajectory of Section 3 as the input block (B-T2). Section 4.1 fixes what "exists" means here. Section 4.2 states the uniform bound (H-U) that the blocks must deliver, and Sections 4.3–4.4 state the theorems. Section 4.5 proves them modulo the named blocks, and Section 4.6 gives the dependency table, which is the single status table of the paper. Section 5 then collects the numerical support for the hypotheses.

Source: `ASSEMBLY.md` (2026-10-27), verified in `ASSEMBLY_VERIFY.md`. The statements carry the verifier's corrections (Section 4.8). The theorem is a reduction to named blocks. It is not unconditional. The notation is that of Sections 2.1 and 2.5.

## 4.1 What "exists" means here

Jaffe–Witten footnote 2 [15] [VERIFIED QUOTE] (`reports/clay_ym.txt` line 619; anchor: the sentence "Such a result presumably would enable the study of the limit as $T^4 \to \mathbb{R}^4$", line 582): "We specifically exclude weak-existence (compactness) as the solution to the existence part of the Millennium problem, unless one also uses other techniques to establish properties of the limit (such as the existence of a mass gap and the axioms)."

Reading used here. "Exists" means: (i) the family $\{S_{n,\varepsilon}\}$ is relatively compact (uniform bounds); (ii) at least one limit point exists along a sequence $\varepsilon_k \to 0$; (iii) every limit point has the properties (OS0)–(OS3) below, proved on the limit, not assumed. No uniqueness is claimed. Full-sequence convergence needs E13 (comparison of two trajectories; `SOURCES.md` (E14-d), D6), which is not in this paper.

This reading is weaker than the footnote's parenthesis. The theorem below gives no gap and only the torus versions of four axioms. It is NOT an admissible Clay solution. It is the finite-torus step of Jaffe–Witten §6.5 [VERIFIED QUOTE]: "One must then verify the existence of limits of appropriate

expectations of gauge-invariant observables as the lattice spacing tends to zero and as the volume tends to infinity" (only the first limit is done here).

The OS axioms are not stated in `reports/clay_ym.txt`. It cites them as OS (CMP 31 (1973), CMP 42 (1975)) [19, 20] and requires "axiomatic properties at least as strong as those cited" (lines 279–280) [VERIFIED QUOTE]. The forms used follow the template's application of OS II (`reports/o3_continuum.txt` lines 4987–5003) [VERIFIED QUOTE]: "The linear growth condition requires a bound $|S_n(F)| \le \sigma_n |F|_{s_n}$ with fixed $s$ and $\sigma_n \le A(n!)^D$ for fixed $A$, $D$", and restriction to Schwartz tests flat on the coincidence diagonals gives the test class of that reference. OS II itself was not re-read [CITED].

### 4.2 Hypothesis delivered by the blocks

**Definition 4.1** ((H-U) uniform bound)**.** There are $s \in \mathbb{N}$ and $A, D < \infty$ such that for all $K \ge K_0$, all $n \ge 1$, and all $F \in \mathcal{D}_n$:

$$|S_{n,\varepsilon}(F)| \le \sigma_n \|F\|_{sn}, \qquad \sigma_n := A(n!)^D, \quad \varepsilon = L^{-K}.$$

The derivative order $sn$ grows linearly in $n$. The OS linear growth condition admits this. (H-U) is the output of blocks (B-T2) + (B-S) + the bridge (B-N1) of Section 4.5.

### 4.3 Theorem (finite-torus existence, compactness form)

**Theorem 4.2** (finite-torus existence, compactness form [PROVED MODULO NAMED BLOCKS])**.** *Let $G = SU(2)$, $d = 4$, $L$ odd and large, $\ell \in 2\mathbb{N}$ fixed. Let $\gamma > 0$ be Balaban's small constant and $g \in ]0, \gamma]$ small. For $K \ge K_0$ let $\varepsilon = L^{-K}$ and $g_0 = g_0(\varepsilon, g)$ be a shooting solution (Lemma 3.4) whose effective couplings satisfy $g_k \in ]0, \gamma]$ for $0 \le k \le K$ and $g_K = g$. Assume (H-U). When (H-U) is taken from the blocks (Section 4.2) rather than assumed, the chain also assumes the $\kappa \to \kappa/2$ decay recovery of D2 [R, B22] (not stated, `SOURCES_D2_VERIFY.md` (5)), which does not follow from the other hypotheses. Then:*

*(a) Compactness. Every sequence $K_m \to \infty$ has a subsequence $K_{m_i}$ such that, with $\varepsilon_i = L^{-K_{m_i}}$, the limit $S_n(F) := \lim_i S_{n,\varepsilon_i}(F)$ exists for every $n \ge 1$ and every $F \in \mathcal{D}_n$.*

*(b) For every limit family $(S_n)_{n \ge 1}$ obtained in (a):*

- *(OS0) Regularity. $S_n$ is linear on $\mathcal{D}_n$, and $|S_n(F)| \le A(n!)^D \|F\|_{sn}$. In particular $S_n$ restricted to $C_c^\infty(T^{4n} \setminus \bigcup_{i<j}\{x_i = x_j\})$ is a distribution of order $\le sn$.*
- *(OS1) Invariance. $S_n(F \circ \tau_a^{\times n}) = S_n(F)$ for every $a \in T_\ell^4$ (all continuous translations), and $S_n(F \circ R^{\times n}) = S_n(F)$ for every $R$ in the hypercubic group $B_4$ (signed permutations of the axes, acting about a lattice site).*
- *(OS2) Reflection positivity. Let $\theta(x_0, \vec{x}) = (-x_0, \vec{x})$ on $T_\ell^4$ and $T_+ := \{x : x_0 \in ]0, \ell/2[\}$. For every finite family $F_0 \in \mathbb{C}$, $F_n \in \mathcal{D}_n$ with $\operatorname{supp} F_n \subset T_+^n$ $(1 \le n \le N)$:*

$$\sum_{n,m=0}^{N} S_{n+m}(\Theta F_n \otimes F_m) \ge 0, \quad (\Theta F_n)(x_1, \dots, x_n) := \overline{F_n(\theta x_n, \dots, \theta x_1)}, \quad S_0 := 1.$$

- *(OS3) Symmetry and reality. $S_n(F \circ \pi) = S_n(F)$ for every permutation $\pi$ of the $n$ arguments, and $S_n(F) \in \mathbb{R}$ for real $F$.*

*(c) Nontriviality is a separate statement, Theorem 4.3.*

The sourced blocks use the unit-scale condition (U), $\rho_1 \le c_* r_0/M^a$ at $\lambda_K = 1$ ($r_0$ the Cauchy margin of the constant-source family, `SOURCES_D2C.md` (3.5)), as a constant choice (C23), made once after $r_0$ and before $K$; B-N1 meets it by the rescaling $\lambda_i = 2N_U(f_i)$ at a $C^n$ cost, $s$ and $D$ unchanged (`SOURCES_UN1.md`, `SOURCES_UN1_VERIFY.md`). This (U) is not the Route B hypothesis (U). *What is claimed about measures.* No probability measure is claimed. The limit is a family of Schwinger functions on $\mathcal{D}_n$. A measure on distributions would need finite moments of smeared fields, and $\mathrm{Var}\sum_p f(x_p)\varphi_p \sim \varepsilon^{-4}$ (Section 2.5). (OS0)–(OS3) on $\mathcal{D}_n$ are the inputs of OS reconstruction.

## 4.4 Theorem (nontriviality), conditional on (B-NT)

**Theorem 4.3** (nontriviality [CONDITIONAL ON (NT) OF SECTION 4.5.9])**.** *Under the hypotheses of Theorem 4.2 and (NT), every limit family satisfies: (i) $S_2 \neq 0$: there are $f_1, f_2 \in C^\infty(T^4)$ with disjoint supports and $S_2(f_1 \otimes f_2) \neq 0$. (ii) $S_2$ is not the two-point function of a scale-invariant (in particular free Maxwell) composite: $d^8 S_2(x,y)$, $|x-y| = d$, is not asymptotically constant as $d \to 0$.*

Part (ii) is stated here in the form the proof supports (two-sided envelope only); see Section 4.8, item 3. The source stated (ii) as "behaves like $c\,g(d)^4$ with $g(d)^2 \sim (\beta \log(1/d))^{-1}$", which needs $\beta_j \to \beta_\infty$ and is not proved.

## 4.5 Proof modulo blocks

Structure. (B-T2) gives the trajectory. (B-S) gives sourced bounds uniform in $\varepsilon$. (B-N1) turns them into (H-U). (B-C) gives (a). (B-REG), (B-TI), (B-RP), and (OS3) give (b). (B-NT) gives Theorem 4.3. Only (B-C), (B-REG), (B-TI), (B-RP), and (OS3) are proved in this section; they use nothing beyond (H-U) and exact lattice symmetries.

### 4.5.1 (B-T2) Theorem 2: the running coupling

Needed. The hypothesis of Balaban's Theorem 1 (CMP 122 II p. 355), $g_k \in ]0,\gamma]$ for all $k$, quoted in Section 1.2 [VERIFIED QUOTE]. It was to be discharged by Theorem 2 of CMP 109 p. 259 [8]: for small $g$ there is $g_0(\varepsilon, g)$ with all $g_k \in ]0,\gamma]$, $g_K = g$, and (0.31): $1/g^2 + \beta \log(L^k\varepsilon)^{-1} \le 1/g_k^2 \le 1/g^2 + \beta' \log(L^k\varepsilon)^{-1}$. Balaban: "has not been published yet" [VERIFIED QUOTE].

Status from Section 3 (and §10 of `THEOREM2_PROOF_v4.md`). The weak form, Theorem 2w, is [PROVED+VERIFIED, CONDITIONAL] by Route K (Theorem K + Lemma D″) on (H2) and the fixed-level inputs (PI), (F), ($M_j$), V1, (G2), plus L5 and item 7′ ((H0) at odd $L > 11$, or (G-gain) with the running-maximum lift, verified (`SOURCES_GGAIN_NONMONO_VERIFY`)); item 9 ($K$-independence) is closed. It suffices for Theorem 1 by the cross-scale sweep [PROVED, UNVERIFIED]. Literal (0.31) is not obtained (it needs eventual positivity of $\beta_j(0)$). Route B (Theorem B of `BETA_FP_PW`) is a formally distinct second route; its logic is [PROVED+VERIFIED, CONDITIONAL], its inputs (U), (DF), (T1) are [PROVED, UNVERIFIED] and rest on (H2). The reduction pieces are Lemma D′, Lemma D″, and Lemmas D, E (the one-argument forms in Lean).

Conclusion. At $L = 13$, (H2) is certified (Theorem 5.1); (B-T2) remains conditional on the fixed-level inputs, L5, and item 7′; see the dependency table (Section 4.6).

### 4.5.2 (B-S) Sourced bounds E14-a..g

Needed. A version of Balaban's Theorem 1 with the (C1) source, i.e. with the local bare coupling $w_p = 1/g_0^2 - f(x_p)$, $f$ real, $|\partial^m f| \le r$ for $m \le 5$ (class $F_r$), uniformly in $\varepsilon$. Output form (`SOURCES.md` (N1)): $\log Z_\varepsilon[f] - W_\varepsilon[f]$ is $C^\infty$ in source amplitudes $z$ with $|\partial_z^\alpha(\log Z_\varepsilon[f] - W_\varepsilon[f])| \le C^{|\alpha|}\alpha!\, r^{-|\alpha|}$

for $f = \sum_i z_i f_i$ with disjoint supports. $W_\varepsilon[f]$ is a local field-independent contact functional (allowed to diverge; it contributes zero to every mixed derivative of order $\geq 2$ for disjoint supports, `SOURCES_VERIFY` item 5(a)).

Estimates needed (`SOURCES.md` §2.5): (E14-a) sourced small-field step; (E14-b) sourced large-field step and R operation; (E14-c) derivative bounds in $f$. Not needed: (E14-d) fixed-torus convergence (= E13); (E14-g) tensor sources. (E14-e) RP and symmetry transfer is done in this section (Sections 4.5.6, 4.5.7). (E14-f) nontriviality at a fixed layer is needed only for Theorem 4.3. The current status of each sub-block is in the dependency table (Section 4.6); (B-S) remains a [HYPOTHESIS].

### 4.5.3 (B-N1) Bridge: source-form bounds ⇒ (H-U) [OPEN]

`SOURCES_VERIFY` item 5(b) [VERIFIED QUOTE]: the source form "does not give $|S_n(F)| \leq C^n n! \|F\|_{s,n}$ for $F$ flat on diagonals with a fixed $s$ ... Mark N1 for general $F$ as a gap." This paper does not close the gap. It gives a route [DERIVED SKETCH], which shows that the gap is removed by the OS index growth $sn$.

1. Moments from cumulants. $\langle \prod_i \phi(f_i) \rangle = \sum_{\text{partitions}} \prod_{\text{blocks}} \kappa(\cdot)$. Singleton cumulants vanish ($\varphi$ is centred). For disjoint supports the source form bounds every cumulant by $C^{|B|}|B|!\, r^{-|B|}$, so the moment by $C^n n!\, r^{-n}$.

2. Scale-$\delta$ tests. A bump of side $\delta \leq 1$ and amplitude $a$ has $|\partial^5 f| \sim a\delta^{-5}$; it lies in $F_r$ if $a \leq cr\delta^5$. So $n$ disjoint unit-amplitude scale-$\delta$ bumps have $|\text{moment}| \leq C^n n!\, \delta^{-5n}$. This is lossy but polynomial in $\delta^{-1}$.

3. Whitney decomposition. Split $T^{4n}$ by a smooth dyadic partition in $\rho(x) := \min_{i<j} |x_i - x_j| \in [\delta, 2\delta]$, $\delta = 2^{-m} \geq \varepsilon$. On each piece cover by cubes of side $\delta/100$; cube tuples are pairwise disjoint. Expand $F\cdot$(cutoffs) in a Fourier series on each cube tuple; the coefficients decay faster than any power; each term is a product of disjointly supported scale-$\delta$ bumps. Flatness gives $|\partial^\beta F(x)| \leq C_k \|F\|_{k+|\beta|} \rho(x)^k / k!$. The count is $\leq C^n \delta^{-4n}$ cube tuples. Derivatives of the cutoffs cost at most $(Cn^2/\delta)^M$; with $M = O(n)$ this is $\leq C^n (n!)^c \delta^{-M}$.

4. Bound per dyadic scale: $\leq C^n (n!)^{D'} \delta^{k-9n-M} \|F\|_{k+M}$. Choose $k = sn - M$ with $s$ large enough ($s \geq 20$ suffices on this count) so the $\delta$-exponent is $\geq n$; the sum over dyadic $\delta \in [\varepsilon, 1]$ converges uniformly in $\varepsilon$.

5. Sub-lattice region $\rho(x) \lesssim \varepsilon$: $\leq C^n \varepsilon^{-4n}$ plaquette tuples, each moment $\leq 2^n$, and $|F| \leq \|F\|_k \varepsilon^k$; this is $\leq C^n \|F\|_{sn}$ for $k = sn \geq 4n$.

Result (if the steps hold): (H-U) with some fixed $s$ and $D$. To check: uniformity of the source bound in the position of the bumps and in $\varepsilon$; the $C^\infty$ real-source form of (E14-c) with $n$-th derivatives for all $n$; the cutoff-derivative count in step 3. The verifier lists further gaps: complex Fourier modes against real sources (a factor $2^n$), the extra derivatives of Fourier summation (absorbed in $s$), and the smooth min-distance partition. Later work (`SOURCES_D4D5`, graded in `SOURCES_D4D5_VERIFY`) confirms "($S_{src}$) ⇒ (H-U) with $s = 6$, $D > 1$" given (FX) (see Section 4.6). `ASSEMBLY_N1.md` closes the three gaps given (S) (confirmed in `SOURCES_HHIST_N1_VERIFY.md`) and, since 2026-10-28, uses the verified form: $C^1$ norm $\max(\|f\|_\infty/r, \|\nabla f\|_\infty/\rho_1)$, order $(n!)^{1+\eta}$, $s = 6$, $D > 1$ ($D > 2$ under the fallback order $2 + \eta$); the $\partial^5$, $\delta^{-5n}$, and $s \geq 20$ counts in the sketch above are superseded.

### 4.5.4 (B-C) Compactness [PROVED HERE, GIVEN (H-U)]

**Proposition 4.4.** *Under (H-U), every sequence $\varepsilon_m \to 0$ has a subsequence along which $S_{n,\varepsilon}(F)$ converges for every $n$ and every $F \in \mathcal{D}_n$. The limit $S_n$ is linear and satisfies $|S_n(F)| \le \sigma_n \|F\|_{sn}$.*

*Proof.* 1. Separability. $C^{sn}(T^{4n})$ with $\|\cdot\|_{sn}$ is separable (trigonometric polynomials with rational coefficients are dense; Fejér means converge in $C^k$). A subset of a separable metric space is separable. So $\mathcal{D}_n$ has a countable subset $Q_n$ dense for $\|\cdot\|_{sn}$. 2. Diagonal extraction. $Q := \bigcup_n Q_n \times \{n\}$ is countable; enumerate it as $(F^{(1)}, n_1), (F^{(2)}, n_2), \dots$. Each sequence $S_{n_t,\varepsilon_m}(F^{(t)})$ is bounded by $\sigma_{n_t}\|F^{(t)}\|_{sn_t}$ (H-U), so it has a convergent subsequence. Extract successively and take the diagonal subsequence $\varepsilon_i$. 3. Extension. Let $F \in \mathcal{D}_n$, $\eta > 0$, and $G \in Q_n$ with $\|F - G\|_{sn} \le \eta$. By (H-U) and linearity, $|S_{n,\varepsilon_i}(F) - S_{n,\varepsilon_j}(F)| \le 2\sigma_n\eta + |S_{n,\varepsilon_i}(G) - S_{n,\varepsilon_j}(G)|$. The last term $\to 0$. So the sequence is Cauchy; call the limit $S_n(F)$. 4. Linearity passes to pointwise limits, and $|S_n(F)| = \lim |S_{n,\varepsilon_i}(F)| \le \sigma_n\|F\|_{sn}$. □

Remarks. (i) The same subsequence serves all $n$; this matters for (OS2). (ii) Nothing here needs E13. (iii) The bounded normalizations $B_K$ and $c_\varepsilon$ can be included in the extraction.

### 4.5.5 (B-REG) Regularity (OS0) [PROVED HERE, GIVEN (H-U)]

By step 4 of the proof of Proposition 4.4 the limit satisfies $|S_n(F)| \le \sigma_n\|F\|_{sn}$, $\sigma_n = A(n!)^D$. On the compact torus the Schwartz seminorm is replaced by $\|F\|_p$ (the weights control decay at infinity on $\mathbb{R}^{dn}$; the torus has none). This is the quoted OS II form. Distribution property: for $F \in C_c^\infty$ of the off-diagonal open set, $F \in \mathcal{D}_n$ and the bound is continuity in the $C^{sn}$ topology, so $S_n$ is a distribution of order $\le sn$ there.

Caveat. OS II is formulated on $\mathbb{R}^{dn}$. On $T^{4n}$ the axioms are used in their torus form (periodic test functions). This is a change of setting, not a consequence of OS II.

### 4.5.6 (B-TI) Translation and hypercubic invariance (OS1) [PROVED HERE, GIVEN (H-U)]

*Lattice, exact.* Put the origin at a lattice site.

- Translations $a \in \varepsilon\mathbb{Z}^4$. The map $U \mapsto U^a$, $U^a(b) := U(b-a)$, permutes the links, preserves the product Haar measure and the Wilson action, and $P_p(U^a) = P_{p-a}(U)$. So $\langle \prod_i \varphi_{p_i+a}\rangle_\varepsilon = \langle \prod_i \varphi_{p_i}\rangle_\varepsilon$ and $S_{n,\varepsilon}(F \circ \tau_a^{\times n}) = S_{n,\varepsilon}(F)$. The coupling $g_0(\varepsilon)$ is one number, so nothing in $\mu_\varepsilon$ depends on position.

- Hypercubic $R \in B_4$ (signed permutation matrices, $|B_4| = 384$). $R$ maps sites to sites, links to links (possibly reversed), plaquettes to plaquettes, and $x_{Rp} = Rx_p$. Define $(RU)(b) := U(R^{-1}b)$ with $U(-b) = U(b)^{-1}$. Haar measure is invariant under $U \mapsto U^{-1}$ and $\operatorname{Re}\operatorname{tr} W^{-1} = \operatorname{Re}\operatorname{tr} W$ for $W \in SU(2)$, so $A$ and the measure are invariant and $P_{Rp}(RU) = P_p(U)$. Hence $S_{n,\varepsilon}(F \circ R^{\times n}) = S_{n,\varepsilon}(F)$.

- These identities hold at every $\varepsilon$. $F \circ \tau_a^{\times n}$ and $F \circ R^{\times n}$ are again in $\mathcal{D}_n$. They pass to the limit along the subsequence of Proposition 4.4: $S_n(F \circ R^{\times n}) = S_n(F)$, and $S_n(F \circ \tau_a^{\times n}) = S_n(F)$ for $a \in \varepsilon_i\mathbb{Z}^4$ for all large $i$.

*Continuous translations.* Let $a \in T_\ell^4$. Pick $a_i \in \varepsilon_i\mathbb{Z}^4$ with $|a - a_i| \le \varepsilon_i$ (componentwise rounding gives $\le \varepsilon_i\sqrt{4}/2 = \varepsilon_i$). Write $F_a := F \circ \tau_a^{\times n}$. The mean value theorem along the diagonal direction gives $\|F_a - F_{a_i}\|_{sn} \le 4n|a - a_i|\,\|F\|_{sn+1}$. Then

$$|S_n(F_a) - S_n(F)| \le |S_n(F_a) - S_{n,\varepsilon_i}(F_a)| + |S_{n,\varepsilon_i}(F_a) - S_{n,\varepsilon_i}(F_{a_i})| + |S_{n,\varepsilon_i}(F_{a_i}) - S_{n,\varepsilon_i}(F)| + |S_{n,\varepsilon_i}(F) - S_n(F)|.$$

The first and last terms $\to 0$ by Proposition 4.4. The third is 0 by lattice invariance. The second is $\le \sigma_n \cdot 4n\,\varepsilon_i \|F\|_{sn+1}$ by (H-U) and linearity. Let $i \to \infty$. $F$ is smooth, so $\|F\|_{sn+1} < \infty$. Rotations outside $B_4$ are not available (Section 4.7).

### 4.5.7 (B-RP) Reflection positivity (OS2)

*Lattice theorem* [CITED]: Osterwalder–Seiler [21], Ann. Phys. 110 (1978) 440–471. The paper is not in the project folder; no verbatim quote. Content as usually stated: for the Wilson action with compact gauge group and any $\beta \ge 0$, and $\theta$ the reflection in a lattice hyperplane that bisects the links orthogonal to it, $\langle \Theta(\Phi)\Phi \rangle \ge 0$ for every function $\Phi$ of the link variables in the positive half-lattice, $(\Theta\Phi)(U) := \overline{\Phi(\theta U)}$. Jaffe–Witten [15] line 546 [VERIFIED QUOTE]: "Reflection positivity holds for the Wilson approximation [36], a major advantage". The template uses the same input (`o3_continuum.txt` line 4902) [VERIFIED QUOTE]: "At every cutoff the nearest-neighbour measure is reflection positive across microscopic site and link seams." Proof idea (standard): fix the crossing links to 1 by a gauge transformation; the action splits as $A_+ + \theta A_+ + \sum_{\text{crossing}} \operatorname{Re}\operatorname{tr}(X_p \theta(X_p)^*)$, and $\exp(\beta \operatorname{Re}\operatorname{tr}(XY^*))$ is a positive-type kernel (positive combination of characters), which gives a sum of squares. On the periodic lattice there are two fixed planes, $x_0 = \varepsilon/2$ and $x_0 = \varepsilon/2 + \ell/2$; both are link planes because $N_\varepsilon$ is even ($\ell$ even, Section 2.5). The periodic two-seam version is standard [CITED]; its text was not checked. For site-plane reflections see Menotti–Pelissetto [17] (Section 4.8, item 1).

*Passage to the limit* [PROVED HERE, GIVEN (H-U) AND THE LATTICE THEOREM]. Setup. $\theta_\varepsilon(x) := \theta(x) + \varepsilon e_0 = \theta(x - \varepsilon e_0)$, fixed planes $x_0 = \varepsilon/2,\ \varepsilon/2 + \ell/2$. The positive half $\Lambda_+$ consists of links with both ends in $\{x_0 \in \{\varepsilon, 2\varepsilon, \dots, \ell/2\}\}$. Fix $F_0 \in \mathbb{C}$ and $F_n \in \mathcal{D}_n$, $\operatorname{supp} F_n \subset T^n_+$, $1 \le n \le N$. Supports are compact in the open set, so there is $\delta > 0$ with $x_0 \in [\delta, \ell/2 - \delta]$ on all supports.

Step 1 (lattice form). For $\varepsilon < \delta/2$ every plaquette with centre in a support has all four links in $\Lambda_+$. Define $\Phi := F_0 + \sum_{m=1}^N \sum_{p_1 \dots p_m} F_m(x_{p_1}, \dots, x_{p_m}) \prod_i \varphi_{p_i}$. $\Phi$ depends only on $\Lambda_+$ links. Since $\varphi_p$ is real and $P_p(\theta_\varepsilon U) = P_{\theta_\varepsilon p}(U)$, re-indexing gives $\Theta\Phi = \bar{F}_0 + \sum_n \sum_q \overline{F_n}(\theta_\varepsilon x_{q_1}, \dots, \theta_\varepsilon x_{q_n}) \prod \varphi_{q_i}$. Hence

$$0 \le \langle \Theta\Phi\, \Phi \rangle_\varepsilon = \sum_{n,m=0}^{N} S_{n+m,\varepsilon}(\Theta_\varepsilon F_n \otimes F_m), \quad \Theta_\varepsilon F_n := (\Theta F_n) \circ \tau^{\times n}_{-\varepsilon e_0},\ S_{0,\varepsilon} := 1.$$

(Argument order inside $\Theta F_n$ is immaterial by (OS3).) Each $\Theta_\varepsilon F_n \otimes F_m$ lies in $\mathcal{D}_{n+m}$: its two factors have supports at positive distance and each factor is flat on its own diagonals.

Step 2 (seam shift). $\|\Theta_\varepsilon F_n \otimes F_m - \Theta F_n \otimes F_m\|_{s(n+m)} \le 2^{s(n+m)} \cdot 4n\,\varepsilon\, \|F_n\|_{s(n+m)+1} \|F_m\|_{s(n+m)}$. By (H-U) the difference of the $S_{n+m,\varepsilon}$ values is $\le \sigma_{n+m}$ times this, which $\to 0$.

Step 3 (limit). Along the subsequence of Proposition 4.4, $S_{n+m,\varepsilon_i}(\Theta F_n \otimes F_m) \to S_{n+m}(\Theta F_n \otimes F_m)$ for each of the finitely many $(n, m)$. So $\sum_{n,m} S_{n+m}(\Theta F_n \otimes F_m) = \lim_i \sum_{n,m} S_{n+m,\varepsilon_i}(\Theta_{\varepsilon_i} F_n \otimes F_m) \ge 0$.

**Lemma 4.5** (positivity of a finite form survives pointwise limits [PROVED HERE])**.** *Let $a^{(i)} \in \mathbb{C}^{(N+1)\times(N+1)}$ with $\sum_{n,m} \bar{c}_n c_m a^{(i)}_{nm} \ge 0$ for all $c$ and all $i$, and $a^{(i)}_{nm} \to a_{nm}$ for every $(n, m)$. Then $\sum \bar{c}_n c_m a_{nm} \ge 0$ for all $c$.*

*Proof.* For fixed $c$ the left side is a finite linear combination of the entries, so it converges to $\sum \bar{c}_n c_m a_{nm}$; $[0, \infty)$ is closed. □

Step 3 is this lemma with $a^{(i)}_{nm} := S_{n+m,\varepsilon_i}(\Theta_{\varepsilon_i} F_n \otimes F_m)$ (replace $F_n$ by $c_n F_n$), plus Step 2.

#### 4.5.8 (OS3) Symmetry and reality [PROVED HERE]

$S_{n,\varepsilon}(F \circ \pi) = S_{n,\varepsilon}(F)$: the $\varphi_p$ are commuting real random variables; re-index the sum. For real $F$, $S_{n,\varepsilon}(F) \in \mathbb{R}$. Both identities are linear and pass to the limit. $S_0 = 1$ by definition.

#### 4.5.9 (B-NT) Nontriviality: what is needed

- A two-point function alone never excludes a Gaussian: every positive-definite $S_2$ is the covariance of some Gaussian generalized field.
- The plaquette density is a composite ($\approx \frac{1}{4} \operatorname{tr} F^2 \times \varepsilon^4$). Even in the free theory its connected $n \geq 3$ functions are nonzero (Wick squares). So $S_n^c \neq 0$ for some $n \geq 3$ does not prove interaction.
- The meaningful test excludes (a) the zero theory and (b) the free Maxwell composite. The free composite has, at short distance, the scale-invariant form $c_0 d^{-8}$. Asymptotic freedom predicts $c\, g(d)^4 d^{-8}$ with a running coupling.

*Free-theory prediction along this sequence.* At tree level with the bare coupling, $\langle \varphi_p \varphi_{p'} \rangle \approx c\, \varepsilon^8 g_0^4 |x - y|^{-8}$. Since $g_0(\varepsilon)^2 \leq (1/g^2 + \beta \log \varepsilon^{-1})^{-1} \to 0$ by (0.31), the bare-coupling prediction for $S_2$ at fixed separation tends to 0. The claim is that the true $S_2$ does not.

*Statement (NT) needed* [HYPOTHESIS]; owned by (E14-f), a corollary of (E14-a) at a fixed layer. There are $c > 0$ and $C$ such that for all $K$, all $f_1, f_2 \in C^\infty(T^4)$ with disjoint supports at distance $\geq d$, $d \geq L\varepsilon$, and $k(d)$ the layer with $L^k \varepsilon \leq d < L^{k+1}\varepsilon$:

$$S_{2,\varepsilon}(f_1 \otimes f_2) = c \iint f_1(x) f_2(y)\, g^4_{k(|x-y|)} |x - y|^{-8}\, dx\, dy\, (1 + R), \quad |R| \leq C(g^2_{k(d)} + (d/\ell)^2).$$

$c$ is the one-loop (tree-level in the composite) coefficient: $c \propto \sum (\text{free field-strength covariance})^2$, a sum of squares, so $c > 0$ [DERIVED, STANDARD]; its value must be computed in Balaban's normalization.

*Derivation of Theorem 4.3 from (NT) and (B-T2)* [PROVED HERE, GIVEN (NT)]. (i) Take $f_1, f_2 \geq 0$, not identically 0, with supports at distance $\geq d_0$. Corrected condition (Section 4.8, item 3): the support diameters are $\ll \ell$, so that $g^2 + (|x-y|/\ell)^2$ is small on the whole support pair and $|R| \leq \frac{1}{2}$. By (0.31) at layer $k(d)$ with $d \in [d_0, \ell]$: $1/g_k^2 \leq 1/g^2 + \beta' \log(L^k \varepsilon)^{-1} \leq 1/g^2 + \beta' \log(L/d_0)$, uniformly in $\varepsilon$. So $S_{2,\varepsilon}(f_1 \otimes f_2) \geq \frac{1}{2} c (1/g^2 + \beta' \log(L/d_0))^{-2} \iint f_1 f_2 |x - y|^{-8} > 0$ uniformly in $\varepsilon$, and the limit inherits the bound. (ii) For bumps concentrated at separation $d \to 0$ (with $d$ a fixed multiple of their width), $d^8 S_2 \leq (1 + C g^2_{k(d)})\, c\, (1/g^2 + \beta \log(1/d))^{-2} \to 0$ as $d \to 0$ by the lower bound in (0.31), while $d^8 S_2 > 0$ by (i). A scale-invariant composite would give a nonzero constant limit. So the limit is neither zero nor the free composite. Only the two-sided bound (0.31) is used. Caveat. (ii) needs (NT) uniformly down to $d \geq L\varepsilon$ with the remainder controlled by $g^2_{k(d)}$, i.e. the fixed-layer expansion at every layer. If (E14-f) is proved only at a fixed layer $j$, (i) holds, and (ii) holds only in the weaker form "$d^8 S_2(d)$ is bounded between the two (0.31) envelopes for $d \in [L^{-j}, 1]$".

### 4.6 Dependency table

The table gives the status of every block of Theorem B and of every named hypothesis of Theorem A. It follows the state recorded in `PLAN.md` (through the entry `SOURCES_MA2_VERIFY.md`, 2026-10-28) and §10 of `THEOREM2_PROOF_v4.md`. Global conditions carried by every row: (GC1) Balaban's

published proofs CMP 95–122 (1984–89) are correct as published; nothing in this paper re-checks them line by line. (GC2) Balaban's inductive assumptions (CMP 109 pp. 263–264) hold with the stated uniformity. (GC3) $L$ odd, $L > 11$, $M$ large, $\gamma$ small as Balaban requires; $\ell$ even (sufficient, Section 2.5). Rows "item $n$" are the named hypotheses of Section 3; $n$ is their item in Section 6. Their file columns use the short names of `THEOREM2_PROOF_v2_VERIFY` §K (KAPPA = `KAPPA_BOUNDED`, H_VERIFY = `BETA_FP_H_VERIFY`, PW_VERIFY = `BETA_FP_PW_VERIFY`, v1 verifier = `THEOREM2_PROOF_VERIFY`).

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-T2 | Theorem 2w (Theorem 3.1): trajectory $g_0(\varepsilon, g)$, (b), (c′), (S) | [PROVED+VERIFIED, CONDITIONAL] (Route K = Theorem K + Lemma D″). Literal (0.31) not obtained | `THEOREM2_PROOF_v4.md`, `KAPPA_BOUNDED.md` | (H2), (PI), (F), $(M_j)$, V1, (G2), L5, item 7′ (from (H0) or (G-gain), verified (`SOURCES_GGAIN_NONMONO_VERIFY`)) |
| B-T2.a | Lemmas D, D′, D″, E (shooting) | [PROVED+VERIFIED; LEAN] (D′, D, E, D″ one-argument form; history form open) | `LEMMA_DPRIME.md`, `lean/` | continuity of $\beta_{k+1}$ in history |
| B-T2.b | Sweep: (c′)+(S) ⇒ Balaban's Theorem 1 inputs (Section 3.4.3; item 10) | [PROVED, UNVERIFIED]; 4 of 19 rows verified (LEMMA_DPRIME_VERIFY Steps 4–5, rows 11, 12, 16, 18) | `LEMMA_DPRIME_SWEEP.md` | Balaban readings (unchecked list) |
| B-T2.1 | U3 Thm 4.1 (Gaussian RG, rate $L^{-2j}$) + O5 | [PROVED+VERIFIED] | `U3_GAUSSIAN_RG.md`, `U3_REPAIRS.md` | trivial background |
| B-T2.2 | U3 Lemma 5.1; (A), (W) | neutral sector [PROVED+VERIFIED]. (W) numerically exact; (A) confirmed qualitatively and quantitatively (`U3_AW_VERIFY.md`); (HS) [REFUTED] at $L = 3$. Not on the path of Theorem 2w after (H2); needed for the convergence route only | `U3_LEMMA51.md`, `U3_AW.md` | (A) in full, (N) (plausibly false) |
| B-T2.3 | O6 bordered-matrix invertibility | one-step [PROVED+VERIFIED]; composed constraint reduced to one-step O6 + (A) | `U3_REPAIRS.md` | — |
| B-T2.4 | $c = 11N^2/(12\pi^2)$, anchor-free identity Prop. 2.1, telescoping | [PROVED+VERIFIED]. $\beta_\infty = c \log L$ itself replaced by window sums (Theorem K, Route B) | `BETA_FP_LEMMA43.md`, `BETA_FP_LEMMA43_VERIFY.md` | (F) [CITED, SOURCE CHECKED] |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-T2.5 | $\kappa_k$ bounded (Theorem K) | [PROVED+VERIFIED, CONDITIONAL ON (B2)]; (B2) [PROVED+VERIFIED, CONDITIONAL ON (H2)] | `KAPPA_BOUNDED.md`, `KAPPA_BOUNDED_VERIFY.md`, `B2_TWOSTEP.md`, `B2_TWOSTEP_VERIFY.md` | (H2), (PI), $(M_j)$ |
| B-T2.H2 | (H2) uniform two-step inequality $\rho_2 < L$ (Section 3.7.3; item H) | [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)] at $L = 13$: $\rho_2 \le \sup \rho_1^2 \le 10 < 13$ (Theorem 5.1); residue: reading of (0.4), gauge choice, Prop. A at $M \ge 27$, Lemma G beyond point checks, `bb_arb.py` audited only. At $L = 3, 5, 9$ [NUMERICAL]; point enclosure $[1.9993, 2.1273] < 3$ at $L = 3$ [INTERVAL ENCLOSURE, VERIFIED] | `H1_ANALYTIC.md`, `H1_ANALYTIC_VERIFY.md`, `H1_ANALYTIC_VERIFY2.md`, `H2_CERTIFY_P5_VERIFY.md`, `H2_LARGE_L_VERIFY.md`, `H2_CERTIFY_VERIFY.md`; B2_TWOSTEP §2 | — (box sweep at $L = 13$ running as a cross-check) |
| B-T2.H1 | one-step route at $L = 13$: $\sup \rho_1^2 < L \Rightarrow$ (H2) | [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]: $\sup_{j,K} \rho_1^2 \le 9.9995 < 13$ by the majorant $B_{13}$ and `bb_arb.py`; factorization [PROVED+VERIFIED; LEAN] (`rho2_le`); majorant core [LEAN] (`Tower.lean`) | `H1_ANALYTIC.md`, `H1_ANALYTIC_VERIFY.md`, `H1_ANALYTIC_VERIFY2.md`, `H2_CERTIFY_VERIFY.md` | — |
| B-T2.B | Route B (Theorem $B_{PW}$; item B8) | logic [PROVED+VERIFIED, CONDITIONAL] (tag decision, `PLAN.md`; PW_VERIFY §6 items 1–4); inputs (U), (DF), (T1) [PROVED, UNVERIFIED] | `BETA_FP_PW.md`, `BETA_FP_PW_VERIFY.md`, `BETA_FP_U.md` | (H2), $(B\theta)$, A6, LW (5.7), (H0) |
| item 1 | (B2) / (B2w) (Section 3.7) | open; reduced to (H2) | KAPPA §2, Rem. 2.3; KAPPA_VERIFY §7–§9 | (H2) |
| item 1 | (H2) $\Rightarrow$ $(C1\sigma)$ $\Rightarrow$ (B2) (Theorems 3.10, 3.11) | proved+verified, conditional on (H2) | B2_TWOSTEP §2–§3; B2_TWOSTEP_VERIFY grades 1, 3, 4 | (H2) |
| item 1 | $(C1\sigma)$ + (h) $\Rightarrow$ (B2) | implication verified; $(C1\sigma) \Leftarrow$ (H2); (h) numerical $L = 3, 5$ | KAPPA Prop. 4.1; KAPPA_VERIFY §7 | — |
| item 1 | (H″) $\Rightarrow$ (A) $\Rightarrow$ (B2) | implication argued; not needed after (H2) | U3_AW §4.7; no verifier (U3_AW_VERIFY grade 5 covers (H′) only) | — |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| — | (HS) (Section 3.7.2) | refuted $L = 3$ | U3_AW (H′); U3_AW_VERIFY §6 | — |
| item H | $\rho_1^2 < L \Rightarrow$ (H2) (one step, $L \geq 5$) | arithmetic; $\rho_1^2/L = 0.771$ at $L = 5$, numerical | H2_LARGE_L §3; H2_LARGE_L_VERIFY grade 4 (measurement) | — |
| item H | certification tail $j_0$ | derivation confirmed; $j_0 \approx 8$ ($L = 3$), $\approx 6$ ($L = 5$), corrected | H2_LARGE_L §4.1; H2_LARGE_L_VERIFY grade 6 | — |
| item 2 | (PI) | argued | BETA_FP_H §3.1; H_VERIFY §5 | — |
| item 3 | (F) | cited, source checked | BETA_FP_LEMMA43; H_VERIFY §4 | — |
| item 4 | ($M_j$): (S1)–(S3), Prop. 4.4 | open / argued | BETA_FP_M §4 (no verifier); H_VERIFY §5 refuted the old step | (S1)–(S3) |
| item 5 | V1 | open | R2; U3_VERIFY C4 (order $p^2$) | — |
| item 6 | (G2) | formal | BETA_FP_LEMMA43 | — |
| item 7′ | $\lvert\beta_{k+1}(h,g) - \beta_{k+1}(0)\rvert \leq C_\beta g$ on every past (was (H0), uniform in $j$ and history) | from (H0) [Balaban], odd $L > 11$, or from (G-gain) with the running-maximum lift [proved+verified] | CMP 109 (1.22); `SOURCES_GGAIN_NONMONO.md`; KUNIF_VERIFY item 3 | (G-gain) inputs if taken from it |
| item 8 | L5 history form and domain (Section 3.4.5) | open | LEMMA_DPRIME Rem. 3; LEMMAS_DE_VERIFY §2.1; v2_VERIFY §F | — |
| item 9 | $\beta_j(g)$ independent of $K$ | closed as a reduction [verified]: volume statement, $\beta_j$ whole-lattice by definition, D′/D″ at fixed $K$; residual volume question = ledger B8, an input of Theorem 1 | `SOURCES_T2_KUNIF.md`, `SOURCES_T2_KUNIF_VERIFY.md` | (K1), (K2) = item 7′, (K3) |
| item 10 | BALABAN_CHECK item 11 readings | unchecked | BALABAN_EXCERPTS | — |
| item 11 | odd $L$; $L > 11$ in Balaban | input | CMP 109 pp. 251–252; v1 verifier Step 6 | — |
| item 12 | SU(2) | scope | — | — |
| item 13 | Lean history form | open | — | — |
| B1, B2′ | Route B: (H2) + (T1) ⇒ (U) (Theorem 2.4 of BETA_FP_U) | proved, unverified | BETA_FP_U §2.4; no verifier | (H2), (T1) |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B1, B2′ | Route B: (U) | hypothesis; $\Leftarrow$ (T1) + (H2) (BETA_FP_U); proved, unverified | BETA_FP_PW §1.2; PW_VERIFY §3 | (T1), (H2) |
| B1 | Route B: (T1) | argued | BETA_FP_U §2.3; no verifier | — |
| B3 | Route B: $(B\theta) = \gamma_{rec} + \mathfrak{d}_F$ | $\gamma_{rec}$ bounded given Thm 2.4 (proved, unverified); (DF) argued | BETA_FP_PW §3.3; BETA_FP_U §3; PW_VERIFY §6 (logic of Theorem $B_{PW}$); none for BETA_FP_U | (DF) |
| B4 | Route B: LW (5.7) | cited | cited; PW_VERIFY §3 (regime correct) | — |
| B5 | Route B: A6 ($\beta$ identification, uniform in $k$) | argued | SOURCES_D1_VERIFY 8; PW_VERIFY §3; BETA_FP_U §4.2 | — |
| B7 | Route B: $\theta_* < 1$ | trivial condition | PW_VERIFY §3 | — |
| B-T2.6 | certified early steps $\beta_j(0) > 0$ | needed only for literal (0.31) at every $K$, not for Theorem 1; [NUMERICAL], no certificate; $\beta_1(0)$ computed to $L = 11$ | `numerics/NUMERICS.md`, `UMK_BETA1.md` | $L > 11$ is Balaban's range |
| B-S | sourced Theorem 1 (E14-a,b,c) | [OPEN] | `SOURCES.md` §2.5 | B-T2 for the sourced trajectory |
| B-S.D1 | one sourced step, Gaussian level | [PROVED+VERIFIED] (with corrections) | `SOURCES_D1.md`, `SOURCES_D1_VERIFY.md` | — |
| B-S.D2 | sourced small-field step (E14-a) | (a), (b), (d), (f) confirmed; (c) rejected; (e) reduced | `SOURCES_D2.md`, `SOURCES_D2_VERIFY.md` | analyticity in input activities; $\kappa \to \kappa/2$ [R, verified, B22; uses A23]; (U) [C23, constant choice] |
| B-S.D2C | redo of (c): frozen source per domain, Lemma A, $h \equiv 0$ | Lemma A and recursion confirmed; (I) corrected (exact only with adapted gauge-fixing weight) | `SOURCES_D2C.md`, `SOURCES_D2C_VERIFY.md` | domain hypothesis (D); (H$\beta$) |
| B-S.H$\beta$ | two-loop bound $\sup \sum_j \lvert B'_j \rvert \le \frac{1}{2}$ | [PROVED+VERIFIED, CONDITIONAL] on (I), (H-hist) | `SOURCES_HBETA.md`, `SOURCES_HBETA_VERIFY.md` | (I), (H-hist) |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-S.hist | (H-hist) history sensitivity | [VERIFIED REDUCTION] on (W4), Facts A/B, (I)/L3 and (G-gain) in both slots (SOURCES_SM.md; verifier: items 1–4 confirmed, 5–6 corrected); the (G-gain) chain continues in row B-S.Ggain (G3, (Q)); with the large-field half (SOURCES_GGAIN_LF_VERIFY) the real uniform (H-hist) is a [VERIFIED REDUCTION] in both halves (ledger B13); Lemma HH: SOURCES_HHIST.md, SOURCES_HHIST_N1_VERIFY.md | SOURCES_SM.md, SOURCES_SM_VERIFY.md | (G-gain), small and large field |
| B-S.Ggain | (G-gain): $g_k\lvert B\rvert$ gain in the small-field step | pointwise Cauchy and uniform $C_\beta$ on $]0,\gamma]$ [PROVED+VERIFIED] (small-field, incl. vacuum subtraction); Lemma G4 off-chain (SOURCES_G4.md); $\chi_k$ surface term = Lemma G3, proved given (Q) [PROVED+VERIFIED, CONDITIONAL] (SOURCES_G3_VERIFY); (Q) reduced to CMP 99 (3.132), ($E_U$), and the localization lemma LOC [VERIFIED REDUCTION] (SOURCES_Q_VERIFY, SOURCES_CDC_VERIFY, SOURCES_EU_VERIFY, SOURCES_LOC_VERIFY); large field [VERIFIED REDUCTION] (one Cauchy step on the large-field disc, SOURCES_GGAIN_LF_VERIFY); non-monotone pasts: running-maximum lift (SOURCES_GGAIN_NONMONO.md, verified (SOURCES_GGAIN_NONMONO_VERIFY)) | SOURCES_GGAIN.md, SOURCES_GGAIN_VERIFY.md, SOURCES_GGAIN_LF.md | (Q)/$c_-$, ($M_r$), CMP 102 (189), (M-B) bookkeeping; not at $g = 0$ |
| B-S.G3 | Lemma G3: $\chi_k$ surface term, uniform in $g$ | [PROVED+VERIFIED, CONDITIONAL] on (Q) (Route 2 surface estimate; the B/B′ cut closed in SOURCES_Q) | SOURCES_G3.md, SOURCES_G3_VERIFY.md | (Q) |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-S.Q | (Q): covariance lower bound $c_- > 0$ | [VERIFIED REDUCTION] to CMP 99 (3.132) (printed; ledger A1, kernel-decay form now a verified reduction via Lemma CT to CMP 99 Thms 3.1, 3.3, 3.12, `SOURCES_S2_FORM_VERIFY`), ($\mathrm{E}_U$), and LOC; Woodbury identity confirmed (CDC); free-case constants [NUMERICAL] (`numerics/kappa_d.py`) | `SOURCES_Q.md`, `SOURCES_Q_VERIFY.md`, `SOURCES_CDC.md`, `SOURCES_CDC_VERIFY.md`, `numerics/KAPPA_D.md` | A1 residue (torus-side hypothesis, conventions, $G_1$ P-sketch); two [I] bridges |
| B-S.EU | ($\mathrm{E}_U$): uniform block extension | $K_0(4) = 195.19$ ($N = \infty$) [NUMERICAL, REPRODUCED]; $K \le K_0(1 + c_{2/3}(M\alpha_0)^{2/3} + c_1 M\alpha_0)$ [DERIVED SKETCH]; ($\mathrm{E}_U$) [VERIFIED REDUCTION] to kernel-moment bounds Bałaban does not print | `SOURCES_EU.md`, `SOURCES_EU_VERIFY.md`, `numerics/k0_d.py` | $k$-uniform kernel moments; unprinted $C_1'$ |
| B-S.LOC | LOC: localized covariance $C^{(k)}(Z_0, 0)$ | [PROVED+VERIFIED, CONDITIONAL] (short lemma; symmetry (S) not needed) | `SOURCES_LOC.md`, `SOURCES_LOC_VERIFY.md` | $R_2$ bound (2.16) (ledger A2, verified reduction to (W) + A1w + A7 + (An)); $\gamma_0$ for general $U$ (ledger A3, verified reduction); $\alpha_0, \alpha_1$ small |
| B-S.thr | (H-thr) threshold shell sum | Lemma T and trajectory sum [PROVED+VERIFIED], conditional on (H-hist2); (H-hist2) [VERIFIED REDUCTION] to the real uniform (H-hist), no new input (`SOURCES_HHIST2`, `SOURCES_HHIST2_VERIFY`) | `SOURCES_HTHR.md`, `SOURCES_HTHR2.md`, `SOURCES_HTHR2_VERIFY.md` | (H-hist2), M3 weights, frozen integers (L3) |
| B-S.hist2 | (H-hist2): mixed second history derivatives | [VERIFIED REDUCTION] to the real uniform (H-hist), no new input (Route 2(a); decay $\lambda_C^{m/2}$; the product form $\lambda^m$ [REFUTED]) | `SOURCES_HHIST2.md`, `SOURCES_HHIST2_VERIFY.md` | real uniform (H-hist) |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-S.D3 | sourced large-field step + R operation (E14-b) | draft verified with corrections (SOURCES_D3_VERIFY, SOURCES_D3R_VERIFY, SOURCES_D3W8_VERIFY; route (a), Lemma F2′ confirmed given (W4), (Eps), F1, (D)). (FX) is a genuine new hypothesis (SOURCES_D4D5_VERIFY.md); its state is in row B-S.FX | SOURCES_D3.md, SOURCES_FX.md, SOURCES_FX_VERIFY.md | (FX), (FX2); (U) [C23, constant choice] |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-S.FX | O1(b), O2, O3, O4, R2 | O1(b) [PROVED+VERIFIED] (Lemma [P], no additive constant); O2 [PROVED+VERIFIED] (the R-denominator is the bare Wilson integral (1.100)); O3 [PROVED+VERIFIED, CONDITIONAL ON N3] (N3 = the open HBETA strip bound $\|\delta w\| \le \frac{3}{2}\tau$); R2 ($\kappa_{fin} \ge \kappa_{final}$) [PROVED+VERIFIED] as a reduction, in volume form, given $R_{K+1} \ge 1$ and $C_c$ independent of $g$; O4 (unbounded components in (1.103)) avoided by keeping Balaban's main form (1.98), whose fallback was Gevrey $2+\eta$. N3 reduced to (H-hol) + real (H-hist) + (H2-hist), condition (a) settled [VERIFIED REDUCTION] (`SOURCES_HBETA_N3_VERIFY`, `SOURCES_N3A_VERIFY`); (H-hol) small field [VERIFIED REDUCTION] (`SOURCES_HHOL_VERIFY`); large field [VERIFIED REDUCTION] to (M-A), (M-B), (M-C), (H-hist2) (`SOURCES_HHOL_LF_VERIFY`); (M-B) [PROVED+VERIFIED] with $p_0 \ge 17/(\gamma_0 A_1^2)$, $\theta_W + \tau_C \le 0.2$; (M-A) reduced to (H−), (M-A′) closed, (H−) rests on CMP 102 (189) [VERIFIED REDUCTION] (`SOURCES_MA_VERIFY`, `SOURCES_MA2_VERIFY`); (M-C) [VERIFIED REDUCTION] to printed CMP 116 §2 conditions (`SOURCES_MC_VERIFY`); $R_{K+1} \ge L$ and $C_c$ free of $g$ are verified readings (`SOURCES_BOOKKEEPING.md`) | `SOURCES_FX_O1.md`, `SOURCES_FX_O1_VERIFY.md`, `SOURCES_FX_O23.md`, `SOURCES_FX_O23_VERIFY.md`, `SOURCES_FX_R2.md`, `SOURCES_FX_R2_VERIFY.md` | residual inputs of (M-C); CMP 102 (189) (ledger A4, verified reduction to A1) |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-S.N3 | N3: HBETA strip bound $\lvert\delta w\rvert \le \frac{3}{2}\tau$ | [VERIFIED REDUCTION] to (H-hol) + real (H-hist) + (H2-hist) | `SOURCES_HBETA_N3.md`, `SOURCES_HBETA_N3_VERIFY.md` | (H-hol), (H-hist), (H2-hist) |
| B-S.N3A | condition (a) of Lemma N3 | settled [PROVED+VERIFIED] (slack $> 5\times$; item 1 corrected) | `SOURCES_N3A.md`, `SOURCES_N3A_VERIFY.md` | constant order $L \to$ Bałaban's constants $\to \gamma \to g \le \gamma/2 \to r_0 \to K$ |
| B-S.hol | (H-hol), small field: $\beta_j$ holomorphic on $\lvert g - x\rvert < \rho_0 x$ | [VERIFIED REDUCTION] to Bałaban's conditions with modified constants | `SOURCES_HHOL.md`, `SOURCES_HHOL_VERIFY.md` | (C1′)–(C4′), (C4″), (2.24′) |
| B-S.holLF | (H-hol), large field (frozen thresholds at $x^*$) | [VERIFIED REDUCTION] to (M-A), (M-B), (M-C), (H-hist2); (M-C) itself a verified reduction | `SOURCES_HHOL_LF.md`, `SOURCES_HHOL_LF_VERIFY.md` | (M-A)–(M-D) |
| M-A | equipartition $\langle A - A_{\min}\rangle_u \le (n_i/u)(1+\eta_x)$ | [PROVED+VERIFIED] given (H−); (M-A′) closed [PROVED+VERIFIED]; (H−) [VERIFIED REDUCTION] to CMP 102 (189), an existing trust item | `SOURCES_MA.md`, `SOURCES_MA_VERIFY.md`, `SOURCES_MA2.md`, `SOURCES_MA2_VERIFY.md` | CMP 102 (189); (C1), (C2) |
| M-B | (1.80)–(1.98) bookkeeping, changed constants | [PROVED+VERIFIED] with $p_0 \ge 17/(\gamma_0 A_1^2)$, $\theta_W + \tau_C \le 0.2$ | `SOURCES_MB.md`, `SOURCES_MB_VERIFY.md` | (M-A), (M-C) |
| M-C | complex small-field step | [VERIFIED REDUCTION] to printed CMP 116 §2 (B1–B23 + 4), Lemma 2, (1.36), p. 12 (2.5)–(2.6); shrinks $\varepsilon_1/(4\lambda_V)$, $\gamma/(8\lambda_V)$; (I-X) resolved | `SOURCES_MC.md`, `SOURCES_MC_VERIFY.md` | (I-$\varepsilon$), (I-V0) residual, (M1) |
| M-D | frozen map vs Bałaban's map | not a convention: needs the frozen map's derivative at $x^*$, which is (H-hist) + (H-thr) [VERIFIED REDUCTION] | `SOURCES_HHOL_LF_VERIFY.md` | (H-hist), (H-hist2) |
| B-S.D4 | real-source derivative bounds (E14-c) | D4.1, D4.3, D4.4 and Lemma C (order $1+\eta$, given (FX)) [PROVED+VERIFIED]; ($S_{\mathrm{src}}$) $\Rightarrow$ (H-U), $s = 6$, $D > 1$, confirmed given (FX) | `SOURCES_D4D5.md`, `SOURCES_D4D5_VERIFY.md` | (FX), (FX2), S1 at all scales |

| Block | Content | Current tag | Files | Hypotheses carried |
|---|---|---|---|---|
| B-N1 | source form $\Rightarrow$ (H-U) on $\mathcal{D}_n$ | derived given the sourced Gevrey bound (S) (`ASSEMBLY_N1.md`; confirmed, `SOURCES_HHIST_N1_VERIFY.md`); ($S_{\mathrm{src}}$) $\Rightarrow$ (H-U), $s = 6$, $D > 1$, confirmed given (FX); uniformity of (S) in support size and $\varepsilon$ [VERIFIED REDUCTION] ($h$-independent; `SOURCES_B21_S_VERIFY`); ASSEMBLY_N1 aligned 2026-10-28 with the $C^1$ norm, $(n!)^{1+\eta}$, $s = 6$, $D > 1$ ($D > 2$ if the coarsest-scale (FX2) radius fails); open: the (E14-c) clause | this paper, `SOURCES_D4D5.md`, `SOURCES_B21_S.md` | (FX) main form, coarsest-scale (FX2) radius, S1 at all scales, Lemma C |
| B-C | compactness (diagonal extraction) | [PROVED HERE]; verifier: correct | Proposition 4.4 | (H-U) |
| B-REG | (OS0) | [PROVED HERE]; verifier: correct | Section 4.5.5 | (H-U); torus form of OS II |
| B-TI | (OS1) all of $T^4$ and $B_4$ | [PROVED HERE]; verifier: correct | Section 4.5.6 | (H-U); origin at a site |
| B-RP | (OS2) | lattice: [CITED]; passage: [PROVED HERE]; verifier: correct | Section 4.5.7 | $\ell$ even (sufficient); (H-U) |
| OS3 | permutation symmetry, reality | [PROVED HERE] | Section 4.5.8 | — |
| B-NT | nontriviality (Theorem 4.3) | [OPEN]: needs (NT) = (E14-f); derivation from (NT) [PROVED HERE], corrected statement | Section 4.5.9 | (NT) at every layer for (ii); $c > 0$ in Balaban's normalization |
| E13 | full-sequence convergence | not needed here; [OPEN] | `SOURCES.md` (E14-d), D6 | — |

Critical path to an unconditional (modulo GC1–GC3) Theorem 4.2: (H2) is certified at $L = 13$ (Theorem 5.1); B-T2 still needs (PI), (F), ($M_j$), V1, (G2), L5, and item 7′; then B-S: the open inputs of the real uniform (H-hist) ((Q)/$c_-$, ($M_r$), CMP 102 (189), (M-B) bookkeeping), A1w and the torus-side hypothesis under A1–A4, the residual inputs of (M-C), and the (E14-c) clause of B-N1. The blocks proved in this section are short and are not on the critical path.

## 4.7 What the theorem does not give

Each item needs a separate argument that this paper does not give.

- Rotation invariance. Only $B_4$ and the torus translations are proved. Full $O(4)$ is not a symmetry of $T^4_\ell$. `LEMMAS.md` ledger item 5 [VERIFIED QUOTE]: on the torus the inclined comparison "gives only the tilted-axis reflection symmetry $R_{2\theta}s$, never a rotation"; the template reaches rotations only after the torus $\to$ plane limit, which it controls with its mass gap.

- Infinite volume. $\ell$ is fixed. Jaffe–Witten tie the limit $T^4 \to \mathbb{R}^4$ to a volume-uniform gap [VERIFIED QUOTE]: "New ideas are needed to prove the existence of a mass gap that is uniform in the volume of space-time. Such a result presumably would enable the study of the limit as $T^4 \to \mathbb{R}^4$." No public mechanism without a gap is known to us (`LEMMAS.md` ledger item 8).

- Mass gap. Not addressed. On a finite torus a "gap" of the transfer matrix is not the Clay mass gap.

- Uniqueness. Different subsequences may give different Schwinger functions. Uniqueness needs E13, which Balaban does not have. Also not given: uniqueness of the vacuum after reconstruction (no clustering on a torus).

- Vacuum state. Time is periodic, so OS reconstruction from torus reflection positivity gives a transfer matrix on $T^3$ and a thermal state at inverse temperature $\ell$, not a vacuum state (Section 4.8, item 4).

- Also not given: clustering (OS4); a probability measure for the density field; Wilson-loop observables (class C3); the stress tensor and the operator product expansion (class C2 and beyond); independence of the limit from the block-averaging choices of Balaban's RG (only the lattice measure enters $S_{n,\varepsilon}$, but $g_0(\varepsilon)$ is chosen through Balaban's flow).

- Status of the whole statement. Even with every block closed, the theorem is conditional on Balaban's published 1984–89 proofs (GC1). Under Jaffe–Witten footnote 2 it is a compactness ("weak-existence") result on a torus with the torus forms of OS0–OS3; it is not a solution of the Millennium problem.

### 4.8 Corrections after verification

The verifier confirmed the theorem statement, all in-chapter proofs (B-C, B-REG, B-TI, $B_4$, the RP passage, OS3, the finite-form lemma), the quotes of Jaffe–Witten, and the open status of B-N1. Four items were corrected (`ASSEMBLY_VERIFY.md`). The statements above carry the corrected wording; the original claims are listed here.

1. *$\ell$ even is sufficient, not necessary* (verifier §5, grade 3). The parity count is right: $L$ odd gives $N_\varepsilon = \ell L^K$ even iff $\ell$ even; for odd $N_\varepsilon$ every time reflection has one site seam and one link seam. Osterwalder–Seiler (1978) [21] treat link-plane (between-site) reflections. Site-plane reflections were added by Menotti–Pelissetto, CMP 113 (1987) 369–373 [17] [VERIFIED QUOTE] (abstract, per the verifier): "We extend the proof by Osterwalder and Seiler of reflection positivity of lattice gauge theories with fermions to the case in which the reflection is performed with respect to planes containing sites." and "Our proof applies to all observables and relies on the particular structure of the Wilson action." The mixed case (one site seam, one link seam on a periodic lattice) is very likely routine, since each mechanism is local to its seam, but no citation states it. So odd $\ell$ is not excluded; the hypothesis $\ell$ even only avoids the mixed case. The source's phrase "plausible but not cited" stands. The periodic two-seam link version attributed to Osterwalder–Seiler is [CITED], not [VERIFIED QUOTE]. Two minor points from the verifier: the support distance after the seam shift is $\geq 2\delta - \varepsilon$ (the source's $2\delta - 2\varepsilon$ is conservative); the factor $2^{s(n+m)}$ in Step 2 is unneeded but harmless.

2. *Reference title.* The Jaffe–Witten reference list gives the title "Gauge theories on the lattice"; the bibliography entry follows the Annals of Physics title.

3. *Nontriviality needs the two-sided envelope only and support diameters $\ll \ell$* (verifier §6, grade 6). In Theorem 4.3(i), $|R| \le \frac{1}{2}$ needs $g^2 + (|x-y|/\ell)^2$ small on the whole support pair, so the support diameters must be $\ll \ell$, not only the minimum distance $d_0$. In Theorem 4.3(ii) the source stated the asymptotic $d^8 S_2 \sim c\, g(d)^4$ with $g(d)^2 \sim (\beta \log(1/d))^{-1}$. The proof uses only the two-sided bound (0.31); the asymptotic needs $\beta_j \to \beta_\infty$ and is not proved. The statement of Theorem 4.3(ii) above is the supported one. The overclaim was in the statement, not in the proof.

4. *Periodic time gives a thermal state* (verifier §6, grade 8). OS reconstruction from torus reflection positivity gives a transfer matrix on $T^3$ and a thermal state at inverse temperature $\ell$, not a vacuum. This remark is added to the list of non-results (Section 4.7).

# 5 Numerical and certified results

This section collects the numerical results that support the hypotheses of Section 3. Sections 5.1–5.3 give the one-loop increments and the per-level quantities of the Route K proof. Sections 5.4–5.6 give the values of (H2), the one-step values, and one rigorous point enclosure. Section 5.7 proves (H2) at $L = 13$ (Theorem 5.1, certified). Section 5.8 gives the original plan for a cell certificate, now an independent cross-check at $L = 13$. SU(2), infinite volume, Balaban's normalized-trace units. Except for Sections 5.6 and 5.7, none of this is certified. Only odd $L$ is inside the proofs; $L = 2, 4$ use corner blocks and are outside them.

## 5.1 First increment $\beta_1(0)$

Sources: `NUMERICS` §6–§7; `UMK_BETA1.md`.

| $L$ | $\beta_1(0)$ | how | $b = c \log L$ | $\beta_1 - b$ | status |
|---|---|---|---|---|---|
| 2 | 0.630 | $N_g = 12$, $p \to 0$ fit | 0.2575 | +0.372 | [NUMERICAL, REPRODUCED] (`KAPPA_BOUNDED_VERIFY` §4: 0.6296) |
| 3 | 0.664 | $N_g = 8$, $p \to 0$ fit (0.6625–0.6661) | 0.4082 | +0.256 | [NUMERICAL, REPRODUCED]; $N_g$ $6 \to 8$ shift $\approx 0.015$ |
| 4 | 0.757 | $N_g = 6$, single $p = 0.2$; $O((Lp)^2)$ bias $\approx 0.01$ | 0.5150 | +0.242 | [NUMERICAL] |
| 5 | 0.749 ($N_g = 12$, $p = 0.1$); fit 0.743 | umklapp code | 0.5979 | +0.145 | [NUMERICAL] |
| 7 | 0.803 | umklapp code, $N_g = 8$, $Lp = 0.5$ | 0.7229 | +0.080 | [NUMERICAL] |
| 9 | 0.833 ($Lp = 0.5$); 0.846 ($Lp = 0.9$), 0.851 ($Lp = 1.0$) | umklapp code, Windows, $N_g = 8$ | 0.8163 | $\approx +0.02$ | [NUMERICAL] |

| | | | | | |
|---|---|---|---|---|---|
| 11 | 0.850 ($Lp = 0.5$); 0.868 ($Lp = 1.0$) | umklapp code, Windows, $N_g = 8$; $N_g = 8$ grid adequacy at $L = 11$ unchecked | 0.8908 | $\approx -0.04$ | [NUMERICAL] |

The first-step constant $\beta_1 - c \log L$ is $+0.37$, $+0.26$, $+0.24$, $+0.15$, $+0.08$, $+0.02$, $-0.04$ for $L = 2, 3, 4, 5, 7, 9, 11$. It decreases with $L$ and changes sign between $L = 9$ and $L = 11$. $\beta_1(0)$ itself stays positive at every $L$ computed. The umklapp code was validated against `ym_momentum` to $10^{-12}$ per $K$ by its author; no independent verifier for the $\beta_1$ values at $L \geq 5$. The queue continues to $L = 15$ on Windows. No value at $L > 11$ exists yet.

### 5.2 Later increments ($L = 2$ only)

| quantity | value | status |
|---|---|---|
| $\beta_1 + \beta_2$ (composed, $N_g = 4/6/8$) | 1.148 / 0.967 / 0.953 | [NUMERICAL]; single $p = 0.2$, $Lp$ bias label |
| per-level $\beta_2(0)$ | $\approx 0.32$ ($0.953 - 0.630$). Per-level recursion 0.317–0.341 ($N = 8$–12, single $p = 0.2$). Symmetric Nyquist: 0.33334 ($N = 8$, $p_1 = 0.2$), 0.31679 ($N = 12$, $p_1 = 0.2$), 0.32283 ($N = 12$, $p_1 = 0.4$) (`BETA_FP_VERIFY` §7; `BETA_FP_PW_VERIFY` §5–§6). Level-1 values carry $N$-aliasing of about $\pm 5\%$. | [NUMERICAL]; per-level recursion reproduced in `BETA_FP_VERIFY` §7 |
| $\beta_3(0)$ | 0.40 ($N = 8$), 0.286 ($N = 12$), 0.261 (verifier, symmetric Nyquist, $N = 8$) | unconverged; not usable |
| abelian sector $S_k^{ab}$ | $(1 - L^{-4k})/4$, matched to all digits | [PROVED+VERIFIED]; [LEAN] counting part (`AbelianCount.lean`) |
| (U) test | $\lvert\beta_{j+1}(p) - \beta_{j+1}(0)\rvert \leq 0.2p^2$ at every computed point ($L = 2$ levels 0–1, $L = 3$ level 0, $p \leq 0.8$); uniformity in $j$ not testable with two levels | [NUMERICAL, RUN BY THE VERIFIER] (`BETA_FP_PW_VERIFY` §6) |

All computed increments are positive.

### 5.3 Quantities of the Route K proof

| quantity | value | status |
|---|---|---|
| $d_1 = b - \beta_1$, $L = 3$ | $-0.256 \pm 0.01$ (fits $-0.254 \cdots - 0.258$) | [NUMERICAL, REPRODUCED] |
| $d_1$, $\kappa_1$, $L = 2$ | $-0.372$, 1.114 | [NUMERICAL, REPRODUCED] |
| $d_2$, $d_3$, $L = 2$ | $\approx -0.44$, $\approx -0.45$ ($d_3$ uses unconverged $\beta_3$) | [NUMERICAL]; bounded-looking, not a test |

| | | |
|---|---|---|
| $\rho_1 = \max_K \|H_j(K)\|/L$, $L = 3$, $j = 1, 2, 3, \infty$ | 1.652, 1.736, 1.757, 1.760 | [NUMERICAL, RUN BY THE VERIFIER, `U3_AW_VERIFY` §6]; refutes (HS); $h^2/L^4 \approx 0.34 < 1$; one-step $C^1$ factor $\rho_1^2/L > 1$ for $j \geq 2$ |
| $\rho_1$, $L = 5$, $j = 1, 2, \infty$ | 1.860, 1.9565, 1.963; $\rho_1^2/L = 0.771$ | [NUMERICAL]; $j = 1$ and the bound measured by the verifier (`H2_LARGE_L_VERIFY` §6) |
| $\rho_2$ ((H2)), $L = 3, 5, 7, 9$ | see Section 5.4: 2.0634, 2.108, $\geq$ 2.1131, $\geq$ 2.1142 at $j = \infty$ against thresholds 3, 5, 7, 9 | [NUMERICAL, REPRODUCED] at $L = 3, 5$; [NUMERICAL, RUN BY THE VERIFIER] at $L = 7, 9$ |
| $a_*$, $L = 5$ | 0.0107 (binding mode near $K = 0$, gauge/ghost sector) | [NUMERICAL, RUN BY THE VERIFIER] (`H2_LARGE_L_VERIFY` §6) |
| (H″), $n = 2$, $L = 3$ | $2.056 < 2.828$ | [NUMERICAL, RUN BY THE VERIFIER], coarse grid, one level pair |
| $\sigma_j^{(1)}(0)$ $J$-part, $j = 0, 1, 2$ ($d = 2$ proxy) | 0.750000, 0.75015, 0.75051 vs 0.75 | [NUMERICAL, RUN BY THE VERIFIER] supports Theorem W |
| $\sigma_j^{(2)}(K)$ scan | not run: no $d = 4$ constant-field per-momentum code | — |

### 5.4 Values of (H2) [NUMERICAL; (H2) AT $L = 13$ IS CERTIFIED IN SECTION 5.7]

Code: `numerics/h2_scan.py` (umklapp Schur pieces, one colour, Gram form; `H2_LARGE_L` §1). Code structure CONFIRMED (`H2_LARGE_L_VERIFY` grade 1): the one-colour reduction is exact at $U = 1$, the $H$ and $\Delta_{out}$ blocks and the Gram identity are correct, the separable Prop. A form is exact. Sign slip in `H2_LARGE_L` §4 ($H = +[\mathfrak{B}^{-1}]_{f,c}$); harmless.

| $L$ | $j$ | $\rho_2$ (scan max) | threshold $L$ | status |
|---|---|---|---|---|
| 3 | 2 | 2.00857 | 3 | [NUMERICAL, REPRODUCED] (`B2_TWOSTEP_VERIFY` §4; `H2_LARGE_L_VERIFY` §1) |
| 3 | 3 | 2.05601 | 3 | [NUMERICAL, REPRODUCED] (spot value 2.05599 at the argmax) |
| 3 | $\infty$ | 2.0634 (2.06332–2.06377 over $l_{max}$ 2–5) | 3 | [NUMERICAL, REPRODUCED]; margin 31%; $q_1 \leq 0.473$, $q_2 \leq 0.053$ |
| 5 | 2, 3, 4, $\infty$ | 2.098, 2.108, 2.108, 2.108 | 5 | [NUMERICAL, REPRODUCED]: independent dense code gives 2.098283 at $j = 2$; the verifier's own search gives 2.107921 at $j = \infty$, same argmax; margin 58% |
| 7 | $\infty$ | $\geq$ 2.1131 (face diagonal only) | 7 | [NUMERICAL, RUN BY THE VERIFIER] (spot 2.11309) |

| | | | | |
|---|---|---|---|---|
| 9 | $\infty$ | $\geq$ 2.1142 (one point, $(2.6, 2.6, \pi, \pi)$) | 9 | [NUMERICAL, RUN BY THE VERIFIER]; the $j = 2$ point of `H2_LARGE_L` gives $\geq 2.1104$ |

- Shape. The argmax lies on the face $K_3 = K_4 = \pi$, on the diagonal $K_1 = K_2$: $\approx 2.22 \to 2.40$ ($L = 3$, $j = 2 \to \infty$), $\approx 2.54$ ($L = 5$). Off-face points are lower: best off-face 1.995 ($L = 3$, $j = 2$), 2.004 ($L = 3$, $j = \infty$), 2.088 ($L = 5$, $j = \infty$). The corner $(\pi, \pi, \pi, \pi)$ stays below the face peak, but the gap shrinks: 0.0070, 0.0028, $\geq 0.0025$, $\geq 0.0023$ ($L = 3, 5, 7, 9$). The corner is the obvious place for a competing maximum at large $L$.

- Level dependence. At $L = 5$, $\rho_2(\infty) - \rho_2(j) = 9.64 \cdot 10^{-3}$, $4.0 \cdot 10^{-4}$, $1.6 \cdot 10^{-5}$ ($j = 2, 3, 4$); ratios 24.1, 25.0, against $L^2 = 25$. At $L = 3$ the ratios are 7.5, 8.7. Levels $j = 4, \ldots, \infty - 1$ are not sampled at $L = 3$.

- $L$-trend. $j = \infty$: 2.0633, 2.1079, $\geq 2.1131$, $\geq 2.1142$ ($L = 3, 5, 7, 9$). This is consistent with a bounded $\rho_2$ near 2.11–2.14. The rate claim "at least as fast as $L^{-2}$" is CORRECTED (`H2_LARGE_L_VERIFY` grade 5): it used a lower bound at $L = 7$ as if it were the sup. "(H2) gets easier with $L$" is a measured trend for $L \leq 9$, not a result for $L > 11$.

- All values are lower estimates of a sup (grid plus local search).

### 5.5 One-step values

The arithmetic of Section 3.7.4 shows that $\rho_1^2 < L$ implies (H2). The measured one-step values are as follows.

- $L = 3$: $\rho_1^2 = 3.10 > 3$. The one-step route fails (Section 3.7.2); the two-step map is needed.

- $L = 5$: $\rho_1 = 1.860$ ($j = 1$, Wilson fine kernel, dense, measured by the verifier), 1.9565 ($j = 2$), 1.963 ($j = \infty$). So $\rho_1^2 \leq 3.853 < 5$ and $\rho_1^2/L = 0.771$ [NUMERICAL, CONFIRMED AS A MEASUREMENT, UNCERTIFIED; `H2_LARGE_L_VERIFY` GRADE 4].

- $L = 7$: $\rho_1 \geq 2.032$ (diagonal only), $\rho_1^2/L \approx 0.59$ [NUMERICAL].

- $L = 13$: measured $\sup \rho_1^2 = 4.37$; certified bound $\sup_{j,K} \rho_1^2 \leq 10 < 13$ (Theorem 5.1).

So at $L \geq 5$ the one-step map suffices for Theorem C1, as far as measured. For certification this replaces $L^8$ blocks per $K$ (two-step map) by $L^4$ blocks per $K$ (625× fewer at $L = 5$).

### 5.6 A rigorous point enclosure at $L = 3$ [INTERVAL ENCLOSURE, VERIFIED]

Source: `H2_CERTIFY.md` with the prototype `numerics/h2cert_proto.py`; verifier `H2_CERTIFY_VERIFY.md`. At $L = 3$, $j = \infty$, and the face point $K^* = (2.4035, 2.4035, \pi, \pi)$ near the argmax of Section 5.4, ball arithmetic (python-flint) gives

$$\rho_2(\infty, K^*) \in [1.9993,\ 2.1273] < 3.$$

The width comes from the umklapp tail and scales as $N^{-4}$ in the truncation $N$. The verifier reran the code at $N = 12$ and obtained [1.5475, 2.5792], which contains the floating-point value 2.063361 and confirms the $N^{-4}$ scaling; at $N = 6$ the code stops with an assertion instead of returning a wrong interval. The verdict is CONFIRMED given the premises P1–P3 of `H2_CERTIFY_VERIFY`; P1

(the loop-family coefficients of the (0.4) row are exact rationals $k/648$, $k$ even) is proved there, and the fidelity of the loop family to Bałaban's paper stays inside P3. Further results of the same two files:

- One-step factorization: $\rho_2(j,K) \le \rho_1(j,K)\cdot \max_l \rho_1(j-1,k_l)$, exact [PROVED+VERIFIED]. So two-step work is needed only where the right side is $\ge L$, near the face diagonal.
- Oishi–Rump bound $\|\mathfrak{B}^{-1}\| \in [42.7,\ 43.6]$ at $K^*$ (bounds CONFIRMED). The values 177 and 195 of Section 3.9 use the full $\Delta$ instead of $\frac{1}{4}\Delta$; this factor reconciles them. No $\|\mathfrak{B}^{-1}\|$ enters $\rho_2$.
- The umklapp tail enclosure is CONFIRMED for $M=\infty$. For finite $M$ it is CORRECTED: its lower bound must be 0, with class-wise factors $(\pi/2)^{12}$ for two or more large components.
- A uniform Lipschitz grid is refuted as a route: $\Lambda_K \approx 2.8\cdot 10^4$ (CONFIRMED) would need about $10^{13}$ cells. The replacement is a per-cell invertibility test with a mean-value enclosure.
- Cost: `H2_CERTIFY` estimates about $1.5\cdot 10^4$ core-hours for a full certification at $L=3$. The verifier grades this estimate UNSUPPORTED: at the tested sizes the arb overhead dominates, and the box-sum speedup is overstated.

This is one point at one level. It does not certify the supremum over the Brillouin zone or over $j$. Section 5.7 does this at $L=13$.

## 5.7 (H2) at $L=13$ [CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]

Sources: `H1_ANALYTIC.md` §0–§8 (statement, lemmas, majorant); verifier `H1_ANALYTIC_VERIFY.md` V0–V9 with `numerics/h1analytic/bb_arb.py`; operator check `H1_ANALYTIC_VERIFY2.md` with `numerics/h1analytic/indep_H.py`; symmetry check `H2_CERTIFY_P5_VERIFY.md`. Here $\rho_1(j,K) = \|H_j(K)\|/L$ with $H_j(K)$ the one-step gauge-fixed minimizer map of Section 3.7.

**Theorem 5.1** ((H2) at $L=13$)**.** *For $L=13$, every level $j\ge 1$ including $j=\infty$, and every coarse momentum $K\in \mathrm{BZ}$,*

$$\rho_1(j,K)^2 \le 10 < 13.$$

*Hence $\rho_2 \le \sup_{j,K}\rho_1(j,K)^2 \le 10 < L$ by the factorization $\rho_2 \le \rho_1(j)\max_l \rho_1(j-1,k_l)$ (Lean `rho2_le`, Section 3.7.4). That is, (H2) holds at $L=13$.*

*Proof sketch.* Write $s=L^{-2}$, $k_l=(K+2\pi l)/L$, $M=L^{j-1}$, $d^M$ for the Prop. A symbol at level $M$, $P_\perp$ for the projection onto $\hat K^\perp$, and $T$, $T_2$, $\sigma$, $Z=(P_\perp T P_\perp)^+$ as in `H1_ANALYTIC` §1.

1. Exact split. $\rho_1(j,K)^2L^2 = \lambda_{\max}\big(ZT_2Z + cc^\dagger/(\sigma|\hat K|^4)\big)$, $c^\dagger = \hat K^\dagger(1-TZ)$: a transverse part and an orthogonal pure-gauge part.
2. Three exact lemmas. Lemma S: the averaging symbol is $E_q(k_l) = E_l^0 - \hat K g_l^\dagger$, with $E_l^0$ diagonal (block average times straight path), $|E^0_{l,\mu\mu}|$ built from $a_{l,\rho} = \sin(K_\rho/2)/(L\sin(k_{l,\rho}/2))$, and $\sum_l \prod_\rho a_{l,\rho}^2 = 1$ (Fejér). Tower identity: $d_\mu^{ML}(K) = L^{-2}\sum_l w_{l,\mu} d_\mu^M(k_l)$, hence $P_\perp T P_\perp = P_\perp \operatorname{diag}(d^{ML}(K))P_\perp$ and $Z$ is the next-level kernel of Prop. A. Lemma G: a closed form of the gauge row $g_l$ from the path average, so the averaging operator is explicit.

3. Majorant. Lemma T (transverse): $\lambda_{\max}(ZT_2Z)/L^2 \le \gamma_L F(K)$, $F(K) = 1/(|K|^2 \underline{d}(K))$, $\gamma_L = \mathrm{sinc}(\pi/2L)^{-2}$, from $T_2 \le d_{\max} T$, $ZTZ = Z$, $d_{\max} \le L^2\gamma_L/|K|^2$, and $\|Z\| \le 1/\underline{d}$. Lemma Gauge: the gauge part is $\le (1+\tau)^2/(|\hat{K}|^2 S_L)$ with explicit sums $V$, $U$ and $\tau = (V + |\hat{K}|U)/\underline{d}$. So $\rho_1(j,K)^2 \le B_L(K) := \gamma_L F(K) + (1+\tau)^2/(|\hat{K}|^2 S_L)$. By Cauchy–Schwarz $F \le F_{CS}$, a closed form made of one-dimensional factors in each $K_\rho$.

4. Uniformity in $j$. $B_L$ does not depend on $j$. For every $M \in \{1, L, L^2, \dots, \infty\}$ the fine symbol satisfies $d^M \le 1/|\tilde{q}_0^M|^2 \le 1/|\hat{k}|^2$, because the Fejér weights sum to 1. The coarse symbol $d^{ML}$ dominates the truncated $M = \infty$ sum termwise. No monotonicity in $j$ and no O5 rate is used.

5. Branch-and-bound. $B_L$ is invariant under the 384 signed permutations of $K$ by construction, so the sup over the sorted chamber $0 \le K_1 \le \cdots \le K_4 \le \pi$ is the sup over BZ. The program `numerics/h1analytic/bb_arb.py` encloses $\gamma_L F_{CS} + \bar{G}_L$ on boxes $K = \pi u$ with exact dyadic $u$, evaluates every one-dimensional value in arb (python-flint), and rounds outward by one ulp after every floating-point operation, with Higham bounds on every sum. Result: $\sup_{\text{chamber}} B_{13} \le 9.9995277$, 25,169 boxes, 8,075 gauge evaluations, 0 failed leaves, 58 s. Boxes that contain $K = 0$ are enclosed directly; $\rho_1(j,0) = 1$.

Margins. The measured $\sup \rho_1^2$ is 4.37. The majorant is tight to 2.6% at $|K| \approx 1.5$ ($\min B_{13}/\rho_1^2 = 1.026$ over 896 cases and a local search). The certificate refines boxes only until each is $\le 10$, so its margin to 10 is $5 \cdot 10^{-5}$; its margin to the threshold 13 is 23%.

Verification. `H1_ANALYTIC_VERIFY` rederives the split, Lemmas S, G, the tower identity, Lemma T, and Lemma Gauge on paper and checks them against the production objects to $\le 5.3 \cdot 10^{-13}$ (V1–V2); it checks $B_{13} \ge \rho_1^2$ at 896 $(j,K)$ cases (V3–V4); it grades the original float program CORRECTED (no directed rounding) and replaces it by `bb_arb.py` (V5–V8). `H1_ANALYTIC_VERIFY2` rebuilds $H_j(K)$ from the definitions (position-space (0.4) paths, sparse KKT, ghost-domain gauge) and reproduces the production $\rho_1$ to $5.8 \cdot 10^{-9}$ ($10^{-14}$ with the same $d$ symbol); the two parts of the split agree to $3 \cdot 10^{-8}$, and the Lemma S conventions are derived by hand and confirmed to $1.2 \cdot 10^{-13}$. `H2_CERTIFY_P5_VERIFY` proves and checks the invariance of $\rho_1$ under the 384 signed permutations (P5a, P5b); the analytic certificate does not need it, the box sweep does. Lean: `lean/YmLean/Tower.lean` formalizes the algebraic core of the majorant (tower identity in weighted form, Rayleigh and rank-one bounds, `majorant_core`, `lemmaT_core`); `BlockNorm.rho2_le` is the one-step $\Rightarrow$ two-step factorization.

**Residue.** The theorem is certified given the following, which no second source has checked.

- The reading of Bałaban's (0.4) (the path family that defines the averaging operator) and the gauge choice (`NUMERICS` §5). The certificate is a statement about $H_j$ so defined; both implementations use the same reading.
- Prop. A at $M \ge 27$ and $M = \infty$: checked only by a second implementation of the same formula (independent recursion at $M = 3, 9$).
- Lemma G and the bound chain: proved on paper and rederived by the verifier, checked numerically at points only. The Fourier/umklapp derivation, Lemmas S and G, and the identification of the split are not in Lean.
- `bb_arb.py`: audited line by line and checked against the float program on 30 boxes; not formally verified. It trusts python-flint arb and IEEE correctly rounded $+, -, \times, \div, \sqrt{}$.
- Independent cross-check: the box sweep of the cell certificate (Section 5.8) is running; its result is not in yet.

### 5.8 Certification plan and $j_0$ [DESIGNED; SUPERSEDED AT $L = 13$ BY SECTION 5.7]

At $L = 13$ the analytic route of Section 5.7 replaces this plan: it needs no tail level $j_0$, no $\|\mathfrak{B}^{-1}\|$ bound, and no Lipschitz constant. The cell certificate below is run at $L = 13$ only as an independent cross-check (box sweep, in progress). `B2_TWOSTEP_VERIFY` grade 6 (CORRECTED in size): certifying (H2) is a computer-assisted proof project, not a check. `H2_LARGE_L` §4 gives the design; `H2_LARGE_L_VERIFY` grades 6–8 check it. Three parts.

(a) Tail $j \geq j_0$. From the resolvent identity, $\rho_2(j) \leq R_\infty + (\rho_1^2/a_*)(\delta_{j-2} + \delta_{j-1})$, where $R_\infty = \rho_2$ at $j = \infty$, $a_* = \inf_K a_j(K)$ with $\|[\mathfrak{B}^{-1}]_{f,f}\| = 1/a_j(K)$ exactly, and $\delta_m = \sup_K \|\Delta_{L^m} - \Delta_\infty\|$ (O5 decay) [derivation CONFIRMED]. Inputs CORRECTED by the verifier: the 2-norm sup of $\delta_m$ is at the corner, $\delta = 179, 92.0, 18.7, 2.39, 0.317$ ($M = 1, 3, 9, 27, 81$), 45–90× the assumed value at small $m$; $a_*(L = 5) = 0.0107$, not 0.05. With these inputs, $j_0 \approx 8$ ($L = 3$) and $\approx 6$ ($L = 5$). With the O6 coercivity constant, $j_0 = 25$, which cannot be used in practice (`B2_TWOSTEP_VERIFY` §5 estimated $j \gtrsim 45$ with $\|\mathfrak{B}^{-1}\| \leq b(L) \sim 10^{19}$).

(b) Levels $2 \leq j < j_0$ and $j = \infty$. A certified sup over the BZ of $\|H_{(2),j}(K)\|$ (at $L \geq 5$: of $\|H_j(K)\|$, Section 3.7.4). Plan: interval arithmetic with local mean-value enclosures on boxes in the sorted chamber (1/384 of the BZ), $4 \times 4$ and $5 \times 5$ verified inverses, one $4 \times 4$ $\lambda_{max}$ per $K$; about $2.5 \cdot 10^3$ boxes at half-width 0.05 at $L = 3$; the analytic form of $\Delta$ near $K = 0$, where $\rho_2(0) = 1$. At $j = \infty$ also a certified tail bound for the Prop. A sum ($l_{max}$ truncation; observed effect $4 \cdot 10^{-4}$). There is no explicit global Lipschitz constant today: the A0/O5 strip width is existential, and the constant that can be made explicit is $\sim 10^{40}$. Monotonicity in $j$ is not proved and is not used.

(c) Inputs to certify first: $\|\mathfrak{B}_j(K)^{-1}\| \leq B_{num}$ for all $j$, $K$ (the O6 constant $b(L)$ is unusable), the O5 constant $C_0$ explicitly (or certified $\delta_m$), certified $a_*$, $\rho_1$, $R_\infty$.

The run plan (`H2_LARGE_L_VERIFY` grade 8) uses atomic JSON writes and pre-copied profile caches; the profile build time at $L \geq 11$ is not budgeted. At $L = 13$ the cell certificate runs as the box sweep (`numerics/h1cert_cell2.py`, sorted chamber, valid by P5), an independent cross-check of Theorem 5.1; its result is not in yet.

## 6 What remains

This section lists every open item of the paper in one place. The items are grouped by the kind of work they need: certification, fixed-level and structural inputs of Theorem 2w, items of the sources block, and verification. The item labels H, 1–13, and B1–B8 are those of the dependency table (Section 4.6). For Theorem 2w ⇒ Theorem 1 the list covers §10 of `THEOREM2_PROOF_v4.md`. The trust ledger (Section 6.7, `TRUST_LEDGER.md`) supersedes this list as the complete record of what the two main theorems take on trust: Bałaban-level statements, external results, our open items, and conventions.

### 6.1 Certification

Single uniform hypothesis (both routes):

H. (H2) $\rho_2 = \sup_{j \geq 2} \sup_{K \text{ real}} \|(\oplus_l H_{j-1}(k_l)) H_j(K)\|/L^2 < L$. [DONE AT $L = 13$: CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)], Theorem 5.1: $\rho_1(j, K)^2 \leq 10 < 13$ for all $j \geq 1$ (including $\infty$) and all $K$, hence $\rho_2 \leq 10 < 13$ by `rho2_le`. Route K: (H2) ⇒ ($C1\sigma$) ⇒ (B2) [PROVED+VERIFIED, `B2_TWOSTEP_VERIFY` GRADE 3]. Route B: (H2) + (T1) ⇒ (U) [PROVED, UNVERIFIED, `BETA_FP_U` THEOREM 2.4]. Cross-check in progress: the box sweep of the cell certificate at $L = 13$ (Section 5.8). At $L = 3, 5, 9$ (H2) stays [NUMERICAL].

H′. Residue of Theorem 5.1 (Section 5.7): the reading of Bałaban's (0.4) and the gauge choice (single source); Prop. A at $M \geq 27$ and $M = \infty$ (second implementation of the same formula only); Lemma G and the bound chain (point checks only beyond the paper proof); `bb_arb.py` audited, not formally verified.

At $L = 3$ one point is enclosed: $j = \infty$, $K^* = (2.4035, 2.4035, \pi, \pi)$, $\rho_2 \in [1.9993, 2.1273]$ [INTERVAL ENCLOSURE, VERIFIED] (Section 5.6). Certified early steps $\beta_1(0), \ldots, \beta_{j_*}(0) > 0$ at odd $L > 11$ are needed only for literal (0.31) at every $K$ (Section 3.9.2).

## 6.2 Fixed-level and structural inputs of Theorem 2w

Route K hypothesis set (all must hold):

1. (B2) $\sup_j \sup_K |\sigma_j^{(2)}| < \infty$ (or (B2w)). Reduced to H [VERIFIED REDUCTION]. Older reductions kept: $(C1\sigma)$ + (h) [PROVED+VERIFIED]; (H″) [ARGUED]. (HS) is refuted.
2. (PI) probe independence, at each fixed level [ARGUED].
3. (F) lattice = continuum log coefficient at constant field [CITED SOURCE CHECKED (SAVVIDY [22], LEDGER X5); LATTICE TRANSFER STANDARD, NOT PROVED HERE].
4. $(M_j)$ formal = true $f^2$ coefficient at each fixed level: (S1) $\mathcal{K}_t(f) \geq -C_1|f|$, (S2) density of states $\leq C_2|f|$ near 0, (S3) relative bound of the level difference, all on the lattice [OPEN]; Prop. 4.4 [ARGUED].
5. V1 order-$g^0$ completeness of CMP 109 (1.3)–(1.5): $\beta_{j+1}(0)$ is the Maxwell coefficient of the per-step Gaussian determinant [OPEN; SUPPORTED AT ORDER $p^2$].
6. (G2) per-cell infinite-volume densities [FORMAL].

Route B hypothesis set (alternative to 1–5; per `BETA_FP_U` §4; all [PROVED, UNVERIFIED] as implications, no verifier file for `BETA_FP_U`):

B1. (T1) $\sup_j \sup_{q \text{ real}} |c_{T,j}^{(1)}(q)| < \infty$, the plane-wave form of Theorem A3 [ARGUED; THE LANDAU-TO-SYMMETRIC GAUGE CONVERSION AND STEPS 2–5 OF A3 FOR (2.1) ARE NOT WRITTEN].

B2′. (H2) = item H.

B3. (DF) $\sup_k |\mathfrak{d}_F(\theta_*; L^k)| < \infty$, the gauge-fixing difference at the plane wave [HYPOTHESIS, ARGUED]. With the exact split $\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$ (`BETA_FP_U` Prop. 3.1) there is no $Y$-defect; all non-stationarity sits in $\mathfrak{d}_F$. This replaces the earlier $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ items for this route (Section 3.6.2).

B4. Lüscher–Weisz (5.7) [CITED].

B5. A6 (the label of v4; not ledger A6): the plane-wave increment equals Balaban's $\beta$ up to the expansion point, uniformly in $k$ [ARGUED]. A uniform-in-$k$ input, of the same kind as (DF).

B6. (H0) = item 7′ (Lemma D″ (i)).

B7. $\theta_* < 1$, so that $c_2 \ln(1/\theta_*^2) \geq 0$ in $C_B$ [TRIVIAL CONDITION ON THE CHOICE OF $\theta_*$].

Route B also needs 6 ((G2)), item 7′, and the items 8 and 10–13 (item 9 is closed). It does not need (PI), $(M_j)$, V1; it needs (DF) and A6 instead.

Common to both routes:

7′. The own-slot bound $|\beta_{k+1}(h,g) - \beta_{k+1}(0)| \le C_\beta g$ on $]0,\gamma]^k \times [0,\gamma]$, for every past $h$, monotone or not [PROVED+VERIFIED]. Item 7 asked for (H0), Bałaban's smoothness in $g$ with bounds uniform in $j$ and history (CMP 109 inductive assumption at odd $L > 11$; "uniform in $j$" is our reading of p. 264, not printed, `R2_VERIFY` row 12; uniformity in the history is not printed; ledger A5). The shooting evaluates $\beta$ on clamped, non-monotone pasts, and `SOURCES_T2_KUNIF_VERIFY` found that (G-gain) proves $C_\beta$ only on monotone pasts. `SOURCES_GGAIN_NONMONO.md` lifts the restriction with no change of constant: the only use of monotonicity is a comparison of bound sizes in Lemma F2′, and the running maximum $G_k = \max_{i \le k} g_i$ closes the induction for every past. So 7′ follows from (H0) or from (G-gain) with this lift. The proof uses the Lipschitz form on $]0,\gamma]$ only.

8. L5 history dependence [OPEN]. (a) The clamp proof of Lemmas D′/D″ is valid only if $\beta_{k+1}(g_0,\dots,g_k)$ is defined and continuous on the whole cube $]0,\gamma]^{k+1}$; Balaban does not supply this. (b) Balaban's $\beta_{k+1}$ exists only on histories where his induction has run; that domain must carry (c′) + (S), not only $g_i \in ]0,\gamma]$. Clamped histories can violate the growth side of (c′). (c) True trajectories with all $u_i \ge \gamma^{-2}$ inherit (c′) from (a″). A nested-interval variant of Lemma D″ on these trajectories would avoid the gap [NOT WRITTEN]; the variant is written only for Lemma D (`LEMMAS_DE_VERIFY` §1.5). Lean covers one-argument $\beta$ only (Lemmas D′, D″, D, E).

9. $K$-independence of $\beta_j(g)$ [CLOSED AS A REDUCTION; VERIFIED] (`SOURCES_T2_KUNIF.md` + VERIFY). Item 9 is a volume statement, not a history identity: $K$ enters $\beta_j$ only through the torus size. CMP 119 (2.24), p. 249, and p. 279 define $\beta_j$ as the whole-lattice coefficient, so the $\beta_j$ of the recursion is $K$-free by definition. Lemmas D′/D″ run at one fixed $K$ and never compare trajectories. The window bound needs only (K1) (history-free at zero own coupling), (K2) $C_\beta$ (item 7′), and (K3) the window bound (5.1) of Theorem K. The residual volume question (the finite-volume step equals the whole-lattice one, ledger A15, B8) is an input of Theorem 1, not of the shooting.

11. Odd $L$. All one-loop proofs need odd $L \ge 3$ (centered blocks: U3 Lemma 1.2, O5/Prop. A, evenness, Lemma 4.1 of LEMMA43). Balaban assumes $L$ odd $> 11$ throughout CMP 109, so (H0) and $\beta_j(g)$ at $g > 0$ exist only there. Theorem 2w is stated for odd $L > 11$, or with (H0) at this $L$ as a hypothesis. Numerics: $\beta_1(0)$ up to $L = 11$ (Section 5.1); (H2) at $L = 3, 5$ (scans), 7 (face diagonal), 9 (one point); $\rho_1$ at $L = 3, \dots, 13$. Only $L = 13$ is inside Bałaban's range; there (H2) is certified (Theorem 5.1).

12. SU(2) only. The charged/neutral split of Lemma 5.1 and (W) are SU(2) statements; $c = 11N^2/(12\pi^2)$ is general but the chain is not checked for $N > 2$.

### 6.3 Items of the sources block

These items stand between the trajectory of Section 3 and the bound (H-U) of Section 4.2. Tags and files are those of the dependency table.

- (B-S) The sourced version of Bałaban's Theorem 1, (E14-a, b, c), is [OPEN] as a whole.
- (H-hist) via (G-gain). The small-field step is done: pointwise Cauchy and uniform $C_\beta$ on $]0,\gamma]$ [PROVED+VERIFIED]. Lemma G3 (the $\chi_k$ surface term) holds given (Q); (Q) is a verified reduction to CMP 99 (3.132), ($E_U$), and LOC. The large-field half is a [VERIFIED REDUCTION] (`SOURCES_GGAIN_LF.md` + VERIFY): one Cauchy step on the large-field disc, with the factor

$1/g$ absorbed by the exponent of $p_0$ (CMP 119 p. 244). The real uniform (H-hist) is therefore reduced [R] in both halves; open inputs: (Q)/$c_-$, the margin ($M_r$), CMP 102 (189), and the (M-B) bookkeeping. Lemma G4 is off-chain (`SOURCES_G4.md`). Second derivatives: (H-hist2).

- Bałaban-level core (round 3). The propagator and covariance inputs A1–A4 of the ledger reduce to published theorems:
  - A1, kernel-decay form: [VERIFIED REDUCTION] via Lemma CT to the printed CMP 99 Theorems 3.1, 3.3, and the $G$ part of 3.12, with (KE′) proved by citation (CMP 95 (1.11)/(1.18), CMP 96 (2.35)), ($H_Q$) proved, and S2 closed for $G$. Residue: a stated torus-side hypothesis (side $\geq c \cdot RM$ coarsest blocks), two unprinted conventions, and the $G_1$ clause at P-sketch level.
  - A1w, walk-expansion form of A1 (CMP 99 Theorem 3.15, "analyzed in the same way"): [OPEN]. Lemma CT gives kernel decay only, not a walk expansion.
  - A2 ($R_2$, $R_3$ of CMP 116 (2.16)): [VERIFIED REDUCTION] to the walk bounds (W) = CMP 99 (3.107)–(3.108) + A1w, decay of $C^*\Delta_k C$ at real $U$ (A7), and analyticity on the $\alpha'$-space (An); $R_3$ in weaker forms sufficient for (2.21).
  - A3 ($C^*\Delta_k C \geq \gamma_0$): [VERIFIED REDUCTION] (P-sketch) to the printed CMP 99 Theorems 3.1, 3.3, 3.11–3.12, with $\gamma_0 \geq (48d^2)^{-1}L^{-d-1}$ and no $M$; the $\sigma$-localized clause reduces to (W) + (N), without symmetry.
  - A4 (CMP 102 (189)): [VERIFIED REDUCTION] to A1 + published CMP 102 statements + A18 + one bookkeeping item.

  So the Bałaban-level core now rests on published propagator theorems, A1w, the two conventions, and the torus-side hypothesis.

- (H-hist2), on which the threshold-shell bound (H-thr) depends: [VERIFIED REDUCTION] to the real uniform (H-hist), no new input; with it the M3 history weights and the frozen-integer convention (L3).

- (FX). O1(b), O2, and R2 are [PROVED+VERIFIED]; O3 is [PROVED+VERIFIED, CONDITIONAL ON N3]; O4 is avoided by Bałaban's main form (1.98). N3, the HBETA strip bound $|\delta w| \leq \frac{3}{2}\tau$, is reduced to (H-hol) + real (H-hist) + (H2-hist) (condition (a) settled); (H-hol) is a verified reduction to (M-A) (via (H−) and CMP 102 (189)), (M-B) (verified), and (M-C). (M-C) is a [VERIFIED REDUCTION] (`SOURCES_MC.md` + VERIFY) to the printed conditions of CMP 116 §2 (B1–B23 and four more), Lemma 2, (1.36), and p. 12 (2.5)–(2.6), with shrinks $\varepsilon_1/(4\lambda_V)$, $\gamma/(8\lambda_V)$; open: (I-$\varepsilon$), the residual of (I-V0), and "$M$ large" (M1). The two minor inputs of R2 ($R_{K+1} \geq L$, $C_c$ independent of $g$) are verified readings (`SOURCES_BOOKKEEPING.md`). (FX2) is conditional on O3, so also on N3.

- (B-S.D2C, B-S.H$\beta$) The domain hypothesis (D), and the conditions (I) and (H-hist) of the two-loop bound.

- (B-S.D2, B-S.D3) The recovery of the decay loss $\kappa \to \kappa/2$ [VERIFIED REDUCTION] by $L/2$-rescaling, $\kappa \geq 1536$ at $d = 4$ (ledger B22; it uses the unprinted large-field T-step rate loss, ledger A23). The scale condition (U) ($\rho_1 \leq c_* r_0/M^a$ at the unit scale) is a constant choice (ledger C23), verified compatible with (B-N1) (`SOURCES_UN1.md`), and does not appear in Theorem B.

- (B-N1) The bridge from the source form to (H-U) derived given (S) (`ASSEMBLY_N1.md`, confirmed); the uniformity of (S) in support size and in $\varepsilon$ is a [VERIFIED REDUCTION] (`SOURCES_B21_S.md` + VERIFY): every factor of $C^n(n!)^{1+\eta}$ is independent of $h$, given Lemma C, S1 at all scales, (FX) in the main form, and the coarsest-scale (FX2) radius; open: the (E14-c) clause. Given (FX), the step $(\mathrm{S}_{\mathrm{src}}) \Rightarrow$ (H-U) with $s = 6$, $D > 1$ is confirmed.

- (NT) = (E14-f), for nontriviality (Theorem 4.3); at every layer for part (ii). The constant $c > 0$ must be computed in Bałaban's normalization.

- E13, full-sequence convergence: not needed for Theorem B; [OPEN].

### 6.4 Verification debt

10. Balaban readings: the sweep (Section 3.4.3) is grep plus page images, not line-by-line, with 4 of 19 rows checked by a second reader; "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) not located; "$\beta_0$ can be chosen arbitrarily small" is on CMP 119 p. 255, and $\beta_0$ is fixed before $\gamma$, as used (verified reading, `SOURCES_BOOKKEEPING.md`; former ledger A12); the readings of `BALABAN_EXCERPTS_CHECK` item 11 (II (4.37)/(5.1), CMP 119 (2.42), (2.12)/(2.23), the "$L$ odd" page) are verified, with corrections, in `SOURCES_BOOKKEEPING.md` (former ledger A22); the CMP 119 Sect. 3 residue is the unprinted T-step rate loss (ledger A23); Theorem 1 itself rests on Balaban's published proof, which uses the Group A items of Section 6.7.

B8. Verification: `BETA_FP_U` has no verifier file. `BETA_FP_PW_VERIFY` §6 items 1–4 grade the logic of Theorem B [PROVED+VERIFIED, CONDITIONAL] (tag decision, `PLAN.md`); the inputs (U), (DF), (T1) stay [PROVED, UNVERIFIED].

13. Lean formalization of the history form of Lemmas D′/D″ [OPEN]. Lemma D″ in the one-argument form is done (Section 3.4.2).

- Results tagged [PROVED, UNVERIFIED] need a verifier pass; Appendix B lists them. They include the sweep, the step Theorem 2w ⇒ Theorem 1, the `BETA_FP_U` reductions, `BETA_FP_M` Prop. 4.3, the Prop. 3.6 assembly, and H2_CERTIFY_L13 (its cell plan is superseded at $L = 13$ by Theorem 5.1; the box sweep runs as a cross-check). H2_CERTIFY_P4 has partial support: identity (D) is proved in Lean (`P4_divergence`), Lemma S is rederived in `H1_ANALYTIC_VERIFY.md` V1, and the rebuilt $H_j$ of `H1_ANALYTIC_VERIFY2.md` confirms it numerically; the Fourier step has no independent verifier.

- The paper is not verified as one document. Its statements carry the tags of the project files.

### 6.5 Open for stronger statements

Not needed for Theorem 1, open for the stronger statements: (A) in full and Lemma 5.1 for $m = 2$ (convergence, rate); $C^2$, $C^3$ of $\sigma^{(1)}$ under (H2) (not written); certified $\beta_1(0), \dots, \beta_{j_*}(0) > 0$ at odd $L > 11$ (literal (0.31) at every $K$); explicit $j_*$; (N).

### 6.6 Why the mass gap is not on this route

Write $\beta_W$ for the coefficient of $\sum_p (1 - \frac{1}{N} \operatorname{Re} \operatorname{Tr} U_p)$ in the Wilson action. Bałaban's action is $g^{-2} \sum_p (1 - \operatorname{Re} \operatorname{tr} U_p)$ with the normalized trace (CMP 109 (0.2)), so $\beta_W = 1/g_B^2$. Shen, Zhu, and Zhu [24] prove a mass gap for SU($N$) lattice Yang–Mills with action $N\beta \operatorname{Re} \sum_p \operatorname{Tr} Q_p$ (their (1.2))

when $|\beta| < 1/(16(d-1))$ (their (1.3)); with $\beta_W = N^2\beta$, $N = 2$, and $d = 4$, this is $\beta_W < 1/12$. Bałaban's construction requires $-2(1+\beta_0)^{-1}p_0(g_k) + 1 + 2\kappa(100R_k)^d \le -\frac{3}{2}p_0(g_k)$ (CMP 122 II p. 389). With $d = 4$ and $R_K \ge 1$, this gives $p_0(g_K) > 4\kappa \cdot 10^8 + 2$ for every $\beta_0 > 0$ (twice this at Bałaban's $\beta_0 = 1/7$). The unit-scale threshold $g_K p_0(g_K)$ must be below 2, the largest value of $\|U - 1\|$ on SU(2); with $\kappa \ge 1$ this gives $\beta_W = 1/g_K^2 > p_0(g_K)^2/4 > 4 \times 10^{16}$ at the unit scale. The two regimes are therefore separated by at least seventeen orders of magnitude in $\beta_W$, and no choice of $L$ narrows this, because $L$ does not enter the condition. A proof of a mass gap needs a mechanism outside the renormalization-group construction, and Bałaban's papers do not address it.

### 6.7 Trust ledger

The file `TRUST_LEDGER.md`, corrected by `TRUST_LEDGER_VERIFY.md` (Steps 1–28), lists every statement that the two main theorems use and that this project does not prove with a verifier's confirmation. The three tables below condense it to one row per item. Group A holds Bałaban-level statements and external results (X), Group B holds our own open items, and Group C holds conventions and choices. The IDs are those of the ledger. Ledger B1–B22 are not the Route B items B1–B7 or item B8 of this section, and ledger A6 is not the item "A6" of Route B (that label comes from v4 §4.2a). Sources are cited by paper and page; "[I]" marks an inference step of ours that no verifier proved.

**Group A**: Bałaban-level statements and external results

| ID | statement | source | status |
|---|---|---|---|
| A1 | Kernel bounds on $(QGQ^*)^{-1}$, $(QG_1Q^*)^{-1}$, uniform in $k$ and regular $U$ | CMP 99 p. 422 (3.132) | asserted, "not repeated"; kernel-decay form a verified reduction via Lemma CT to CMP 99 Thms 3.1, 3.3, 3.12 + (KE′) + S2; residue: torus-side hypothesis, two conventions, $G_1$ P-sketch |
| A1w | Walk-expansion form of A1 (builds the walk expansion of $C^{(k)}(Z_0)$, bounds (3.107)–(3.108)) | CMP 99 Thm 3.15, p. 416 | asserted, "analyzed in the same way"; open |
| A2 | $R_2$ and $R_3$ obey (2.16) | CMP 116 p. 16 | asserted, "similarly"; verified reduction to (W) + A1w + A7 + (An); $R_3$ in weaker forms |
| A3 | $C^*\Delta_k C \ge \gamma_0 > 0$, uniform in $k$ and $U$ | CMP 99 p. 428 | asserted, "we can prove"; verified reduction (P-sketch) to CMP 99 Thms 3.1, 3.3, 3.11–3.12 + A18, no $M$; $\sigma$-localized clause to (W) + (N) |
| A4 | Kernel decay (189) of $\delta^2 V/\delta A'^2$ and its consequence (190) | CMP 102 p. 308, Prop. 9 | announced; verified reduction to A1 + published CMP 102 + A18 + one bookkeeping item |

**Group A**: Bałaban-level statements and external results

| ID | statement | source | status |
|---|---|---|---|
| A5 | (H0): $E^{(j)}$, $\beta_{j+1}$ smooth in $g$ with bounded derivatives | CMP 109 pp. 263–264 | inductive assumption; "uniform in $j$" is our reading, uniformity in the history not printed; Theorem 2w uses only item 7′ (Lipschitz form on $]0,\gamma]$, every past), which also follows from (G-gain) with the running-maximum lift [verified (`SOURCES_GGAIN_NONMONO_VERIFY`)] |
| A6 | $E^{(j)}$, $\beta_j$ analytic in the couplings | CMP 109 pp. 264, 266 | asserted; not used |
| A7 | $\|C^*\Delta_k C\| \le \gamma_1$ and uniform decay of $C^*\Delta_k C$ | CMP 116 p. 13 (2.7); CMP 99 p. 428 | premise; reduced to A1 + A18 |
| A8 | Positive Gaussian measure for the real pair $(U,0)$; complex case "perturbative" | CMP 116 p. 15 | asserted |
| A10 | Lower-order bracket terms "bounded similarly as in (1.24)" | CMP 116 pp. 8–9 | asserted, "similarly" |
| A11 | "Similar inequalities hold for other constants" | CMP 119 p. 256 | asserted; the later inequalities not located |
| A13 | Theorem 1 | CMP 122 II p. 355; CMP 116, 119, 122 I–II | proved, conditional on $g_k \in ]0,\gamma]$; uses A1–A4, A1w, A7, A8, A10, A14, A15, A17, A18 |
| A14 | Induction closure recovers $E_0$ | CMP 109 p. 280 | asserted, constant unspecified; inside A13 only |
| A15 | Small-field terms and $\beta_j$ are whole-lattice quantities, independent of $\Lambda_{k+1}$ | CMP 119 p. 278, (2.24), p. 249; CMP 109 (4.35)–(4.37) | asserted, small-field part only; R-term part is B8 |
| A16 | Propagators may be replaced by operators with the same bounds | CMP 116 p. 3 | asserted for (1.3)–(1.4); our use for $C_f$ is an extension [I] |
| A17 | $\operatorname{supp}\chi_k$ lies in the analyticity domain of (1.34) | used in CMP 116 Lemma 1 | implicit; inside A13; C9 changes it |
| A18 | Backgrounds lie in the regular class of CMP 99/102; C-maps agree | used in CMP 109, 116, 122 I–II | implicit; pieces unsupported on the read pages |
| A19 | (190) at complex $B$ | CMP 102 Prop. 9 (silent) | implicit |
| A21 | Uniform-in-$k$ moments of the free extension kernel $E_0$ | not printed | implicit; numerical at $N = 3, 5$; alternative route only |
| A23 | Large-field T-step rate loss | CMP 119 Sect. 3 (defers to CMP 109 Sects. 6–7, CMP 116 (2.33)–(2.41)) | not printed; use consistent; used by B22 |
| X1 | Lüscher–Weisz (5.7) one-loop coefficient | Lüscher–Weisz [16] | cited; Route B only |
| X2 | OS II growth condition and reconstruction from torus data | OS II [20] | cited |
| X3 | Reflection positivity of the Wilson action, link planes | Osterwalder–Seiler [21] | cited, text not fetched |
| X4 | Convergence criterion of the final cluster expansion | Cammarota, CMP 85 (1982) | cited |
| X5 | (F): continuum log coefficient at constant field | Savvidy [22] | cited, source checked |

**Group B**: our own open items

| ID | statement | source | status |
|---|---|---|---|
| B1 | (H2) $\rho_2 < L$ | none | certified at $L = 13$ (Theorem 5.1); residue: reading of (0.4), gauge choice |
| B2 | (PI) probe independence | none | argued |
| B3 | (F) lattice coefficient = continuum coefficient | X5 | source checked; lattice transfer not proved |
| B4 | ($M_j$) formal = true $f^2$ coefficient, (S1)–(S3) | none | open; Prop. 4.4 argued |
| B5 | V1: $\beta_{j+1}(0)$ is the Maxwell coefficient | CMP 109 (1.3)–(1.5) | open; supported at order $p^2$ |
| B6 | (G2) per-cell infinite-volume densities | → A15 | formal |
| B7 | L5: $\beta_{k+1}$ on the clamped cube, or nested-interval Lemma D″ | CMP 109 p. 298 | open |
| B8 | $K$-independence of $\beta_j(g)$, R-term part: finite-volume step = whole-lattice step | → A15 | open; input of Theorem 1, not of the shooting (item 9 closed) |
| B9 | Route B inputs (T1), (DF), v4 "A6", `BETA_FP_U` Theorem 2.4 | none | argued; Route B only |
| B10 | Proved, unverified results in the chains (sweep, `BETA_FP_M` Prop. 4.3, Prop. 3.6, N3 in D″ form) | → A11 | proved, unverified |
| B11 | (B-S): sourced Theorem 1, (E14-a, b, c) | → A13 | open as a whole; parts B12–B20, B22 |
| B12 | (I): $B_j(c) = \beta_j(u(c))$ | none | by construction (C1, C10); not written as a lemma |
| B13 | Real uniform (H-hist) | → A1–A3, A1w, A7, A8, A10, A17, A18 | verified reduction, both halves (large field: `SOURCES_GGAIN_LF`); open inputs (Q)/$c_-$, ($M_r$), CMP 102 (189), (M-B) bookkeeping; G4 off-chain |
| B14 | (H-hist2) | none | verified reduction |
| B15 | (H-thr) threshold mismatch | CMP 122 II p. 383 | reduced to M2′, M4, M5 |
| B16 | (H-hol), small field | conditions (C1)–(C5) | verified reduction |
| B17 | (H-hol), large field | → A4, A18, A19 | verified reduction; (M-C) reduced to printed CMP 116 §2 conditions + Lemma 2; open (I-$\varepsilon$), (I-V0) residual, (M1) |
| B18 | N3 strip bound | none | proved+verified given B12–B17 |
| B19 | (FX) final Mayer expansion | not in CMP 116–122 | reduced; remaining input B18 |
| B20 | Sourced step bookkeeping: (W4), (P8), F2′, (D), $\Lambda_j$ | → A4, A10, A16 | reduced; $\Lambda_j$ bookkeeping open |
| B21 | (B-N1) bridge and (S) | none | (S) clauses: verified reduction, $h$-independent; (E14-c) clause open; ($S_{src}$) ⇒ (H-U) proved+verified; compatible with (U) (C23) |

**Group B**: our own open items

| ID | statement | source | status |
|---|---|---|---|
| B22 | Recovery of $\kappa \to \kappa/2$ in the sourced step (only; (U) moved to C23) | none | verified reduction [R] by $L/2$-rescaling, $\kappa \geq 1536$; uses A23 |

**Group C**: conventions and choices

| ID | choice | source | status |
|---|---|---|---|
| C1 | Freeze thresholds and partitions at Bałaban's trajectory $x^*$ | `SOURCES_D3`, `SOURCES_HTHR` | free; derivative needs B15 |
| C2 | Option B margin, $\theta_W + \tau_C \leq 0.2$ | `SOURCES_MB_VERIFY` | free; costs a smaller $\gamma$ |
| C3 | $L$ odd, $L > 11$; target $L = 13$ | v4 §1.1 | loses generality |
| C4 | $G = SU(2)$, $d = 4$ | Theorem 2 | loses generality |
| C5 | Order of constants | `SOURCES_N3A` | free; settled |
| C6 | Holomorphy radius $\rho_0$ | `SOURCES_HHOL` | free |
| C7 | Cut on $B$, not $B'$ | CMP 109 (2.9), (2.12) | reading; closed |
| C8 | Sharp $\chi_k$; smooth fallback | `SOURCES_G3` | free |
| C9 | Shrunk $\chi_k$ threshold | `SOURCES_GGAIN_VERIFY` | changes the construction; A13 must hold for it |
| C10 | Gauge-fixing weight $1/u_k(c)$ | `SOURCES_D2C_VERIFY` | changes the construction; A13 must hold for it |
| C11 | $h \equiv 0$ booking; freezing set | `SOURCES_D2C` | free |
| C12 | Additive source normalization | `SOURCES_HBETA_N3` | free |
| C13 | Main R-step form (1.90)–(1.98) | `SOURCES_FX_O23_VERIFY` | free; removes O4 |
| C14 | $\kappa_{\text{final}}$ fixed before $M$ | `SOURCES_FX_R2` | free; not sharp |
| C15 | $2p_1 - (d+5)r_0 > p_0$ | CMP 122 II p. 383 | condition (Bałaban's own) |
| C16 | $32\varepsilon' \leq 1$, $S \geq 1$, $\kappa_0 > 2$ | `SOURCES_D3W8_VERIFY` | changes the construction; A13 must hold for it |
| C17 | (S): $\gamma^{-2} \geq c_{sw} D'$ | v4 §1.2 | condition; needs $\beta_0$ fixed before $\gamma$ (verified reading) |
| C18 | $\ell$ even; thermal state; plaquette observable; test space $D_n$; origin at a site; compactness form | `ASSEMBLY` §0–§1 | loses generality |
| C19 | Coupling domain $]0, \gamma]$ | `SOURCES_GGAIN_VERIFY` | free |
| C20 | Route K primary, Route B alternative | v4 §1.3 | free |
| C21 | $\theta_* < 1$ in Route B | `BETA_FP_PW_VERIFY` | free; trivial |
| C22 | $g_0(K, g)$: any shooting solution | Lemma D | loses generality |
| C23 | (U): $\rho_1 \leq c_* r_0 / M^a$ at $\lambda_K = 1$ | `SOURCES_UN1` (+ VERIFY) | constant choice, not a hypothesis on test functions; verified compatible with B-N1; $C^n$ cost per $n$-point bound |

**Verified readings.** Four former Group A rows are now verified readings and appear in no set (`SOURCES_BOOKKEEPING.md`): A9 (the constant $C_1'(d, L)$ is unprinted and not derivable from the printed lines; its value is not used), A12 ($\beta_0$ "arbitrarily small" is on CMP 119 p. 255; $\beta_0$ is fixed before $\gamma$), A20 ($R_{K+1} \geq L$ follows from CMP 119 (2.5); $C_c$ is free of $g$), and A22 (Π (4.37)/(5.1), CMP 119 (2.42) corrected, (2.12)/(2.23), the "$L$ odd" page; the Sect. 3 residue is A23).

**Binding sets.** *Theorem 2w* (Theorem A(i), Route K) rests on A5, A15, B1–B7, and X5. Item 9 is closed, so B8 is no longer an input of the shooting. Item 7′ comes from A5 or from (G-gain) with the running-maximum lift [PROVED+VERIFIED]. Route B replaces B2–B5 by B9 and X5 by X1.

*Theorem 2w* ⇒ *Theorem 1* (Theorem A(ii)) rests on 4 Group A items (A5, A11, A13, A15), 9 Group B items (B1–B8, B10), and X5. Inside the proof of Theorem 1 (A13), Bałaban also uses A1–A4, A1w, A7, A8, A10, A14, and the items A15, A17, A18, which he uses without a separate proof.

*Theorem B* rests on 17 Group A items (A1–A5, A1w, A7, A8, A10, A11, A13, A15–A19, A23), 4 externals (X2–X5), and 21 Group B items (B1–B8, B10–B22). C9 does not remove A17, because A13 uses A17. Beyond A13, Theorem B adds only A16, A19, A23 to the Bałaban-level trust. Of the 17, A1–A4 are verified reductions to published theorems of CMP 99 and CMP 102; their joint residue is A1w, A7, the analyticity piece of A18, two unprinted conventions, a stated torus-side hypothesis, and the $G_1$ P-sketch.

Counts: Group A has 20 items (A1w added; A9, A12, A20, A22 moved to verified readings; A23 added in the previous pass). 14 are printed by Bałaban, and 13 of these are printed without a proof (11 of the 17 behind Theorem B, 3 of the 4 behind Theorem A); 4 of the 13 (A1–A4) now carry verified reductions. There are 5 externals, 22 Group B items, and 23 Group C items. In Group B, B13, B17, B21 ((S) clauses), and B22 are now verified reductions. Of Group C, C3, C4, C18, C22 lose generality, C23 ((U)) is a constant choice, and C9, C10, C16 change Bałaban's construction.

# A Lean formalization

The shooting lemmas of Section 3.4, the arithmetic of Theorem K and Theorem C1, and the bridge lemmas of the certifier are formalized in Lean 4 with Mathlib (21 files in `lean/YmLean/`); three files formalize lemmas of the sources block (Schur test, Combes–Thomas, interpolation and rescaling). The inventory is as follows.

[LEAN] (no sorry; axioms propext, Classical.choice, Quot.sound)

- Lemma D′ (a)–(d) (`YmLean.lemma_DPrime`), Step 8 (`lemma_DPrime_step8`), Lemma D, Lemma E ($Cg^2$ form), Lemma D+E (`lean/YmLean/LemmaDPrime.lean`, `LemmaDE.lean`). One-argument $\beta$ only.
- Lemma D″ (`YmLean.lemma_DDoublePrime`, `lean/YmLean/LemmaDDoublePrime.lean`, 146 lines): (a″) bracket $(3b/4)(K-k) - 2R \le T(k,K) \le (5b/4)(K-k) + 2R$; (b) shooting for $g \in ]0, g_1]$ with the clamp inactive; (c′) with $\beta = 3b/4$, $B' = 5b/4 + 2R$, $D' = 2R$, for every trajectory with couplings in $]0, \gamma]$; (d) for $n - m \ge 8R/b$, lower constant $b/2$. Hypotheses weaker than the note (continuity, any admissible $\gamma$, $R \ge 0$ derived). One-argument $\beta$ only; L5 not formalized.
- Full inventory (`lean/README.md`; file, main theorems, source note):
  1. `LemmaDPrime`: `lemma_DPrime`; `LEMMA_DPRIME.md` §1–§2.
  2. `LemmaDE`: `lemma_D`, `lemma_E`, `lemma_DE`, `lemma_DPrime_step8`; `LEMMAS.md`.
  3. `LemmaH`: `gronwall_telescope`, `lemma_H`; `SOURCES_HBETA.md` Lemma H.
  4. `DetIdentities`: `det_bordered`, `det_restrict_add_gauge`; `numerics/NUMERICS.md` §1.
  5. `Telescoping`: `telescoping_det`, `telescoping_log`; `R2_TELESCOPING.md` Prop. 3.1.
  6. `AbelianCount`: `su2_count`, `abelian_share`; `U1_INFINITE_LATTICE.md` Prop. 9.1 (used here only as a cross-check, Section 5.2).

7. `LemmaDDoublePrime`: `lemma_DDoublePrime`; `KAPPA_BOUNDED.md` §5.3.
8. `TheoremK`: `telescoping_bound`, `window_of_partial` (the window bound (5.1) from partial sums); `KAPPA_BOUNDED.md` §5.
9. `Gronwall`: `discrete_gronwall`, `contraction_bound` ($N_{j+1} \le qN_j + Z$); `KAPPA_BOUNDED.md` Prop. 4.1.
10. `CubeCount`: `cube_count_sharp`, `cube_count_lower` (Lemma O1-R′ arithmetic); `SOURCES_FX_O1.md`.
11. `BlockNorm`: `rho2_le`, the exact factorization $\rho_2 \le \rho_1(j)\max_l \rho_1(j-1,k_l)$; `H2_CERTIFY.md`.
12. `OishiRump`: `oishi_rump`, `oishi_rump_eta`, `frobenius_power_bound`; `H2_CERTIFY.md` §5.
13. `Envelope`: `envelope_nontrivial`, `envelope_not_scale_invariant` ((B-NT) in log form); `ASSEMBLY.md` §2.9.
14. `Divergence`: `profile_divergence`, `P4_divergence` (identity (D) of premise P4); `H2_CERTIFY_P4.md`.
15. `TheoremC1`: `c1_two_step`, `c1_kappa_bound`, `c1_window`; `B2_TWOSTEP.md` §2.
16. `StripBound`: `strip_bound_linear`, `strip_bound_quadratic` (the $\frac{3}{2}\tau$ bootstrap); `SOURCES_HBETA_N3.md` §4.
17. `CertBridge`: `cell_invertible`, `cell_sup_bound`, `vertex_bound` (interval-to-norm bridge of the cell certifier); `numerics/h1cert_cell.py`, `h1cert_cell2.py`.
18. `Tower`: `tower_bounds`, `I1_compress`, `majorant_core`, `lemmaT_core` (algebraic core of the one-step majorant, Theorem 5.1); `H1_ANALYTIC.md`.
19. `Schur`: `schur_test_weighted`, `schur_test` (weighted Schur test for the spectral norm); `SOURCES_LOC.md` §3.3, `SOURCES_S2_FORM.md` §2, `SOURCES_A3_SIGMA.md` 2.6.
20. `CombesThomas`: `norm_entry_le`, `coercive_inv`, `combes_thomas`, `combes_thomas_exp`, `combes_thomas_decay`, `combes_thomas_schur` (the proof structure of Lemma CT, the kernel-decay reduction of ledger A1); `SOURCES_A1_KERNEL.md` §7. The identification with the block operators, the kernel-to-row-sum step, (K)/(C), the rescaling, and the block-valued Schur test are not formalized (`lean/VERIFY_CombesThomas.md`).
21. `Interp`: `multilinear_convex`, `monomial_affine` (`SOURCES_A3_SIGMA.md` 3.2); `window_perturb`, `window_perturb_partial` (`SOURCES_T2_KUNIF.md` 5.2, chained to Theorem K); `rescale_exp`, `rescale_rate`, `K1_iff`, `K1_numeric` (`SOURCES_T2_KAPPA2.md`); `xexp_le`, `xexp_absorb` (`SOURCES_GGAIN_LF.md` §2).

Not formalized: interval arithmetic itself, the enclosure premises P1′–P4, the identity $\rho_1^2 L^2 = \lambda_{\max}(G)$, and the history form of the shooting lemmas.

# B Index of results by status

This appendix indexes the results of the paper by status tag; the tags match the verifier files. Here v1–v4 denote the successive versions `THEOREM2_PROOF.md`, `THEOREM2_PROOF_v2.md`, `THEOREM2_PROOF_v3.md`, and `THEOREM2_PROOF_v4.md` of the Theorem 2 notes. The statements that the main theorems take on trust are listed by item in the trust ledger `TRUST_LEDGER.md`, condensed in Section 6.7; ledger IDs are given below where an entry corresponds to a ledger item.

[PROVED+VERIFIED]

- Lemma D′ (a)–(d), $K_0$, Remark 4 (`LEMMA_DPRIME_VERIFY` grades 1, 2, 5).
- Lemma D (corrected interval), Lemma E ($Cg^2$ form), nested-interval variant of Lemma D (`LEMMAS_DE_VERIFY`). Lemma E $C^1$ form: trivial variant, same proof.

- Lemma D″ (tail-sum form, no $J$) (`KAPPA_BOUNDED_VERIFY` §6).
- Telescoping $S_k = \sum \beta_j(0)$ for the $p \to 0$ coefficient (`BETA_FP_VERIFY` item 1, corrected scope).
- Prop. 2.1 anchor-free identity; exact at the constant field (`BETA_FP_LEMMA43_VERIFY` item 1; `BETA_FP_H_VERIFY` item 1).
- $c = 11N^2/(12\pi^2)$, normalization chain; (F) source (Savvidy eq. (1.4); ledger X5) (`BETA_FP_LEMMA43_VERIFY` item 2; `BETA_FP_H_VERIFY` §4).
- KAPPA Prop. 1.1 as an identity given its fixed-level inputs; Lemma 2.1 (integration by parts); Prop. 2.2; level-independence of the degree $-4$ part; Theorem K and Theorem 2w as conditional reductions (`KAPPA_BOUNDED_VERIFY` (i)–(v)).
- $\sup_j |\sigma_j^{(2)}(0)| < \infty$; Prop. 4.1 algebra (`KAPPA_BOUNDED_VERIFY` (vi), reason for A4 failure corrected: obstruction = $\partial\sigma^{(1)}$ in $Q_j$ and in (W-2), §7 there).
- `B2_TWOSTEP` two-step recursion; Theorem C1 ((H2) $\Rightarrow$ ($C1\sigma$), factor $\rho_2^2/L^2$) and Theorem B2 ((H2) $\Rightarrow$ (B2), factor $\rho_2^2/L^4$), conditional on (H2); threshold $\rho_2 < L$ (no strip loss); (H2) weaker than (H″) at $n = 2$ (`B2_TWOSTEP_VERIFY` grades 1, 3, 4). Form of $Q_j$ by read-off (grade 8; `U3_LEMMA51` (3.3) not re-derived).
- `h2_scan` code structure, one-colour reduction, $H$ and $\Delta_{out}$ blocks, Gram identity, separable Prop. A form (`H2_LARGE_L_VERIFY` grade 1); tail-bound derivation (4.1)–(4.3) (grade 6, inputs corrected).
- Logic of Theorem B and Lemma 1.3 of `BETA_FP_PW`, as conditional implications (`BETA_FP_PW_VERIFY` §6 items 1–4 CONFIRMED: doubling identity and Lemma 1.3; Prop. 2.2 / Lemma 2.3; Theorems A, B; Lemma 4.1, conditional). Its inputs (U), (DF), (T1) are listed under [PROVED, UNVERIFIED] and [ARGUED].
- Prop. E evenness, Theorem W0/W, Theorem A3, Lemma A4, A0, Prop. A2 (`U3_AW_VERIFY` grades 1–4).
- U3 Theorem 4.1 + O5, Prop. A, Prop. B, O6, Theorem 6.1 (given V1; Taylor-data norm), Lemma 5.1 neutral value bound, single-block $m = 2$ failure in $d = 4$, one-step recursion, R2 midpoint identity (`U3_VERIFY`, `U3_REPAIRS_VERIFY`, `U3_LEMMA51_VERIFY`, `U3_LEMMA51_R23_VERIFY`).
- Exact one-step factorization $\rho_2 \le \rho_1(j) \max_l \rho_1(j-1, k_l)$ (`H2_CERTIFY_VERIFY`; Lean `rho2_le`).
- Sources block: Lemma G3 given (Q) (`SOURCES_G3_VERIFY`); LOC (`SOURCES_LOC_VERIFY`); (M-A) given (H−) and (M-A′) (`SOURCES_MA_VERIFY`, `SOURCES_MA2_VERIFY`); (M-B) with $p_0 \ge 17/(\gamma_0 A_1^2)$ (`SOURCES_MB_VERIFY`); condition (a) of Lemma N3 (`SOURCES_N3A_VERIFY`); the sourced large-field draft B-S.D3, with corrections (`SOURCES_D3_VERIFY`, `SOURCES_D3R_VERIFY`, `SOURCES_D3W8_VERIFY`); (B-N1) given (S) (`SOURCES_HHIST_N1_VERIFY`).
- Cesàro: given convergence, $\kappa_k = o(k) \Leftrightarrow \beta_\infty = c \log L$ (equivalence, not a weaker form) (`THEOREM2_PROOF_VERIFY` Point 4).

[PROVED, UNVERIFIED]

- `BETA_FP_U`, no verifier file: Theorem 2.4 ((U) $\Leftarrow$ (T1) + (H2)), Prop. 3.1 ($\Gamma^{ex} = \gamma_{rec} + \mathfrak{d}_F$), Prop. 3.2 ($\gamma_{rec}$ bounded), Corollary 4.1, Theorem 2w by the plane-wave route given (T1), (H2), (DF), LW (5.7), A6, (H0); the argued step "(H2) closes ($C1\sigma$)" of §3.5 there is superseded by `B2_TWOSTEP` Theorem C1 [PROVED+VERIFIED].
- `LEMMA_DPRIME_SWEEP`: 19 cross-scale uses covered by (b) + (c′) + (S) (4 of 19 table rows, 11, 12, 16, 18, verified on page images by `LEMMA_DPRIME_VERIFY` comparisons A–E).
- Step "Theorem 2w ⇒ Balaban's Theorem 1" (depends on the sweep and on L5 in the sharpened form of Section 3.4.5).
- Prop. 3.6 assembly with $C(1+j)^2 L^{-2j}$; neutral-sector $K$-derivative paragraph; $\sigma^{(1)}$ convergence sketch $jL^{-2j}$.
- Necessity of individual positivity for literal (0.31) at every $K$ (elementary).
- `BETA_FP_M` Prop. 4.3 ($\log|\det|$ bound in the N–O window, given (S1)–(S3)); no verifier file.
- Item 7′: the running-maximum lift of $C_\beta$ to non-monotone pasts (`SOURCES_GGAIN_NONMONO.md`); verified (`SOURCES_GGAIN_NONMONO_VERIFY`).
- Premise P4 of the one-step certificate (`H2_CERTIFY_P4`; identity (D) in Lean `P4_divergence`); the $L = 13$ certificate sizing after fix G1 (`H2_CERTIFY_L13_G1`).

[NUMERICAL]

- $\rho_1(\infty) = 1.760, \ldots, 2.0895$ at $L = 3, \ldots, 13$; at $L = 13$ the measured $\sup \rho_1^2 = 4.37$ (`H2_CERTIFY_L13`); the certified bound is below.

[CERTIFIED (INTERVAL ARITHMETIC, OUTWARD ROUNDING; VERIFIED)]

- (H2) at $L = 13$ (Theorem 5.1; ledger B1): $\rho_1(j,K)^2 \le 10 < 13$ for all $j \ge 1$ (including $\infty$) and all $K$, hence $\rho_2 \le 10 < 13$ by `rho2_le` (`H1_ANALYTIC`; `H1_ANALYTIC_VERIFY` V0–V9, `H1_ANALYTIC_VERIFY2`, `H2_CERTIFY_P5_VERIFY`). Residue as in Section 5.7.

[VERIFIED REDUCTION]

- (Q) to CMP 99 (3.132), ($E_U$), and LOC (`SOURCES_Q_VERIFY`, `SOURCES_CDC_VERIFY`); ($E_U$) to kernel-moment bounds Bałaban does not print (`SOURCES_EU_VERIFY`; the bound $K \le K_0(1 + \ldots)$ is a [DERIVED SKETCH]).
- N3 to (H-hol) + real (H-hist) + (H2-hist) (`SOURCES_HBETA_N3_VERIFY`).
- (H-hol), small field, to Bałaban's conditions with modified constants (`SOURCES_HHOL_VERIFY`); large field to (M-A), (M-B), (M-C), (H-hist2) (`SOURCES_HHOL_LF_VERIFY`); (M-D) to (H-hist) + (H-thr).
- (H−) to CMP 102 (189), an existing trust item (`SOURCES_MA2_VERIFY`).
- (H-hist2) to the real uniform (H-hist), no new input (`SOURCES_HHIST2_VERIFY`).

- Round 3 (ledger A1–A4, B13, B17, B21, B22): A1 kernel-decay form via Lemma CT to CMP 99 Thms 3.1, 3.3, 3.12 + (KE′) + ($H_Q$) + S2 (`SOURCES_QE_KE_HQ_VERIFY`, `SOURCES_S2_FORM_VERIFY`); A2 to (W) + A1w + A7 + (An) (`SOURCES_A2_R2_VERIFY`); A3 and its $\sigma$-localized clause (`SOURCES_A3_GAMMA0_VERIFY`, `SOURCES_A3_SIGMA_VERIFY`); A4 to A1 + CMP 102 (`SOURCES_A1_A4_VERIFY`); large-field (H-hist) (`SOURCES_GGAIN_LF_VERIFY`); (M-C) to the printed CMP 116 §2 conditions (`SOURCES_MC_VERIFY`); the (S) clauses of B21 (`SOURCES_B21_S_VERIFY`); the $\kappa \to \kappa/2$ recovery B22 (`SOURCES_T2_KAPPA2_VERIFY`).
- Item 9 ($K$-independence in the shooting) closed: $\beta_j$ is the whole-lattice coefficient by definition, Lemmas D′/D″ run at fixed $K$ (`SOURCES_T2_KUNIF_VERIFY`).

[INTERVAL ENCLOSURE, VERIFIED]

- Point enclosure $\rho_2(\infty, K^*) \in [1.9993, 2.1273] < 3$ at $L = 3$; P1 (loop-family coefficients exact rationals); Oishi–Rump bound on $\|\mathfrak{B}^{-1}\|$ (`H2_CERTIFY_VERIFY`).

[ARGUED]

- (PI); ($M_j$) (Prop. 4.4 of `BETA_FP_M`); (U) (now a constant choice, ledger C23; the $\kappa \to \kappa/2$ recovery B22 is a verified reduction); (T1) (plane-wave form of A3, `BETA_FP_U` §2.3); (DF) (`BETA_FP_U` §3.4); A6 (v4 label, not ledger A6) = plane-wave vs constrained-minimizer expansion point (uniform in $k$); $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ bounds; (H″) ⇒ (A) in full ⇒ (B2) (`U3_AW` §4.7 sketch); $C^2$, $C^3$ of $\sigma^{(1)}$ under (H2) (not written); $j_*$ size; the rate $O(\Lambda^{-2} \log \Lambda)$.

[OPEN]

- (H2) at $L \neq 13$ (numerical at $L = 3, 5, 9$; certified at $L = 13$, see above); (B2) is reduced to it [VERIFIED REDUCTION]. A1w, the walk-expansion form of A1 (CMP 99 Thm 3.15). The residual inputs of (M-C): (I-$\varepsilon$), (I-V0), (M1). (T1), (DF), A6 (v4 label) for Route B (ledger B9). V1. (S1)–(S3). L5 (domain of $\beta$; nested-interval D″ [NOT WRITTEN]). The finite-volume identification of the step, R-term part (ledger B8; an input of Theorem 1). The (E14-c) clause of B21. (A) in full. See Section 6.

[CORRECTED BY VERIFIERS SINCE V3]

- `B2_TWOSTEP` G1: the certification of (H2) is a computer-assisted proof project, not a check (`B2_TWOSTEP_VERIFY` grade 6).
- `H2_LARGE_L`: $L$-trend rate claim (grade 5); $j_0$ inputs, $\delta_m$ 45–90× larger, $a_*(L = 5) = 0.0107$ (grade 6), giving $j_0 \approx 8$ ($L = 3$), $\approx 6$ ($L = 5$).
- `BETA_FP_PW`: comparison and numerics (`BETA_FP_PW_VERIFY` grades 5, 6).

[REFUTED]

- (HS) $\sup \|H_j\| < L\sqrt{L-1}$ at $L = 3$.
- Not a refuted statement, but a failed method: the one-step sup-norm $C^1$ recursion at $L = 3$ (factor 1.005–1.033 for $j \geq 2$; Section 3.7.2).
- `U3_LEMMA51` Theorem 4.1 for $m = 2$ with a single block-$M$ constraint in $d = 4$.

- v1 claims withdrawn: "every early step must be certified for Theorem 1" (false: Lemma D′/D″); "(H′) is weaker than $\beta_\infty = c \log L$" (it is equivalent); "the whole $\log \Lambda$ sits in the fine piece" as a fact (it is equivalent to the open statement); "one-loop parts hold for SU($N$)" (Lemma 5.1's charged/neutral split is the SU(2) Cartan split); (3.2) with $Cj^2L^{-2j}$ (0 at $j = 0$); the (M) resolvent bound $\|\mathcal{K}_t^{-1}\| \le C/|f|$ (Nielsen–Olesen zero); (H) with IR weights as stated (UNSUPPORTED).

[Balaban]

- (H0) smoothness of $\beta_{j+1}$ in $g$ (CMP 109 (1.22), pp. 263–264), at odd $L > 11$ only; uniformity in $j$ is our reading and uniformity in the history is not printed (ledger A5); Theorem 1 and the inductive bounds of CMP 109, 116, 119, 122 and the propagator papers CMP 95, 96, 98, 99, 102. Uniform level bounds partly cited from CMP 95/96/99 (O6 replaced the main citation). Each Bałaban input that is printed without proof, or used without being printed, is a Group A row of the trust ledger.

[verified quote]

- Theorem 2 and Theorem 1 wording, "separate paper", p. 298 history sentence, (0.1) with $1/g_k^2$ and $\chi_k$, (1.3)–(1.5) (print misprint $E^{(j+k)}$ read as $E^{(j+1)}$), (2.6)–(2.9) of CMP 119, items A–E of CMP 122 (`BALABAN_EXCERPTS_CHECK`; `LEMMA_DPRIME_VERIFY`; `LEMMA_DPRIME_SWEEP` page images). `BALABAN_EXCERPTS_CHECK` items 10 (CMP 116 has no $C^m$-in-$g$ statement) and 11 (unchecked readings) are unchanged since v3.

[numerical] see Section 5.

# C Label map

The numbered environments of this paper carry the names used in the project files. The first table maps them. The second table maps the section numbers of `THEOREM2_PROOF_v4.md` ("v4") and `ASSEMBLY.md` to the sections of this paper; section references inside the project files use the old numbers.

| this paper | name in the project files | source |
|---|---|---|
| Theorem A | Route K summary: Theorem 2w ⇒ Theorem 1 | `THEOREM2_PROOF_v4` §6, §10 |
| Theorem B | Theorem 1.3 (summary) | `ASSEMBLY` §1.3 |
| Theorem 3.1 | Theorem 2w | `THEOREM2_PROOF_v4` §1.3 |
| Lemma 3.2 | Lemma D′ | `LEMMA_DPRIME`; v4 §2.1 |
| Remark 3.3 | Remark 4 | `LEMMA_DPRIME`; v4 §2.1 |
| Lemma 3.4 | Lemma D″ | `KAPPA_BOUNDED` §5.3; v4 §2.2 |
| Lemma 3.5 | Lemma D | `LEMMAS`; v4 §2.4 |
| Lemma 3.6 | Lemma E | `LEMMAS`; v4 §2.4 |
| Proposition 3.7 | Prop. 2.1 (anchor-free identity) | `BETA_FP_LEMMA43`; v4 §3.2 |
| Theorem 3.8 | Theorem K | `KAPPA_BOUNDED`; v4 §4.1 |
| Theorem 3.9 (Theorem $B_{PW}$) | Theorem B = Theorem 3.2 | `BETA_FP_PW`; v4 §4.2 |
| Theorem 3.10 | Theorem C1 | `B2_TWOSTEP`; v4 §5.3 |
| Theorem 3.11 | Theorem B2 | `B2_TWOSTEP`; v4 §5.3 |

| this paper | name in the project files | source |
|---|---|---|
| Definition 4.1 | (H-U) | `ASSEMBLY` §1.2 |
| Theorem 4.2 | Theorem 1.3 | `ASSEMBLY` §1.3 |
| Theorem 4.3 | Theorem 1.4 | `ASSEMBLY` §1.4 |
| Proposition 4.4 | Proposition of (B-C) | `ASSEMBLY` §2.4 |
| Lemma 4.5 | finite-form lemma of (B-RP) | `ASSEMBLY` §2.7 |
| (GC1)–(GC3) | (G1)–(G3) of the dependency table | `ASSEMBLY` §3 |
| ($S_{src}$) | (S), final sourced bound | `SOURCES_D4D5` |
| items H, 1–13, B1–B8 | items of the open list | `THEOREM2_PROOF_v4` §10 |

| old section | this paper |
|---|---|
| v4 §1.1 | Section 1.2 |
| v4 §1.2 | Section 3.1 |
| v4 §1.3 | Section 3.2 |
| v4 §1.4 | Section 3.3 |
| v4 §2 (notation) | Section 2.3 |
| v4 §2.1 | Section 3.4.1 |
| v4 §2.2 | Section 3.4.2 |
| v4 §2.3 | Section 3.4.3 |
| v4 §2.4 | Section 3.4.4 |
| v4 §2.4 (L5) | Section 3.4.5 |
| v4 §3.1 | Section 3.5.1 |
| v4 §3.2 | Section 3.5.2 |
| v4 §3.3 | Section 3.5.3 |
| v4 §3.4 | Section 3.5.4 |
| v4 §3.5 | Section 3.5.5 |
| v4 §4 | Section 3.6 |
| v4 §4.1 | Section 3.6.1 |
| v4 §4.2 | Section 3.6.2 |
| v4 §4.2a | Section 3.6.3 |
| v4 §4.3 | Section 3.6.4 |
| v4 §5 | Section 3.7 |
| v4 §5.1 | Section 3.7.1 |
| v4 §5.2 | Section 3.7.2 |
| v4 §5.3 | Section 3.7.3 |
| v4 §5.4 | Section 5.4 |
| v4 §5.5 (arithmetic) | Section 3.7.4 |
| v4 §5.5 (values) | Section 5.5 |
| v4 §5.6 | Section 5.8 |
| v4 §5.7 | Section 3.7.5 |
| v4 §6 | Section 3.8 |
| v4 §7 | Section 3.9 |
| v4 §8.1 | Section 5.1 |
| v4 §8.2 | Section 5.2 |
| v4 §8.3 | Section 5.3 |
| v4 §9 (Lean) | Appendix A |
| v4 §9 (other tags) | Appendix B |
| v4 §10 (open list) | Section 6 |
| v4 §10 (hypothesis table) | Section 4.6 |
| v4 §11 | Section 1.5 |
| v4 preamble (tag legend) | Section 1.4 |

| old section | this paper |
|---|---|
| v4 preamble (range of $L$) | Section 2.2 |
| v4 change logs | Appendix D |
| ASSEMBLY §0 (notation) | Section 2.1 |
| ASSEMBLY §0 (observables) | Section 2.5 |
| ASSEMBLY §1.1 | Section 4.1 |
| ASSEMBLY §1.2 | Section 4.2 |
| ASSEMBLY §1.3 | Section 4.3 |
| ASSEMBLY §1.4 | Section 4.4 |
| ASSEMBLY §2 | Section 4.5 |
| ASSEMBLY §2.1 | Section 4.5.1 |
| ASSEMBLY §2.2 | Section 4.5.2 |
| ASSEMBLY §2.3 | Section 4.5.3 |
| ASSEMBLY §2.4 | Section 4.5.4 |
| ASSEMBLY §2.5 | Section 4.5.5 |
| ASSEMBLY §2.6 | Section 4.5.6 |
| ASSEMBLY §2.7 | Section 4.5.7 |
| ASSEMBLY §2.8 | Section 4.5.8 |
| ASSEMBLY §2.9 | Section 4.5.9 |
| ASSEMBLY §3 | Section 4.6 |
| ASSEMBLY §4 | Section 4.7 |
| ASSEMBLY corrections | Section 4.8 |

# D Revision history

- *Theorem 2 notes, v3.* Built on THEOREM2_PROOF_v2.md with the corrections of THEOREM2_PROOF_v2_VERIFY (cited there as v2_VERIFY): the range of $L$ (odd, $L > 11$, or (H0) at this $L$ as a hypothesis); L5 sharpened (the clamped shooting leaves Bałaban's domain; the nested-interval variant of Lemma D$''$ is [NOT WRITTEN]); (H$''$) $\Rightarrow$ (B2) tagged [ARGUED]; "independent" routes replaced by "formally distinct", with the points $\theta_* < 1$, $k$-uniform $\beta$ identification, and the first-order vertex smoothness of (U); the independence record completed (ONLINE_SWEEP §2 as the first channel; the O(3) paper cited); open list extended (item 9, B5, item 10, hypothesis table); sweep count "4 of 19 table rows"; $\gamma_*$ depends on $R$ through $B'$. At that time BETA_FP_PW_VERIFY.md had no grade, and Route B stayed [PROVED, UNVERIFIED]. The grades of LEMMA_DPRIME_VERIFY and KAPPA_BOUNDED_VERIFY §9(iv) were cited explicitly.

- *Theorem 2 notes, v4.* Added the material verified after v3: B2_TWOSTEP.md with B2_TWOSTEP_VERIFY.md, H2_LARGE_L.md with H2_LARGE_L_VERIFY.md, lean/README.md (Lemma D$''$ in Lean), BETA_FP_U.md (Route B inputs), and numerics/NUMERICS.md §7. The section on (B2) was rewritten around (H2) (Theorems C1, B2; threshold $\rho_2 < L$; one-step sufficiency at $L \geq 5$; certification plan; corrected $L$-trend and $j_0$ inputs). The ledger and open list were updated, with item H as the single uniform hypothesis of both routes. KAPPA_BOUNDED_VERIFY states the (B2) obstruction in its §7 and §9(vi). BALABAN_EXCERPTS_CHECK had no change since v3.

- *Tag change in the LaTeX version of v4.* The logic of Theorem $\mathrm{B_{PW}}$ is tagged [PROVED+VERIFIED, CONDITIONAL], citing BETA_FP_PW_VERIFY §6 items 1–4, per the tag decision in PLAN.md (2026-10-28). The Markdown v4 kept [PROVED, UNVERIFIED]. The inputs (U), (DF), (T1) from BETA_FP_U stay [PROVED, UNVERIFIED].

- *Assembly.* `ASSEMBLY.md` with the corrections of `ASSEMBLY_VERIFY.md` (Section 4.8). The verifier also found the dependency table stale in five rows (grade 7; no overclaim): B-T2.2, B-T2.3, B-T2.4, B-S.D2C, B-S.D3 lagged `PLAN.md`. The table was rebuilt with a tag for every block and the sub-blocks added later (Route K, Route B, (H2), (H-thr), (G-gain), (FX)). Minor items: `SOURCES.md` §1.3 gives $c_j = 1 + O(g_j^4)$, while this paper uses $O(g^3)$ per `SOURCES_VERIFY` item 8 (the corrected value); the variance estimate drops a harmless $g_0^4$ (logarithmic) factor.

- *This version.* The Theorem 2 chapter and the assembly chapter are merged into one article: one notation section, one status legend, numbered environments (Appendix C), one numerical section, one list of open items, and the hypothesis table of v4 §10 merged into the dependency table. Three names were changed to avoid collisions (Section 2.6). The dependency table and the open list were brought to the current state of `PLAN.md`: (FX) is reduced to N3 (O2 and R2 [PROVED+VERIFIED], O3 [PROVED+VERIFIED, CONDITIONAL ON N3], O4 avoided by the main form; previously O2, O3 [PROVED, UNVERIFIED] and R2, O4 [OPEN]), and the point enclosure of (H2) at $L = 3$ is verified (Section 5.6). No theorem, lemma, or proof changed.

## References


[1] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. I. *Comm. Math. Phys.*, 95:17–40, 1984.

[2] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. II. *Comm. Math. Phys.*, 96:223–250, 1984.

[3] Tadeusz Bałaban. Averaging operations for lattice gauge theories. *Comm. Math. Phys.*, 98:17–51, 1985.

[4] Tadeusz Bałaban. Propagators for lattice gauge theories in a background field. *Comm. Math. Phys.*, 99:389–434, 1985.

[5] Tadeusz Bałaban. Spaces of regular gauge field configurations on a lattice and gauge fixing conditions. *Comm. Math. Phys.*, 99:75–102, 1985.

[6] Tadeusz Bałaban. Ultraviolet stability of three-dimensional lattice pure gauge field theories. *Comm. Math. Phys.*, 102:255–275, 1985.

[7] Tadeusz Bałaban. The variational problem and background fields in renormalization group method for lattice gauge theories. *Comm. Math. Phys.*, 102:277–309, 1985.

[8] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. I. generation of effective actions in a small field approximation and a coupling constant renormalization in four dimensions. *Comm. Math. Phys.*, 109:249–301, 1987.

[9] Tadeusz Bałaban. Convergent renormalization expansions for lattice gauge theories. *Comm. Math. Phys.*, 119:243–285, 1988.

[10] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. II. cluster expansions. *Comm. Math. Phys.*, 116:1–22, 1988.

[11] Tadeusz Bałaban. Large field renormalization. I. the basic step of the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:175–202, 1989.

[12] Tadeusz Bałaban. Large field renormalization. II. localization, exponentiation, and bounds for the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:355–392, 1989.

[13] Roger Dashen and David J. Gross. The relationship between lattice and continuum definitions of the gauge theory coupling. *Phys. Rev. D*, 23:2340–2348, 1981.

[14] Elvin. yang-mills: project repository (notes/ directory holds the project files cited as `FILE.md`), 2026. Local repository; no public remote.

[15] Arthur Jaffe and Edward Witten. Quantum Yang–Mills theory. In *The Millennium Prize Problems*, pages 129–152. Clay Mathematics Institute, American Mathematical Society, 2006. Official problem statement, Clay Mathematics Institute.

[16] Martin Lüscher and Peter Weisz. Background field technique and renormalization in lattice gauge theory. *Nucl. Phys. B*, 452:213–233, 1995.

[17] Paolo Menotti and Andrea Pelissetto. General proof of Osterwalder–Schrader positivity for the Wilson action. *Comm. Math. Phys.*, 113:369–373, 1987.

[18] OpenAI. The canonical massive continuum limit of the two-dimensional O(3) model. `https://github.com/openai/math`, family 215, preprints folder, 2026. October 4, 2026.

[19] Konrad Osterwalder and Robert Schrader. Axioms for Euclidean Green's functions. *Comm. Math. Phys.*, 31:83–112, 1973.

[20] Konrad Osterwalder and Robert Schrader. Axioms for Euclidean Green's functions. II. *Comm. Math. Phys.*, 42:281–305, 1975.

[21] Konrad Osterwalder and Erhard Seiler. Gauge field theories on a lattice. *Ann. Phys.*, 110:440–471, 1978.

[22] G. K. Savvidy. Infrared instability of the vacuum state of gauge theories and asymptotic freedom. *Phys. Lett. B*, 71:133–134, 1977.

[23] George Savvidy. From Heisenberg–Euler Lagrangian to the discovery of chromomagnetic gluon condensation. *Eur. Phys. J. C*, 80:165, 2020. arXiv:1910.00654.

[24] Hao Shen, Rongchan Zhu, and Xiangchan Zhu. A stochastic analysis approach to lattice Yang–Mills at strong coupling. *Comm. Math. Phys.*, 400(2):805–851, 2023.

[25] Ethan Sussman and Levent Alpöge. SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality. ym4_manuscript_zeroth_draft.pdf, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310, 2026. Draft dated 6 October 2026. Accessed 2026-10-27.

[26] Ethan Sussman and Levent Alpöge. The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0. LaTeX files theorem_0_complete_account_raw_{master,I,II,III}.tex, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310 (2026-10-27), 2026. Written largely by LLM, with Claude; not vetted by the repository authors. Accessed 2026-10-27.