Theorem 2 for lattice Yang–Mills theory: proof chapter, version 3

(shortest route; Route K reduction verified)

Written 2026-10-27/28

Contents

1 Statement **3**
- 1.1 Balaban's statements [verified quote] . . . 3
- 1.2 What Balaban's Theorem 1 actually needs (`LEMMA_DPRIME_SWEEP` §2, §7) . . . 3
- 1.3 The weak form proved conditionally in this chapter . . . 4
- 1.4 Literal (0.31) . . . 4

2 Shooting lemmas **4**
- 2.1 Lemma D′ (shooting without early positivity) [proved+verified; Lean] . . . 5
- 2.2 Lemma D″ (tail-sum form; no index J) [proved+verified] . . . 5
- 2.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [proved, unverified (sweep); 4 of 19 table rows verified] . . . 6
- 2.4 Lemma D and Lemma E [proved+verified; Lean] . . . 6

3 The exact structure of the one-loop sums **7**
- 3.1 Telescoping [proved+verified, corrected scope] . . . 7
- 3.2 Prop. 2.1, anchor-free identity [proved+verified] . . . 7
- 3.3 The constant c [proved+verified] . . . 8
- 3.4 The decomposition $S_k = ck \log L + \kappa_W^F - \kappa_k$ . . . 8
- 3.5 Level-uniform symbol facts used below [proved+verified] . . . 9

4 Bounded window sums: two formally distinct conditional routes **9**
- 4.1 Route K: Theorem K [proved+verified, conditional on (B2) and the fixed-level inputs] 9
- 4.2 Route B: Theorem B of `BETA_FP_PW` [proved, unverified] . . . 10
- 4.3 Comparison . . . 11

5 The hypothesis (B2) of Route K **12**
- 5.1 Proved . . . 12
- 5.2 Refuted . . . 12
- 5.3 Live routes and the precise remaining inequality . . . 12

6 Conclusion **13**

7 The convergence route: what it adds, and its status **14**
- 7.1 Components and status . . . 14
- 7.2 Early steps and j_* . . . 15

8 Numerics **15**
8.1 First increment $\beta_1(0)$ (`NUMERICS` §6–§7; `UMK_BETA1.md`) 15
8.2 Later increments ($L = 2$ only) 16
8.3 Quantities of the Route K proof 16

9 Ledger (tags match the verifier files) **16**

10 Open list (complete for Theorem 2w ⇒ Theorem 1) **18**

11 Citation and independence **21**

A Change log (v2 → v3) **22**

Preamble

Written 2026-10-27/28 (Opus). v3 = v2 (`THEOREM2_PROOF_v2.md`[1]) with the corrections of `THEOREM2_PROOF_v2_VERIFY.md` §A–§N applied. Nothing else is changed. It does not edit v1 or v2. It is not itself verified as a whole document. The change log is at the end (Appendix).

Tag legend.

- [proved+verified]: proved in a project file and confirmed by an independent verifier file.
- [proved+verified, conditional on X]: the implication is confirmed; X is open or assumed.
- [proved, unverified]: proved in a project file; no verifier has graded it yet.
- [Lean]: formalized in Lean 4 + Mathlib, no `sorry`, axioms propext, Classical.choice, Quot.sound only.
- [verified quote]: read on the page image of Balaban's paper (`reports/BALABAN_EXCERPTS_CHECK.md` or a verifier file).
- [numerical, reproduced]: floating-point result reproduced by a verifier. [numerical, run by the verifier]: run once, by a verifier. [numerical]: not reproduced. None is certified.
- [argued]: a plausibility argument, not a proof. [open]: no proof. [refuted]: shown false.
- [Balaban]: an input taken from Balaban's published papers, used as stated there.
- [not written]: a proof plan; no project file contains the proof.

Range of L. Balaban assumes L odd, $L > 11$, as a standing assumption for the whole of CMP 109 (pp. 251–252) [8]. His construction of $\beta_j(g)$ for $g > 0$ and his inductive assumption (H0) are therefore available only at odd $L > 11$. Every statement below that uses (H0) or $\beta_j(g)$ at $g > 0$ is stated for odd $L > 11$, or carries "(H0) and Balaban's construction at this L" as an explicit hypothesis. Statements about the one-loop quantities $\beta_j(0)$, S_k, and the symbols σ_j hold for odd $L \geq 3$.

[1] Project files named in this chapter live in the `notes/` directory of the `yang-mills` repository.

1 Statement

1.1 Balaban's statements [verified quote]

Theorem 2 (CMP 109 (1987), p. 259): "Let $d = 4$, $G = SU(2)$, and let γ be a sufficiently small positive constant, then for a sufficiently small positive g there exists a bare coupling constant $g_0 = g_0(\varepsilon, g)$ such that the sequence of the effective coupling constants g_k is contained in the interval $]0, \gamma]$, and $g_K = g$. Moreover, there exist constants β, β', $0 < \beta \leq \beta'$, such that g_k satisfy the inequality $1/g^2 + \beta \log(L^k\varepsilon)^{-1} \leq 1/g_k^2 \leq 1/g^2 + \beta' \log(L^k\varepsilon)^{-1}$. (0.31)" Balaban adds that the proof "will be given in a separate paper". It was never published (CMP 122 II p. 355, verbatim: "has not been published yet, so we have Theorem 1 with the assumption").

Theorem 1 (CMP 122 II p. 355): "If the sequence of the effective coupling constants is contained in an interval $]0, \gamma]$ with a sufficiently small positive γ, then the effective densities ρ_k have the form, and satisfy all the conditions and bounds, described in Sect. 2 [III]." The bound (0.1) there reads $\chi_k \exp[-(1/g_k^2)A(U_k) - E_-|T_\eta|] \leq \rho_k \leq \exp E_+|T_\eta|$ (corrected reading, `BALABAN_EXCERPTS_CHECK` item 3).

Setting. $\varepsilon = L^{-K}$, so $\log(L^k\varepsilon)^{-1} = (K-k)\log L$. $u_k := 1/g_k^2$. Recursion (CMP 109 (0.18)/(0.20)): $u_{k+1} = u_k - \beta_{k+1}(g_k)$. The increment β_{k+1} depends on the whole history $g_0, \ldots, g_k$ (CMP 109 p. 298, [verified quote]). $\beta_j(0)$ is the one-loop (Gaussian) increment; it is history-independent by definition. Balaban assumes L odd, $L > 11$, for the whole paper (CMP 109 pp. 251–252, OCR line 124, verbatim confirmed in `THEOREM2_PROOF_VERIFY` Step 6).

1.2 What Balaban's Theorem 1 actually needs (`LEMMA_DPRIME_SWEEP` §2, §7)

The quoted hypothesis of Theorem 1 is only $g_k \in]0, \gamma]$. The proof uses more. CMP 119 Theorem 1 (p. 262) assumes "(I.0.33)", a misnumbered reference to (0.31) of CMP 109. Its working form is CMP 119 p. 255–256 (2.6)–(2.9), for scales $m < n$:

- (2.6) $g_n \leq (1 + g_n^2\beta'(n-m))^{1/2} g_m$ and $g_m \leq (1+\beta_0) g_n$;
- (2.7) $(\log g_n^{-2})^p \leq (1+\beta_0)(\log g_m^{-2})^p$ and $(\log g_m^{-2})^p \leq (1 + g_n^2\beta'(n-m))^{\beta_0}(\log g_n^{-2})^p$;
- (2.8), (2.9) the same for $\varepsilon_j = g_j p_0(g_j)$ and R_j.

Here β_0 is Balaban's small constant (1/7 in CMP 122 II p. 389), not a coupling. [verified quote] for (2.6)–(2.9): page images, `LEMMA_DPRIME_SWEEP` §1; the sweep itself is single-author, see §2.3.

In u-variables these follow from two pairwise bounds:

- (P-up) $u_m \leq u_n + B'(n-m)$, with no additive constant;
- (P-low) $u_m \geq u_n + \beta(n-m) - D'$, with $\beta > 0$, $D' \geq 0$ fixed,

plus the smallness condition

- (S) $\gamma^{-2} \geq c_{sw} D'$, $c_{sw} = \max(2, 1/(1-(1+\beta_0)^{-2})) = 64/15$ at $\beta_0 = 1/7$, and $\gamma \leq \gamma_*(\beta_0, \beta, B', p_0, p_1, r, \nu, \kappa_0)$ (the conditions that literal (0.31) also needs). γ_* is free of the D'-ratio conditions, but it depends on R through B' (it contains $\gamma^2 B' \leq (1+\beta_0)^2 - 1$ with $B' = 5b/4 + 2R$). The order of constants in §6 (R before γ) handles this.

β_0 must be fixed before γ.

1.3 The weak form proved conditionally in this chapter

Theorem (Theorem 2w (weak form)). *Let $G = SU(2)$, $d = 4$, L odd, $L > 11$ (equivalently: L odd ≥ 3 with (H0) and Balaban's construction of $\beta_j(g)$ at this L as an explicit hypothesis). There are $\gamma > 0$, $g_1 \in]0, \gamma]$, $\beta > 0$, $B' \geq \beta$, $D' \geq 0$ such that for every $K \geq 1$ and every $g \in]0, g_1]$ there is a bare coupling $g_0 = g_0(K, g)$ with*

- *(b) $g_k \in]0, \gamma]$ for $0 \leq k \leq K$, and $g_K = g$;*
- *(c′) for all $0 \leq m \leq n \leq K$: $u_n + \beta(n - m) - D' \leq u_m \leq u_n + B'(n - m)$;*
- *(S) holds for γ, D'.*

Consequence: Balaban's (0.31) holds up to the additive constant D' on the lower side, with constants $\beta/\log L$ and $B'/\log L$. The two-sided bound fixes $1/g_k^2$ up to constants, as required.

Status: [proved+verified, conditional on (B2) and the fixed-level inputs of §4] by Route K (Theorem K + Lemma D″). Second route: [proved, unverified, conditional on (U), ($B\theta$), and the plane-wave defects] by Route B (`BETA_FP_PW` Theorem 3.2 + Lemma D″). Both routes also need (H0) at this L and L5 (§2.4). See §4.

1.4 Literal (0.31)

- (L1) Literal (0.31) for every $K \geq K_0$ and every $g \in]0, \gamma]$. Needs eventual uniform positivity: there are J and $\beta_* > 0$ with $\beta_j(0) \geq \beta_*$ for all $j > J$, plus $\sup_j |\beta_j(0)| < \infty$ (Lemma D′ (d), §2.1). [proved+verified; Lean] as an implication. Theorem K and Theorem B do NOT give eventual positivity: they bound window sums, not single increments. Eventual positivity follows from the convergence route (§7), which is open.
- (L2) Literal (0.31) for every $K \geq 1$. Needs $\inf_j \beta_j(0) > 0$ and $\sup_j \beta_j(0) < \infty$ (Lemmas D + E). If $\beta_j(0) \to \beta_\infty > 0$, then "every $\beta_j(0) > 0$" is equivalent to $\inf_j \beta_j(0) > 0$ (Lemma D′ Step 8). [proved+verified; Lean] as an implication. Inputs open.
- Literal (0.31) is not needed for Theorem 1: §1.2 shows the proof uses (c′) + (S). Literal (0.31) is used directly in one place only, the alternative choice of ε_0 in CMP 119 p. 248, which Balaban does not adopt.
- Necessity. Literal (0.31) at every $K \geq 1$ forces $\beta_K(0) \geq \beta \log L > 0$ for every K (window $k = K - 1$, let $g \to 0$ using (H0); so L odd > 11). [proved, unverified]; elementary; `THEOREM2_PROOF_VERIFY` Point 5(a) agrees. So certified early steps are needed only for (L2), not for Theorem 1.

2 Shooting lemmas

Notation: $u_k = 1/g_k^2$, $t = 1/g^2$, $T(k, K) := \sum_{i=k+1}^{K} \beta_i(g_{i-1})$. Clamp $\varphi(u) := \max(u, 1/\gamma^2)^{-1/2}$. Lemmas D′, D″, D, E are abstract statements about a sequence of functions β_j. Applied to Balaban's $\beta_j(g)$, they need L odd > 11 (or (H0) at this L) and L5.

2.1 Lemma D′ (shooting without early positivity) [proved+verified; Lean]

Sources: `LEMMA_DPRIME.md` §1–§2; `LEMMA_DPRIME_VERIFY.md` grades 1, 2, 5 (CONFIRMED). Step 7 of that file also holds grades 3, 4, 6, which are CORRECTED (Remark 1 overclaim, misreads in §5, ledger); this chapter uses none of them except through the sweep (§2.3). Lean: `lean/YmLean/LemmaDPrime.lean` (`YmLean.lemma_DPrime`, 405 lines), statement check `lean/VERIFY_LemmaDPrime.md`.

Lemma (Lemma D′)**.** Hypotheses. *(i)* β_{k+1} *continuous on* $[0,\gamma_0]$ *and* $|\beta_{k+1}(g)-\beta_{k+1}(0)|\le C_1 g$, C_1 *independent of* k, K. *(ii)* $\sup_j|\beta_j(0)|\le B$. *(iii′) there is* J *with* $\beta_j(0)\ge\beta_\infty/2$ *for all* $j>J$ *(this is all the proof uses of "*$\beta_j(0)\to\beta_\infty>0$*").*

Constants. $\gamma:=\min(\gamma_0,\gamma_{Bal},\beta_\infty/(4C_1))$; $B':=B+\beta_\infty/4$; $D:=J\cdot B'$; $D':=J(B+\beta_\infty/2)$; $g_1:=(1/\gamma^2+D)^{-1/2}$; $K_0:=J+\lceil 8D'/\beta_\infty\rceil$.

Conclusions.

- *(a)* $T(k,K)\ge -D$ *for all* $k<K$ *along any couplings in* $]0,\gamma]$.
- *(b) For every* $K\ge 1$ *and* $g\in]0,g_1]$ *there is* g_0 *with all* $g_k\in]0,\gamma]$ *and* $g_K=g$.
- *(c)* $u_K+(\beta_\infty/4)(K-k)-D'\le u_k\le u_K+B'(K-k)$. *The upper side has no constant.*
- *(d) For* $K\ge K_0$*: literal (0.31) with lower constant* $\beta_\infty/8$*, and shooting for every* $g\in]0,\gamma]$. *If every* $\beta_j(0)>0$*: literal (0.31) for all* K *(Step 8 = Lemmas D + E).*

Proof idea. Clamped flow $\tilde u_{k+1}=\tilde u_k-\beta_{k+1}(\varphi(\tilde u_k))$ is continuous in u_0. Its tail sums lie in $[-D,B'(K-k)]$ for every u_0, because clamped couplings lie in $]0,\gamma]$. IVT on $[t-D,t+B'K]$ gives $\tilde u_K=t$. Then $\tilde u_k\ge t-D\ge 1/\gamma^2$ for $g\le g_1$, so the clamp is never active. The bound is derived for the clamped flow first; there is no circularity. For history-dependent β this proof needs β_{k+1} defined and continuous on the whole cube $]0,\gamma]^{k+1}$; see L5 in §2.4.

Lean. The Lean hypotheses are weaker (ContinuousOn instead of C^1; (iii′) instead of convergence). Step 8 is `lemma_DPrime_step8` (`lean/YmLean/LemmaDE.lean`). Lean covers one-argument $\beta_j(g)$ only, not the history form.

Remark (Remark 4). Without $g\le g_1$, (b) can fail at small K ($\beta_1=\beta_2=\beta_3=-0.8$, $K=3$, $g=\gamma$). [proved+verified]

2.2 Lemma D″ (tail-sum form; no index J) [proved+verified]

Source: `KAPPA_BOUNDED.md` §5.3; `KAPPA_BOUNDED_VERIFY` §6 (CONFIRMED: "No J needed"); hand rederivation in `THEOREM2_PROOF_v2_VERIFY` §A.

Lemma (Lemma D″)**.** Hypotheses. *(i) as in 2.1, and the window bound*

$$\textit{(5.1)}\quad T_0(k,K):=\sum_{i=k+1}^{K}\beta_i(0)\in[\,b(K-k)-2R,\ b(K-k)+2R\,]\quad \textit{for all } 0\le k<K,$$

with $b:=c\log L>0$ *and some* $R\ge 0$. *(Route K gives* $R=\kappa_*$*; Route B gives* $R=C_B$*.)*

Constants. $\gamma:=\min(\gamma_0,\gamma_{Bal},b/(4C_1))$; $\beta:=3b/4$; $B':=5b/4+2R$; $D=D':=2R$; $g_1:=(1/\gamma^2+D)^{-1/2}$.

Conclusions. *(b) of 2.1, and the pairwise bound*

$$\textit{(c′)}\quad u_n+\beta(n-m)-D'\le u_m\le u_n+B'(n-m),\quad 0\le m\le n\le K,$$

upper side without constant.

Proof. $|T - T_0| \le C_1\gamma(K-k) \le (b/4)(K-k)$. Add (5.1). Steps 2–5 of Lemma D′ use only continuity and the bracket $\tilde{T} \in [-D, B'(K-k)]$; this bracket now comes from (5.1). The pairwise form is the same window argument with $K \to n$. The proof is the clamped-flow proof of 2.1; for history-dependent β the L5 caveat of §2.4 applies.

Not in Lean (the Lean lemma has hypothesis (iii′), not (5.1)). The reduction is short; formalizing it is a small task. [open as Lean work]

Literal (0.31) from (5.1) alone holds only for windows with $n - m \ge 8R/b$; short windows near K are not controlled (see §1.4).

2.3 Lemma D″ meets Balaban's proof: the cross-scale sweep [proved, unverified (sweep); 4 of 19 table rows verified]

Source: `LEMMA_DPRIME_SWEEP.md` (all 12 Balaban files; OCR grep plus page image for every non-single-scale hit). `LEMMA_DPRIME_VERIFY` Steps 4–5 checked comparisons A–E on page images (CMP 122 I p. 192, p. 198 (1.94); II p. 381 ×2, p. 386) and corrected two misreadings of A (factor $(1+\beta_0)^2(k-j)^{\beta_0}$; $\delta'_j = g_j A_1 p_1(g_j)$). The five comparisons map to four table rows: A = row 11, E = row 12, C and D = row 16, B = row 18.

Result. 19 cross-scale uses of the coupling sequence (table in SWEEP §7), all in CMP 119 and CMP 122 I–II; none in CMP 95, 96, 98, 99, 102, 109, 116. Each is covered by (b) + (c′) + (S) of §1.2:

- items 1, 7, 10, 12, 17 need (P-up) with no constant (e.g. (2.6) line 1 is exactly (P-up));
- items 2, 3, 11, 13, 15, 16, 18 need (P-low) with fixed multiplicative slack $(1+\beta_0)$ or $(1+\beta_0)^2$; item 2 ((2.6) line 2) gives the constant 64/15; item 11 needs only $2.42D'$;
- items 4 ((2.8)–(2.9), ε and R ratios), 6 (CMP 119 (3.7), (3.8), (3.12), (3.13)), and 9 (CMP 122 I (1.52)), the last two of the form $\varepsilon_{k+1} \le (1+\beta_0)\varepsilon_k$, use both sides; they follow as items 1 and 3 do (item 4 from items 1–3);
- item 14 (CMP 122 I p. 200, R_{k-N_0+1} vs R_k) uses the lower side; it follows as item 3 does;
- item 5 (CMP 119 (2.31), (2.46): $\sum_j g_j^{\kappa_0} \le g_n^{\kappa_0-6}$) needs linear growth (P-low) and $\gamma^{-2} \ge 2D'$;
- item 0 is CMP 119 Theorem 1 assuming (0.31); its proof works with (2.6)–(2.9);
- item 8 (CMP 119 p. 284) is in a sketch, not load-bearing.

Caveats [open]: (1) the sweep is a grep plus page images, not a line-by-line read; (2) "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) was not located; (3) Balaban says β_0 "can be chosen arbitrarily small, if g is sufficiently small"; under (P-low) the floor is $\beta_0 \gtrsim D\gamma^2/2$, so β_0 must be fixed first; no step was found that needs $\beta_0 \to 0$; (4) the sweep has no independent verifier beyond rows 11, 12, 16, 18; (5) `BALABAN_EXCERPTS_CHECK` item 11 lists readings not checked on page images: II (4.37)/(5.1), CMP 119 (2.42), the (2.12)/(2.23) ambiguity, the "L odd" page, the Sect. 3 estimates, and the 1989 bound lists.

2.4 Lemma D and Lemma E [proved+verified; Lean]

Sources: `LEMMAS.md` (corrected), `LEMMAS_DE_VERIFY.md` §1.4, §1.5, §2.4; `lean/YmLean/LemmaDE.lean` (`lemma_D`, `lemma_E`, `lemma_DE`).

- **Lemma D.** If $b \le \beta_j(g) \le b'$ on $]0,\gamma]$ with $b > 0$, then for every K and $g \in]0,\gamma]$ shooting succeeds and $t + b(K-k) \le u_k \le t + b'(K-k)$. This is literal (0.31). The first IVT interval in the draft was wrong and was refuted by the verifier; the corrected interval is $[t + bK,\ t + b'K]$. The nested-interval variant for β defined only on admissible histories (`LEMMAS_DE_VERIFY` §1.5) is [proved+verified], not in Lean.

- **Lemma E.** If $\beta_0 \le \beta_j(0) \le \beta_0'$ and $|\beta_j(g) - \beta_j(0)| \le Cg^2$, then for $\gamma \le \sqrt{\beta_0/(2C)}$: $\beta_0/2 \le \beta_j(g) \le \beta_0' + \beta_0/2$. The verifier checked this Cg^2 form; the C^1 form ($C_1 g$, $\gamma \le \beta_0/(2C_1)$) is the same two-line proof [proved, trivial variant]. Lean has the Cg^2 form.

- **History dependence (ledger item L5)** [open]. Balaban's β_{k+1} depends on $(g_0, \dots, g_k)$. Lemma D′ Remark 3 states the cube-continuity version (same proof). The gap is sharper than v2 stated (`THEOREM2_PROOF_v2_VERIFY` §F):

  - *Domain.* Balaban defines β_{k+1} only on histories for which his Theorem 1 induction has run to step k (`LEMMAS_DE_VERIFY` §2 item 1). By the sweep (§2.3), that induction uses (2.6)–(2.9), that is (c′) + (S), on the history, not only $g_i \in]0,\gamma]$.
  - *The new gap.* The clamped shooting of §2.1–§2.2 evaluates β on clamped histories $\tilde{g}_i = \varphi(\tilde{u}_i)$. A clamped history can violate the growth side of (c′): from $\max(a,c) - \max(b,c) \ge \min(a-b, 0)$ one gets only $u_m \ge u_n - D'$, not $u_m \ge u_n + \beta(n-m) - D'$. So the clamped shooting evaluates β on histories outside Balaban's domain. The clamp proof is valid only under the cube hypothesis, which Balaban does not supply.
  - *A variant that avoids it* [not written]. Use a nested-interval proof in the style of `LEMMAS_DE_VERIFY` §1.5, with domain $D_k = \{u_0, \dots, u_k \ge \gamma^{-2}\}$ of true (unclamped) trajectories. On such trajectories, (c′) up to step k follows from the tail-sum bound (a″) of Lemma D″, and (S) holds by the choice of γ. So every evaluation of β stays inside Balaban's domain. At each step k, keep the set of u_0 whose trajectory stays in D_k; show by IVT, step by step, that this set contains a closed interval on which u_k sweeps a range that contains the next target interval. For Lemma D (all increments positive) this is written. For Lemma D″ the increments can be negative, so the nested intervals need new envelopes, built from the window bounds $[b(n-m) - 2R - (b/4)(n-m), \dots]$. The adaptation is plausible; it is not written and not verified.
  - Lean is one-argument only.

3 The exact structure of the one-loop sums

3.1 Telescoping [proved+verified, corrected scope]

$S_k := p \to 0$ F^2-coefficient of the composed k-step gauge-fixed one-loop integral (U1 (1.2)). Then $S_k = \sum_{j=1}^{k} \beta_j(0)$, $S_0 = 0$ (`BETA_FP` Prop. 1.1; `BETA_FP_VERIFY` item 1). Scope: exact for the $p \to 0$ coefficient at fixed k, not for $\log Z_k$ at fixed p (the plane wave is off-shell).

3.2 Prop. 2.1, anchor-free identity [proved+verified]

Source: `BETA_FP_LEMMA43` Prop. 2.1; `BETA_FP_LEMMA43_VERIFY` items 1, 3–4 (hand rederivation; lattice check with the real CMP 109 averaging at random SU(2) backgrounds, relative difference 10^{-14}).

Under (G1) = $\{\Delta D = 0,\ \tilde{Q}D = D_c \iota_a^T\}$:

$$\log Z_\Lambda(V) = \big[\log\det X - \tfrac{1}{2}\log\det K\big] - \big[\log\det(D_c^T D_c) - \tfrac{1}{2}\log\det(\Delta_\Lambda(V) + D_c D_c^T)\big].$$

In words: constrained one-loop = fine Wilson one-loop in background Feynman gauge minus the one-loop of the coarse Gaussian theory with kernel $\Delta_\Lambda(V)$. All point-anchor objects cancel; the constant is exactly 0.

- $\tilde{Q}D = D_c\iota_a^T$ is exact at any background (lattice 10^{-15}). $\Delta D = 0$ is load-bearing (raw Hessian gives $O(1)$ errors).
- At the constant abelian field V_f the background is critical, so $\Delta D = 0$ is exact and Prop. 2.1 holds exactly (`BETA_FP_H_VERIFY` §2, CONFIRMED). At a plane wave it holds up to the defect $\mathfrak{d}_Y$ (`BETA_FP_PW` Lemma 2.3) [proved, unverified].
- Per-cell infinite-volume densities only: on a torus the constant zero modes must be removed. This is input (G2) [formal].

3.3 The constant c [proved+verified]

$c = 11N^2/(12\pi^2) = 2N\cdot b_0/(8\pi^2)$, $b_0 = 11N/3$; SU(2): $c = 0.371511$. Normalization $1/g_B^2 = 2N/g^2$ (`R2_VERIFY`; `BETA_FP_LEMMA43_VERIFY` §5). The log coefficient of the fine piece at constant field is $c_2 = c/2$ per $\ln(1/\varphi)$, checked against Savvidy, arXiv:1910.00654 eq. (1.4) (`BETA_FP_H_VERIFY` §4) = input (F) [cited, source checked; lattice = continuum log coefficient is standard, not proved in project files].

$b := c\log L$: 0.25751 ($L = 2$), 0.40815 (3), 0.51502 (4), 0.59793 (5), 0.72293 (7).

3.4 The decomposition $S_k = ck\log L + \kappa_W^F - \kappa_k$

- As bookkeeping the split is empty: any c' gives an exact identity if κ_k is defined as the remainder (`U1_VERIFY`). Boundedness of a remainder-defined κ_k is a restatement, not progress.
- Content comes from defining κ_k by the coarse theory alone. `KAPPA_BOUNDED` Prop. 1.1 [proved+verified, conditional on (PI), (F), (M_j), V1, (G2)]: with the Wilson level $j = 0$ as reference,

$$S_j = cj\log L - d_j, \qquad \kappa_j = \kappa_W^F + d_j,$$

$$d_j := -\frac{1}{2n_2}\int_0^1 dt \int_{\mathrm{BZ}} \frac{dK}{(2\pi)^4}\ \mathrm{tr}\big[b_{t,2}\,\delta\sigma_j^{(0)} + b_{t,1}\,\delta\sigma_j^{(1)} + b_{t,0}\,\delta\sigma_j^{(2)}\big],$$

  where $\delta\sigma_j^{(m)} = \sigma_j^{(m)} - \sigma_0^{(m)}$ are differences of the f^m Taylor symbols of the coarse kernel at level j and level 0, and $b_{t,m}$ are the Moyal inverse coefficients of the interpolated coarse operator. No limit symbols σ_∞ are used.
- Checks [numerical, reproduced]: $\rho(1) = \kappa_W^F$ (Lüscher–Weisz (5.7) [16]); $L = 2$: $d_1 = -0.372$, $\kappa_1 = 1.114$; $L = 3$: $d_1 = c\log 3 - \beta_1 = -0.256 \pm 0.01$ (`KAPPA_BOUNDED_VERIFY` §4: fits give $-0.254\cdots-0.258$). This checks the definition of d_j through S_j; it does not evaluate (1.1) from the symbols independently.
- Cor. 3.1 of `BETA_FP_LEMMA43` ($\mathcal{C}(\Lambda) = c\log\Lambda + \kappa_W^F - \rho(\Lambda)$) is exact as an identity. "The whole $\log\Lambda$ sits in the fine piece" is equivalent to $\rho(\Lambda) = o(\log\Lambda)$. It is not a separate fact (`BETA_FP_LEMMA43_VERIFY` item 2; v1 overclaim removed).

- Given convergence $\beta_j(0) \to \beta_\infty$, Cesàro gives $\beta_\infty = c \log L - \lim \kappa_k/k$. So v1's (H′) "$\kappa_k = o(k)$" is EQUIVALENT to $\beta_\infty = c \log L$, not weaker (`THEOREM2_PROOF_VERIFY` Point 4). Theorem K (§4) gives κ_k bounded, hence $\kappa_k = o(k)$ trivially.

3.5 Level-uniform symbol facts used below [proved+verified]

Source: `U3_AW.md`, `U3_AW_VERIFY.md` (grades 1–4 CONFIRMED); `U3_REPAIRS_VERIFY`.

- O5/A0: Δ_M real-analytic on a fixed strip, $|\partial^\alpha(\Delta_M - \Delta_\infty)| \le CM^{-2}$, uniform to $K = 0$ (odd L, centered blocks). Prop. B: $c_0|\hat{K}|^{-2}P_T' \le P_T' S_M P_T' \le C_1|\hat{K}|^{-2}P_T'$, $c_0 = (2/\pi)^{12}$, $C_1 \le 1363$. Existential constants.

- Prop. E (evenness): $\sigma_j(f; -K) = \sigma_j(f; K)$ in the midpoint trivialization, all f, j. Tested in $d = 2$ only.

- Theorem W: $\sigma_j^{(1)}(0) = \tau_c = (3ie/2)(E_{10} - E_{01})$ for every $j \ge 0$, from the exact Ward identity $\Delta_j(f)D_f = 0$. Numerically 0.750000, 0.75015, 0.75051 at $j = 0, 1, 2$ vs 0.75 ($d = 2$ proxy).

- Theorem A3: $|\sigma_j^{(1)}(K) - \tau_c| \le C_A|K|^2$, real K, uniform in j (C^0 only).

- Prop. A2: $\sigma_j^{(m)}$, $m \le 2$, analytic on the same strip at every fixed j (not uniform in j for $m = 1$ derivatives, $m = 2$).

- Lemma M: Maxwell coefficient exactly 1 at every level.

4 Bounded window sums: two formally distinct conditional routes

Both routes give the window bound (5.1) of §2.2, $|S_K - S_k - b(K-k)| \le 2R$, with $b = c \log L$. Lemma D″ then gives Theorem 2w (at odd $L > 11$, or with (H0) at this L as a hypothesis, and with L5). Neither route uses convergence of $\beta_j(0)$, a rate, (N), or certified early steps. The two routes have formally distinct named hypotheses. Their mathematical content is not shown to be independent; both plausibly need a uniform first-order smoothness of the level-j vertex (`BETA_FP_PW_VERIFY` §3; §4.3 below).

4.1 Route K: Theorem K [proved+verified, conditional on (B2) and the fixed-level inputs]

Source: `KAPPA_BOUNDED.md`; `KAPPA_BOUNDED_VERIFY.md` (items (i)–(v) and (vii) CONFIRMED; (vi) CORRECTED in reasoning, conclusion unchanged).

Theorem (Theorem K)**.** *L odd ≥ 3. Assume (B2) (§5) and the fixed-level inputs below. Then $\sup_k |d_k| \le \kappa_* < \infty$, so $\sup_k |\kappa_k| \le |\kappa_W^F| + \kappa_*$ and $|S_k - ck \log L| \le \kappa_*$ for all k. Window sums satisfy (5.1) with $R = \kappa_*$. (Theorem K is a statement about one-loop quantities. Its use in Lemma D″ for Balaban's $\beta_j(g)$ needs L odd > 11 or (H0) at this L.)*

Proof structure.

1. Prop. 1.1 (§3.4): $S_j = cj \log L - d_j$. [proved+verified, conditional on (PI), (F), (M_j), V1, (G2)]

2. Lemma 2.1: the only K-derivative of $\sigma^{(1)}$ in d_j enters linearly with f^0 coefficients. Integration by parts moves it onto f^0 data; boundary terms vanish (periodic up to a common constant conjugation; inner sphere $O(r^2)$). So $\sigma^{(1)}$ is needed only in C^0, which Theorem A3 gives. [proved+verified]

3. Prop. 2.2: with (E0) f^0 bounds (Prop. B, O5/A0, Lemma M), (E1) Theorem A3 + Theorem W, and (B2), every term of the d_j integrand is $\le C|K|^{-2}$, uniformly in j and t. In $d = 4$ this has two powers of infrared margin. $|d_j| \le \kappa_* := (C(L)/n_2)(1 + Y_1^2 + C_A(1 + Y_1) + Y_2) \cdot I_2$. [proved+verified, conditional on (B2)]

4. Infrared: the degree -4 part of the coarse f^2 integrand is the same at every level (Lemma M + Theorem W), so the $\tau_c \log$ cancels exactly in level differences. [proved+verified]

Constants are existential; κ_* is not evaluated.

Fixed-level inputs of Route K (needed at each j separately; no uniformity in j):

- (PI) probe independence: plane-wave p^2 coefficient = constant-field f^2 coefficient [argued; per-cell limit for the linear potential not written].
- (F) fine log coefficient c_2 at constant field [cited, source checked, §3.3].
- (M_j) the formal f^2 coefficient d_j equals the true one at fixed j. The resolvent step of the first argument was false (Nielsen–Olesen zero at $k_\perp^2 = |ef|$; `BETA_FP_H_VERIFY` §5). `BETA_FP_M` repairs it with a direct $\log|\det|$ bound given spectral bounds (S1)–(S3) [open on the lattice]; Prop. 4.4 there is [argued]; `BETA_FP_M` has no verifier file. The t-subdivision of KAPPA §1.3 removes the small-ε requirement [argued, CONFIRMED as plausible]. Status: [argued].
- V1: the Maxwell coefficient of the per-step determinant equals $\beta_{j+1}(0)$, i.e. order-g^0 completeness of CMP 109 (1.3)–(1.5) [open; supported at order p^2 by `U3_VERIFY` §6].
- (G2) per-cell infinite-volume densities [formal].
- (H0) Balaban's inductive assumption: β_{j+1} smooth on $[0, \gamma]$, "uniformly bounded on this interval together with all derivatives" (CMP 109 pp. 263–264, (1.22)) [Balaban; available at odd $L > 11$ only]. CMP 116 has no theorem statement and no C^m-in-g bound in words (`BALABAN_EXCERPTS_CHECK` item 10); the smoothness in g is CMP 109's inductive assumption, asserted by CMP 109 Theorem 3, whose proof CMP 116 completes.
- L5 history-dependent β, its domain, and its continuity (§2.4) [open].

4.2 Route B: Theorem B of `BETA_FP_PW` [proved, unverified]

Source: `BETA_FP_PW.md` §1–§5 (Theorem 3.2 = Theorem B, Theorem 5.1). `BETA_FP_PW_VERIFY.md` (checked 2026-10-27) has §1–§5 and no final grade. It rederives the Lemma 1.3 arithmetic and the Lemma 2.3 / Prop. 2.2 algebra correctly, confirms the Prop. 2.2 determinant identity on a 4^4 lattice (4e−13), and notes that identity 3.0 is a tautology ($\Gamma^{ex} := \hat{\Pi} - S$), so all content sits in $(B\theta)/(C\theta)$. The tag stays [proved, unverified].

Theorem (Theorem B)**.** *L odd ≥ 3. Assume*

- *(U) there are $p_* > 0$, C_U with $|\beta_{j+1}(p) - \beta_{j+1}(0)| \le C_U p^2$ for all $j \ge 0$, $0 < p \le p_*$* [HYPOTHESIS; ARGUED FROM UNIFORM ANALYTICITY IN p OF THE ONE-STEP PLANE-WAVE DATA PLUS ONE-STEP O6]*;*

- *($B\theta$) there is $\theta_* \in (0, \min(p_*, 1))$ with $B_* := \sup_k |\Gamma^{ex}(\theta_*; L^k)| < \infty$, where Γ^{ex} is the coarse remainder at fixed coarse momentum θ_** [HYPOTHESIS]*;*

- *Lüscher–Weisz (5.7) [16]* [CITED].

Then $|S_k - ck \log L| \le C_B := B_* + c_2 \ln(1/\theta_*^2) + |\kappa_W^F| + \sup_{p \le \theta_*} |\eta(p)| + C_U \theta_*^2/(L^2 - 1)$, and window sums satisfy (5.1) with $R = C_B$. $\theta_* < 1$ is required so that $c_2 \ln(1/\theta_*^2) \ge 0$; otherwise the term must be replaced by its absolute value (`BETA_FP_PW_VERIFY` §3).

Key step. Lemma 1.3: $|S(\theta; L^k) - S_k| \le C_U \theta^2/(L^2 - 1)$ from (U). The massless coarse side is then needed only at a fixed $\theta_* > 0$, where there is no infrared problem. No (W), (PI), (M), IR weights, or spectral theory.

What ($B\theta$) reduces to (`BETA_FP_PW` §4) [argued]: Lemma 4.1(i) (trivial-background convergence, from O5, odd L) [proved]; vertex bounds at fixed θ_* (the uniform bounds cited by U3 Theorem 7.1 to Balaban; the fixed-θ data hypothesis ($D\theta$) is needed only in its "bounded" form); and boundedness in k of the defect $\mathfrak{d} = \mathfrak{d}_Y + \mathfrak{d}_{gf}$.

- $\mathfrak{d}_Y$ (Prop. 2.1 defect at the non-stationary plane wave): $O(a\Lambda^{-2})$ relative at fixed θ, from zero capacity of point anchors in $d = 4$ [argued; region $|K| \lesssim \Lambda^{-1}$ not written].

- $\mathfrak{d}_{gf}$ (composite gauge vs D_k^T gauge at the plane wave): $O(\theta^2)$ relative; uniformity in k not shown [argued]. Data: 0.953 (direct) vs 0.9595 (recursion) at $L = 2$, $p_0 = 0.2$; difference below the tabulation error, so unresolved.

Identification. Route B's $\beta_j(0)$ is the per-level plane-wave increment (the quantity the code computes). It matches Balaban's second-moment β up to the expansion point (fine wave vs constrained minimizer), argued to change S by $O(\theta^2)$ relative (`SOURCES_D1_VERIFY` item 8) [argued]. This identification and the $\mathfrak{d}_{gf}$ bound need expansion-point bounds uniform in k. So the (PI)-type content of Route K is not removed in Route B; it moves into a uniform-in-k input (`BETA_FP_PW_VERIFY` §3).

Three points from `BETA_FP_PW_VERIFY` §3 (no grade yet):

1. $\theta_* < 1$ is required (above).

2. The β identification (B4) is a uniform-in-k input, not a fixed-level input.

3. (U) is a uniform-in-j analyticity of the level data. It plausibly needs the plane-wave analogue of a uniform C^1 bound on $\sigma^{(1)}$: uniform p-smoothness of the a^1 vertex. That is the same uniform first-order vertex smoothness as Route K's open core ($C1\sigma$).

4.3 Comparison

	Route K	Route B
uniform-in-level input	(B2): $\sup_j \sup_K \lvert\sigma_j^{(2)}\rvert < \infty$	(U), ($B\theta$), and the β identification and $\mathfrak{d}_{gf}$ bound, both with k-uniform expansion-point bounds
fixed-level inputs	(PI), (F), (M_j), V1, (G2)	LW (5.7), $\mathfrak{d}_Y$, $\theta_* < 1$
uses (W), Theorem A3	yes	no

verification	verified (conditional)	unverified (`BETA_FP_PW_VERIFY` has no grade)
common	(H0) at odd $L > 11$, L5, (G2), V1-type identification, Balaban's proofs, SU(2)	

Shared inputs: (H0), L5, (G2), odd L (> 11 for (H0)), and the identification of the computed $\beta_j(0)$ with Balaban's $\beta_j(0)$ (V1; in Route B also the expansion point). The uniform-in-level hypotheses ((B2) vs (U) + $(B\theta)$) are formally distinct as named hypotheses. Their content plausibly overlaps: both need a uniform first-order smoothness of the level-j vertex ($(C1\sigma)$ for Route K, uniform p-smoothness of the a^1 vertex for (U)). One closed route suffices for Theorem 2w.

5 The hypothesis (B2) of Route K

(B2): $\sup_{j\geq 0}\sup_{K\in \mathrm{BZ}} |\sigma_j^{(2)}(K)| < \infty$. Weaker form that suffices, (B2w): $\sup_j \int_{\mathrm{BZ}} |K|^{-2}|\sigma_j^{(2)}(K)|\, d^4K < \infty$ ($\sigma^{(2)}$ enters d_j only undifferentiated; `KAPPA_BOUNDED` Remark 2.3) [proved+verified: not graded by name; implied by `KAPPA_BOUNDED_VERIFY` §9(iv)].

$\sigma^{(2)}$ obeys the one-step recursion (`U3_AW` (1.2)): $\sigma_j^{(2)} = L^{-2}\mathcal{A}_j\sigma_{j-1}^{(2)} + Q_j[\sigma_{j-1}^{(1)}] + g_j^{(2)}$, with $\mathcal{A}_j s = L^{-2}H_j^\dagger(\oplus_l s(k_l))H_j$ and $H_j(K)$ the one-step gauge-fixed minimizer map.

5.1 Proved

- $K = 0$ bound [proved+verified]: $\sup_j |\sigma_j^{(2)}(0)| \leq C(|e|C_A + e^2)$. Theorem A3 + scaling give $|\partial^2\sigma_j^{(1)}(0)| \leq 2C_A$ (polarization; `KAPPA_BOUNDED_VERIFY` §7), and the slaving identity (W-2) expresses $\sigma_j^{(2)}(0)$ through these second derivatives.

- Each fixed j [proved+verified]: $\sigma_j^{(2)}$ is analytic on a fixed strip (Prop. A2), hence bounded. The content of (B2) is uniformity in j.

- Prop. 4.1 [proved+verified, conditional]: if (h) $h := \sup_{j,K}\|H_j(K)\| < L^2$ and $(C1\sigma)$ $\sup_j\sup_K |\partial_K\sigma_j^{(1)}(K)| < \infty$, then (B2) holds. Proof: in the form norm $\pm s \leq N(\Delta + 1)$, $N_j \leq (h^2/L^4)N_{j-1} + Z_2$, and Z_2 is finite by Theorem A3, $(C1\sigma)$, and fixed f^0 vertices.

5.2 Refuted

- (HS) $h < L\sqrt{L-1}$ (named (H′) in `U3_AW`; renamed to avoid the clash with v1's (H′)) [refuted at $L = 3$]. Measured $\max_K \|H_j(K)\|/L = 1.652, 1.736, 1.757, 1.760$ for $j = 1, 2, 3, \infty$ (`U3_AW_VERIFY` §6); the threshold is $\sqrt{2} = 1.414$. The maximum lies in a BZ face interior, $\approx (1.9, 1.9, \pi, \pi)$. Gauge choice cannot help: H_j is already minimal-norm in its gauge orbit. $L \geq 5$ (threshold ≥ 2) is untested.

5.3 Live routes and the precise remaining inequality

- Why the Lemma A4 mechanism alone fails [proved+verified, reason corrected by `KAPPA_BOUNDED_VERIFY` §7]: the obstruction is the missing uniform C^1 bound on $\sigma^{(1)}$, which enters twice: in Q_j (term $L^{-3}\partial_K\sigma_{j-1}^{(1)}$) and in the longitudinal part of $\sigma^{(2)}$ given by (W-2). Theorem A3 is C^0 only.

- Condition (h) at $L = 3$ [numerical, run by the verifier, `U3_AW_VERIFY` §6]: $h/L \approx 1.76$, so $h^2/L^4 \approx 0.34 < 1$ for $j \le 3$ and $j = \infty$ (sup over all j inferred from the observed convergence in j). A proof of (h) is [open]; $h \ge L$ is known ($H_j(0) = Le_{l=0}$).

- (H″) (n-step form): $\sup_{n,j,K} \|H_{(n),j}(K)\|/L^n < \infty$ with $L^n - 1$ above its square [open]. At $n = 2$ it passes on a coarse grid: $2.056 < 2.828$ (`U3_AW_VERIFY` §9, one level pair) [numerical, run by the verifier]. (H″) ⇒ (A) in full ⇒ (B2) and ($C1\sigma$) [argued]: the implication is a sketch in `U3_AW` §4.7; `U3_AW_VERIFY` grade 5 covers only the reduction to (H′); no verifier graded (H″) ⇒ (B2).

- Two-step recursion route (`B2_TWOSTEP.md`): an unverified candidate proof of ($C1\sigma$). It claims (H2) (a two-step norm condition) ⇒ uniform C^1 bound on $\sigma^{(1)}$ ⇒ (B2), and that a plain sup-norm contraction needs only $\rho_1 < L$, i.e. (h). No verifier file exists. Nothing from it is used here.

The precise remaining inequality for Route K, at a fixed odd $L \ge 3$, is

$$(C1\sigma)\ \sup_{j\ge 0}\sup_{K\in \mathrm{BZ}} |\partial_K \sigma_j^{(1)}(K)| < \infty, \qquad \text{together with} \qquad (h)\ \sup_{j\ge 0}\sup_{K\in \mathrm{BZ}} \|H_j(K)\| < L^2,$$

or, alternatively, (H″). ($C1\sigma$) + (h) ⇒ (B2) is [proved+verified]; (H″) ⇒ (B2) is [argued]. (B2w) directly is also enough. No per-momentum $d = 4$ constant-field code exists, so (B2) itself has not been scanned numerically (`KAPPA_BOUNDED_VERIFY` §8).

6 Conclusion

Theorem 2w ⇐ (Route K inputs) or (Route B inputs). Precisely:

Route K. (B2) [open; reduces to ($C1\sigma$) + (h) (verified reduction), or (H″) (argued reduction)] + (PI), (F), (M_j), V1, (G2) [fixed-level] + (H0) [Balaban, odd $L > 11$] + L5 [open] ⇒ Theorem K ($R = \kappa_*$) ⇒ (5.1) ⇒ Lemma D″ ⇒ Theorem 2w. Status of the implication: [proved+verified, conditional].

Route B. (U) + ($B\theta$) with $\theta_* < 1$ + LW (5.7) + bounded $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ + identification of β (uniform in k) + (G2) + (H0) [odd $L > 11$] + L5 ⇒ Theorem B ($R = C_B$) ⇒ (5.1) ⇒ Lemma D″ ⇒ Theorem 2w. Status: [proved, unverified, conditional].

Then Balaban's Theorem 1. Theorem 2w gives (b) and (c′). With γ shrunk so that (S) holds, every one of the 19 cross-scale uses in CMP 119 and CMP 122 I–II is covered (§2.3), and the single-scale uses need only $g_k \le \gamma$. So Theorem 1 holds for the bare coupling $g_0(K, g)$, for every K and every $g \in]0, g_1]$, without Balaban's coupling assumption. This needs L5 in the sharpened form of §2.4: β must exist on every history the shooting visits. Status: [proved, unverified (sweep), conditional on Route K or Route B inputs, on L5, and on Balaban's published proofs].

Order of constants. Fix $\beta_0, p_0, p_1, r, \nu, \kappa_0$ (Balaban). Fix L (odd, > 11), then R from Route K or B, then $b = c \log L$, then $\gamma \le \min(\gamma_0, \gamma_{Bal}, b/(4C_1), \gamma_*)$ with $\gamma^{-2} \ge c_{sw} \cdot 2R$. Then $g_1 = (\gamma^{-2} + 2R)^{-1/2}$. Balaban's phrase "for a sufficiently small positive g" fits $g \le g_1$.

What this does not give.

- Literal (0.31): only for windows of length $\ge 8R/b$. Literal (0.31) at every K needs (L1) or (L2) of §1.4, which need individual increments, not window sums. Not needed for Theorem 1.

- The value $\lim \beta_j(0)$ and any rate: §7.

- Even L, or odd $L \leq 11$ without (H0) as an extra hypothesis (Balaban assumes L odd > 11 throughout CMP 109); SU(N) with $N > 2$.
- Monotone couplings: g_m can exceed g_n for $m < n$ by the bounded amount $u_m \geq u_n - D'$. §2.3 shows Balaban's proof tolerates this under (S); a line-by-line read was not done.
- A continuum limit (`ASSEMBLY.md` treats the assembly separately; not claimed here).

7 The convergence route: what it adds, and its status

Not needed for Theorem 1. It adds: (i) existence of $\beta_\infty = \lim \beta_j(0)$; (ii) its value $c \log L$; (iii) a rate; (iv) eventual positivity, hence literal (0.31) for $K \geq K_0$ (Lemma D′ (d)); (v) with certified early steps, literal (0.31) at every K (§1.4 (L2)). It also implies (B2), since convergence of $\sigma_j^{(2)}$ implies boundedness. So it needs strictly more than Route K.

7.1 Components and status

- U3 Theorem 4.1 with O5: $|\partial_K^\alpha(\Delta_M - \Delta_\infty)| \leq CM^{-2}$, $|\alpha| \leq 2$, uniform to $K = 0$, odd L [proved+verified] (`U3_VERIFY`, `U3_REPAIRS_VERIFY`). Numerically $|\Delta_M - \Delta_\infty| = 1.007, 0.129, 0.0147$ at $M = 3, 9, 27$.
- O6: $\|\mathfrak{B}_j(K)^{-1}\| \leq b(L)$ for all j, K, one-step constraint, trivial background [proved+verified]. Crude constant ($b(3) \approx 7.5 \cdot 10^{33}$); measured corner values 103.5, 176.7, 193.0, 195.3 ($j = 0, 1, 2, \infty$, $L = 3$) use a surrogate Λ (`U3_REPAIRS_VERIFY` §5).
- Theorem 6.1: $\beta_{j+1}(0) = \Phi(\mathcal{D}_j)$ is Lipschitz in the level data [proved+verified, conditional on V1]. The norm is the Taylor-data norm (`U3_VERIFY` P8 correction).
- Lemma 5.1 (convergence of $\sigma_j^{(1)}$, $\sigma_j^{(2)}$ in C^2, $|\alpha| \leq 2$):
  - neutral sector, rate L^{-4j} [proved+verified for the value bound; the K-derivative paragraph is proved, unverified];
  - (W) [proved+verified] (`U3_AW` Theorem W); it removes the marginal direction of the f^1 recursion;
  - (A) qualitative and Theorem A3 [proved+verified]; (A) in full (strip/C^3 for $m = 1$, all of $m = 2$) $\Leftarrow$ (H″) [open]; the route via (HS) is [refuted] at $L = 3$;
  - $\sigma^{(1)}$ convergence at rate jL^{-2j} [sketch in `U3_AW` §4.7, unverified]; $\sigma^{(2)}$ convergence [open, needs (A) in full];
  - the single-block-constraint version of Lemma 5.1 is false for $m = 2$ in $d = 4$ [proved+verified]; Balaban's composed constraint is not affected.
- Prop. 3.6: $|\beta_{j+1}(0) - \beta_\infty| \leq C(1 + j)^2 L^{-2j}$ [proved, unverified as an assembly; conditional on V1, (A) in full]. (v1 wrote Cj^2L^{-2j}, which is 0 at $j = 0$; corrected.) The identification of σ-Taylor data with the data of (6.1) is cited from `U3_VERIFY` §7, not re-shown. (N) (clean rate L^{-2j}) is not needed and is judged plausibly false.
- Value of the limit. Given convergence, Theorem K or Theorem B gives $\beta_\infty = c \log L$ at once: $S_k/k \to \beta_\infty$ by Cesàro and $S_k/k = c \log L + O(1/k)$ [proved+verified as an implication: `THEOREM2_PROOF_VERIFY` Point 4; `BETA_FP_PW` Theorem 3.2(iii)]. Older routes to the value:

`BETA_FP_H` Theorem 3.1′ with (H) restated at the constant field, which given (W) reduces to Lemma 5.1 (1.2) alone (`BETA_FP_H_VERIFY` items (1)–(4)), plus (PI), (F), (M) [conditional; (M) argued]; and `BETA_FP_PW` Theorem 3.1 (limit form) given (U), ($C\theta$), LW (5.7) [proved, unverified]. The original (H) with IR weights (`BETA_FP_LEMMA43` (4.2)) was graded UNSUPPORTED and is not used.

- Rate of $\mathcal{C}(\Lambda)$: $O(\Lambda^{-2}\log\Lambda)$ given Lemma 5.1 with rates, via `U3_AW` §3.4(b) slaving [argued, `BETA_FP_H_VERIFY` §5].

7.2 Early steps and j_*

Under Prop. 3.6 and $\beta_\infty = c\log L$, $\beta_{j+1}(0) \geq (c\log L)/2$ for $j \geq j_* := \min\{J : C(1+j)^2 L^{-2j} \leq (c\log L)/2 \text{ for } j \geq J\}$. Heuristic size [argued]: $C \sim b^3$ gives $j_* \approx 110$ ($L = 3$), ≈ 70 ($L = 13$); $C \sim b^6$ (one-step inverses inside C_{data}) gives ≈ 218, ≈ 138 (`THEOREM2_PROOF_VERIFY` Point 5(e)). With the measured $\|\mathfrak{B}^{-1}\| \approx 195$: $j_* \approx 10$ ($L = 3$), ≈ 4 ($L = 13$). Certifying $\beta_1(0), \ldots, \beta_{j_*}(0) > 0$ needs interval arithmetic at odd $L \geq 13$ (blocks $\approx 3.4 \cdot 10^5$ rows per momentum by dense methods; the umklapp code reduces this to 12×12 blocks at L^4 momenta). It is needed only for (L2), not for Theorem 1.

8 Numerics

SU(2), infinite volume, Balaban's normalized-trace units. None of this is certified. Only odd L is inside the proofs; $L = 2, 4$ use corner blocks and are outside them.

8.1 First increment $\beta_1(0)$ (`NUMERICS` §6–§7; `UMK_BETA1.md`)

L	$\beta_1(0)$	how	$b = c\log L$	$\beta_1 - b$	status
2	0.630	$N_g = 12$, $p \to 0$ fit	0.2575	+0.372	[numerical, reproduced] (`KAPPA_BOUNDED_VERIFY` §4: 0.6296)
3	0.664	$N_g = 8$, $p \to 0$ fit (0.6625–0.6661)	0.4082	+0.256	[numerical, reproduced]; N_g 6 → 8 shift ≈ 0.015
4	0.757	$N_g = 6$, single $p = 0.2$; $O((Lp)^2)$ bias ≈ 0.01	0.5150	+0.242	[numerical]
5	0.749 ($N_g = 12$, $p = 0.1$); fit 0.743	umklapp code	0.5979	+0.145	[numerical]
7	0.803	umklapp code, $N_g = 8$, $Lp = 0.5$	0.7229	+0.080	[numerical]

The first-step scheme constant $\beta_1 - b$ decreases with L. The umklapp code was validated against `ym_momentum` to 10^{-12} per K by its author; no independent verifier. Queue continues to $L = 15$ on Windows. No value at $L > 11$ exists yet.

8.2 Later increments ($L = 2$ only)

quantity	value	status
$\beta_1 + \beta_2$ (composed, $N_g = 4/6/8$)	1.148 / 0.967 / 0.953	[numerical]; single $p = 0.2$, Lp bias label
$\beta_2(0)$	≈ 0.32 ($0.953 - 0.630$); per-level recursion 0.317–0.341 ($N = 8$–12, single $p = 0.2$); 0.333 (verifier, symmetric Nyquist, $N = 8$)	[numerical]; per-level recursion reproduced in `BETA_FP_VERIFY` §7
$\beta_3(0)$	0.40 ($N = 8$), 0.286 ($N = 12$), 0.261 (verifier, symmetric Nyquist, $N = 8$)	unconverged; not usable
abelian sector S_k^{ab}	$(1 - L^{-4k})/4$, matched to all digits	[proved+verified]

All computed increments are positive.

8.3 Quantities of the Route K proof

quantity	value	status
$d_1 = b - \beta_1$, $L = 3$	-0.256 ± 0.01 (fits $-0.254 \cdots - 0.258$)	[numerical, reproduced]
d_1, κ_1, $L = 2$	-0.372, 1.114	[numerical, reproduced]
d_2, d_3, $L = 2$	≈ -0.44, ≈ -0.45 (d_3 uses unconverged β_3)	[numerical]; bounded-looking, not a test
$\max_K \|H_j(K)\|/L$, $L = 3$, $j = 1, 2, 3, \infty$	1.652, 1.736, 1.757, 1.760	[numerical, run by the verifier, `U3_AW_VERIFY` §6]; refutes (HS); $h^2/L^4 \approx 0.34 < 1$
(H″), $n = 2$, $L = 3$	$2.056 < 2.828$	[numerical, run by the verifier], coarse grid, one level pair
$\sigma_j^{(1)}(0)$ J-part, $j = 0, 1, 2$ ($d = 2$ proxy)	0.750000, 0.75015, 0.75051 vs 0.75	[numerical, run by the verifier] supports Theorem W
$\sigma_j^{(2)}(K)$ scan	not run: no $d = 4$ constant-field per-momentum code	—

9 Ledger (tags match the verifier files)

[Lean] (no sorry; axioms propext, Classical.choice, Quot.sound)

- Lemma D′ (a)–(d) (`YmLean.lemma_DPrime`), Step 8 (`lemma_DPrime_step8`), Lemma D, Lemma E (Cg^2 form), Lemma D+E (`lean/YmLean/LemmaDPrime.lean`, `LemmaDE.lean`). One-argument β only.
- The Lean folder also has Telescoping, DetIdentities, AbelianCount, and LemmaH files (`THEOREM2_PROOF_v2_VERIFY` §M item 8); this chapter does not use them.

[proved+verified]

- Lemma D′ (a)–(d), K_0, Remark 4 (`LEMMA_DPRIME_VERIFY` grades 1, 2, 5).

- Lemma D (corrected interval), Lemma E (Cg^2 form), nested-interval variant of Lemma D (`LEMMAS_DE_VERIFY`). Lemma E C^1 form: trivial variant, same proof.
- Lemma D″ (tail-sum form, no J) (`KAPPA_BOUNDED_VERIFY` §6).
- Telescoping $S_k = \sum \beta_j(0)$ for the $p \to 0$ coefficient (`BETA_FP_VERIFY` item 1, corrected scope).
- Prop. 2.1 anchor-free identity; exact at the constant field (`BETA_FP_LEMMA43_VERIFY` item 1; `BETA_FP_H_VERIFY` item 1).
- $c = 11N^2/(12\pi^2)$, normalization chain; (F) source (Savvidy eq. (1.4)) (`BETA_FP_LEMMA43_VERIFY` item 2; `BETA_FP_H_VERIFY` §4).
- KAPPA Prop. 1.1 as an identity given its fixed-level inputs; Lemma 2.1 (integration by parts); Prop. 2.2; level-independence of the degree -4 part; Theorem K and Theorem 2w as conditional reductions (`KAPPA_BOUNDED_VERIFY` (i)–(v)).
- $\sup_j |\sigma_j^{(2)}(0)| < \infty$; Prop. 4.1 algebra (`KAPPA_BOUNDED_VERIFY` (vi), reason for A4 failure corrected).
- Prop. E evenness, Theorem W0/W, Theorem A3, Lemma A4, A0, Prop. A2 (`U3_AW_VERIFY` grades 1–4).
- U3 Theorem 4.1 + O5, Prop. A, Prop. B, O6, Theorem 6.1 (given V1; Taylor-data norm), Lemma 5.1 neutral value bound, single-block $m = 2$ failure in $d = 4$, one-step recursion, R2 midpoint identity (`U3_VERIFY`, `U3_REPAIRS_VERIFY`, `U3_LEMMA51_VERIFY`, `U3_LEMMA51_R23_VERIFY`).
- Cesàro: given convergence, $\kappa_k = o(k) \Leftrightarrow \beta_\infty = c \log L$ (equivalence, not a weaker form) (`THEOREM2_PROOF_VERIFY` Point 4).

[proved, unverified]

- Theorem B and Lemma 1.3 of `BETA_FP_PW` (`BETA_FP_PW_VERIFY` has no grade; its checks of Lemma 1.3, Lemma 2.3, and Prop. 2.2 are correct; it adds $\theta_* < 1$ and the uniform-in-k β identification).
- `LEMMA_DPRIME_SWEEP`: 19 cross-scale uses covered by (b) + (c′) + (S) (4 of 19 table rows, 11, 12, 16, 18, verified on page images by `LEMMA_DPRIME_VERIFY` comparisons A–E).
- Step "Theorem 2w ⇒ Balaban's Theorem 1" (depends on the sweep and on L5 in the sharpened form of §2.4).
- Prop. 3.6 assembly with $C(1+j)^2 L^{-2j}$; neutral-sector K-derivative paragraph; $\sigma^{(1)}$ convergence sketch jL^{-2j}.
- Necessity of individual positivity for literal (0.31) at every K (elementary).
- `BETA_FP_M` Prop. 4.3 ($\log|\det|$ bound in the N–O window, given (S1)–(S3)); no verifier file.
- `B2_TWOSTEP` (H2) ⇒ ($C1\sigma$) ⇒ (B2) (candidate; no verifier file).

[argued]

- (PI); (M_j) (Prop. 4.4 of `BETA_FP_M`); (U); $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ bounds; plane-wave vs constrained-minimizer expansion point (uniform in k); (H″) ⇒ (A) in full ⇒ (B2) (`U3_AW` §4.7 sketch); j_* size; the rate $O(\Lambda^{-2}\log\Lambda)$.

[open]

- (B2), reduced here to $(C1\sigma)$ + (h) [verified reduction], or (H″) [argued reduction]. $(B\theta)$ and (U) for Route B. V1. (S1)–(S3). L5 (domain of β; nested-interval D″ [not written]). K-independence of $\beta_j(g)$. (A) in full. Lemma D″ in Lean. See §10.

[refuted]

- (HS) $\sup\|H_j\| < L\sqrt{L-1}$ at $L = 3$.
- `U3_LEMMA51` Theorem 4.1 for $m = 2$ with a single block-M constraint in $d = 4$.
- v1 claims withdrawn: "every early step must be certified for Theorem 1" (false: Lemma D′/D″); "(H′) is weaker than $\beta_\infty = c\log L$" (it is equivalent); "the whole $\log\Lambda$ sits in the fine piece" as a fact (it is equivalent to the open statement); "one-loop parts hold for SU(N)" (Lemma 5.1's charged/neutral split is the SU(2) Cartan split); (3.2) with Cj^2L^{-2j} (0 at $j = 0$); the (M) resolvent bound $\|\mathcal{K}_t^{-1}\| \le C/|f|$ (Nielsen–Olesen zero); (H) with IR weights as stated (UNSUPPORTED).

[Balaban]

- (H0) smoothness and uniform bounds of β_{j+1} in g (CMP 109 (1.22), pp. 263–264), at odd $L > 11$ only; Theorem 1 and the inductive bounds of CMP 109, 116, 119, 122 and the propagator papers CMP 95, 96, 98, 99, 102. Uniform level bounds partly cited from CMP 95/96/99 (O6 replaced the main citation).

[verified quote]

- Theorem 2 and Theorem 1 wording, "separate paper", p. 298 history sentence, (0.1) with $1/g_k^2$ and χ_k, (1.3)–(1.5) (print misprint $E^{(j+k)}$ read as $E^{(j+1)}$), (2.6)–(2.9) of CMP 119, items A–E of CMP 122 (`BALABAN_EXCERPTS_CHECK`; `LEMMA_DPRIME_VERIFY`; `LEMMA_DPRIME_SWEEP` page images).

[numerical] see §8.

10 Open list (complete for Theorem 2w ⇒ Theorem 1)

Route K hypothesis set (all must hold):

1. (B2) $\sup_j \sup_K |\sigma_j^{(2)}| < \infty$ (or (B2w)). Precise remaining form: $(C1\sigma)$ $\sup_{j,K} |\partial_K \sigma_j^{(1)}| < \infty$ and (h) $\sup_{j,K} \|H_j(K)\| < L^2$ [reduction proved+verified]; or (H″) [reduction argued]. (h) holds numerically at $L = 3$ ($h^2/L^4 \approx 0.34$); (HS) is refuted. `B2_TWOSTEP.md` is an unverified candidate proof of $(C1\sigma)$ from a two-step norm condition (H2); it has no verifier file.

2. (PI) probe independence, at each fixed level [argued].

3. (F) lattice = continuum log coefficient at constant field [cited source checked; lattice transfer standard, not proved here].

4. (M_j) formal = true f^2 coefficient at each fixed level: (S1) $\mathcal{K}_t(f) \geq -C_1|f|$, (S2) density of states $\leq C_2|f|$ near 0, (S3) relative bound of the level difference, all on the lattice [open]; Prop. 4.4 [argued].

5. V1 order-g^0 completeness of CMP 109 (1.3)–(1.5): $\beta_{j+1}(0)$ is the Maxwell coefficient of the per-step Gaussian determinant [open; supported at order p^2].

6. (G2) per-cell infinite-volume densities [formal].

Route B hypothesis set (alternative to 1–4; still needs 5–6 and the common items):

B1. (U) $|\beta_{j+1}(p) - \beta_{j+1}(0)| \leq C_U p^2$, uniform in j, $0 < p \leq p_*$ [hypothesis; argued from uniform analyticity in p of the one-step data; plausibly needs uniform p-smoothness of the a^1 vertex, the same kind of input as $(C1\sigma)$].

B2′. $(B\theta)$ $\sup_k |\Gamma^{ex}(\theta_*; L^k)| < \infty$ at one $\theta_* > 0$; it needs fixed-θ vertex bounds ($(D\theta)$ in bounded form) and bounded defects $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$ [hypothesis/argued].

B3. Lüscher–Weisz (5.7) [cited].

B4. Identification of the plane-wave increment with Balaban's β up to the expansion point [argued]. This is a uniform-in-k input, not a fixed-level input.

B5. $\theta_* < 1$, so that $c_2 \ln(1/\theta_*^2) \geq 0$ in C_B [trivial condition on the choice of θ_*].

B6. Independent verification of `BETA_FP_PW` (`BETA_FP_PW_VERIFY` has §1–§5, no grade).

Common to both routes:

7. (H0) Balaban's smoothness in g with uniform bounds, $|\beta_j(g) - \beta_j(0)| \leq C_1 g$ uniform in j and history [Balaban; CMP 109 inductive assumption at odd $L > 11$; no explicit proof of the g-derivative bound located in CMP 116].

8. L5 history dependence [open]. (a) The clamp proof of Lemmas D′/D″ is valid only if $\beta_{k+1}(g_0, \ldots, g_k)$ is defined and continuous on the whole cube $]0, \gamma]^{k+1}$; Balaban does not supply this. (b) Balaban's β_{k+1} exists only on histories where his induction has run; that domain must carry (c′) + (S), not only $g_i \in]0, \gamma]$. Clamped histories can violate the growth side of (c′). (c) True trajectories with all $u_i \geq \gamma^{-2}$ inherit (c′) from (a″). A nested-interval variant of Lemma D″ on these trajectories would avoid the gap [not written]; the variant is written only for Lemma D (`LEMMAS_DE_VERIFY` §1.5). Lean covers one-argument β only.

9. K-independence of $\beta_j(g)$ [open]. The shooting needs $\beta_j(g)$, γ, and the constants independent of $K = -\log_L \varepsilon$. After rescaling, each step is the same unit-lattice block step, so β_j is K-independent only in the infinite-volume limit; a finite volume brings K-dependence back. This infinite-volume identification is not proved (`LEMMAS_DE_VERIFY` §2 item 2; `THEOREM2_PROOF_VERIFY` Point 7). It is used implicitly through (G2) and (H0).

10. Balaban readings: the sweep (§2.3) is grep plus page images, not line-by-line, with 4 of 19 rows checked by a second reader; "Similar inequalities hold for other constants, which will be introduced later" (CMP 119 p. 256) not located; β_0 must be fixed before γ, while Balaban says β_0 "can be chosen arbitrarily small, if g is sufficiently small" (not checked line by line); `BALABAN_EXCERPTS_CHECK` item 11 lists unchecked readings (Π (4.37)/(5.1), CMP 119 (2.42), (2.12)/(2.23), the "L odd" page, Sect. 3 estimates, 1989 bound lists); Theorem 1 itself rests on Balaban's published proofs.

11. Odd L. All one-loop proofs need odd $L \geq 3$ (centered blocks: U3 Lemma 1.2, O5/Prop. A, evenness, Lemma 4.1 of LEMMA43). Balaban assumes L odd > 11 throughout CMP 109, so (H0) and $\beta_j(g)$ at $g > 0$ exist only there. Theorem 2w is stated for odd $L > 11$, or with (H0) at this L as a hypothesis. No numerics exist at $L > 11$ (umklapp queue to $L = 15$ running); (h) and (H″) are tested at $L = 3$ only, outside the range of Theorem 2w, so they are supportive only.

12. SU(2) only. The charged/neutral split of Lemma 5.1 and (W) are SU(2) statements; $c = 11N^2/(12\pi^2)$ is general but the chain is not checked for $N > 2$.

13. Lean formalization of Lemma D″ and of the history form [open, small].

Not needed for Theorem 1, open for the stronger statements: (A) in full and Lemma 5.1 for $m = 2$ (convergence, rate); certified $\beta_1(0), \ldots, \beta_{j_*}(0) > 0$ at odd $L > 11$ (literal (0.31) at every K); explicit j_*; (N).

Hypothesis table (from `THEOREM2_PROOF_v2_VERIFY` §K, with the v3 status).

hypothesis	introduced	verified / checked in	status	§10 item
(B2) / (B2w)	KAPPA §2, Rem. 2.3	KAPPA_VERIFY §7–§9	open	1
$(C1\sigma)$ + (h) ⇒ (B2)	KAPPA Prop. 4.1	KAPPA_VERIFY §7	implication verified; $(C1\sigma)$ open; (h) numerical $L = 3$	1
(H″) ⇒ (A) ⇒ (B2)	U3_AW §4.7	none (U3_AW_VERIFY grade 5 covers (H′) only)	implication argued; (H″) numerical $n = 2$ only	1
(HS)	U3_AW (H′)	U3_AW_VERIFY §6	refuted $L = 3$	§5.2
B2_TWOSTEP (H2) ⇒ $(C1\sigma)$	B2_TWOSTEP §2–§3	none	proved, unverified (candidate)	1
(PI)	BETA_FP_H §3.1	H_VERIFY §5	argued	2
(F)	BETA_FP_LEMMA43	H_VERIFY §4	cited, source checked	3
(M_j): (S1)–(S3), Prop. 4.4	BETA_FP_M §4	none for BETA_FP_M; H_VERIFY §5 refuted old step	open / argued	4

V1	R2	U3_VERIFY C4 (order p^2)	open	5
(G2)	BETA_FP_LEMMA43	—	formal	6
$\beta_j(g)$ independent of K	LEMMAS_DE_VERIFY §2 item 2	v1 verifier Point 7	open	9
(H0), uniform in j and history	CMP 109 (1.22)	BALABAN_CHECK item 10	[Balaban], odd $L > 11$	7
L5 history form and domain	LEMMA_DPRIME Rem. 3; LEMMAS_DE_VERIFY §2.1; v2_VERIFY §F	—	open	8
sweep reading	SWEEP §7	LEMMA_DPRIME_VERIFY Steps 4–5 (rows 11, 12, 16, 18)	proved, unverified	10
odd L; $L > 11$ in Balaban	CMP 109 pp. 251–252	v1 verifier Step 6	input	11
SU(2)	—	—	scope	12
Lean D″, history	—	—	open	13
Route B: (U)	BETA_FP_PW §1.2	PW_VERIFY §3	hypothesis	B1
Route B: $(B\theta)$, $\mathfrak{d}_Y$, $\mathfrak{d}_{gf}$	BETA_FP_PW §3.3, §4.3	PW_VERIFY (no grade)	hypothesis / argued	B2′
Route B: β identification, uniform in k	SOURCES_D1_VERIFY 8; PW_VERIFY §3	—	argued	B4
Route B: $\theta_* < 1$	PW_VERIFY §3	—	trivial condition	B5
LW (5.7)	cited	PW_VERIFY §3 (regime correct)	cited	B3
BALABAN_CHECK item 11 readings	BALABAN_EXCERPTS	—	unchecked	10

11 Citation and independence

Related work 1. E. Sussman and L. Alpöge (with Claude; README: "written largely by LLM … Novel mathematical ideas should be attributed to Claude (under the guidance of LA)"), "The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0" (LaTeX, files `theorem_0_complete_account_raw_{master,I,II,III}.tex`, 424k lines), and "SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality" (`ym4_manuscript_zeroth_draft.pdf`, dated 6 October 2026) [22, 21]. Repository: `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit `1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310` (2026-10-27T05:45:28-05:00), accessed 2026-10-27. Repository created 2026-10-06. Disclaimer in README: "This repository contains work-in-progress, written largely by LLM. Its content has not been satisfactorily vetted. All claims should be understood as tentative, pending verification." Suggested keys: SA26-T0 (account), SA26-YM4

(draft).

Related work 2. OpenAI, "The canonical massive continuum limit of the two-dimensional O(3) model" (October 4, 2026), `github.com/openai/math`, family 215 (preprints folder; read locally from the PDF, `reports/ONLINE_SWEEP.md` §1.2; text in `reports/o3_continuum.txt`) [18]. Suggested key: OAI26-O3.

How content from these works reached this project. The record (file times, `PLAN.md`, read lists; `THEOREM2_PROOF_v2_VERIFY` §I) shows the following.

1. This chapter was written without reading the Sussman–Alpöge repository. Only the citation block above (`reports/SUSSMAN_ALPOGE_MAP.md` §5.1) was read.

2. First channel. On the night of 2026-10-27, `reports/ONLINE_SWEEP.md` §2 (about 20:20) read the Sussman–Alpöge repository directly. It reported that they prove a weak two-sided flow with an additive constant, never need per-step positivity, use a first-passage cutoff, and invoke only containment $g_k \le \gamma$. This is the first project file with Sussman–Alpöge content.

3. Second channel. `reports/SUSSMAN_ALPOGE_MAP.md` (about 20:28) was written by a separate agent under a firewall. Through it, the bounded-κ idea entered the project: the weak form of Theorem 2 needs only κ_k bounded, not convergence (their Lemma 4.19 structure; `PLAN.md` line 146). `KAPPA_BOUNDED.md` (about 20:39) received this idea through its prompt. Its read list excludes the map and ONLINE_SWEEP.

4. Lemma D′ and the cross-scale sweep predate any Sussman–Alpöge content in project files. The shooting-without-positivity lemma and the band with additive constant appear in `THEOREM2_PROOF_VERIFY` Point 5(b) (about 20:05) and `LEMMA_DPRIME.md` (about 20:10). `LEMMA_DPRIME_SWEEP.md` (about 20:16–20:23) overlaps ONLINE_SWEEP in time, but its read list is LEMMA_DPRIME and its verifier only; it did not read ONLINE_SWEEP. Lemma D″ is in `KAPPA_BOUNDED.md`, written after the map.

5. Theorem K. Its result form, $|S_k - ck\log L| \le \kappa_*$, matches Sussman–Alpöge Lemma 4.2(b), $|\sum_l \beta_l^0 - kc_0| \le c_2$ (as reported in `PLAN.md` line 146; cited there, not proved). The proof in this project (Prop. 1.1, Lemma 2.1, Prop. 2.2 of `KAPPA_BOUNDED`) was written by firewalled agents from Balaban's papers and project files.

6. O(3) paper. Route B's doubling comparison follows the template of OAI26-O3 Lemma 6.1 (doubling difference; `BETA_FP_PW` uses the "O(3) Lemma 6.1 form"). The idea of a coupling band in place of per-step positivity also appears there (ONLINE_SWEEP §1.2(a)). `BETA_FP_PW` read ONLINE_SWEEP §1.2, §5, §8 and o3_continuum Lemma 6.1, and did not read ONLINE_SWEEP §2.

7. Route B (other than the O(3) template), the numerics, and the symbol results of §3.5 and §5 were derived by firewalled agents from Balaban's papers and project files.

Any write-up must cite SA26-T0, SA26-YM4, and OAI26-O3, and state these overlaps.

A Change log (v2 → v3)

1. Range of L: new paragraph after the tag legend; §1.1 wording "Balaban assumes L odd, $L > 11$, for the whole paper"; Theorem 2w stated for odd $L > 11$, or odd $L \ge 3$ with (H0) at this L as a hypothesis; same restriction noted in §1.4 necessity, §2 notation, §4, §4.1 (H0),

§6 order of constants and "what this does not give", §9 [Balaban], §10 items 7 and 11. "his constants need odd $L > 11$" replaced (`v2_VERIFY` §D, §L(7)).

2. L5 sharpened in §2.4, §2.1–§2.2 remarks, §6, §9, and §10 item 8: the clamped shooting evaluates β on histories outside Balaban's domain; the nested-interval variant of Lemma D″ on true trajectories is described and tagged [not written] (`v2_VERIFY` §F).

3. (H″) ⇒ (B2) tagged [argued] in §5.3, §6, §9, §10 (`v2_VERIFY` §C).

4. "independent" / "disjoint" replaced by "formally distinct" in §4 title, §4 intro, §4.3; Route B points added: $\theta_* < 1$ (§4.2 statement, B5), β identification uniform in k (§4.2, §4.3 table row moved, B4), (U) plausibly needs the same uniform first-order vertex smoothness as $(C1\sigma)$ (§4.2, §4.3, B1) (`v2_VERIFY` §G).

5. §11 completed: ONLINE_SWEEP §2 named as the first channel; map named as the channel of the bounded-κ idea; Lemma D′ and the sweep dated before any Sussman–Alpöge content; Theorem K's result form compared with their Lemma 4.2(b); OpenAI O(3) paper (family 215) cited for the doubling difference and the coupling band (`v2_VERIFY` §I).

6. §10: added K-independence of $\beta_j(g)$ (item 9), $\theta_* < 1$ (B5), B2_TWOSTEP as an unverified candidate proof of $(C1\sigma)$ (item 1; also §5.3, §9), unchecked Balaban items (item 10: CMP 119 p. 256 "other constants", β_0 before γ, `BALABAN_EXCERPTS_CHECK` item 11); hypothesis table of `v2_VERIFY` §K added; items renumbered (`v2_VERIFY` §J, §K).

7. §2.3: "4 of 19 table rows verified (rows 11, 12, 16, 18)" replaces "5 of 19 items"; sweep items 4, 6, 9, 14 added to the bullet list (`v2_VERIFY` §E).

8. §1.2: "D-free" replaced; γ_* depends on R through B' (`v2_VERIFY` §A).

9. Status tags: `BETA_FP_PW_VERIFY.md` checked 2026-10-27; it has §1–§5 and no grade. Route B stays [proved, unverified]; "verifier running" replaced by "no grade". "Theorem 2w ⇒ Theorem 1" stays [proved, unverified], now also conditional on L5 (`v2_VERIFY` §N).

10. Smaller wording fixes (`v2_VERIFY` §A, §B, §E, §J, §L(9)): title no longer says "verified route"; §2.1 cites `LEMMA_DPRIME_VERIFY` grades 1, 2, 5 and notes the CORRECTED grades 3, 4, 6 of Step 7; (B2w) tag cites `KAPPA_BOUNDED_VERIFY` §9(iv); (h) and (H″) numerics tagged [numerical, run by the verifier]; B2_TWOSTEP no longer described as nonexistent; §9 notes the other Lean files.

Status: complete (2026-10-28). Not verified as a whole document.

References

[1] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. I. *Comm. Math. Phys.*, 95:17–40, 1984.

[2] Tadeusz Bałaban. Propagators and renormalization transformations for lattice gauge theories. II. *Comm. Math. Phys.*, 96:223–250, 1984.

[3] Tadeusz Bałaban. Averaging operations for lattice gauge theories. *Comm. Math. Phys.*, 98:17–51, 1985.

[4] Tadeusz Bałaban. Propagators for lattice gauge theories in a background field. *Comm. Math. Phys.*, 99:389–434, 1985.

[5] Tadeusz Bałaban. Spaces of regular gauge field configurations on a lattice and gauge fixing conditions. *Comm. Math. Phys.*, 99:75–102, 1985.

[6] Tadeusz Bałaban. Ultraviolet stability of three-dimensional lattice pure gauge field theories. *Comm. Math. Phys.*, 102:255–275, 1985.

[7] Tadeusz Bałaban. The variational problem and background field in renormalization group method for lattice gauge theories. *Comm. Math. Phys.*, 102:277–309, 1985.

[8] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. I. generation of effective actions in a small field approximation and a coupling constant renormalization in four dimensions. *Comm. Math. Phys.*, 109:249–301, 1987.

[9] Tadeusz Bałaban. Convergent renormalization expansions for lattice gauge theories. *Comm. Math. Phys.*, 119:243–285, 1988.

[10] Tadeusz Bałaban. Renormalization group approach to lattice gauge field theories. II. cluster expansions. *Comm. Math. Phys.*, 116:1–22, 1988.

[11] Tadeusz Bałaban. Large field renormalization. I. the basic step of the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:175–202, 1989.

[12] Tadeusz Bałaban. Large field renormalization. II. localization, exponentiation, and bounds for the $\mathcal{R}$ operation. *Comm. Math. Phys.*, 122:355–392, 1989.

[13] Roger Dashen and David J. Gross. The relationship between lattice and continuum definitions of the gauge-field coupling. *Phys. Rev. D*, 23:2340–2349, 1981.

[14] Elvin. yang-mills: project repository (notes/ directory holds the project files cited as `FILE.md`), 2026. Local repository; no public remote.

[15] Arthur Jaffe and Edward Witten. Quantum Yang–Mills theory. In *The Millennium Prize Problems*, pages 129–152. Clay Mathematics Institute, American Mathematical Society, 2006. Official problem statement, Clay Mathematics Institute.

[16] Martin Lüscher and Peter Weisz. Background field technique and renormalization in lattice gauge theory. *Nucl. Phys. B*, 452:213–233, 1995.

[17] Paolo Menotti and Andrea Pelissetto. Osterwalder–Schrader positivity and continuum limit for the lattice gauge theories. *Comm. Math. Phys.*, 113:369–373, 1987.

[18] OpenAI. The canonical massive continuum limit of the two-dimensional O(3) model. `https://github.com/openai/math`, family 215, preprints folder, 2026. October 4, 2026.

[19] Konrad Osterwalder and Erhard Seiler. Gauge field theories on a lattice. *Ann. Phys.*, 110:440–471, 1978.

[20] G. K. Savvidy. Infrared instability of the vacuum state of gauge theories and asymptotic freedom. *Phys. Lett. B*, 71:133–134, 1977.

[21] Ethan Sussman and Levent Alpöge. SU(2) Yang–Mills theory in four dimensions via Bałaban's renormalisation group: construction, axioms and non-triviality. ym4_manuscript_zeroth_draft.pdf, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310, 2026. Draft dated 6 October 2026. Accessed 2026-10-27.

[22] Ethan Sussman and Levent Alpöge. The Yang–Mills continuum limit along Bałaban's renormalization group: a complete account of Theorem 0. LaTeX files theorem_0_complete_account_raw_{master,I,II,III}.tex, repository `https://github.com/ethanwsussman/Yang--Mills-LLM-experiment`, commit 1b92bd0347e3b8763cd8ecc1f96dc3c2a92dd310 (2026-10-27), 2026. Written largely by LLM, with Claude; not vetted by the repository authors. Accessed 2026-10-27.