

1

University of Warsaw

2

Faculty of Mathematics, Informatics and Mechanics

3

Katarzyna Kowalska

Student no. 371053

4

Approximation and Parameterized Algorithms for Segment Set Cover

5

6

Master's thesis

7

in COMPUTER SCIENCE

8

Supervisor:

dr hab Michał Pilipczuk

Institute of Informatics

9

June 2022

10 **Supervisor's statement**

11 Hereby I confirm that the presented thesis was prepared under my supervision and
12 that it fulfils the requirements for the degree of Master of Computer Science.

13 Date

Supervisor's signature

14 **Author's statement**

15 Hereby I declare that the presented thesis was prepared by me and none of its contents
16 was obtained by means that are against the law.

17 The thesis has never before been a subject of any procedure of obtaining an academic
18 degree.

19 Moreover, I declare that the present version of the thesis is identical to the attached
20 electronic version.

21 Date

Author's signature

22

Abstract

23 In this thesis we study approximation and parameterized algorithms for a variant of the
 24 SET COVER problem, where the universe of elements to cover are points in the plane, and
 25 sets to cover objects with are segments. We call this problem SEGMENT SET COVER. We
 26 also consider the problem relaxed with δ -extension, where we need to cover the points by
 27 segments, which are extended by a tiny fraction, but we compare the solution size to the
 28 optimum solution without extension. We prove that SEGMENT SET COVER is APX-hard
 29 even if we restrict segments to be axis-parallel and allow $\frac{1}{2}$ -extension. We provide FPT algo-
 30 rithms for unweighted SEGMENT SET COVER parameterized by the size of the solution k and
 31 for WEIGHTED SEGMENT SET COVER with δ -extension. Finally, we prove that WEIGHTED
 32 SEGMENT SET COVER is W[1]-hard and there does not exist an algorithm running in time
 33 $f(k) \cdot n^{o(\sqrt{k})}$ solving this problem even if we restrict the segments to 3 directions.

34

Keywords

35 geometric set cover, weighted set cover, FPT, W[1]-hard, APX-hard

36

Thesis domain (Socrates-Erasmus subject area codes)

37 11.3 Informatyka

38

39

Subject classification

40 Theory of computation \rightarrow Design and analysis of algorithms \rightarrow Parameterized complexity
 41 and exact algorithms

42 Theory of computation \rightarrow Design and analysis of algorithms \rightarrow Approximation algorithms
 43 analysis \rightarrow Packing and covering problems

44

45

Tytuł pracy w języku polskim

46 Algorytmy aproksymacyjne i parametryzowane dla problemu pokrywania punktów
 47 odcinkami na płaszczyźnie

Contents

49	1. Introduction	5
50	1.1. Background	5
51	1.2. Our contribution	6
52	2. Preliminaries	9
53	2.1. GEOMETRIC SET COVER	9
54	2.2. Parameterization	9
55	2.3. Approximation	10
56	2.4. δ -extension	10
57	2.5. WEIGHTED GEOMETRIC SET COVER	11
58	3. APX-hardness of SEGMENT SET COVER	13
59	3.1. MAX-(3,3)-SAT	13
60	3.2. Statement of reduction	13
61	3.3. Construction of the SEGMENT SET COVER instance	15
62	3.3.1. VARIABLE-gadget	15
63	3.3.2. OR-gadget	16
64	3.3.3. CLAUSE-gadget	18
65	3.3.4. Summary	20
66	3.4. Proof that the reduction is correct	23
67	4. Fixed-parameter tractable algorithm for geometric set cover problem	27
68	4.1. Fixed-parameter tractable algorithm for unweighted segments	27
69	4.1.1. Axis-parallel segments	27
70	4.1.2. Segments in arbitrary directions	28
71	4.2. Fixed-parameter tractable algorithm for weighted segments with δ -extension	30
72	4.2.1. Dense subsets	30
73	4.2.2. Algorithm	32
74	5. W[1]-hardness for axis-parallel weighted segments	35
75	5.1. Grid Tiling	35
76	5.2. Statement of reduction	36
77	5.3. Construction of the Geometric Set Cover instance	37
78	5.3.1. Points	38
79	5.3.2. Segments	38
80	5.4. Proof that the reduction is correct	39

Chapter 1

Introduction

1.1. Background

Some problems in Computer Science are known to be NP-complete, meaning that assuming $P \neq NP$ there is no polynomial-time algorithm that can solve these problems. Even so, they still can be amenable to different approaches, such as approximation or parameterization.

Definition 1.1. In the **SET COVER** problem we are given a set of elements (universe) \mathcal{C} and a family of sets \mathcal{P} that are subsets of the universe \mathcal{C} and sum up to the whole \mathcal{C} . Our task is to find a set $\mathcal{R} \subseteq \mathcal{P}$ such that $\bigcup \mathcal{R} = \mathcal{C}$ and the size of \mathcal{R} is minimum possible.

SET COVER is a classical example of an NP-complete problem, which has been proven in [Dinur and Steurer, 2014] to be inapproximable with factor $(1 - o(1)) \ln n$ assuming $P \neq NP$ (which is a stronger result than APX-hardness), and W[2]-complete with the natural parameterization, see Theorem 13.21 in [Cygan et al., 2015]. However, restricting the problem to various specialized settings can lead to more tractable special cases. In this thesis we take a closer look at the GEOMETRIC SET COVER problem in the plane, where elements to cover are points in the plane and sets to cover them with are geometric objects.

Definition 1.2. **SEGMENT SET COVER** is GEOMETRIC SET COVER where objects that we cover the points with are segments in the plane.

Approximation Over the years there has been a lot of work related to approximation algorithms for GEOMETRIC SET COVER. Notably, GEOMETRIC SET COVER with unweighted unit disks admits a PTAS (see Corollary 1.1 in [Mustafa and Ray, 2010]). When we consider the same problem with weighted unit disks (or unit squares), the problem admits a QPTAS [Mustafa et al., 2014], see also [Pilipczuk et al., 2020]. On the other hand, [Chan and Grant, 2014] proved that GEOMETRIC SET COVER with unweighted axis-parallel fat rectangles is APX-hard; they also show similar hardness for GEOMETRIC SET COVER with many other standard geometric objects.

Parameterization We consider GEOMETRIC SET COVER parameterized by the size of solution. GEOMETRIC SET COVER with unit squares was first proven to be W[1]-hard in [Marx, 2005] (Theorem 5). A later follow-up work [Marx and Pilipczuk, 2015] shows that there is an algorithm running in time $n^{\mathcal{O}(\sqrt{k})}$ that solves GEOMETRIC SET COVER with unit squares or disks and that there is no algorithm running in time $f(k) \cdot n^{o(\sqrt{k})}$ for any computable f under the Exponential-Time Hypothesis, so this is a tight bound for this problem.

We also consider parameterization of weighted problems. There does not seem to be a consensus of what parameterization in the weighted setting is exactly; there was an attempt to introduce a quite complicated general framework of weighted parameterized setting in [Shachnai and Zehavi, 2017]. Kernels for several well-known weighted problems such as WEIGHTED SUBSET SUM or WEIGHTED KNAPSACK are presented in [Etscheid et al., 2017]. Another work [Kim et al., 2021] considers weighted parameterization of WEIGHTED DIRECTED FEEDBACK SET and WEIGHTED *st*-CUT.

δ -extension In this paper, we focus on SEGMENT SET COVER with δ -extension. δ -extension is a problem relaxation method based on the δ -shrinking model which was introduced in [Adamaszek et al., 2015] to provide interesting results for the MAXIMUM WEIGHT INDEPENDENT SET OF RECTANGLES problem. In this problem one is given a family of weighted rectangles and needs to find a set of non-overlapping rectangles with the largest possible total weight. In the δ -shrinking relaxed problem the returned set of rectangles must be non-overlapping after all the rectangles are shrunk by a tiny fraction δ towards the centre of symmetry. This problem is easier, because we compare the weight of the obtained solution to the optimum result before the shrinking. It might even lead to finding a set with result better than the optimum for the original problem. The authors in [Adamaszek et al., 2015] present a PTAS for MAXIMUM WEIGHT INDEPENDENT SET OF RECTANGLES with δ -shrinking, which was later improved to an EPTAS in [Pilipczuk et al., 2016], alongside with presenting a new FPT algorithm for this problem with the natural parameterization. Later, a similar δ -shrinking model was used in [Wiese, 2018] to present a PTAS for MAXIMUM WEIGHT INDEPENDENT SET OF POLYGONS with δ -shrinking.

Definition 1.3. For any $\delta > 0$ and a centre-symmetric convex object L with centre of symmetry $S = (x_s, y_s)$, the **δ -extension** of L is the open set of points:

$$L^{+\delta} = \text{Int}\{(1 + \delta) \cdot (x - x_s, y - y_s) + (x_s, y_s) : (x, y) \in L\},$$

where Int denotes the interior of a set of points. That is, $L^{+\delta}$ is the interior of the image of L under homothety centred at S with scale $(1 + \delta)$.

Analogous to δ -shrinking, δ -extension provides a framework for relaxing GEOMETRIC SET COVER problems, where we allow the returned set of objects \mathcal{R} to *almost* cover the points in the universe by requiring that they are covered by \mathcal{R} after δ -extension, i.e. by the set $\mathcal{R}^{+\delta}$. The same concept could be used for GEOMETRIC HITTING SET problems.

For a longer discussion of this concept see Section 2.4.

Similar model is used to prove that GEOMETRIC SET COVER with fat polygons relaxed with δ -extension admits an EPTAS [Har-Peled and Lee, 2009]. The δ -extension model presented there is well-defined only for fat polygons. An object P is extended by all the points that are at distance to the closest point in the object P no larger than $\delta \cdot \text{rad}(P)$, where $\text{rad}(P)$ is the largest radius of a circle inscribed into P . Since segments do not have any circle inscribed into them, the definition presented there cannot be utilized for the setting of segments considered here. Polygon extended by δ -extension defined in Definition 1.3 covers a superset of points that the polygon extended by δ -extension defined in [Har-Peled and Lee, 2009] covers. Since our definition is more permissive for any polygon, the EPTAS from [Har-Peled and Lee, 2009] also works for polygons extended according to our definition of δ -extension.

1.2. Our contribution

In this thesis we make the following contributions.

154 We show that SEGMENT SET COVER is APX-hard, even if segments are axis-parallel and
 155 we relax the problem with $\frac{1}{2}$ -extension, (Theorem 1.1).

156 **Theorem 1.1. (SEGMENT SET COVER is APX-hard).** *SEGMENT SET COVER is APX-*
 157 *hard even when relaxed with $\frac{1}{2}$ -extension and segments are axis-parallel. That is, assuming*
 158 *$P \neq NP$, there does not exist a PTAS for this problem.*

159 Theorem 1.1 implies the following. Note that segments are just degenerated rectangles.

160 **Corollary 1.1. (GEOMETRIC SET COVER with rectangles is APX-hard).** *GEOMET-*
 161 *RIC SET COVER with axis-parallel rectangles is APX-hard even when relaxed with $\frac{1}{2}$ -extension.*

162 This expands the previous result of [Chan and Grant, 2014] that GEOMETRIC SET COVER
 163 with axis-parallel fat rectangles is APX-hard, we improved the result that rectangles no longer
 164 have to be fat (Corollary 1.1) and it holds when the problem is relaxed with $\frac{1}{2}$ -extension. It
 165 also proves that the assumption in [Har-Peled and Lee, 2009] about polygons being fat is
 166 necessary, because covering with arbitrary polygons with $\frac{1}{2}$ -extension is APX-hard.

167 We also provide two FPT algorithms for parameterized SEGMENT SET COVER (Theorem 1.2) and WEIGHTED SEGMENT SET COVER relaxed with δ -extension (Theorem 1.3).

169 **Theorem 1.2. (FPT for SEGMENT SET COVER).** *There exists an algorithm that given*
 170 *a family \mathcal{P} of segments (in any direction), a set of points \mathcal{C} and a parameter k , runs in time*
 171 *$k^{\mathcal{O}(k)}(|\mathcal{C}| \cdot |\mathcal{P}|)^2$, and outputs a solution $\mathcal{R} \subseteq \mathcal{P}$ such that $|\mathcal{R}| \leq k$ and \mathcal{R} covers all points*
 172 *in \mathcal{C} , or determines that such a set \mathcal{R} does not exist.*

173 **Theorem 1.3. (FPT for WEIGHTED SEGMENT SET COVER with δ -extension).** *There*
 174 *exists an algorithm that given a family \mathcal{P} of n weighted segments (in any direction), a set of*
 175 *m points \mathcal{C} , and parameters k and $\delta > 0$, runs in time $f(k, \delta) \cdot (nm)^c$ for some computable*
 176 *function f and a constant c and outputs a set \mathcal{R} such that:*

- 177 • $\mathcal{R} \subseteq \mathcal{P}$,
- 178 • $|\mathcal{R}| \leq k$,
- 179 • $\mathcal{R}^{+\delta}$ covers all points in \mathcal{C} ,
- 180 • the weight of \mathcal{R} is not greater than the weight of an optimum solution of size at most k
- 181 for this problem without δ -extension,

182 or determines that there is no set \mathcal{R} with $|\mathcal{R}| \leq k$ such that \mathcal{R} covers all points in \mathcal{C} .

183 On the other hand, we prove that WEIGHTED SEGMENT SET COVER is $W[1]$ -hard even
 184 when segments are limited to 3 directions (Theorem 1.4) and assuming ETH there does not
 185 exist an algorithm for this problem that runs in time $f(k)(|\mathcal{C}| + |\mathcal{P}|)^{o(\sqrt{k})}$. See Figure 1.1 for
 186 a summary of parameterized results for WEIGHTED SEGMENT SET COVER. Similar table for
 187 the unweighted problem is shown in Figure 1.2.

188 **Theorem 1.4. (WEIGHTED SEGMENT SET COVER is $W[1]$ -hard).** *Consider the prob-*
 189 *lem of covering a set \mathcal{C} of points by selecting at most k segments from a set of segments \mathcal{P}*
 190 *with non-negative weights $w : \mathcal{P} \rightarrow \mathbb{R}^+$ so that the weight of the cover is minimal. Then this*
 191 *problem is $W[1]$ -hard when parameterized by k and assuming ETH, there is no algorithm for*
 192 *this problem with running time $f(k) \cdot (|\mathcal{C}| + |\mathcal{P}|)^{o(\sqrt{k})}$ for any computable function f . Moreover,*
 193 *this holds even if all segments in \mathcal{P} are axis-parallel or right-diagonal.*

See Section 2.1 for exact definitions of axis-parallel and right-diagonal segments.

This result is particularly interesting, because the problem without weights is FPT, while the weighted variant is W[1]-hard. Moreover, δ -extension allowed us to provide an FPT algorithm for the problem which is W[1]-hard otherwise.

Note that the result of Theorem 1.4 is not tight: there exists a simple algorithm running in time $f(k)(|\mathcal{C}| + |\mathcal{P}|)^k$. So the question whether there exists an algorithm for this problem running in time $f(k) \cdot (|\mathcal{C}| + |\mathcal{P}|)^{o(k)}$ is still open.

Permissive FPT is a relaxed FPT problem, where we need to find a solution of *any* size in FPT-time, but we compare it to the optimum solution of size at most k . Idea for permissive FPT in local search was presented in [Marx and Schlotter, 2011], [Gaspers et al., 2012]. Theorem 1.4 can be improved to show that a permissive FPT algorithm does not exist. This is formulated precisely in Theorem 5.2.

	exact	δ -extension
axis-parallel	?	FPT*
3 directions	W[1]-hard	FPT*
any direction	W[1]-hard*	FPT

Figure 1.1: Our results for WEIGHTED SEGMENT SET COVER parameterized by the size of a solution. Results marked with * are not explicitly given in this thesis, but they trivially follow from stronger results shown in the other cells of the table.

	exact	δ -extension
axis-parallel	FPT*	FPT*
3 directions	FPT*	FPT*
any direction	FPT	FPT*

Figure 1.2: Our results for unweighted SEGMENT SET COVER parameterized by the size of a solution. Results marked with * are not explicitly given in this thesis, but they trivially follow from stronger results shown in the other cells of the table.

Future work. There are two aforementioned problems that relate to Theorem 1.4 and were not solved in this thesis. We have given a W[1]-hardness proof for WEIGHTED SEGMENT SET COVER where segments are limited to 3 directions, but the segments in the construction may be also right-diagonal. However, it may be possible to improve this construction to use segments in 2 directions instead of 3 directions. The other question is what is the tight bound for this problem. The simple algorithm solving this problem is running in time $f(k)(|\mathcal{C}| + |\mathcal{P}|)^k$, while our lower bound refutes running time $f(k)(|\mathcal{C}| + |\mathcal{P}|)^{\sqrt{k}}$.

Another problem to consider is whether GEOMETRIC HITTING SET relaxed with δ -extension can yield some better results.

Acknowledgments. Words cannot express my gratitude to my supervisor, dr hab Michał Pilipczuk, for his patience and eager help with research and editing. I am also grateful to Krzysztof Maziarz for countless hours of proofreading this thesis.

218 Chapter 2

219 Preliminaries

220 In this chapter we present some basic definitions that will be used later.

221 2.1. GEOMETRIC SET COVER

222 Whenever speaking about GEOMETRIC SET COVER, we consider it in the 2-dimensional
223 plane.

224 In the GEOMETRIC SET COVER problem we are given \mathcal{P} — a set of objects, which
225 are connected subsets of the plane and \mathcal{C} — a set of points in the plane. The task is to choose
226 $\mathcal{R} \subseteq \mathcal{P}$ such that every point in \mathcal{C} is inside some object from \mathcal{R} and $|\mathcal{R}|$ is minimized. We
227 will mostly consider the case where \mathcal{P} consists of segments in the plane.

228 In the weighted setting, there is some given weight function $f : \mathcal{P} \rightarrow \mathbb{R}^+$ and we would
229 like to find a solution \mathcal{R} that minimizes $\sum_{R \in \mathcal{R}} f(R)$.

230 **Definition 2.1.** A segment is **axis-parallel** if it lies on a line that is either horizontal $y = c$
231 or vertical $x = c$.

232 **Definition 2.2.** A line is **right-diagonal** if it is described by the linear function $x + y = d$
233 for some $d \in \mathbb{R}$. Segment is **right-diagonal** if its direction is a right-diagonal line.

234 2.2. Parameterization

235 In the parameterized setting of the GEOMETRIC SET COVER for a given k , our task is to
236 either find a solution \mathcal{R} such that $|\mathcal{R}| \leq k$ or decide that there is no such solution.

237 **Definition 2.3.** A **Fixed-parameter Tractable (FPT)** algorithm for a problem with pa-
238 rameter k and instance size n is an algorithm running in time $f(k) \cdot n^c$ for some constant c
239 and some computable function f .

240 **Definition 2.4.** Boolean formula is in **conjunctive normal form (CNF)** if it is a con-
241 junction of one or more formulas, which are disjunction of literals. **k -CNF** formula is a CNF
242 formula, where every disjunction consists of at most k literals.

243 **Definition 2.5.** **k -SAT** problem is a Boolean satisfiability problem of k -CNF formulas.
244 Given k -CNF formula, one must answer if there exists any variable assignment that satisfies
245 the formula.

Definition 2.6. For $k \geq 3$ set us define S_k as a set of constants σ such that there exists an algorithm solving k -SAT running in time $\mathcal{O}^*(2^{\sigma n})$. Let s_k be the infimum of the set S_k .

Exponential Time Hypothesis (ETH) asserts that $s_3 > 0$. This conjecture implies that there does not exist an algorithm solving 3-SAT running in time $2^{o(n)}$.

To see the definition of a W[1]-hard problem and W hierarchy, see Chapter 13.3 of [Cygan et al., 2015]. When proving that a problem is W[1]-hard, we are going to use Theorem 5.1, which was proved in [Marx, 2007].

2.3. Approximation

Let us recall some definitions related to optimization problems.

Definition 2.7. A **polynomial-time approximation scheme (PTAS)** for a minimization problem Π is a family of algorithms \mathcal{A}_ϵ for every $\epsilon > 0$ such that \mathcal{A}_ϵ takes an instance I of Π and in polynomial time finds a solution that is within a factor of $(1 + \epsilon)$ of being optimal. This means that the reported solution has weight at most $(1 + \epsilon)\text{opt}(I)$, where $\text{opt}(I)$ is the weight of an optimal solution to I .

Definition 2.8. A problem Π is **APX-hard** if assuming $P \neq NP$, there exists $\epsilon > 0$ such that there is no polynomial-time $(1 + \epsilon)$ -approximation algorithm for Π .

2.4. δ -extension

Another idea presented here, which can be utilized only when considering the problems with geometric objects, is δ -extension. We define it specifically for the GEOMETRIC SET COVER problem with convex centre-symmetric objects.

Intuitively, we consider a problem with slightly larger objects, which makes the instance more permissive. However, we aim to find a solution that is not larger than the optimum solution to the original problem, so this is substantially easier than just solving the problem for the larger objects. It may even be the case that we are able to find a solution of size smaller than the optimum solution to the original problem.

Formal definition of δ -extended objects is present in Definition 1.3.

The GEOMETRIC SET COVER with δ -extension is a version of GEOMETRIC SET COVER with the following modifications.

- We need to cover all the points in \mathcal{C} by selecting objects from $\{P^{+\delta} : P \in \mathcal{P}\}$ (which always include no fewer points than the objects before δ -extension).
- We look for a solution that is not larger than the optimum solution to the original problem. Note that it does not need to be an optimal solution in the modified problem.

Formally, we have the following.

Definition 2.9. The **GEOMETRIC SET COVER problem with δ -extension** is the problem where for an input instance $I = (\mathcal{P}, \mathcal{C})$ of GEOMETRIC SET COVER, the task is to output a solution $\mathcal{R} \subseteq \mathcal{P}$ such that the δ -extended set $\{R^{+\delta} : R \in \mathcal{R}\}$ covers \mathcal{C} and is not larger than the optimal solution to the problem without extension, i.e. $|\mathcal{R}| \leq |\text{opt}(I)|$.

At last, we formulate a definition of the polynomial-time approximation scheme (PTAS) for a problem with δ -extension.

285 **Definition 2.10.** A PTAS for GEOMETRIC SET COVER with δ -extension is a family
 286 of algorithms $\{\mathcal{A}_{\delta,\epsilon}\}_{\delta,\epsilon>0}$ that each takes as an input instance $I = (\mathcal{P}, \mathcal{C})$ of GEOMETRIC SET
 287 COVER where objects are centre-symmetric and convex, and in polynomial-time outputs a
 288 solution $\mathcal{R} \subseteq \mathcal{P}$ such that the δ -extended set $\{R^{+\delta} : R \in \mathcal{R}\}$ covers \mathcal{C} and is within a $(1 + \epsilon)$
 289 factor of the optimal solution to this problem without extension, i.e. $(1 + \epsilon)|\mathcal{R}| \leq |\text{opt}(I)|$.

290 2.5. WEIGHTED GEOMETRIC SET COVER

291 In this thesis we also consider a WEIGHTED GEOMETRIC SET COVER problem, which is a
 292 combination of the weighted and parameterized setting described in Section 2.1. We already
 293 argued in the introduction that there is no consensus of how it is defined, but when we discuss
 294 the weighted parameterized setting we will consider the following definition. There is a given
 295 weight function $f : \mathcal{P} \rightarrow \mathbb{R}^+$ and we would like to find a solution \mathcal{R} such that $|\mathcal{R}| \leq k$ and
 296 $\sum_{R \in \mathcal{R}} f(R)$ is minimum possible among such sets \mathcal{R} .

297 **Definition 2.11.** The WEIGHTED GEOMETRIC SET COVER problem with δ -extension
 298 is the problem where for an input instance $I = (\mathcal{P}, \mathcal{C}, f)$ of WEIGHTED GEOMETRIC SET
 299 COVER, the task is to output a solution $\mathcal{R} \subseteq \mathcal{P}$ such that the δ -extended set $\{R^{+\delta} : R \in \mathcal{R}\}$
 300 covers \mathcal{C} and it has weight not larger than the optimal solution to the problem without ex-
 301 tension, i.e. $\sum_{R \in \mathcal{R}} f(R) \leq |\text{opt}(I)|$.

302 We also consider weighted parameterized setting with δ -extension, which we formally
 303 define below.

304 **Definition 2.12.** The WEIGHTED GEOMETRIC SET COVER problem with δ -extension
 305 parameterized by the size of a solution is a problem where for an input instance
 306 $I = (\mathcal{P}, \mathcal{C}, f, k)$ of WEIGHTED GEOMETRIC SET COVER parameterized by the size of a so-
 307 lution k , the task is to output a solution $\mathcal{R} \subseteq \mathcal{P}$ such that the δ -extended set $\{R^{+\delta} : R \in \mathcal{R}\}$
 308 covers \mathcal{C} , uses no more than k sets, i.e. $|\mathcal{R}| \leq k$ and it has weight not larger than the optimal
 309 solution to the problem without extension, i.e. $\sum_{R \in \mathcal{R}} f(R) \leq |\text{opt}(I)|$.

Chapter 3

APX-hardness of SEGMENT SET COVER

In this section we analyze whether there exists a PTAS for GEOMETRIC SET COVER for rectangles. We show that SEGMENT SET COVER is APX-hard even if we can restrict this problem to a very simple setting: segments parallel to axes and allow $\frac{1}{2}$ -extension.

Our result can be summarized in the following theorem and this section aims to prove it.

Theorem 1.1. (SEGMENT SET COVER is APX-hard). *SEGMENT SET COVER is APX-hard even when relaxed with $\frac{1}{2}$ -extension and segments are axis-parallel. That is, assuming $P \neq NP$, there does not exist a PTAS for this problem.*

We prove Theorem 1.1 by taking a problem that is APX-hard and showing a reduction. For this problem we choose MAX-(3,3)-SAT which we define below.

3.1. MAX-(3,3)-SAT

See Definition 2.4 for the definition of the k -CNF formula.

Definition 3.1. MAX-3SAT is the following maximization problem. We are given a 3-CNF formula, and we need to find a Boolean assignment of variables that satisfies the most clauses.

Definition 3.2. MAX-(3,3)-SAT is a variant of MAX-3SAT with an additional restriction that every variable appears in exactly 3 clauses and every clause contains exactly 3 literals of 3 different variables. Note that thus, the number of clauses is equal to the number of variables.

In our proof of Theorem 1.1 we use hardness of approximation of MAX-(3,3)-SAT proved in [Håstad, 2001] and described in Theorem 3.1 below.

Definition 3.3. MAX-3SAT formula with m clauses is **at most α -satisfiable**, if every assignment of variables satisfies no more than αm clauses.

Theorem 3.1 ([Håstad, 2001]). *For any $\epsilon > 0$, it is NP-hard to distinguish satisfiable (3,3)-SAT formulas from at most $(\frac{7}{8} + \epsilon)$ -satisfiable (3,3)-SAT formulas.*

3.2. Statement of reduction

The following lemma encapsulates the properties of the reduction described in this section, and it allows us to prove Theorem 1.1.

Lemma 3.1. *Given an instance S of MAX-(3,3)-SAT with n variables and optimum value $\text{opt}(S)$, we can construct an instance $(\mathcal{C}, \mathcal{P})$ of SEGMENT SET COVER with axis-parallel segments in 2D such that:*

(1) *For every solution to instance S that satisfies k clauses, there exists a solution to $(\mathcal{C}, \mathcal{P})$ of size $15n - k$.*

(2) *For every solution \mathcal{R} to instance $(\mathcal{C}, \mathcal{P})$, there exists a solution to S that satisfies at least $15n - |\mathcal{R}|$ clauses.*

(3) *For every $\mathcal{R} \subseteq \mathcal{P}$, if $\mathcal{R}^{+\frac{1}{2}}$ is a solution to $(\mathcal{C}, \mathcal{P})$, then \mathcal{R} is also a solution to $(\mathcal{C}, \mathcal{P})$.*

Therefore, the optimum size of a solution to $(\mathcal{C}, \mathcal{P})$ is $\text{opt}((\mathcal{C}, \mathcal{P})) = 15n - \text{opt}(S)$.

We prove Lemma 3.1 in subsequent sections. Section 3.3 describes the proposed instance $(\mathcal{C}, \mathcal{P})$. Property (1) is proved by Lemma 3.11, (2) by Lemma 3.13, and finally (3) trivially follows from Lemma 3.10. Firstly let us prove Theorem 1.1 using Lemma 3.1 and Theorem 3.1.

Proof of Theorem 1.1. Consider any $0 < \epsilon < \frac{1}{15.8}$.

Let us assume that there exists a polynomial-time $(1 + \epsilon)$ -approximation algorithm for unweighted SEGMENT SET COVER with axis-parallel segments in 2D with $\frac{1}{2}$ -extension. We construct an algorithm that solves the problem stated in Theorem 3.1, thereby proving that $P = NP$.

Take an instance S of MAX-(3,3)-SAT to be distinguished and construct an instance of SEGMENT SET COVER $(\mathcal{C}, \mathcal{P})$ using Lemma 3.1. We now use the $(1 + \epsilon)$ -approximation algorithm for SEGMENT SET COVER relaxed with $\frac{1}{2}$ -extensions on $(\mathcal{C}, \mathcal{P})$. Denote the size of the solution returned by this algorithm as $\text{approx}^*((\mathcal{C}, \mathcal{P}))$. We prove that if in S one can satisfy at most $(\frac{7}{8} + \epsilon)n$ clauses, then $\text{approx}^*((\mathcal{C}, \mathcal{P})) \geq 15n - (\frac{7}{8} + \epsilon)n$, and if S is satisfiable, then $\text{approx}^*((\mathcal{C}, \mathcal{P})) < 15n - (\frac{7}{8} + \epsilon)n$.

Assume S satisfiable. From the definition of S being satisfiable, we have:

$$\text{opt}(S) = n.$$

From Lemma 3.1 we have:

$$\text{opt}((\mathcal{C}, \mathcal{P})) = 14n.$$

Therefore,

$$\begin{aligned} \text{approx}^*((\mathcal{C}, \mathcal{P})) &\leq (1 + \epsilon)\text{opt}((\mathcal{C}, \mathcal{P})) = 14n(1 + \epsilon) = 14n + 14\epsilon \cdot n = \\ &= 14n + (15\epsilon - \epsilon)n < 14n + \left(\frac{1}{8} - \epsilon\right)n = 15n - \left(\frac{7}{8} + \epsilon\right)n. \end{aligned}$$

Assume S is at most $(\frac{7}{8} + \epsilon)$ satisfiable. From the definition of S being at most $(\frac{7}{8} + \epsilon)n$ satisfiable, we have:

$$\text{opt}(S) \leq \left(\frac{7}{8} + \epsilon\right)n$$

From Lemma 3.1 we have:

$$\text{opt}((\mathcal{C}, \mathcal{P})) \geq 15n - \left(\frac{7}{8} + \epsilon\right)n$$

363 Since a solution to $(\mathcal{C}, \mathcal{P})$ with $\frac{1}{2}$ -extension is also a solution without any extension, by
 364 Lemma 3.1 (3), we have:

$$\text{approx}^*((\mathcal{C}, \mathcal{P})) \geq \text{opt}((\mathcal{C}, \mathcal{P})) = 15n - \left(\frac{7}{8} + \epsilon\right)n$$

365 Therefore, by using the assumed $(1 + \epsilon)$ -approximation algorithm, it is possible to distin-
 366 guish the case when S is satisfiable from the case when it is at most $(\frac{7}{8} + \epsilon)n$ satisfiable: it
 367 suffices to compare $\text{approx}^*((\mathcal{C}, \mathcal{P}))$ with $15n - (\frac{7}{8} + \epsilon)n$. Hence, the assumed approximation
 368 algorithm cannot exist, unless $P = NP$. \square

369 3.3. Construction of the SEGMENT SET COVER instance

370 We proceed to the proof of Lemma 3.1. That is, we show a reduction from the MAX-(3,3)-SAT
 371 problem to SEGMENT SET COVER with segments parallel to axes. Moreover, the obtained
 372 instance of SEGMENT SET COVER will be robust to $\frac{1}{2}$ -extension (have the same optimal
 373 solution after $\frac{1}{2}$ -extension).

374 The construction will be composed of 2 types of gadgets: **VARIABLE-gadgets** and
 375 **CLAUSE-gadgets**. **CLAUSE-gadgets** will be constructed using two **OR-gadgets** connected
 376 together.

377 3.3.1. VARIABLE-gadget

378 **VARIABLE-gadget** is responsible for choosing the value of a variable in a CNF formula. It
 379 allows two minimum solutions of size 3 each. These two choices correspond to the two Boolean
 380 values of the variable corresponding to this gadget.

381 **Points.** Define points a, b, c, d, e, f, g, h as follows, where $L = 22n$:



Figure 3.1: **VARIABLE-gadget**. We denote the set of points marked with black circles as pointsVariable_i , and they need to be covered (are part of the set \mathcal{C}). Note that some of the points are not marked as black dots and exists only to name segments for further reference. We denote the set of red segments as $\text{chooseVariable}_i^{\text{false}}$ and the set of blue segments as $\text{chooseVariable}_i^{\text{true}}$.

382

$$\begin{array}{llll} a := (-3L, 0) & b := (-2L, 0) & c := (-L, 0) & d := (-3L, 1) \\ e := (-2L, 1) & f := (-2L, 2) & g := (L, 0) & h := (L, 2) \end{array}$$

Let us define:

$$\text{pointsVariable} := \{a, b, c, d, e, f\}$$

and, for any $1 \leq i \leq n$,

$$\text{pointsVariable}_i := \text{pointsVariable} + (0, 4i).$$

383 We denote $a_i := a + (0, 4i)$ etc.

384 **Segments.** Let us define:

$$\text{chooseVariable}_i^{\text{true}} := \{(a_i, d_i), (b_i, f_i), (c_i, g_i)\},$$

$$\text{chooseVariable}_i^{\text{false}} := \{(a_i, c_i), (d_i, e_i), (f_i, h_i)\},$$

$$\text{segmentsVariable}_i := \text{chooseVariable}_i^{\text{true}} \cup \text{chooseVariable}_i^{\text{false}}.$$

385 We also name two of these segments for future reference: $\text{xTrueSegment}_i := (c_i, g_i)$,
386 $\text{xFalseSegment}_i := (f_i, h_i)$.

387 **Lemma 3.2.** *For any $1 \leq i \leq n$, points in pointsVariable_i can be covered using 3 segments*
388 *from $\text{segmentsVariable}_i$.*

389 *Proof.* We can use either set $\text{chooseVariable}_i^{\text{true}}$ or $\text{chooseVariable}_i^{\text{false}}$. □

390 **Lemma 3.3.** *For any $1 \leq i \leq n$, points in pointsVariable_i can not be covered with fewer than*
391 *3 segments from $\text{segmentsVariable}_i$.*

392 *Proof.* No segment of $\text{segmentsVariable}_i$ covers more than one point from $\{d_i, f_i, c_i\}$, therefore
393 pointsVariable_i can not be covered with fewer than 3 segments. □

394 **Lemma 3.4.** *For every set $A \subseteq \text{segmentsVariable}_i$ such that A covers pointsVariable_i and*
395 *$\text{xTrueSegment}_i, \text{xFalseSegment}_i \in A$, it holds that $|A| \geq 4$.*

396 *Proof.* No segment from $\text{segmentsVariable}_i$ covers more than one point from $\{a_i, e_i\}$, therefore
397 $\text{pointsVariable}_i - \{c_i, f_i\}$ can not be covered with fewer than 2 segments. □

398 3.3.2. OR-gadget

399 An OR-gadget connects input and output segments (see Figure 3.2) in a way that is supposed
400 to simulate the binary disjunction.

401 Input segments are the only segments that cover points outside of the gadget, as their left
402 ends lie outside of it. Point $v_{i,j}$ is the only one that can be covered by segments that do not
403 belong to the gadget.

404 The OR-gadget has the property that every set of segments that covers all the points in
405 the gadget uses at least 3 segments from it. Moreover, the output segment belongs to the
406 solution of size 3 only if at least one of the input segments belongs to the solution. Therefore,
407 optimum solutions restricted to the OR-gadget behave like a binary disjunction for the input
408 segments.



Figure 3.2: **OR-gadget**. Segments from $\text{chooseOr}_{i,j}^{\text{false}}$ are **red**, segments from $\text{chooseOr}_{i,j}^{\text{true}}$ are blue (both **light blue** and **dark blue**), segments from $\text{orMoveVariable}_{i,j}$ are **green** and **yellow**. **Dark blue** segment is the *output* segment. Grey segments input_x and input_y are input segments that are not part of $\text{segmentsOr}_{i,j}$.

409 **Points.** We define

$$\begin{aligned}
 l_0 &:= (0, 0) & m_0 &:= (0, 1) & n_0 &:= (0, 2) & o_0 &:= (0, 3) \\
 p_0 &:= (0, 4) & q_0 &:= (1, 1) & r_0 &:= (1, 3) & s_0 &:= (2, 1) \\
 t_0 &:= (2, 2) & u_0 &:= (2, 3) & v_0 &:= (3, 2)
 \end{aligned}$$

$$vec_{i,j} := (20i + 3 + 3j, 4(n + 1) + 2j)$$

411 For integers i, j , define $\{l_{i,j}, m_{i,j}, \dots, v_{i,j}\}$ as $\{l_0, m_0, \dots, v_0\}$ shifted by $vec_{i,j}$, i.e. $l_{i,j} = l_0 + vec_{i,j}$
 412 etc.

413 Note that $v_{i,0} = l_{i,1}$ (see Figure 3.3). Next, let

$$\text{pointsOr}_{i,j} := \{l_{i,j}, m_{i,j}, n_{i,j}, o_{i,j}, p_{i,j}, q_{i,j}, r_{i,j}, s_{i,j}, t_{i,j}, u_{i,j}\}$$

414 Note that $\text{pointsOr}_{i,j}$ does not include the point $v_{i,j}$.

415 **Segments.** We define the set of segments in several parts:

$$\begin{aligned}
 \text{chooseOr}_{i,j}^{\text{false}} &:= \{(q_{i,j}, r_{i,j}), (s_{i,j}, u_{i,j})\}, \\
 \text{chooseOr}_{i,j}^{\text{true}} &:= \{(m_{i,j}, s_{i,j}), (o_{i,j}, u_{i,j}), (t_{i,j}, v_{i,j})\}, \\
 \text{orMoveVariable}_{i,j} &:= \{(l_{i,j}, n_{i,j}), (n_{i,j}, p_{i,j})\}.
 \end{aligned}$$

416 Finally all segments on OR-gadget are defined as:

$$\text{segmentsOr}_{i,j} := \text{chooseOr}_{i,j}^{\text{false}} \cup \text{chooseOr}_{i,j}^{\text{true}} \cup \text{orMoveVariable}_{i,j}$$

417 **Lemma 3.5.** For any $1 \leq i \leq n, j \in \{0, 1\}$ and $x \in \{l_{i,j}, p_{i,j}\}$, points in $\text{pointsOr}_{i,j} - \{x\} \cup \{v_{i,j}\}$
 418 can be covered with 4 segments from $\text{segmentsOr}_{i,j}$.

419 *Proof.* We can do this using one segment from $\text{orMoveVariable}_{i,j}$, the one that does not cover
 420 x , and all segments from $\text{chooseOr}_{i,j}^{\text{true}}$. \square

421 **Lemma 3.6.** For any $1 \leq i \leq n, j \in \{0,1\}$, points in $\text{pointsOr}_{i,j}$ can be covered with 4
 422 segments from $\text{segmentsOr}_{i,j}$.

423 *Proof.* We can do this using segments from $\text{orMoveVariable}_{i,j} \cup \text{chooseOr}_{i,j}^{\text{false}}$. \square

424 3.3.3. CLAUSE-gadget

425 A CLAUSE-gadget is responsible for determining whether variable values assigned in variable
 426 gadgets satisfy the corresponding clause in the input formula ϕ . It has a minimum solution
 427 of size w if and only if the clause is satisfied, i.e. at least one of the respective variables is
 428 assigned the correct value. Otherwise, its minimum solution has size $w + 1$. In this way, by
 429 analyzing the size of the minimum solution to the entire constructed instance, we will be able
 430 to tell how many clauses it is possible to satisfy in an optimum solution to ϕ .



Figure 3.3: **CLAUSE-gadget for a clause $a \vee b \vee \neg c$.** Every green rectangle is an OR-gadget. y -coordinates of $x_{i,0}, y_{i,0}$ and $z_{i,0}$ depend on the variables in the i -th clause. Grey segments corresponds to the values of variables satisfying the i -th clause.

431 **Points.** First, we define auxiliary functions for literals. For a literal w , let $\text{idx}(w)$ be the
 432 index of the variable in w , and $\text{neg}(w)$ be the Boolean value (0 or 1) whether the variable is
 433 negated in w or not.

$$\begin{aligned} \text{idx}(w) &:= i \text{ when } w = x_i \\ \text{neg}(w) &:= \begin{cases} 0 & \text{if } w = x_i \\ 1 & \text{if } w = \neg x_i \end{cases} \end{aligned}$$

434 Let us assume that clause $C_i = a \vee b \vee c$ for any literals a, b, c . Then, we define points in
 435 the gadget as:

$$\begin{aligned}
 436 \quad x_{i,0} &:= (20i, 4 \cdot \text{id}\mathbf{x}(a) + 2 \cdot \text{neg}(c)), & x_{i,1} &:= (20i, 4(n+1)), \\
 y_{i,0} &:= (20i+1, 4 \cdot \text{id}\mathbf{x}(b) + 2 \cdot \text{neg}(b)), & y_{i,1} &:= (20i+1, 4(n+1)+4), \\
 z_{i,0} &:= (20i+2, 4 \cdot \text{id}\mathbf{x}(c) + 2 \cdot \text{neg}(c)), & z_{i,1} &:= (20i+2, 4(n+1)+6).
 \end{aligned}$$

437 We are now ready to define the set of points in a CLAUSE-gadget:

$$\begin{aligned}
 \text{moveVariablePoints}_i &:= \{x_{i,j} : j \in \{0,1\}\} \cup \{y_{i,j} : j \in \{0,1\}\} \cup \{z_{i,j} : j \in \{0,1\}\}, \\
 \text{pointsClause}_i &:= \text{moveVariablePoints}_i \cup \text{pointsOr}_{i,0} \cup \text{pointsOr}_{i,1} \cup \{v_{i,1}\}.
 \end{aligned}$$

438 Note that these two points are equal: $v_{i,0} = l_{i,1}$. This translates to the fact that the
 439 output of the first OR-gadget is an input to the second OR-gadget. This creates an *or* of 3
 440 Boolean values.

441 **Segments.** We also define segments for the CLAUSE-gadget as below:

$$\begin{aligned}
 \text{moveVariableSegments}_i &:= \{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1}), (x_{i,1}, l_{i,0}), (y_{i,1}, p_{i,0}), (z_{i,1}, p_{i,1})\} \\
 \text{segmentsClause}_i &:= \text{moveVariableSegments}_i \cup \text{segmentsOr}_{i,0} \cup \text{segmentsOr}_{i,1}.
 \end{aligned}$$

442 The CLAUSE-gadgets consist of two OR-gadgets. Ideally, we would place the i -th CLAUSE-
 443 gadget close to the $\mathbf{xTrueSegment}_{j_1}$ or $\mathbf{xFalseSegment}_{j_1}$ segments corresponding to the literals
 444 that occur in the i -th clause. It would be inconvenient to position them there, because be-
 445 tween these segments there may be additional $\mathbf{xTrueSegment}_{j_2}$ or $\mathbf{xFalseSegment}_{j_2}$ segments
 446 corresponding to the other literals.

447 Instead, we use simple auxiliary gadgets to *transfer* whether the segment is in a solution,
 448 i.e. segments $(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1})$. Each transfer gadget consists of two segments
 449 $(x_{i,0}, x_{i,1}), (x_{i,1}, a)$. These are the only segments that can cover $x_{i,1}$. We place $x_{i,0}$ on a
 450 segment that we want to transfer (i.e. segment responsible for choosing the variable value
 451 satisfying the corresponding literal). If in some solution $x_{i,0}$ is already covered by this segment,
 452 then we can cover $x_{i,1}$ by $(x_{i,1}, a)$, thus also covering a . If $x_{i,0}$ is not covered by this segment,
 453 then the only way to cover $x_{i,0}$ is to use segment $(x_{i,0}, x_{i,1})$. Intuitively, in any optimal
 454 solution the two segments *transfer* the state of whether $x_{i,0}$ is covered onto whether a is
 455 covered. Therefore, the number of segments in the optimal solution is increased by one, and
 456 we get a point a that was effectively placed on some segment s , but it can be placed anywhere
 457 in the plane instead, consequently simplifying the construction.

458 **Lemma 3.7.** *For any $1 \leq i \leq n$ and $a \in \{x_{i,0}, y_{i,0}, z_{i,0}\}$, there is a set $\text{solClause}_i^{\text{true},a} \subseteq$
 459 segmentsClause_i with $|\text{solClause}_i^{\text{true},a}| = 11$ that covers all points in $\text{pointsClause}_i - \{a\}$.*

460 *Proof.* For $a = x_{i,0}$ (analogous proof for $y_{i,0}$): First we use Lemma 3.5 twice with excluded
 461 $x = l_{i,0}$ and $x = l_{i,1} = v_{i,0}$, resulting with 8 segments in $\text{chooseOr}_{i,0}^{\text{true}} \cup \text{chooseOr}_{i,1}^{\text{true}}$ which
 462 cover all required points apart from $x_{i,1}, y_{i,0}, y_{i,1}, z_{i,0}, z_{i,1}, l_{i,0}$. We cover those using additional
 463 3 segments: $\{(x_{i,1}, l_{i,0}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1})\}$.

464 For $a = z_{0,i}$: Using Lemma 3.6 and Lemma 3.5 with $x = p_{i,1}$, we obtain 8 segments in
 465 $\text{chooseOr}_{i,0}^{\text{false}} \cup \text{chooseOr}_{i,1}^{\text{true}}$ which cover all required points apart from $x_{i,0}, x_{i,1}, y_{i,0}, y_{i,1}, z_{i,1}, p_{i,1}$.
 466 We cover those using additional 3 segments: $\{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,1}, p_{i,1})\}$. \square

467 **Lemma 3.8.** For any $1 \leq i \leq n$ there is a set $\text{solClause}_i^{\text{false}} \subseteq \text{segmentsClause}_i$ with
 468 $|\text{solClause}_i^{\text{false}}| = 12$ that covers all points in pointsClause_i .

469 *Proof.* Using Lemma 3.6 twice we can cover $\text{pointsOr}_{i,0}$ and $\text{pointsOr}_{i,1}$ with 8 segments. To
 470 cover the remaining points we additionally use: $\{(x_{i,0}, x_{i,1}), (y_{i,0}, y_{i,1}), (z_{i,0}, z_{i,1}), (t_{i,1}, v_{i,1})\}$.
 471 \square

472 **Lemma 3.9.** For any $1 \leq i \leq n$:

- 473 (1) points in pointsClause_i can not be covered using any subset of segments from segmentsClause_i
 474 of size smaller than 12;
 475 (2) points in $\text{pointsClause}_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$ can not be covered using any subset of segments
 476 from segmentsClause_i of size smaller than 11.

Proof of (1). No segment in segmentsClause_i covers more than 1 point from

$$\{x_{i,0}, y_{i,0}, z_{i,0}, l_{i,0}, p_{i,0}, q_{i,0}, u_{i,0}, v_{i,0} = l_{i,1}, p_{i,1}, q_{i,1}, u_{i,1}, v_{i,1}\}.$$

477 Therefore we need to use at least 12 segments. \square

Proof of (2). We can define disjoint sets X, Y, Z such that

$$X \cup Y \cup Z \subseteq \text{pointsClause}_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$$

478 and there are no segments in segmentsClause_i covering points from different sets. And we
 479 prove a lower bound for each of these sets. First, let:

$$X := \{x_{i,1}, y_{i,1}, z_{i,1}\}.$$

480 No two points in X can be covered with one segment of segmentsClause_i , so it must be
 481 covered with 3 different segments. Next we define other sets:

$$Y := \text{pointsOr}_{i,0} - \{l_{i,0}, p_{i,0}\},$$

$$Z := \text{pointsOr}_{i,1} - \{l_{i,1}, p_{i,1}\}.$$

482 For both Y and Z we can check all of the subsets of 3 segments of segmentsClause_i to
 483 conclude that none of them cover the considered, so both Y and Z have to be covered with
 484 disjoint sets of 4 segments each.

485 Therefore, $\text{pointsClause}_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$ must be covered with at least $3 + 4 + 4 = 11$
 486 segments from segmentsClause_i . \square

487 3.3.4. Summary

Finally we define the set of points and segments for the constructed instance:

$$\mathcal{C} := \bigcup_{1 \leq i \leq n} \text{pointsVariable}_i \cup \text{pointsClause}_i,$$

$$\mathcal{P} := \bigcup_{1 \leq i \leq n} \text{segmentsVariable}_i \cup \text{segmentsClause}_i.$$

488 **Lemma 3.10** (Robustness to $\frac{1}{2}$ -extension). For every segment $s \in \mathcal{P}$, s and $s^{+\frac{1}{2}}$ cover the
 489 same points from \mathcal{C} .



Figure 3.4: **Scheme of the whole construction.**
General layout of VARIABLE-gadgets and CLAUSE-gadgets and how they interact with each other.

In order to prove this lemma we will define a bounding rectangle R for every gadget, with the following property: R fits both segments and points from the gadget and $R^{\frac{1}{2}}$ (R after $\frac{1}{2}$ -extension) does not cover any points outside of R . Checking that the property from the above lemma holds for points and segments within the same gadget can be easily done using the figures above as references.

Note that the claims stated below also encapsulate the interaction between the gadgets, which are also mentioned in the helper lemmas above, and prove that gadgets are independent otherwise.

First let us define points to cover inside of rectangle R as:

$$\text{points}(R) := \text{points from } \mathcal{C} \text{ that lie in rectangle } R.$$

Claim 3.1. For any $1 \leq i \leq n$, $\text{segmentsVariable}_i$ fit in rectangle defined by points a_i and h_i from *VARIABLE-gadget*:

$$R := [-3L, L] \times [4i, 4i + 2].$$

(1) The only points in R are pointsVariable_i and $x_{j,0}, y_{j,0}$ or $z_{j,0}$ points from *CLAUSE-gadgets*:

$$\text{pointsVariable}_i \subseteq \text{points}(R) \subseteq \text{pointsVariable}_i \cup \{x_{j,0}, y_{j,0}, z_{j,0} : 1 \leq j \leq n\}.$$

(2) R covers the same points from \mathcal{C} before and after $\frac{1}{2}$ -extension, i.e. $\text{points}(R) = \text{points}(R^{+\delta})$.

(3) All segments $\text{segmentsVariable}_i$ fit fully inside of R .

Claim 3.2. For any $1 \leq i \leq n$, pointsVariable_i fit in rectangle from *VARIABLE-gadget*:

$$R := [-3L, -L] \times [4i, 4i + 2].$$

(1) The only points in R are pointsVariable_i : $\text{points}(R) = \text{pointsVariable}_i$.

(2) R covers the same points from \mathcal{C} before and after $\frac{1}{2}$ -extension, i.e. $\text{points}(R) = \text{points}(R^{+\delta})$.

(3) All segments $\text{segmentsVariable}_i - \{\text{xTrueSegment}_i, \text{xFalseSegment}_i\}$ fit fully inside of R .

Claim 3.3. For any $1 \leq i \leq n$ and $j \in \{0, 1\}$, points from *OR-gadget* $\text{pointsOr}_{i,j}$ and segments $\text{segmentsOr}_{i,j} - \{(t_{i,j}, v_{i,j})\}$ fit in rectangle defined as:

$$R := [x, x + 2] \times [y, y + 4], \text{ where } x = 20i + 3j + 3, y = 4(n + 1) + 2j.$$

(1) R covers only $\text{pointsOr}_{i,j}$, i.e. $\text{points}(R) = \text{pointsOr}_{i,j}$.

(2) R covers the same points from \mathcal{C} before and after $\frac{1}{2}$ -extension, i.e. $\text{points}(R) = \text{points}(R^{+\delta})$.

(3) All segments $\text{segmentsOr}_{i,j} - \{(t_{i,j}, v_{i,j})\}$ fit fully inside of R .

Claim 3.4. For any $1 \leq i \leq n$, segmentsClause_i and pointsClause_i fit in rectangle:

$$R := [20i, 20i + 9] \times [0, 4(n + 1) + 6].$$

(1) R covers only pointsClause_i , i.e. $\text{points}(R) = \text{pointsClause}_i$.

(2) R covers the same points from \mathcal{C} before and after $\frac{1}{2}$ -extension, i.e. $\text{points}(R) = \text{points}(R^{+\delta})$.

(3) All segments segmentsClause_i fit fully inside of R .

Proof of Lemma 3.10. First, we check one by one for every segment within every *VARIABLE-gadget* and *OR-gadget* that if it covers some point after $\frac{1}{2}$ -extension, then it covered that point before extension. In other words, every segment does not cover any new point from the same gadget after $\frac{1}{2}$ -extension.

Next, we consider interactions of segments and points from different gadgets.

514 **VARIABLE-gadget** Let us fix $1 \leq i \leq n$ and consider segments from the i -th VARIABLE-
515 gadget. We use Claim 3.1 and name the resulting rectangle R_1 . $\text{segmentsVariable}_i$ do not cover
516 any point outside of R_1 after $\frac{1}{2}$ -extension. However, some points from pointsClause_j for some j
517 can lie within R_1 , hence we use Claim 3.2 and name the resulting rectangle R_2 . R_2 covers only
518 points from pointsVariable_i (even after $\frac{1}{2}$ -extension), then all points from CLAUSE-gadgets
519 inside of R_1 lie on either xTrueSegment_i or xFalseSegment_i , and it is enough to check that these
520 segments cover exactly the same points from CLAUSE-gadgets before and after $\frac{1}{2}$ -extension.
521 They both cover all points from any CLAUSE-gadget that are collinear with these segments,
522 so they cover exactly the same set of points after extension.

523 **CLAUSE-gadget** Let us fix $1 \leq i \leq n$ and consider segments from the i -th CLAUSE-
524 gadget. We use Claim 3.3 for $j \in \{0, 1\}$ to get rectangles R_0 and R_1 respectively. We need to
525 check whether segments $\text{moveVariableSegments}_i \cup \{(t_{i,j}, v_{i,j}) : j \in \{0, 1\}\}$ cover any new points
526 from pointsClause_i after $\frac{1}{2}$ -extension, because their interaction is not considered by Claim 3.3
527 for R_0 and R_1 .

528 Then we use Claim 3.4 to conclude that no segment from segmentsClause_i after $\frac{1}{2}$ -extension
529 covers any point from different CLAUSE-gadget or any VARIABLE-gadget. \square

530 3.4. Proof that the reduction is correct

531 In order to prove Lemma 3.1 we introduce several auxiliary lemmas proving properties of the
532 construction described in the previous section.

533 Consider an instance S of MAX-(3,3)-SAT of size n with optimum solution satisfying
534 k clauses. Let us construct an instance $(\mathcal{C}, \mathcal{P})$ of SEGMENT SET COVER as described in
535 Section 3.3 for the instance S of MAX-(3,3)-SAT.

536 **Lemma 3.11.** *The instance $(\mathcal{C}, \mathcal{P})$ of SEGMENT SET COVER admits a solution of size $15n - k$.*

537 *Proof.* Let the clauses in S be c_1, c_2, \dots, c_n and the variables be x_1, x_2, \dots, x_n . Let the variable
538 assignment in the optimum solution to S be $\phi : \{x_1, x_2, \dots, x_n\} \rightarrow \{\text{true}, \text{false}\}$.

539 We cover every VARIABLE-gadget with solution described in Lemma 3.2, where in the
540 i -th gadget we choose the set of segments corresponding to the value of $\phi(x_i)$.

541 For every clause that is satisfied, say c_i , let us name the variable that is **true** in it as x_i
542 and the point corresponding to x_i in pointsClause_i as a . Points in pointsClause_i are covered
543 with set $\text{solClause}_i^{\text{true}, a}$ described in Lemma 3.7. For every clause that is not satisfied, say c_j ,
544 points in pointsClause_j are covered with set $\text{solClause}_j^{\text{false}}$ described in Lemma 3.8.

545 Formally, we define sets responsible for choosing variable assignment and satisfying clauses,
546 R_i and C_i respectively, as following:

$$\begin{aligned}
 R_i &:= \begin{cases} \text{chooseVariable}_i^{\text{true}} & \text{if } \phi(x_i) = \text{true} \\ \text{chooseVariable}_i^{\text{false}} & \text{if } \phi(x_i) = \text{false} \end{cases} \\
 C_i &:= \begin{cases} \text{solClause}_i^{\text{true}, a} & \text{if } c_i \text{ satisfied by the literal corresponding to point } a \\ \text{solClause}_i^{\text{false}} & \text{if } c_i \text{ not satisfied} \end{cases} \\
 \mathcal{R} &:= \bigcup_{i=1}^n \{R_i \cup C_i : 1 \leq i \leq n\}.
 \end{aligned}$$

547 This set covers all the points from \mathcal{C} , because the sets R_i , C_i individually cover their
548 corresponding gadgets, as proved in the respective lemmas.

549 All of these sets are disjoint, so the size of the obtained solution is:

$$|\mathcal{R}| = \sum_{i=1}^n R_i + \sum_{i=1}^n C_i = 3n + 11k + 12(n - k) = 15n - k. \quad \square$$

550 **Lemma 3.12.** *Suppose we have a solution \mathcal{R} of the instance $(\mathcal{C}, \mathcal{P})$ of SEGMENT SET COVER.*
 551 *Then there exists a solution \mathcal{R}' such that $|\mathcal{R}'| \leq |\mathcal{R}|$ and \mathcal{R}' contains at most one of the*
 552 *segments xTrueSegment_i and xFalseSegment_i from each VARIABLE-gadget.*

553 *Proof.* Assume that we have $\{\text{xTrueSegment}_i, \text{xFalseSegment}_i\} \subseteq \mathcal{R}$ for some i . We will show
 554 how to modify \mathcal{R} into \mathcal{R}' , such that the number of such i decreases, while \mathcal{R}' is still a valid
 555 solution to $(\mathcal{C}, \mathcal{P})$, and $|\mathcal{R}'| \leq |\mathcal{R}|$. Then, by repeating this procedure, we can eventually
 556 construct a solution satisfying the property from the Lemma.

557 To construct \mathcal{R}' , we first remove from \mathcal{R} all segments belonging to $\text{segmentsVariable}_i$.
 558 Recall that the i -th VARIABLE-gadget corresponds to variable x_i in S . As every variable in
 559 S is used in exactly 3 clauses, then one literal x_i or $\neg x_i$ must appear in at least 2 clauses. If
 560 that literal is x_i , then we add to the constructed solution all segments from $\text{chooseVariable}_i^{\text{true}}$,
 561 otherwise we add all segments from $\text{chooseVariable}_i^{\text{false}}$.

562 Now, there exists at most one CLAUSE-gadget which needs adjustment to make \mathcal{R}' valid;
 563 assuming it is the j -th clause, then one of the points $x_{j,0}, y_{j,0}$ or $z_{j,0}$ for this CLAUSE-gadget
 564 might be not covered, say $y_{j,0}$. We amend the solution by adding $(y_{j,0}, y_{j,1})$ to \mathcal{R}' .

565 By Lemma 3.4 we know that \mathcal{R} used at least 4 segments from $\text{segmentsVariable}_i$. Therefore,
 566 we removed at least 4 segments and added at most 4 segments, so $|\mathcal{R}'| \leq |\mathcal{R}|$. \square

567 **Lemma 3.13.** *Suppose we have a solution \mathcal{R} of the instance $(\mathcal{C}, \mathcal{P})$ of SEGMENT SET COVER.*
 568 *Then there exists a solution to S that satisfies at least $15n - |\mathcal{R}|$ clauses.*

569 *Proof.* Let the clauses in S be c_1, c_2, \dots, c_n and the variables be x_1, x_2, \dots, x_n . Given a
 570 solution \mathcal{R} of the instance $(\mathcal{C}, \mathcal{P})$ of SEGMENT SET COVER, we use Lemma 3.12 to modify
 571 \mathcal{R} so that for any i , \mathcal{R} contains at most one of xTrueSegment_i and xFalseSegment_i ; this may
 572 decrease the size of \mathcal{R} , but that does not matter in the subsequent construction. To simplify
 573 notation, in the remainder of this proof we use \mathcal{R} to refer to the modified solution.

Given \mathcal{R} , we construct a solution to S by defining an assignment of variables:

$$\phi : \{x_1, x_2, \dots, x_n\} \rightarrow \{\text{true}, \text{false}\}$$

574 that satisfies at least $15n - |\mathcal{R}|$ clauses in S .

575 **Definition of ϕ .** Recall that due to Lemma 3.12, \mathcal{R} contains at most one of xTrueSegment_i
 576 and xFalseSegment_i .

We define the value $\phi(x_i)$ for the variable x_i as follows:

$$\phi(x_i) := \begin{cases} \text{true} & \text{if } \text{xTrueSegment}_i \in \mathcal{R}, \\ \text{false} & \text{otherwise} \end{cases}$$

577 Moreover, from Lemma 3.3 we get $|\text{segmentsVariable}_i \cap \mathcal{R}| \geq 3$ for every i .

578 **Clauses satisfied with the chosen variable assignment.** For a clause c_i , \mathcal{R} needs
 579 to use at least 11 segments to cover $\text{pointsClause}_i - \{x_{i,0}, y_{i,0}, z_{i,0}\}$ in the i -th CLAUSE-gadget
 580 (Lemma 3.9).

581 Moreover, if none of the points $\{x_{i,0}, y_{i,0}, z_{i,0}\}$ are covered by the segments from $\mathcal{R} \cap$
 582 $\text{segmentsVariable}_i$, then \mathcal{R} needs to cover pointsClause_i with at least 12 segments by Lemma 3.9.

583 Let a be the number of clauses c_i for which none of the points $x_{i,0}, y_{i,0}, z_{i,0}$ in pointsClause_i
 584 are covered by segments from $\mathcal{R} \cap \text{segmentsVariable}_j$ for any $1 \leq j \leq n$.

585 Consider a clause c_i for which at least one of the points $x_{i,0}, y_{i,0}, z_{i,0}$ in pointsClause_i is
 586 covered by segments from $\mathcal{R} \cap \text{segmentsVariable}_j$ for some $1 \leq j \leq n$. Denote this point
 587 as t and say it corresponds to literal q and variable x_j . Point t can be only covered in
 588 $\text{segmentsVariable}_j$ by a corresponding segment xTrueSegment_j or xFalseSegment_j (depending
 589 on whether the literal q is negated or not). From the definition of ϕ and the fact that one of
 590 this segment is in \mathcal{R} , we know that $\phi(j)$ has the value that evaluates q to be **true**. Therefore,
 591 clause c_i is satisfied.

592 Consequently, ϕ satisfies all but at most a clauses in S .

593 To conclude, given a solution \mathcal{R} to $(\mathcal{C}, \mathcal{P})$ we constructed a variable assignment ϕ that
 594 satisfies at least $n - a$ clauses of S . Finally, note that

$$|\mathcal{R}| \geq 3n + 11(n - a) + 12a = 3n + 11n + a = 14n + a,$$

hence

$$15n - |\mathcal{R}| \leq 15n - 14n - a = n - a.$$

595 Therefore, ϕ satisfies at least $15n - |\mathcal{R}|$ clauses of S . □

Chapter 4

Fixed-parameter tractable algorithm for geometric set cover problem

In this chapter we show fixed-parameter tractable algorithms for the geometric set cover problem in two different settings. Section 4.1 shows a fixed-parameter tractable algorithm for geometric set cover with unweighted segments. The remainder of the chapter presents a fixed-parameter tractable algorithm for geometric set cover with weighted segments with δ -extension. We show an algorithm for the setting with δ -extension, because the original problem with weights is W[1]-hard, as we show in Chapter 5.

We start with a shared definition for this problem. We define *extreme points* for a set of collinear points.

Definition 4.1. For a set of collinear points C in the plane, **extreme points** of C are the endpoints of the smallest segment that covers all points from set C .

If C consists of one point or is empty, then there are 1 or 0 extreme points respectively.

4.1. Fixed-parameter tractable algorithm for unweighted segments

In this section we consider fixed-parameter tractable algorithms for unweighted geometric set cover with segments. The setting where segments are required to be axis-parallel (or limited to a constant number of directions) has a trivial FPT algorithm. We present an FPT algorithm for geometric set cover with unweighted segments, where segments are in arbitrary directions.

4.1.1. Axis-parallel segments

Theorem 4.1. (*FPT for segment cover with axis-parallel segments*). There exists an algorithm that given a family \mathcal{P} of axis-parallel segments, a set of points C and a parameter k , runs in time $\mathcal{O}(2^k)$, and outputs a solution $\mathcal{R} \subseteq \mathcal{P}$ such that $|\mathcal{R}| \leq k$ and \mathcal{R} covers all points in C , or determines that such a set \mathcal{R} does not exist.

Proof. We show an $\mathcal{O}(2^k)$ -time branching algorithm. In each step, the algorithm selects a point a which is not yet covered, branches to choose one of the two directions, and greedily chooses a segment a in that direction to cover. This proceeds until either all points are covered or k segments are chosen.

Let us take the point $a = (x_a, y_a)$ which is the smallest among points that are not yet covered in the lexicographic ordering of points in \mathbb{R}^2 . We need to cover a with some of the remaining segments.

Branch over the choice of one of the coordinates (x or y); without loss of generality, let us assume we chose x . Among the segments lying on line $x = x_a$, we greedily add to the solution the one that covers the most points. As a was the smallest in the lexicographical order, all points on the line $x = x_a$ have the y -coordinate larger than y_a . Therefore, if we denote the greedily chosen segment as s , then any other segment on the line $x = x_a$ that covers a can only cover a subset of points covered by s . Thus, greedily choosing s is optimal.

In each step of the algorithm we add one segment to the solution, thus the recursion can be stopped at depth k . If no branch finds a solution, then this means that a solution of size at most k does not exist. \square

Note that the same algorithm can be used for segments in d directions, where we branch over d choices of directions, and it runs in complexity $\mathcal{O}(d^k)$.

4.1.2. Segments in arbitrary directions

In this section we consider the setting where segments are not constrained to a constant number of directions. We present a fixed-parameter tractable algorithm, parameterized by the size of the solution.

Theorem 1.2. (FPT for SEGMENT SET COVER). *There exists an algorithm that given a family \mathcal{P} of segments (in any direction), a set of points \mathcal{C} and a parameter k , runs in time $k^{\mathcal{O}(k)}(|\mathcal{C}| \cdot |\mathcal{P}|)^2$, and outputs a solution $\mathcal{R} \subseteq \mathcal{P}$ such that $|\mathcal{R}| \leq k$ and \mathcal{R} covers all points in \mathcal{C} , or determines that such a set \mathcal{R} does not exist.*

We will need the following lemmas proving properties of any instance of the problem.

Lemma 4.1. *Given an instance $(\mathcal{P}, \mathcal{C})$ of the segment cover problem, without loss of generality we can assume that no segment covers a superset of what another segment covers. That is, for any distinct $A, B \in \mathcal{P}$, we have $A \cap \mathcal{C} \not\subseteq B \cap \mathcal{C}$ and $A \cap \mathcal{C} \not\supseteq B \cap \mathcal{C}$.*

Proof. Assume towards a contradiction that there is an instance $(\mathcal{P}, \mathcal{C})$, and two distinct subsets of \mathcal{P} , A, B , such that $A \cap \mathcal{C} \subseteq B \cap \mathcal{C}$.

We construct a set $\mathcal{P}' := \mathcal{P} - \{A\}$. We prove that for any solution \mathcal{R} of $(\mathcal{P}, \mathcal{C})$, we can construct a solution $\mathcal{R}' \subseteq \mathcal{P}'$, such that $|\mathcal{R}'| \leq |\mathcal{R}|$. Let us take any solution \mathcal{R} of $(\mathcal{P}, \mathcal{C})$. If $A \in \mathcal{R}$, then $\mathcal{R}' := \mathcal{R} \cup \{B\} - \{A\}$, otherwise $\mathcal{R}' := \mathcal{R}$. Let us consider the case when $A \in \mathcal{R}$, because the other case is trivial. Since $A \cap \mathcal{C} \subseteq B \cap \mathcal{C}$, then $\mathcal{R} \cup \{B\} - \{A\}$ covers any point from \mathcal{C} that was covered by \mathcal{R} . Also, $|\mathcal{R} \cup \{B\} - \{A\}| \leq |\mathcal{R}|$. \square

Lemma 4.2. *Given an instance $(\mathcal{P}, \mathcal{C})$ of the segment cover problem transformed by Lemma 4.1, if there exists a line L with at least $k + 1$ points on it, then there exists a subset $A \subseteq \mathcal{P}$, of size at most k , such that every solution \mathcal{R} with $|\mathcal{R}| \leq k$ satisfies $|A \cap \mathcal{R}| \geq 1$. Moreover, such a subset can be found in polynomial time.*

Proof. Let us enumerate the points from \mathcal{C} that lie on L as x_1, x_2, \dots, x_t in the order in which they appear on L . Our proposed set is defined as:

$$A := \{\text{segment collinear with } L \text{ that covers } x_i \text{ and does not cover } x_{i-1} : i \in \{1, \dots, k\}\},$$

where for $i = 1$ we just take a segment that covers x_1 . If such a segment does not exist for any point x as above, then x does not give rise to any segment in A .

664 We prove the lemma by contradiction. Let us assume that there exists a solution \mathcal{R} of
 665 size at most k such that $\mathcal{R} \cap A = \emptyset$.

666 Let \mathcal{R}_L be the set of segments from \mathcal{R} that are collinear with L .

667 Every segment that is not collinear with L can cover at most one of the points that lie
 668 on this line. Hence, if \mathcal{R}_L was empty, then \mathcal{R} would cover at most k points on line L , but L
 669 had at least $k + 1$ different points from \mathcal{C} on it.

670 Therefore, we know that \mathcal{R}_L is not empty and $|\mathcal{R} - \mathcal{R}_L| \leq k - 1$. Segments from $\mathcal{R} - \mathcal{R}_L$
 671 can cover at most $k - 1$ points among $\{x_1, x_2, \dots, x_k\}$, therefore at least one of these points
 672 must be covered by segments from \mathcal{R}_L . We take the leftmost point from $\{x_1, x_2, \dots, x_k\}$ that
 673 is covered in \mathcal{R}_L and name it a . After the transformation from Lemma 4.1, in \mathcal{R} there is only
 674 one segment that starts in a and is collinear with L , therefore this segment must be in both
 675 \mathcal{R} and A . This contradiction concludes the proof that $|A \cap \mathcal{R}| \geq 1$ for any solution \mathcal{R} of size
 676 at most k . \square

677 We are now ready to prove Theorem 1.2.

678 *Proof of Theorem 1.2.* We will prove this theorem by presenting a branching algorithm that
 679 works in desired complexity. It first branches over the choice of segments to cover the lines
 680 with *many* points and then solves a small instance (where every line has at most k points) by
 681 checking all possible solutions.

682 **Algorithm.** We present a recursive algorithm. Given an instance of the problem:

- 683 (1) Use Lemma 4.1 to remove some redundant segments from our instance.
- 684 (2) If there exists a line with at least $k + 1$ points from \mathcal{C} , we branch over the choice of
 685 adding to the solution one of the at most k possible segments provided by Lemma 4.2;
 686 name this segment s and name the set of points from \mathcal{C} that lie on s as S . By recursion,
 687 we find a solution \mathcal{R} for the instance $(\mathcal{C} - S, \mathcal{P} - \{s\})$, and parameter $k - 1$. We return
 688 $\mathcal{R} \cup \{s\}$. Note that if Lemma 4.2 returned \emptyset , then we respond NO.
- 689 (3) If every line has at most k points on it and $|\mathcal{C}| > k^2$, then answer NO.
- 690 (4) If $|\mathcal{C}| \leq k^2$, solve the problem by brute force: check all subsets of \mathcal{P} of size at most k .

691 **Correctness.** Lemma 4.2 proves that at least one segment that we branch over in (1)
 692 must be present in every solution \mathcal{R} with $|\mathcal{R}| \leq k$. Therefore, the recursive call can find
 693 a solution, provided there exists one.

694 In (2) the answer is no, because every line covers no more than k points from \mathcal{C} , which
 695 implies the same about every segment from \mathcal{P} . Under this assumption we can cover only k^2
 696 points with a solution of size k , which is less than $|\mathcal{C}|$.

697 Checking all possible solutions in (3) is trivially correct.

698 **Complexity.** In the leaves of the recursion we have $|\mathcal{C}| \leq k^2$, so $|\mathcal{P}| \leq k^4$, because
 699 every segment can be uniquely identified by the two extreme points it covers (by Lemma 4.1).
 700 Therefore, there are $\binom{k^4}{k}$ possible solutions to check, each can be checked in time $\mathcal{O}(k|\mathcal{C}|)$.
 701 Thus, (3) takes time $k^{\mathcal{O}(k)}$.

702 In this branching algorithm our parameter k is decreased with every recursive call, so we
 703 have at most k levels of recursion with branching over k possibilities. Candidates to branch
 704 over can be found on each level in time $\mathcal{O}((|\mathcal{C}| \cdot |\mathcal{P}|)^{\mathcal{O}(1)})$.

Reduction from Lemma 4.1 can be implemented in time $\mathcal{O}((|\mathcal{C}| \cdot |\mathcal{P}|)^{\mathcal{O}(1)})$.

It follows that the overall complexity is $\mathcal{O}((|\mathcal{C}| \cdot |\mathcal{P}|)^{\mathcal{O}(1)} \cdot k^{\mathcal{O}(k)})$ \square

4.2. Fixed-parameter tractable algorithm for weighted segments with δ -extension

In this section we consider the geometric set cover problem for weighted segments relaxed with δ -extension. We show that this problem admits an FPT algorithm when parameterized by the size of the solution and δ . In the next chapter we show that the assumption about the problem being relaxed with δ -extension is necessary: we prove that geometric set cover problem for weighted segments (without extension) is W[1]-hard, which means there does not exist any FPT algorithm parameterized by solution size for it, assuming $\text{FPT} \neq \text{W}[1]$.

Theorem 1.3. (FPT for WEIGHTED SEGMENT SET COVER with δ -extension). *There exists an algorithm that given a family \mathcal{P} of n weighted segments (in any direction), a set of m points \mathcal{C} , and parameters k and $\delta > 0$, runs in time $f(k, \delta) \cdot (nm)^c$ for some computable function f and a constant c and outputs a set \mathcal{R} such that:*

- $\mathcal{R} \subseteq \mathcal{P}$,
- $|\mathcal{R}| \leq k$,
- $\mathcal{R}^{+\delta}$ covers all points in \mathcal{C} ,
- the weight of \mathcal{R} is not greater than the weight of an optimum solution of size at most k for this problem without δ -extension,

or determines that there is no set \mathcal{R} with $|\mathcal{R}| \leq k$ such that \mathcal{R} covers all points in \mathcal{C} .

4.2.1. Dense subsets

To solve this problem we will introduce a lemma about choosing a *dense* subset of points. A dense subset of points for a set of collinear points C and parameters k and δ is a subset of C such that if we cover it with at most k segments, these segments after δ -extension will cover all of the points from C . We will prove that such set of size bounded by some function $f(k, \delta)$ always exists (Lemma 4.3). Later, Lemma 4.3 will allow us to find a kernel for our original problem.

Definition 4.2. For a set of collinear points C , a subset $A \subseteq C$ is (k, δ) -**dense** if for any set of segments R that covers A and such that $|R| \leq k$, it holds that $R^{+\delta}$ covers C .

Lemma 4.3. *For any set of collinear points C , $\delta > 0$ and $k \geq 1$, there exists a (k, δ) -dense set $A \subseteq C$ of size at most $(2 + \frac{2}{\delta})^k$. Moreover, there exists an algorithm that computes the (k, δ) -dense set in time $\mathcal{O}(|C| \cdot (2 + \frac{2}{\delta})^k)$.*

Proof. We prove this for a fixed δ by induction on k .

Inductive hypothesis. For any set of collinear points C , there exists a set A such that:

- A is subset of C ,
- A is (ℓ, δ) -dense for every $1 \leq \ell \leq k$,
- $|A| \leq (2 + \frac{2}{\delta})^k$,
- the extreme points of C are in A .

743 **Base case for $k = 1$.** It is sufficient that A consists of the extreme points of C .
744 If they are covered with one segment, it must be a segment that includes the extreme
745 points from C , so it covers the whole set C .
746 There are at most 2 extreme points in C and $2 < 2 + \frac{2}{\delta}$.

747 **Inductive step.** Assuming inductive hypothesis for any set of collinear points C and
748 for parameter k , we will prove it for $k + 1$.
749 Let s be the minimal segment that includes all points from C . That is, the extreme points
750 of C are endpoints of s .
751 We define $M = \lceil 1 + \frac{2}{\delta} \rceil$ subsegments of s by splitting s into M closed segments of equal
752 length. We name these segments v_i , note that $|v_i| = \frac{|s|}{M}$ for each $1 \leq i \leq M$.
753 Let C_i be the subset of C consisting of points lying on v_i .
754 Let t_i be the segment with endpoints being the extreme points of C_i . It might be a
755 degenerate segment if C_i consists of one point, or t_i might be empty if C_i is empty.
756 Figure 4.1 presents an example of such segments v_i and t_i .




Figure 4.1: **Example of segments v_i and t_i .**

Example for $M = 7$ and some set of points (marked with black circles). The top panel shows segments v_i and the bottom panel shows segments t_i on the same set of points. a and b are the extreme points and therefore segment s ends at a and b . Red segments depict the split into M segments of equal length v_i . Blue segments depict the segments t_i . t_5 is an empty segment, because there are no points that lie on segment v_5 . Segments t_3 and t_7 are degenerated to one point – c and d , respectively. Segments t_1 and t_2 share one point b .

757 We use the inductive hypothesis to choose (k, δ) -dense sets A_i for sets C_i . Note that if
758 $|C_i| \leq 1$, then $A_i = C_i$ and it is still a (k, δ) -dense set for C_i .
759 Then we define $A = \bigcup_{i=1}^M A_i$. Thus A includes the extreme points of C , because they are
760 included in the sets A_1 and A_M .

The size of each A_i is at most $(2 + \frac{2}{\delta})^k$ from the inductive hypothesis, therefore size of A is at most:

$$M \left(2 + \frac{2}{\delta}\right)^k = \left\lceil 1 + \frac{2}{\delta} \right\rceil \cdot \left(2 + \frac{2}{\delta}\right)^k \leq \left(2 + \frac{2}{\delta}\right)^{k+1}.$$

761 **Proof that A is $(k+1, \delta)$ -dense for C .** Let us take any cover of A with $k+1$ segments
762 and call it \mathcal{R} .

763 For every segment t_i , if there exists a segment x in \mathcal{R} that is disjoint with t_i , then we have
764 a cover of A_i with at most k segments using $\mathcal{R} - \{x\}$. Since A_i is (k, δ) -dense for t_i and C_i ,
765 $(\mathcal{R} - \{x\})^{+\delta}$ covers C_i . So $\mathcal{R}^{+\delta}$ covers C_i as well.
766 If there exists a segment t_i for which a segment x as defined above does not exist, then
767 all $k + 1$ segments that cover A_i intersect t_i . An example of such segments is depicted in
768 Figure 4.2. Let us consider any such t_i . By the inductive hypothesis, the endpoints of s
769 are in A_1 and A_M respectively, so \mathcal{R} must cover them. For each endpoint of s , there exists
770 a segment that contains this endpoint and intersects t_i . Let us call these two segments y
771 and z . It follows that: $|y| + |z| + |t_i| \geq |s|$. Since $|t_i| \leq |v_i| = \frac{|s|}{M} \leq \frac{|s|}{1+\frac{2}{\delta}} = \frac{|s|\delta}{\delta+2}$, we have
772 $\max(|y|, |z|) \geq |s|(1 - \frac{\delta}{\delta+2})/2 = \frac{|s|}{\delta+2}$.



Figure 4.2: **Example of all $k + 1$ segments intersecting one segment t_i .**
Both panels show the same set \mathcal{C} (black circles), the same as in Figure 4.1. The top panel shows blue segments t_i for $M = 7$. The bottom panel shows green segments – solution \mathcal{R} of size 4. All segments from \mathcal{R} intersect t_4 . Segments z and y are named in the figure.

After δ -extension, the longer of these segments will expand at both ends by at least:

$$\max(|y|, |z|)\delta \geq \frac{|s|\delta}{\delta+2} = \frac{|s|}{1+\frac{2}{\delta}} \geq \frac{|s|}{M} = |v_i| \geq |t_i|.$$

773 Therefore, the longer of segments y and z will cover the whole segment t_i after δ -extension.
774 We conclude that $\mathcal{R}^{+\delta}$ covers C_i .
775 Since $C = \bigcup_{i=1}^M C_i$, it follows that $\mathcal{R}^{+\delta}$ covers C .

Algorithm. We can simulate the inductive proof presented above by a recursive algorithm with the following complexity:

$$O\left(|C| + \frac{1}{\delta}\right) + O\left(|C| \cdot \left(2 + \frac{2}{\delta}\right)^k\right).$$

776

□

777 4.2.2. Algorithm

778 Let us now formulate some claims about the properties for the problem parameterized by the
779 solution size. These properties provide bounds for different objects in the problem instance,
780 which help us to find a small kernel for the problem or conclude that the optimum solution
781 to this instance must be, in terms of size, above some threshold.

782 **Definition 4.3.** A line in the plane is **long** if there are at least $k + 1$ points from \mathcal{C} on it.

783 **Claim 4.1.** *If there are more than k different long lines, then \mathcal{C} can not be covered with k*
 784 *segments.*

785 *Proof.* We prove the claim by contradiction. Let us assume that we have at least $k + 1$ different
 786 long lines in our instance of the problem and there is a solution \mathcal{R} of size at most k covering
 787 points \mathcal{C} .

788 Choose any long line L . Every segment from \mathcal{R} which is not collinear with L , covers at
 789 most one point that lies on L . L is long, so there are at least $k + 1$ points from \mathcal{C} that lie on
 790 L . This implies that there must be a segment in \mathcal{R} that is collinear with L .

791 Since we have at least $k + 1$ different long lines, there are at least $k + 1$ segments in \mathcal{R}
 792 collinear with different lines. This contradicts with the assumption that $|\mathcal{R}| \leq k$. \square

793 **Claim 4.2.** *If there are more than k^2 points from \mathcal{C} that do not lie on any long line, then \mathcal{C}*
 794 *can not be covered with k segments.*

795 *Proof.* We prove the claim by contradiction. Let us assume that we have at least $k^2 + 1$ points
 796 from \mathcal{C} that do not lie on any long line, call this set A , and a solution \mathcal{R} of size at most k
 797 covering all points in \mathcal{C} .

798 Every segment s from \mathcal{R} covers at most k points from A . This is because if s covered at
 799 least $k + 1$ points from A , then the line in the direction of s would be a long line and that
 800 contradicts the definition of A .

801 If every segment from \mathcal{R} covers at most k points from A and $|\mathcal{R}| \leq k$, then at most k^2
 802 points from A are covered by \mathcal{R} and that contradicts the fact that \mathcal{R} is a solution to the given
 803 geometric set cover instance. \square

804 We are now ready to give a proof of Theorem 1.3.

805 *Proof of Theorem 1.3.* Our goal is to either answer NO or to find a kernel $(\mathcal{C}', \mathcal{P}')$ of size
 806 bounded by $f(k)$ for some function f , such that:

- 807 • (*Property 1*) for every solution \mathcal{R} to $(\mathcal{C}, \mathcal{P})$ of size at most k , there exists a set $\mathcal{R}_1 \subseteq \mathcal{P}'$
 808 such that $|\mathcal{R}_1| \leq k$, the weight of \mathcal{R}_1 is not greater than the weight of \mathcal{R} , and \mathcal{R}_1 covers
 809 \mathcal{C}' ;
- 810 • (*Property 2*) for every set $\mathcal{R}_2 \subseteq \mathcal{P}'$ such that $|\mathcal{R}_2| \leq k$ and \mathcal{R}_2 covers all points in \mathcal{C}' ,
 811 $\mathcal{R}_2^{+\delta}$ covers all points in the original set \mathcal{C} .

812 If we found such sets $(\mathcal{C}', \mathcal{P}')$, using *Property 1* we know that an optimum solution of size
 813 at most k to $(\mathcal{C}', \mathcal{P}')$ has no greater weight than an optimum solution of size at most k to
 814 $(\mathcal{C}, \mathcal{P})$. Using *Property 2* we know that any solution to $(\mathcal{C}', \mathcal{P}')$ after δ -extension covers \mathcal{C} .

815 Therefore, finding such sets and solving the instance $(\mathcal{C}', \mathcal{P}')$ by iterating over all of the
 816 subsets of \mathcal{P}' of size at most k in desired complexity is sufficient to prove Theorem 1.3.

817 **Definition of \mathcal{C}' and \mathcal{P}' .** Let us name the number of different long lines as l . Applying
 818 Claims 4.1 and 4.2, if we have more than k different long lines or more than k^2 points from
 819 \mathcal{C} that do not lie on any long line, then we answer NO, because these lemmas prove that there
 820 is no solution of size at most k to this instance.

821 Otherwise, we can split \mathcal{C} into at most $k + 1$ sets:

- 822 • D : points that do not lie on any long line, $|D| \leq k^2$;

823 • C_i for $1 \leq i \leq l$: points that lie on the i -th long line, $|C_i| > k$.

824 Note that sets C_i do not need to be disjoint.

825 Then, for every set C_i we can use Lemma 4.3 to obtain a (k, δ) -dense set A_i for C_i with
826 $|A_i| \leq (2 + \frac{2}{\delta})^k$.

827 We define $\mathcal{C}' := D \cup (\bigcup A_i)$. \mathcal{C}' has size at most $k^2 + k(2 + \frac{2}{\delta})^k$. We define \mathcal{P}' as
828 follows: for every pair of points \mathcal{C}' , we choose one segment from \mathcal{P} that has the lowest weight
829 among segments that cover these points or decide that there is no segment that covers them.
830 There are at most $|\mathcal{C}'|^2$ different segments in \mathcal{P}' , therefore both \mathcal{P}' and \mathcal{C}' have size bounded
831 by $\mathcal{O}((k^2 + k(2 + \frac{2}{\delta}))^2)$.

832 **Proof of Property 2.** Firstly, we prove that for every set $\mathcal{R}_2 \subseteq \mathcal{P}'$ such that $|\mathcal{R}_2| \leq k$ and
833 \mathcal{R}_2 covers points in \mathcal{C}' , $\mathcal{R}_2^{+\delta}$ covers points in the original instance \mathcal{C} .

834 Let us take such a set \mathcal{R}_2 .

835 \mathcal{C} is partitioned into several parts – sets D and C_i . Points from D are covered by \mathcal{R}_2 ,
836 because D is part of \mathcal{C}' . Each point from any A_i is covered, because A_i is a part of \mathcal{C}' ; A_i is a
837 (k, δ) -dense set for C_i , therefore $\mathcal{R}_2^{+\delta}$ covers all points in C_i . Therefore, $\mathcal{R}_2^{+\delta}$ covers all points
838 in \mathcal{C} .

839 **Proof of Property 1.** Secondly, we prove that for every solution \mathcal{R} to $(\mathcal{C}, \mathcal{P})$ of size at
840 most k , there exists a set $\mathcal{R}_1 \subseteq \mathcal{P}'$ such that $|\mathcal{R}_1| \leq k$, the weight of \mathcal{R}_1 is not greater than
841 the weight of \mathcal{R} and \mathcal{R}_1 covers \mathcal{C}' .

842 For every segment in \mathcal{R} , say s , let us look at the points from \mathcal{C}' that lie on s and call this
843 set of points F . F is of course a set of collinear points. We can cover F with any segment
844 that covers extreme points of F , because all other points lie on the segment between these
845 points. Therefore, we can replace s with a segment s' that has lowest weight among the points
846 that cover the extreme points of F . Such a segment belongs to \mathcal{P}' , because this is how it was
847 defined. Segment s' has weight no greater than the weight of s , because s also covers F .

848 Therefore, we produced the set \mathcal{R}_1 that has size not greater than the size of \mathcal{R} (because
849 some segments s can map to the same segment s'), weight not greater than \mathcal{R} , and it covers
850 \mathcal{C}' .

851 **Complexity** We find a solution of $(\mathcal{C}', \mathcal{P}')$ by iterating over all the possible subsets of
852 \mathcal{P}' . Finding sets \mathcal{P}' and \mathcal{C}' and then solving problem for kernel has overall complexity
853 $(|\mathcal{P}| + |\mathcal{C}|)^{\mathcal{O}(1)} \mathcal{O}((2 + \frac{2}{\delta})^k) + \mathcal{O}((k^2 + k(2 + \frac{2}{\delta})^k)^k)$. \square

Chapter 5

W[1]-hardness for axis-parallel weighted segments

In this chapter we consider the geometric set cover problem with axis-parallel or right-diagonal weighted segments. In Theorem 1.4 below, we prove that this problem is W[1]-hard when parameterized by the size of the solution.

We believe that the below construction can be improved to only utilize the axis-parallel segments.

Theorem 1.4. (WEIGHTED SEGMENT SET COVER is W[1]-hard). *Consider the problem of covering a set \mathcal{C} of points by selecting at most k segments from a set of segments \mathcal{P} with non-negative weights $w : \mathcal{P} \rightarrow \mathbb{R}^+$ so that the weight of the cover is minimal. Then this problem is W[1]-hard when parameterized by k and assuming ETH, there is no algorithm for this problem with running time $f(k) \cdot (|\mathcal{C}| + |\mathcal{P}|)^{o(\sqrt{k})}$ for any computable function f . Moreover, this holds even if all segments in \mathcal{P} are axis-parallel or right-diagonal.*

5.1. Grid Tiling

In order to prove Theorem 1.4 we will show a reduction from a W[1]-hard problem: grid tiling. This problem was introduced in [Marx, 2007] (the author called it matrix tiling instead). It was originally described as an approximation problem, but W[1]-hardness follows directly from the theorems stated there. For a more contemporary description of this problem and a proof of W[1]-hardness, see Chapter 14 of [Cygan et al., 2015].

Definition 5.1. We define the **powerset** of a set A , denoted as $\text{Pow}(A)$, as the set of all subsets of A , i.e. $\text{Pow}(A) = \{B : B \subseteq A\}$.

Definition 5.2. In the **grid tiling** problem we are given integers n and k , and a function $f : \{1, \dots, k\} \times \{1, \dots, k\} \rightarrow \text{Pow}(\{1, \dots, n\} \times \{1, \dots, n\})$ specifying the set of allowed tiles for each cell of a $k \times k$ grid. The task is to decide whether there exist functions $x, y : \{1, \dots, k\} \rightarrow \{1, \dots, n\}$ that assign colors from $\{1, \dots, n\}$ to respectively columns and rows of the grid, so that $(x(i), y(j)) \in f(i, j)$ for all $i, j \in \{1, \dots, k\}$.

In short, in the grid tiling problem one needs to assign numbers to rows and columns in such a way that for every pair of a row and a column, the pair of colors assigned to the row and column belongs to the allowed set of tiles for this pair. The next theorem describes the complexity of this problem, which is W[1]-hard when parameterized by the size of the grid.

	$x(1) = 3$	$x(2) = 1$	$x(3) = 3$	$x(4) = 7$
$y(4) = 1$	$(\mathbf{2}, \mathbf{1}); (2, 2);$ $(\mathbf{3}, \mathbf{1}); (3, 9)$	$(1, 1); (3, 1)$	$(\mathbf{3}, \mathbf{1}); (7, 2)$	$(\mathbf{2}, \mathbf{1}); (\mathbf{7}, \mathbf{1})$
$y(3) = 1$	$(\mathbf{2}, \mathbf{1}); (\mathbf{3}, \mathbf{1});$ $(4, 2); (8, 2)$	$(1, 1); (1, 3)$	$(\mathbf{3}, \mathbf{1}); (4, 3)$	$(\mathbf{2}, \mathbf{2}); (\mathbf{7}, \mathbf{1})$
$y(2) = 6$	$(\mathbf{2}, \mathbf{6}); (\mathbf{3}, \mathbf{6})$	$(1, 2); (\mathbf{1}, \mathbf{6});$ $(2, 6)$	$(2, 6); (\mathbf{3}, \mathbf{6})$	$(\mathbf{2}, \mathbf{6}); (\mathbf{7}, \mathbf{6})$
$y(1) = 4$	$(\mathbf{2}, \mathbf{4}); (2, 6);$ $(\mathbf{3}, \mathbf{4}); (\mathbf{3}, \mathbf{9})$	$(1, 4); (\mathbf{1}, \mathbf{9})$	$(\mathbf{3}, \mathbf{4}); (\mathbf{3}, \mathbf{9})$	$(\mathbf{2}, \mathbf{9}); (\mathbf{7}, \mathbf{4})$

Figure 5.1: **Example of a grid tiling instance and its solution.**

In the first row and column of the table you can see the solution: functions x and y . The tiles used in this solution are marked in **bold**. If we instead chose the tiles marked in **blue** (whenever there is one, taking the tile marked in **bold** otherwise), then that corresponds to setting $x(1) = 2$, and would also form a correct solution. On the other hand, if we instead chose the tiles marked in **red** (as before), then this corresponds to setting $y(1) = 9$ and $x(4) = 2$ and that would **not** form a correct solution. Even though the first row is correct, the cell with coordinates $(3, 4)$ requires tile $(2, 1)$, not $(2, 2)$ (marked in **bold red**).

Theorem 5.1. [Marx, 2007] *Grid tiling is $W[1]$ -hard when parameterized by k and assuming ETH, there is no $f(k) \cdot n^{o(k)}$ -time algorithm solving the grid tiling problem for any computable function f .*

The remainder of this section is devoted to proving Theorem 1.4 by a reduction from a grid tiling problem instance with parameter k (number of rows in the grid) to a geometric set cover instance with parameter k^2 (size of solution). This reduction is described in Lemma 5.1. This proves the $W[1]$ -hardness of the geometric set cover problem, because if we could solve it with an FPT algorithm, then we could also solve the grid tiling problem (which we reduced to the geometric set cover). Therefore, geometric set cover with setting described in Theorem 1.4 is at least as hard as the grid tiling problem.

5.2. Statement of reduction

Let us denote an instance of grid tiling problem as (n, k, f) consisting of:

- the number of colors n ,
- the size of the grid k ,
- the function specifying the allowed tiles $f : \{1, \dots, k\} \times \{1, \dots, k\} \rightarrow \text{Pow}(\{1, \dots, n\} \times \{1, \dots, n\})$.

Let us also define constants:

$$\begin{aligned} \epsilon &:= \frac{1}{2k^2} \\ \delta &:= \frac{1}{4k^4} \\ W_{\text{hv}} &:= 2k^2(n^2 + 1) - 4k^2\epsilon - 4k(1 - \epsilon) \end{aligned}$$

which are going to be used when defining the weight of the constructed instance of geometric set cover with weighted segments.

904 **Lemma 5.1.** *Given an instance (n, k, f) of the grid tiling problem, we can construct an*
 905 *instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ of geometric set cover with weighted segments such that:*

- 906 (1) *if the answer to (n, k, f) is YES, then there exists a solution to $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ of*
 907 *weight at most $W_{\text{hv}} + k^2\delta$;*
- 908 (2) *if there exists a solution to $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ of weight at most $W_{\text{hv}} + k^2\delta$, then the*
 909 *answer to (n, k, f) is YES.*

910 First, let us prove Theorem 1.4 using Lemma 5.1.

911 *Proof of Theorem 1.4.* Let us take any instance (n, l, f) of the grid tiling problem. We prove
 912 the theorem by contradiction, therefore we assume that geometric set cover with weighted
 913 segments parameterized by solution size k admits a $g(k) \cdot n^{o(\sqrt{k})}$ -time algorithm for some
 914 computable function g .

915 Using Lemma 5.1 let us construct an instance I for (n, l, f) . Let us assume that the
 916 optimum solution of size at most k to the instance I has weight u . Using (2) we know that if
 917 $u \leq W_{\text{hv}} + k^2\delta$, then the answer to (n, l, f) is YES. If $u > W_{\text{hv}} + k^2\delta$, then using (1) we know
 918 that the answer to (n, l, f) must be NO.

919 Therefore if we could find the solution in time $g(k) \cdot n^{o(\sqrt{k})}$, then we could solve the grid
 920 tiling problem in time $g(l) \cdot n^{o(l)}$ by constructing an instance of the set cover with weighted
 921 segments, solving it for parameter $k = 3l^2 + 2l$ in time $n^{o(\sqrt{3l^2+2l})}$ and then answering based
 922 on the weight of the optimum solution. As $\mathcal{O}(n^{o(l)}) \subseteq \mathcal{O}(n^{o(\sqrt{3l^2+2l})})$, the existence of this
 923 algorithm contradicts Theorem 5.1. Hence such an algorithm can not exist. \square

924 We prove Lemma 5.1 in subsequent sections. First, we define a constructed instance I ,
 925 later property (1) is proved by Lemma 5.2 and property (2) is proved by Lemma 5.6.

926 In the proof of Lemma 5.6 we do not use the assumption that the solution is bounded
 927 by the size, which the problem is parameterized by, $3k^2 + 2k$. If we had a permissive FPT
 928 algorithm that finds a solution of any size that still has weight no more than $W_{\text{hv}} + k^2\delta$, then
 929 we still would have a contradiction with grid tiling being W[1]-hard in proof of Theorem 1.4.
 930 Thus, this reduction proves that the problem is not only W[1]-hard, but assuming ETH there
 931 also does not exist permissive FPT algorithm for this problem. Formally we state this in the
 932 Theorem 5.2.

933 **Theorem 5.2. (Permissive FPT does not exist).** *Consider the problem of covering a*
 934 *set \mathcal{C} of points using segments from a set \mathcal{P} with non-negative weights $w : \mathcal{P} \rightarrow \mathbb{R}^+$ so that*
 935 *the weight of the cover is minimal. Let \mathcal{R}^k be the optimum solution to this problem of size at*
 936 *most k . The task is to find a solution \mathcal{R} of any size such that weight of \mathcal{R} is not greater than*
 937 *the weight of \mathcal{R}^k .*

938 *Assuming ETH, there is no algorithm for this problem with running time $f(k) \cdot (|\mathcal{C}| +$
 939 $|\mathcal{P}|)^{o(\sqrt{k})}$ for any computable function f . Moreover, this holds even if all segments in \mathcal{P} are*
 940 *axis-parallel or right-diagonal.*

941 5.3. Construction of the Geometric Set Cover instance

942 We construct an instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ of geometric set cover as follows.

943 First, let us choose any bijection $\text{order} : \{1, \dots, n^2\} \rightarrow \{1, \dots, n\} \times \{1, \dots, n\}$.

Define $\text{match}_v(i, j)$ and $\text{match}_h(i, j)$ as Boolean functions denoting whether two points share x or y coordinate:

$\text{match}_v(i, j)$ is **true** \iff $\text{order}(i)$ and $\text{order}(j)$ have the same x coordinate,

$\text{match}_h(i, j)$ is **true** \iff $\text{order}(i)$ and $\text{order}(j)$ have the same y coordinate.

944 5.3.1. Points

For $1 \leq i, j \leq k$ and $1 \leq t \leq n^2$ define points:

$$h_{i,j,t} := (i \cdot (n^2 + 1) + t, j \cdot (n^2 + 1)),$$

$$v_{i,j,t} := (i \cdot (n^2 + 1), j \cdot (n^2 + 1) + t).$$

Let us define sets H and V as:

$$H := \{h_{i,j,t} : 1 \leq i, j \leq k, 1 \leq t \leq n^2\},$$

$$V := \{v_{i,j,t} : 1 \leq i, j \leq k, 1 \leq t \leq n^2\}.$$

Let us recall that $\epsilon = \frac{1}{2k^2}$. For a point $p = (x, y)$ we define points:

$$p^L := (x - \epsilon, y),$$

$$p^R := (x + \epsilon, y),$$

$$p^U := (x, y + \epsilon),$$

$$p^D := (x, y - \epsilon).$$

Then we define the point set as follows:

$$\mathcal{C} := H \cup \{p^L : p \in H\} \cup \{p^R : p \in H\} \cup V \cup \{p^U : p \in V\} \cup \{p^D : p \in V\}.$$

945 **Definition 5.3.** For every point $p \in H$, we name point p^L its **left guard** and point p^R its
946 **right guard**.

947 Similarly for every points $p \in V$, we name point p^D its **lower guard** and point p^U its
948 **upper guard**.

949 5.3.2. Segments

950 For $1 \leq i, j \leq k$ and $1 \leq t, t_1, t_2 \leq n^2$ define segments:

$$\text{hor}_{i,j,t_1,t_2} := (h_{i,j,t_1}^R, h_{i+1,j,t_2}^L),$$

$$\text{ver}_{i,j,t_1,t_2} := (v_{i,j,t_1}^U, v_{i,j+1,t_2}^D),$$

$$\text{horBeg}_{i,t} := (h_{1,i,1}^L, h_{1,i,t}^L),$$

$$\text{horEnd}_{i,t} := (h_{k,i,t}^R, h_{k,i,n^2}^R),$$

$$\text{verBeg}_{i,t} := (v_{i,1,1}^D, v_{i,1,t}^D),$$

$$\text{verEnd}_{i,t} := (v_{i,k,t}^U, v_{i,k,n^2}^U).$$

951 Next, we define sets of vertical and horizontal segments:

$$\begin{aligned} \text{HOR} &:= \{ \text{hor}_{i,j,t_1,t_2} : 1 \leq i < k, 1 \leq j \leq k, 1 \leq t_1, t_2 \leq n^2, \text{match}_h(t_1, t_2) \text{ holds} \} \\ &\cup \{ \text{horBeg}_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2 \} \\ &\cup \{ \text{horEnd}_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2 \}, \end{aligned}$$

952

$$\begin{aligned} \text{VER} &:= \{ \text{ver}_{i,j,t_1,t_2} : 1 \leq i \leq k, 1 \leq j < k, 1 \leq t_1, t_2 \leq n^2, \text{match}_v(t_1, t_2) \text{ holds} \} \\ &\cup \{ \text{verBeg}_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2 \} \\ &\cup \{ \text{verEnd}_{i,t} : 1 \leq i \leq k, 1 \leq t \leq n^2 \}. \end{aligned}$$

953 An example is depicted in Figure 5.3.

Finally, we also define a set of right-diagonal segments:

$$\text{DIAG} := \{ (h_{i,j,t}, v_{i,j,t}) : 1 \leq i, j \leq k, 1 \leq t \leq n^2, \text{order}(t) \in f(i, j) \}.$$

954 An example of such segments is depicted in Figure 5.2.

955 Every segment in **DIAG** connects points $(i(n^2+1)+t, j \cdot (n^2+1))$ and $(i \cdot (n^2+1), j(n^2+1) + t)$
 956 for some $1 \leq i, j \leq k, 1 \leq t \leq n^2$. The line on which it lies can be described by linear equation
 957 $x + y = t + (i + j)(n^2 + 1)$, thus these segments are in fact right-diagonal.

958 The constructed segment set is defined as:

$$\mathcal{P} := \text{HOR} \cup \text{VER} \cup \text{DIAG}.$$

959 The weight of each segment in $\text{HOR} \cup \text{VER}$ is equal to its length, while every segment in
 960 **DIAG** has weight δ .

$$w(s) = \begin{cases} \text{length}(s) & \text{if } s \in \text{HOR} \cup \text{VER} \\ \delta & \text{if } s \in \text{DIAG} \end{cases}$$

961 5.4. Proof that the reduction is correct

962 Now, we prove that the constructed instance of geometric set cover with weighted segments
 963 indeed gives a correct and sound reduction of the grid tiling problem. Lemma 5.2 proves that
 964 if a solution to the instance of the grid tiling instance exists, then there exists a solution with
 965 suitably bounded size and weight of the constructed instance of geometric set cover. Then
 966 Lemma 5.6 proves that if there is a solution to the geometric set cover instance with bounded
 967 weight, then there exists a solution to the original grid tiling instance.

968 **Lemma 5.2.** *If there exists a solution to the grid tiling instance $(f_{i,j})$, then there exists*
 969 *a solution to the instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ of geometric set cover with weight $W_{\text{hv}} + k^2\delta$.*

970 *Proof.* Suppose there exists a solution x, y of the instance $(f_{i,j})$ of the grid tiling problem.

971 We define the proposed solution $\mathcal{R} \subseteq \mathcal{P}$ of the instance of geometric set cover in three



Figure 5.2: **Vertices and segments in DIAG.**

This is an example of constructed points any $1 \leq i, j \leq k$. Points from H and V are marked in black, their guards are marked in blue. You can also see segments from DIAG with their weights (equal to δ).



Figure 5.3: **Vertices and segments in HOR.**

This is an example for $n = 2$ and any $1 \leq j \leq k$. Points from H are marked in black, their guards are marked in light blue. $t_{i,j}$ is a notation that we use for $\text{order}^{-1}(i, j)$. Segments are represented as arcs between endpoints. You can see $\text{horBeg}_{j,t}$ segments in red. $\text{horBeg}_{j,1}$ is degenerated to a single point at $h_{1,1,t_{1,1}}^L$. Segments $\text{hor}_{i,j,t_{x_1,y},t_{x_2,y}}$ are marked in blue and green. Blue segments connect $t_{x_1,y}$ and $t_{x_2,y}$ such that they share y-coordinate equal to 1, for green segments it is equal to 2.

972 parts: $D \subseteq \text{DIAG}$, $A \subseteq \text{HOR}$ and $B \subseteq \text{VER}$:

$$\begin{aligned}
 D &:= \{(v_{i,j,t}, h_{i,j,t}) : 1 \leq i, j \leq k, t = \text{order}^{-1}(x(i), y(j))\}, \\
 A &:= \{\text{horBeg}_{i, \text{order}^{-1}(x(1), y(i))} : 1 \leq i \leq k\} \\
 &\quad \cup \{\text{horEnd}_{i, \text{order}^{-1}(x(k), y(i))} : 1 \leq i \leq k\} \\
 &\quad \cup \{\text{hor}_{i,j, \text{order}^{-1}(x(i), y(j)), \text{order}^{-1}(x(i+1), y(j))} : 1 \leq i < k, 1 \leq j \leq k\}, \\
 B &:= \{\text{verBeg}_{i, \text{order}^{-1}(x(i), y(1))} : 1 \leq i \leq k\} \\
 &\quad \cup \{\text{verEnd}_{i, \text{order}^{-1}(x(i), y(k))} : 1 \leq i \leq k\} \\
 &\quad \cup \{\text{ver}_{i,j, \text{order}^{-1}(x(i), y(j)), \text{order}^{-1}(x(i), y(j+1))} : 1 \leq i \leq k, 1 \leq j < k\}, \\
 \mathcal{R} &:= D \cup A \cup B.
 \end{aligned}$$

973 Since $\mathcal{C} = H \cup V$, we show that \mathcal{R} covers the whole set H ; the proof for V is analogous.

974 Fix any $1 \leq j \leq k$ and define $t_i := \text{order}^{-1}(x(i), y(j))$. The two leftmost segments in A
 975 for this j are $\text{horBeg}_{j,t_1} = (h_{1,j,1}^L, h_{1,j,t_1}^L)$ and $\text{hor}_{1,j,t_1,t_2} = (h_{1,j,t_1}^R, h_{2,j,t_2}^L)$. Therefore, points
 976 $h_{1,j,x}, h_{1,j,x}^L$ and $h_{1,j,x}^R$ for all $1 \leq x \leq n^2$ are covered by horBeg_{j,t_1} and hor_{1,j,t_1,t_2} , excluding
 977 point h_{1,j,t_1} .

978 Analogously for $2 \leq i \leq k-1$, the two consecutive segments $\text{hor}_{i-1,j,t_{i-1},t_i}$ and $\text{hor}_{i,j,t_i,t_{i+1}}$
 979 cover points $h_{i,j,x}, h_{i,j,x}^L$ and $h_{i,j,x}^R$ for all $1 \leq x \leq n^2$, excluding point h_{i,j,t_i} .

980 Finally $\text{hor}_{k-1,j,t_{k-1},t_k}$ and horEnd_{j,t_k} cover all points $h_{k,j,x}, h_{k,j,x}^L$ and $h_{k,j,x}^R$ for $1 \leq x \leq n^2$,
 981 excluding point h_{k,j,t_k} .

982 D covers all points h_{i,j,t_i} and v_{i,j,t_i} . As j was chosen arbitrarily, all points in H are covered.
 The size of this proposed solution is:

$$|\mathcal{R}| = |D| + |A| + |B| = k^2 + (k+1)k + (k+1)k = 3k^2 + 2k.$$

983 Then, we need to compute the total weight of the solution \mathcal{R} . First, we compute the sum
 984 of weights of segments in A . Fix $1 \leq j \leq k$ and consider segments collinear with the j -th
 985 horizontal line. All points $h_{i,j,t}, h_{i,j,t}^L$ and $h_{i,j,t}^R$ for every $1 \leq i \leq k$ and $1 \leq t \leq n^2$ are covered
 986 by A excluding points $h_{i,j, \text{order}^{-1}(x(i), y(j))}$. Every such point leaves a gap of length 2ϵ between
 987 $h_{i,j, \text{order}^{-1}(x(i), y(j))}^L$ and $h_{i,j, \text{order}^{-1}(x(i), y(j))}^R$. Therefore, the total weight of segments in A that
 988 lie on the line in question equals the length of the segment $(h_{1,1,1}^L, h_{i,k,n^2}^R)$ minus $2\epsilon k$, which is

989 $k(n^2 + 1) - 2(1 - \epsilon) - 2k\epsilon$. We need to multiply that by k , as we consider all possible values
 990 of j .

991 Computation for vertical segments is analogous and yields the same result. Every segment
 992 in D has weight δ , therefore the sum of all weights is equal to:

$$2k(k(n^2 + 1) - 2(1 - \epsilon) - 2k\epsilon) + k^2\delta = W_{\text{hv}} + k^2\delta. \quad \square$$

993 Now we present a few additional properties of the constructed instance of the geometric
 994 set cover that help us to prove Lemma 5.6.

995 **Claim 5.1.** *In any solution to the instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$:*

- 996 • *the left and right guards of points in H (points in $\{p^L : p \in H\} \cup \{p^R : p \in H\}$) have*
 997 *to be covered with segments from HOR,*
- 998 • *the lower and upper guards of points in V (points in $\{p^D : p \in V\} \cup \{p^U : p \in V\}$) have*
 999 *to be covered with segments from VER.*

1000 *Proof.* We prove the claim for the points from H as the proof for points from V is analogous.
 1001 Every segment in VER is vertical and has x-coordinate equal to $i(n^2 + 1)$ for some $1 \leq i \leq k$,
 1002 so they all have different x-coordinate than any left or right guard of points in H .

1003 For every point x which is a left or right guard of a point in H , there are kn^2 segments
 1004 from DIAG that intersect with the horizontal line that goes through x . All of these segments
 1005 intersect with this line in points from set H , therefore none of them covers any of the guards.

1006 Therefore none of the segments from VER or DIAG covers any of the guards of the points
 1007 in H . \square

1008 **Claim 5.2.** *For any $1 \leq i, j \leq n$ and any solution to the instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$, all*
 1009 *but at most one point $h_{i,j,t}$ and at most one point $v_{i,j,t}$ for $1 \leq t \leq n^2$ must be covered with*
 1010 *segments from HOR or VER.*

1011 *Proof.* We prove the claim for horizontal segments, as the proof for vertical segments is anal-
 1012 ogous.

1013 We prove this by contradiction. Assume that we have two points $h_{i,j,t_1}, h_{i,j,t_2}, 1 \leq t_1 <$
 1014 $t_2 \leq n^2$, such that they are not covered with segments from HOR.

1015 Point h_{i,j,t_1}^R has to be covered with a segment from HOR by Claim 5.1. Every segment in
 1016 HOR covering h_{i,j,t_1}^R , but not h_{i,j,t_1} must start at h_{i,j,t_1}^R and all such segments cover also h_{i,j,t_2} .
 1017 This contradicts the assumption, which concludes the proof. \square

1018 **Lemma 5.3.** *For every solution \mathcal{R} to the instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$, the sum of weights of*
 1019 *segments chosen from sets HOR and VER is at least W_{hv} .*

1020 *Proof.* Let us fix $1 \leq i \leq k$.

1021 We provide a lower bound for the sum of lengths of vertical segments from $\mathcal{R} \cap \text{VER}$. This
 1022 bound is the same for each i and is the same for horizontal lines, thus we need to multiply
 1023 such a bound by $2k$.

(1) The total length between $v_{i,1,1}^D$ and v_{i,k,n^2}^U is:

$$(k(n^2 + 1) + n^2 + \epsilon) - ((n^2 + 1) + 1 - \epsilon) = k(n^2 + 1) - 2(1 - \epsilon).$$

1024 (2) For every $1 \leq j \leq k$ there exists at most one $1 \leq t \leq n^2$ such that $v_{i,j,t}$ is not covered
 1025 by segments from **VER** (Claim 5.2). Its guards (see Definition 5.3) $v_{i,j,t}^U$ and $v_{i,j,t}^D$ have
 1026 to be covered in **VER** (Claim 5.1). Therefore, at most k spaces of length 2ϵ can be left
 1027 not covered by segments from **VER** between $v_{i,1,1}^D$ and v_{i,k,n^2}^U .

The sum of these lower bounds for vertical and horizontal lines is:

$$2k(k(n^2 + 1) - 2k\epsilon - 2(1 - \epsilon)) = 2k^2(n^2 + 1) - 4k^2\epsilon - 4k(1 - \epsilon) = W_{\text{hv}}. \quad \square$$

1028 **Lemma 5.4.** *Let \mathcal{R} be a solution to a constructed instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ with weight at*
 1029 *most $W_{\text{hv}} + k^2\delta$. Then for every $1 \leq i, j \leq k$ there exists $1 \leq t \leq n^2$ such that:*

- 1030 (1) $v_{i,j,t}, h_{i,j,t}$ are not covered by segments from **VER** or **HOR**;
- 1031 (2) segment $(v_{i,j,t}, h_{i,j,t})$ is in solution \mathcal{R} ;
- 1032 (3) $\text{order}(t) \in f(i, j)$, that is, $\text{order}(t)$ is an allowed tile for (i, j) ;
- 1033 (4) for every $1 \leq s \leq n^2$, $s \neq t$, $v_{i,j,s}$ is covered in **VER**;
- 1034 (5) for every $1 \leq s \leq n^2$, $s \neq t$, $h_{i,j,s}$ is covered in **HOR**.

1035 *Proof.* At most one of the points $\{h_{i,j,t_x} : 1 \leq t_x \leq n^2\}$ and one of the points $\{v_{i,j,t_y} : 1 \leq$
 1036 $t_y \leq n^2\}$ is covered with **DIAG** (Claim 5.2).

1037 Moreover, exactly one such point h_{i,j,t_x} and one such point v_{i,j,t_y} is covered with **DIAG**,
 1038 because if none of them were covered, then the solution would have to have weight at least
 1039 $W_{\text{hv}} + 2\epsilon$ (see the proof of Lemma 5.3), which is more than $W_{\text{hv}} + k^2\delta$.

1040 We observe that points h_{i,j,t_x} and v_{i,j,t_y} have to be covered with the same segment from
 1041 **DIAG**. Indeed we need to use at least k^2 of them to use exactly one **DIAG** segment for every
 1042 pair of $1 \leq i, j \leq k$, if we used 2 segments from **DIAG** for one pair (i, j) , then we would have
 1043 used total weight at least $W_{\text{hv}} + k^2\delta + \delta$ (Lemma 5.3), which is more than $W_{\text{hv}} + k^2\delta$. Since
 1044 points h_{i,j,t_x} and v_{i,j,t_y} are covered by a single segment from **DIAG**, we have $t_x = t_y$.

1045 Therefore $t_x = t_y$ and $\text{order}(t_x)$ is an allowed tile for (i, j) because the corresponding
 1046 segment is in **DIAG**. \square

1047 We refer to the function mapping $1 \leq x \leq k$ to t_x from Lemma 5.4 as **diagonal** : $\{1, \dots, k\} \times$
 1048 $\{1, \dots, k\} \rightarrow \{1, \dots, n^2\}$.

1049 **Lemma 5.5.** *Let \mathcal{R} be any solution of a constructed instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ with weight*
 1050 *at most $W_{\text{hv}} + k^2\delta$. Then:*

- 1051 1. for any $1 \leq i < k, 1 \leq j \leq k$, $\text{match}_h(\text{diagonal}(i, j), \text{diagonal}(i + 1, j))$ is **true**;
- 1052 2. for any $1 \leq i \leq k, 1 \leq j < k$, $\text{match}_v(\text{diagonal}(i, j), \text{diagonal}(i, j + 1))$ is **true**.

1053 *Proof.* We prove (1) by contradiction, the proof of (2) is analogous.

1054 Let us take any $1 \leq i < k, 1 \leq j \leq k$ and name $t_1 = \text{diagonal}(i, j)$ and $t_2 = \text{diagonal}(i +$
 1055 $1, j)$. We also assume that $\text{match}_h(t_1, t_2)$ is **false**, which is equivalent to the fact that segment
 1056 $(h_{i,j,t_1}^R, h_{i+1,j,t_2}^L)$ is not in set **HOR**.

1057 Therefore h_{i,j,t_1} and h_{i+1,j,t_2} are not covered by segments from **HOR** (Lemma 5.4), while
 1058 h_{i,j,t_1}^R and h_{i+1,j,t_2}^L have to be covered by segments from **HOR** (Claim 5.1).

1059 Every segment from **HOR** either:

- 1060 • starts at point h_{x,y,z_1}^R and ends at point h_{x+1,y,z_2}^L for some $1 \leq x < k, 1 \leq y \leq k$ and
 1061 $1 \leq z_1, z_2 \leq n^2$; or
- 1062 • is $\text{horBeg}_{y,z}$ and starts at $h_{1,y,1}^L$ and ends at h_{1,y,n^2}^L for some $1 \leq y \leq k$ and $1 \leq z \leq n^2$;
 1063 or
- 1064 • is $\text{horEnd}_{y,z}$ and starts at $h_{k,y,z}^R$ and ends at h_{k,y,n^2}^R for some $1 \leq y \leq k$ and $1 \leq z \leq n^2$.

1065 All of the points between h_{i,j,t_1}^R and h_{i+1,j,t_2}^L are covered by segments in HOR and there is no
 1066 segment $(h_{i,j,t_1}^R, h_{i+1,j,t_2}^L)$ in HOR. Hence, there are at least two different segments covering
 1067 them. If both of these segments are neither $\text{horBeg}_{y,z}$ nor $\text{horEnd}_{y,z}$, then one of them must
 1068 begin at h_{i,j,t_1}^R and end at h_{i+1,j,z_2}^L and there must be other one that begins at h_{i,j,z_1}^R and ends
 1069 at h_{i+1,j,t_2}^L for some $1 \leq z_1, z_2 \leq n^2$.

1070 Thus, the space between h_{i,j,z_1}^R and $h_{i,j+1,z_2}^L$ would be covered twice and is longer than ϵ .
 1071 The case when one of them is $\text{horBeg}_{y,z}$ or $\text{horEnd}_{y,z}$ is analogous. Note that they cannot be
 1072 both $\text{horBeg}_{y,z}$ or $\text{horEnd}_{y,z}$.

1073 By the proof of Lemma 5.3, the lower bound for weight of such a solution is $W_{\text{hv}} + \epsilon$ which
 1074 is more than $W_{\text{hv}} + k^2\delta$.

1075 Therefore h_{i,j,t_1}^R and h_{i+1,j,t_2}^L must be covered by one segment from HOR, namely $(h_{i,j,t_1}^R, h_{i+1,j,t_2}^L)$.
 1076 Hence $(h_{i,j,t_1}^R, h_{i+1,j,t_2}^L)$ is a segment in HOR and $\text{match}_h(t_1, t_2)$ is **true**. \square

1077 **Lemma 5.6.** *If there exists a solution to instance $(\mathcal{C}, \mathcal{P}, w, 3k^2 + 2k)$ with weight at most*
 1078 *$W_{\text{hv}} + k^2\delta$, then there exists a solution to the grid tiling instance $(f_{i,j})$.*

1079 *Proof.* Take **diagonal** function from Lemma 5.4.

1080 To define the x function for every $1 \leq i \leq k$ set $x(i) := x_i$ where $(x_i, a) = \text{order}(v_{i,1})$.
 1081 Similarly, to define the y function, for every $1 \leq i \leq k$ set $y(i) := y_i$ where $(b, y_i) = \text{order}(h_{1,i})$

1082 To prove that this is a correct solution to grid tiling, we need to prove that for every
 1083 $1 \leq i, j \leq k$, $(x(i), y(j))$ is in the allowed tiles set $f(i, j)$.

1084 Let us take any $1 \leq i, j \leq k$. By Lemma 5.5 and simple induction, we know that
 1085 $\text{match}_h(\text{diagonal}(1, j), \text{diagonal}(i, j))$ and $\text{match}_v(\text{diagonal}(i, 1), \text{diagonal}(i, j))$ are **true**. There-
 1086 fore $\text{order}(\text{diagonal}(i, j)) = (x(i), y(j))$. By Lemma 5.4 we know that $\text{order}(\text{diagonal}(i, j))$ is in
 1087 $f(i, j)$. Therefore $(x(i), y(j))$ is in $f(i, j)$. \square

Bibliography

- [Adamaszek et al., 2015] Adamaszek, A., Chalermsook, P., and Wiese, A. (2015). How to tame rectangles: Solving independent set and coloring of rectangles via shrinking. In Garg, N., Jansen, K., Rao, A., and Rolim, J. D. P., editors, *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX/RANDOM 2015, August 24-26, 2015, Princeton, NJ, USA*, volume 40 of *LIPICs*, pages 43–60. Schloss Dagstuhl - Leibniz-Zentrum für Informatik.
- [Chan and Grant, 2014] Chan, T. M. and Grant, E. (2014). Exact algorithms and approximation hardness results for geometric packing and covering problems. *Comput. Geom.*, 47(2):112–124.
- [Cygan et al., 2015] Cygan, M., Fomin, F. V., Kowalik, L., Lokshtanov, D., Marx, D., Pilipczuk, M., Pilipczuk, M., and Saurabh, S. (2015). *Parameterized Algorithms*. Springer.
- [Dinur and Steurer, 2014] Dinur, I. and Steurer, D. (2014). Analytical approach to parallel repetition. In Shmoys, D. B., editor, *Symposium on Theory of Computing, STOC 2014, New York, NY, USA, May 31 - June 03, 2014*, pages 624–633. ACM.
- [Etscheid et al., 2017] Etscheid, M., Kratsch, S., Mnich, M., and Röglin, H. (2017). Polynomial kernels for weighted problems. *J. Comput. Syst. Sci.*, 84:1–10.
- [Gaspers et al., 2012] Gaspers, S., Kim, E. J., Ordyniak, S., Saurabh, S., and Szeider, S. (2012). Don’t be strict in local search! In Hoffmann, J. and Selman, B., editors, *Proceedings of the Twenty-Sixth AAAI Conference on Artificial Intelligence, July 22-26, 2012, Toronto, Ontario, Canada*. AAAI Press.
- [Har-Peled and Lee, 2009] Har-Peled, S. and Lee, M. (2009). Weighted geometric set cover problems revisited. *Journal of Computational Geometry*, 3.
- [Håstad, 2001] Håstad, J. (2001). Some optimal inapproximability results. *J. ACM*, 48(4):798–859.
- [Kim et al., 2021] Kim, E. J., Kratsch, S., Pilipczuk, M., and Wahlström, M. (2021). Directed flow-augmentation. *CoRR*, abs/2111.03450.
- [Marx, 2005] Marx, D. (2005). Efficient approximation schemes for geometric problems? In Brodal, G. S. and Leonardi, S., editors, *Algorithms – ESA 2005*, pages 448–459, Berlin, Heidelberg. Springer Berlin Heidelberg.
- [Marx, 2007] Marx, D. (2007). On the optimality of planar and geometric approximation schemes. In *48th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2007), October 20-23, 2007, Providence, RI, USA, Proceedings*, pages 338–348. IEEE Computer Society.

- 1122 [Marx and Pilipczuk, 2015] Marx, D. and Pilipczuk, M. (2015). Optimal parameterized algo-
1123 rithms for planar facility location problems using voronoi diagrams. *CoRR*, abs/1504.05476.
- 1124 [Marx and Schlotter, 2011] Marx, D. and Schlotter, I. (2011). Stable assignment with couples:
1125 Parameterized complexity and local search. *Discret. Optim.*, 8(1):25–40.
- 1126 [Mustafa et al., 2014] Mustafa, N. H., Raman, R., and Ray, S. (2014). Settling the apx-
1127 hardness status for geometric set cover. In *55th IEEE Annual Symposium on Foundations
1128 of Computer Science, FOCS 2014, Philadelphia, PA, USA, October 18-21, 2014*, pages
1129 541–550. IEEE Computer Society.
- 1130 [Mustafa and Ray, 2010] Mustafa, N. H. and Ray, S. (2010). Improved results on geometric
1131 hitting set problems. *Discret. Comput. Geom.*, 44(4):883–895.
- 1132 [Pilipczuk et al., 2016] Pilipczuk, M., van Leeuwen, E. J., and Wiese, A. (2016). Approxima-
1133 tion and parameterized algorithms for geometric independent set with shrinking. *CoRR*,
1134 abs/1611.06501.
- 1135 [Pilipczuk et al., 2020] Pilipczuk, M., van Leeuwen, E. J., and Wiese, A. (2020). Quasi-
1136 polynomial time approximation schemes for packing and covering problems in planar
1137 graphs. *Algorithmica*, 82(6):1703–1739.
- 1138 [Shachnai and Zehavi, 2017] Shachnai, H. and Zehavi, M. (2017). A multivariate framework
1139 for weighted FPT algorithms. *J. Comput. Syst. Sci.*, 89:157–189.
- 1140 [Wiese, 2018] Wiese, A. (2018). Independent set of convex polygons: From n^ϵ to $1 + \epsilon$ via
1141 shrinking. *Algorithmica*, 80(3):918–934.