

Physiological biomarkers of stream fish from the Atlantic Forest within a Conservation Unit and its surroundings

Subtitle 1 Subtitle 2

Adamastor Coutinho Pinto Alice da Silva Barros
Mayara Mirelly da Silva Monteiro Ellen Gomes da Silva
Enelise Marcelle Amado Elvio Sergio Figueredo Medeiros

2026-03-08

Abstract

Table of contents

1	Introduction	2
2	Material and Methods	5
2.1	Collection and animal maintenance.	5
2.2	Plasmatic protein dosage.	5
2.3	Moisture content.	6
2.4	MXR phenotype activity.	6
3	Results	7
3.1	Plasmatic protein dosage.	7
3.2	Moisture content.	7
3.3	MXR phenotype activity.	8
3.4	Abiotic data.	8
3.5	Environmental variables	9
3.6	Community structure	9
4	Discussion	9
	Conclusions	9

Acknowledgements	9
Authorship contribution statement	9
Ethical approval	9
References	9
Figures and Tables	11
Figures	11
Tables	11
Apendices	12
Non-used figures and tables	12

List of Figures

1 Etapas do desenvolvimento da pesquisa científica.	11
---	----

List of Tables

1 Introduction

The devastation of rivers and streams from the Atlantic forest, biogeographical province ((Udvardy, 1975)) included in the list of the world's Hotspots (high biodiversity rates, endemism and anthropic pressure - (Myers et al., 2000)), is caused mainly by pollution, deforestation, damming and introduction of exotic species. Pollution decreases abundance and richness of local fish, being aggravated by loss of riparian vegetation when there is deforestation or change of river course due to damming. These population reductions are also caused in the presence of invasive species, introduced accidentally or intentionally by humans, modifying the original competition/prey relation (Miranda, 2012). With about 2-5% of the original vegetation preserved in Atlantic forest, one of the main strategies for conserving and preserving biodiversity from this and other biogeographical provinces is the delimitation of areas known as Conservative Units (CU), regulated in Brazil by the National System of Conservative Unit (SNUC) and implemented by Chico Mendes Institute (ICMBio) to be managed at federal, state and local levels (Arruda et al., 2013; Miranda, 2012). Ecological data, such as richness and species abundance, are the most studied and represent how aquatic animals deal with general pollution, but the toxicity responsible for behavioral change depends on the organism, time of exposure and particle type, which demonstrates the need for physiological studies,

specially in natural environments (Amoatey & Baawain, 2019). As physiological, morphological and behavioural aspects represent how their functions are performed in the ecosystem, they are nominated functional characteristics, commonly studied by parameters of strength and resilience. Changes on these parameters reflect environmental changes that could not be seen by parameters of taxonomical diversity, which happens when different species possess similar functions, highlighting how these, still scarce studies, can be promising (Júnior, 2021). Taking it into account, as also the relevance of Conservation Units for the Atlantic forest species, this study brings up the possibility to compare the ecological health from these waterbodies by identifying physiological stress markers from the fish within and near CU's. One of the main physiological functions affected by pollutants (such as pesticides, heavy metals, drugs and microplastics) is the osmoregulation one, that, as well as the pH control, is predominantly done by gills. Being also critical to gas exchange, ionic regulation and nitrogen excretion, its functions are partially performed by hormonal control of blood flow (carried out by catecholamines, which reduce vascular resistance in the gills and increase blood flow in the efferent artery via beta and alpha-adrenoreceptors, in that order) and mainly by the transport of ions, probably made in the chloride cells from the filament epithelium, that possess transport enzymes called sodium/potassium-dependent ATPases (Evans, 1987). The action of toxic compounds in the gills leads to a series of histopathologies in response to stress, such as necrosis and epithelial desquamation, frequently generated by flaws in cellular osmoregulation due to ATPase inhibition and physiological failure of the chloride cells or gills as a whole. The effects of this action on the organism can be measured through a few blood markers, mainly lower levels of Chlorine and Sodium ions, obtained by freshwater fish through exchange with internal ions (Evans, 1987), and higher levels of plasmatic proteins (Sabae & Mohamed, 2015). The hydration level of gills and muscles is another physiological parameter tied to the animal's capacity of dealing with salinity variation, and, for that, suffers mensurable effects from the action of toxic compounds in the environment. This active control parameter was suggested by Ayrapetyan (2012) as a universal biomarker for detecting pollution in the environment, being upheld by David et al. (2018) in a study with osmoconforming and euryhaline oysters under effect of different - but common - levels of salinity, and, more discretely, by Sabae & Mohamed (2015) in a study with freshwater fish. Another relevant biomarker is the abnormal activity of the multixenobiotic resistance (MXR) phenotype, an innate gene of many aquatic species that is expressed in organs such as gills, kidneys, blood-brain barrier, liver and pancreas. It is activated by exposure to pollutants and results in the production of P-glycoproteins (Pgp) in the membrane, which act to remove endogenous or exogenous toxic compounds out of the cell, protecting DNA and inducing its excretion. Although the MXR activity explains how some species are more resistant to the presence of pollutants, some substances can reverse its effectiveness, being called chemosensitizers (or simply MXR-inhibitors). Those compounds can cancel the Pgp activity, alter the mechanism regulation, saturate or even reverse MXR activity, acting, in this case, by saturating the Pgp pump and allowing other xenobiotics to enter the cell and increase its toxicity, even if the chemosensitizers themselves are not toxic. Precisely because they are mostly organic, natural, primarily harmless and very present in the materials emitted anthropogenic in nature, its detection should also be a priority in environmental risk studies (Smital & Kurelec, 1998). The hypothesis of markers linked to physiological stress

of fish providing evidence of the environmental stress levels occurring inside and outside of a Conservation Unity was tested in the Guaribas Environmental Reserve (ReBio Guaribas), one of the 22 CU's inside the Atlantic forest area of Paraíba state, located in the Zona da Mata, wettest and most economically relevant mesoregion from the state (Fig. 1). Grouped, according to SNUC, as one of the integral protection areas (aimed to conserve biodiversity, being minority within the biogeographical province – 33,45% of the CU's), the definition of environmental reserves prohibits human interference inside their limits, except for management actions that aim its preservation and recovery. Despite being one of the most restricted types of Conservation Units, there are 22 populations that influence ReBio Guaribas due to historic factors of occupation and absence of an effective buffer zone around the three areas that make up the reserve. Those populations are mainly rural and cause direct impact, by using wood and plants from the reserve, and indirect, causing siltation of rivers and contaminating groundwater through the use of septic tanks and releasing agricultural chemicals into the soil, promoting possible comparable ecophysiological responses in the ichthyofauna inside and around the CU (Arruda et al., 2013; Ministério do Meio Ambiente, 2025). The ecophysiological evaluation of human impact on the forest, which is close to areas of monoculture and natural vegetation modified for livestock, was mainly guided by the fish species inventory of ReBio Guaribas made by Gouveia et al. (2017), in a study that lists 18 species from five different orders: Characiformes (12 species), Perciformes (3 species), Synbranchiformes (1 species), Siluriformes (1 species) e Cyprinodontiformes (1 species). From those, two are exotic - *Oreochromis niloticus*, second most farmed fish species in the world and tolerant to osmotic variation, and *Poecilia reticulata*, largely used in mosquito control and resistant to stressful environments (Miranda, 2012) - and none are listed as threatened. The predominance of small fish inhabiting small basins explains the higher level of endemism in streams, resulting from the allopatric speciation generated by the low dispersion made by these fish and the consequent isolation of populations (Gouveia et al., 2017). Having in mind that these stream fish are phylogenetic close, this paper offers representative data for the entire local ichthyofauna using species of *Hemigrammus* genus, of the Characiformes order, characterized by a black conspicuous spot delimited on the dorsal fin. Its type species, *Hemigrammus unilineatus*, has a compressed and slightly elongated body and exhibits wide geographical distribution both in Central and South America, with registers in other Brazilian states, such as Alagoas and Pernambuco (Serra, 2010). The identification of physiological stress biomarkers in the collected fish aims to evaluate the ecological health of sites inside and surrounding the CU, through quantitative comparison of the influence of anthropogenic action with data from a legally protected area (statistical analysis made in the integrated programming environment R and R Studio - R Core Team, 2017; R Studio Team, 2022). This promotes a more tangible insight in contrast with other scientific papers that only make qualitative assessments of Conservative Units, and also using more specific markers than articles that only explore the ecological consequences of pollution exposure.

2 Material and Methods

2.1 Collection and animal maintenance.

The sampling sites (Fig. 2) were cataloged and numbered by the co-author of this paper (Prof. Dr. Elvio Sérgio Figueredo Medeiros) in previous works, using three data points inside the Reserva Biológica Guaribas (n=92 fish collected) and three in the surrounding area (n=97), numbered, in order, P5 (n=29), P6 (n=31) and P7 (n=32), and P8 (n=28), P9 (n=35) and P10 (n=34), between October and November 2023. In order to uniform the data, only fish from *Hemigrammus* Gill 1858 (Fricke et al., 2026) genera were analyzed, being found in every sampling site and in abundance, captured in the shallow waters of streams using a hand net in sweeping motion. To facilitate handling, the specimens were transported in plastic bags with air and local water to the Animal Ecophysiology Laboratory (LEFA) from UEPB - campus V (João Pessoa, Brazil). To analyze the osmoregulatory function of the fish, blood samples were collected from some specimens (38 fish from inside the reserve and 40 from outside, a total of 78). For that, those specimens were anesthetized with eugenol, and a blood aliquot was obtained by puncturing the heart using a pipette with a heparinized tip. The samples were kept on ice, and plasma was subsequently separated from cells by centrifugation. Due to low volume of obtained plasma, because of the small size of fish, the chlorine ions assay could not be performed. From those same specimens, gill and muscle were collected in order to calculate the moisture content of tissues. Plasma and tissues samples were stored in a freezer at -20°C until the analyses were performed. The other part of collected specimens (54 fish from inside the reserve and 57 from outside, a total of 111) was used in the multixenobiotic resistance (MXR) phenotype assay. Abiotic data, such as water temperature, dissolved oxygen, pH and multiple nutrients were simultaneously collected with the fish, allowing additional statistical analysis that may point out possible correlations between physiological biotic data and these environment components (analysis performed in the integrated programming environment R and R Studio - R Core Team, 2017; R Studio Team, 2022).

2.2 Plasmatic protein dosage.

The plasma samples were defrosted and an aliquot of 2 μ L of each was diluted with 8 μ L of distilled water (1:4 proportion) to perform the total protein dosage by the Bradford (1976) method, in which the dye was also diluted in 1:4 proportion. The intensity of each sample coloration was measured with a microplate reader (SpectraMax i3) at 595 nm wave length, and compared to a standard curve (BSA protein). The results were transferred to a spreadsheet, where the average value in milligrams of protein dosage was calculated for each sampling site, and a Kruskal-Wallis test was performed to determine the existence of differences between the averages of the points inside and outside the Reserva Biológica Guaribas, calculated in the R Studio program (R Studio Team, 2022). Next, an analysis of variance (ANOVA) was performed to determine where there were differences between each sampling site.

2.3 Moisture content.

The tissues samples (gill and muscle) were individually stored in 2 mL eppendorf tubes (previously weighted), initially weighted with the tissue moist, and, after 24 hours inside a 60°C oven, weighted with the dry tissue. The results were transferred to a spreadsheet, where it was calculated the percentual content of tissue water content and the average valor for each sampling site. A T-test (gill) or Krustal-Wallis test (muscle) was performed to determine if there were significant differences between the average values inside and outside the Conservative Unity, followed by an ANOVA to determine the difference between all six sampling sites.

2.4 MXR phenotype activity.

The multixenobiotic resistance (MXR) gene activity was analysed through the rhodamine B accumulation assay adapted from Smital & Kurelec (1998). Rhodamine B is a substrate of P-glycoprotein, the molecular basis of the MXR phenotype. Thus, after exposing the fish to this substrate, the analysis of the amount of rhodamine accumulated in the animal's tissues (mainly the gills) reflects the activity of the MXR phenotype: the greater the accumulation, the lower the activity, and the lower the accumulation, the greater the activity. Therefore, specimens of fish from each sampling site (three inside the Conservative Unity and three outside, total of six sampling sites) were transferred to plastic aquariums containing dechlorinated water and rhodamine B at 2,5 µM concentration, staying in this condition for one hour. The aquarium remained under constant aeration and protected from direct light. After exposition, the fish were anesthetized with eugenol and sacrificed for gill sampling. The tissue samples were allocated in Eppendorfs tubes and weighted in an analytical scale. The weight of moist tissue was obtained by subtracting the weight of the previously weighted empty eppendorf tubes. The tissues were then homogenized with 500 µL of distilled water and the supernatant was transferred in triplicate of 100 µL to a 96-well microplate. The fluorescence intensity of the supernatant (corresponding to the intracellular rhodamine B fluorescence in the tissue) was measured in the microplate reader (SpectraMax i3) at 544 nm excitation and 590 nm emission. The fluorescence value was then normalized by the moist tissue weight in milligrammes, and a Kruskal-Wallis test was calculated in R Studio (R Studio Team, 2022) to determine if there were differences between the averages inside and outside the Conservative Unity, followed by an ANOVA to visualize the differences between each sampling site.

Here you can add your statistical analyses and follow the text with the RStudio code. Following the model in https://elviomedeiros.github.io/Bentos2006_Q/.

```
1 plot(1:10)
```

3 Results

3.1 Plasmatic protein dosage.

The average plasmatic protein concentration, parameter that is positively correlated to environmental stress, was of 32,2 mg in the sampling sites within the Reserva Biológica Guaribas ($n = 18$ plasma samples) and 80,1 mg in the outside sites ($n=26$), revealing an almost 150% higher concentration in the sites with greater human influence. The total number of plasmatic protein dosages ($n=44$) was smaller than the total number of collected fish for this part of the analysis ($n=78$) because of the difficulty on collecting blood from smaller specimens, in addition to the need to collect a minimum amount of plasma required by the Bradford method. After checking the non-normality of variance of the total protein data, a non parametric Kruskal-Wallis test confirmed the existence of statistically significant difference ($p\text{-value}<0,005$) between the data collected from fish inside and outside the Conservative Unity (Fig. 3). The subsequent ANOVA test confirmed the uniformity of values from the sites inside the reserve (all grouped as “a”), differently from the outside sites (Fig. 4). The increase of total plasmatic protein concentration in the animals exposed to pollution is consistent with the Sabae and Mohamed (2015) study, which attributes this increase to five possibilities: activation of metabolic systems as answer to pollutants exposure, degradation of cellular material in the liver, severe pathological conditions in the liver and kidneys, loss of water in the plasma, and induction of proteic synthesis in the liver.

3.2 Moisture content.

The average values of moisture content, parameter related to the osmoregulatory function, were 84,3% for the gill and 80,3% for the muscle of fish inside the ReBio Guaribas ($n=38$ fish). Regarding the fish from sampling sites surrounding the reserve ($n=40$), the averages were 88,9% for the gills and 82,3% for the muscle, increasing, in order, only about 5,6% and 2,5% the values from inside, results that are not very expressive, like the ones described by Haredi (2020). However, through T test (gills, which data had normal variation) and Kruskal-Wallis test (muscle, with non-normal variation data), it was detected statistical difference for both tissues between fish from the sampling sites inside and outside the Guaribas (Fig. 5). The ANOVA test demonstrated similarity between data from sites inside ReBio Guaribas (P5, P6 and P7) and P8, and a lesser similarity with P9, that was also close to P10 (Fig. 6). The overall statistical analysis performed for moisture content reinforces this as a more discreet and inefficient parameter to make direct quantitative comparison for animals exposed to low salinity variation from the environment, but still accurate in determining the uniformity degree of more representative groups (i.e., fish from inside and outside the Conservative Unity).

3.3 MXR phenotype activity.

The average values of normalized fluorescence measured in fish gills were 25.449 rfu/mg (relative fluorescence units/milligramme of tissue) for inside the Conservative Unity (n=54 fish) and 67.138,3 rfu/mg for outside (n=57), value nearly 2,64 times bigger. This difference between averages is opposite to the expected (higher fluorescence signals a lower MXR phenotype activity, meaning a lower exposition to pollutants), that can be explained by the presence of MXR phenotype inhibitors, which were highly correlated to more polluted waters by Kurelec et al. (2000). Among the possible pollutants presents, pesticides, fragrances, microbial degradation products and natural inhibitors from invasive species were described as compounds with high affinity to the Pgp or inhibiting the PKC (protein kinase C) modulator, resulting in high accumulation of rhodamine B (Smital, 2004) Despite the unexpected result, the variance difference between data from the MXR phenotype activity inside and outside the ReBio Guaribas (confirmed by the Kruskal-Wallis test and illustrated in Fig. 7) was also the most conclusive when analysed by ANOVA test, being the only physiological parameter that shows total uniformity for both relationship with the reserve (Fig. 8).

3.4 Abiotic data.

The abiotic data from every sampling site water was measured and reunited in Tab. 1. The abnormal proportion of ammonia indicates the presence of effluents in the water outside the Reserva Biológica Guaribas, and, as discussed by Piedras et al. (2006), this elevated concentration generates metabolic ammonia retention, causing toxicity. The same paper also points out that the toxicity of non-ionized ammonia in the aquatic environment is greater for smaller organisms, but depends on the interaction with other environmental variables, such as temperature, pH and salinity, demanding specific studies with the target species to determine their tolerance to ammonia. The Pearson correlation analysis among biotic data showed only one significant correlation, between Total Protein and MXR phenotype activity (0,6 - moderate and positive), and among abiotic data only two significance, between Nitrate e Phosphor (0,69 - moderate and positive) and between Dissolved Oxygen and Ammonia (-0,82 - strong and negative). It was also detected significant correlation among both data, between Gill Moisture Content and Ammonia (0,55 - moderate and positive), between Gill Moisture Content and Dissolved Oxygen (-0,62 - moderate and negative), and between MXR activity and Nitrate (-0,62 - moderate and negative).

The results section should follow the model in https://elviomedeiros.github.io/Bentos2006_Q/.

3.5 Environmental variables

3.6 Community structure

4 Discussion

Conclusions

Acknowledgements

Authorship contribution statement

Authorship of this paper is based on CRediT (2026).

Author1: Conceptualization, Data curation, Formal analysis, Investigation, Software, Visualization, Writing – review & editing, Writing – original draft. Author2: Data curation, Formal analysis, Software, Writing – review & editing. Author3: Data curation, Formal analysis, Software, Writing – review & editing. Author4: Data curation, Formal analysis, Software, Writing – review & editing. Author5: Conceptualization, Data curation, Methodology, Project administration, Resources, Supervision, Validation, Writing – review & editing. Author6: Conceptualization, Data curation, Formal analysis, Funding acquisition, Methodology, Project administration, Resources, Software, Supervision, Validation, Writing – review & editing.

Ethical approval

This study did not require ethical approval as it did not involve human participants or sensitive data. Field surveys were conducted based on collection permit number (CEUA-048/2023 and SISBIO-89415-2).

References

- Amoatey, P., & Baawain, M. S. (2019). Effects of pollution on freshwater aquatic organisms. *Water Environment Research*, 91(10), 1272–1287. <https://doi.org/10.1002/wer.1221>
- Arruda, D. B., Cunha, B. P., & Rêgo, K. M. C. (2013). CONFLITOS ENTRE ReBio GUARIBAS E COMUNIDADES LOCAIS: (in)justiça ambiental e ecologia política. *Revista Direitos Emergentes na Sociedade Global*, 2(2).

- Ayrapetyan, S. (2012). Cell hydration as a universal marker for detection of environmental pollution. *The Environmentalist*, 32(2), 210–221. <https://doi.org/10.1007/s10669-011-9380-3>
- Bradford, M. M. (1976). *A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding*.
- CReditT. (2026). CRediT - contributor role taxonomy. *NISO/CRediT Standing Committee*. <https://credit.niso.org/>
- David, D. D., Lima, O. G., Nóbrega, A. M. C. D. S., & Amado, E. M. (2018). Capacity of tissue water regulation is impaired in an osmoconformer living in impacted estuaries? *Ecotoxicology and Environmental Safety*, 166, 375–382. <https://doi.org/10.1016/j.ecoenv.2018.09.111>
- Evans, D. H. (1987). *The Fish Gill: Site of Action and Model for Toxic Effects of Environmental Pollutants*.
- Gouveia, R. S. D., Lira, G. L. D. A., Anselmo Ramos, T. P., & Medeiros, E. S. F. (2017). Ichthyofauna of the Reserva Biológica Guaribas and surrounding areas, state of Paraíba, Brazil. *Check List*, 13(5), 581–590. <https://doi.org/10.15560/13.5.581>
- Júnior, M. M. C. (2021). *Funcionamento de Rios Intermitentes sob Influência da Transposição do Rio São Francisco*.
- Miranda, J. (2012). Ameaças aos peixes de riachos da mata atlântica. *Ameaças Aos Peixes de Riachos Da Mata Atlântica*.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A. B., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403(6772), 853–858. <https://doi.org/10.1038/35002501>
- Sabae, S. Z., & Mohamed, F. A. S. (2015). *Effect of Environmental Pollution on the Health of Tilapia spp. from Lake Qarun*.
- Serra, P. P. (2010). *Análise filogenética das espécies de Hemigrammus Gill, 1858 (Characiformes, Characidae)* [PhD thesis]. <https://doi.org/10.5016/DT000610176>
- Smital, T., & Kurelec, B. (1998). The chemosensitizers of multixenobiotic resistance mechanism in aquatic invertebrates: a new class of pollutants. *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis*, 399(1), 43–53. [https://doi.org/10.1016/S0027-5107\(97\)00265-0](https://doi.org/10.1016/S0027-5107(97)00265-0)
- Udvardy, M. (1975). *A classification of the biogeographical provinces of the world* [Unpublished manuscript].

Figures and Tables

Figures



Figure 1: Etapas do desenvolvimento da pesquisa científica.

Tables

Apendices

Non-used figures and tables

 Non-used codes