

Contrasting conservation conditions shape the spatial partitioning of zooplankton diversity in Atlantic Forest streams

Contrasting conservation conditions shape the spatial partitioning of zooplankton diversity in Atlantic Forest streams ON Acta Limnologica Brasiliensia

Ellen Gomes da Silva Mayara Mirelly da Silva Monteiro
Marcela Vitória Bernardo da Silva
Ludmilla Cavalcanti Antunes Lucena Thais Xavier de Melo
Elvio Sergio Figueredo Medeiros

Invalid Date

Abstract

Table of contents

1	Introduction	2
1.1	Objectives	3
2	Material and methods	3
2.1	Study area and sampling design	3
2.2	Environmental data collection	5
2.3	Zooplankton sampling	5
2.4	Statistical analyses	6
3	Results	15
3.1	Environmental variables and nutrient concentration	15
3.2	Community structure	34
3.3	Diversity Partitioning	49
4	Discussion	60
	Conclusions	60

Acknowledgements	60
Authorship contribution statement	60
Ethical approval	61
Declaration of generative AI and AI-assisted technologies in the manuscript preparation process	61
References	61
Figures and Tables	66
Figures	67
Tables	68
Apendices	77
Non-used figures and tables	77

List of Figures

1	67
2	68
3	69
4	70
5	71
6	72
7	73

List of Tables

1 Environment data GT table.	74
2 Nutrient concentrations at the sampling sites.	74
3 Density (ind/m ²) taxa GT table.	75
4 Partition data GT table.	76

1 Introduction

Essas abordagens são baseadas nos conceitos de turnover and nestedness. Species turnover and nestedness are two distinct components used to break down and understand beta diversity—

which measures how much species composition changes between different communities or sites (Baselga, 2010), where, turnover is the replacement of some species by different species from one site to another, while nestedness is the loss or gain of species where poorer sites contain a strict subset of the species found at richer sites.

CITAR: (Ávila et al., 2011; Baselga, 2010, 2012; Crist et al., 2003; Gering et al., 2003; Ligeiro et al., 2010; Shell & Collier, 2018; Veech et al., 2003)

1.1 Objectives

This study describes the richness and density of zooplankton, comparatively between two small head water streams subject to different levels of environmental pressures. One originates within a protected area - Biological Reserve, or REBIO, according to Brasil (2000) - and its course extends beyond its administrative boundaries (MMA, 2003), and a second one that flows entirely outside of the protected area. Within this framework, we use hierarchical additive partitioning to evaluate the relative contribution of different spatial and temporal scales to the diversity of zooplankton assemblages. Given the importance of the Atlantic Forest to the biodiversity (Pereira et al., 2026) and its fragmented stated north of the São Francisco river basin (J. M. C. Silva & Casteleti, 2003), we aim with this study, (1) to infer environmental degradation from species turnover (Baselga, 2010) and (2) to associate beta-diversity to the ecological health of sites inside and surrounding a protected area of Atlantic Forest. Given the expected different levels of environmental pressures in the area (Gouveia et al., 2017), we hypothesize that the diversity is not be homogeneously distributed among the different spatial scales but concentrated at the finer levels and seasonal variation will be limited to species composition, but no spacial variation is expected between dry and wet seasons. Some environmental features (reach morphology, habitat structure, and water quality) are also described to explore possible explaining factors for the observed patterns.

2 Material and methods

2.1 Study area and sampling design

This study was conducted in the Guaribas Biological Reserve (REBIO Guaribas), a protected area of Atlantic Forest in northeastern Brazil. This protected area is located in a transitional zone between the Caatinga and Atlantic Forest ecoregions. The land use of the study area is based on the occupation of traditional human populations and an increasing use of land for large scale monocultures, mostly sugar cane (Heinrichs et al., 2017). Other than these, small scale agricultural activities, fishing, and the modification of the natural vegetation for livestock are important anthropogenic pressures that affect the buffer zone of the REBIO (Arruda et al., 2013; MMA, 2026). Other common uses of the soil in the area are, coconut plantations, orchards and subsistence agriculture. These land uses represent important sources

of pollution, either through agrochemicals runoff or direct and indirect domestic effluents (such as septic tanks). These nonpoint source pollutants contaminate underground water and create a spatial gradient of environmental disturbance (Mehla et al., 2026). Therefore, the mainly rural nature of the land use associated with the management of vegetation, will cause direct impact to aquatic systems, by the siltation of streams and contamination of groundwater. The hydrographic network of the study region is composed predominantly of small catchments containing first- and second-order streams (Strahler, 1957 classification) (Figure 1). These watercourses are tributaries of the Camaratuba River and are sustained by numerous springs distributed throughout the region. The two main stream systems in the study area are the Barro Branco and Caiana catchments. The Barro Branco stream is approximately 8.8 km long and originates within the protected area. Its upper section remains within the boundaries of the REBIO, whereas its middle and lower sections extend into the surrounding landscape. Despite this transition, riparian vegetation is preserved along most of the stream. The Caiana stream is approximately 9.8 km long and also originates within the REBIO, although most of its course lies outside the protection area.

The climate in the region is classified as As' (Peel et al., 2007), characterized by a mean annual temperature of approximately 25 °C, with maximum temperatures occurring mainly between February and March, during the dry period, as well as by low annual thermal amplitude. Annual precipitation ranges from 1,400 to 1,800 mm, concentrated between March and July, whereas the lowest rainfall levels occur between September and December. In the study area, rainfall is concentrated between March and July, beginning in autumn and extending into winter. This pattern is attributed to the influence of cold air masses from the South Atlantic ocean that affect the region. The hydrographic network is composed of small headwater streams fed by rain (Gouveia et al., 2017).

A total of 15 sampling sites distributed longitudinally across the two Atlantic Forest streams were selected to capture spatial variation in environmental conditions and habitat characteristics. The Barro Branco stream included sampling sites located within a protected area (REBIO Guaribas) and sites located outside its boundaries, whereas all sampling sites in the Caiana stream were located outside the protected area. This sampling design allowed the assessment of zooplankton diversity across nested spatial scales and contrasting conditions of environmental protection and human influence (Figure 1). Sampling included temporal (dry and wet seasons) and spatial scales. The spatial scales of zooplankton diversity were represented by the total gamma () diversity, encompassing both study streams. At each stream, several sites were selected according to their position in the main channel of the catchment basin and, at each site, three habitats were sampled to represent different habitat types (pool, riffle, near underwater structures) (Frissell et al., 1986). We also considered a further spatial scale being the upper reaches (portion of the Barro Branco stream within the conservation unit and the Caiana stream before its confluence with the Pipina stream) and the lower reaches (portion of the Barro Branco stream outside the conservation unit and the Caiana stream after its confluence with the Pipina stream) (Figure 1). Sampling was conducted during the dry (January-February) and wet (June-July) seasons of 2024.

2.2 Environmental data collection

Environmental conditions were characterized using four groups of variables, following procedures adapted by Medeiros et al. (2008) from the protocols proposed by Pusey et al. (2004) and Mugodo et al. (2006): site morphology, water quality, substrate composition, and marginal habitat structure. Morphological characteristics were assessed from three randomly selected transects at each site, where stream width (cm) and water depth (cm) were measured. Bank slope was classified into four categories according to its inclination: 1, 20° ; 2, $> 20\text{--}30^\circ$; 3, $> 30\text{--}50^\circ$; and 4, $> 50^\circ$. Broader-scale landscape characteristics, including elevation and stream length, were obtained from handheld GPS records and satellite imagery. Water quality was characterized using physical and chemical measurements obtained with portable instruments, including water temperature ($^\circ\text{C}$) and dissolved oxygen concentration (mg/L). Turbidity was subsequently measured in the laboratory with a turbidimeter using water samples collected at each sampling site. Current velocity (m/s) was estimated using the float method, as described by Maitland (1990). Substrate composition and the physical structure of marginal habitats were evaluated from 9 to 12 quadrats (1 m^2) distributed along the terrestrial-aquatic interface at each sampling site. Within each quadrat, the proportional cover of different substrate categories was visually estimated, including mud, sand, cobbles, small and large gravel and rocks. The same quadrats were used to characterize littoral and submerged habitat features. Recorded habitat components included macrophytes, marginal grasses, leaf litter, attached and filamentous algae, overhanging and submerged vegetation, small and large woody debris, and root masses (Medeiros et al., 2008). For subsequent analyses and presentation, environmental measurements were summarized as mean values for each sampling site.

Three 250 mL replicate samples of water from each site were taken and refrigerated for concentration analysis (g L^{-1}) of ammonia, nitrite, nitrate, orthophosphate (soluble reactive phosphorus) and total phosphorus. Nutrient concentrations were obtained based on Standard Methods for the Examination of Water (1998) and Mackereth et al. (1978).

2.3 Zooplankton sampling

Zooplankton was sampled at habitats within each sampling site, hereafter referred to as habitat samples. Collections were performed in the main channel of the streams at the surface water. At each sampling habitat, a plankton net (30 cm mouth diameter, 70 cm length, and 60 μm mesh size) was used to filter 50 L of water. The net was rinsed between successive samples to prevent clogging and carryover of organisms. Each sample was concentrated to a final volume of 80 mL for laboratory analyses. Following collection, zooplankton samples were anesthetized with commercial sparkling water and subsequently preserved in 4% formalin (EPA, 2020). Sucrose was added to the preserved sample to prevent female cladocerans from losing eggs and to minimize carapace distortion (Haney & Hall, 1973). In the laboratory, samples had their zooplankton identified and counted using a Sedgewick–Rafter counting chamber. The analyses focused on representatives of Rotifera, Cladocera, and Copepoda. Taxonomic identification was

based on specialized literature and confirmed by specialists. To evaluate whether the sampling effort adequately represented the zooplankton assemblage, species accumulation curves were calculated (Lucena et al., 2015; T. X. D. Melo et al., 2014).

2.4 Statistical analyses

Environmental variables were screened for multicollinearity and to identify redundant descriptors. Variables representing similar environmental information were also combined when appropriate, thereby reducing redundancy in the dataset (Legendre & Legendre, 1998).

```
1  #ZOOOREBIO23_PART - ORGANIZANDO DADOS AMBIENTAIS-----
2  #####....-----
3
4  dev.off() #apaga os graficos, se houver algum
5  rm(list=ls(all=TRUE)) #limpa a memória
6  cat("\014") #limpa o console
7
8  ##CARREGANDO MATRIZES BRUTAS-----
9
10 library(openxlsx)
11 habitat <- read.xlsx("dados/rebio23-habitat.xlsx",
12                     rowNames = T,
13                     colNames = T,
14                     sheet = "ambiente",
15                     startRow = 2)
16 # Identify columns containing only NA values
17 all.na <- names(habitat)[colSums(!is.na(habitat)) == 0]
18 all.na <- c(all.na, "w.Temp_C", "w.DO_mg.L")
19 # Remove those columns by name
20 m_hab_part <- habitat[, !names(habitat) %in% all.na, drop = FALSE]
21 # Removendo linhas por nome
22 rownames(m_hab_part)
23 del_rows <- c("C-C-P11-H1", "C-C-P11-H2", "C-C-P11-H3",
24             "S-C-P11-H1", "S-C-P11-H2", "S-C-P11-H3")
25 del_rows
26 m_hab_part <- m_hab_part[!(row.names(m_hab_part) %in% c(del_rows)),]
27 m_hab_part
28 # Remove columns with at least one NA
29 m_hab_part <- m_hab_part[, colSums(is.na(m_hab_part)) == 0, drop = FALSE]
30
31 ###REMOVENDO COLUNAS ZERADAS DE HABITAT-----
32
```

```

33 sum <- colSums(m_hab_part)
34 sum
35 zero_sum <- names(which(colSums(m_hab_part) == 0))
36 zero_sum #nomes das colunas zeradas
37 m_part_cols <- m_hab_part[(colSums(m_hab_part) != 0)] #em != a exclamação inverte o sentido
38 zero_sum2 <- names(which(colSums(m_part_cols) == 0))
39 zero_sum2 #nomes das colunas zeradas
40 sum<-colSums(m_part_cols)
41 sum
42
43 t_grps <- read.xlsx("dados/rebio23-habitat.xlsx",
44                   rowNames = T,
45                   colNames = T,
46                   sheet = "grupos",
47                   startRow = 2)
48
49 ###REMOVENDO LINHAS ZERADAS DE GRUPOS----
50
51 rownames(t_grps)
52 del_rows
53 t_grps <- t_grps[!(row.names(t_grps) %in% c(del_rows)),]
54 t_grps
55
56 ###SEPARANDO NUTRIENTES----
57
58 nutri <- grep("^n\\.\\.", colnames(habitat), value = TRUE)
59 nutri
60 nutrientes <- habitat[, nutri, drop = FALSE]
61 nutrientes <- nutrientes[, colSums(!is.na(nutrientes)) > 0, drop = FALSE]
62 nutrientes <- na.omit(nutrientes)
63
64 ###SALVANDO MATRIZES FINAIS----
65
66 write.table(m_part_cols, "res/m_hab_part.csv",
67             sep = ";", dec = ".", #"\\t",
68             row.names = TRUE,
69             quote = TRUE,
70             append = FALSE)
71 write.table(nutrientes, "res/t_nutri.csv",
72             sep = ";", dec = ".", #"\\t",
73             row.names = TRUE,
74             quote = TRUE,
75             append = FALSE)

```

```

76 write.table(t_grps, "res/t_grps.csv",
77             sep = ";", dec = ".", #"\t",
78             row.names = TRUE,
79             quote = TRUE,
80             append = FALSE)
81 t_grps <- read.csv("res/t_grps.csv",
82                  sep = ";", dec = ".",
83                  row.names = 1,
84                  header = TRUE,
85                  na.strings = NA)
86 t_nutri <- read.csv("res/t_nutri.csv",
87                   sep = ";", dec = ".",
88                   row.names = 1,
89                   header = TRUE,
90                   na.strings = NA)
91 m_hab <- read.csv("res/m_hab_part.csv",
92                  sep = ";", dec = ".",
93                  row.names = 1,
94                  header = TRUE,
95                  na.strings = NA)
96
97 #ZOOEBIO23_PART - ORGANIZANDO DADOS DA COMUNIDADE----
98 #####....----
99
100 dev.off() #apaga os graficos, se houver algum
101 rm(list=ls(all=TRUE)) #limpa a memória
102 cat("\014") #limpa o console
103
104 library(openxlsx)
105
106 ##CARREGANDO MATRIZES BRUTAS----
107
108 densidade <- read.xlsx("dados/rebio23-zoopl.xlsx",
109                      rowNames = T,
110                      colNames = T,
111                      sheet = "densidade_hab_indL")
112 densidade[1:5,1:5] #[1:5,1:5] mostra apenas as linhas e colunas de 1 a 5.
113
114 ##COLUNAS ZERADAS DE DENSIDADE----
115 sum <- colSums(densidade)
116 sum
117 zero_sum <- names(which(colSums(densidade) == 0))
118 zero_sum #nomes das espécies zeradas

```

```

119 m_part_cols <- densidade[(colSums(densidade) != 0)] #em != a exclamação inverte o sentido
120 zero_sum2 <- names(which(colSums(m_part_cols) == 0))
121 zero_sum2 #nomes das espécies zeradas
122 sum<-colSums(m_part_cols)
123 sum
124
125 ###LINHAS ZERADAS DE DENSIDADE-----
126 m_part2 <- as.data.frame(t(m_part_cols))
127 sum <- colSums(m_part2)
128 sum
129 zero_sum <- names(which(colSums(m_part2) == 0))
130 zero_sum #nomes das espécies zeradas
131 m_part_rows <- m_part2[(colSums(m_part2) != 0)] #em != a exclamação inverte o sentido
132 zero_sum2 <- names(which(colSums(m_part_rows) == 0))
133 zero_sum2 #nomes das espécies zeradas
134 sum<-colSums(m_part_rows)
135 sum
136 m_dens <- as.data.frame((t(m_part_rows)))
137
138 ##SALVANDO MATRIZES FINAIS REMOVIDAS AS LINHAS/COLUNAS ZERADAS----
139
140 write.table(m_dens, "res/m_dens.csv",
141             sep = ";", dec = ".", #"\\t",
142             row.names = TRUE,
143             quote = TRUE,
144             append = FALSE)
145 m_dens <- read.csv(res/"m_dens.csv",
146                   sep = ";", dec = ".",
147                   row.names = 1,
148                   header = TRUE,
149                   na.strings = NA)
150
151 #ZOOEBIO23_PART - MATRIZ DE CONTAGEM----
152 #####....-----
153
154 library(openxlsx)
155 contagem <- read.xlsx("dados/rebio23-zoopl.xlsx",
156                      rowNames = T,
157                      colNames = T,
158                      sheet = "rebio23_80ml",
159                      startRow = 2)
160 contagem[1:5,1:5] #[1:5,1:5] mostra apenas as linhas e colunas de 1 a 5.
161

```

```

162 ###COLUNAS ZERADAS DE CONTAGEM----
163 sum <- colSums(contagem)
164 sum
165 zero_sum <- names(which(colSums(contagem) == 0))
166 zero_sum #nomes das espécies zeradas
167 m_part_cols <- contagem[(colSums(contagem) != 0)] #em != a exclamação inverte o sentido
168 zero_sum2 <- names(which(colSums(m_part_cols) == 0))
169 zero_sum2 #nomes das espécies zeradas
170 sum<-colSums(m_part_cols)
171 sum
172
173 ###LINHAS ZERADAS DE CONTAGEM----
174 m_part2 <- as.data.frame(t(m_part_cols))
175 sum <- colSums(m_part2)
176 sum
177 zero_sum <- names(which(colSums(m_part2) == 0))
178 zero_sum #nomes das espécies zeradas
179 m_part_rows <- m_part2[(colSums(m_part2) != 0)] #em != a exclamação inverte o sentido
180 zero_sum2 <- names(which(colSums(m_part_rows) == 0))
181 zero_sum2 #nomes das espécies zeradas
182 sum<-colSums(m_part_rows)
183 sum
184 m_cont <- (t(m_part_rows))
185 m_cont
186 sum(m_cont)
187 ncol(m_cont)
188
189 ##SALVANDO MATRIZ FINAL DE CONTAGEM----
190
191 write.table(m_cont, "res/m_cont.csv",
192             sep = ";", dec = ".", #"\t",
193             row.names = TRUE,
194             quote = TRUE,
195             append = FALSE)
196 m_cont <- read.csv("res/m_cont.csv",
197                   sep = ";", dec = ".",
198                   row.names = 1,
199                   header = TRUE,
200                   na.strings = NA)
201 m_cont
202
203 #ZOOBIO23_PART - ORGANIZANDO ESCALAS DA COMUNIDADE----
204 #####. ....----

```

```

205
206 dev.off() #apaga os graficos, se houver algum
207 rm(list=ls(all=TRUE)) #limpa a memória
208 cat("\014") #limpa o console
209
210 library(openxlsx)
211
212 ##CARREGANDO MATRIZES BRUTAS-----
213
214 escalas <- read.xlsx("dados/rebio23-zoopl.xlsx",
215                     rowNames = T,
216                     colNames = T,
217                     sheet = "escalas_final",
218                     startRow = 2)
219
220 rownames(escalas)
221 del_rows <- c("C-C-P11-H1", "C-C-P11-H2", "C-C-P11-H3", "S-C-P11-H1", "S-C-P11-H2", "S-C-P11-H3")
222 del_rows
223 escalas <- escalas[!(row.names(escalas) %in% c(del_rows)),]
224
225 write.table(escalas, "res/m_esc.csv",
226             sep = ";", dec = ".", #"\\t",
227             row.names = TRUE,
228             quote = TRUE,
229             append = FALSE)
230 m_esc <- read.csv("res/m_esc.csv",
231                  sep = ";", dec = ".",
232                  row.names = 1,
233                  header = TRUE,
234                  na.strings = NA)
235 m_esc

```



```

1  #REBIO23-part-----
2
3  ##ORGANIZANDO DADOS-----
4
5  dev.off()
6  rm(list=ls(all=TRUE))
7  cat("\014")
8
9  t_grps <- read.csv("res/t_grps.csv",
10                   sep = ";", dec = ".",
11                   row.names = 1,

```

```

12         header = TRUE,
13         na.strings = NA)
14 m_hab <- read.csv("res/m_hab_part.csv",
15                 sep = ";", dec = ".",
16                 row.names = 1,
17                 header = TRUE,
18                 na.strings = NA)
19
20 ##CORRELOGRAMA E REMOÇÃO DE VARIÁVEIS REDUNDANTES OU DESNECESSÁRIAS----
21
22 library(psych)
23 colnames(m_hab)
24 png("res/fig-h.hab_pairs.png")
25 pairs.panels(m_hab[,1:8],
26             method = "pearson", # correlation method
27             scale = FALSE, lm = FALSE,
28             hist.col = "#00AFBB", pch = 19,
29             density = TRUE, # show density plots
30             ellipses = TRUE, # show correlation ellipses
31             alpha = 0.5)
32 dev.off()
33
34 cor <- cor(m_hab)
35 cor
36
37 library(corrplot)
38 png("res/fig-hab_corrplot.png")
39 corrplot(cor, method = "circle")
40 dev.off()
41
42 ##DELETANDO COLINEARES OU INDESEJADAS----
43
44 sink(file = "res/colineares.txt", append = F, split = T)
45 colnames(m_hab)
46 del_cols <- c("", "")
47 #del_cols <- grep("^n\\. ", colnames(m_hab), value = TRUE)
48 m_hab_part <- m_hab[, !(colnames(m_hab) %in% del_cols)]
49
50 ##SOMANDO REDUNDANTES----
51
52 m_hab_part$h.Algae <- m_hab_part$h.Algae_f + m_hab_part$h.Algae_a
53 m_hab_part <- m_hab_part[, !(colnames(m_hab_part)
54                             %in% c("h.Algae_f", "h.Algae_a"))]

```

```

55 m_hab_part$h.Debris <- m_hab_part$h.Debris_l + m_hab_part$h.Debris_s
56 m_hab_part <- m_hab_part[, !(colnames(m_hab_part)
57                               %in% c("h.Debris_l", "h.Debris_s"))]
58
59 colnames(m_hab_part)
60 m_hab_part
61 sink()
62
63 write.table(m_hab_part, "res/m_hab_part.csv",
64             sep = ";", dec = ".", #"\\t",
65             row.names = TRUE,
66             quote = TRUE,
67             append = FALSE)
68 m_hab_part <- read.csv("res/m_hab_part.csv",
69                        sep = ";", dec = ".",
70                        row.names = 1,
71                        header = TRUE,
72                        na.strings = NA)
73
74 colnames(m_hab_part)
75 library(PerformanceAnalytics)
76 chart.Correlation(m_hab_part[, c(1:20)], histogram=TRUE)
77 # ***p<0.001, **p<0.01, *p<0.05, .p<0.10
78
79 library(Hmisc)
80 mat <- as.matrix(m_hab_part[, 1:20])
81 res <- rcorr(mat, type = "pearson")
82 cor_mat <- res$r
83 p_mat <- res$P
84
85 stars <- ifelse(p_mat < 0.001, "****",
86                ifelse(p_mat < 0.01, "***",
87                ifelse(p_mat < 0.05, "*", "")))
88
89 cor_star <- ifelse(p_mat < 0.05,
90                   paste0(round(cor_mat, 2), stars),
91                   NA)
92
93 cor_star <- matrix(
94   cor_star,
95   nrow = nrow(cor_mat),
96   dimnames = dimnames(cor_mat)
97 )

```

```

25
26 cor_star[upper.tri(cor_star, diag = TRUE)] <- NA
27 cor_star <- noquote(cor_star)
28 print(cor_star, na.print = "")

```

These data were then analyzed using Principal Component Analysis (PCA) to summarize multivariate patterns of environmental variation among sampling sites. Prior to PCA, site morphology and water-quality variables were square-root transformed (Sokal & Rohlf, 1995). Variables describing sediment composition and marginal habitat structure, expressed as percentages, were relativized by their column totals and subsequently subjected to the arcsine square-root transformation (McCune & Grace, 2002). PCA was conducted in R using the **FactoMineR** package (Lê et al., 2008), with all variables centered and standardized to unit variance before analysis. The water quality parameters temperature and dissolved oxygen, and all nutrients were used to characterize differences between streams but were not used in the PCA because they were not measured for both seasons. A Permutational Multivariate Analysis of Variance (PERMANOVA, 999 permutations) was performed to test whether the overall set of environmental variables measured differed across the seasons and streams (Anderson & Walsh, 2013; Freiry et al., 2020).

Differences in assemblage composition across the spatial hierarchy and seasons were explored using the Unweighted Pair Group Method with Averages (UPGMA) (Belbin & McDonald, 1993; Diniz-Reis et al., 2022), based on Bray-Curtis dissimilarities calculated from the zooplankton density matrix (Ávila et al., 2011). Prior to the analyses, density data were relativized and square-root transformed to reduce the influence of highly abundant taxa and improve the distributional properties of the dataset (Rocha et al., 2012). To focus on spatial variation across sampling sites, habitat density data were pooled for this analysis. To assess how well the UPGMA dendrograms represented the original Bray-Curtis dissimilarities among sampling sites, the Cophenetic Correlation Coefficient (CC) was calculated (Saraçlı et al., 2013; Sneath & Sokal, 1973; Sokal & Rohlf, 1962), where a cophenetic matrix was constructed from each seasonal dendrogram using the *cophenetic* function in the **vegan** package in R. The cophenetic distance matrix was then correlated with the corresponding original Bray-Curtis dissimilarity matrix using Pearson’s product-moment correlation coefficient (Farris, 1969; A. R. Silva & Dias, 2013; Sokal & Rohlf, 1962). To evaluate the strength of the hierarchical clustering structure obtained from the UPGMA dendrograms, the Agglomerative Coefficient (AC) was calculated for each season using the *coef.hclust* function from the **cluster** package in R (Kaufman & Rousseeuw, 1990; Maechler et al., 2026). The AC ranges from 0 to 1, where values closer to 1 indicate a strong, well-defined clustering structure and values near 0 reflect weak or indistinct grouping.

Additive diversity partitioning was used to quantify the contribution of within- and between-unit diversity to total zooplankton diversity. In this framework, alpha diversity (α) represents diversity occurring within sampling units, whereas beta diversity (β) represents the contribution of differences among units at successive levels of the spatial hierarchy (Crist et al., 2003; Lande, 1996). The hierarchical structure included three spatial scales. The finest scales consisted of the

habitat samples within each site ($n = 3$), and of each site ($n = 6$, for the Barro Branco stream, and $n = 9$, for the Caiana stream). The intermediate scales corresponded to the upper and lower stream sections ($n = 2$), whereas the broadest scale represented differences among the two streams. Accordingly, the total diversity (H') was decomposed into within-habitat diversity (α_1) and the successive beta components associated with differences among habitats (β_1), among sites within upper and lower stream reaches (β_2), and among streams (β_3). The contribution of each component was expressed as a percentage of total diversity (Crist et al., 2003; Farias et al., 2020). To investigate seasonal variation in the hierarchical structure of diversity, additive partitioning was performed separately for each season and for each stream (Farias et al., 2020; T. X. Melo & Medeiros, 2013).

The overall diversity was taken to be the sum of the relative contributions of each scale in the hierarchical sampling design. Analysis of significance of the contribution of each hierarchical level was performed through individual-based randomizations (type I null model) (see Freiry et al., 2020; T. X. Melo & Medeiros, 2013). The randomization process was repeated 999 times to obtain null distributions of the H' and H'' diversity at each hierarchical level, and used to test whether the observed values were higher or lower than expected by chance (Crist et al., 2003). The probabilities obtained from randomization tests were interpreted as P -values, as a low P -value ($\text{Prop}_{exp} > \text{Prop}_{obs} < 0.05$) indicates an observed diversity significantly different than that expected under the null model. On the other hand, a high P -value ($\text{Prop}_{exp} > \text{Prop}_{obs} > 0.95$) indicates an observed diversity not different than that expected by chance (Crist et al., 2003; Freiry et al., 2020; Gering et al., 2003; Veech et al., 2002). The results of the analysis were subject to a Monte Carlo permutation test (999 permutations) for validation. The additive partitioning of the diversity was carried out with the function *adipart* from package *vegan* (Oksanen et al., n.d.). All analyses were conducted in the statistical environment R (R Core Team, 2017).

3 Results

3.1 Environmental variables and nutrient concentration

Water velocity ranged from 0.022 to 0.293 m/s, with the highest values recorded at site 5. Width varied from 1.2 to 7.2 m, while depth ranged from 13.7 to 44.2 cm. Waters were, on average, well oxygenated (from 5.4 to 8.1 mg/L), and cool, with average temperatures from 25.2 to 27.8 °C. Conductivity was high, between 129.0 and 163.3 S/cm, and pH, neutral to slightly acidic (from 5.5 to 7.0). Turbidity was generally low, although higher values were observed at sites 5 and 6, reaching 17.5 and 29.6 NTU, respectively. Elevation ranged from 41.5 to 112.8 m a.s.l., while the slope reached 50° for some sites. Mud and sand were the main substrate components across all sampled sites, with average cover ranging from 30 to 100% and from 0 to 70%, respectively. Habitat structure varied along the stream course. The reference site (site 1) was characterized by high contributions of overhanging vegetation (90%) and exclusively muddy substrates. Site 2 exhibited similar characteristics, whereas site 3 showed a more reduced

habitat complexity. Sites 4–6 displayed greater variation in habitat composition. Overall, the habitat structures contributing most were overhanging vegetation, debris, and root masses, although these contributions varied among sites. Aquatic macrophytes and littoral grass were particularly abundant at site 4, contributing on average 35.8% and 39.2%, respectively, whereas root masses represented the dominant habitat structure at site 5 (76.7%). Sand cover increased downstream, reaching 70% at site 6, where overhanging vegetation was markedly reduced (Table 1).

```

1  #TABELA DE HABITAT----
2
3  m_hab_part <- read.csv("res/m_hab_part.csv",
4                        sep = ";", dec = ".",
5                        row.names = 1,
6                        header = TRUE,
7                        na.strings = NA)
8  t_grps <- read.csv("res/t_grps.csv",
9                   sep = ";", dec = ".",
10                  row.names = 1,
11                  header = TRUE,
12                  na.strings = NA)
13
14  library(dplyr)
15  library(tidyr)
16  m_trab <- m_hab_part %>%
17    rename_with(~ gsub("_", ".", .))
18  m <- m_trab %>%
19    group_by(PontoN = t_grps$PontoN) %>%
20    summarise(across(where(is.numeric),
21                     list(mean = mean, min = min, max = max)),
22             .groups = 'drop') %>%
23    pivot_longer(
24      cols = -c(PontoN),
25      names_to = c("Variable", ".value"),
26      names_sep = "_"
27    )
28
29  m <- as.data.frame(m)
30  m_wide <- m %>%
31    mutate(stat_string = ifelse(Variable == c("m.Vel.m.s"),
32                                paste0(round(mean, 3), " (", round(min, 3), "-", round(max, 3),
33                                paste0(round(mean, 1), " (", round(min, 1), "-", round(max, 1),
34    unite("Location", PontoN, sep = "_") %>%
35    select(Variable, Location, stat_string) %>%

```

```

36   pivot_wider(names_from = Location, values_from = stat_string)
37
38   m_wide
39   m_wide <- as.data.frame(m_wide)
40   cols <- colnames(m_wide)
41   col_order <- c(
42     "Variable",
43     cols[cols != "Variable"][
44       order(as.numeric(gsub("Ponto", "", cols[cols != "Variable"])))]
45   ]
46 )
47 m_wide <- m_wide[, c(col_order)] #sequência das colunas
48 row_order <- c(
49   "w.Cond.uS.cm",
50   "w.Turb.ntu",
51   "w.pH",
52   "m.Elev.m",
53   "m.Vel.m.s",
54   "m.Slope",
55   "m.Depth.avg.cm",
56   "m.Depth.mar.cm",
57   "s.Mud",
58   "s.Sand",
59   "s.Gravel.f",
60   "s.Rocks",
61   "h.Macrophytes",
62   "h.Algae",
63   "h.Subm.veg",
64   "h.Overh.veg",
65   "h.Grass",
66   "h.Leaf.l",
67   "h.Root.masses",
68   "h.Debris"
69 )
70 m_wide <- m_wide[match(row_order, m_wide$Variable), ]
71 m_wide
72
73 # Salvado m_wide
74 write.table(m_wide, "res/m_wide_hab.txt",
75             sep = ";", dec = ".", #"\\t",
76             row.names = TRUE,
77             quote = TRUE,
78             append = FALSE)

```

```

79 m_wide_hab <- read.table("res/m_wide_hab.txt",
80                           sep = ";", dec = ".",
81                           row.names = 1,
82                           header = TRUE,
83                           na.strings = NA)
84
85 # Exportando dados para Excel
86 library(openxlsx)
87 #write.xlsx(m_wide, file = "res/tabela de habitat.xlsx", rowNames = FALSE)
88 wb <- loadWorkbook("res/tabela de habitat.xlsx")
89 writeData(wb, sheet = "Sheet 1", x = m_wide)
90 saveWorkbook(wb, "res/tabela de habitat.xlsx", overwrite = TRUE)
91
92 # Tabela final ajustada no Excel
93 library(readxl)
94 hab_gtttable <- read_excel(
95   path = "res/tabela de habitat.xlsx",
96   sheet = "tabela_final",
97   range = "A1:P21") #ajustar para o tamanho da tabela final
98 hab_gtttable
99 library(gt)
100 hab_gtttable <- gt(hab_gtttable)
101 hab_gtttable <- sub_missing(hab_gtttable,
102                             columns = everything(), missing_text = "")
103 #hab_gtttable <- fmt_number(hab_gtttable, columns = "F.O.(%)", decimals = 1)
104 hab_gtttable
105 gtsave(hab_gtttable, "res/gt-hab_gtttable.xlsx.html")
106 saveRDS(hab_gtttable, "res/gt-hab_gtttable.xlsx.rds")

```



```

1 # Escolher sumário de uma variavel
2 m
3 var <- "w.Temp.C"
4 m[m$Variable == var, "mean"] #cada valor de var
5 summary(m[m$Variable == var, "mean"]) #sumário dos valores de var
6
7 # Escolher sumário de um grupo de variáveis do df m
8 vars <- unique(grep("^h\\.\\.", m$Variable, value = TRUE))
9 summaries <- list() #criar uma lista vazia para guardar os sumários
10 # Loop para cada variável do grupo e guarda em summaries
11 for (var in vars) {
12   summaries[[var]] <- summary(m[m$Variable == var, "mean"])
13 }
14 #var is a temporary variable used in the for loop to iterate through

```

```

15  #each variable name that starts with "h."
16
17  summaries
18  summary_table <- do.call(rbind, lapply(summaries, as.data.frame.list))
19  round(summary_table, 2)
20  #sink(file = "summary_h.txt", split = TRUE)
21  round(summary_table[order(summary_table$Mean, decreasing = FALSE), ], 2)
22  #sink()
23
24  # Tabela limpa
25  #summary_table <- cbind(Variable = rownames(summary_table), summary_table)
26  #rownames(summary_table) <- NULL
27  #colnames(summary_table) <- c("Variable", "Min", "Q1", "Median", "Mean", "Q3", "Max")
28  #summary_table

```

Principal Component Analysis (PCA) was used to summarize environmental variation among sampling sites and to identify variables contributing to their differences (Figure 2 and Figure 3). The first two principal components explained 37% of the total variance, with the first axis accounting for 21.1%. PCA confirmed the expected quality of sites, separating site types (Figure 2), where site types I and II were associated with higher elevation, the presence of algae and leaf litter, and where dominated by muddy substrate. Sites type III, tended to have higher water velocity and the presence of gravel and sand as substrate. The presence of root masses and greater inclination on the margins were also important features of these sites. Sites type IV showed a wide range of environmental characteristics, certainly due to their greater number. Nonetheless, these sites may be associated with greater depths, the presence of rocks in the substrate, and higher pH and turbidity. Comparison of the two study streams showed that, the Barro Branco stream tended to have a greater presence of algae, macrophytes, leaf litter and submerged vegetation as major habitat structures, a muddy substrate and an overall higher elevation. The Caiana stream was characterized by sandy substrate, with occasionally gravel and rocks, higher depths, and higher pH, turbidity, and water velocity. Dry and rainy seasons did not show clear segregation in the PCA.

```

1  #PCA fviz package----
2  #browseURL("https://www.sthda.com/english/articles/31-principal-component-methods-in-r-pract
3
4  #ORGANIZANDO DADOS----
5
6  dev.off()
7  rm(list=ls(all=TRUE))
8  cat("\014")
9
10 t_grps <- read.csv("res/t_grps.csv",
11                  sep = ";", dec = ".",

```

```

12         row.names = 1,
13         header = TRUE,
14         na.strings = NA)
15 m_hab_trns <- read.csv("res/m_hab_trns.csv",
16                       sep = ";", dec = ".",
17                       row.names = 1,
18                       header = TRUE,
19                       na.strings = NA)
20
21 #PCA-----
22
23 library("FactoMineR")
24 library("factoextra")
25
26 pca <- PCA(m_hab_trns, scale.unit = TRUE, ncp = 5, graph = TRUE)
27 print(pca)
28
29 eig.val <- get_eigenvalue(pca)
30 eig.val
31
32 #sink(file = "res/cor_matrix.txt", append = FALSE)
33 pca$var$cor
34 #sink()
35
36 fviz_eig(pca, addlabels = TRUE, ylim = c(0,30))
37
38 var <- get_pca_var(pca)
39 var
40
41 fviz_pca_var(pca, col.var = "black")
42
43 library("corrplot")
44 corrplot(var$cos2, is.corr=FALSE)
45
46 fviz_cos2(pca, choice = "var", axes = 1:2)
47
48 fviz_pca_var(pca, col.var = "cos2",
49             gradient.cols = c("#00AFBB", "#E7B800", "#FC4E07"),
50             repel = TRUE # Avoid text overlapping
51 )
52
53 fviz_pca_var(pca, alpha.var = "cos2")
54

```

```

55 corrplot(var$contrib, is.corr=FALSE)
56
57 fviz_contrib(pca, choice = "var", axes = 1, top = 10)
58 fviz_contrib(pca, choice = "var", axes = 2, top = 10)
59 fviz_contrib(pca, choice = "var", axes = 1:2, top = 10)
60
61 fviz_pca_var(pca, col.var = "contrib",
62               gradient.cols = c("#00AFBB", "#E7B800", "#FC4E07")
63 )
64 fviz_pca_var(pca, alpha.var = "contrib")
65
66 ##GROUPING BY KMEANS----
67
68 set.seed(123)
69 res.km <- kmeans(var$coord, centers = 3, nstart = 25)
70 grp <- as.factor(res.km$cluster)
71 # Color variables by groups
72 fviz_pca_var(pca, col.var = grp,
73               palette = c("#0073C2FF", "#EFC000FF", "#868686FF"),
74               legend.title = "Cluster")
75
76 ind <- get_pca_ind(pca)
77 ind
78 ind$contrib
79
80 ##BIPLOTS-----
81
82 fviz_pca_ind(pca)
83
84 fviz_pca_ind(pca, col.ind = "cos2",
85               gradient.cols = c("#00AFBB", "#E7B800", "#FC4E07"),
86               repel = TRUE #avoid text overlapping (slow if many points)
87 )
88
89 fviz_pca_ind(pca, pointsize = "cos2",
90               pointshape = 21, fill = "#E7B800",
91               repel = TRUE #avoid text overlapping (slow if many points)
92 )
93
94 fviz_pca_ind(pca, col.ind = "cos2", pointsize = "cos2",
95               gradient.cols = c("#00AFBB", "#E7B800", "#FC4E07"),
96               repel = TRUE #avoid text overlapping (slow if many points)
97 )

```

```

98
99 fviz_cos2(pca, choice = "ind")
100
101 fviz_contrib(pca, choice = "ind", axes = 1:2)
102
103 # Create a random continuous variable of length 23,
104 # Same length as the number of active individuals in the PCA
105 t_grps <- t_grps[rownames(pca$ind$coord), ]
106 colnames(t_grps)
107 set.seed(123)
108 # Escolhendo variável categorica
109 my.cont.var <- t_grps$Rio
110 my.cont.var
111 # Color individuals by the continuous variable
112 fviz_pca_ind(
113   pca,
114   habillage = my.cont.var,
115   palette = c("blue", "yellow", "red", "pink"),
116   repel = FALSE
117 )
118
119 fviz_pca_ind(pca,
120             geom.ind = "point", #show points only (nbut not "text")
121             col.ind = as.factor(my.cont.var), #color by groups
122             palette = "grey",
123             addEllipses = TRUE, #concentration ellipses
124             legend.title = "Groups"
125 )
126
127 length(unique(my.cont.var))
128 fviz_pca_ind(pca,
129             geom.ind = "point",
130             col.ind = my.cont.var,
131             palette = "grey",
132             addEllipses = TRUE, ellipse.type = "confidence",
133             mean.point = TRUE,
134             legend.title = "Groups")
135
136 fviz_pca_biplot(pca, repel = TRUE,
137                col.var = "#2E9FDF", #variables color
138                col.ind = "#696969" #individuals color
139 )
140

```

```

141 fviz_pca_biplot(pca,
142                 col.ind = my.cont.var, palette = "jco",
143                 addEllipses = TRUE, ellipse.type = "t", #t-distribution confidence ellipses
144                 ellipse.level = 0.95,
145                 shape.ind = my.cont.var,
146                 label = "var",
147                 col.var = "black", repel = TRUE,
148                 legend.title = "Species") +
149 scale_color_manual(values = c(
150   "#1B9E77",
151   "#D95F02",
152   "#7570B3",
153   "#E7298A"
154 )) +
155 scale_shape_manual(values = c(
156   16, # circle
157   17, # triangle
158   15, # square
159   18 # diamond
160 ))
161
162 ### FILTRANDO AS VARIÁVEIS
163 var_coord <- pca$var$coord[,1:2]
164 keep_vars <- apply(abs(var_coord) > 0.5,1,any)
165 vars_to_keep <- rownames(var_coord)[keep_vars]
166 pca_filt <- pca
167 pca_filt$var$coord <- pca$var$coord[vars_to_keep,,drop=FALSE]
168 pca_filt$var$cos2 <- pca$var$cos2[vars_to_keep,,drop=FALSE]
169 pca_filt$var$contrib <- pca$var$contrib[vars_to_keep,,drop=FALSE]
170
171 ### AJUSTE MANUAL DAS COORDENADAS DOS VETORES
172 var_coords <- as.data.frame(pca_filt$var$coord)
173 var_coords$Dim.1 <- var_coords$Dim.1*4.2
174 var_coords$Dim.2 <- var_coords$Dim.2*4.2
175
176 ## Caso queira editar manualmente
177 #fix(var_coords) #nomes das colunas
178 #write.table(var_coords,"coords_var.txt")
179
180 var_coords <- read.table("coords_var.txt",
181                        header=TRUE,
182                        row.names=1)
183

```

```

184 ### AJUSTE MANUAL DOS INDIVÍDUOS
185 ind_coords <- as.data.frame(pca_filt$ind$coord)
186
187 ## Caso queira editar manualmente
188 #fix(ind_coords) #nomes das linhas
189 #write.table(ind_coords,"coords_ind.txt")
190
191 ind_coords <- read.table("coords_ind.txt",
192                         header=TRUE,
193                         row.names=1)
194
195 # GRÁFICO FINAL
196
197 library(factoextra)
198 library(ggplot2)
199
200 t_grps$Ref_site <- as.factor(t_grps$Ref_site)
201
202 #png("fig-hab_PCA_fviz.png", width=11, height=9, units="in", res=400)
203
204 library(ggplot2)
205
206 ggplot() +
207
208   geom_hline(yintercept = 0, linetype = 2) +
209   geom_vline(xintercept = 0, linetype = 2) +
210
211   geom_point(
212     data = ind_coords,
213     aes(Dim.1, Dim.2, shape = t_grps$Ref_site),
214     size = 3,
215     colour = "black"
216   ) +
217
218   geom_segment(
219     data = var_coords,
220     aes(x = 0, y = 0,
221         xend = Dim.1,
222         yend = Dim.2),
223     colour = "red",
224     arrow = arrow(length = unit(0.25,"cm"))
225   ) +
226

```

```

227 geom_text(
228   data = transform(
229     var_coords,
230     nudge_y = ifelse(Dim.2 >= 0, 0.25, -0.25)
231   ),
232   aes(
233     x = Dim.1,
234     y = Dim.2 + nudge_y,
235     label = rownames(var_coords)
236   ),
237   colour = "red",
238   size = 4,
239   nudge_x = 0.19
240 ) +
241
242 scale_shape_manual(values = c(15,16,0,1,2,17)) +
243
244 labs(
245   x = paste0("Dim1 (", round(pca$eig[1,2],1), "%)"),
246   y = paste0("Dim2 (", round(pca$eig[2,2],1), "%)"),
247   shape = "Sites"
248 ) +
249
250 coord_equal() +
251 theme_minimal()
252
253 #dev.off()

```

```

1  #ORGANIZANDO DADOS----
2
3  dev.off()
4  rm(list=ls(all=TRUE))
5  cat("\014")
6
7  t_grps <- read.csv("res/t_grps.csv",
8                    sep = ";", dec = ".",
9                    row.names = 1,
10                   header = TRUE,
11                   na.strings = NA)
12  m_hab_trns <- read.csv("res/m_hab_trns.csv",
13                       sep = ";", dec = ".",
14                       row.names = 1,
15                       header = TRUE,

```

```

16         na.strings = NA)
17
18 colnames(t_grps)
19 m_trab <- m_hab_trns
20 library(vegan)
21
22 #MRPP----
23
24 #TRABALHANDO COM MULTIPLAS VARIAVEIS----
25
26 # Run MRPP for multiple variables
27 fatores <- c("Periodo", "Rio", "Ref_site")
28 set.seed(123)
29 mrpp_results <- lapply(fatores, function(v) {
30     grp <- as.factor(t_grps[[v]])
31     mrpp(
32         dat = m_trab,
33         grouping = grp,
34         distance = "bray"
35     )
36 })
37 names(mrpp_results) <- fatores
38 mrpp_results
39 # Print one result
40 mrpp_results$Ref_site
41 # Loop through summaries
42 sink(file = "res/mrpp.txt", split = TRUE)#, append = TRUE)
43 for (v in names(mrpp_results)) {
44     cat("\n=====\\n")
45     cat("MRPP for:", v, "\\n")
46     print(mrpp_results[[v]])
47 }
48 sink()
49 # Extract key statistics into a table
50 mrpp_summary <- data.frame(
51     Variable = fatores,
52     Delta = sapply(mrpp_results, function(x) x$delta),
53     Expected_Delta = sapply(mrpp_results, function(x) x$E.delta),
54     A = sapply(mrpp_results, function(x) x$A),
55     P_value = sapply(mrpp_results, function(x) x$Pvalue)
56 )
57 sink("res/mrpp.txt", append = TRUE)
58 mrpp_summary

```

```

59 sink()
60
61 ##TRABALHANDO COM UMA UNICA VARIABEL----
62
63 # Run MRPP for a single variable
64 grp <- as.factor(t_grps$Ref_site) #UA, mes, estacao, area, ambiente
65 set.seed(123)
66 mrpp <- mrpp(dat = m_trab, grouping = grp, distance = "bray")
67 mrpp
68 summary(mrpp)
69
70 # Teste de hipótese para uma unica variavel
71 str(mrpp)
72 p_value <- mrpp$Pvalue
73 print(sprintf("p-value: %.10f", p_value))
74 # Conditional statement to check the p-value and print the appropriate message
75 if (p_value < 0.05) {
76   print(sprintf("Existe diferença significativa porque o valor de p é %.10f, sendo menor que
77 } else {
78   print(sprintf("Não existe diferença significativa porque o valor de p é %.10f, sendo maior
79 }
80
81 ##TESTES PAREADOS----
82
83 # Todos os pares de níveis do fator
84 pairs <- combn(levels(as.factor(grp)), 2, simplify = FALSE)
85 pairs
86 # Aplicar mrpp para cada par
87 res_list <- lapply(pairs, function(p) {
88   sel <- grp %in% p
89   mrpp(dat = m_trab[sel, ], grouping = droplevels(grp[sel]), distance = "bray")
90 })
91 # Dar nomes aos resultados
92 names(res_list) <- sapply(pairs, function(p) paste(p, collapse = "_vs_"))
93 # visualizar
94 res_list
95 # Resumo
96 summary_tab <- data.frame(
97   Comparison = names(res_list),
98   A = sapply(res_list, function(x) x$A),
99   Delta = sapply(res_list, function(x) x$delta),
100   Pvalue = sapply(res_list, function(x) x$Pvalue)
101 )

```

```

102 sink("res/mrpp.txt", append = TRUE)
103 summary_tab
104 sink()
105
106 ###MATRIX-STYLE TABLE----
107 library(dplyr)
108 library(tidyr)
109 library(knitr)
110 library(kableExtra)
111 # Extract unique group names from the comparisons
112 groups1 <- sort(unique(unlist(strsplit(summary_tab$Comparison, "_vs_"))))
113 # Initialize empty matrix
114 pval_matrix <- matrix(NA, nrow = length(groups1), ncol = length(groups1),
115                       dimnames = list(groups1, groups1))
116 # Fill matrix with Pvalues
117 for (i in 1:nrow(summary_tab)) {
118   comp <- unlist(strsplit(summary_tab$Comparison[i], "_vs_"))
119   g1 <- comp[1]
120   g2 <- comp[2]
121   p <- summary_tab$Pvalue[i]
122
123   pval_matrix[g1, g2] <- p
124   pval_matrix[g2, g1] <- p #make symmetric
125 }
126
127 # Format p-values: red if <0.05
128 pval_colored <- ifelse(is.na(pval_matrix), "",
129                       ifelse(pval_matrix < 0.05,
130                             paste0("<span style='color:red;'>", sprintf("%.3f", pval_matrix),
131                                     sprintf("%.3f", pval_matrix)))
132
133 # Convert to data.frame for kable
134 pval_df <- as.data.frame(pval_colored)
135 pval_df <- tibble::rownames_to_column(pval_df, "Group")
136 # Render HTML table
137 matrixstyle <- pval_df %>%
138   kable("html", escape = FALSE) %>%
139   kable_styling(full_width = FALSE)
140 matrixstyle
141 save_kable(matrixstyle, "res/kb-mrpp_matrixstyle.html")
142
143 ##GRÁFICO DA MRPP----
144

```

```

145 png("res/fig-mrpp.png", width = 15, height = 10, units = "in", res = 300)
146 def.par <- par(no.readonly = TRUE)
147 layout(matrix(1:2,nr=1))
148 plot(ord <- metaMDS(m_trab, distance="bray"), type="text", display="sites")
149 with(t_grps, ordihull(ord, grp))
150 with(mrpp, {
151   fig.dist <- hist(boot.deltas, xlim=range(c(delta,boot.deltas)),
152                   main="Test of Differences Among Groups")
153   abline(v=delta);
154   text(delta, 2*mean(fig.dist$counts), adj = -0.5,
155         expression(bold(delta)), cex=1.5 ) }
156 )
157 par(def.par)
158 dev.off()
159
160 ###AVALIANDO AS DISTÂNCIAS MÉDIAS----
161 md <- with(t_grps, meandist(vegdist(m_trab), grp))
162 md
163 summary(md)
164 par(mfrow=c(1,2))
165 plot(md)
166 plot(md, kind="histogram")
167 par(mfrow=c(1,1))
168
169 #PerMANOVA----
170
171 #browseURL("https://riffomonas.org/code_club/2021-03-18-vegan")
172 #browseURL("https://riffomonas.org/code_club/2021-03-22-reproducibility")
173
174 library(tidyverse)
175 library(ggtext)
176 library(ggplot2)
177 library(vegan)
178
179 adonis2(m_trab ~ Perodo, data = t_grps, method = "bray")
180 set.seed(123)
181 perm <- 999
182
183 all_test <- adonis2(m_trab ~ Perodo, data = t_grps, method = "bray", permutations = perm)
184 all_test
185
186 adonis2(m_trab ~ Perodo*Rio, data = t_grps)
187 adonis2(m_trab ~ Perodo*Rio, data = t_grps, by = "terms") #ou by=NULL remove as interações

```

```

188
189 all_p <- all_test$`Pr(>F)`[1]
190 all_p
191
192 count(t_grps, Periodo)
193 colnames(t_grps)
194
195 # Pairwise comparisons
196 #browseURL("https://doi.org/10.1007/s10641-018-0751-1")
197 #Lanés, L.E.K., Reichard, M., de Moura, R.G. et al. 2018.
198 #Environmental predictors for annual fish assemblages in subtropical
199 #grasslands of South America: the role of landscape and habitat
200 #characteristics. Environ Biol Fish 101, 963-977.
201
202 library(dplyr)
203 #library(combninat) # or base R combn()
204
205 set.seed(123)
206
207 # Get unique levels of UA
208 groups <- unique(t_grps$Ref_site)
209 groups
210 # All pairwise combinations
211 pairs <- combn(groups, 2, simplify = FALSE)
212 pairs
213 # Store results
214 pairwise_results <- lapply(pairs, function(g) {
215   # Subset data
216   sub_data <- t_grps %>% filter(Ref_site %in% g)
217   sub_dist <- m_trab[rownames(sub_data), ]
218
219   # Run adonis2
220   test <- adonis2(sub_dist ~ Ref_site, data = sub_data,
221     method = "bray", permutations = perm)
222
223   tibble(
224     group1 = g[1],
225     group2 = g[2],
226     F_value = test$F[1],
227     R2 = test$R2[1],
228     p_value = test$`Pr(>F)`[1]
229   )
230 })

```

```

231 # Bind all results
232 pairwise_results <- bind_rows(pairwise_results)
233 pairwise_results
234 pairwise_p <- numeric()
235 pairwise_p <- pairwise_results$p_value
236 pairwise_p
237 adj_p_values <- p.adjust(pairwise_p, method="bonferroni") #c("holm", "hochberg", "hommel", "
238 adj_p_values
239 pairwise_results$adj_p_values <- adj_p_values
240 pairwise_results
241 # Get unique group names
242 groups2 <- sort(unique(c(pairwise_results$group1, pairwise_results$group2)))
243 # Initialize empty matrix
244 pval_matrix <- matrix(NA, nrow = length(groups2), ncol = length(groups2),
245                       dimnames = list(groups2, groups2))
246 # Fill in p-values
247 for (i in 1:nrow(pairwise_results)) {
248   g1 <- pairwise_results$group1[i]
249   g2 <- pairwise_results$group2[i]
250   p <- pairwise_results$p_value[i]
251
252   pval_matrix[g1, g2] <- p
253   pval_matrix[g2, g1] <- p #make symmetric
254 }
255 pval_matrix
256
257 library(knitr)
258 library(kableExtra)
259
260 # Make a copy of the matrix
261 pval_colored <- pval_matrix
262 # Replace numeric values with formatted HTML strings
263 pval_colored[] <- ifelse(is.na(pval_matrix), "",
264                          ifelse(pval_matrix < 0.05,
265                                paste0("<span style='color:red;'>", sprintf("%.3f", pval_matr
266                                sprintf("%.3f", pval_matrix)))
267 # Convert to data frame with rownames
268 pval_df <- as.data.frame(pval_colored)
269 pval_df <- tibble::rownames_to_column(pval_df, "Group")
270 # Render table
271 pval_df %>%
272   kable("html", escape = FALSE) %>%
273   kable_styling(full_width = FALSE)

```

```

274 save_kable(matrixstyle, "res/kb-perma_matrixstyle.html")
275
276 ## AVALIANDO INTERAÇÕES-----
277
278 adonis2(m_trab ~ Período, data = t_grps, method = "bray")
279 adonis2(m_trab ~ Período*Rio, data = t_grps,
280        method = "bray") #+ no lugar de * tira os residuos
281 adonis2(m_trab ~ Período*Rio, data = t_grps,
282        method = "bray", by = "terms", strata = NULL)

```

Among the averaged nutrient concentrations (Table 2), ammonia ranged from 49.2 to 243.7 $\mu\text{g L}^{-1}$. The Caiana stream tended to have greater values, overall no clear trend can be observed when comparing both streams. Nitrite and nitrate, on the other hand, show greater concentrations at the Caiana stream, the latter varying from 166.4 to 2003.0 $\mu\text{g L}^{-1}$, compared with the maximum of 18.8 $\mu\text{g L}^{-1}$ at the Barro Branco stream. Nitrite was too low to be detectable at the Barro Branco stream. The same pattern was observed for orthophosphate concentrations, being barely detectable at Barro Branco stream, and varying from 4.0 to 13.0 $\mu\text{g L}^{-1}$ at the Caiana stream. Total phosphorus, however, tended to have higher concentrations at Barro Branco stream, reaching 50.1 $\mu\text{g L}^{-1}$ at Site 3, but with highly variable values (from non detectable to 38.7 $\mu\text{g L}^{-1}$, as the second highest value).

```

1  #TABELA DE NUTRIENTES----
2
3  ##ORGANIZANDO DADOS----
4
5  dev.off()
6  rm(list=ls(all=TRUE))
7  cat("\014")
8
9  t_nutri <- read.csv("res/t_nutri.csv",
10                   sep = ";", dec = ".",
11                   row.names = 1,
12                   header = TRUE,
13                   na.strings = NA)
14
15  t_grps <- read.csv("res/t_grps.csv",
16                   sep = ";", dec = ".",
17                   row.names = 1,
18                   header = TRUE,
19                   na.strings = NA)
20
21  nutri <- t_nutri
22

```

```

23 library(dplyr)
24 library(tidyr)
25 t_grps <- t_grps[seq_len(nrow(nutri)), , drop = FALSE]
26 m_trab <- nutri %>%
27   rename_with(~ gsub("_", ".", .))
28 m <- m_trab %>%
29   group_by(PontoN = t_grps$PontoN) %>%
30   summarise(across(where(is.numeric),
31                     list(mean = mean, min = min, max = max)),
32             .groups = 'drop') %>%
33   pivot_longer(
34     cols = -c(PontoN),
35     names_to = c("Variable", ".value"),
36     names_sep = "_"
37   )
38
39 m <- as.data.frame(m)
40 colnames(m_trab)
41 m_wide <- m %>%
42   mutate(stat_string = ifelse(Variable == c("Ammonia.NH3.ug.L"),
43                                paste0(round(mean, 1), " (", round(min, 1), "-", round(max, 1)
44                                paste0(round(mean, 1), " (", round(min, 1), "-", round(max, 1)
45   unite("Location", PontoN, sep = "_") %>%
46   select(Variable, Location, stat_string) %>%
47   pivot_wider(names_from = Location, values_from = stat_string)
48
49 m_wide
50 m_wide <- as.data.frame(m_wide)
51 cols <- colnames(m_wide)
52 col_order <- c(
53   "Variable",
54   cols[cols != "Variable"][
55     order(as.numeric(gsub("Ponto", "", cols[cols != "Variable"])))
56   ]
57 )
58 m_wide <- m_wide[, c(col_order)] #sequência das colunas
59 m_wide
60
61 # Salvado m_wide
62 write.table(m_wide, "res/m_wide_nutri.txt",
63             sep = ";", dec = ".", #"\\t",
64             row.names = TRUE,
65             quote = TRUE,

```

```

66         append = FALSE)
67 m_wide_nutri <- read.table("res/m_wide_nutri.txt",
68         sep = ";", dec = ".",
69         row.names = 1,
70         header = TRUE,
71         na.strings = NA)
72
73 # Exportando dados para Excel
74 library(openxlsx)
75 #write.xlsx(m_wide, file = "res/tabela de nutrientes.xlsx", rowNames = FALSE)
76 wb <- loadWorkbook("res/tabela de nutrientes.xlsx")
77 writeData(wb, sheet = "Sheet 1", x = m_wide_nutri)
78 saveWorkbook(wb, "res/tabela de nutrientes.xlsx", overwrite = TRUE)
79
80 # Tabela final ajustada no Excel
81 library(readxl)
82 nutri_gtable <- read_excel(
83     path = "res/tabela de nutrientes.xlsx",
84     sheet = "tabela_final",
85     range = "A1:P6") #ajustar para o tamanho da tabela final
86 nutri_gtable
87 library(gt)
88 nutri_gtable <- gt(nutri_gtable)
89 nutri_gtable <- sub_missing(nutri_gtable,
90     columns = everything(), missing_text = "")
91 nutri_gtable
92 gtsave(nutri_gtable, "res/gt-nutri_gtable.xlsx.html")
93 saveRDS(nutri_gtable, "res/gt-nutri_gtable.xlsx.rds")

```

3.2 Community structure

```

1 #ZOOEBIO_part - CURVA DE ACUMULAÇÃO----
2 #####....----
3
4 ##ORGANIZANDO DADOS----
5
6 dev.off() #apaga os graficos, se houver algum
7 rm(list=ls(all=TRUE)) #limpa a memória
8 cat("\014") #limpa o console
9
10 m_cont <- read.csv("res/m_cont.csv",
11     sep = ";", dec = ".",

```

```

12         row.names = 1,
13         header = TRUE,
14         na.strings = NA)
15 m_trab <- m_cont
16
17 library(tidyverse)
18
19 m_trab_r <- mutate(m_trab, across(everything(), ceiling)) #arredonda a matriz original para
20
21 colSums(t(m_trab_r))
22 colnames(t(m_trab_r))
23
24 ca_abund <- colSums(m_trab_r[grepl("-C-", rownames(m_trab_r)), ])
25 bb_abund <- colSums(m_trab_r[grepl("-B-", rownames(m_trab_r)), ])
26
27 m_rios <- rbind("Caiana" = ca_abund, "Barro Branco" = bb_abund)
28 m_rios
29
30 rarefa <- t(m_rios)
31
32 library(iNEXT)
33 out <- iNEXT(rarefa, q = 0,
34             datatype = "abundance",
35             size = NULL,
36             endpoint = 15000, #define o comprimento de eixo x
37             knots = 40,
38             se = TRUE,
39             conf = 0.95,
40             nboot = 50)
41
42 library(ggplot2)
43 grafico3 <- ggiNEXT(out, type = 1, facet.var="None") + #ver como fica com facet.var="Assembl
44   theme_bw() +
45   labs(fill = "Áreas") +
46   xlab("Número de indivíduos") +
47   ylab("Riqueza de espécies") +
48   scale_linetype_discrete(
49     labels = c("Interpolation", "Extrapolation")) +
50   theme(legend.title = element_blank(),
51         legend.position = "bottom",
52         legend.text = element_text(size = 14),
53         axis.title = element_text(size = 14),
54         axis.text = element_text(size = 12))

```

```

55 )
56 grafico3
57 ggsave(grafico3, width = 8, height = 6, dpi = 300, filename = "res/fig-rare1.png")

1  #ZOOEBIO_part - ESTR. DA COMUNIDADE-----
2  #####....-----
3
4  ##ORGANIZANDO DADOS-----
5
6  dev.off() #apaga os graficos, se houver algum
7  rm(list=ls(all=TRUE)) #limpa a memória
8  cat("\014") #limpa o console
9
10 t_grps <- read.csv("res/t_grps.csv",
11                   sep = ";", dec = ".",
12                   row.names = 1,
13                   header = TRUE,
14                   na.strings = NA)
15 m_dens <- read.csv("res/m_dens.csv",
16                  sep = ";", dec = ".",
17                  row.names = 1,
18                  header = TRUE,
19                  na.strings = NA)
20 m_hab_part <- read.csv("res/m_hab_part.csv",
21                      sep = ";", dec = ".",
22                      row.names = 1,
23                      header = TRUE,
24                      na.strings = NA)
25 m_cont <- read.csv("res/m_cont.csv",
26                   sep = ";", dec = ".",
27                   row.names = 1,
28                   header = TRUE,
29                   na.strings = NA)
30
31 library(tibble)
32 library(tidyverse)
33 library(forcats)
34 library(openxlsx)
35 library(Rcpp)
36
37 m_dens
38 str(m_dens)
39 m_trab <- m_dens

```

```

40
41 m_trab <- (m_dens) #SITES
42 m_trab <- t(m_dens) #SPP
43
44 ##TAMANHO DA MATRIZ-----
45
46 range(m_trab) #menor e maior valores
47 length(m_trab) #no. de colunas
48 ncol(m_trab) #no. de N colunas
49 nrow(m_trab) #no. de M linhas
50 sum(lengths(m_trab)) #soma os nos. de colunas
51 length(as.matrix(m_trab)) #tamanho da matriz m x n
52 sum(m_trab == 0) #número de observações igual a zero
53 sum(m_trab > 0) #número de observações maiores que zero
54 #calculando a proporção de zeros na matriz
55 zeros <- (sum(m_trab == 0)/length(as.matrix(m_trab)))*100
56 zeros
57
58 ##ESTRUTURA DA COMUNIDADE-----
59
60 Sum <- rowSums(m_trab)
61 #ou
62 Sum <- apply(m_trab,1,sum)
63 Sum
64 # Abundância relativa (%)
65 RA <- (Sum / sum(Sum)) * 100 # percentage
66 # Media
67 Mean <- rowMeans(m_trab)
68 Mean
69 # Ou
70 Mean <- apply(m_trab,1,mean)
71 Mean
72 # Desvio padrão
73 DP <- apply(m_trab,1,sd)
74 DP
75 # Máximo
76 Max <- apply(m_trab,1,max)
77 Max
78 # Mínimo
79 Min <- apply(m_trab,1,min)
80 Min
81 # Mínimo não-zero
82 MinZ <- apply(m_trab, 1, function(row) {

```

```

83   non_zero_values <- row[row > 0] # Filter out zero values
84   if (length(non_zero_values) == 0) {
85     return(0) # If all values are zero, return 0
86   } else {
87     return(min(non_zero_values)) # Return the minimum of non-zero values
88   }
89 })
90 MinZ
91
92 m_pa <- m_trab
93 m_pa[m_pa != 0] <- 1
94 rowSums(m_pa)
95 library(vegan)
96 bin <- decostand(m_trab,"pa")
97 bin[1:10, 1:10]
98 S <- apply(bin,1,sum)
99 S
100 #OU
101 Riqueza <- specnumber(m_trab)
102 Riqueza
103 Riqueza_total <- specnumber(colSums(m_trab))
104 Riqueza_total
105
106 FO <- rowSums(m_trab > 0) / ncol(m_trab) * 100
107 FO
108
109 H <- diversity(m_trab, index = "shannon")
110 H
111 D <- diversity(m_trab, "simpson")
112 D
113 D[is.na(D)] <- 0 #substitui NA ou NaN por 0
114 D
115 E <- H/log(specnumber(m_trab))
116 E
117 E[is.na(E)] <- 0 #substitui NA ou NaN por 0
118 E
119
120 ###DESCRITORES-----
121 Descritores1 <- cbind(Sum, RA, Mean, DP, Max, Min, MinZ, FO, S, E, H, D)
122 Descritores1 <- as.data.frame(Descritores1)
123 Descritores1
124 Descritores1 <- Descritores1[order(-Descritores1$Mean), ] #ordena do maior para o menor "-"
125 #Descritores1 <- Descritores1 %>% rownames_to_column(var="Espécies") #da nome a primeira col.

```

```

126 SomaTotalD <- apply(Descritores1,2,sum)
127 SomaTotalD
128 MediaTotalD <- apply(Descritores1,2,mean)
129 MediaTotalD
130 DPTotalD <- apply(Descritores1,2,sd)
131 DPTotalD
132 Descritores2 <- cbind(SomaTotalD, MediaTotalD, DPTotalD)
133 Descritores2 <- as.data.frame(Descritores2)
134 Descritores2 <- t(Descritores2)
135 Descritores2
136 DescritoresFinal <- rbind(Descritores1, Descritores2)
137 DescritoresFinal
138 DescritoresFinal <- round (DescritoresFinal, 2)
139 DescritoresFinal
140
141 # Salvando DescritoresFinal
142 write.table(data.frame("SPP"=rownames(DescritoresFinal),
143                       DescritoresFinal),
144             "res/DescritoresSPP.txt",
145             row.names=FALSE,
146             sep="\t")
147
148 ###TABELA FINAL DESCRITORES----
149 library(gt)
150 df <- DescritoresFinal
151 ncol(df); nrow(df) #no. de N colunas x M linhas
152 df <- cbind(SPP = rownames(df), df)
153 gt <- gt(df, rowname_col = "SPP",
154         caption = "Descritores da diversidade por espécie (colunas). Sum, soma; RA, abundância re")
155 gt
156 gtsave(gt, "res/DescritoresSPP.html")
157 gtsave(gt, "res/DescritoresSPP.png")
158
159 ###NORMALIDADE----
160 library(moments)
161 Assimetria <- apply(m_trab,1,skewness)
162 Assimetria
163 Curtose <- apply(m_trab,1,kurtosis)
164 Curtose
165 Normalidade1 <- cbind(Assimetria, Curtose)
166 Normalidade1 <- as.data.frame(Normalidade1)
167 Normalidade1
168 SomaTotalN <- apply(Normalidade1,2,sum)

```

```

169 SomaTotalN
170 MediaTotalN <- apply(Normalidade1,2,mean)
171 MediaTotalN
172 DPTotalN <- apply(Normalidade1,2,sd)
173 DPTotalN
174 Normalidade2<-cbind(SomaTotalN, MediaTotalN, DPTotalN)
175 Normalidade2<-as.data.frame(Normalidade2)
176 Normalidade2 <- t(Normalidade2) #"t" transpoe a matriz
177 Normalidade2
178 NormalidadeFinal <- rbind(Normalidade1, Normalidade2)
179 NormalidadeFinal
180 NormalidadeFinal <- round(NormalidadeFinal, 2)
181 NormalidadeFinal
182
183 # Salvando NormalidadeFinal
184 write.table(data.frame("SPP"=rownames(NormalidadeFinal),
185                        NormalidadeFinal),
186             "res/NormalidadeSPP.txt",
187             row.names=FALSE,
188             sep="\t")
189
190 ###TABELA FINAL NORMALIDADE----
191 nf <- NormalidadeFinal
192 ncol(nf); nrow(nf) #no. de N colunas x M linhas
193 nf <- cbind(SPP = rownames(nf), nf)
194 gt <- gt(nf, rowname_col = "SPP", caption = "Descritores da normalidade por SPP (COLUNA)")
195 gt
196 gtsave(gt, "res/NormalidadeSPP.html")
197
198 #DIVERSIDADE EM DENSIDADE MATRIZ NÃO-TRANSPOSTA----
199
200 m_trab <- (m_dens) #HILL
201
202 # Números de Hill
203 Hill <- renyi(m_trab,
204              scales = c(0:5),
205              hill = TRUE)
206 Hill
207 library(ggplot2)
208 library(tidyr)
209 library(tidyverse)
210 grafico1 <- Hill %>%
211   rownames_to_column() %>%

```

```

212   pivot_longer(-rowname) %>%
213   mutate(name = factor(name, name[1:length(Hill)])) %>%
214   ggplot(aes(x = name, y = value, group = rowname,
215             col = rowname)) +
216   geom_point(size = 2) +
217   geom_line(size = 1) +
218   xlab("Parâmetro de ordem de diversidade (q)") +
219   ylab("Diversidade") +
220   labs(col = "Locais") +
221   theme_bw() +
222   theme(text = element_text(size = 10)) #ajustar a fonte caso nao caiba no output html
223   grafico1
224   ggsave(grafico1, dpi = 300, filename = "res/fig-hill.png")
225
226   # Gráfico de abundância
227   abund <- colSums(m_trab)
228   #abund <- m_trab[1, ] #escolhe a primeira linha para a distribuição de abundância
229   df <- data.frame(sp = colnames(m_trab),
230                   abun = abund)
231   grafico2 <- ggplot(df, aes(fct_reorder(sp, -abun),
232                                abun, group = 1)) +
233   geom_col() +
234   geom_line(col = "red", linetype = "dashed") +
235   geom_point(col = "red") +
236   xlab("Espécies") +
237   ylab("Abundância") +
238   theme_bw() +
239   theme(axis.text.x = element_text(
240     angle = 45,
241     hjust = 1,
242     face = "italic",
243     size = 10),
244     axis.text.y = element_text(size = 10),
245     axis.title.x = element_text(size = 12),
246     axis.title.y = element_text(size = 12))
247   grafico2
248   ggsave(grafico2, dpi = 300, filename = "res/fig-abun.png")
249
250   # Curvas de rarefação
251   m_trab_r <- mutate(as.data.frame(m_trab), across(everything(), ceiling)) #arredonda a matriz
252   colSums(t(m_trab_r))
253   colnames(t(m_trab_r))
254   #rarefa <- t(m_trab_r[c("S-R-CT2", "S-R-CP2", "S-A-TA2", "B-A-MU2"),]) #curva de rarefação para

```

```

255 class(m_trab)
256
257 # Curva de rarefação para todos os sítios
258 rarefa <- t(m_trab_r)
259 ## Curva de rarefação para as 8 UA's com maior soma
260 #m_trab_r <- as.data.frame(t(m_trab_r)) #transpõe a matriz
261 #col_sums <- colSums(m_trab_r)
262 #largest_columns <- names(sort(col_sums, decreasing = TRUE)[1:8])
263 #rarefa <- m_trab_r[largest_columns] #curva de rarefação para as 8 UA's com maior soma
264 library(iNEXT)
265 out <- iNEXT(rarefa, q = 0,
266             datatype = "abundance",
267             size = NULL,
268             endpoint = 2000, #define o comprimento de eixo x
269             knots = 40,
270             se = TRUE,
271             conf = 0.95,
272             nboot = 50)
273 grafico3 <- ggiNEXT(out, type = 1, facet.var="None") +
274   theme_bw() +
275   labs(fill = "Áreas") +
276   xlab("Número de indivíduos") +
277   ylab("Riqueza de espécies") +
278   theme(legend.title=element_blank()) #ver como fica com facet.var="Assemblage"
279 #grafico3
280 #ggsave(grafico3, dpi = 300, filename = "fig-rare1.png")
281
282 # Curvas de acumulação de espécies
283 acumula <- specaccum(m_trab,
284                     method = "random")
285
286 acumula
287 plot(acumula)
288 plot_data <- data.frame("UAs" = c(0, acumula$sites),
289                         "Riqueza" = c(0, acumula$richness),
290                         "lower" = c(0, acumula$richness - acumula$sd),
291                         "upper" = c(0, acumula$richness + acumula$sd))
292 gLocais <- ggplot(plot_data, aes(x = UAs, y = Riqueza)) +
293   geom_point(color = "blue", size = 4) +
294   geom_line(color = "blue", lwd = 2) +
295   geom_ribbon(aes(ymin = lower, ymax = upper),
296             linetype=2, alpha=0.3, fill = "yellow") +
297   ylab("Riqueza acumulada") +
298   theme_classic() +

```

```

298   theme(text = element_text(size = 16))
299 gLocais
300 ggsave(gLocais, dpi = 300, filename = "res/fig-acum.png")
301
302 plot_data <- data.frame("Individuos" = c(0, acumula$individuals),
303                        "Riqueza" = c(0, acumula$richness),
304                        "lower" = c(0, acumula$richness - acumula$sd),
305                        "upper" = c(0, acumula$richness + acumula$sd))
306 gInd <- ggplot(plot_data, aes(x = Individuos, y = Riqueza)) +
307   geom_point(color = "blue", size = 4) +
308   geom_line(color = "blue", lwd = 2) +
309   geom_ribbon(aes(ymin = lower, ymax = upper),
310             linetype=2, alpha=0.3, fill = "yellow") +
311   ylab("Riqueza acumulada") +
312   theme_classic() +
313   theme(text = element_text(size = 16))
314 gInd
315
316 #RIQUEZA RAREFACIONADA----
317
318 # Rarefação baseada na matriz de contagem
319 library(vegan)
320 data <- m_cont #raremax precisa de numeros inteiros
321 S <- specnumber(data) #observed number of species
322 S
323 #raremax <- min(rowSums(data))
324 raremax <- round(mean(rowSums(data)))
325 #raremax <- 1000
326 raremax
327 Srare <- rarefy(data, raremax)
328 Srare
329 plot(S, Srare, xlab = "Observed No. of Species", ylab = "Rarefied No. of Species")
330 abline(0, 1)
331
332 #png("res/fig-rarecurve.png")
333 rarecurve(data, step = 20,
334           sample = raremax, #linhas de rarefação
335           col = "blue", cex = 0.6,
336           xlab = "Sample size", ylab = "Taxa", xlim = c()) #c(0,5000)
337 #dev.off()
338
339 ##TABELA DA COMUNIDADE----
340

```

```

341 library(dplyr)
342 library(tidyr)
343 m_trab <- m_dens %>%
344   rename_with(~ gsub("_", ".", .))
345 m <- m_trab %>%
346   group_by(Rio = t_grps$Rio,
347             Período = t_grps$Período,
348             Ponto = t_grps$PontoN) %>%
349   summarise(across(where(is.numeric),
350                     list(mean = mean, sd = sd)),
351             .groups = 'drop') %>%
352   pivot_longer(
353     cols = -c(Rio, Período, Ponto),
354     names_to = c("Variable", ".value"),
355     names_sep = "_"
356   )
357 m
358 m <- as.data.frame(m)
359 m_wide <- m %>%
360   mutate(stat_string = ifelse(
361     Variable == "nome_da_variavel", #escolhe uma variável pra ter 3 casas decimais
362     paste0(round(mean, 3), " (", round(sd, 3), ")"),
363     paste0(round(mean, 1), " (", round(sd, 1), ")")
364   )) %>%
365   unite("Location", Rio, Período, Ponto, sep = "_") %>%
366   dplyr::select(Variable, Location, stat_string) %>% #dplyr dá conflito com MASS
367   pivot_wider(names_from = Location, values_from = stat_string)
368
369 m_wide
370 m_wide <- as.data.frame(m_wide)
371 m_wide
372
373 # Salvado m_wide
374 write.table(m_wide, "res/m_wide_spp.txt",
375             sep = ";", dec = ".", #"\\t",
376             row.names = TRUE,
377             quote = TRUE,
378             append = FALSE)
379 m_wide_spp <- read.table("res/m_wide_spp.txt",
380                          sep = ";", dec = ".",
381                          row.names = 1,
382                          header = TRUE,
383                          na.strings = NA)

```

```

384
385 # Exportando dados para Excel
386 library(openxlsx)
387 #write.xlsx(m_wide_spp, file = "res/tabela da comunidade.xlsx", rowNames = FALSE)
388 wb <- loadWorkbook("res/tabela da comunidade.xlsx")
389 writeData(wb, sheet = "Sheet 1", x = m_wide_spp)
390 saveWorkbook(wb, "res/tabela da comunidade.xlsx", overwrite = TRUE)
391
392 # Tabela final ajustada no Excel
393 library(readxl)
394 spp_gtttable <- read_excel(
395   path = "res/tabela da comunidade.xlsx",
396   sheet = "tabela_final",
397   col_names = T,
398   range = "A1:AE61") #ajustar para o tamanho da tabela final
399 spp_gtttable
400 library(gt)
401 spp_gtttable <- gt(spp_gtttable)
402 spp_gtttable <- sub_missing(spp_gtttable,
403   columns = everything(), missing_text = "")
404 #spp_gtttable <- fmt_number(spp_gtttable,
405 #   columns = "F.O.(%)", decimals = 1)
406 spp_gtttable
407 gtsave(spp_gtttable, "res/gt-spp_gtttable_xlsx.html")
408 saveRDS(spp_gtttable, "res/gt-spp_gtttable_xlsx.rds")
409
410 1 # Escolher sumário de uma variavel
411 2 m
412 3 var <- "h.roots"
413 4 m[m$Variable == var, "mean"] #cada valor de var
414 5 summary(m[m$Variable == var, "mean"]) #sumário dos valores de var
415 6
416 7 # Escolher sumário de um grupo de variáveis do df m
417 8 vars <- unique(grep("^h\\.", m$Variable, value = TRUE))
418 9 summaries <- list() #cria uma lista vazia para guardar os sumários
419 10 # Loop para cada variável do grupo e guarda em summaries
420 11 for (var in vars) {
421 12   summaries[[var]] <- summary(m[m$Variable == var, "mean"])
422 13 }
423 14 #var is a temporary variable used in the for loop to iterate through
424 15 #each variable name that starts with "h."
425 16
426 17 summaries

```

```

18 summary_table <- do.call(rbind, lapply(summaries, as.data.frame.list))
19 round(summary_table, 2)
20 #sink(file = "summary_h.txt", split = TRUE)
21 round(summary_table[order(summary_table$Mean, decreasing = FALSE), ], 2)
22 #sink()
23
24 # Tabela limpa
25 #summary_table <- cbind(Variable = rownames(summary_table), summary_table)
26 #rownames(summary_table) <- NULL
27 #colnames(summary_table) <- c("Variable", "Min", "Q1", "Median", "Mean", "Q3", "Max")
28 #summary_table

1 #ZOOEBIO_part - UPGMA----
2 #####....-----
3
4 ##ORGANIZANDO DADOS----
5
6 dev.off() #apaga os graficos, se houver algum
7 rm(list=ls(all=TRUE)) #limpa a memória
8 cat("\014") #limpa o console
9
10 t_grps <- read.csv("res/t_grps.csv",
11                   sep = ";", dec = ".",
12                   row.names = 1,
13                   header = TRUE,
14                   na.strings = NA)
15 m_dens <- read.csv("res/m_dens.csv",
16                   sep = ";", dec = ".",
17                   row.names = 1,
18                   header = TRUE,
19                   na.strings = NA)
20 m_hab_part <- read.csv("res/m_hab_part.csv",
21                       sep = ";", dec = ".",
22                       row.names = 1,
23                       header = TRUE,
24                       na.strings = NA)
25 m_cont <- read.csv("res/m_cont.csv",
26                   sep = ";", dec = ".",
27                   row.names = 1,
28                   header = TRUE,
29                   na.strings = NA)
30
31 m_dens

```

```

32 str(m_dens)
33
34 #SEPARANDO OS PERÍODOS
35
36 library(vegan)
37 library(cluster)
38
39 # Fazendo a média por ponto
40 ponto_id <- sub("-H[0-9]+$", "", rownames(m_dens))
41 m_dens_media <- aggregate(m_dens, by = list(Ponto = ponto_id), FUN = mean)
42 rownames(m_dens_media) <- m_dens_media$Ponto
43 m_dens_media$Ponto <- NULL
44 # Separar por período
45 seco_med <- m_dens_media[startsWith(rownames(m_dens_media), "S-"), ]
46 chuv_med <- m_dens_media[startsWith(rownames(m_dens_media), "C-"), ]
47 # Filtrar COLUNAS zeradas (espécies ausentes no período)
48 seco_med <- seco_med[, colSums(seco_med) > 0]
49 chuv_med <- chuv_med[, colSums(chuv_med) > 0]
50 # Filtrar LINHAS zeradas (pontos sem organismos)
51 seco_med <- seco_med[rowSums(seco_med) > 0, ]
52 chuv_med <- chuv_med[rowSums(chuv_med) > 0, ]
53 # Transformação
54 seco_trns <- asin(sqrt(decostand(seco_med,
55                                method="total", MARGIN = 2)))
56 chuv_trns <- asin(sqrt(decostand(chuv_med,
57                                method="total", MARGIN = 2)))
58 # Distâncias de Bray-Curtis e UPGMA
59 dist_seco <- vegdist(seco_trns, method = "bray")
60 dist_chuv <- vegdist(chuv_trns, method = "bray")
61 upgma_seco <- hclust(dist_seco, method = "average")
62 upgma_chuv <- hclust(dist_chuv, method = "average")
63
64 # Coeficientes de aglomeração
65 agg_coef_seco <- coef.hclust(upgma_seco)
66 agg_coef_chuv <- coef.hclust(upgma_chuv)
67 # Gerar a matriz de distância cofenética a partir da árvore UPGMA
68 cofe_seco <- cophenetic(upgma_seco)
69 cofe_chuv <- cophenetic(upgma_chuv)
70 # Calcular a correlação com a matriz de distância original (Bray-Curtis)
71 cor_seco <- cor(dist_seco, cofe_seco)
72 cor_chuv <- cor(dist_chuv, cofe_chuv)
73 # Visualizar os resultados
74 cat("Correlação Cofenética (Seco):", round(cor_seco, 3), "\n")

```

```

75 cat("Correlação Cofenética (Chuvoso):", round(cor_chuv, 3), "\n")
76
77 # Plot dos dendrogramas agrupados por ponto
78 upgma_seco$labels <- sub("[A-Z]-", "", upgma_seco$labels)
79 upgma_chuv$labels <- sub("[A-Z]-", "", upgma_chuv$labels)
80 dend_seco <- as.dendrogram(hclust(dist_seco, method = "average"))
81 dend_chuv <- as.dendrogram(hclust(dist_chuv, method = "average"))
82
83 #png("res/fig-upgma.png", width=11, height=9, units="in", res=400)
84 par(mfrow = c(1, 2), mar = c(5, 4, 2, 6)) #B, L, T, R
85 plot(as.dendrogram(upgma_seco),
86      horiz = TRUE,
87      xlab = "Bray-Curtis dissimilariry",
88      xlim = c(max(attr(dend_seco, "height")), 0),
89      cex.lab = 1)
90 title(main = "Dry", line = 0, cex.main = 1.2)
91 mtext(paste0("Agglomerative Coefficient = ", round(agg_coef_seco, 2)),
92      side = 2,                # Lado esquerdo (eixo Y)
93      line = 1,                # Distância da margem
94      at = 7,                  # Posição na base do eixo
95      las = 0,                 # Orientação do texto
96      cex = 0.8)               # Tamanho da fonte
97 mtext(paste0("Cophenetic Coefficient = ", round(cor_seco, 2)),
98      side = 2,                # Lado esquerdo (eixo Y)
99      line = 2,                # Distância da margem
100     at = 7,                  # Posição na base do eixo
101     las = 0,                 # Orientação do texto
102     cex = 0.8)               # Tamanho da fonte
103
104 plot(as.dendrogram(upgma_chuv),
105      horiz = TRUE,
106      xlab = "Bray-Curtis dissimilariry",
107      xlim = c(max(attr(dend_chuv, "height")), 0),
108      cex.lab = 1)
109 title(main = "Wet", line = 0, cex.main = 1.2)
110 mtext(paste0("Agglomerative Coefficient = ", round(agg_coef_chuv, 2)),
111      side = 2,                # Lado esquerdo (eixo Y)
112      line = 1,                # Distância da margem
113      at = 7,                  # Posição na base do eixo
114      las = 0,                 # Orientação do texto
115      cex = 0.8)               # Tamanho da fonte
116 mtext(paste0("Cophenetic Coefficient = ", round(cor_chuv, 2)),
117      side = 2,                # Lado esquerdo (eixo Y)

```

```

118     line = 2,          # Distância da margem
119     at = 7,           # Posição na base do eixo
120     las = 0,          # Orientação do texto
121     cex = 0.8)        # Tamanho da fonte
122 par(mfrow = c(1, 1))
123 #dev.off()
124
125 #OUTRA FORMA
126 #plot(as.dendrogram(upgma_chuv),
127 #     horiz = TRUE,
128 #     main = "Wet",
129 #     xlab = "Bray-Curtis dissimilarity",
130 #     xlim = c(max(attr(dend_chuv, "height")), 0))
131 # Adiciona o texto no canto inferior esquerdo
132 #text(x = max(attr(dend_chuv, "height")) * 0.9,
133 #     y = 3,
134 #     labels = paste0("Agglomerative Coefficient = ", round(agg_coef_chuv, 2)),
135 #     adj = c(0, 0),
136 #     cex = 0.85,
137 #     srt = 90) # srt = 90 gira o texto para ficar vertical
138 #OU
139 #mtext(paste0("Cophenetic Coefficient = ", round(cor_seco, 2)),
140 #     side = 2,      # 1 = bottom margin
141 #     line = 2.5,    # Line distance below axis (increase if it overlaps numbers)
142 #     adj = 0.36,    # Posição horizontal
143 #     cex = 0.85)   # Font size

```

The UPGMA cluster analyses revealed low-to-moderate spatial clustering strength for the zooplankton communities across both periods, as indicated by the Agglomerative Coefficients ($AC_{dry} = 0.34$ and $AC_{wet} = 0.39$). While community composition exhibited continuous variation across sites, spatial structuring was slightly more pronounced during the wet season than during the dry season.

3.3 Diversity Partitioning

```

1 #ZOOREBIO_part - ADDPART----
2 #####....----
3
4 ##ORGANIZANDO DADOS----
5
6 dev.off() #apaga os graficos, se houver algum
7 rm(list=ls(all=TRUE)) #limpa a memória

```

```

8  cat("\014") #limpa o console
9
10 m_esc <- read.csv("res/m_esc.csv",
11                  sep = ";", dec = ".",
12                  row.names = 1,
13                  header = TRUE,
14                  na.strings = NA)
15 m_dens <- read.csv("res/m_dens.csv",
16                  sep = ";", dec = ".",
17                  row.names = 1,
18                  header = TRUE,
19                  na.strings = NA)
20 m_cont <- read.csv("res/m_cont.csv",
21                  sep = ";", dec = ".",
22                  row.names = 1,
23                  header = TRUE,
24                  na.strings = NA)
25
26 #BUSCANDO AS ESCALAS
27
28 # Partição total
29 colnames(m_esc)
30 pt.total <- m_esc[, c("pt.rio", "pt.periodo", "pt.rebio")]
31 head(pt.total)
32 dim(pt.total)
33
34 # Partição seco
35 pt.seco <- m_esc[, c("s.habitat", "s.ponto", "s.rio", "s.rebio")]
36 pt.seco <- pt.seco[grepl("^S-", rownames(pt.seco)), ]
37 head(pt.seco)
38 dim(pt.seco)
39 m_cont_s <- m_cont[grepl("^S-", rownames(m_cont)), ]
40
41 # Partição chuvoso
42 pt.chuva <- m_esc[, c("c.habitat", "c.ponto", "c.rio", "c.rebio")]
43 pt.chuva <- pt.chuva[grepl("^C-", rownames(pt.chuva)), ]
44 head(pt.chuva)
45 dim(pt.chuva)
46 m_cont_c <- m_cont[grepl("^C-", rownames(m_cont)), ]
47
48 # Partição bb.seco (A)
49 bb.seco <- m_esc[, c("bb.habitat", "bb.ponto", "bb.rio")]
50 bb.seco <- bb.seco[grepl("S-B-", rownames(bb.seco)), ]

```

```

51 head(bb.seco)
52 dim(bb.seco)
53 m_cont_bb.seco <- m_cont[grepl("S-B-", rownames(m_cont)), ]
54
55 # Partição bb.chuva (B)
56 bb.chuva <- m_esc[, c("bb.habitat", "bb.ponto", "bb.rio")]
57 bb.chuva <- bb.chuva[grepl("C-B-", rownames(bb.chuva)), ]
58 head(bb.chuva)
59 dim(bb.chuva)
60 m_cont_bb.chuva <- m_cont[grepl("C-B-", rownames(bb.chuva)), ]
61
62 # Partição ca.seco (C)
63 ca.seco <- m_esc[, c("ca.habitat", "ca.ponto", "ca.rio")]
64 ca.seco <- ca.seco[grepl("S-C-", rownames(ca.seco)), ]
65 head(ca.seco)
66 dim(ca.seco)
67 m_cont_ca.seco <- m_cont[grepl("S-C-", rownames(ca.seco)), ]
68
69 # Partição ca.chuva (D)
70 ca.chuva <- m_esc[, c("ca.habitat", "ca.ponto", "ca.rio")]
71 ca.chuva <- ca.chuva[grepl("C-C-", rownames(ca.chuva)), ]
72 head(ca.chuva)
73 dim(ca.chuva)
74 m_cont_ca.chuva <- m_cont[grepl("C-C-", rownames(ca.chuva)), ]
75
76 #PARTIÇÃO
77
78 #pt.seco (m_cont_s), pt.chuva (m_cont_c)
79 #bb.seco(A) (m_cont_bb.seco), bb.chuva(B) (m_cont_bb.chuva)
80 #ca.seco(C) (m_cont_ca.seco), ca.chuva(D) (m_cont_ca.chuva)
81
82 part1 <- ca.chuva
83 cont <- m_cont_ca.chuva
84
85 library(vegan)
86
87 sink(file = "res/part-ca.chuva.txt")
88 #VALORES ABSOLUTOS
89 part.a <- adpart(y = cont, x = part1, index = "richness", relative = FALSE,
90                 weights = "unif", nsimul = 999)
91 part.a
92
93 #EM PORCENTAGEM

```

```

94 part.p <- adipart(y = cont, x = part1, index = "richness", relative = TRUE,
95                  weights = "unif", nsimul = 999)
96 part.p
97
98 #str(part.p)
99
100 # Extrair valores do objeto part.p
101
102 part2 <- part.p
103
104 statistic <- part2$statistic
105 SES <- part2$oecosimu$z
106 mean <- part2$oecosimu$means
107 lower_2.5 <- apply(part2$oecosimu$simulated, 1, quantile, probs = 0.025)
108 median_50 <- apply(part2$oecosimu$simulated, 1, quantile, probs = 0.50)
109 upper_97.5 <- apply(part2$oecosimu$simulated, 1, quantile, probs = 0.975)
110 Pr_sim <- part2$oecosimu$pval
111
112 # Definir função para adicionar significância
113 get_significance <- function(p) {
114   if (p <= 0.001) {
115     return("****")
116   } else if (p <= 0.01) {
117     return("***")
118   } else if (p <= 0.05) {
119     return("*")
120   } else if (p <= 0.1) {
121     return(".")
122   } else {
123     return(" ")
124   }
125 }
126 significance <- sapply(Pr_sim, get_significance)
127
128 # Criar dataframe com todas as informações
129
130 diversity_data <- data.frame(
131   Component = names(statistic),
132   Observed = round(statistic, 2),
133   SES = round(SES, 2),
134   Mean = round(mean, 2),
135   Lower_2.5 = round(lower_2.5, 2),
136   Median_50 = round(median_50, 2),

```

```

137   Upper_97.5 = round(upper_97.5, 2),
138   Pr_sim = Pr_sim,
139   Sign = significance
140 )
141
142 # Exibir a tabela
143 library(gt)
144 print(diversity_data)
145 #str(part2)
146 gt(diversity_data)
147
148 # Criar dataframe com valores observados e esperados
149 observed_values <- part2$statistic
150 observed_values
151 expected_values <- part2$oeocosimu$means
152 expected_values
153
154 # Combine observed and expected values into a dataframe
155 df <- data.frame(
156   Component = rep(names(observed_values), 2), # Use names from observed_values
157   Type = rep(c("Observed", "Expected"), each = length(observed_values)),
158   Value = c(observed_values, expected_values) # Observado primeiro, Esperado depois
159 )
160
161 df$Type <- factor(df$Type, levels = c("Observed", "Expected"))
162 df
163 df_filtered <- df[!df$Component %in% c("alpha.2", "alpha.3", "gamma"), ]
164 df_filtered$Component <- factor(df_filtered$Component, levels = c("beta.3", "beta.2", "beta.1"))
165 df_filtered
166
167 diversity_data <- df_filtered
168 diversity_data
169
170 # Calcular porcentagens para cada componente
171 diversity_data <- diversity_data %>%
172   group_by(Type) %>%
173   mutate(Percentage = Value / sum(Value) * 100)
174
175 hier <- hiersimu(y = cont, x = part1, FUN = diversity)
176 hier
177 sink()
178
179 #GRÁFICOS

```

```

180
181 library(ggplot2)
182 library(dplyr)
183
184 # Partição total
185
186 ggplot_pt.total <- ggplot(diversity_data, aes(fill = Component, y = Percentage, x = Type)) +
187   geom_bar(position = "stack", stat = "identity") +
188   labs(title = "Total",
189        y = "% of the total diversity",
190        x = "") +
191   scale_fill_manual(values = c("alpha.1" = "black",
192                                "beta.1" = "darkgray",
193                                "beta.2" = "gray",
194                                "beta.3" = "lightgray")) +
195   theme_minimal() +
196   theme(
197     axis.text.x = element_text(angle = 0, hjust = 0.5),
198     panel.grid = element_blank(),
199   )
200 ggplot_pt.total
201 ggsave(ggplot_pt.total, dpi = 300, filename = "res/fig-part_total.png")
202
203 # Partição seco
204
205 ggplot_pt.seco <- ggplot(diversity_data, aes(fill = Component, y = Percentage, x = Type)) +
206   geom_bar(position = "stack", stat = "identity") +
207   labs(title = "Seco",
208        y = "% of the total diversity",
209        x = "") +
210   scale_fill_manual(values = c("alpha.1" = "black",
211                                "beta.1" = "darkgray",
212                                "beta.2" = "gray",
213                                "beta.3" = "lightgray")) +
214   theme_minimal() +
215   theme(
216     axis.text.x = element_text(angle = 0, hjust = 0.5),
217     panel.grid = element_blank(),
218   )
219 ggplot_pt.seco
220 ggsave(ggplot_pt.seco, dpi = 300, filename = "res/fig-part_seco.png")
221
222 # Partição chuvoso

```

```

223
224 ggplot_pt.chuva <- ggplot(diversity_data, aes(fill = Component, y = Percentage, x = Type)) +
225   geom_bar(position = "stack", stat = "identity") +
226   labs(title = "Chuvoso",
227         y = "% of the total diversity",
228         x = "") +
229   scale_fill_manual(values = c("alpha.1" = "black",
230                                "beta.1" = "darkgray",
231                                "beta.2" = "gray",
232                                "beta.3" = "lightgray")) +
233   theme_minimal() +
234   theme(
235     axis.text.x = element_text(angle = 0, hjust = 0.5),
236     panel.grid = element_blank(),
237   )
238 ggplot_pt.chuva
239 ggsave(ggplot_pt.total, dpi = 300, filename = "res/fig-part_chuva.png")
240
241 # bb.seco (A)
242
243 ggplot_bb.seco <- ggplot(diversity_data, aes(fill = Component, y = Percentage, x = Type)) +
244   geom_bar(position = "stack", stat = "identity") +
245   labs(title = "(A)",
246         y = "% of the total diversity",
247         x = "") +
248   scale_fill_manual(values = c("alpha.1" = "black",
249                                "beta.1" = "darkgray",
250                                "beta.2" = "gray",
251                                "beta.3" = "lightgray")) +
252   theme_minimal() +
253   theme(
254     axis.text.x = element_text(angle = 0, hjust = 0.5),
255     panel.grid = element_blank(),
256   )
257 ggplot_bb.seco
258 ggsave(ggplot_bb.seco, dpi = 300, filename = "res/fig-bb.seco.png")
259
260 # bb.chuva (B)
261
262 ggplot_bb.chuva <- ggplot(diversity_data, aes(fill = Component, y = Percentage, x = Type)) +
263   geom_bar(position = "stack", stat = "identity") +
264   labs(title = "(B)",
265         y = "% of the total diversity",

```

```

266     x = "") +
267   scale_fill_manual(values = c("alpha.1" = "black",
268                               "beta.1" = "darkgray",
269                               "beta.2" = "gray",
270                               "beta.3" = "lightgray")) +
271   theme_minimal() +
272   theme(
273     axis.text.x = element_text(angle = 0, hjust = 0.5),
274     panel.grid = element_blank(),
275   )
276 ggplot_bb.chuva
277 ggsave(ggplot_bb.chuva, dpi = 300, filename = "res/fig-bb.chuva.png")
278
279 # ca.seco (C)
280
281 ggplot_ca.seco <- ggplot(diversity_data, aes(fill = Component, y = Percentage, x = Type)) +
282   geom_bar(position = "stack", stat = "identity") +
283   labs(title = "(C)",
284        y = "% of the total diversity",
285        x = "") +
286   scale_fill_manual(values = c("alpha.1" = "black",
287                               "beta.1" = "darkgray",
288                               "beta.2" = "gray",
289                               "beta.3" = "lightgray")) +
290   theme_minimal() +
291   theme(
292     axis.text.x = element_text(angle = 0, hjust = 0.5),
293     panel.grid = element_blank(),
294   )
295 ggplot_ca.seco
296 ggsave(ggplot_ca.seco, dpi = 300, filename = "res/fig-ca.seco.png")
297
298 # ca.chuva (D)
299
300 ggplot_ca.chuva <- ggplot(diversity_data, aes(fill = Component, y = Percentage, x = Type)) +
301   geom_bar(position = "stack", stat = "identity") +
302   labs(title = "(D)",
303        y = "% of the total diversity",
304        x = "") +
305   scale_fill_manual(values = c("alpha.1" = "black",
306                               "beta.1" = "darkgray",
307                               "beta.2" = "gray",
308                               "beta.3" = "lightgray")) +

```

```

309   theme_minimal() +
310   theme(
311     axis.text.x = element_text(angle = 0, hjust = 0.5),
312     panel.grid = element_blank(),
313   )
314 ggplot_ca.chuva
315 ggsave(ggplot_ca.chuva, dpi = 300, filename = "res/fig-ca.chuva.png")
316
317 # PAINEL DE GRÁFICOS
318
319 library(gridExtra)
320
321 # 1. Organizar os gráficos em 2 linhas e 2 colunas
322
323 painel_completo <- grid.arrange(
324   ggplot_bb.seco, ggplot_bb.chuva,
325   ggplot_ca.seco, ggplot_ca.chuva,
326   ncol = 2, nrow = 2
327 )
328
329 # 2. Salvar o painel em alta resolução (300 DPI)
330 ggsave(
331   filename = "res/fig-painel_partição.png",
332   plot = painel_completo,
333   width = 10,
334   height = 8,
335   dpi = 300
336 )

```

1 #TABELA DE PARTIÇÃO----

2 #Feita no Gemini, conferir

3

4 # -----

5 # 1. FUNÇÃO DE EXTRAÇÃO

6 # -----

```

7 extrair_linha_particao <- function(part_obj, sistema, periodo) {
8
9   mapa_componentes <- c(
10     "alpha.1" = " 1 (micro-habitat)",
11     "alpha.2" = " 2 (ponto)",
12     "alpha.3" = " 3 (riacho)",
13     "beta.1"  = " 1 (entre micro-habitats)",
14     "beta.2"  = " 2 (entre pontos)",

```

```

15     "beta.3" = " 3 (entre riachos)",
16     "gamma"  = "  (Riqueza total)"
17 )
18
19 stat <- part_obj$statistic
20 comp_names <- names(stat)
21
22 df <- data.frame(
23     Sistema      = sistema,
24     Período      = período,
25     Componente   = ifelse(comp_names %in% names(mapa_componentes),
26                           mapa_componentes[comp_names], comp_names),
27     Observado    = sprintf("%.2f%%", stat * 100),
28     Esperado     = sprintf("%.2f%%", part_obj$oecosimu$means * 100),
29     SES          = sprintf("%.2f", part_obj$oecosimu$z),
30     p_val        = part_obj$oecosimu$pval,
31     stringsAsFactors = FALSE
32 )
33
34 df$p_valor <- ifelse(df$p_val <= 0.001, "< 0.001*",
35                     ifelse(df$p_val <= 0.05, sprintf("%.3f*", df$p_val),
36                           sprintf("%.3f", df$p_val)))
37
38 return(df %>% select(-p_val))
39 }
40
41 # -----
42 # 2. PROCESSAMENTO E RUNDOWN DOS OBJETOS ADIPART
43 # -----
44 # Rodando o adipart para cada cenário e gerando os objetos corretos
45
46 p_ca_seco <- adipart(y = m_cont_ca.seco, x = ca.seco, index = "richness", relative = TRUE)
47 p_ca_chuva <- adipart(y = m_cont_ca.chuva, x = ca.chuva, index = "richness", relative = TRUE)
48
49 p_bb_seco <- adipart(y = m_cont_bb.seco, x = bb.seco, index = "richness", relative = TRUE)
50 p_bb_chuva <- adipart(y = m_cont_bb.chuva, x = bb.chuva, index = "richness", relative = TRUE)
51
52 p_pt_seco <- adipart(y = m_cont_s, x = pt.seco, index = "richness", relative = TRUE)
53 p_pt_chuva <- adipart(y = m_cont_c, x = pt.chuva, index = "richness", relative = TRUE)
54
55 # -----
56 # 3. CONSOLIDAÇÃO DA TABELA 1
57 # -----

```

```

58 tabela_1_raw <- rbind(
59   extrair_linha_particao(p_ca_seco, "Riacho Caiana (CA)", "Seco"),
60   extrair_linha_particao(p_ca_chuva, "Riacho Caiana (CA)", "Chuvoso"),
61   extrair_linha_particao(p_bb_seco, "Riacho Barro Branco (BB)", "Seco"),
62   extrair_linha_particao(p_bb_chuva, "Riacho Barro Branco (BB)", "Chuvoso"),
63   extrair_linha_particao(p_pt_seco, "Partição Total (PT)", "Seco"),
64   extrair_linha_particao(p_pt_chuva, "Partição Total (PT)", "Chuvoso")
65 )
66
67 # -----
68 # 4. IMPRESSÃO DA TABELA GT
69 # -----
70 tabela_1_final <- tabela_1_raw %>%
71   gt(groupname_col = "Sistema") %>%
72   tab_header(
73     title = md("**Tabela 1.** Partição aditiva da riqueza de espécies zooplanctônicas nas di.
74   ) %>%
75   cols_label(
76     Período = "Período",
77     Componente = "Componente Hierárquico",
78     Observado = "Observado (%)",
79     Esperado = "Esperado (%)",
80     SES = "SES",
81     p_valor = "p-valor"
82   ) %>%
83   cols_align(
84     align = "center",
85     columns = c("Período", "Observado", "Esperado", "SES", "p_valor")
86   )
87 tabela_1_final
88
89 # Formatar nomes das colunas diretamente
90 tabela_1_simples <- tabela_1_raw
91 colnames(tabela_1_simples) <- c("Sistema", "Período", "Componente", "Obs (%)", "Esp (%)", "SES", "p-valor")
92
93 # Imprimir no console
94 print(tabela_1_simples, row.names = FALSE)
95
96 # Exportando dados para Excel
97 library(openxlsx)
98 #write.xlsx(tabela_1_simples, file = "res/tabela de partição.xlsx", rowNames = FALSE)
99 wb <- loadWorkbook("res/tabela de partição.xlsx")
100 writeData(wb, sheet = "Sheet 1", x = tabela_1_simples)

```

```

101 saveWorkbook(wb, "res/tabela de partição.xlsx", overwrite = TRUE)
102
103 # Tabela final ajustada no Excel
104 library(readxl)
105 part_gtttable <- read_excel(
106   path = "res/tabela de partição.xlsx",
107   sheet = "tabela_final",
108   range = "A1:G35") #ajustar para o tamanho da tabela final
109 part_gtttable
110 library(gt)
111 part_gtttable <- gt(part_gtttable)
112 part_gtttable <- sub_missing(part_gtttable,
113                               columns = everything(), missing_text = "")
114 part_gtttable
115 gtsave(part_gtttable, "res/gt-part_gtttable.xlsx.html")
116 saveRDS(part_gtttable, "res/gt-part_gtttable.xlsx.rds")

```

4 Discussion

Conclusions

Acknowledgements

We are greatly indebted to the staff of the REBIO Guaribas for field work assistance. LCA is grateful to Programa de Pós-Graduação em Ecologia e Conservação (PPGEC-UEPB) and CAPES for scholarship granted. EGS, MMSM and MVBS are grateful to Programa de Iniciação Científica (PIBIC-UEPB) and CNPq for scholarship granted.

Authorship contribution statement

Authorship of this paper is based on CRediT (2026).

Ellen Gomes da Silva: Data curation, Formal analysis, Investigation, Software, Visualization, Writing – original draft.

Mayara Mirelly da Silva Monteiro: Data curation, Formal analysis, Investigation.

Ludmilla Cavalcanti Antunes Lucena: Supervision, Validation.

Thais Xavier de Melo: Supervision, Validation.

Elvio Sergio Figueredo Medeiros: Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing.

Ethical approval

Activities inside the REBIO Guaribas were authorized by the Instituto Chico Mendes de Conservação da Biodiversidade (license no. SISBIO-89415-1, SISBIO-89429-2 and SISBIO-89415-2).

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this manuscript, the authors used ChatGPT (OpenAI) to assist with English language editing and improve the clarity and fluency of the writing. No AI was used for data analysis, interpretation, or generation of scientific conclusions. The authors carefully reviewed and revised all AI-generated suggestions and take full responsibility for the accuracy and content of the final manuscript (CNPq, 2026).

References

- Anderson, M. J., & Walsh, D. C. I. (2013). PERMANOVA, ANOSIM, and the mantel test in the face of heterogeneous dispersions: What null hypothesis are you testing? *Ecological Monographs*, 83(4), 557–574. <https://doi.org/10.1890/12-2010.1>
- APHA. (1998). *Standard methods for the examination of water and wastewater*. <https://www.standardmethods.org/>
- Arruda, D. B., Cunha, B. P., & Rêgo, K. M. C. (2013). Conflitos entre REBIO Guaribas e comunidades locais: (in)justiça ambiental e ecologia política. *Revista Direitos Emergentes na Sociedade Global*, 2(2), 280–304. <https://doi.org/10.5902/2316305410852>
- Ávila, A. C., Stenert, C., & Maltchik, L. (2011). Partitioning macroinvertebrate diversity across different spatial scales in southern brazil coastal wetlands. *Wetlands*, 31(3), 459–469. <https://doi.org/10.1007/s13157-011-0178-3>
- Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, 19(1), 134–143. <https://doi.org/10.1111/j.1466-8238.2009.00490.x>
- Baselga, A. (2012). The relationship between species replacement, dissimilarity derived from nestedness, and nestedness. *Global Ecology and Biogeography*, 21(12), 1223–1232. <https://doi.org/10.1111/j.1466-8238.2011.00756.x>

- Belbin, L., & McDonald, C. (1993). Comparing three classification strategies for use in ecology. *Journal of Vegetation Science*, 4(3), 341–348. <https://doi.org/10.2307/3235592>
- Brasil. (2000). *Lei no 9.985, de 18 de julho de 2000. Regulamenta o art. 225, § 1o, incisos i, II, III e VII da constituição federal, institui o sistema nacional de unidades de conservação da natureza e dá outras providências.* https://www.planalto.gov.br/ccivil_03/leis/19985.htm
- CNPq. (2026). *Portaria 2664/2026 de 6 de março de 2026 - política de integridade na atividade científica do CNPq.* http://memoria2.cnpq.br/web/guest/view/-/journal_content/56_INSTANCE_0oED/10157/23142775
- CReditT. (2026). *CReditT - contributor role taxonomy.* <https://credit.niso.org/>
- Crist, Thomas O., Veech, Joseph A., Gering, Jon C., & Summerville, Keith S. (2003). Partitioning species diversity across landscapes and regions: A hierarchical analysis of , , and diversity. *The American Naturalist*, 162(6), 734–743. <https://doi.org/10.1086/378901>
- Diniz-Reis, T. R., Augusto, F. G., Abdalla Filho, A. L., Araújo, M. G. D. S., Chaves, S. S. F., Almeida, R. F., Perez, E. B., Simon, C. D. P., De Souza, J., Da Costa, C. F. G., Gomes, T., Martinez, M. G., Soltangheisi, A., Mariano, E., Vanin, A. S., Andrade, T., Boesing, A., Costa, F. J. V., Fortuna, M. D., ... Martinelli, L. A. (2022). SIA-BRA: A database of animal stable carbon and nitrogen isotope ratios of brazil. *Global Ecology and Biogeography*, 31(4), 611–620. <https://doi.org/10.1111/geb.13449>
- EPA. (2020). *Standard operating procedure for zooplankton sample collection and preservation.* https://www.epa.gov/system/files/documents/2021-08/lg402a.v15-zooplankton-sampling_rfa.pdf
- Farias, R. L., Stenert, C., Maltchik, L., & Medeiros, E. S. F. (2020). Partitioning of macroinvertebrate assemblages across temporary pools in an intermittent dryland river. *Inland Waters*, 10(4), 480–492. <https://doi.org/10.1080/20442041.2020.1738841>
- Farris, J. S. (1969). On the cophenetic correlation coefficient. *Systematic Zoology*, 18(3), 279. <https://doi.org/10.2307/2412324>
- Freiry, R. F., Weber, V., Bonecker, C. C., Lansac-Tôha, F. A., Pires, M. M., Stenert, C., & Maltchik, L. (2020). Additive partitioning of the diversity of the dormant zooplankton communities in intermittent ponds along a forest–grassland transition. *Hydrobiologia*, 847(5), 1327–1342. <https://doi.org/10.1007/s10750-020-04187-0>
- Frissell, C. A., Liss, W. J., Warren, C. E., & Hurley, M. D. (1986). A hierarchical framework for stream habitat classification: Viewing streams in a watershed context. *Environmental Management*, 10, 199–214. <https://doi.org/10.1007/BF01867358>
- Gering, J. C., Crist, T. O., & Veech, J. A. (2003). Additive partitioning of species diversity across multiple spatial scales: Implications for regional conservation of biodiversity. *Conservation Biology*, 17(2), 488–499. <https://doi.org/10.1046/j.1523-1739.2003.01465.x>
- Gouveia, R. S. D., Lira, G. L. D. A., Ramos, T. P. A., & Medeiros, E. S. F. (2017). Ichthyofauna of the reserva biológica guaribas and surrounding areas, state of paraíba, brazil. *Check List*, 13(5), 581–590. <https://doi.org/10.15560/13.5.581>
- Haney, J. F., & Hall, D. J. (1973). Sugar-coated *Daphnia* : A preservation technique for Cladocera1. *Limnology and Oceanography*, 18(2), 331–333. <https://doi.org/10.4319/lo.1973.18.2.0331>
- Heinrichs, R., Otto, R., Magalhães, A., & Meirelles, G. C. (2017). *Importance of sugarcane in*

- brazilian and world bioeconomy* (S. Dabbert, I. Lewandowski, J. Weiss, & A. Pyka, Eds.; pp. 205–217). Springer International Publishing. https://doi.org/10.1007/978-3-319-58374-7_11
- Kaufman, L., & Rousseeuw, P. J. (1990). *Agglomerative nesting (program AGNES)* (1st ed., pp. 199–252). Wiley. <https://doi.org/10.1002/9780470316801.ch5>
- Lande, R. (1996). Statistics and partitioning of species diversity, and similarity among multiple communities. *Oikos*, 76(1), 5. <https://doi.org/10.2307/3545743>
- Lê, S., Josse, J., & Husson, F. (2008). FactoMineR: An r package for multivariate analysis. *Journal of Statistical Software*, 25(1), 1–18.
- Legendre, P., & Legendre, L. (1998). *Numerical ecology* (2nd English ed).
- Ligeiro, R., Melo, A. S., & Callisto, M. (2010). Spatial scale and the diversity of macroinvertebrates in a neotropical catchment. *Freshwater Biology*, 2010(55), 424–435. <https://doi.org/10.1111/j.1365-2427.2009.02291.x>
- Lucena, L. C. A., Melo, T. X., & Medeiros, E. S. F. (2015). Zooplankton community of parnaíba river, northeastern brazil. *Acta Limnologica Brasiliensia*, 27(1), 118–129. <https://doi.org/10.1590/S2179-975X3214>
- Mackereth, F. J. H., Heron, J., & Talling, J. F. (1978). Water analysis: Some revised methods for limnologists.(scientific publication n. 36) freshwater biological association. *Cumbria and Dorset, England*.
- Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., & Hornik, K. (2026). *Cluster: Cluster analysis basics and extensions*. <https://CRAN.R-project.org/package=cluster>
- Maitland, P. S. (1990). *Biology of fresh waters* (1st ed.). Springer Netherlands. <https://doi.org/10.1007/978-94-011-7852-5>
- McCune, B., & Grace, J. B. (2002). *Analysis of ecological communities*. MjM Software Design.
- Medeiros, E. S. F., Silva, M. J., & Ramos, R. T. C. (2008). Application of catchment- and local-scale variables for aquatic habitat characterization and assessment in the Brazilian semi-arid region. *Neotropical Biology and Conservation*, 3, 13–20. <https://doi.org/10.4013/5440>
- Mehla, M. K., Jeong, J., Cho, G., Wolfe, J. E., & Kan, E. (2026). How agricultural field management modulates downstream transport of antibiotics, bacteria and nutrients in runoff water using edge-of-field monitoring. *Water Research*, 307, 126704. <https://doi.org/10.1016/j.watres.2026.126704>
- Melo, T. X. D., Lourenço, L. D. J. S., & Medeiros, E. S. F. (2014). Checklist of zooplankton from the upper ipanema river (pernambuco), an intermittent river in semi-arid brazil. *Check List*, 10(3), 524–528. <https://doi.org/10.15560/10.3.524>
- Melo, T. X., & Medeiros, E. S. F. (2013). Spatial distribution of zooplankton diversity across temporary pools in a semiarid intermittent river. *International Journal of Biodiversity*, 2013, 1–13. <https://doi.org/10.1155/2013/946361>
- MMA. (2003). *Plano de Manejo da Reserva Biológica Guaribas*.
- MMA. (2026). *Plano de manejo integrado do fogo 2026-2030. Reserva biológica guaribas* (p. 56). https://www.gov.br/icmbio/pt-br/centrais-de-conteudo/publicacoes/planos-de-manejo-integrado-do-fogo/Plano_de_Manejo_Integrado_do_Fogo_da_Reserva_Biologica_Guaribas.pdf
- Mugodo, J., Kennard, M. J., Liston, P., Nichols, S., Linke, S., Norris, R. H., & Lintermans,

- M. (2006). Local stream habitat variables predicted from catchment scale characteristics are useful for predicting fish distribution. *Hydrobiologia*, 572(1), 59–70. <https://doi.org/10.1007/s10750-006-0252-7>
- Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Borman, T., Carvalho, G., Chirico, M., Caceres, M., ... Weedon, J. (n.d.). *Vegan: Community ecology package*. <https://doi.org/10.32614/CRAN.package.vegan>
- Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007). Updated world map of the köppen-geiger climate classification. *Hydrology and Earth System Sciences*, 11(5), 1633–1644. <https://doi.org/10.5194/hess-11-1633-2007>
- Pereira, K. M. G., Cordeiro, N. G., Suzuki, L. P. Z. L., & Berg, E. van den. (2026). How important are small fragments for restoration in the central portion of the atlantic forest? *Biological Conservation*, 321, Article 111958. <https://doi.org/https://doi.org/10.1016/j.biocon.2026.111958>
- Pusey, B., Kennard, M. J., & Arthington, A. (2004). *Study area, data collection, analysis and presentation* (B. Pusey, M. J. Kennard, & A. Arthington, Eds.; pp. 26–48). CSIRO Publishing.
- R Core Team. (2017). *R: A language and environment for statistical computing*. <https://www.R-project.org/>
- Rocha, L. G., Medeiros, E. S. F., & Andrade, H. T. A. (2012). Influence of flow variability on macroinvertebrate assemblages in an intermittent stream of semi-arid brazil. *Journal of Arid Environments*, 85, 33–40. <https://doi.org/10.1016/j.jaridenv.2012.04.001>
- Saraçlı, S., Doğan, N., & Doğan, İ. (2013). Comparison of hierarchical cluster analysis methods by cophenetic correlation. *Journal of Inequalities and Applications*, 2013(1), 203. <https://doi.org/10.1186/1029-242X-2013-203>
- Shell, T. M., & Collier, K. J. (2018). Partitioning of macroinvertebrate communities in a large new zealand river highlights the role of multiple shore-zone habitat types. *River Research and Applications*, 34(8), 993–1002. <https://doi.org/10.1002/rra.3344>
- Silva, A. R., & Dias, C. T. S. (2013). A cophenetic correlation coefficient for tocher's method. *Pesquisa Agropecuária Brasileira*, 48(6), 589–596. <https://doi.org/10.1590/S0100-204X2013000600003>
- Silva, J. M. C., & Casteleti, C. H. M. (2003). *Status of the biodiversity of the atlantic forest of brazil* (C. Galindo-Leal & I. G. Câmara, Eds.; pp. 43–59). CABS; Island Press.
- Sneath, P. H. A., & Sokal, R. R. (1973). *Numerical taxonomy: The principles and practice of numerical classification* (1st ed.). W H Freeman & Co.
- Sokal, R. R., & Rohlf, F. J. (1995). *Biometry: The principles and practice of statistics in biological research* (3rd ed.). W.H. Freeman; Company.
- Sokal, R. R., & Rohlf, F. J. (1962). The comparison of dendrograms by objective methods. *Taxon*, 11(2), 33–40. <https://doi.org/10.2307/1217208>
- Strahler, A. N. (1957). Quantitative analysis of watershed geomorphology. *Eos, Transactions American Geophysical Union*, 38(6), 913–920. <https://doi.org/10.1029/TR038i006p00913>
- Veech, J. A., Crist, T. O., & Summerville, K. S. (2003). Intraspecific aggregation decreases local species diversity of arthropods. *Ecology*, 84(12), 3376–3383. <https://doi.org/10.1890/02->

3127

Veech, J. A., Summerville, K. S., Crist, T. O., & Gering, J. C. (2002). The additive partitioning of species diversity: Recent revival of an old idea. *Oikos*, 99(1), 3–9. <https://doi.org/10.1034/j.1600-0706.2002.990101.x>

Figures and Tables

Figure 1: Location of sampling sites along the Barro Branco Stream (Mamanguape, PB) (Site 1 06°43'06"S and 35°10'54"W; Site 2 06°42'38"S and 35°10'38"W; Site 3 06°43'06"S and 35°10'54"W; Site 4 06°41'46"S and 35°10'36"W; Site 5 06°40'56"S and 35°10'27"; Site 6 06°40'18"S and 35°10'34"W).

Figure 2: Principal Component Analysis (PCA) biplot of the sampling units (SUs) and environmental variables in the Barro Branco and Caiana streams during the study period, categorized by site type. Points represent the SUs, whereas vectors represent the environmental variables associated with habitat structure. Ellipses represent 95% confidence regions for the distribution of the groups of points, calculated based on the t-distribution.

Figure 3: Principal Component Analysis (PCA) biplot of the sampling units (SUs) and environmental variables in the Barro Branco and Caiana streams during the study period, categorized by stream. Points represent the SUs, whereas vectors represent the environmental variables associated with habitat structure. Ellipses represent 95% confidence regions for the distribution of the groups of points, calculated based on the t-distribution.

Figure 4: Principal Component Analysis (PCA) biplot of the sampling units (SUs) and environmental variables in the Barro Branco and Caiana streams during the study period, categorized by season. Points represent the SUs, whereas vectors represent the environmental variables associated with habitat structure. Ellipses represent 95% confidence regions for the distribution of the groups of points, calculated based on the t-distribution.

Figure 5: Individual-based extrapolation and rarefaction curves for the richness of resting stages of zooplanktonic taxa in each study stream. Shaded areas indicate 95% confidence intervals.

Figure 6: Classification analysis (UPGMA using Bray-Curtis dissimilarity) of wetland macroinvertebrate samples obtained in a. permanent and b. intermitent wetlands. The first number of the sample label indicates the wetland. The second number indicates the habitat type (1=hydrophytes; 2=reed-like emergent plants; and 3=leafy emergent plants). The third number indicates the individual sweeps realized in each habitat (1–3)UPGMA Hierarchical flexible beta clustering ($\alpha = -0.25$) based on Bray-Curtis distance of (a) macroinvertebrates and (b) Chironomidae in the Paraíba River during the 2015 wet season. Tag codes indicate different sections of the river: U = upper segment, M = middle segment, and T = Taperoá River segment. R = reaches (1–3), S = sites (1–3), and Hab = habitat types (1–3). Percent chaining <4%.

Figure 7: Observed and expected diversity, partitioned into one alpha and three beta components, expressed as percent of total species richness. Numbers indicate the proportion of randomized samples containing more species than the observed sample for each partition. 1: subsample species richness, 1: beta diversity among sub-samples; 2: beta diversity among pools or habitats within pool, and 3: beta diversity among river reaches.

Table 1: Environmental variables averaged for sampling sites (minimum-maximum) for the Barro Branco stream (Mamanguape, PB).

Table 2: Nutrient concentration (g L ⁻¹) averaged for sampling sites (minimum-maximum) for the Barro Branco stream (Mamanguape, PB).

Table 3: Density (ind/m²) and frequency of occurrence (FO%) of zooplankton collected during the 2024 hydrological cycle.

Table 4: Contribution (expressed in percentage values) of each spatial scale assessed to the total diversity (richness) of the dormant stages of the zooplankton. Numbers in ‘P value’ column indicate the proportion of randomized samples containing more species than the observed sample for each partition.

Figures

Figure 1




Figure 1

Figure 2




Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Tables

Table 1

Table 2

Table 3

Table 4




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

Table 1: Environment data GT table.

Variable	Ponto5	Ponto6	Ponto7	Ponto8
w.Cond.uS.cm	137.5 (119.2-164.4)	148 (134.1-156.8)	137.5 (98.3-160.2)	154.2 (141.6-161.8)
w.Turb.ntu	5 (2-10.7)	4.9 (2.1-14.1)	4.9 (1.5-9.2)	5.7 (4-6.8)
w.pH	5.9 (4.7-7.5)	5.7 (4.6-7.7)	5.1 (4.5-6)	5.5 (5-6)
m.Elev.m	107.1 (107.1-107.1)	112.8 (112.8-112.8)	89 (89-89)	89.1 (89.1-89.1)
m.Vel.m.s	0.073 (0.025-0.161)	0.086 (0.035-0.24)	0.135 (0.022-0.207)	0.036 (0-0.115)
m.Slope	1.4 (1-2)	2.7 (2.5-3)	2.4 (1.5-3)	1.5 (1-2)
m.Depth.avg.cm	12.8 (9-14)	23.7 (16.3-31.7)	23.8 (16-41)	43.3 (28.3-51.3)
m.Depth.mar.cm	11.8 (3-16)	21.7 (8-30)	18.8 (7-42)	48.3 (16-87)
s.Mud	100 (100-100)	100 (100-100)	95 (90-100)	74.2 (5-100)
s.Sand	0 (0-0)	0 (0-0)	5 (0-10)	25.8 (0-95)
s.Gravel.f	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
s.Rocks	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
h.Macrophytes	5.8 (5-10)	10 (0-30)	0 (0-0)	17.9 (0-60)
h.Grass	3.3 (0-10)	8.3 (5-15)	6 (0-15)	27.1 (15-50)
h.Subm.veg	7.5 (5-15)	8.3 (5-15)	5.8 (0-10)	5.8 (0-10)
h.Overh.veg	68.3 (40-90)	70.8 (50-85)	76.7 (60-90)	31.7 (25-50)
h.Leaf.l	7.5 (5-10)	11 (1-25)	15.8 (5-30)	18.3 (10-30)
h.Root.masses	5 (0-15)	10.8 (5-15)	12.5 (5-20)	4.2 (0-10)
h.Algae	2.5 (0-5)	0.2 (0-1)	5.5 (0-21)	19.2 (15-25)
h.Debris	30.8 (20-45)	12.7 (5-20)	17.7 (5-30)	21.7 (5-35)

Table 2: Nutrient concentrations at the sampling sites.

Variable	Ponto5	Ponto6	Ponto7	Ponto8
n.Ammonia.NH3.ug.L	54.8 (39.3-64.3)	58.9 (47.5-77)	72 (33.4-115.7)	81.6 (68.9-93)
n.Nitrite.NO2.ug.L	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
n.Nitrate.NO3.ug.L	6 (0-9.7)	8.8 (6.9-12.6)	5.5 (0-12.6)	18.8 (11.1-31.1)
n.Orthophosphate.SRP.ug.L	0 (0-0)	0 (0-0)	1 (0-3)	0 (0-0)
n.Phosphorus.TP.ug.L	0 (0-0)	4.6 (0.3-10.3)	50.1 (0-150.3)	9 (0-27)

Table 3: Density (ind/m²) taxa GT table.

Stream	Barro Branco Stream	...3	...4	...5	...6	...7	...8
Season	Wet						Dr
Site	Ponto5	Ponto6	Ponto7	Ponto8	Ponto9	Ponto10	Po
Alo.dadayi	0 (0)	0 (0)	0 (0)	5.3 (6.6)	0 (0)	0.4 (0.3)	0 (0)
Alo.hamulata	0 (0)	0 (0)	0 (0)	1.1 (1.3)	0 (0)	0.1 (0.1)	0 (0)
Alonella.sp	0 (0)	0 (0)	0 (0)	0.4 (0.5)	0 (0)	0 (0.1)	0 (0)
Bdelloidea	1.7 (0.5)	1.3 (0)	1.5 (0.5)	3.4 (0.8)	2 (0.8)	2.9 (1.3)	0.6
Bra.patulus	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Cephalodella.sp	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0.1 (0.1)	0 (0)
Colurella.sp	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Copepodite	0.5 (0.4)	0.1 (0)	0.1 (0)	12.1 (14.5)	0 (0)	0.5 (0.8)	0 (0)
Chydorus.sp	0 (0)	0 (0)	0 (0)	3.2 (3.5)	0 (0)	0.2 (0.1)	0 (0)
Dicranophoroides.sp	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Dia.birgei	0 (0)	0 (0)	0 (0)	0.1 (0.1)	0 (0)	0.1 (0.1)	0 (0)
Dia.spinulosum	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Dis.aculeata	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Euc.dilatata	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Fil.longiseta	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Ily.spinifer	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0.1 (0.1)	0 (0)
Lec.aculeata	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.bulla	0 (0)	0 (0)	0 (0)	0 (0.1)	0 (0)	0 (0)	0 (0)
Lec.cornuta	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.crepida	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.curvicornis	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.furcata	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lecane.sp1	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.leontina	0 (0)	0 (0)	0 (0)	0.5 (0.5)	0 (0)	0 (0.1)	0 (0)
Lec.lunaris	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.luna	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.monostyla	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lecane.sp2	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.papuana	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lec.quadridentata	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lecane.sp3	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Leydigia.sp	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lepadella.sp1	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lepadella.sp2	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lep.ovalis	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Lep.patella	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Mesocyclops.sp	0 (0)	0 (0)	0 (0)	0.6 (0.6)	0 (0)	0 (0)	0 (0)
Macrocylops.sp	0 (0)	0 (0)	0 (0)	0.2 (0.2)	0 (0)	0 (0)	0 (0)
Macrochaetus.sp	0 (0)	0 (0)	0 (0)	0.2 (0.3)	0 (0)	0 (0)	0 (0)
Macrothrix.sp	0 (0)	75 0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Mac.mira	0 (0)	0 (0)	0 (0)	0.2 (0.3)	0 (0)	0 (0)	0 (0)
Mac.laticornis	0 (0)	0 (0)	0 (0)	0.1 (0.1)	0 (0)	0 (0)	0 (0)
Moi.minuta	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Myt.crassipes	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Mytilina.sp	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Myt.bisulcata	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
N.elli	0.2 (0.2)	0.1 (0.1)	0.1 (0)	2.2 (2.6)	0 (0.1)	0.2 (1.2)	0 (0)

Table 4: Partition data GT table.

Sistema	Período	Componente	Obs (%)	Esp (%)	SES	p-valor
Riacho Caiana (CA)	Seco	1 (micro-habitat)	16.40%	21.17%	-12.27	< 0.00
Riacho Caiana (CA)	Seco	2 (ponto)	28.84%	37.52%	-9.32	< 0.00
Riacho Caiana (CA)	Seco	(Riqueza total)	100.00%	100.00%	0.00	1.000
Riacho Caiana (CA)	Seco	1 (entre micro-habitats)	12.43%	16.35%	-4.84	< 0.00
Riacho Caiana (CA)	Seco	2 (entre pontos)	71.16%	62.48%	9.32	< 0.00
Riacho Caiana (CA)	Chuvoso	1 (micro-habitat)	16.40%	21.18%	-12.12	< 0.00
Riacho Caiana (CA)	Chuvoso	2 (ponto)	28.84%	37.52%	-9.29	< 0.00
Riacho Caiana (CA)	Chuvoso	(Riqueza total)	100.00%	100.00%	0.00	1.000
Riacho Caiana (CA)	Chuvoso	1 (entre micro-habitats)	12.43%	16.33%	-4.70	< 0.00
Riacho Caiana (CA)	Chuvoso	2 (entre pontos)	71.16%	62.48%	9.29	< 0.00
Riacho Barro Branco (BB)	Seco	1 (micro-habitat)	20.23%	26.63%	-11.64	< 0.00
Riacho Barro Branco (BB)	Seco	2 (ponto)	35.04%	46.86%	-9.03	< 0.00
Riacho Barro Branco (BB)	Seco	(Riqueza total)	100.00%	100.00%	0.00	1.000
Riacho Barro Branco (BB)	Seco	1 (entre micro-habitats)	14.81%	20.23%	-4.66	< 0.00
Riacho Barro Branco (BB)	Seco	2 (entre pontos)	64.96%	53.14%	9.03	< 0.00
Riacho Barro Branco (BB)	Chuvoso	1 (micro-habitat)	20.23%	26.64%	-11.86	< 0.00
Riacho Barro Branco (BB)	Chuvoso	2 (ponto)	35.04%	46.88%	-9.39	< 0.00
Riacho Barro Branco (BB)	Chuvoso	(Riqueza total)	100.00%	100.00%	0.00	1.000
Riacho Barro Branco (BB)	Chuvoso	1 (entre micro-habitats)	14.81%	20.24%	-4.88	< 0.00
Riacho Barro Branco (BB)	Chuvoso	2 (entre pontos)	64.96%	53.12%	9.39	< 0.00
Partição Total (PT)	Seco	1 (micro-habitat)	12.45%	19.41%	-21.58	< 0.00
Partição Total (PT)	Seco	2 (ponto)	22.18%	34.41%	-16.50	< 0.00
Partição Total (PT)	Seco	3 (riacho)	73.28%	84.12%	-5.60	< 0.00
Partição Total (PT)	Seco	(Riqueza total)	100.00%	100.00%	0.00	1.000
Partição Total (PT)	Seco	1 (entre micro-habitats)	9.73%	15.00%	-8.68	< 0.00
Partição Total (PT)	Seco	2 (entre pontos)	51.09%	49.72%	0.69	0.499
Partição Total (PT)	Seco	3 (entre riachos)	26.72%	15.88%	5.60	< 0.00
Partição Total (PT)	Chuvoso	1 (micro-habitat)	23.72%	51.72%	-45.73	< 0.00
Partição Total (PT)	Chuvoso	2 (ponto)	35.50%	69.45%	-35.41	< 0.00
Partição Total (PT)	Chuvoso	3 (riacho)	83.75%	96.48%	-9.42	< 0.00
Partição Total (PT)	Chuvoso	(Riqueza total)	100.00%	100.00%	0.00	1.000
Partição Total (PT)	Chuvoso	1 (entre micro-habitats)	11.78%	17.73%	-7.46	< 0.00
Partição Total (PT)	Chuvoso	2 (entre pontos)	48.25%	27.04%	13.64	< 0.00
Partição Total (PT)	Chuvoso	3 (entre riachos)	16.25%	3.52%	9.42	< 0.00

Apendices

Non-used figures and tables

 Non-used codes

1 `table(1:10)`