

Master Program in *Data Science and Business Informatics*

Statistics for Data Science

Lesson 01 - Probabilities and independence

Salvatore Ruggieri


Department of Computer Science

University of Pisa, Italy

salvatore.ruggieri@unipi.it

Why Statistics in Data Science?

We need grounded means for reasoning about data generated from real world with some degree of randomness.



What will you learn?

- Probability: properties of data generated by a known/assumed randomness model
- Statistics: properties of a randomness model that could have generated given data
- The R programming language

Sample spaces and events

- An **experiment** is a measurement of a random process
- The **outcome** of a measurement takes values in some set Ω , called the **sample space**.

Examples:

- ▶ Tossing a coin: $\Omega = \{H, T\}$ ne kete slide ka shpjegu pak ne gjenarle termin e bashkesise dhe popullimin , bashkesite ku marrin vlerat ngjarjet, risultati eshte shuma e te gjitha munsive qe mund te dodhin **[Finite]**
- ▶ Month of birthdays $\Omega = \{\text{Jan}, \dots, \text{Dec}\}$ **[Finite]**
- ▶ Population of a city $\Omega = \mathbb{N} = \{0, 1, 2, \dots, \}$ **[Countably infinite]**
- ▶ Length of a street $\Omega = \mathbb{R}^+ = (0, \infty)$ **[Uncountably infinite]**
- ▶ Tossing a coin twice: $\Omega = \{H, T\} \times \{H, T\} = \{(H, H), (H, T), (T, H), (T, T)\}$
- ▶ Testing for Covid-19 (univariate): $\Omega = \{+, -\}$ in questo caso abbiamo i dati binari che possiamo dare solo due possibilita,
- ▶ Testing for Covid-19 (multivariate): $\Omega = \{f, m\} \times \mathbb{N} \times \{+, -\}$, e.g., $(f, 25, -) \in \Omega$

Look at seeing-theory.brown.edu

- An **event** is some subset of $A \subseteq \Omega$ of possible outcomes of an experiment.
 - ▶ $L = \{\text{Jan}, \text{March}, \text{May}, \text{July}, \text{August}, \text{October}, \text{December}\}$ *a long month with 31 days*
- We say that an event A **occurs** if the outcome of the experiment belongs to the set A .
 - ▶ If the outcome is Jan then L occurs

Probability functions on finite sample space

A **probability function** is a mapping from events to **real numbers** that satisfies certain axioms. *Intuition: how likely is an event to occur.*

ne kete rast kemi def te probabilitetit , qe shpjegohet , ne qoftese kemi nje hapsire mundeshish omega probabiliteti i merr vlerat nga [0:1]
dhe probabiliteti e bashkimit te dy bashkesive eshte sa shuma e probabiliteteve, ne qoftese a e b sono disjoint

DEFINITION. A *probability function* P on a finite sample space Ω assigns to each event A in Ω a number $P(A)$ in $[0,1]$ such that

- (i) $P(\Omega) = 1$, and
- (ii) $P(A \cup B) = P(A) + P(B)$ if A and B are disjoint.

The number $P(A)$ is called the probability that A occurs.

- Fact: $P(\{a_1, \dots, a_n\}) = P(\{a_1\}) + \dots + P(\{a_n\})$

[Generalized additivity]

- ▶ Assigning probability to a singleton is enough

- Examples:

in this example we have probability two possibility,

if the second one we have a probability of a month with a number of large day on the month. the sum of a probability is a sum of single one probability.

- ▶ $P(\{H\}) = P(\{T\}) = 1/2$
 - ▶ $P(\{\text{Jan}\}) = 31/365, P(\{\text{Feb}\}) = 28/365, \dots P(\{\text{Dec}\}) = 31/365$
 - ▶ $P(L) = 7/12$ or $31 \cdot 7/365$?

- $P(\{a\})$ often abbreviated as $P(a)$, e.g., $P(\text{Jan})$ instead of $P(\{\text{Jan}\})$

Properties of probability functions

- $P(A^c) = 1 - P(A)$

- $P(\emptyset) = 0$

in this slide we have a criterion of the probability, if probability of an event is zero, then the probability of the complement is one. if an event is empty the probability is zero, then the probability of the complement is one. if two events are disjoint the probability of their union is the sum of their probabilities. if two events are not disjoint the probability of their union is the sum of their probabilities minus the probability of their intersection.

[Impossible event]

- $A \subseteq B \Rightarrow P(A) \leq P(B)$

- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

[Inclusion-exclusion principle]

- $P(A \cup B) = P(A) + P(B \setminus A)$

- probability that at least one coin toss over two lands head?

Defining probability functions

Assigning probability is **NOT** an easy task: a prob. function can be an approximation of reality


- **Frequentist** interpretation: probability measures a “*proportion of outcomes*” .
 - ▶ A fair coin lands on heads 50% of times
 - ▶ $P(A) = |A|/\Omega$ [Counting]
 - ▶ $P(\{\text{at least one H in two coin tosses}\}) = |\{(H, H), (H, T), (T, H)\}|/4 = 3/4$
- **Bayesian** (or epistemological) interpretation: probability measures a “*degree of belief*” .
 - ▶ Iliad and Odissey were composed by the same person at 90%

Counting: The Monty Hall problem

<https://math.andyou.com/tools/montyhallsimulator/montysim.htm>

(See also Exercise 2.14 of textbook [T])

Tree-based sequential description of probability function



Probability functions on countably infinite sample space

DEFINITION. A *probability function* on an infinite (or finite) sample space Ω assigns to each event A in Ω a number $P(A)$ in $[0, 1]$ such that

- (i) $P(\Omega) = 1$, and
- (ii) $P(A_1 \cup A_2 \cup A_3 \cup \dots) = P(A_1) + P(A_2) + P(A_3) + \dots$
if A_1, A_2, A_3, \dots are disjoint events.

- (ii) is called **countable additivity**. It is equivalent to σ -additivity: for $A_1 \subseteq A_2 \subseteq \dots$

$$P\left(\lim_{n \rightarrow \infty} A_i\right) = \lim_{n \rightarrow \infty} P(A_i)$$

- Example

- ▶ Experiment: we toss a coin repeatedly until H turns up.
- ▶ Outcome: the number of tosses needed.
- ▶ $\Omega = \{1, 2, \dots\} = \mathbb{N}^+$
- ▶ Suppose: $P(H) = p$. Then: $P(n) = (1 - p)^{n-1}p$
- ▶ Is it a probability function? $P(\Omega) = \dots$

Conditional probability

- Long months and months with 'r'
 - ▶ $L = \{ \text{Jan, Mar, May, July, Aug, Oct, Dec} \}$ *a long month with 31 days*
 - ▶ $R = \{ \text{Jan, Feb, Mar, Apr, Sep, Oct, Nov, Dec} \}$ *a month with 'r'*
 - ▶ $P(L) = 7/12$ $P(R) = 8/12$
- Anna is born in a long month. What is the probability she is born in a month with 'r'?

$$P(R|L) = \frac{P(L \cap R)}{P(L)} = \frac{P(\{ \text{Jan, Mar, Oct, Dec} \})}{P(L)} = \frac{4/12}{7/12} = \frac{4}{7}$$

- **Intuition:** probability of an event in the restricted sample space $\Omega \cap L$
 - ▶ *a-priori* probability $P(R) = 8/12$
 - ▶ *a-posteriori* probability $P(R|L) = 4/7 < 8/12$
- **Example (classification):** probab. of Covid given gender=f and age ≥ 60 : $P(C|G \cap A)$
 - ▶ $\Omega = \{f, m\} \times \mathbb{N} \times \{+, -\}$
 - ▶ $C = \{(-, -, +) \in \Omega\}$ $G = \{(f, -, -) \in \Omega\}$ $A = \{(-, a, -) \in \Omega \mid a \geq 60\}$
 - ▶ naming triples with features (gender, age, covid): $P(\text{covid} = + | \text{gender} = f, \text{age} \geq 60)$

Another example at seeing-theory.brown.edu

Conditional probability

DEFINITION. The **conditional probability** of A given C is given by:

$$P(A|C) = \frac{P(A \cap C)}{P(C)},$$

provided $P(C) > 0$.

Properties:

- $P(A|C) \neq P(C|A)$, in general
- $P(\Omega|C) = 1$
- if $A \cap B = \emptyset$ then $P(A \cup B|C) = P(A|C) + P(B|C)$ $P(\cdot|C)$ is a probability function

THE MULTIPLICATION RULE. For any events A and C :

$$P(A \cap C) = P(A|C) \cdot P(C).$$


More generally, the **Chain Rule**:

$$P(A_1 \cap A_2 \cap A_3 \dots \cap A_n) = P(A_1) \cdot P(A_2|A_1) \cdot P(A_3|A_1 \cap A_2) \cdot \dots \cdot P(A_n|\cap_{i=1}^{n-1} A_i)$$

Example: no coincident birthdays

- $B_n = \{n \text{ different birthdays}\}$
- For $n = 1$, $P(B_1) = 1$
- For $n > 1$,

$$\begin{aligned}P(B_n) &= P(B_{n-1}) \cdot P(\{\text{the } n\text{-th person's birthday differs from the other } n-1\} | B_{n-1}) \\&= P(B_{n-1}) \cdot (1 - \frac{n-1}{365}) = \dots = \prod_{i=1}^{n-1} (1 - \frac{i}{365})\end{aligned}$$



Example: case-based reasoning

Factory 1's light bulbs work for over 5000 hours in 99% of cases.

Factory 2's bulbs work for over 5000 hours in 95% of cases.

Factory 1 supplies 60% of the total bulbs on the market and Factory 2 supplies 40% of it.

What is the chance that a purchased bulb will work for longer than 5000 hours?

- $A = \{\text{bulbs working for longer than 5000 hours}\}$
- $C = \{\text{bulbs made by Factory 1}\}$, hence $C^c = \{\text{bulbs made by Factory 2}\}$
- Since $A = (A \cap C) \cup (A \cap C^c)$ with $(A \cap C)$ and $(A \cap C^c)$ disjoint:

$$P(A) = P(A \cap C) + P(A \cap C^c)$$

- and then by the multiplication rule:

$$P(A) = P(A|C) \cdot P(C) + P(A|C^c) \cdot P(C^c)$$

Answer: $P(A) = 0.99 \cdot 0.6 + 0.95 \cdot 0.4 = 0.974$

The law of total probability

THE **LAW OF TOTAL PROBABILITY**. Suppose C_1, C_2, \dots, C_m are disjoint events such that $C_1 \cup C_2 \cup \dots \cup C_m = \Omega$. The probability of an arbitrary event A can be expressed as:

$$P(A) = P(A | C_1)P(C_1) + P(A | C_2)P(C_2) + \dots + P(A | C_m)P(C_m).$$

- **Intuition:** case-based reasoning




Fig. 3.2. The law of total probability (illustration for $m = 5$).

Exercise: Prisoners and guard dilemma

3 prisoners, 2 of which will be released.

$A1 = \{ \text{Prisoner 1 is released} \}$, $A2 = \{ \text{Prisoner 2 is released} \}$, $A3 = \{ \text{Prisoner 3 is released} \}$

$$P(A1^c \cap A2 \cap A3) = P(A1 \cap A2^c \cap A3) = P(A1 \cap A2 \cap A3^c) = 1/3$$

You are Prisoner 1:

- at your's present state of knowledge, the probability of being released is $2/3$
 - ▶ $P(A1) = P(A1 \cap A2^c \cap A3) + P(A1 \cap A2 \cap A3^c) = 2/3$
- if you ask a friendly guard to tell you who is the prisoner other than yourself that will be released, your probability of being released will become $1/2$
 - ▶ $P(A1|A2) = P(A1 \cap A2)/P(A2) = (1/3)/(2/3) = 1/2$

What is wrong with this line of reasoning?

- $P(A1) = P(A1|A2)P(A2) + P(A1|A3)P(A3) - P(A1|A2 \cap A3)P(A2 \cap A3) = 1/2 \cdot 2/3 + 1/2 \cdot 2/3 - 0 \cdot 1/3 = 2/3$

Independence of events

Intuition: whether one event provides any information about another.

Independence

An event A is independent of B , if $P(B) = 0$ or

$$P(A|B) = P(A)$$

- For $P(R|L) = 4/7 \neq 8/12 = P(R)$ - knowing Anna was born in a long month change the probability she was born in a month with 'r'!
- Tossing 2 coins:
 - ▶ A_1 is "H on toss 1" and A_2 is "H on toss 2"
 - ▶ $P(A_1) = P(A_2) = 1/2$
 - ▶ $P(A_2|A_1) = P(A_2 \cap A_1)/P(A_1) = 1/4/1/2 = 1/2 = P(A_2)$
- Physical and stochastic independence
- Properties:
 - ▶ A independent of B iff $P(A \cap B) = P(A) \cdot P(B)$
 - ▶ A independent of B iff B independent of A
 - ▶ A independent of B iff A^c independent of B

[Symmetry]

Conditional independence of events

Intuition: whether one event provides any information about another given a third event occurred. Technically, consider $P(\cdot|C)$ in independence.

Conditional independence

An event A is conditionally independent of B given C such that $P(C) > 0$, if $P(B|C) = 0$ or

$$P(A|B \cap C) = P(A|C)$$

- Properties:
 - ▶ A conditionally independent of B iff $P(A \cap B|C) = P(A|C) \cdot P(B|C)$
 - ▶ A conditionally independent of B iff B conditionally independent of A
- **Exercise at home.** Prove or disprove:
 - ▶ If A is independent of B then A is conditionally independent of B given C

[Symmetry]

Independence of two or more events

INDEPENDENCE OF TWO OR MORE EVENTS. Events A_1, A_2, \dots, A_m are called independent if

$$P(A_1 \cap A_2 \cap \dots \cap A_m) = P(A_1) P(A_2) \dots P(A_m)$$

and this statement *also* holds when any number of the events A_1, \dots, A_m are replaced by their complements throughout the formula.

Alternative definition

Events A_1, A_2, \dots, A_m are called independent if for every $J \subseteq \{1, \dots, m\}$:

$$P\left(\bigcap_{i \in J} A_i\right) = \prod_{i \in J} P(A_i)$$

- **Exercise at home:** show the two definitions are equivalent

Independence of two or more events

Alternative definition

Events A_1, A_2, \dots, A_m are called independent if for every $J \subseteq \{1, \dots, m\}$:

$$P\left(\bigcap_{i \in J} A_i\right) = \prod_{i \in J} P(A_i)$$

- It is **stronger** than **pairwise independence**

$$P(A_i \cap A_j) = P(A_i) \cdot P(A_j) \text{ for } i \neq j \in \{1, \dots, m\}$$

- Example: what is the probability of at least one head in the first 10 tosses of a coin?
 $A_i = \{\text{head in } i\text{-th toss}\}$

$$P\left(\bigcup_{i=1}^{10} A_i\right) = 1 - P\left(\bigcap_{i=1}^{10} A_i^c\right) = 1 - \prod_{i=1}^{10} P(A_i^c) = 1 - \prod_{i=1}^{10} (1 - P(A_i))$$