Category 3

Mild

- Stay at home
- Self care
- 111-Online if health needs

Category 2

Moderate

- Needs assessment (phone/video)
 Or
- "Hot Site" review for examination:

Hot Site: 07899 097 302

Category 1

Severe - needs admission:

- Clearly very unwell
- RR >25
- HR >131
- Sats <92%
- <u>Call 999</u> and advise them <u>suspected COVID</u>

Category 2A

Telephone assessment

- Completing full sentences
- No SOB
- No chest pain
- · Able to get out of bed
- · Able to do usual activities
- ROTH Score* >8s
 (*may underestimate severity)

Face-to face assessment (adults):

- RR 14-20
- HR 50-100
- Sats≥96%

Category 2B

Telephone assessment

- Completing full sentences
- Some SOB (new)
- No chest pain
- Able to do some activities but with some difficulty; Lethargic
- ROTH Score* 6-8s
 (*may underestimate severity)

Face-to face assessment (adults):

- RR 21-22
- HR 100-110
- Sats 94-96%

Category 2C

Telephone assessment

- Completing full sentences
- SOB on exertion (new)
- Mild chest tightness
- Able to do some activities but with some difficulty; lethargic
- ROTH Score* 6-8s
 (*may underestimate severity)

Face-to face assessment (adults):

- RR 23-24
- HR 110-130
- Sats 92-93%

Category 2A

Management:

- · Rest and make safe
- Oral fluids
- Paracetamol
- SAFETY-NETTING and consider AKI

Category 2B

Management:

- Rest and make safe
- Oral fluids
- Paracetamol

<u>Consider antibiotics to treat/prevent secondary bacterial pneumonia</u>:

- Doxycycline 200mg *stat*, 100mg *od*, 5 days in total

<u>OR</u>

- Amoxicillin 500mg *tds*, 5 days in total

If known Asthma/COPD do not use nebulisers unless no choice in severe exacerbations. (Increase SABA / similar)

Only use oral steroids if sure not COVID

SAFETY-NETTING and consider AKI

Category 2C

Management:

- · Rest and make safe
- Oral fluids
- Paracetamol

Consider antibiotics to treat/prevent secondary bacterial pneumonia:

- Doxycycline 200mg *stat*, 100mg *od*, 5 days in total

OR

- Amoxicillin 500mg *tds,* 5 days in total

If frail, significant chronic disease or immunocompromised [and not a candidate for palliative care] referral to medical team on-call

 SAFETY-NETTING: review in 24-48h (or earlier if unwell), and consider AKI