

THE CURE PROJECT

CURING TOBACCO ADDICTION IN LEICESTER, LEICESTERSHIRE AND RUTLAND



UNIVERSITY HOSPITALS OF LEICESTER NHS TRUST

Stop Smoking Medication

Patient Guidance

Nicotine Lozenges



Your Stop Smoking Medication

Nicotine Lozenges

This lozenge is designed to reduce your urge to smoke by giving a dose of nicotine within 10 to 15 minutes.

1. Avoid acidic drinks such as coffee, juice and soft drinks for up to 15 minutes before a lozenge is used.
2. Do not eat or drink while the lozenge is in your mouth.
3. Put one lozenge in your mouth and allow it to dissolve slowly moving from one side of your mouth to the other from time to time. It should take 20-30 minutes to dissolve.
4. The nicotine is absorbed through the soft skin in your mouth, so it is important not to chew the lozenge or swallow it too quickly. Once the nicotine is in your stomach it is not so effective.
5. If you suck or chew the lozenge too quickly, you may experience hiccups or indigestion. If this happens try putting it under your tongue or down the side of your mouth between your cheek and gum, to let it dissolve more slowly.
6. To begin with you can take one lozenge every 1-2 hours.
7. Do not use more than 15 lozenges per day.

This guidance is applicable for all brands of this product.