

ED Adult Burns Proforma

‘Lund and Browder’ Chart

An aid for documenting burn extent and calculating the percentage of burnt total body surface area (TBSA)

Patient details

Full name


DoB


Unit number

(use sticker if available)

Do not record epidermal (superficial) burns

The surface area of the patient’s palm plus fingers equates to around 1% of their TBSA. Making a paper cutout of similar dimensions can help to estimate the size of larger burns.



	 % superficial or deep partial thickness	 % full thickness
Head		
Neck		
Front		
Back		
R arm		
L arm		
Buttocks		
Genitalia		
R leg		
L leg		
TBSA burnt		

ED Adult Burns Proforma

Fluid prescribing aid for the first 24h

For use only in patients with a burnt TBSA >15%.
Calculations are based on the 'Parkland Formula'.

Patient details

Full name

DoB

Unit number

(use sticker if available)

1. Record % burnt TBSA
2. Record patient's weight
3. Record time of burn injury
4. Record current time
5. Work out how many hours have passed since burn (interval between **A** and **B**; rounded to nearest hour)
6. Work out how many hours remain of the initial 8-hour post-injury period (8 minus **C**)
7. Work out total crystalloid needs for the first 24h ($4 \times \% \text{ burnt TBSA} \times \text{weight in kg}$)
8. Work out rate of infusion so that the first half of the total amount is given within 8h of the burn injury (**E** divided by 2 divided by **D**)
9. Work out rate of infusion so that the second half of the total amount is given within the remaining 16h (**E** divided by 2 divided by 16)
10. Prescribe fluid needs on a drug chart as shown below

	Your patient	Worked example
	%	17 %
	kg	83kg
A	HH:MM use 24h clock	19:45 use 24h clock
B	HH:MM use 24h clock	20:50 use 24h clock
C		1
D		7
E		$4 \times 17 \times 83$ $= 5644\text{ml}$
		$5644 \div 2 \div 7$ $= 403.1\text{ml/h}$
		$5644 \div 2 \div 16$ $= 176.4\text{ml/h}$

Crystalloid prescription for the first 24h after burn injury as per example above										
Date	Infusion fluid		Additions to infusion		IV or SC	Line	Start Time	Time to run or ml/hr	Fluid Batch No.	
	Type/strength	Volume	Drug	Dose						Prescriber
DD/MM/YY	Hartmann	1L			IV		20:50	} 403mL/h until 03:45, from then 176mL/h		Dr.'s Name
DD/MM/YY	Hartmann	1L			IV					Dr.'s Name
DD/MM/YY	Hartmann	1L			IV					Dr.'s Name
DD/MM/YY	Hartmann	1L			IV					Dr.'s Name
DD/MM/YY	Hartmann	1L			IV					Dr.'s Name
DD/MM/YY	Hartmann	1L			IV					Dr.'s Name

Calculations completed by

Print name

Signature

Role

Date

Time