

PATIENT LABEL	Hospital		INREACH LABEL (WHERE APPLICABLE)
	Ward		
	Indication		
	Duration		
	Target Range		
Tait and Sefcick Algorithm For Warfarin Initiation			

1. Check Baseline Bloods (FBC, UE's, LFT's INR & APTT)
2. Prescribe Warfarin on Drug Chart daily
3. Consider Heparin cover for higher risk patients and should continue until INR is in range for two consecutive tests.
4. Heparin cover should be for a minimum of 5 days
5. Commence Warfarin at 5mg Daily for FOUR DAYS and check INR on Day 5

INR on day 5	Dose for days 5-7	INR on day 8	Dose from day 8
<=1.7	5mg	<=1.7 1.8-2.4 2.5-3.0 >3.0	6mg x 7 days 5mg x 7 days 4mg x 7 days 3mg x 4 days
1.8-2.2	4mg	<=1.7 1.8-2.4 2.5-3.0 3.1-3.5 >3.5	5mg x 7 days 4mg x 7 days 3.5mg x 7 days 3mg x 4 days 2.5mg x 4 days
2.3-2.7	3mg	<=1.7 1.8-2.4 2.5-3.0 3.1-3.5 >3.5	4mg x 7 days 3.5mg x 7 days 3mg x 7 days 2.5mg x 4 days 2mg x 4 days
2.8-3.2	2mg	<=1.7 1.8-2.4 2.5-3.0 3.1-3.5 >3.5	3mg x 7 days 2.5mg x 7 days 2mg x 7 days 1.5mg x 4 days 1mg x 4 days
3.3-3.7	1mg	<=1.7 1.8-2.4 2.5-3.0 3.1-3.5 >3.5	2mg x 7 days 1.5mg x 7 days 1mg x 7 days 0.5mg x 4 days omit x 4 days
>3.7	0 mg	<2.0 2.0-2.9 3.0-3.5	1.5mg x 4 days 1mg x 4 days 0.5mg x 4 days

UHL Anticoagulation Service

1. Available for advice, guidance.
2. Available for INReach, ward based service offering advice on dosing and patient education. (where available)
3. Discharge advice and information and monitoring in primary care.
4. Outpatient Clinics Available at all three sites offering education and dosing and smooth transfer to primary care.
5. Ward referrals taken via ICE,
6. Verbal referrals not accepted

Contact Details:

anticoagulation@UHL-tr.nhs.uk

Patient Tel: 0116 2585069

Clinician Tel: 6720 Hotline: 07960779941

Bleep: (GH) 2712, (LRI) 7158

BLEEDING WHILST ANTICOAGULATED: 4 CATEGORIES

1, Life threatening haemorrhage Stop Warfarin 5mg Vitamin K by slow injection (refer to IV policy)	3, INR high without haemorrhage INR >4.5 withhold Warfarin for 1-2 days and review INR >8.0 give 0.5mg Vitamin k by slow injection
2, Less Severe haemorrhage Withhold Warfarin for 1 or more days and review And/or give 0.5mg vitamin K by slow injection	4, Unexpected bleeding at therapeutic levels Investigate underlying cause 5, Contact the UHL Anticoagulation Service for further advice if required.
NB: Take care in reversing anticoagulation in patients with prosthetic valves	

Completed By (BLOCK CAPITALS):	Signature:	Date:
--------------------------------	------------	-------

[illegible]