

Emergency Decision Unit Pathway

NAC treatment

Date

DD/MM/YY

Patient details

Full name

DoB

Unit number

(use sticker if available)

Inclusion criteria	ED/EDU patients requiring a course of NAC as per Adult ED paracetamol poisoning proforma
Exclusion criteria	- Abnormal initial blood results (admit to AMU) - Co-ingestion of drugs with actual or expected serious cardiovascular or central nervous side-effects (i.e. tricyclic antidepressants or beta-blockers)

Notes to doctor completing this pathway (EPIC to ensure compliance)

- This pathway must only be used in conjunction with the [Adult ED paracetamol poisoning proforma](#)
- Ensure that proforma was fully completed
- If the decision to start NAC is made for a pt admitted on the EDU toxicology pathway, start **THIS** pathway on EDU
 - Prescribe complete SNAP regimen including PRN medicines for any adverse reactions:
 - If patient weighs 40kg and over, go to NC Meds > Emergency Medicine (ED) > Antidotes (ED) > N-Acetylcysteine > [select weight range] > Complete SNAP regimen
 - If less than 40kg, use NAC dosing table from [Children’s ED paracetamol poisoning proforma](#)

EDU plan

- Deliver complete SNAP regimen (1st bag; 100mg/kg over 2h then 2nd bag, 200mg/kg over 10h)
- Starting from within last hour of the infusion, manage as per flowchart on the back
- If CBG in diabetics proves difficult to control while receiving the 2nd (10h) bag, stop the infusion, go to N-Acetylcysteine as above > [select weight range] > Repeat 10hr NAC infusions and select 0.9% NaCl option instead

Planned & agreed by

Print name

Signature

Referring Clinician

NB: Pathway must be filled in and signed by the referring clinician and then discussed with, and 'EDU admission approved' written by, a consultant or the EPIC in the Nervecentre 'ED Progress Notes'

Patients will NOT be accepted until this has been done

EDU management of paracetamol OD after SNAP regimen completed

