


Adult (over 18 years old) Gentamicin Once Daily Treatment Regimen Prescription

Section 1: Complete and sign checklist prior to administration of 1st dose. Tick each box to indicate completed/reviewed

- | | | |
|--|---|--------------------------|
| 1 Contact ward / on-call pharmacist for dosing and assessment of risk factors, document pharmacist name below | <i>Not applicable if patient on ITU and prescriber familiar with prescribing Gentamicin</i> | ☐ |
| 2 Do not use in any of the following conditions: endocarditis, pregnancy, cystic fibrosis, major burns, ascites, dialysis | <i>Use conventional divided doses regimen if any of these exclusions apply</i> | ☐ |
| 3 Do not use if patient's creatinine clearance is less than 30mls/min | <i>CrCl electronic calculator, and dosing advice in renal impairment available on Antimicrobial intranet site</i> | ☐ |
| 4 Caution if previous courses of gentamicin in the last 3 months | <i>If respiratory patient review with respiratory Consultant</i> | ☐ |
| 5 Consider risk factors for toxicity: <u>age>65 years</u> , other drugs (NSAIDS, ACE inhibitors, diuretics), dehydration, sepsis | <i>Check renal function daily if risk factors, until renal function and gentamicin levels stable.</i> | ☐ |
| If a patient only has one kidney, where possible use an alternative suitable antibiotic (having only one kidney, with normal GFR, is not a contra-indication for gentamicin administration, but caution is advised). | | ☐ |
| 6 Inform patients of possible adverse effects | | ☐ |
| 7 Prescribers Signature: | Pharmacist Name: | |

Section 2: Calculate First Dose

- Calculate dose according to patient weight. Complete the flow diagram below.
- DO NOT ESTIMATE weight, if patient cannot be weighed use Ideal Body Weight (IBW)**



Ideal Body Weight and Overweight Chart					
Height		MEN		WOMEN	
ft. inches	cm	IB W (kg)	Overweight if >kg	IBW (kg)	Overweight if >kg
5.0	152.40	50	60	45.5	54.5
5.1	154.94	52.3	62.5	47.8	57
5.2	157.48	54.6	65.5	50.1	60
5.3	160.02	56.9	68	52.4	63
5.4	162.56	59.2	71	54.7	65.5
5.5	165.10	61.5	74	57.0	68.5
5.6	167.64	63.8	76.5	59.3	71
5.7	170.18	66.1	79	61.6	74
5.8	172.72	68.4	82	63.9	76.5
5.9	175.26	70.7	85	66.2	79.5
5.10	177.80	73.0	87.5	68.5	82.2
5.11	180.34	75.3	90	70.8	85
6.0	182.88	77.6	93	73.1	87.5
6.1	185.42	79.9	96	75.4	90
6.2	187.96	82.2	98.5	77.7	93
6.3	190.50	84.5	101.5	80.0	96
6.4	193.04	86.8	104	82.3	98.5
6.5	195.58	89.1	107	84.6	101.5

Section 3: Prescribing

- On the patients main prescription chart **prescribe and sign for** gentamicin, write in the dose section 'give as per gentamicin chart'
- Each dose of gentamicin to be prescribed and signed for on this chart.
- If Gentamicin injection is not stocked on the ward, order from pharmacy for individual patients
- In ePMA prescribe for 5 days – document 'give as per Gentamicin chart' in the qualifier

Adult (over 18 years old) Gentamicin Once Daily Treatment Regimen Prescription

Section 1 to 3 MUST be completed BEFORE completing section 4


<u>Patient Addressograph</u> Hospital No: Name: D.O.B:	Ward: Consultant: Indication: Intended duration / Review date:
---	---

Section 4: Monitoring (refer to bottom of page for more details) and Assay Sampling Lab and Transport Information

- Take a single blood sample at any time between 6-14 hours after the start of infusion of 1st dose.
- Further levels should be monitored on day 3 and 7, or the day after any change of duration.
- Samples must be sent to the **fast track chemistry laboratory, Level 4, Sandringham Building, LRI** without delay. Samples will be analysed on receipt and results will be available immediately after analysis on iLAB. Please do not phone the laboratory for results

* Before administration check if levels are needed after this dose or if any previous levels have been taken and actioned appropriately											
Prescription Give in 100ml 0.9% sodium chloride / 5% glucose as an IV infusion over 1 hour						Blood levels	Administration*		Blood levels		Advice
Day	Date	Dose (usual MAX dose 640mg)	Serum Creatinine	Time Dose Due	Signature of prescriber / Print name	Date / Time level to be taken (6-14 hrs after start of infusion)	Date and Time given	Given by / Check by	Date and Time blood taken	Level result (mg/l)	New dose if change needed from nomogram and Signature
1											
2											
3											
4											
5											

Re-prescribe on new chart and obtain microbiology verification code (not required if duration approved in guidelines e.g bronchiectasis) Code:



Monitoring

- Take a single blood sample (**brown top serum gel tube Z/4.9 ml** or 2.7ml paediatric) at any time between **6-14 hours after the start of infusion of 1st dose**. It is **vital** to record the exact times of commencing infusion and taking the blood level on this prescription chart. Always write the sample time on the **combined chemistry/haematology request** form before sending for processing.
- Use the **Hartford nomogram above** to determine at what time intervals subsequent doses should be given. All dosages remain the same. If the level falls in the area designated Q24h, Q36h or Q48h the interval should be 24, 36 or 48 hours respectively. If the plotted point falls on a line, choose the longer time interval. If plotted point is above 48hrs line do not give further dose without further levels.
- Further gentamicin levels should be monitored on day 3 and 7 or the day after any change of duration.
- Monitor renal function by measuring serum creatinine three times a week or daily if unstable. If serum creatinine rises by more than 45 micromoles/L, stop gentamicin and discuss with microbiology / pharmacy.
- Close monitoring of auditory and vestibular functions