

LRI Emergency Department

HIV post-exposure prophylaxis following sexual exposure (PEPSE)

Version 77

PEPSE is time-critical

Assign DPS 2 on arrival to facilitate rapid risk assessment and appropriate treatment

Disclaimer:
This is a clinical template; clinicians should always use judgment when managing individual patients

Created by Martin Wiese with Benjamin Feist, Helen Wiggins and Hannah Church

Re-approved by ED guidelines committee on 28May25
Review due May 2028 . Trust Ref: C194/2016

Patient details

Full name

DoB

Unit number

(use sticker if available)

① Finding the applicable Sexual Health Service

Go to the 'Find your local council' page on GOV.UK to find the council serving your patient

Leicester City
[Leicester Sexual Health](#) (LSH)

Mon 08:15 - 18:15

Tue 08:15 - 18:15

Wed 08:15 - 18:15

Thu 08:15 - 16:00

Fri 08:15 - 18:15

0300 124 0102

Leicestershire and Rutland
[Leicestershire and Rutland Sexual Health](#) (SH:24)

Mon 08:15 - 16:30

Tue 08:15 - 18:30


Wed 08:15 - 18:30

Thu 08:15 - 17:30

Fri 08:15 - 16:30

Sat 09:15 - 14:30

0300 247 0068



Patient seen by

Print name

Signature

Role

④ Sexual health service referral email template

Copy & paste the applicable service's email address into the 'To' line
For Leicester City
LLRHealthAdvisingTeam@mpft.nhs.uk
For Leicestershire and Rutland
HCRG.LRSexualhealthhub@nhs.net

Copy & paste into the subject line
Referral from LRI ED

Copy all of the text on the right into the email body (use the 'Paste text only' option to preserve format). Add/delete [including help text] as needed.

Dear colleague, your patient presented to our ED today following potential sexual exposure to HIV. [Consider adding detail in one sentence].

NHS number:

NAME:

DOB:

ADDRESS:

PHONE:

Date of exposure:

Date PEPSE started:

[Below, delete YES or NO as applicable].

Permission for voice message given? YES NO

Permission for text message given? YES NO

Hepatitis B vaccine dose given? YES NO

② Is source from a high HIV prevalence group?

Tick relevant prevalence (in %)

Men who have sex with men

England 2.30

London 3.21

Elsewhere in the UK 2.09

Heterosexual men

Black African 0.58

Non-black African 0.02

Heterosexual women

Black African 0.87

Non-Black African 0.01

People who inject drugs

All 0.67

Men 0.53

Women 1.15

☐ **Yes** – prevalence in group ≥1%

☐ **No** – prevalence in group <1%

③ Need for PEP cautions?

☐ **YES**, as at least one of the below

Pregnancy (consider testing)

Known or suspected eGFR<50

Source HIV positive **AND** on antiretroviral therapy

Exposed patient is taking

Phenobarbital,

Rifampicin

Carbamazepine

Phenytoin

Oxcarbazepine

Calcium

Magnesium

Iron

Aluminium (often found in multi-vitamin supplements or indigestion remedies)

Exposed patient has been taking PrEP (pre-exposure prophylaxis)

☐ **NO**, as none of the above

⑤ HIV transmission risk assessment

tick applicable sexual activities and source individual's HIV status group

HIV status of source individual / sexual partner / assailant

Positive	Unknown	Positive	Unknown
<i>One or more of the below</i> • Viral load detectable • Viral load unknown • Undetectable HIV viral load (<200 copies/mL) since less than 6 months • On antiretroviral therapy since less than 6 months • Uncertainty about adherence to antiretroviral therapy	From group where prevalence is ≥1% (see box 2)	ALL of the below • Undetectable HIV viral load (<200 copies/mL) when last tested and within the last 6 months • AND on antiretroviral therapy for at least the last 6 months • AND good adherence to antiretroviral therapy	From group where prevalence is <1% (see box 2)
☐	☐	☐	☐

Sexual activity

☐ Receptive anal sex
☐ Insertive anal sex
☐ Sharing injecting equipment (AKA 'slamming' or 'chemsex')
☐ Receptive vaginal sex
☐ Insertive vaginal sex
☐ Bite (= breakage of the skin of either person with passage of blood) with ALL THREE of the conditions below ☐ 'Biter' is known or suspected to have an HIV viral load >3.0 log copies/mL AND ☐ 'Biter's saliva was visibly contaminated with blood AND ☐ Bite has resulted in severe and/or deep tissue injuries
☐ Bite (= breakage of the skin of either person with passage of blood) without any of the risk factors above
☐ Receptive oral sex (giving fellatio)
☐ Insertive oral sex (receiving fellatio)
☐ Splash of semen into eye
☐ Cunnilingus (oral sex involving the female genitalia)

Start PEPSE	Start PEPSE
Start PEPSE	seek advice
Start PEPSE	seek advice
Start PEPSE	seek advice
seek advice	seek advice
seek advice	

PEPSE not indicated and advice from Sexual Health not required

Transmission risk of HIV is less than 1 in 10.000