


Slides developed by Mine Çetinkaya-Rundel of OpenIntro
Translated from LaTeX to Google Slides by Curry W. Hilton of OpenIntro.
The slides may be copied, edited, and/or shared via the [CC BY-SA license](https://creativecommons.org/licenses/by-sa/4.0/)

To make a copy of these slides, go to *File > Download as > [option]*, as shown below. Or if you are logged into a Google account, you can choose *Make a copy...* to create your own version in Google Drive.



Negative binomial distribution

Negative binomial distribution

- The negative binomial distribution describes the probability of observing the k th success on the n th trial.
- The following four conditions are useful for identifying a negative binomial case:
 1. The trials are independent.
 2. Each trial outcome can be classified as a success or failure.
 3. The probability of success (p) is the same for each trial.
 4. The last trial must be a success.

Note that the first three conditions are common to the binomial distribution.

Negative binomial distribution

$$P(k^{th} \text{ success on the } n^{th} \text{ trial}) = \binom{n-1}{k-1} p^k (1-p)^{n-k},$$

where p is the probability that an individual trial is a success. All trials are assumed to be independent.

Practice

A college student working at a psychology lab is asked to recruit 10 couples to participate in a study. She decides to stand outside the student center and ask every 5th person leaving the building whether they are in a relationship and, if so, whether they would like to participate in the study with their significant other. Suppose the probability of finding such a person is 10%. What is the probability that she will need to ask 30 people before she hits her goal?

Practice

A college student working at a psychology lab is asked to recruit 10 couples to participate in a study. She decides to stand outside the student center and ask every 5th person leaving the building whether they are in a relationship and, if so, whether they would like to participate in the study with their significant other. Suppose the probability of finding such a person is 10%. What is the probability that she will need to ask 30 people before she hits her goal?

Given: $p = 0.10$, $k = 10$, $n = 30$. We are asked to find the probability of 10th success on the 30th trial, therefore we use the negative binomial distribution.

Practice

A college student working at a psychology lab is asked to recruit 10 couples to participate in a study. She decides to stand outside the student center and ask every 5th person leaving the building whether they are in a relationship and, if so, whether they would like to participate in the study with their significant other. Suppose the probability of finding such a person is 10%. What is the probability that she will need to ask 30 people before she hits her goal?

Given: $p = 0.10$, $k = 10$, $n = 30$. We are asked to find the probability of 10th success on the 30th trial, therefore we use the negative binomial distribution.

$$P(10^{th} \text{ success on the } 30^{th} \text{ trial})$$

Practice

A college student working at a psychology lab is asked to recruit 10 couples to participate in a study. She decides to stand outside the student center and ask every 5th person leaving the building whether they are in a relationship and, if so, whether they would like to participate in the study with their significant other. Suppose the probability of finding such a person is 10%. What is the probability that she will need to ask 30 people before she hits her goal?

Given: $p = 0.10$, $k = 10$, $n = 30$. We are asked to find the probability of 10th success on the 30th trial, therefore we use the negative binomial distribution.

$$P(10^{th} \text{ success on the } 30^{th} \text{ trial}) = \binom{29}{9} \times 0.10^{10} \times 0.90^{20}$$

Practice

A college student working at a psychology lab is asked to recruit 10 couples to participate in a study. She decides to stand outside the student center and ask every 5th person leaving the building whether they are in a relationship and, if so, whether they would like to participate in the study with their significant other. Suppose the probability of finding such a person is 10%. What is the probability that she will need to ask 30 people before she hits her goal?

Given: $p = 0.10$, $k = 10$, $n = 30$. We are asked to find the probability of 10th success on the 30th trial, therefore we use the negative binomial distribution.

$$\begin{aligned} P(10^{th} \text{ success on the } 30^{th} \text{ trial}) &= \binom{29}{9} \times 0.10^{10} \times 0.90^{20} \\ &= 10,015,005 \times 0.10^{10} \times 0.90^{20} \end{aligned}$$

Practice

A college student working at a psychology lab is asked to recruit 10 couples to participate in a study. She decides to stand outside the student center and ask every 5th person leaving the building whether they are in a relationship and, if so, whether they would like to participate in the study with their significant other. Suppose the probability of finding such a person is 10%. What is the probability that she will need to ask 30 people before she hits her goal?

Given: $p = 0.10$, $k = 10$, $n = 30$. We are asked to find the probability of 10th success on the 30th trial, therefore we use the negative binomial distribution.

$$\begin{aligned}P(10^{th} \text{ success on the } 30^{th} \text{ trial}) &= \binom{29}{9} \times 0.10^{10} \times 0.90^{20} \\&= 10,015,005 \times 0.10^{10} \times 0.90^{20} \\&= 0.00012\end{aligned}$$

Binomial vs negative binomial

How is the negative binomial distribution different from the binomial distribution?

Binomial vs negative binomial

How is the negative binomial distribution different from the binomial distribution?

- In the binomial case, we typically have a fixed number of trials and instead consider the number of successes.
- In the negative binomial case, we examine how many trials it takes to observe a fixed number of successes and require that the last observation be a success.

Practice

Which of the following describes a case where we would use the negative binomial distribution to calculate the desired probability?

- (a) Probability that a 5 year old boy is taller than 42 inches.
- (b) Probability that 3 out of 10 softball throws are successful.
- (c) Probability of being dealt a straight flush hand in poker.
- (d) Probability of missing 8 shots before the first hit.
- (e) Probability of hitting the ball for the 3rd time on the 8th try.

Practice

Which of the following describes a case where we would use the negative binomial distribution to calculate the desired probability?

- (a) Probability that a 5 year old boy is taller than 42 inches.
- (b) Probability that 3 out of 10 softball throws are successful.
- (c) Probability of being dealt a straight flush hand in poker.
- (d) Probability of missing 8 shots before the first hit.
- (e) Probability of hitting the ball for the 3rd time on the 8th try.*