

GENERAL PHYSICS PH1110

Nam Le, Dr.

School of Engineering Physics

Hanoi University of Science and Technology

2. THERMODYNAMICS

2.1 KINETIC THEORY AND LAWS OF DISTRIBUTION

- 1 **STARTING POINTS**
- 2 **EQUATION OF STATE**
- 3 **KINETIC THEORY OF GASES**
 - Exercises
- 4 **MAXWELL DISTRIBUTION**
- 5 **BOLTZMANN DISTRIBUTION**

1. Starting points

1. Starting points

♣ States of matter

1. Starting points

♣ States of matter

- All matter consists of atoms or molecules (groups of atoms).

1. Starting points

♣ States of matter

- All matter consists of atoms or molecules (groups of atoms).
- The spacing of these molecules and the forces between them determine whether the matter is a **solid**, a **liquid** or a **gas**.

1. Starting points

♣ States of matter

- All matter consists of atoms or molecules (groups of atoms).
- The spacing of these molecules and the forces between them determine whether the matter is a **solid**, a **liquid** or a **gas**.

♣ Gases

1. Starting points

♣ States of matter

- All matter consists of atoms or molecules (groups of atoms).
- The spacing of these molecules and the forces between them determine whether the matter is a **solid**, a **liquid** or a **gas**.

♣ **Gases** have no fixed volume and no fixed shape.

1. Starting points

♣ States of matter

- All matter consists of atoms or molecules (groups of atoms).
- The spacing of these molecules and the forces between them determine whether the matter is a **solid**, a **liquid** or a **gas**.

♣ Gases have no fixed volume and no fixed shape.

- In gases, the separation of the molecules is large and the interaction between the molecules is small.

1. Starting points

♣ States of matter

- All matter consists of atoms or molecules (groups of atoms).
- The spacing of these molecules and the forces between them determine whether the matter is a **solid**, a **liquid** or a **gas**.

♣ Gases have no fixed volume and no fixed shape.

- In gases, the separation of the molecules is large and the interaction between the molecules is small.
- The **parameters** of a gaseous state are

1. Starting points

♣ States of matter

- All matter consists of atoms or molecules (groups of atoms).
- The spacing of these molecules and the forces between them determine whether the matter is a **solid**, a **liquid** or a **gas**.

♣ Gases have no fixed volume and no fixed shape.

- In gases, the separation of the molecules is large and the interaction between the molecules is small.
- The **parameters** of a gaseous state are volume, temperature and pressure.

1. Starting points

♣ Recall

One mole of a substance is

The Avogadro constant is

1. Starting points

♣ Recall

One mole of a substance is the amount of that substance which contains the same number of particles as there are in 0.012 kg of carbon-12.

The Avogadro constant is

1. Starting points

♣ Recall

One mole of a substance is the amount of that substance which contains the same number of particles as there are in 0.012 kg of carbon-12.

The Avogadro constant is the number of particles that are contained in the amount of substance given by one mole.

1. Starting points

♣ Recall

One mole of a substance is the amount of that substance which contains the same number of particles as there are in 0.012 kg of carbon-12.

The Avogadro constant is the number of particles that are contained in the amount of substance given by one mole.

- ▶ So, the Avogadro constant is equal to the number carbon atoms in 0.012 kg of carbon-12.

1. Starting points

♣ Recall

One mole of a substance is the amount of that substance which contains the same number of particles as there are in 0.012 kg of carbon-12.

The Avogadro constant is the number of particles that are contained in the amount of substance given by one mole.

- So, the Avogadro constant is equal to the number carbon atoms in 0.012 kg of carbon-12.

$$N_A = 6.02214 \times 10^{23} \text{ mol}^{-1}$$

2. THERMODYNAMICS

2.1 KINETIC THEORY AND LAWS OF DISTRIBUTION

- 1 **STARTING POINTS**
- 2 **EQUATION OF STATE**
- 3 **KINETIC THEORY OF GASES**
 - Exercises
- 4 **MAXWELL DISTRIBUTION**
- 5 **BOLTZMANN DISTRIBUTION**

2. Equation of state

2. Equation of state

♣ Experimental laws for gases

2. Equation of state

♣ Experimental laws for gases

Boyle's law:

Charles' law:

Gay-Lussac's law (law of pressures):

2. Equation of state

♣ Experimental laws for gases

Boyle's law: The volume of a gas is inversely proportional to its pressure, provided that the temperature is held constant.

Charles' law:

Gay-Lussac's law (law of pressures):

2. Equation of state

♣ Experimental laws for gases

Boyle's law: The volume of a gas is inversely proportional to its pressure, provided that the temperature is held constant.

Charles' law: The volume of a gas is directly proportional to its thermodynamic temperature, provided that the pressure is held constant.

Gay-Lussac's law (law of pressures):

2. Equation of state

♣ Experimental laws for gases

Boyle's law: The volume of a gas is inversely proportional to its pressure, provided that the temperature is held constant.

Charles' law: The volume of a gas is directly proportional to its thermodynamic temperature, provided that the pressure is held constant.

Gay-Lussac's law (law of pressures): The pressure of a gas is directly proportional to its thermodynamic temperature, provided that the volume is held constant.

2. Equation of state

- Boyle's law: If T is constant then $V \propto 1/p$.
- Charles' law: If p is constant then $V \propto T$.
- Gay-Lussac law: If V is constant then $p \propto T$.

2. Equation of state

- Boyle's law: If T is constant then $V \propto 1/p$.
- Charles' law: If p is constant then $V \propto T$.
- Gay-Lussac law: If V is constant then $p \propto T$.

♣ These three laws can be combined into a single equation:

2. Equation of state

- Boyle's law: If T is constant then $V \propto 1/p$.
- Charles' law: If p is constant then $V \propto T$.
- Gay-Lussac law: If V is constant then $p \propto T$.

♣ These three laws can be combined into a single equation:

$$pV \propto T$$

2. Equation of state

- Boyle's law: If T is constant then $V \propto 1/p$.
- Charles' law: If p is constant then $V \propto T$.
- Gay-Lussac law: If V is constant then $p \propto T$.

♣ These three laws can be combined into a single equation:

$$\boxed{pV \propto T} \quad \text{or} \quad \boxed{\frac{pV}{T} = \text{constant}}$$

2. Equation of state

- Boyle's law: If T is constant then $V \propto 1/p$.
- Charles' law: If p is constant then $V \propto T$.
- Gay-Lussac law: If V is constant then $p \propto T$.

♣ These three laws can be combined into a single equation:

$$\boxed{pV \propto T} \quad \text{or} \quad \boxed{\frac{pV}{T} = \text{constant}}$$

- The value of the constant is dependent of the amount of gas.

2. Equation of state

- Boyle's law: If T is constant then $V \propto 1/p$.
- Charles' law: If p is constant then $V \propto T$.
- Gay-Lussac law: If V is constant then $p \propto T$.

♣ These three laws can be combined into a single equation:

$$\boxed{pV \propto T} \quad \text{or} \quad \boxed{\frac{pV}{T} = \text{constant}}$$

- The value of the constant is dependent of the amount of gas. It is proportional to the number n of moles in the gas.

2. Equation of state

$$\frac{pV}{T} \propto n$$

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.
- The **Boltzmann constant**:

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.
- The **Boltzmann constant**: $k_B = \frac{R}{N_A}$

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.
- The **Boltzmann constant**: $k_B = \frac{R}{N_A} = 1.38 \times 10^{-23} \text{ J K}^{-1}$

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.
- The **Boltzmann constant**: $k_B = \frac{R}{N_A} = 1.38 \times 10^{-23} \text{ J K}^{-1}$

♣ Equation of state

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.
- The **Boltzmann constant**: $k_B = \frac{R}{N_A} = 1.38 \times 10^{-23} \text{ J K}^{-1}$

♣ Equation of state

$$pV = nRT$$

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.
- The **Boltzmann constant**: $k_B = \frac{R}{N_A} = 1.38 \times 10^{-23} \text{ J K}^{-1}$

♣ Equation of state

$$pV = nRT$$

or

$$pV = Nk_B T$$

2. Equation of state

$$\frac{pV}{T} \propto n \quad \rightarrow \quad \frac{pV}{T} = Rn, \quad \text{where } R \text{ is a constant.}$$

- $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ – is called the **molar gas constant**.
- The **Boltzmann constant**: $k_B = \frac{R}{N_A} = 1.38 \times 10^{-23} \text{ J K}^{-1}$

♣ Equation of state

$$pV = nRT$$

or

$$pV = Nk_B T$$

- Here, $N = nN_A$ is the total number of particles in the gas.

2. Equation of state

♣ Ideal gases

2. Equation of state

♣ Ideal gases

- The gas laws can approximately be applied to **real gases** at low pressure and high temperature.

2. Equation of state

♣ Ideal gases

- The gas laws can approximately be applied to **real gases** at low pressure and high temperature.
- No real gases obey the laws at any pressures, volumes and temperatures.

2. Equation of state

♣ Ideal gases

- The gas laws can approximately be applied to **real gases** at low pressure and high temperature.
- No real gases obey the laws at any pressures, volumes and temperatures.

A gas which obeys the equation of state, $pV = nRT$, for all pressures, volumes and temperatures

2. Equation of state

♣ Ideal gases

- The gas laws can approximately be applied to **real gases** at low pressure and high temperature.
- No real gases obey the laws at any pressures, volumes and temperatures.

A gas which obeys the equation of state, $pV = nRT$, for all pressures, volumes and temperatures is called an **ideal gas**.

2. THERMODYNAMICS

2.1 KINETIC THEORY AND LAWS OF DISTRIBUTION

- 1 STARTING POINTS
- 2 EQUATION OF STATE
- 3 KINETIC THEORY OF GASES
 - Exercises
- 4 MAXWELL DISTRIBUTION
- 5 BOLTZMANN DISTRIBUTION

3. Kinetic theory of gases

3. Kinetic theory of gases

♣ The first evidence for the existence of atoms/molecules is

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

- ♣ The basic idea for explaining the gas pressure is

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

- ♣ The basic idea for explaining the gas pressure is that the gas consists of molecules randomly moving about at great speed.

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

- ♣ The basic idea for explaining the gas pressure is that the gas consists of molecules randomly moving about at great speed.
 - When a molecule hits the wall

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

- ♣ The basic idea for explaining the gas pressure is that the gas consists of molecules randomly moving about at great speed.
 - When a molecule hits the wall → rebounds

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

- ♣ The basic idea for explaining the gas pressure is that the gas consists of molecules randomly moving about at great speed.
 - When a molecule hits the wall → rebounds → experiences an impulse

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

- ♣ The basic idea for explaining the gas pressure is that the gas consists of molecules randomly moving about at great speed.
- When a molecule hits the wall → rebounds → experiences an impulse → (Newton's third law) gives an impulse to the wall.

3. Kinetic theory of gases

- ♣ The first evidence for the existence of atoms/molecules is the **Brownian motion**, such as **motion of pollen grains in water**.

Brownian motion is the random motion of particles suspended in a fluid resulting from their collision with the fast-moving molecules in the fluid.

- ♣ The basic idea for explaining the gas pressure is that the gas consists of molecules randomly moving about at great speed.
 - When a molecule hits the wall → rebounds → experiences an impulse → (Newton's third law) gives an impulse to the wall.
 - The gas pressure cannot be explained if the gas is continuous.

3. Kinetic theory of gases

♣ The kinetic theory of an ideal gas

3. Kinetic theory of gases

♣ The kinetic theory of an ideal gas

- 1 A gas contains a very large number of particles (atoms or molecules).

3. Kinetic theory of gases

♣ The kinetic theory of an ideal gas

- 1 A gas contains a very large number of particles (atoms or molecules).
- 2 The forces between particles are negligible, except during collisions.

3. Kinetic theory of gases

♣ The kinetic theory of an ideal gas

- 1 A gas contains a very large number of particles (atoms or molecules).
- 2 The forces between particles are negligible, except during collisions.
- 3 The volume of the particles is negligible compared to the volume occupied by the gas.

3. Kinetic theory of gases

♣ The kinetic theory of an ideal gas

- 1 A gas contains a very large number of particles (atoms or molecules).
- 2 The forces between particles are negligible, except during collisions.
- 3 The volume of the particles is negligible compared to the volume occupied by the gas.
- 4 The collisions of particles with each other and with the container are perfectly elastic.

3. Kinetic theory of gases

♣ The kinetic theory of an ideal gas

- 1 A gas contains a very large number of particles (atoms or molecules).
- 2 The forces between particles are negligible, except during collisions.
- 3 The volume of the particles is negligible compared to the volume occupied by the gas.
- 4 The collisions of particles with each other and with the container are perfectly elastic.
- 5 The particles mostly move in straight lines at constant velocities. The collision time is negligible compared with the time between collisions.

3. Kinetic theory of gases


♣ How pressure and volume relate to molecular motion?

3. Kinetic theory of gases

- ♣ How pressure and volume relate to molecular motion?
- Suppose the container is a cube of side L .

3. Kinetic theory of gases


- ♣ How pressure and volume relate to molecular motion?
- Suppose the container is a cube of side L .



3. Kinetic theory of gases

♣ How pressure and volume relate to molecular motion?


- Suppose the container is a cube of side L .
- Consider a molecule with x -component of velocity c_x and mass m .



3. Kinetic theory of gases

♣ How pressure and volume relate to molecular motion?


- Suppose the container is a cube of side L .
- Consider a molecule with x -component of velocity c_x and mass m .
- Time between two consecutive collisions with one wall



3. Kinetic theory of gases

♣ How pressure and volume relate to molecular motion?


- Suppose the container is a cube of side L .
- Consider a molecule with x -component of velocity c_x and mass m .
- Time between two consecutive collisions with one wall = $2L/c_x$.



3. Kinetic theory of gases

♣ How pressure and volume relate to molecular motion?


- Suppose the container is a cube of side L .
- Consider a molecule with x -component of velocity c_x and mass m .
- Time between two consecutive collisions with one wall = $2L/c_x$.
- Change in momentum in each collision



3. Kinetic theory of gases

♣ How pressure and volume relate to molecular motion?

- Suppose the container is a cube of side L .
- Consider a molecule with x -component of velocity c_x and mass m .
- Time between two consecutive collisions with one wall = $2L/c_x$.
- Change in momentum in each collision = $2mc_x$.



3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}}$$

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x}$$

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

- The total force acting by N particles in the container is

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

- The total force acting by N particles in the container is

$$N \left\langle \frac{mc_x^2}{L} \right\rangle$$

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

- The total force acting by N particles in the container is

$$N \left\langle \frac{mc_x^2}{L} \right\rangle = \frac{Nm}{L} \langle c_x^2 \rangle$$

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

- The total force acting by N particles in the container is

$$N \left\langle \frac{mc_x^2}{L} \right\rangle = \frac{Nm}{L} \langle c_x^2 \rangle$$

- Pressure is force divided by area:

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

- The total force acting by N particles in the container is

$$N \left\langle \frac{mc_x^2}{L} \right\rangle = \frac{Nm}{L} \langle c_x^2 \rangle$$

- Pressure is force divided by area:

$$p = \frac{Nm \langle c_x^2 \rangle / L}{L^2}$$

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

- The total force acting by N particles in the container is

$$N \left\langle \frac{mc_x^2}{L} \right\rangle = \frac{Nm}{L} \langle c_x^2 \rangle$$

- Pressure is force divided by area:

$$p = \frac{Nm \langle c_x^2 \rangle / L}{L^2} = \frac{Nm \langle c_x^2 \rangle}{L^3}$$

3. Kinetic theory of gases

- The average force exerted by each molecule to a wall in the x direction is equal to

$$\frac{\text{change in momentum}}{\text{time between collisions}} = \frac{2mc_x}{2L/c_x} = \frac{mc_x^2}{L}$$

- The total force acting by N particles in the container is

$$N \left\langle \frac{mc_x^2}{L} \right\rangle = \frac{Nm}{L} \langle c_x^2 \rangle$$

- Pressure is force divided by area:

$$p = \frac{Nm \langle c_x^2 \rangle / L}{L^2} = \frac{Nm \langle c_x^2 \rangle}{L^3} = \frac{Nm \langle c_x^2 \rangle}{V}$$

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.
- Since $c^2 = c_x^2 + c_y^2 + c_z^2$

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.
- Since $c^2 = c_x^2 + c_y^2 + c_z^2 \rightarrow \langle c^2 \rangle = \langle c_x^2 \rangle + \langle c_y^2 \rangle + \langle c_z^2 \rangle$

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.
- Since $c^2 = c_x^2 + c_y^2 + c_z^2 \rightarrow \langle c^2 \rangle = \langle c_x^2 \rangle + \langle c_y^2 \rangle + \langle c_z^2 \rangle = 3\langle c_x^2 \rangle$.

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.
- Since $c^2 = c_x^2 + c_y^2 + c_z^2 \rightarrow \langle c^2 \rangle = \langle c_x^2 \rangle + \langle c_y^2 \rangle + \langle c_z^2 \rangle = 3\langle c_x^2 \rangle$.
- Finally:

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.
- Since $c^2 = c_x^2 + c_y^2 + c_z^2 \rightarrow \langle c^2 \rangle = \langle c_x^2 \rangle + \langle c_y^2 \rangle + \langle c_z^2 \rangle = 3\langle c_x^2 \rangle$.

- Finally:
$$pV = \frac{1}{3}Nm\langle c^2 \rangle$$

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.
- Since $c^2 = c_x^2 + c_y^2 + c_z^2 \rightarrow \langle c^2 \rangle = \langle c_x^2 \rangle + \langle c_y^2 \rangle + \langle c_z^2 \rangle = 3\langle c_x^2 \rangle$.

- Finally:
$$pV = \frac{1}{3}Nm\langle c^2 \rangle$$

Remember that, in this relation, m is the molecular mass

3. Kinetic theory of gases

- Therefore, $pV = Nm\langle c_x^2 \rangle$.
- Because the number of particles is very large and the motion of the particles is random $\rightarrow \langle c_x^2 \rangle = \langle c_y^2 \rangle = \langle c_z^2 \rangle$.
- Since $c^2 = c_x^2 + c_y^2 + c_z^2 \rightarrow \langle c^2 \rangle = \langle c_x^2 \rangle + \langle c_y^2 \rangle + \langle c_z^2 \rangle = 3\langle c_x^2 \rangle$.

- Finally:
$$pV = \frac{1}{3}Nm\langle c^2 \rangle$$

Remember that, in this relation, m is the molecular mass and $\langle c^2 \rangle$ is the **mean-square speed** of the molecules.

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

- The average value of kinetic energy:

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

- The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

- The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$
- Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle$

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

- The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$
- Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle \rightarrow m \langle c^2 \rangle = 3k_B T$

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

- The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$
- Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle \rightarrow m \langle c^2 \rangle = 3k_B T$
- Therefore:

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

○ The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$

○ Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle \rightarrow m \langle c^2 \rangle = 3k_B T$

● Therefore: $\langle E_k \rangle = \frac{3}{2} k_B T$

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

○ The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$

○ Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle \rightarrow m \langle c^2 \rangle = 3k_B T$

● Therefore: $\boxed{\langle E_k \rangle = \frac{3}{2} k_B T}$ (a crucial relation)

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

○ The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$

○ Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle \rightarrow m \langle c^2 \rangle = 3k_B T$

● Therefore: $\boxed{\langle E_k \rangle = \frac{3}{2} k_B T}$ (a crucial relation)

This is the relation between kinetic energy and temperature for systems of **monatomic molecules**.

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

○ The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$

○ Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle \rightarrow m \langle c^2 \rangle = 3k_B T$

● Therefore: $\boxed{\langle E_k \rangle = \frac{3}{2} k_B T}$ (a crucial relation)

This is the relation between kinetic energy and temperature for systems of **monatomic molecules**. The position of each molecule is determined by

3. Kinetic theory of gases

♣ Average kinetic energy and thermodynamic temperature

○ The average value of kinetic energy: $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$

○ Since $pV \equiv Nk_B T = \frac{1}{3} N m \langle c^2 \rangle \rightarrow m \langle c^2 \rangle = 3k_B T$

● Therefore: $\boxed{\langle E_k \rangle = \frac{3}{2} k_B T}$ (a crucial relation)

This is the relation between kinetic energy and temperature for systems of **monatomic molecules**. The position of each molecule is determined by three independent parameters.

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

- For monoatomic gas: $i = 3$.

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

- For monoatomic gas: $i = 3$.
- For diatomic gas:

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

- For monoatomic gas: $i = 3$.
- For diatomic gas: $i = 5$.

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

- For monoatomic gas: $i = 3$.
- For diatomic gas: $i = 5$.
- For polyatomic gas (three or more atoms):

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

- For monoatomic gas: $i = 3$.
- For diatomic gas: $i = 5$.
- For polyatomic gas (three or more atoms): $i = 6$.

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

- For monoatomic gas: $i = 3$.
- For diatomic gas: $i = 5$.
- For polyatomic gas (three or more atoms): $i = 6$.

♣ **Equipartition theorem** (law of equipartition of energy)

3. Kinetic theory of gases

The number of degrees of freedom, i , of a gas is the number of independent parameters used to determine the position of each molecule in the gas.

- For monoatomic gas: $i = 3$.
- For diatomic gas: $i = 5$.
- For polyatomic gas (three or more atoms): $i = 6$.

♣ **Equipartition theorem** (law of equipartition of energy)

In thermal equilibrium, the average energy is shared equally to every degree of freedom.

3. Kinetic theory of gases

- For monoatomic gas:

3. Kinetic theory of gases

- For monoatomic gas: $\langle E_k \rangle = \frac{3}{2} k_B T$

3. Kinetic theory of gases

- For monoatomic gas: $\langle E_k \rangle = \frac{3}{2} k_B T$ and $i = 3$.

3. Kinetic theory of gases

- For monoatomic gas: $\langle E_k \rangle = \frac{3}{2} k_B T$ and $i = 3$.
- For each degree of freedom:

3. Kinetic theory of gases

- For monoatomic gas: $\langle E_k \rangle = \frac{3}{2} k_B T$ and $i = 3$.
- For each degree of freedom: $\langle E_k \rangle = \frac{1}{2} k_B T$.

3. Kinetic theory of gases

- For monoatomic gas: $\langle E_k \rangle = \frac{3}{2} k_B T$ and $i = 3$.
- For each degree of freedom: $\langle E_k \rangle = \frac{1}{2} k_B T$.
- For gas of number of degrees of freedom i :

3. Kinetic theory of gases

- For monoatomic gas: $\langle E_k \rangle = \frac{3}{2} k_B T$ and $i = 3$.
- For each degree of freedom: $\langle E_k \rangle = \frac{1}{2} k_B T$.
- For gas of number of degrees of freedom i :

$$\langle E_k \rangle = \frac{i}{2} k_B T$$

3. Kinetic theory of gases

♣ **Root-mean-square speed**

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity is zero.

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity is zero. But the mean speed

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero.

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

$$\langle c \rangle \neq 0 \quad \text{and} \quad \langle c \rangle^2 \neq \langle c^2 \rangle$$

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

$$\langle c \rangle \neq 0 \quad \text{and} \quad \langle c \rangle^2 \neq \langle c^2 \rangle$$

The root-mean-square speed

3. Kinetic theory of gases

♣ **Root-mean-square speed**

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

$$\langle c \rangle \neq 0 \quad \text{and} \quad \langle c \rangle^2 \neq \langle c^2 \rangle$$

The root-mean-square speed is defined as

$$c_{\text{r.m.s.}} = \sqrt{\langle c^2 \rangle}$$

3. Kinetic theory of gases

♣ Root-mean-square speed

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

$$\langle c \rangle \neq 0 \quad \text{and} \quad \langle c \rangle^2 \neq \langle c^2 \rangle$$

The root-mean-square speed is defined as

$$c_{\text{r.m.s.}} = \sqrt{\langle c^2 \rangle}$$

- $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$

3. Kinetic theory of gases

♣ Root-mean-square speed

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

$$\langle c \rangle \neq 0 \quad \text{and} \quad \langle c \rangle^2 \neq \langle c^2 \rangle$$

The root-mean-square speed is defined as

$$c_{\text{r.m.s.}} = \sqrt{\langle c^2 \rangle}$$

- $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$ and $\langle E_k \rangle = \frac{3}{2} k_B T$

3. Kinetic theory of gases

♣ Root-mean-square speed

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

$$\langle c \rangle \neq 0 \quad \text{and} \quad \langle c \rangle^2 \neq \langle c^2 \rangle$$

The root-mean-square speed is defined as

$$c_{\text{r.m.s.}} = \sqrt{\langle c^2 \rangle}$$

- $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$ and $\langle E_k \rangle = \frac{3}{2} k_B T \rightarrow$

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}}$$

3. Kinetic theory of gases

♣ Root-mean-square speed

- In the random motion of the molecules, the mean velocity is zero. But the mean speed is not zero. Moreover, the square of mean speed is not the same as the mean-square speed.

$$\langle c \rangle \neq 0 \quad \text{and} \quad \langle c \rangle^2 \neq \langle c^2 \rangle$$

The root-mean-square speed is defined as

$$c_{\text{r.m.s.}} = \sqrt{\langle c^2 \rangle}$$

- $\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$ and $\langle E_k \rangle = \frac{3}{2} k_B T \rightarrow$

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}}$$

- Normally: $c_{\text{r.m.s.}} \approx 1.1 \times \langle c \rangle.$

3. Kinetic theory of gases

♣ Internal energy

3. Kinetic theory of gases

♣ Internal energy

Internal energy of a thermodynamic system

3. Kinetic theory of gases

♣ Internal energy

Internal energy of a thermodynamic system is determined by the state of the system

3. Kinetic theory of gases

♣ Internal energy

Internal energy of a thermodynamic system is determined by the state of the system and can be expressed as the sum of a random distribution of kinetic and potential energies associated with the molecules of the system.

3. Kinetic theory of gases

♣ Internal energy

Internal energy of a thermodynamic system is determined by the state of the system and can be expressed as the sum of a random distribution of kinetic and potential energies associated with the molecules of the system.

$$\boxed{U = \sum E_k + \sum E_p} \quad \left\{ \begin{array}{l} E_k \rightarrow \text{random motions of molecules} \\ E_p \rightarrow \text{interactions between molecules} \end{array} \right.$$

3. Kinetic theory of gases

♣ Internal energy

Internal energy of a thermodynamic system is determined by the state of the system and can be expressed as the sum of a random distribution of kinetic and potential energies associated with the molecules of the system.

$$\boxed{U = \sum E_k + \sum E_p} \quad \left\{ \begin{array}{l} E_k \rightarrow \text{random motions of molecules} \\ E_p \rightarrow \text{interactions between molecules} \end{array} \right.$$

- It does not include the kinetic energy in motion of the system as a whole,

3. Kinetic theory of gases

♣ Internal energy

Internal energy of a thermodynamic system is determined by the state of the system and can be expressed as the sum of a random distribution of kinetic and potential energies associated with the molecules of the system.

$$\boxed{U = \sum E_k + \sum E_p} \quad \left\{ \begin{array}{l} E_k \rightarrow \text{random motions of molecules} \\ E_p \rightarrow \text{interactions between molecules} \end{array} \right.$$

- It does not include the kinetic energy in motion of the system as a whole, nor the potential energy due to external forces.

3. Kinetic theory of gases

♣ **Internal energy**

- For ideal gases, since the molecules do not interact each other (except during collisions),

3. Kinetic theory of gases

♣ Internal energy

- For ideal gases, since the molecules do not interact each other (except during collisions), the potential energy term is zero.

3. Kinetic theory of gases

♣ Internal energy

- For ideal gases, since the molecules do not interact each other (except during collisions), the potential energy term is zero.

$$U = \sum E_k$$

3. Kinetic theory of gases

♣ Internal energy

- For ideal gases, since the molecules do not interact each other (except during collisions), the potential energy term is zero.

$$U = \sum E_k = N \langle E_k \rangle$$

3. Kinetic theory of gases

♣ Internal energy

- For ideal gases, since the molecules do not interact each other (except during collisions), the potential energy term is zero.

$$U = \sum E_k = N \langle E_k \rangle = \frac{m}{\mu} N_A \frac{i}{2} k_B T$$

3. Kinetic theory of gases

♣ Internal energy

- For ideal gases, since the molecules do not interact each other (except during collisions), the potential energy term is zero.

$$U = \sum E_k = N \langle E_k \rangle = \frac{m}{\mu} N_A \frac{i}{2} k_B T$$

$$U = \frac{m}{\mu} \frac{i}{2} RT$$

3. Kinetic theory of gases

♣ Internal energy

- For ideal gases, since the molecules do not interact each other (except during collisions), the potential energy term is zero.

$$U = \sum E_k = N \langle E_k \rangle = \frac{m}{\mu} N_A \frac{i}{2} k_B T$$

$$U = \frac{m}{\mu} \frac{i}{2} RT$$

- Here, μ is the molar mass and m is the mass of the system.

3. Kinetic theory of gases

- 1** The speed of seven molecules in a gas are numerically equal to 2, 4, 6, 8, 10, 12, and 14 units. Find the numerical values of
- (a) the mean speed $\langle c \rangle$,
 - (b) the mean speed squared $\langle c \rangle^2$,
 - (c) the mean-square speed $\langle c^2 \rangle$,
 - (d) the r.m.s. speed $\sqrt{\langle c^2 \rangle}$.

3. Kinetic theory of gases

- 1** The speed of seven molecules in a gas are numerically equal to 2, 4, 6, 8, 10, 12, and 14 units. Find the numerical values of
- (a) the mean speed $\langle c \rangle$, [8.0 units]
 - (b) the mean speed squared $\langle c \rangle^2$,
 - (c) the mean-square speed $\langle c^2 \rangle$,
 - (d) the r.m.s. speed $\sqrt{\langle c^2 \rangle}$.

3. Kinetic theory of gases

- 1** The speed of seven molecules in a gas are numerically equal to 2, 4, 6, 8, 10, 12, and 14 units. Find the numerical values of
- (a) the mean speed $\langle c \rangle$, [8.0 units]
 - (b) the mean speed squared $\langle c \rangle^2$, [64 units²]
 - (c) the mean-square speed $\langle c^2 \rangle$,
 - (d) the r.m.s. speed $\sqrt{\langle c^2 \rangle}$.

3. Kinetic theory of gases

- 1** The speed of seven molecules in a gas are numerically equal to 2, 4, 6, 8, 10, 12, and 14 units. Find the numerical values of
- (a) the mean speed $\langle c \rangle$, [8.0 units]
 - (b) the mean speed squared $\langle c \rangle^2$, [64 units²]
 - (c) the mean-square speed $\langle c^2 \rangle$, [80 units²]
 - (d) the r.m.s. speed $\sqrt{\langle c^2 \rangle}$.

3. Kinetic theory of gases

- 1** The speed of seven molecules in a gas are numerically equal to 2, 4, 6, 8, 10, 12, and 14 units. Find the numerical values of
- (a) the mean speed $\langle c \rangle$, [8.0 units]
 - (b) the mean speed squared $\langle c \rangle^2$, [64 units²]
 - (c) the mean-square speed $\langle c^2 \rangle$, [80 units²]
 - (d) the r.m.s. speed $\sqrt{\langle c^2 \rangle}$. [8.9 units]

3. Kinetic theory of gases

- 2** Find the total kinetic energy of molecules in one mole of ideal gas at standard temperature (273 K).

Take $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$.

- 3** The mass of a nitrogen molecule is $4.6 \times 10^{-26} \text{ kg}$. Find the root-mean-square speed of molecules in nitrogen gas at 27°C .

Take $k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$.

3. Kinetic theory of gases

- 2** Find the total kinetic energy of molecules in one mole of ideal gas at standard temperature (273 K).

Take $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$.

$$E_k^{(\text{tot})} = N_A \langle E_k \rangle = N_A \left(\frac{3}{2} k_B T \right) = \frac{3}{2} RT = \frac{3}{2} \times 8.3 \times 273 = 3400 \text{ J}$$

- 3** The mass of a nitrogen molecule is $4.6 \times 10^{-26} \text{ kg}$. Find the root-mean-square speed of molecules in nitrogen gas at 27°C .

Take $k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$.

3. Kinetic theory of gases

- 2** Find the total kinetic energy of molecules in one mole of ideal gas at standard temperature (273 K).

Take $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$.

$$E_k^{(\text{tot})} = N_A \langle E_k \rangle = N_A \left(\frac{3}{2} k_B T \right) = \frac{3}{2} RT = \frac{3}{2} \times 8.3 \times 273 = 3400 \text{ J}$$

- 3** The mass of a nitrogen molecule is $4.6 \times 10^{-26} \text{ kg}$. Find the root-mean-square speed of molecules in nitrogen gas at 27°C .

Take $k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$.

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}} = \sqrt{\frac{3 \times 1.38 \times 10^{-23} \times 300}{4.6 \times 10^{-26}}} = 520 \text{ m s}^{-1}$$

2. THERMODYNAMICS

2.1 KINETIC THEORY AND LAWS OF DISTRIBUTION

- 1 **STARTING POINTS**
- 2 **EQUATION OF STATE**
- 3 **KINETIC THEORY OF GASES**
 - Exercises
- 4 **MAXWELL DISTRIBUTION**
- 5 **BOLTZMANN DISTRIBUTION**

4. Maxwell distribution

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed:

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i ,

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.
- Probability for an arbitrary particle to have speed c_i :

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.
- Probability for an arbitrary particle to have speed c_i :

$$p_i = \frac{N_i}{N}$$

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.
- Probability for an arbitrary particle to have speed c_i :

$$\boxed{p_i = \frac{N_i}{N}} \quad \rightarrow \quad \sum_{i=1}^n p_i =$$

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots, n$.
- Probability for an arbitrary particle to have speed c_i :

$$p_i = \frac{N_i}{N}$$

 \rightarrow

$$\sum_{i=1}^n p_i = 1$$

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.
- Probability for an arbitrary particle to have speed c_i :

$$p_i = \frac{N_i}{N}$$

 \rightarrow

$$\sum_{i=1}^n p_i = 1$$

(normalized condition)

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.
- Probability for an arbitrary particle to have speed c_i :

$$\boxed{p_i = \frac{N_i}{N}} \quad \rightarrow \quad \boxed{\sum_{i=1}^n p_i = 1} \quad \text{(normalized condition)}$$

- Mean speed of the particles:

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.
- Probability for an arbitrary particle to have speed c_i :

$$\boxed{p_i = \frac{N_i}{N}} \quad \rightarrow \quad \boxed{\sum_{i=1}^n p_i = 1} \quad \text{(normalized condition)}$$

- Mean speed of the particles:

$$\langle c \rangle = \frac{1}{N} \sum N_i c_i$$

4. Maxwell distribution

♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.

- Probability for an arbitrary particle to have speed c_i :

$$\boxed{p_i = \frac{N_i}{N}} \quad \rightarrow \quad \boxed{\sum_{i=1}^n p_i = 1} \quad \text{(normalized condition)}$$

- Mean speed of the particles:

$$\langle c \rangle = \frac{1}{N} \sum N_i c_i = \sum \frac{N_i}{N} c_i$$

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots, n$.
- Probability for an arbitrary particle to have speed c_i :

$$\boxed{p_i = \frac{N_i}{N}} \quad \rightarrow \quad \boxed{\sum_{i=1}^n p_i = 1} \quad \text{(normalized condition)}$$

- Mean speed of the particles:

$$\langle c \rangle = \frac{1}{N} \sum N_i c_i = \sum \frac{N_i}{N} c_i \quad \rightarrow \quad \boxed{\langle c \rangle = \sum p_i c_i}$$

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots, n$.
- Probability for an arbitrary particle to have speed c_i :

$$\boxed{p_i = \frac{N_i}{N}} \quad \rightarrow \quad \boxed{\sum_{i=1}^n p_i = 1} \quad \text{(normalized condition)}$$

- Mean speed of the particles:

$$\langle c \rangle = \frac{1}{N} \sum N_i c_i = \sum \frac{N_i}{N} c_i \quad \rightarrow \quad \boxed{\langle c \rangle = \sum p_i c_i}$$

The mean value of any function $f(c)$:

4. Maxwell distribution

- ♣ Imagine a system of N particles which have discrete values of speed: N_i particles have speed c_i , $i = 1, 2, 3, \dots n$.
- Probability for an arbitrary particle to have speed c_i :

$$\boxed{p_i = \frac{N_i}{N}} \quad \rightarrow \quad \boxed{\sum_{i=1}^n p_i = 1} \quad \textbf{(normalized condition)}$$

- Mean speed of the particles:

$$\langle c \rangle = \frac{1}{N} \sum N_i c_i = \sum \frac{N_i}{N} c_i \quad \rightarrow \quad \boxed{\langle c \rangle = \sum p_i c_i}$$

The mean value of any function $f(c)$:

$$\boxed{\langle f \rangle = \sum p_i f(c_i)}$$

4. Maxwell distribution

♣ For a real gas system,

4. Maxwell distribution

- ♣ For a real gas system, the speed of the particles is continuously distributed

4. Maxwell distribution


- ♣ For a real gas system, the speed of the particles is continuously distributed (can have any value).

4. Maxwell distribution

- ♣ For a real gas system, the speed of the particles is continuously distributed (can have any value). The distribution depends on the temperature and properties of the system.

4. Maxwell distribution

- ♣ For a real gas system, the speed of the particles is continuously distributed (can have any value). The distribution depends on the temperature and properties of the system.



4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$.

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles:

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range:

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,
 - and the value c of the speed at which the range is chosen.

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,
 - and the value c of the speed at which the range is chosen.
- Therefore:

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,
 - and the value c of the speed at which the range is chosen.
- Therefore: $dN = F(c)Ndc$

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,
 - and the value c of the speed at which the range is chosen.
- Therefore: $dN = F(c)Ndc$
- The value of $F(c)dc$

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,
 - and the value c of the speed at which the range is chosen.
- Therefore:
$$dN = F(c)Ndc$$
- The value of $F(c)dc = \frac{dN}{N}$

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,
 - and the value c of the speed at which the range is chosen.
- Therefore:
$$dN = F(c)Ndc$$
- The value of $F(c)dc = \frac{dN}{N}$ is the probability for any particle of the system to have speed in the range $(c, c + dc)$.

4. Maxwell distribution

- Suppose that there are dN particles which have speed in the range $(c, c + dc)$. The value of dN depends on:
 - the total number N of particles: $dN \propto N$,
 - the width of the range: $dN \propto dc$,
 - and the value c of the speed at which the range is chosen.
- Therefore:
$$dN = F(c)Ndc$$
- The value of $F(c)dc = \frac{dN}{N}$ is the probability for any particle of the system to have speed in the range $(c, c + dc)$.
- The function $F(c)$ is called the **distribution function** of the molecules with speed.

4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} =$$

4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1$$

4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^\infty F(c)dc = 1}$$

4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^\infty F(c)dc = 1} \quad (\text{normalized condition})$$

4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^\infty F(c)dc = 1} \quad (\text{normalized condition})$$


- Maxwell distribution function**

4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^{\infty} F(c) dc = 1} \quad (\text{normalized condition})$$

- Maxwell distribution function**




4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^{\infty} F(c) dc = 1} \quad (\text{normalized condition})$$

- Maxwell distribution function**
 - The function depends on kinetic energy,




4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^{\infty} F(c) dc = 1} \quad (\text{normalized condition})$$

- Maxwell distribution function**

- The function depends on kinetic energy, i.e. depends on c^2 .




4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^{\infty} F(c) dc = 1} \quad (\text{normalized condition})$$

- Maxwell distribution function**

- The function depends on kinetic energy, i.e. depends on c^2 .
- First part $\propto c^2$.




4. Maxwell distribution

- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^{\infty} F(c) dc = 1} \quad (\text{normalized condition})$$

- Maxwell distribution function**

- The function depends on kinetic energy, i.e. depends on c^2 .
- First part $\propto c^2$.
- Second part decays rapidly.



4. Maxwell distribution


- For the whole system:

$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^{\infty} F(c) dc = 1} \quad (\text{normalized condition})$$

- **Maxwell distribution function**

- The function depends on kinetic energy, i.e. depends on c^2 .
- First part $\propto c^2$.
- Second part decays rapidly.

$$F(c) = A c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$



4. Maxwell distribution

- For the whole system:


$$\int \frac{dN}{N} = 1 \quad \rightarrow \quad \boxed{\int_0^{\infty} F(c) dc = 1} \quad (\text{normalized condition})$$

- **Maxwell distribution function**

- The function depends on kinetic energy, i.e. depends on c^2 .
- First part $\propto c^2$.
- Second part decays rapidly.

$$F(c) = A c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$

A and α are positive constants.



4. Maxwell distribution

- The constant α depends on temperature

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^{\infty} F(c)dc$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^{\infty} F(c)dc = \int_0^{\infty} Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^{\infty} F(c)dc = \int_0^{\infty} Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^{\infty} F(c)dc = \int_0^{\infty} Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**,

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^{\infty} F(c)dc = \int_0^{\infty} Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^{\infty} e^{-\alpha x^2/2} dx$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy,$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^{\infty} F(c)dc = \int_0^{\infty} Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^{\infty} e^{-\alpha x^2/2} dx = \int_0^{\infty} e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} =$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^{\infty} F(c)dc = \int_0^{\infty} Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^{\infty} e^{-\alpha x^2/2} dx = \int_0^{\infty} e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

$$I_0^2 =$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

$$I_0^2 = \int_{x=0}^\infty \int_{y=0}^\infty e^{-\alpha(x^2+y^2)/2} dx dy$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

$$I_0^2 = \int_{x=0}^\infty \int_{y=0}^\infty e^{-\alpha(x^2+y^2)/2} dx dy = \int_{r=0}^\infty \int_{\phi=0}^{\pi/2} e^{-\alpha r^2/2} r dr d\phi$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

$$I_0^2 = \int_{x=0}^\infty \int_{y=0}^\infty e^{-\alpha(x^2+y^2)/2} dx dy = \int_{r=0}^\infty \int_{\phi=0}^{\pi/2} e^{-\alpha r^2/2} r dr d\phi = \frac{\pi}{2\alpha}$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

$$I_0^2 = \int_{x=0}^\infty \int_{y=0}^\infty e^{-\alpha(x^2+y^2)/2} dx dy = \int_{r=0}^\infty \int_{\phi=0}^{\pi/2} e^{-\alpha r^2/2} r dr d\phi = \frac{\pi}{2\alpha}$$

$$I_0 = \sqrt{\frac{\pi}{2\alpha}}$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

$$I_0^2 = \int_{x=0}^\infty \int_{y=0}^\infty e^{-\alpha(x^2+y^2)/2} dx dy = \int_{r=0}^\infty \int_{\phi=0}^{\pi/2} e^{-\alpha r^2/2} r dr d\phi = \frac{\pi}{2\alpha}$$

$$I_0 = \sqrt{\frac{\pi}{2\alpha}} \quad \rightarrow \quad I_2 = -2\frac{\partial I_0}{\partial \alpha}$$

4. Maxwell distribution

- The constant α depends on temperature and the constant A is determined by the normalized condition.

$$1 = \int_0^\infty F(c)dc = \int_0^\infty Ac^2 \exp\left(-\frac{\alpha c^2}{2}\right)dc = AI_2$$

- Use the **gamma function**, or

$$I_0 = \int_0^\infty e^{-\alpha x^2/2} dx = \int_0^\infty e^{-\alpha y^2/2} dy, \quad \frac{\partial I_0}{\partial \alpha} = -\frac{1}{2}I_2$$

$$I_0^2 = \int_{x=0}^\infty \int_{y=0}^\infty e^{-\alpha(x^2+y^2)/2} dx dy = \int_{r=0}^\infty \int_{\phi=0}^{\pi/2} e^{-\alpha r^2/2} r dr d\phi = \frac{\pi}{2\alpha}$$

$$I_0 = \sqrt{\frac{\pi}{2\alpha}} \quad \rightarrow \quad I_2 = -2\frac{\partial I_0}{\partial \alpha} = \sqrt{\frac{\pi}{2\alpha^3}}$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}}$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}}$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}} = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}} = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi$$

$$F(c) = \sqrt{\frac{2\alpha^3}{\pi}} c^2 \exp\left(-\frac{\alpha c^2}{2}\right) = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}} = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi$$

$$F(c) = \sqrt{\frac{2\alpha^3}{\pi}} c^2 \exp\left(-\frac{\alpha c^2}{2}\right) = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$

♣ Mean values of a function $f(c)$:

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}} = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi$$

$$F(c) = \sqrt{\frac{2\alpha^3}{\pi}} c^2 \exp\left(-\frac{\alpha c^2}{2}\right) = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$

♣ Mean values of a function $f(c)$:

$$\langle f \rangle = \sum f(c_i) p_i$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}} = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi$$

$$F(c) = \sqrt{\frac{2\alpha^3}{\pi}} c^2 \exp\left(-\frac{\alpha c^2}{2}\right) = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$

♣ Mean values of a function $f(c)$:

$$\langle f \rangle = \sum f(c_i) p_i \quad \rightarrow \quad \langle f \rangle = \int f(c) dp$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}} = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi$$

$$F(c) = \sqrt{\frac{2\alpha^3}{\pi}} c^2 \exp\left(-\frac{\alpha c^2}{2}\right) = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$

♣ Mean values of a function $f(c)$:

$$\langle f \rangle = \sum f(c_i) p_i \quad \rightarrow \quad \langle f \rangle = \int f(c) dp = \int f(c) \frac{dN}{N}$$

4. Maxwell distribution

$$1 = A \sqrt{\frac{\pi}{2\alpha^3}} \quad \rightarrow \quad A = \sqrt{\frac{2\alpha^3}{\pi}} = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi$$

$$F(c) = \sqrt{\frac{2\alpha^3}{\pi}} c^2 \exp\left(-\frac{\alpha c^2}{2}\right) = \left(\frac{\alpha}{2\pi}\right)^{3/2} 4\pi c^2 \exp\left(-\frac{\alpha c^2}{2}\right)$$

♣ Mean values of a function $f(c)$:

$$\langle f \rangle = \sum f(c_i) p_i \quad \rightarrow \quad \langle f \rangle = \int f(c) dp = \int f(c) \frac{dN}{N}$$

$$\langle f \rangle = \int_0^\infty f(c) F(c) dc$$

4. Maxwell distribution

- Mean-square speed:

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle =$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

- Average kinetic energy:

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

- Average kinetic energy:

$$\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

- Average kinetic energy:

$$\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle = \frac{3}{2} k_B T$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

- Average kinetic energy:

$$\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle = \frac{3}{2} k_B T \quad \rightarrow \quad \frac{1}{2} m \frac{3}{\alpha} = \frac{3}{2} k_B T$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

- Average kinetic energy:

$$\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle = \frac{3}{2} k_B T \quad \rightarrow \quad \frac{1}{2} m \frac{3}{\alpha} = \frac{3}{2} k_B T \quad \rightarrow \quad \alpha = \frac{m}{k_B T}$$

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

- Average kinetic energy:

$$\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle = \frac{3}{2} k_B T \quad \rightarrow \quad \frac{1}{2} m \frac{3}{\alpha} = \frac{3}{2} k_B T \quad \rightarrow \quad \alpha = \frac{m}{k_B T}$$

- Explicit formula of the Maxwell distribution function:

4. Maxwell distribution

- Mean-square speed:

$$\langle c^2 \rangle = \int_0^\infty c^2 F(c) dc = \sqrt{\frac{2\alpha^3}{\pi}} \int_0^\infty c^4 e^{-\alpha c^2/2} dc = \frac{3}{\alpha}$$

- Average kinetic energy:

$$\langle E_k \rangle = \frac{1}{2} m \langle c^2 \rangle = \frac{3}{2} k_B T \quad \rightarrow \quad \frac{1}{2} m \frac{3}{\alpha} = \frac{3}{2} k_B T \quad \rightarrow \quad \alpha = \frac{m}{k_B T}$$

- Explicit formula of the Maxwell distribution function:

$$F(c) = \left(\frac{m}{2\pi k_B T} \right)^{3/2} 4\pi c^2 \exp\left(-\frac{mc^2}{2k_B T} \right)$$

4. Maxwell distribution

♣ Typical speeds

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle =$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_{\text{B}}T}{\pi m}}$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_{\text{B}}T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}}$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_{\text{B}}T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_{\text{B}}T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_{\text{B}}T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} =$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_B T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}}$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_{\text{B}}T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_{\text{B}}T}{m}} = \sqrt{\frac{3RT}{\mu}}$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_B T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}} = \sqrt{\frac{3RT}{\mu}}$$

- **Most probable speed**

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_B T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}} = \sqrt{\frac{3RT}{\mu}}$$

- **Most probable speed** (at which $F(c)$ is maximum):

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_B T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}} = \sqrt{\frac{3RT}{\mu}}$$

- **Most probable speed** (at which $F(c)$ is maximum):

$$c_p =$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_B T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}} = \sqrt{\frac{3RT}{\mu}}$$

- **Most probable speed** (at which $F(c)$ is maximum):

$$c_p = \sqrt{\frac{2k_B T}{m}}$$

4. Maxwell distribution

♣ Typical speeds

- Mean speed:

$$\langle c \rangle = \sqrt{\frac{8k_B T}{\pi m}} = \sqrt{\frac{8RT}{\pi \mu}} \quad (\mu \text{ is the molar mass})$$

- Root-mean-square speed:

$$c_{\text{r.m.s.}} = \sqrt{\frac{3k_B T}{m}} = \sqrt{\frac{3RT}{\mu}}$$


- **Most probable speed** (at which $F(c)$ is maximum):

$$c_p = \sqrt{\frac{2k_B T}{m}} = \sqrt{\frac{2RT}{\mu}}$$

4. Maxwell distribution

♣ Typical speeds

$$f(v) = 4\pi \left[\frac{M}{2\pi RT} \right]^{\frac{3}{2}} v^2 \exp \left[\frac{-Mv^2}{2RT} \right]$$



2. THERMODYNAMICS

2.1 KINETIC THEORY AND LAWS OF DISTRIBUTION

- 1 STARTING POINTS
- 2 EQUATION OF STATE
- 3 KINETIC THEORY OF GASES
 - Exercises
- 4 MAXWELL DISTRIBUTION
- 5 **BOLTZMANN DISTRIBUTION**

5. Boltzmann distribution

5. Boltzmann distribution

♣ Atmospheric pressure

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas.

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp =$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

$$pdV = \frac{dM}{\mu} RT$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

$$pdV = \frac{dM}{\mu} RT \quad \rightarrow \quad \rho = \frac{dM}{dV}$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

$$pdV = \frac{dM}{\mu} RT \quad \rightarrow \quad \rho = \frac{dM}{dV} = \frac{p\mu}{RT}$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

$$p dV = \frac{dM}{\mu} RT \quad \rightarrow \quad \rho = \frac{dM}{dV} = \frac{p\mu}{RT}$$

$$\frac{dp}{p} = -\frac{\mu g}{RT} dz$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

$$p dV = \frac{dM}{\mu} RT \quad \rightarrow \quad \rho = \frac{dM}{dV} = \frac{p\mu}{RT}$$

$$\frac{dp}{p} = -\frac{\mu g}{RT} dz \quad \rightarrow \quad \boxed{p = p_0 \exp\left(-\frac{\mu g z}{RT}\right)}$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

$$p dV = \frac{dM}{\mu} RT \quad \rightarrow \quad \rho = \frac{dM}{dV} = \frac{p\mu}{RT}$$

$$\frac{dp}{p} = -\frac{\mu g}{RT} dz \quad \rightarrow \quad \boxed{p = p_0 \exp\left(-\frac{\mu g z}{RT}\right)} \quad \left\{ \begin{array}{l} \text{Atmospheric} \\ \text{pressure} \\ \text{formula} \end{array} \right.$$

5. Boltzmann distribution

♣ Atmospheric pressure

- Suppose that the gas in the uniform gravitational field of the Earth is an ideal gas. Let p be the pressure at height z :

$$dp = -\rho g dz$$

- For a small amount (mass dM and volume dV) of gas at z :

$$p dV = \frac{dM}{\mu} RT \quad \rightarrow \quad \rho = \frac{dM}{dV} = \frac{p\mu}{RT}$$

$$\frac{dp}{p} = -\frac{\mu g}{RT} dz \quad \rightarrow \quad \boxed{p = p_0 \exp\left(-\frac{\mu g z}{RT}\right)} \quad \left\{ \begin{array}{l} \text{Atmospheric} \\ \text{pressure} \\ \text{formula} \end{array} \right.$$

Here, p_0 is the pressure at $z = 0$.

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n$$

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n \quad \rightarrow \quad n = n_0 \exp\left(-\frac{\mu g z}{RT}\right)$$

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n \quad \rightarrow \quad n = n_0 \exp\left(-\frac{\mu g z}{RT}\right) = n_0 \exp\left(-\frac{m g z}{k_B T}\right)$$

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n \quad \rightarrow \quad n = n_0 \exp\left(-\frac{\mu g z}{RT}\right) = n_0 \exp\left(-\frac{m g z}{k_B T}\right)$$

- Here, μ is the molar mass,

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n \quad \rightarrow \quad n = n_0 \exp\left(-\frac{\mu g z}{RT}\right) = n_0 \exp\left(-\frac{m g z}{k_B T}\right)$$

- Here, μ is the molar mass, m is the molecular mass,

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n \quad \rightarrow \quad n = n_0 \exp\left(-\frac{\mu g z}{RT}\right) = n_0 \exp\left(-\frac{m g z}{k_B T}\right)$$

- Here, μ is the molar mass, m is the molecular mass, and $m g z$ is the gravitational potential energy of the molecule.

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n \quad \rightarrow \quad n = n_0 \exp\left(-\frac{\mu g z}{RT}\right) = n_0 \exp\left(-\frac{m g z}{k_B T}\right)$$

- Here, μ is the molar mass, m is the molecular mass, and $m g z$ is the gravitational potential energy of the molecule.

$$n = n_0 \exp\left(-\frac{E_p}{k_B T}\right)$$

5. Boltzmann distribution

- ♣ Let n be the molecular density (number of molecules per unit volume):

$$p \propto n \quad \rightarrow \quad n = n_0 \exp\left(-\frac{\mu g z}{RT}\right) = n_0 \exp\left(-\frac{mgz}{k_B T}\right)$$

- Here, μ is the molar mass, m is the molecular mass, and mgz is the gravitational potential energy of the molecule.

$$n = n_0 \exp\left(-\frac{E_p}{k_B T}\right)$$

- This is called the **Boltzmann distribution**, the distribution of particles in term of potential energy.